


4C
770
U64
V.26
1972
Abs. no. 45181-50202

NUCLEAR SCIENCE ABSTRACTS



U. of ILL. LIBRARY

DEC 21 1972

CHICAGO CIRCLE

Volume 26 Number 19

Abstracts 45181-47652

October 15, 1972

U. OF I. LIB.
DOCUMENTS

OCT 30 1972

CHICAGO CIRCLE

UNITED STATES ATOMIC ENERGY COMMISSION

REPORT COLLECTIONS IN THE UNITED STATES

The libraries listed below purchase and maintain microfiche collections of USAEC and foreign reports that are abstracted in *NSA*. Their collections also include *NSA* and its indexes, AEC's Technical Progress Reviews, AEC-sponsored translations, conference proceedings published by AEC, and various bibliographic tools. Most of these libraries have microfiche reader-printers or other photocopy facilities with which to reproduce enlarged copies from microfiche. Charges for reproduction services vary.

ALABAMA

Auburn, Auburn University
Tuskegee Institute, Tuskegee Institute

ALASKA

College, University of Alaska

ARIZONA

Tucson, University of Arizona

CALIFORNIA

Davis, University of California
Los Angeles, University of California
Santa Barbara, University of California

COLORADO

Boulder, University of Colorado

DISTRICT OF COLUMBIA

Washington, Library of Congress

FLORIDA

Gainesville, University of Florida

GEORGIA

Atlanta, Georgia Institute of
Technology

HAWAII

Honolulu, University of Hawaii

IDAHO

Pocatello, Idaho State University

ILLINOIS

Urbana, University of Illinois

INDIANA

Lafayette, Purdue University

IOWA

Ames, Iowa State University

KANSAS

Manhattan, Kansas State University

KENTUCKY

Lexington, University of Kentucky
Louisville, University of Louisville

MARYLAND

Baltimore, Johns Hopkins University
College Park, University of Maryland

MASSACHUSETTS

Cambridge, Massachusetts Institute of
Technology
Worcester, Worcester Polytechnic Institute

MICHIGAN

Ann Arbor, University of Michigan

MISSISSIPPI

State College, Mississippi State University

MISSOURI

Columbia, University of Missouri
Kansas City, Linda Hall Library

NEW JERSEY

Princeton, Princeton University

NEW MEXICO

Albuquerque, University of New Mexico

NEW YORK

Albany, New York State Library
Albany, State University at Albany
Ithaca, Cornell University
New York, Columbia University
Rochester, University of Rochester
Syracuse, Syracuse University
Troy, Rensselaer Polytechnic Institute

NORTH CAROLINA

Raleigh, North Carolina State University

OHIO

Cincinnati, University of Cincinnati
Cleveland, Cleveland Public Library

OHIO (Continued)

Columbus, Ohio State University
Toledo, University of Toledo

OKLAHOMA

Norman, University of Oklahoma

PENNSYLVANIA

Philadelphia, University of Pennsylvania
Pittsburgh, Carnegie Library
University Park, Pennsylvania State
University

PUERTO RICO

San Juan, University of Puerto Rico

SOUTH CAROLINA

Columbia, University of South Carolina

TENNESSEE

Knoxville, University of Tennessee
Nashville, Joint University Libraries

TEXAS

Austin, University of Texas
College Station, Texas A & M University
Houston, Rice University

UTAH

Salt Lake City, University of Utah

VIRGINIA

Blacksburg, Virginia Polytechnic Institute
Charlottesville, University of Virginia

WASHINGTON

Pullman, Washington State University
Seattle, University of Washington

WEST VIRGINIA


Morgantown, West Virginia University

WISCONSIN

Madison, University of Wisconsin

The following governmental and international organizations offer for sale USAEC reports abstracted in *Nuclear Science Abstracts*:

National Technical Information Service, U. S. Department of Commerce
Springfield, Virginia 22151
International Atomic Energy Agency, Vienna, Austria
National Lending Library, Boston Spa, England



Volume 26, Number 19, October 15, 1972

NUCLEAR SCIENCE ABSTRACTS

A semimonthly publication of the USAEC Office of Information Services, John A. Harris, Director. Published by the USAEC Technical Information Center, Robert L. Shannon, Manager.

Nuclear Science Abstracts provides the only comprehensive abstracting and indexing coverage of international nuclear science literature. *Nuclear Science Abstracts* covers scientific and technical reports of the U. S. Atomic Energy Commission and its contractors, other U. S. Government agencies, other governments, universities, and industrial and research organizations. In addition, books, conference proceedings, individual conference papers, patents, and journal literature on a worldwide basis are abstracted and indexed. Each issue of *Nuclear Science Abstracts* includes four indexes: subject, personal author, corporate author, and report number.

The following organizations scan and abstract the nuclear science literature in their respective countries:

Atomic Energy of Canada Limited
Australian Atomic Energy Commission
Danish Atomic Energy Commission
Finnish Atomic Energy Commission
French Atomic Energy Commission
Japan Atomic Energy Bureau-Atomic Energy Research Institute
Institut for Atomenergi (Norway)
Aktiebolaget Atomenergi (Sweden)
United Kingdom Atomic Energy Authority
Zentralstelle für Atomkernenergie-Dokumentation (West Germany)

Numerous organizations, societies, and publishers throughout the world provide *Nuclear Science Abstracts* with advance page-proof copy of their publications, e.g., in the United States the American Institute of Physics provides page-proof copy of its publications.

A list of the journals scanned regularly and the abbreviations used appears in Issue 1 (January 15) of each volume of *Nuclear Science Abstracts* and is supplemented in Issues 7, 13, and 19. Issue 1 of each volume also describes the scope and arrangement of the subject matter covered in *Nuclear Science Abstracts*. The subject matter covered is also described in TID-4552, *Subject Scope of Nuclear Science Abstracts*, available free from the USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830. An outline of the subject categories appears in the Contents of each issue.

AVAILABILITY OF PUBLICATIONS ABSTRACTED

Public availability of all U. S. Atomic Energy Commission reports is indicated in the report number indexes. The availability of other reports, when known, is also indicated. The availability of some unnumbered reports is given in the abstract.

Dissertations abstracted in *Dissertation Abstracts International, Part B* and included in *Nuclear Science Abstracts* are available from University Microfilms, Inc., 315 N. First Street, Ann Arbor, Michigan 48103.

Information on the availability of the published literature and unpublished conference papers abstracted in *Nuclear Science*

Abstracts can be obtained from the USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830.

HOW TO OBTAIN NUCLEAR SCIENCE ABSTRACTS

Nuclear Science Abstracts is available to the public on a subscription basis from the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402.

For the 24 regular issues, the yearly rate is \$42.00 for domestic subscribers and \$52.50 for foreign subscribers. A single issue costs \$1.75, domestic rates, or \$2.19, foreign rates. Domestic rates apply to the United States, Canada, Mexico, and Central and South American countries except Argentina, Brazil, Guyana, French Guiana, Surinam, and British Honduras.

For a yearly subscription to the cumulative indexes, the rates are \$34.50 for domestic subscribers and \$43.00 for foreign subscribers.

The following cumulative indexes are available:

From Johnson Reprint Corporation, 111 Fifth Avenue, New York, N. Y. 10003:

Vols. 1-4, Subjects and Authors (1948-1950), \$30.00.

From Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402:

Vols. 5-10, Subjects and Authors (1951-1956), \$5.25.

Vols. 11-15, Subjects (1957-1961), \$19.00.

Vols. 11-15, Personal Authors (1957-1961), \$14.25.

Vols. 11-15, Corporate Authors (1957-1961), \$2.50.

Vols. 16-20, Subjects (1962-1966), \$36.00.

Vols. 16-20, Personal Authors (1962-1966), \$22.50.

Vols. 16-20, Corporate Authors (1962-1966), \$3.75.

Vol. 21, Subjects and Authors (1967), \$16.00.

Vol. 22, Subjects and Authors (1968), \$26.00.

Vol. 23, Subjects and Authors (1969), \$25.75.

Vol. 24, Subjects and Authors (1970), \$22.50.

Vol. 24, Authors (1970), Suppl., \$2.00.

Vol. 25, Subjects and Authors (1971), \$27.50.

Cumulative Report Number Index, Vols. 1-15 Revised, \$4.00.

Cumulative Report Number Index, Vols. 16-20 Revised, \$5.50.

Cumulative Report Number Index, Vols. 21-25, \$5.75.

Nuclear Science Abstracts is available on an exchange basis to universities, research institutions, industrial firms, and publishers of scientific information. Government agencies may obtain *NSA* free of charge. Inquiries should be directed to the USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830.

Nuclear Science Abstracts is also available on 16-mm and 35-mm microfilm in reel or cartridge from Princeton Microfilm Corporation, Alexander Road, Princeton, New Jersey 08540. The cost of Vols. 1-24 (1948-1970) including *Abstracts of Declassified Documents* (1947-1948) is \$625.20. The indexes for *Nuclear Science Abstracts* Vols. 1-24 (1948-1970) are also available for \$320.00. The cost of each volume (one calendar year) of *NSA* and its indexes, including that for the year 1971, is \$95.

INDEXES TO NSA

Comprehensive indexes appear in each issue and are cumulated quarterly and annually. Multivolume cumulations have been issued as indicated above.

All documents abstracted are indexed by subjects and personal authors. Report literature is indexed also by corporate author and report number. Each of these indexes in every issue is preceded by an introduction explaining in detail the organization of the index and the principles by which it is compiled. The reader is referred to these explanations for information not found in the brief summary below.

Personal Authors

Each author's name is indexed in the form appearing on the document abstracted:

NEWCOMB, William A., Stability of a magnetic well, 25:1239
 NEWKIRK, L. R., Superconductivity in the pseudobinary alloys V_3Si-Nb_3Al , 25:1093 (CALT-822-13)
 NEWLAND, E. A., Radiation treatment of sewage, 25:140 (AAEC/LIB/81b-254)

Subjects

The subject index consisting of specific materials, objects, and processes is arranged alphabetically. Modifying phrases specific to these subjects in turn are arranged alphabetically under the subjects.

TRANSURANIUM ELEMENTS
 determination in bone ash, radiochemical method for, 25:40 (IDO-12073, pp 28-36)
 separation by ion exchange and complex-formation elution, 25:251(P)

TRIETHYLENETETRAMINE
 complexes with cobalt(III) and rhodium(III) chlorides, radiation effects on labeled, thermal-neutron, 25:137
 TRITIUM
 content in Cattaraugus Creek near West Valley Processing Plant, July through Dec. 1969, 25:258 (DOCKET-50201-48)
 content in Cattaraugus Creek near West Valley Processing Plant, Jan. to July 1970, 25:259 (DOCKET-50201-49)
 determination in luminescent material by scintillation counting, 25:40 (IDO-12073, pp 28-36)

Corporate Authors

Technical report literature is indexed from the name of the organization or institution responsible for the issuance of the report.

372 (UCRL-72739) CONSTRUCTION OF A UNIQUE UNDERGROUND WHOLE BODY COUNTER. Griffith, R. V.; Anderson, A. L.; Lindeken, C. L. (California Univ., Livermore. Lawrence Radiation Lab.). Oct 1970. 27p. (CONF-701009-1). Dep. NTIS.

is indexed as:

CALIFORNIA UNIV., LIVERMORE. LAWRENCE RADIATION LAB.
 Construction of a unique underground whole body counter, 25:372 (UCRL-72739)

Report Numbers

Technical report literature is also indexed by report numbers. This index includes information on where individual reports may be obtained. While not reports, patents and conference papers are indexed here as a matter of convenience.

ORNL-		
4270(Vol.2)	25-614	Dep. NTIS
4606	25-1530	Dep. NTIS
ORNL-IBP-		
70-3	25-2025	Dep. NTIS
ORNL-IIC-		
21	25-261	Dep. NTIS
30(Pt.2)	25-331	Dep. NTIS
ORNL-NSIC-		
78	25-1912	Dep. NTIS

TABLE OF CONTENTS

Volume 26, Number 19, October 15, 1972

Category	Page	Category	Page
CHEMISTRY	4349	LIFE SCIENCES	4409
Analytical Chemistry	4349	Radiation Effects on Biochemicals (External Source)	4409
Activation and Nuclear Reaction Procedures	4349	Radiation Effects on Cells (External Source)	4410
Chemical and Spectral Procedures	4352	Radiation Effects on Microorganisms (External Source)	4411
Radiometric and Radiochemical Procedures	4356	Radiation Effects on Plants (External Source)	4412
Separation Procedures	4357	Radiation Effects on Animals (External Source)	4415
Inorganic, Organic, and Physical Chemistry	4358	Man	4415
Chemical and Physico-chemical Properties	4358	Vertebrates	4418
Isotope Effects	4358	Invertebrates	4426
Isotope Exchange and Isotope Separation	4360	Radionuclide Effects (Internal Source)	4428
Feed Materials Processing	4360	Man	4428
Radiation Chemistry	4361	Animals	4428
Radiochemistry and Nuclear Chemistry	4366	Plants	4429
Hot-Atom Chemistry	4366	Nuclide Kinetics and Toxicology	4429
Properties of Radioactive Materials	4367	Man	4429
Radioactive Waste Treatment	4369	Animals	4430
Radioisotope Production	4370	Plants	4432
Reactor Fuel Reprocessing	4370	New Tracer Techniques	4432
ENGINEERING	4375	MATERIALS	4433
Energy Conversion	4375	Ceramics and Cermets	4433
Facilities and Equipment	4376	Metals and Alloys	4436
Heat Transfer and Fluid Flow	4378	Corrosion and Erosion	4436
Materials Testing	4379	Preparation and Fabrication	4437
Nuclear Explosions	4380	Properties, Structure, and Phase Studies	4438
Protective Structures and Equipment	4383	Radiation Effects	4444
ENVIRONMENTAL AND EARTH SCIENCES	4384	Other Materials	4448
Minerals and Ores	4384	NUCLEAR MATERIALS AND WASTE MANAGEMENT	4450
Radioactive Effluents	4384	Accountability and Safeguards	4450
Radioactivity Transport and Monitoring	4384	Disposal and Storage	4450
Atmosphere	4385	Handling Equipment and Procedures	4451
Soil	4386	Transportation	4452
Water	4387	PARTICLE ACCELERATORS	4453
Ecosystems and Food Cycles	4388	Design, Development, and Operation	4453
Radiometric Techniques	4391	Auxiliaries and Components	4453
Site Surveys	4392	Ion Sources, Injection, and Extraction	4454
Thermal Effluents	4393	Ion Optics and Field Calculations	4454
INSTRUMENTATION	4395	Storage Rings	4455
Radiation Detection Instruments	4395	Experimental Facilities	4455
General Detectors and Monitors	4395	PHYSICS (ASTROPHYSICS AND COSMOLOGY)	4456
Radiation Dosimeters	4397	Cosmic Radiation	4456
Nuclear Spectroscopic Instrumentation	4399	Stars	4457
High Energy Physics Instrumentation	4400	Quasi-Stellar, Radio, and X-Ray Sources	4461
Radiometric Instruments	4401	Solar Phenomena	4462
Radiation Effects on Instrument Components, Instruments, and Electronic Systems	4401	Galaxies	4464
Miscellaneous Instruments and Components	4403	Cosmology	4466
ISOTOPE AND RADIATION SOURCE TECHNOLOGY	4405	Planetary Phenomena	4467
Isotope Separation (Industrial)	4405	PHYSICS (ATMOSPHERIC)	4468
Isotopic Power Sources	4406	Auroral and Ionospheric Phenomena	4468
Radiation Sources	4407	Particle Kinetics and Trapping	4472
Design and Fabrication	4407	Solar Wind and Cosmic Ray Interactions	4475
Industrial Uses	4408		

TABLE OF CONTENTS (continued)

<i>Category</i>	<i>Page</i>	<i>Category</i>	<i>Page</i>
PHYSICS (ATOMIC AND MOLECULAR)	4478	PHYSICS (RADIATION AND SHIELDING)	4561
Atomic, Electron, Ion, and Molecular		Radiation Physics	4561
Beams and Their Reactions	4478	Radiation Protection Standards	4562
Atomic and Molecular Properties	4478	Radiation Source Calibration and Standardization	4562
Positronium, Muonium, and Mesic Atoms		Shielding Calculations and Theory	4562
and Molecules	4480		
Collision Phenomena	4481	PHYSICS (SOLID-STATE)	4564
Atomic Theory	4485	Radiation Effects	4564
PHYSICS (ELECTROFLUID AND MAGNETOFLUID)	4488	Physical Properties	4566
PHYSICS (HIGH-ENERGY)	4489	Moessbauer Effects and NMR	4569
Particle Interactions and Properties (Experimental)	4489		
Electromagnetic	4489	PHYSICS (THEORETICAL)	4571
Weak	4491		
Strong (Baryon-induced)	4491	REACTOR TECHNOLOGY	4573
Strong (Meson-induced)	4493	Power Reactor Development	4573
Particle Interactions and Properties (Theoretical)	4495	Physics and Core Engineering	4573
General	4495	Control and Instrumentation	4573
Electromagnetic	4496	Coolant System	4574
Weak	4501	Fuels and Fuel Elements	4574
Strong (General)	4502	Operation and Economics	4574
Strong (Baryon No. = 0)	4504	Safety and Siting	4574
Strong (Baryon No. = 1)	4504	Commercial Power Reactors	4575
Strong (Baryon No. ≥ 2)	4507	Physics and Core Engineering	4575
Particle Invariance Principles and Symmetries	4508	Control and Instrumentation	4575
General	4508	Containment Systems, Waste Processing,	
Applications (Electromagnetic and Weak)	4509	and Radiation Protection	4575
Applications (Strong)	4510	Coolant System	4575
Field Theories	4510	Auxiliary Systems and Engineered Safeguards	4575
General	4510	Operation and Economics	4575
Quantum Field Theory	4511	Safety and Siting	4577
Gravitation and General Relativity	4515	Steam and Power Conversion System	4585
Scattering Theory	4517	Propulsion and Auxiliary Power Reactors	4585
PHYSICS (LOW-TEMPERATURE)	4520	Process Heat and Desalination Reactors	4586
Superfluidity	4520	Research and Test Reactors	4586
Superconductivity	4521	Reactor Components, Fuels, and Accessories	4587
Equipment and Devices	4524	Reactor Physics and Core Engineering	4587
PHYSICS (NUCLEAR)	4525	Physics	4587
Neutron Interactions with Matter	4525	Core Thermal Hydraulics	4587
Nuclear Properties and Reactions, $A \leq 5$	4526		
Nuclear Properties and Reactions, $6 \leq A \leq 19$	4527	GENERAL	4588
Nuclear Properties and Reactions, $20 \leq A \leq 38$	4530	Civilian Defense	4588
Nuclear Properties and Reactions, $39 \leq A \leq 58$	4532	Law	4588
Nuclear Properties and Reactions, $59 \leq A \leq 89$	4533	Miscellaneous	4588
Nuclear Properties and Reactions, $90 \leq A \leq 149$	4534		
Nuclear Properties and Reactions, $150 \leq A \leq 189$	4537	CORPORATE AUTHOR INDEX	INDEX-1
Nuclear Properties and Reactions, $190 \leq A \leq 219$	4539		
Nuclear Properties and Reactions, $A \geq 220$	4540	PERSONAL AUTHOR INDEX	INDEX-13
Nuclear Theory	4542		
PHYSICS (PLASMA AND THERMONUCLEAR)	4549	SUBJECT INDEX	INDEX-62
Confinement and Heating	4549		
Diagnostics	4550	REPORT NUMBER INDEX	INDEX-161
Kinetics (General)	4552		
Kinetics (Experimental)	4553	JOURNAL LIST	APPENDIX-1
Kinetics (Theoretical)	4554		
Production	4555		
Stability	4555		
Thermonuclear Processes	4557		
Wave Phenomena	4558		

STAFF OF NUCLEAR SCIENCE ABSTRACTS

John W. Norris

Managing Editor

Donald D. Davis

Assistant Managing Editor

David E. Bost

Production Manager

Sidney F. Lanier

Subject Heading Specialist

J. Paul Meredith

Descriptive Cataloging Specialist

Frederick E. Marsh, Jr.

Computer Index Coordinator

Clara L. Fox

Corporate Author Specialist

Helen W. White

Report Number Specialist

Section Editors

Donald D. Davis

Life Sciences

Everett J. Hoffman

Chemistry

James M. Jacobs

Reactor Technology

Robert C. Kelly

*Engineering; Environmental and Earth Sciences;
Isotope and Radiation Source Technology; Materials;
Nuclear Materials and Waste Management*

Charles E. Stuber

*Instrumentation; Particle Accelerators; High Energy
Physics; Theoretical Physics*

Lee M. Thompson

*Astrophysics and Cosmology; Atmospheric Physics;
Atomic and Molecular Physics; Electrofluid and
Magnetofluid Physics; Low Temperature Physics;
Plasma and Thermonuclear Physics; Radiation and
Shielding Physics; Solid State Physics*

Lawrence T. Whitehead

Nuclear Physics

Evaluators

Thomas S. Chapman

John H. Mitchell

Mary H. Newman

Polly C. Summers

Abstractors

Ellis R. Bedford

Jane G. Buchanan

D. Lamar Cason

David C. Cunningham

Theodore F. Davis

James R. Dulaney

Helga B. Gerstner

Mary C. Grissom

Phyllis C. Hansen

Carolyn D. Houston

W. Hugh Kinser, Jr.

Lisbeth C. Lieberman

William D. Matheny

Dora H. Moneyhun

Jean F. Peckham

Henry D. Raleigh

Mona H. Raridon

Julia S. Redford

Jessie W. Rushing

Robert W. Rutkowski

Francis L. Sachs

Lila B. Smith

Anne G. Ward

Helen L. Ward

Kenneth S. Warren


Milton O. Whitson

Juanita Wilson

Charles R. Bruce

Publishing Coordinator

Communication concerning the editorial policy and content of *Nuclear Science Abstracts* should be addressed to the Managing Editor, *Nuclear Science Abstracts*, USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830.



NUCLEAR SCIENCE ABSTRACTS

CHEMISTRY

Analytical Chemistry

Refer also to abstracts: 45286, 45310, 45405, and 45681.

45181 (CISE-R-172) REACTOR CHEMISTRY: ANALYTICAL WORK. Topical Report No. 6. Cerrai, E.; Triulzi, C.; Trucco, R. (Centro Informazioni Studi Esperienze, Milan (Italy)). Apr 1967. 116p. Dep. NTIS (U. S. Sales Only).

Analytical methods applied to determinations of materials of interest in the field of the CIRENE program are described. Chemical procedures are given for determination of Al, B, Cl, Cr, Co, Cu, Pb, Lu, Mo, Nb, phosphate, sulfate, Th, Sn, Ti, U, and V. Emission spectroscopic methods are discussed for analysis of solutions, stainless steel, and zircaloy. Radiochemical procedures, including preparation of γ standard sources and methods for uranium determination in loop samples, are described. An appendix of miscellaneous applications is included. (M.C.G.)

45182 (JAERI-M-4594) CHEMICAL ANALYSIS PROCEDURES FOR JMTR. Tone, Hirohito; Yamamoto, Katumune; Okagawa, Seigo; Yokouchi, Ichiro; Atomura, Masakatu (Japan Atomic Energy Research Inst., Tokyo). Sep 1971. 211p. (In Japanese). NTIS (U. S. Sales Only).

Chemical analyses at the JMTR are important for the operation of reactor and high-pressure water loops and for the reactor and loop chemistry. The analytical procedures and equipment employed are described. Many of these procedures have been standardized to speed up analytical services in the laboratory. The objects for chemical analyses are active and nonactive nuclides and chemical compounds in the primary coolants, which are concerned with corrosion, fuel failure, plant safety, water treatment in loops and reactor, and water chemistry. (auth) (Japan)

45183 (TID-7029(2nd Ed.)) SELECTED MEASUREMENT METHODS FOR PLUTONIUM AND URANIUM IN THE NUCLEAR FUEL CYCLE. Second Edition. Rodden, Clement J. (ed.) (Office of Information Services (AEC), Washington, D. C.). 1972. 440p. Dep. NTIS.

Separate abstracts were prepared for Part I, General Considerations, and for each of the 4 sections of Part II, Analytical Methods. The sections are: uranium-bearing materials, plutonium-bearing materials, hot process solutions, and apparatus and reagents. (M.C.G.)

For abstracts of individual sections see: 45184-45186, 45190, and 45191.

45184 (TID-7029(2nd Ed.), pp 3-65) GENERAL CONSIDERATIONS. 1972.

In Selected measurement methods for plutonium and uranium in the nuclear fuel cycle.

Standard reference materials, sampling, and bulk measurements are discussed for determination of Pu and U in nuclear materials. (M.C.G.)

45185 (TID-7029(2nd Ed.), pp 69-252) URANIUM-BEARING MATERIAL. 1,000 Series. 1972.

In Selected measurement methods for plutonium and uranium in the nuclear fuel cycle.

Methods are considered for determination of the uranium concentration and its isotopic composition in product material. Thirty-five methods are described. (M.C.G.)

45186 (TID-7029(2nd Ed.), pp 253-329) PLUTONIUM-BEARING MATERIAL. 2,000 Series. 1972.

In Selected measurement methods for plutonium and uranium in the nuclear fuel cycle.

Methods are considered for determination of plutonium and its isotopic composition in fuel cycle product material. Sixteen methods are described. (M.C.G.)

45187 MODERN ANALYTICAL TECHNIQUES FOR METALS AND ALLOYS. PART 2. Bunshah, R. F. (ed.). Techniques of Metals Research. Volume III. New York; John Wiley and Sons, Inc. (1970). 520p.

Separate abstracts were prepared for each of 10 chapters. (M.C.G.)

For abstracts of individual chapters see: 45208-45212, 45246-45249, and 45271.

45188 MASKING AND DEMASKING OF CHEMICAL REACTIONS. THEORETICAL ASPECTS AND PRACTICAL APPLICATIONS. Perrin, D. D. Chemical Analysis, A Series of Monographs on Analytical Chemistry and Its Applications. Volume 33. New York; John Wiley and Sons, Inc. (1970). 222p.

The use of masking and demasking of chemical reactions in analytical chemistry, industry, biology, and medicine is discussed. Applications to analytical chemistry are described for titrimetry, precipitation reactions, spectrophotometry, ion exchange, and masking of chemical reactivity. Masking agents for cations are listed. Anions and neutral molecules are classified by types and by species that are masked. The industrial applications are concerned with the control of undesirable effects due to traces of metal ions, including hardness of water, production of blotches or staining and the development of off-flavors and rancidity. Biological and medical applications comprise such examples as metal-ion buffers and treatments for metal poisoning. (D.H.M.)

Activation and Nuclear Reaction Procedures

Refer also to abstracts: 45309, 45567, 45788, and 46146.

45189 (BFL-1/1972) DEVELOPMENT OF METHODS FOR THE APPLICATION OF NEUTRON ACTIVATION ANALYSIS TO THE DETERMINATION OF TRACE ELEMENTS IN FOODS. Progress Report, October 1-December 31, 1971. Toeroek, G.; Schelenz, R.; Fischer, E. (Bundesforschungsanstalt fuer Lebensmittelfrischhaltung, Karlsruhe (West Germany)). Feb 1972. 23p. (In German). Dep. NTIS (U. S. Sales Only).

The significance of trace element determination in foodstuffs is pointed out and the advantages of using neutron activation analysis for such determinations are described. Arguments are presented for a comprehensive analytical scheme. It consists of a combination of radiochemical separations and gamma spectrometry. First results of the testing of individual steps in this scheme are presented. (auth)

45190 (TID-7029(2nd Ed.), pp 331-411) HOT PROCESS SOLUTIONS. 3,000 Series. 1972.

In Selected measurement methods for plutonium and uranium in the nuclear fuel cycle.

The analysis of process solutions of spent fuels for uranium and plutonium is considered and 12 methods are described. Determinations of U and Pu in waste materials are also discussed. (M.C.G.)

45191 (TID-7029(2nd Ed.), pp 413-33) APPARATUS AND REAGENTS. 4,000 Series. 1972.

In Selected measurement methods for plutonium and uranium in the nuclear fuel cycle.

Thirteen pieces of equipment and 33 reagents for use in determination of plutonium and uranium are considered. (M.C.G.)

45192 (TID-25954, pp 373-5) STABLE CESIUM ANALYSIS BY NEUTRON ACTIVATION OF PLANTS FROM THE NEVADA TEST SITE. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Cesium contents of leaves for several species of plants were determined by neutron activation. Concentrations varied from 30 to 372 nanograms per gram dry weight. (M.C.G.)

45193 (TID-25954, pp 376-8) CYCLING OF STABLE CESIUM IN A DESERT ECOSYSTEM. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Cesium was determined in plant, animal, and soil samples by neutron activation. Potassium was determined by counting ^{40}K by γ spectrometry. Samples were obtained from the Rock Valley area of the Nevada Test Site in 1969 and 1970. (M.C.G.)

45194 RADIOANALYTICAL INVESTIGATION ON THE EXTRACTION BEHAVIOR OF RHODIUM IN THE THIOCYANATE-TBP SYSTEM. Di Casa, M.; Stella, R. (Univ., Pavia, Italy). Radiochem. Radioanal. Lett.; 10: No. 6, 331-7(15 Jul 1972).

A detailed study of various conditions affecting the extraction of small amounts of rhodium into tributyl phosphate from thiocyanate solutions is reported. Neutron activation analysis was used to measure the extraction efficiencies, thus avoiding the use of spectrophotometric or similar relatively cumbersome methods usually applied. (auth)

45195 DETERMINATION OF ASH CONTENT OF COAL BY NEUTRON ACTIVATION ANALYSIS. Loska, L.; Gorski, L. (Inst. of Nuclear Techniques, Krakow). Radiochem. Radioanal. Lett.; 10: No. 6, 315-22(15 Jul 1972).

A method is presented for determining the ash content of coal by 14-MeV neutron activation analysis using a neutron generator. The activation of silicon and aluminum was the basis of the analysis. The total ash content of coal was determined based on the correlation between the chemical composition of the ash (especially Si and Al content) and the ash content of the coal. The presented method allows the determination of the ash content of coal with a mean absolute error of about 1.8%. (auth)

45196 DETERMINATION OF CHROMIUM IN MILK POWDER. Das, H. A.; Erlich, Y.; Hoede, D.; Zonderhuis, J. (Reactor Centrum Nederland, Petten). Radiochem. Radioanal. Lett.; 10: No. 6, 361-6(15 Jul 1972).

Milk powder sample is irradiated by thermal neutrons for 24 hours and allowed to cool for 10 days. After dissolution K_2CrO_4 is added as carrier and the chromium is extracted as peroxychromic acid into ethyl acetate, then back extracted into the aqueous phase. Chromate is precipitated as BaCrO_4 and its activity measured. The time required for an analysis is about 30 minutes. (auth)

45197 NEW METHOD FOR POTASSIUM DETERMINATION BY NEUTRON ACTIVATION IN MINERAL AND ROCK SAMPLES. Soriou, M.; Popescu, G. (Institutul de Fizica Atomica, Bucharest). Ann. Soc. Geol. Belg.; 94: No. 2, 132(1971). (CONF-710953-4).

From European conference of geochronology; Brussels, Belgium (6 Sep 1971).

For potassium determination by neutron activation, the reaction $^{39}\text{K}(n, p)^{39}\text{Ar}$ is used. Depending on their potassium content, samples of 100 or 200 milligrams are irradiated 40 hours at a flux of 5.10^{12} fast neutrons and 10^{13} thermal neutrons/cm² sec. By comparing the potassium contents determined by classical and by neutron activation methods on 12 different mineral samples, the conclusion can be drawn that the precision and the accuracy of the new method are better than $\pm 3\%$. (auth)

45198 ACTIVATION ANALYSIS OF ENVIRONMENTAL POLLUTANTS. Chattopadhyay, A.; Bennett, L. G. I.; Jervis, R. E. (Univ. of Toronto). Can. J. Chem. Eng.; 50: No. 2, 189-93(Apr 1972).

Activation analysis is a highly sensitive and specific technique for the detection of trace heavy metal pollutants in environmental materials. Comparison of natural "baseline" levels with elevated levels provides direct evidence of contamination. Neutron activation analysis (NAA) was used to determine arsenic, zinc, cadmium, and mercury. Photo-nuclear activation analysis (PNAA) was used to determine lead in soils. Holland Marsh, a market-gardening area north of Toronto, was studied as various herbicides and fungicides were used there. The analyses of soil and vegetation

samples, and of human head hair collected from residents are reported. (auth)

45199 ACTIVATION ANALYSIS WITH A Pu-Be NEUTRON SOURCE. Pawaskar, P. B.; Reddy, G. R.; Sankar Das, M. (Bhabha Atomic Research Centre, Trombay, India). Indian J. Technol.; 10: No. 2, 80-2(1972).

Simple procedures are described for the routine activation analysis of manganese in steels and fluorine in fluorites and plant products using a 5 curie Pu-Be neutron source. Manganese analysis is based on $^{55}\text{Mn}(n, \gamma)^{56}\text{Mn}$ reaction using thermal neutrons from the source and counting the 0.84-MeV photopeak of ^{56}Mn . Fluorine is estimated by $^{19}\text{F}(n, \alpha)^{16}\text{N}$ reaction using fast neutrons from the source and counting the 6-7-MeV gamma rays of ^{16}N . The results obtained by the methods described are compared with those obtained by chemical methods. (Indian Sci. Abstr.)

45200 GOLD ASSAY ON GEOLOGICAL SAMPLES BY NEUTRON ACTIVATION WITH PRELIMINARY CRUCIBLE FUSION. Leipunskaya, D. I.; Savosin, S. I.; Drynkin, V. I.; (and others). Ind. Lab. (USSR) (Engl. Transl.); 37: No. 12, 1894-6(Dec 1971).

Translated from Zavod. Lab.; 37: No. 12, 1471-3(Dec 1971).

In crucible melting, gold enters the lead alloy from the ore along with other elements; however, lead and these elements produce very little interference in determination of the gold by neutron activation. The lead alloy is irradiated along with standards and the γ spectra recorded. The limit of detection was found to be 0.01 g/ton. It was found that silver could also be determined by this method. (M.C.G.)

45201 ACTIVATION METHODS OF ANALYSIS. Zmijewska, W. (Inst. of Nuclear Research, Warsaw). Wiad. Chem.; 25: 99-122(1971). (In Polish).

The contemporary techniques of activation analysis are described, particularly thermal and fast neutron activation, photo-activation, charged-particle activation, and techniques based on the measurement of prompt gamma radiation. Examples of application are given. (auth) (NSA of Poland)

45202 SIMPLE AND RAPID PROCEDURE FOR STRONTIUM DETERMINATION IN RAT PLASMA USING NEUTRON ACTIVATION. Maillard, M. J. (CEN de Fontenay-aux-Roses, France); Maziere, B.; Comar, D. Bull. Soc. Chim. Fr.; No. 4, 1679-81 (Apr 1972). (In French).

The determination of strontium in small samples of plasma was effected by destructive activation analysis. The separation of strontium was carried out by double precipitation of oxalates (pH8) followed by filtration through an antimony pentoxide column in order to completely eliminate sodium. A third oxalate precipitation gave a manganese-free sample. A series of measurements, carried out on nine 0.5 ml samples of the same rat plasma, yielded the following data: 78.6 ± 3.8 ng/ml. This method is the most accurate for the low plasma volumes considered. (France)

45203 VANADIUM DETERMINATION IN BIOLOGICAL MATERIALS AT NANOGRAM LEVELS BY A CATALYTIC METHOD. Welch, Ross M.; Allaway, William H. (Plant, Soil, and Nutrition Lab., Ithaca, N. Y. Cornell Univ., Ithaca, N. Y.). Anal. Chem.; 44: No. 9, 1644-7(Aug 1972).

A sensitive and accurate spectrophotometric method for measuring amounts of V as low as 5 ng in biological materials is described. Samples are digested with HNO_3 - HClO_4 and V is separated from interfering elements by complexing with 8-quinolinol in chloroform at pH 4.0. It is then released from the complex by readjusting the pH to 9.4 with an aqueous NH_4NO_3 buffer. Vanadium is then determined in the aqueous extract by its catalytic effect on the rate of oxidation of gallic acid by acid persulfate. The extraction procedure eliminates all interfering elements except Mo. Mo interfered with V determination when it exceeded 10 μg per sample. Comparisons with a neutron activation method were made. Recoveries of V added to plant samples were between 94 and 98%. Recoveries of radioactive ^{48}V added to standard V solutions and carried through the extraction procedure were between 90 and 100%. (auth)

45204 DETERMINATION OF INORGANIC IMPURITIES IN VINYL CHLORIDE BY ACTIVATION ANALYSIS. Bunus, Florin T.; Murgalescu, Sanda (Inst. of Atomic Physics, Bucharest). Talanta; 19: No. 3, 372-7(Mar 1972).

Trace impurities in vinyl chloride were determined by activation analysis in a reactor thermal neutron flux of 1.2×10^8 n/mm²/sec. Gamma-spectrometry was used to reveal the presence of arsenic and bromine. The impurities determined were: arsenic $8.0 \pm 0.6 \times 10^{-4}\%$, bromine $3.07 \pm 0.2 \times 10^{-3}\%$ and phosphorus $2.0 \pm 0.15 \times 10^{-2}\%$. (auth)

45205 PROCEEDINGS OF THE SECOND INTERNATIONAL CLEAN AIR CONGRESS HELD AT WASHINGTON, D. C., DE-

CEMBER 6-11, 1970. Englund, H. M.; Berry, W. T. (eds.). New York; Academic Press, Inc. (1971). 1378p. \$85.00. (CONF-701229).

Separate abstracts were prepared for six of the two hundred fifty papers presented. The remaining papers were not in the subject scope of NSA. (D.H.M.)

For abstracts of individual papers see: 45206, 45207, 45244, 45245, 45514, and 45568.

45206 TECHNIQUES FOR SOURCE DETERMINATION IN THE TUCSON BASIN. Post, Roy G.; Wacks, Morton E. (Univ. of Arizona, Tucson). pp 410-18 of Proceedings of the Second International Clean Air Congress. /Englund, H. M. (ed.). New York; Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

Two approaches to computer processing of data obtained from elemental analysis of airborne particulate matter by neutron activation were successful. The first program, regression analysis, used less computer time and memory, but the second, least squares, gave a more exact synthesis of the sample spectrum from the library spectra. Compared to other types of analysis, such as spectroscopic, gravimetric, or volumetric analysis, activation analysis is potentially at least an order of magnitude more sensitive for elements with high cross sections. Other advantages of activation analysis are its speed, the fact that it does not damage the sample, and that it is less subject to human error. Its major disadvantage is that activation analysis cannot provide molecular information. However, elemental analysis combined with mineral content information and total organic content should provide sufficient information to relate the particulates to their sources. (auth)

45207 MULTI-ELEMENT ANALYSIS OF AIR POLLUTION PARTICULATES BY NONDESTRUCTIVE NEUTRON ACTIVATION. Dams, Richard (Inst. of Nuclear Research, Ghent); Rahn, Kenneth A.; Robbins, J. A.; Nifong, Gordon D.; Winchester, John W. pp 509-16 of Proceedings of the Second International Clean Air Congress. /Englund, H. M. (ed.). New York; Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

Tests of possible applications of neutron activation analysis to the study of trace elements in air pollution particulates indicate that this analytical technique can lead to discovery of unsuspected pollution anomalies such as Cu, meteorologically related time variations that are not the same for all elements, and size distributions that are related to chemical processes of pollution at pollution sources and in the atmosphere. These findings may have immediate usefulness both in pollution monitoring and control and in research on the migration and ultimate removal of pollutants from the atmosphere. (D.H.M.)

45208 ACTIVATION ANALYSIS WITH THERMAL NEUTRONS. Albert, Philippe (CNRS, Vitry-sur-Seine, France). pp 687-754 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Principles of activation analysis, advantages of thermal neutron activation, chemical separations, comparison of radioactivity and check of radiochemical purity, sensitivity, suppression of the risk of contamination, the very high selectivity of the method, examples of systematic analyses, nondestructive methods, and causes of errors of nuclear origin are discussed. (130 references). (M.C.G.)

45209 ACTIVATION ANALYSIS WITH FAST NEUTRONS. Laverlohere, J. (CEN, Grenoble, France). pp 755-75 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

The principle and techniques used, the equipment, the advantages and limitations, and the applications of fast neutron activation analysis are considered. (104 references) (M.C.G.)

45210 ACTIVATION ANALYSIS WITH GAMMA RAYS AND CHARGED PARTICLES. Engelmann, Ch. (CEN de Saclay, Gif-sur-Yvette, France). pp 777-865 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Topics covered include: advantages and inconveniences of γ and charged particle activation analysis, principle of determination, choice of nuclear reaction and experimental conditions, standardization of the determination, examples of use, and future perspectives. (126 references) (M.C.G.)

45211 USE OF COMPUTERS IN ACTIVATION ANALYSIS. Laverlohere, J. (CEN, Grenoble, France). pp 867-80 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

The use of digital computers in the activation analysis method to facilitate optimization and exploitation of this technique and to improve its performance is discussed. Apparatus and technique, applications, and limits and advantages are considered. (50 references) (M.C.G.)

45212 GENERAL CONCLUSIONS ON THE METHODS OF ACTIVATION ANALYSIS. Albert, Ph. (CECM-CNRS, Vitry-sur-Seine, France). pp 881-4 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Choice of an activation method among those that are candidates for solution of an analytical problem is illustrated by the example of the determination of oxygen. (M.C.G.)

45213 DEVELOPMENT AND APPLICATION OF A SYSTEM-ATIC APPROACH TO ELEMENTAL ANALYSIS IN FORENSIC INVESTIGATIONS. Whitworth, James Bradley. Austin, Tex.; Univ. of Texas (1971). 235p. University Microfilms Order No. 72-19,693.

Thesis.

A program of cooperation between the Texas Department of Public Safety (DPS) and the Nuclear Reactor Laboratory (NRL) of the University of Texas at Austin was initiated. The purpose of this program is to apply analytical techniques used in the NRL to actual forensic cases in which the DPS provides samples and guidance. The techniques used in these analyses are neutron activation analysis (NAA) and nondispersive x-ray fluorescence (NDFX). Analysis of data is accomplished with two computer programs, one for scintillation detector spectra and one for solid-state detector spectra. Evidence from several actual cases were analyzed. Reactor neutron activation coupled to a Compton-suppressed Ge(Li) detector and versatile computer program for semiautomatic analysis, is used to investigate some actual criminal cases. In addition to establishing lower limits of detectability with typical forensic evidence, trace analyses of some types of evidence heretofore unreported were obtained. The value of using a Compton suppression device for certain types of samples is shown. Limits of detectability for several elements were determined with a 1000 μg ^{252}Cf neutron source using both Ge(Li) and NaI(Tl) detectors. It was hoped that cyclic activation with ^{252}Cf would add to the information obtained with reactor activation. Preliminary results indicate that the Ge(Li) detector is too inefficient for this work except in very special cases. Nondispersive x-ray fluorescence was used to supplement the neutron activation data of most of the forensic cases attempted. Samples were excited by ^{238}Pu , ^{241}Am , and ^{109}Cd radioisotopic sources and the spectra were acquired on a cooled Si(Li) detector. In several cases the NDFX data substantiated or supplemented results obtained by NAA. The results proved that NDFX is a fast, inexpensive method to analyze forensic samples for ratios of minor elements. In addition to the routine cases, survey projects are in progress on paint chips and metal alloys. These two classes of samples proved to be troublesome for other analytical techniques and this program is establishing the capabilities of NDFX in these areas in anticipation of future cases. (Diss. Abstr. Int., B)

45214 NEUTRON ACTIVATION ANALYSIS OF TRACE ELEMENTS IN HAIR. Huckabee, John W. Laramie, Wyo.; Univ. of Wyoming (1971). 75p. University Microfilms Order No. 72-13,028.

Thesis.

To determine the trace element content of the hair of natural free-ranging mammals, hair samples were collected from several populations in Idaho and Wyoming. To determine the variation in hair trace element content of several sympatric species, small mammals were trapped in an area in Jackson Hole, Wyoming. To determine the uptake by an animal and subsequent deposition in hair of an abundant environmental element, small mammals were trapped for hair analyses in highly mineralized zones in Idaho and Wyoming. Hair analyses were by neutron activation with gamma spectrometry at the NRTS in Idaho. The gamma spectra were analyzed by computerized techniques using an IBM 360/75 and a PDP-8 interfaced to one of the PHAS. Irradiations were performed in the MTR and the ETR during 1969, 1970, and 1971. After irradiation the hair samples were brought into the laboratory and prepared for gamma counting on the spectrometers. All hair tested had Zn, Na, K, and Br. Eighty-two percent had Se, 50% had Fe, 52% had Ag, 31% had Sb, 22% had Hg, 15% had Sc, 9% had Ce, 1% had Au, 0.9% had Rb. No statistically significant differences were found quantitatively or qualitatively at this sample size. (Diss. Abstr. Int., B) (TSS)

45215 TITANIUM AND CALCIUM CONTENT OF METEORITES. van Zelst, L.; Aten, A. H. W. Jr. (Instituut voor Kernfysisch Onderzoek, Amsterdam). Radiochim. Acta; 17: No. 2, 115-16 (Jul 1972).

The determination of titanium and calcium in meteorites by non-destructive proton activation is reported involving the nuclear reactions $^{46}\text{Ti}(p,n)^{46}\text{V}$ and $^{44}\text{Ca}(p,n)^{44}\text{Sc}$. For comparison purposes the calcium content was determined by bremsstrahlung activation involving the reaction $^{44}\text{Ca}(\gamma,p)^{43}\text{K}$. (D.H.M.)

45216 DETERMINATION OF Mn IN HIGH-PURITY IRON BY NEUTRON ACTIVATION ANALYSIS AFTER PRE-IRRADIATION SEPARATION. Steinnes, E. (Institut for Atomenergi, Kjeller, Norway). *Radiochem. Acta*; 17: No. 2, 119-20 (Jul 1972).

A method for determining manganese in high-purity iron by neutron activation analysis is reported that includes a preirradiation separation technique based on the removal of iron by anion exchange separation in strong hydrochloric acid for the elimination of the interference from the reaction $^{56}\text{Fe}(n,p)^{56}\text{Mn}$. (D.H.M.)

45217 TRACE ANALYSIS FOR PLATINUM IN GLASSES BY NEUTRON ACTIVATION. Becker, Donald A. (National Bureau of Standards, Washington, D. C.). *Anal. Chim. Acta*; 61: No. 1, 1-6 (Aug 1972).

Trace amounts of platinum were determined in eight glasses. These glasses were prepared under special conditions, and information on both the surface and interior platinum was desired. Previous microscopic examination of the glass samples had identified platinum on the surface of at least four of the samples. The platinum was analyzed by means of the nuclear reaction $^{198}\text{Pt}(n,\gamma)^{199}\text{Pt}$ - ^{198}Au . The ^{198}Au product has a 3.15-day $T_{1/2}$ and E_{γ} of 158 and 208 keV. Samples were irradiated in the NBSR for 2 h at $1.3 \times 10^{13} \text{ n cm}^{-2} \text{ sec}^{-1}$. After irradiation, the glasses were etched with aqua regia for the determination of surface platinum, and then dissolved in hydrofluoric and perchloric acids for the determination of interior platinum. The radioactive ^{198}Au was separated from the matrix by spontaneous deposition on metallic zinc. Experimental sensitivities of 0.04 μg of platinum were obtained. (auth)

45218 INSTRUMENTAL NEUTRON ACTIVATION ANALYSIS OF GERMANIUM. Adams, F.; Maenhaut, W.; Hoste, J. (Ghent Univ.). *Anal. Chim. Acta*; 61: No. 1, 127-31 (Aug 1972).

The determination of trace impurities (Co, In, Ni, Sb, Zn, Cr, Ir, Se, Sc, Ag, Ta, Hf, and U) in germanium by instrumental neutron activation analysis with a high resolution Ge(Li) detector is described. Measurements less than 3 weeks after the irradiation were precluded by the high matrix activity. (D.H.M.)

Chemical and Spectral Procedures

45219 (ARH-SA-110) SPECTROPHOTOMETRIC DETERMINATION OF PHOSPHATE IN NUCLEAR MATERIALS. Winters, William I. (Atlantic Richfield Hanford Co., Richland, Wash.). Jan 1972. Contract AT(45-1)-2130. 18p. (CONF-720628-3). Dep. NTIS.

From twenty-seventh ACS northwest regional meeting; Corvallis, OR. (15 Jun 1972).

By extracting molybdophosphoric acid into 1-butyl acetate in the presence of ethylenediaminetetraacetic acid and fluoride, a very simple but selective and sensitive spectrophotometric procedure for phosphate is obtained by measuring the absorbance of the extracted heteropoly acid at 310 nanometers. The optimum acid and reagent concentrations for the method were determined and the effect and tolerance levels of 36 interfering ions, including uranium, thorium, and plutonium, were measured. The method is capable of measuring 0.7 μg of phosphorus and has a molar absorptivity of approximately 25,000. The standard deviation for the method is $\pm 5.4\%$. This procedure was used to determine the concentration of phosphate in ^{233}U product solutions, to follow the radiolysis of entrained tributyl phosphate and dibutyl phosphate in ^{233}U product solutions, and to determine the phosphate content of nuclear waste solutions. (auth)

45220 (BARC-593) SPECTROCHEMICAL ANALYSIS OF HIGH PURITY CADMIUM. Murty, P. Sreerama; Kaimal, V. N. P. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 14p. Dep. NTIS (U. S. Sales Only).

A spectrographic method using dc arc excitation is described to estimate twelve trace impurities such as As, Sb, etc., in high-purity cadmium oxide. Cadmium metal is converted to oxide and mixed with pure graphite in the ratio 1:1 by weight. Forty milligrams of the mixture are taken in a graphite electrode and excited at a current of 12 amp dc. The spectra are recorded on an Ebert 3.4-meter plane grating spectrograph. The detection limits obtained for various elements range from 1 to 10 ppm. The precision of the method is ± 8 to 16%. (auth)

45221 (EUR-4822) TRACE LEVEL DETERMINATION OF OXYGEN IN ORGANIC COOLANTS AND ULTRA-MICRO DETERMINATION OF OXYGEN IN C-H-O ORGANIC COMPOUNDS.

Colombo, A. (European Atomic Energy Community, Ispra (Italy). Joint Nuclear Research Center). 1972. 20p. Dep. NTIS (U. S. Sales Only). EUR B.Fr. 40.-.

A method of high sensitivity that requires the use of small samples was developed for determination of the oxygen contained at trace levels in the organic coolants used in nuclear reactors and for the ultra-microanalysis of oxygen-rich organic C-H-O compounds. The weights of the samples used are lower than 30 and 0.5 mg, respectively, and no modification to the apparatus is required for changing from one type of determination to another. The method is based on a catalytic pyrolysis and reduction in hydrogen stream and on the final measurement of the formed water by means of a commercial Keidel electrolytic hygrometer. Although it has not been applied to compounds containing atoms other than carbon, hydrogen, and oxygen, it seems at least theoretically that small and simple variations of the apparatus or of the analytical procedure could render the method able to analyze substances containing other atoms. (auth)

45222 (JEN-229-DQ/I-79) VOLATILIZATION MECHANISM OF CERTAIN ELEMENTS IN A METHOD OF SEMIQUANTITATIVE SPECTROGRAPHIC ANALYSIS. Diaz-Guerra, J. P. (Junta de Energia Nuclear, Madrid (Spain). Division de Quimica). 1972. 18p. (In Spanish). Dep. NTIS (U. S. Sales Only).

The efficiency of the compounds GeO_2 , $\text{Li}_2\text{B}_4\text{O}_7$, Li_2CO_3 , SiO_2 , and SrCO_3 as spectrochemical buffers, in the development of a semiquantitative spectrographic method of analysis that can be applied to the determination of 47 elements in different matrices, was tested. By the use of $\text{Li}_2\text{B}_4\text{O}_7$ or GeO_2 adequate accuracy is obtained, and with the latter the detection of much lower concentrations, especially for the elements Mo, Ti, V, and W, was attained. In order to account for the different behavior of these elements, their volatilization mechanism was studied. MoB_2 , TiB_2 , VB_2 , $\delta\text{-WB}$, and $\epsilon\text{-WB}$ were found by x-ray-diffraction analysis to be reaction products in the case of mixtures with $\text{Li}_2\text{B}_4\text{O}_7$. (auth)

45223 (JEN-232-DQ/I-81) APPLICATION OF SPECTROGRAPHIC ANALYSIS TO THE RADIOISOTOPE PRODUCTION CONTROL. II. ANALYSIS OF ^{45}Ca , ^{46}Sc , ^{63}Ni , AND ^{64}Cu SOLUTIONS. Capdevila, C.; Roca, M. (Junta de Energia Nuclear, Madrid (Spain). Division de Quimica). 1972. 19p. (In Spanish). Dep. NTIS (U. S. Sales Only).

Semiquantitative and quantitative determinations of both the radioactive and the target element in each radioisotope are described. The copper-spark technique was used except for Cu determinations, which need silver or graphite electrodes. Inter-element effects and their compensation through the use of Bi, Ga, In, Mo, Pd, Tl, and Y as reference elements was examined. For determination of Ca in ^{45}Ca samples, Ba, La, Li and Sr were also tested. Good results were achieved with Li for Ca, Y for Sc, Ti, and Ni, and either In or Y for Cu and Zn. (auth)

45224 (LA-DC-72-524) REMOTIZED APPARATUS FOR DETERMINING OXYGEN IN IRRADIATED REACTOR FUELS AND CLADDING MATERIALS. MacDougall, C. S.; Smith, M. E.; Waterbury, G. R. (Los Alamos Scientific Lab., N. Mex.). 1972. 18p. (CONF-720903-8). Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

Two well-tested, reliable methods for determining oxygen were modified for hot cell operations and applied to analyses of irradiated materials. In either method the sample is reacted at high temperature with carbon to convert the oxygen to carbon monoxide and dioxide. One reaction furnace is used with either of two analytical finishes to measure traces of oxygen in carbides and claddings or larger amounts in oxide fuels. Unique features in the modified method include: equipment and operations for grinding and pelletizing oxides, techniques for removing sodium and fuel from cladding prior to analysis, a crucible assembly to maximize heating efficiency, and a reaction furnace tube which facilitates easy access to the crucible assembly and trouble-free maintenance. The method and equipment were evaluated by analyzing uranium oxide, mixed oxide fuels having undergone up to 8.5 at. % burnup, metals, carbides, and alloys. There is no known bias. (auth)

45225 COMPARISON OF PRECISION IN SINGLE AND DOUBLE ISOTOPE DILUTION ANALYSIS. Boelrijk, N. A. I. M. (ZWO Laboratorium voor Isotopen-Geologie, Amsterdam). *Ann. Soc. Geol. Belg.*; 94: No. 2, 105-6 (1971). (CONF-710953-2).

From European conference of geochronology; Brussels, Belgium (6 Sep 1971).

Gain in accuracy, due to correction for mass discrimination by double isotope dilution analysis, is obtained at the expense of precision. Loss of precision may be considerable when an unfavorable mixing ratio between spike and unknown was used. A method is

proposed to decide whether the result evaluated by single or by double isotope dilution calculation should be preferred. The decision takes into account the quality of the mass spectrometric analysis and is not based on assumptions concerning the precision of the isotopic ratio determination. (auth)

45226 SEPARATION AND SPECTROPHOTOMETRIC DETERMINATION OF CERIUM AND MANGANESE BY TTA EXTRACTION. Onishi, Hiroshi; Toita, Yukio (Japan Atomic Energy Research Inst., Tokai). Jap. Anal.; 21: No. 6, 756-9 (Jun 1972). (In Japanese).

Spectrophotometric methods were previously described for the determination of cerium and manganese with TTA. In these methods, cerium and manganese interfered with each other, the absorbances were measured before and after back-extraction cerium. A method was developed for the separation of cerium and manganese by TTA extraction prior to their spectrophotometric determination with TTA. Cerium(III) is extracted with 0.5 M TTA in xylene at pH 4.3 ± 0.2 (acetate solution). Manganese is then extracted with TTA-acetone-xylene at pH 6 to 7. The cerium and manganese are back-extracted with 1 M nitric acid. The back-extracts are evaporated to dryness, and the residues are treated with nitric and perchloric acids. After oxidation with sodium bromate, cerium(IV) and manganese(III) are extracted with 0.5 M TTA in xylene from 0.5 M sulfuric acid solutions. The absorbance of each extract is measured at 440 nm. The recoveries of cerium(III) and manganese after single extraction with 0.5 M TTA in xylene as a function of pH are shown. About 95% of cerium is extracted at pH 4.1 to 4.5. Manganese begins to be extracted at pH 4.6, and about 80% of manganese is extracted at pH about 6. By this procedure 50 μg of cerium can be determined in the presence of up to 1 mg of manganese. Ten μg of manganese can be determined in the presence of up to 100 μg of cerium. Mixtures containing 100 μg each of cerium and manganese were analyzed six times, and the relative standard deviations were 1.0% for cerium and 3.7% for manganese. Effects of foreign ions on the analysis for cerium and manganese are summarized. (auth)

45227 DETERMINATION OF SMALL AMOUNTS OF MOLYBDENUM BASED ON ITS CATALYTIC EFFECT IN THE METHYLENE BLUE-HYDRAZINE SULFATE REACTION. Yamane, Takeshi; Kitamura, Teruo; Fukasawa, Tsutomu; Suzuki, Takashi (Yamanashi Univ., Kofu, Japan). Jap. Anal.; 21: No. 6, 799-803 (Jun 1972). (In Japanese).

A rapid and sensitive method was developed for the determination of small amounts of molybdenum. The method is based on the catalytic activity of molybdenum in the redox reaction between Methylene Blue and hydrazine sulfate. The rates of catalyzed reduction of Methylene Blue by hydrazine sulfate are followed spectrophotometrically by measuring the rate of decolorization of Methylene Blue at 660 nm. A linear relationship $k = k[\text{Mo}]$ was obtained between initial concentration of molybdenum up to 20 μg and the pseudo first order rate constant k evaluated from the plot of $\log(\text{Methylene Blue concentration})$ vs. reaction time. Using this relationship, molybdenum for the range 1 to 20 μg in unknown samples can be determined by measurements of reaction rate. Variables in the determination were studied. Of the many ions examined, W(VI) had an accelerating effect on the reaction and hence caused a larger value, while Fe(III) had a decelerating effect even at a few μg level and caused a smaller value. At the stated level, no interference was observed from the other cations. None of the anions and organic acids examined at 5 to 10 mg level interfered. The time required for an analysis is about 20 minutes. The standard deviation at a molybdenum level of 10 μg is 2.2%. (auth)

45228 COMPARISON OF THEORETICAL AND EXPERIMENTAL LIMITS OF DETECTION IN ATOMIC ABSORPTION SPECTROMETRY USING AIR-ACETYLENE AND NITROUS OXIDE-ACETYLENE FLAMES. Parson, M. L.; McElfresh, P. M. (Arizona State Univ., Tempe). Appl. Spectrosc.; 26: No. 4, 472-80 (1972).

Limits of detection were calculated for the air-acetylene and nitrous oxide-acetylene flames used in atomic absorption spectrometry using the theoretical treatment developed by Winefordner. Values for the parameters were chosen for these calculations to give the lowest possible limits of detection. These calculated values were compared with experimentally determined limits of detection for 57 elements. Values of the atom fraction, β , are also estimated by means of this comparison. The values for the air-acetylene flame are in fair agreement; however, the comparison of the nitrous oxide flame showed anomalous behavior. An attempt to explain this is made, as well as a general discussion of the results. (auth)

45229 FIRE ASSAY-EMISSION SPECTROGRAPHIC DETERMINATION OF IRIDIUM, RUTHENIUM, AND OSMIUM. Broadhead, K. G.; Piper, B. C.; Heady, H. H. (Bureau of Mines, Reno, Nev.). Appl. Spectrosc.; 26: No. 4, 461-7 (1972).

A method is presented for determination of microgram levels of Ir, Ru, and Os in ores, concentrates, and heavy sands. Radio-tracers are used to monitor the recovery of these three metals in a 10-mg Pt bead during the normal fire assaying procedure. The Pt bead is arced at 20 A dc in a graphite electrode, and Pt serves as an internal standard. The precision of the method for Ir, Ru, and Os is about 5, 10, and 22% coefficient of variation, respectively. If high accuracy is not required, empirical correction factors may be applied for the assay losses of Ir and Ru, thereby eliminating the need for radiotracers. (auth)

45230 INFRARED ABSORPTION SPECTRA OF POTASSIUM URANIUM FLUORIDES. Soga, T.; Ohwada, K.; Iwasaki, M. (Japan Atomic Energy Research Inst., Tokai). Appl. Spectrosc.; 26: No. 4, 482-3 (1972).

Studies were made to obtain information on U-F vibrations in the Kf-UF_4 system and to relate the band location of the U-F stretching vibration with the change of valency states of uranium. Observed frequencies and tentative assignments are given. (M.C.G.)

45231 MICRODETERMINATIONS OF TERBIUM BY A LUMINESCENT METHOD WITH THE REAGENT PYROGALLOLSULFONIC ACID. Poluektov, N. S.; Alakaeva, L. A.; Tishchenko, M. A. (Inst. of General and Inorganic Chemistry, Odessa). Ukr. Khim. Zh.; 38: No. 2, 175-9 (Feb 1972). (In Russian).

The suitability of the luminescent reaction of Tb with pyrogallousulfonic acid as a quantitative method was studied. The Tb complex formed from a 1:1 ratio of Tb: reagent could be excited by a Hg spectrum at the lines 289.4 and 296.7 nm. The sensitivity of the method was 2×10^{-10} Tb_4O_7 . Tb could be determined in a rare earth mixture or admixed with Y_2O_3 . (tr-auth)

45232 DETERMINATION OF ALLOYING ADDITIVES IN ARTIFICIAL FLUORITE. Kornienko, T. G.; Endrzhhevskaya, S. N. (Inst. of Problems of Materials, Kiev. Inst. of Geochemistry and Physics of Minerals, Kiev). Ukr. Khim. Zh.; 38: No. 2, 194-6 (Feb 1972). (In Russian).

The entire quantity of rare earth elements found in an original charge of an artificial fluorite was determined by disintegrating the sample in H_2SO_4 and paper chromatography. The small amounts of additives were completely intruded into the crystalline lattice of the fluorite. (K.S.W.)

45233 PHOTOMETRIC DETERMINATION OF ZIRCONIUM IN NATURAL MATERIALS WITH THE REAGENT PICRAMINE-EPSILON. Dedkov, Yu. M.; Ermakov, A. N.; Korsakova, N. V. (Inst. of Geochemistry and Analytical Chemistry, Moscow). Ind. Lab. (USSR) (Engl. Transl.); 37: No. 12, 1811-12 (Dec 1971). Translated from Zavod, Lab.; 37: No. 12, 1411-12 (Dec 1971).

A method is described for determination of zirconium in complex geological materials by use of an excess of picramine-epsilon. Spectrophotometric reaction characteristics were calculated. Hf and Nb interfered with the determination. (M.C.G.)

45234 DETERMINATION OF NIOBIUM IN ROCKS, ORES, AND ALLOYS BY ATOMIC-ABSORPTION SPECTROPHOTOMETRY. Husler, John (Univ. of New Mexico, Albuquerque). Talanta; 19: No. 7, 863-9 (Jul 1972).

Niobium, in concentrations as low as 0.02% Nb_2O_5 , was determined in a variety of materials without separation or enrichment. Chemical and ionization interferences were controlled, and sensitivity increased, by maintaining the iron, aluminium, hydrofluoric acid, and potassium content within certain broad concentration limits. There was close agreement with the results of analyses by emission spectrographic, electron microprobe and x-ray fluorescence methods. (auth)

45235 RAPID MASS-SPECTROMETRIC DETERMINATION OF BORON ISOTOPIC DISTRIBUTION IN BORON CARBIDE. Rein, James E.; Abernathy, Robert M. (Los Alamos Scientific Lab., N. Mex.). Talanta; 19: No. 7, 857-62 (Jul 1972).

Boron isotopic ratios were measured in boron carbide by thermionic ionization mass spectrometry with no prior chemical separation. A powder blend of boron carbide and sodium hydroxide was prepared, a small portion transferred to a tantalum filament, the filament heated to produce sodium borate, and the filament transferred to the mass spectrometer where the $^{11}\text{B}/^{10}\text{B}$ ratio was measured, using the Na_2BO_2^+ ion. Variables investigated for their effect on preferential volatilization of ^{10}B included the sodium hydroxide-boron carbide ratio and the temperature and duration of filament heating. A series of boron carbide pellets containing natural boron, of the type proposed for the control rods of the Fast Flux Test Facility reactor, were analyzed with an apparently unbiased result of 4.0560 for the $^{11}\text{B}/^{10}\text{B}$ ratio (standard deviation 0.0087). The pellets contained over 3% metal impurities typically found in this material. Time of analysis was 45 min per sample, with one analyst. (auth)

45236 FLUOROMETRIC DETERMINATION OF ALUMINUM AND GALLIUM WITH SUPERCHROME GARNET Y. Hiraki, Keizo (Kinki Univ., Osaka). *Bull. Chem. Soc. Jap.*; 45: No. 5, 1395-9 (May 1972).

A sensitive fluorometric determination of aluminum and gallium with Superchrome Garnet Y is described. The reagent reacts with aluminum and gallium to form complexes with reddish yellow fluorescence of emission maximum at 562 and 565 nm, respectively. Several conditions for the determination, such as the pH of the solution, the reagent concentration, the time of standing etc. were investigated, and the composition of the complexes was determined. The maximum fluorescence intensity of the aluminum complex was found at pH 5 (heated for 10 min at 50°C), and that of the gallium complex, at pH 3. The fluorescence was stable at least for 2 hr. By using Superchrome Garnet Y, 0.02 to 4 µg of aluminum and 0.5 to 30 µg of gallium in 25 ml solution could be determined. Scandium and indium also formed fluorescent complexes. Iron(III), cobalt(II), copper(II), zirconium(IV), vanadate(V), and chromate(VI) ions interfered with the determination. (auth)

45237 ACCURACY OF QUANTITATIVE ELECTRON PROBE MICROANALYSIS WITH ENERGY DISPERSIVE SPECTROMETERS. Beamman, D. R.; Solosky, L. F. (Dow Chemical Co., Midland, Mich.). *Anal. Chem.*; 44: No. 9, 1598-1610 (Aug 1972).

The accuracy in the measurement of chemical composition using an energy dispersive spectrometer was determined. After considering a large number of analyses performed by a number of investigators and comparing them with a series of analyses performed, it was concluded that if the concentration is greater than about 20 wt %, a relative error of ±6% of the amount present could be expected. At lower concentration, large positive errors were often encountered. The EDS results were compared with those obtained with a crystal spectrometer. Possible sources of error were examined. (auth)

45238 DETERMINATION OF TRACE QUANTITIES OF VOLATILE FLUORIDE IN URANIUM HEXAFLUORIDE USING AN INFRARED SPECTROPHOTOMETER. Aubeau, Raymond; Blandenet, Gerard; Brogniart, Guy (CEA, Pierrelatte, France). *Anal. Chem.*; 44: No. 9, 1628-36 (Aug 1972).

Determination of volatile fluoride in uranium hexafluoride by infrared spectrophotometry in the range 0.1 to 1 weight parts per million (wppm) is described. The instrument used is a spectrophotometer equipped with a 50-cm-long dual path cell. According to the test accomplished and the sensitivity required, three versions are available: direct analysis; differential analysis; and analysis after absorption on sodium fluoride. These different technologies allow the contents of weight parts per million to be determined with a precision of 5 to 10%. (auth)

45239 SPECTROPHOTOMETRIC MOLYBDENUM DETERMINATION WITH THIOLACTIC ACID. Fritz, J. S.; Beuerman, D. R. (Ames Lab., Iowa). *Talanta*; 19: No. 3, 366-9 (Mar 1972).

A quick, selective method for molybdenum(VI) is described, based on the formation of a yellow thiolactic acid complex. The complex is formed at pH 1.0 to 1.6, and the absorbance is measured at 365 nm. (auth)

45240 PHENYLTHIAZOLYLAZOCHROMOTROPIC ACID AS A PHOTOCOLORIMETRIC REAGENT FOR THORIUM. Pachadzhinov, D. N.; Yusupov, M. Yu. (Inst. of Chemistry, Dushanbe, USSR). *Dokl. Akad. Nauk Tadzh. SSR*; 15: No. 2, 33-5 (1972). (In Russian).

The azo dye, phenylthiazolylazochromotropic acid, is proposed for the photolorimetric assay of thorium in small quantities. The following agents do not interfere: alkali metals, alkaline earths, Cd, Pb, Co, and the common anions; small quantities of Cu, Fe, Al, Zr, Ti, and Hf do interfere. The wavelength of maximum absorption is: 565 nm for the reagent dye, and 600 nm for the dye-Th complex. (K.S.W.)

45241 ALLOWANCE FOR MASS-SPECTROMETER BACKGROUND AND THE REPRODUCIBILITY IN THE DETERMINATION OF ISOTOPE COMPOSITIONS FOR NITROGEN AND OXYGEN. Vitol, E. N.; Orlova, K. B.; Ivanenko, M. A. *Ind. Lab. (USSR) (Engl. Transl.)*; No. 12, 1888-91 (Dec 1971).

Translated from *Zavod. Lab.*; 37: No. 12, 1466-9 (Dec 1971). The effects of background on isotope analyses of nitrogen, oxygen, and hydrogen were examined by reference to the N₂, CO, and H₂ peaks, which coincide with the strongest background peaks. (M.C.G.)

45242 EXTRACTIVE PHOTOMETRIC DETERMINATION OF URANIUM(VI) IN MINE WATERS (EXCHANGE OF EXPERIENCE). Shchemeleva, G. G.; Stepanenko, Yu. V.; Kovalenko, P. N. (Rostov State Univ., USSR). *Ind. Lab. (USSR) (Engl. Transl.)*; 37: No. 12, 1837 (Dec 1971).

Translated from *Zavod. Lab.*; 37: No. 12, 1431 (Dec 1971).

The reaction between the benzoic acid complex anion of uranium and Brilliant Green was studied. The optimum conditions were selected for extraction of the resulting complex with toluene. Beer's law was obeyed over the range of uranium concentration 0.2 to 10.7 µg/ml. Ferric ions interfered, but could be reduced to Fe²⁺ by ascorbic acid. The method could be used for the rapid determination of uranyl in mine waters, without preliminary concentration, for uranium contents from 2 × 10⁻⁴ g/liter upwards. (M.C.G.)

45243 AUTOMATIC ANALYSIS BY ABSORPTION SPECTROMETRY. Delmas, Pierre; Imbert, Pierre (to Commissariat a l'Energie Atomique). French Patent 2,070,481. 16 Aug 1971. Filed 5 Dec 1969. (In French).

This automatic spectrophotometric absorption analyzer is intended for the continuous determination of several constituents in liquid samples. Small quantities of ClO₃F, COF₂, UF₆, SiF₄ and ClF₃ can be determined in nitrogen. The apparatus is provided with a luminous source, a transparent tank for the sample, a device for modulating the beam at a given frequency, a grating monochromator, a photoelectric converter, and an amplifier that receives the ac signal from the converter. A rectifier circuit converts the ac signal at the output of the amplifier into a dc amplitude. (France)

45244 ATOMIC ABSORPTION AND FLAME EMISSION SPECTROSCOPY IN THE ANALYSIS OF TRACE METALS IN ATMOSPHERIC PARTICULATES. Hwang, Jae Y. (Instrumentation Lab., Inc., Lexington, Mass.). pp 352-6 of Proceedings of the Second International Clean Air Congress. /England, H. M. (ed.). New York; Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

Analytical conditions of flame and flameless atomic spectroscopy for trace analysis of samples were developed by performing simultaneous analysis of two elements as an approach to multi-element analysis; improving accuracy of the analytical results by reducing or eliminating chemical interferences by optimizing experimental conditions; improving the precision of the analysis by utilizing internal standardization techniques; increasing the sensitivities and minimizing sample solution for the analysis by a flameless absorption measurement; and increasing the sensitivities of the analysis by improving the nebulization efficiency with the counter jet device. Detection limits are given for eighteen elements in air samples including Mo. (D.H.M.)

45245 STABLE ISOTOPES OF CARBON AS INDICATORS OF AIR POLLUTION LEVELS. Atkins, Patrick R. (Univ. of Texas, Austin). pp 545-9 of Proceedings of the Second International Clean Air Congress. /England, H. M. (ed.). New York; Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

Preliminary studies indicate that the stable carbon isotope ratios found in polluted air samples can be used to determine the degree of pollution in the area and the type of source from which the majority of the pollution originated. The cold trap method of separating the CO₂ fraction from the gas sample works well and gives reproducible results. However, the method is cumbersome and relatively slow. The precipitation method using barium hydroxide is much more convenient and requires less equipment. But the results to date are not reproducible and cannot be interpreted directly. A calibration procedure will probably be required before the precipitation method can be used to produce useful data. (auth)

45246 ELECTRON PROBE MICROANALYSIS. Philibert, J. (Institut de Recherches de la Siderurgie, St. Germain-en-Laye, France). pp 419-531 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Topics reviewed include: production of the electron probe, analysis of the emitted x rays, ancillary techniques, image display, generated intensity of the primary characteristic lines, general outline of correction calculations, correction calculation methods, statistical errors connected with photon emission, errors associated with the line spectrum, counter errors, errors associated with the electron beam, and errors associated with the specimens. (161 references) (M.C.G.)

45247 EMISSION SPECTROSCOPY INCLUDING dc ARC, SPARK, AND OTHER METHODS. Mossotti, Victor G. (Univ. of Illinois, Urbana). pp 533-72 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Topics covered include: origin of spectra, atomic structure, spectral characteristics of families of elements, excitation of

spectra, energy sources, measurement of spectra (external optics, wavelength dispersion, transducers), interpretation of spectra, and practical analytical procedures. (63 references) (M.C.G.)

45248 MASS SPECTROMETRIC TECHNIQUES. Honig, Richard E. (RCA Labs., Princeton, N. J.). pp 573-640 of *Modern Analytical Techniques for Metals and Alloys. Pt. 2.* /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

The various mass spectrometric techniques used in the analysis of metals and alloys are discussed, with special emphasis on recent developments. Topics covered include: ion sources, mass analyzer, ion detectors, analytical methods, experimental limitations, data processing, and applications. (223 references) (M.C.G.)

45249 SECONDARY ION EMISSION ANALYSIS. Philibert, J. (Institut de Recherches de la Siderurgie, St. Germain-en-Laye, France). pp 641-56 of *Modern Analytical Techniques for Metals and Alloys. Pt. 2.* /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

Applications of secondary ion emission to localized analysis of massive specimens are discussed. Three modes of operation can be used: depth analysis, ion microscopy, and ion probe. Instrumentation, characteristics of secondary ion emission analysis, problems associated with qualitative and quantitative chemical analysis, and examples are discussed. (12 references) (M.C.G.)

45250 PHYSICAL METHODS OF CHEMISTRY. PART IIB. Weissberger, Arnold; Rossiter, Bryant W. (eds.). *Techniques of Chemistry. Volume I.* New York; John Wiley and Sons, Inc. (1972). 726p.

Separate abstracts were prepared for two of the eight chapters. The remaining chapters consider infrared spectroscopy, Raman spectroscopy, visible and ultraviolet spectrophotometry, determination of fluorescence and phosphorescence, and spectroscopy of monolayer assemblies. (D.H.M.)

For abstracts of individual chapters see: 45251 and 45340.

45251 ATOMIC ABSORPTION SPECTROSCOPY. Ambrose, Robert T. (Eastman Kodak Co., Rochester, N. Y.). pp 485-520 of *Physical Methods of Chemistry. Pt. IIB.* /Weissberger, Arnold (ed.). New York; John Wiley and Sons, Inc. (1972).

A review of atomic absorption spectroscopy is presented that includes theory, scope of the method (sensitivities and detection limits for 68 elements including Ce, Cs, Dy, Er, Eu, Gd, Ho, La, Lu, Mo, Nd, Nb, Pr, Ru, Sm, Sc, Sr, Th, Ta, Te, Tb, Tm, Y, Yb, U, and Zr), equipment, experimental techniques, and applications. (D.H.M.)

45252 ANALYTICAL EMISSION SPECTROSCOPY. PART I. Grove, E. L. (ed.). *Analytical Spectroscopy Series. Volume 1.* New York; Marcel Dekker, Inc. (1971). 413p. \$26.50.

A treatise concerned with the uses of emission spectroscopy and the new advances, changes, and consolidation of ideas in the field is presented. Chapters on principles and instrumentation include the basic physics of optics and the development of principles and instrumentation from earlier instruments to those in use today. Areas considered are: historical development, some uses and definitions; origins of atomic spectra; prism systems, spectrographs, and spectrometers; gratings and grating instruments; and spectroradiometric principles. (D.H.M.)

45253 ANALYTICAL EMISSION SPECTROSCOPY. PART II. Grove, E. L. (ed.). *Analytical Spectroscopy Series. Volume 1.* New York; Marcel Dekker, Inc. (1972). 579p. \$35.75.

A treatise concerned with the uses of emission spectroscopy and the new advances, changes, and consolidation of ideas in the field is presented. Areas covered are: excitation of spectra; flame spectrometry; qualitative and semiquantitative analysis; and quantitative analysis. (D.H.M.)

45254 DETERMINING BORON IN GLASS BY FLAME EMISSION SPECTROSCOPY. Morrow, Roy W. (Oak Ridge Y-12 Plant, Tenn.). *Anal. Lett.*; 5: No. 6, 371-6(1972).

A method was developed that will determine the percentage quantities of boron in borosilicate glass by flame emission spectroscopy. Raschig rings of borosilicate glass are used as neutron absorbers during the processing and shipping of radioactive solutions. Since the effectiveness of these rings is dependent on the boron content, analytical control is a necessity. The sample is dissolved in hydrofluoric acid, nebulized into a nitrous oxide-acetylene flame, and a measurement made of the BO_2 band emission at 5180 angstroms. Glass containing at least 1.0 wt% boron can be analyzed. The relative standard deviation of the method is 5.1%. Approximately 10 samples (2 aliquots each) can be analyzed in four hours. (auth)

45255 DETERMINATION OF SILICA IN SILICATES CONTAINING PHOSPHORUS, TITANIUM, AND ZIRCONIUM. Ohl-

weiler, O. A.; Meditsch, J. O.; Silveira, Carmem L. P.; Silva, Sandra (Instituto de Quimica, Porto Alegre, Brazil). *Anal. Chim. Acta*; 61: No. 1, 57-65(Aug 1972).

A procedure based on precipitation of oxine 12-molybdosilicate was examined for the determination of silica in silicates containing phosphorus, titanium, and zirconium. Boron trioxide and lithium carbonate are used in the flux. The fusion product is dissolved in 0.01M hydrochloric acid containing EDTA, in contact with Amberlite IR-120 (H^+). The resultant solution, after stirring with Amberlite IRA-400 (HCO_3^-), is used for the precipitation of silica as oxine 12-molybdosilic acid. (auth)

45256 POLAROGRAPHIC DETERMINATION OF TIN IN COPPER-, ALUMINUM-, AND ZIRCONIUM-BASE ALLOYS. Mukoyama, Tomoyuki; Yamane, Takeshi; Kiba, Nobutoshi; Tanaka, Masao (Yamanashi Univ., Takeda, Japan). *Anal. Chim. Acta*; 61: No. 1, 83-85(Aug 1972).

An alternating-current polarographic method is described for determining tin in copper-, aluminum- and zirconium-base alloys. With EDTA as masking agent and coprecipitation with beryllium hydroxide, tin is separated selectively from other metal ions and determined polarographically in a 3M hydrochloric acid as supporting electrolyte. The method is simple, selective and accurate. A single determination takes about 45 min. (auth)

45257 NANOGRAM DETERMINATION OF IODIDE BY MEANS OF ISOTOPE EXCHANGE. Gabrielsson, Ann-Britt; Beronius, Per (Univ. of Umea, Sweden). *Anal. Chim. Acta*; 61: No. 1, 123-6(Aug 1972).

A study is reported on using the isotope exchange reaction $\text{CH}_3\text{I} + \text{Na}^{131}\text{I} \rightleftharpoons \text{CH}_3^{131}\text{I} + \text{NaI}$ to determine nanogram quantities of iodide. Results indicate that amounts down to 50 ng may be determined with only a few percent error. (D.H.M.)

45258 DEPENDENCE OF THE FORMATION OF URANIUM AND THORIUM IONS AND THEIR OXIDE IONS ON VARIOUS CONDITIONS OF THERMAL IONIZATION. Heumann, K. G. (Technische Hochschule, Darmstadt, Ger.). *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 315-24(Aug 1972). (In German).

The dependence of the formation of uranium ions, thorium ions, and their oxide ions on various conditions of thermal ionization was investigated. A single-focusing mass spectrometer with a double-filament thermal ion source was used. The filaments were made of rhenium. The samples were applied in the form of uranyl nitrate and thorium nitrate. The temperatures of the ionization filament and the evaporation filament were varied. It was found that the best conditions to measure the isotopic ratio were achieved if the difference in temperature between the ionization filament and the evaporation filament is as high as possible. The isotopic ratios of uranium and thorium were determined by measuring the ions Th^+ and U^+ rather than the ions ThO^+ and UO^+ . In addition the mass discrimination during the evaporation of the sample was investigated. The precision of the isotopic ratio measurements is shown by the determination of the ratio $^{234}\text{U}/^{238}\text{U}$ in natural uranium and of the ratio $^{230}\text{Th}/^{232}\text{Th}$ in a ^{230}Th solution. (auth)

45259 SYSTEMATIC STUDY OF THE DECOMPOSITION OF VERY SMALL QUANTITIES OF RARE EARTH OXIDES BY MASS SPECTROMETRY. Nguyen Long-Den; de Saint-Simon, M. (Institut de Physique Nucleaire, Orsay, France). *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 299-314(Aug 1972).

The ratio $i(\text{M}^+)/i(\text{MO}^+)$ was measured as a function of temperature by mass spectrometry. Thermal ionization of small quantities of rare earth oxides, which were deposited on a rhenium filament, was used. The results are interpreted starting from the decomposition of the monoxides, their reduction by rhenium, and the thermal ionization of the products formed. The quantities that entered into the calculations were the dissociation energy $D_0^\circ(\text{MO})$ and the atomic number Z of the investigated element. Good agreement was obtained with experimental results. With this method it seems possible to determine the dissociation energy $D_0^\circ(\text{MO})$ and to detect the presence of crystal transformation of an element in very small quantities. As an application, these results should permit one to identify for lanthanides both Z and A simultaneously, since A is determined by the magnetic field of the mass spectrometer. (auth)

45260 ANALYSIS OF CURIUM AND CALIFORNIUM BY SINGLE-EXPOSURE SPARK SOURCE MASS SPECTROMETRY. Chastagner, P.; Tiffany, B. (E. I. du Pont de Nemours and Co., Aiken, S. C.). *Contract AT(07-2)-1.* *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 325-32(Aug 1972).

Single-exposure spark source mass spectrometry is used for routine quality control analyses of ^{244}Cm and ^{252}Cf at the Savannah River Laboratory. All elements above mass 10, except carbon, nitrogen, oxygen, and the noble gases, are determined as are the isotopic distributions of fission products ruthenium, palladium, cerium, neodymium, samarium, europium, gadolinium, dysprosium,

and erbium. Samples of from 1 to 30 μg of curium and 0.0001 to 0.1 μg of californium are mounted on gold electrodes for analysis. Indium is added as an internal standard to californium samples smaller than 0.005 μg , and ytterbium is added as an internal standard to all samples when photometric or refined visual interpretation is required. Visual interpretation usually yields impurity results accurate within a factor of three and relative standard deviations for isotopic distributions of about 35%. (auth)

Radiometric and Radiochemical Procedures

Refer also to abstracts: 45585, 45587, and 45604.

45261 (ICRU-22) MEASUREMENT OF LOW-LEVEL RADIOACTIVITY. (International Commission on Radiation Units and Measurements, Washington, D. C.). 1 Jun 1972. 66p. ICRU Publications, Washington, D. C. \$3.50.

The assay of gamma- and x-ray emitters is discussed under the headings: composition and reduction of background, scintillation counters, semiconductor detectors, proportional counters, special techniques, and analysis of gamma spectra. Topics discussed concerning the assay of beta emitters include gas-discharge counters, plastic scintillation counters, liquid scintillation counters, Cherenkov counters, semiconductor detectors, and procedures for specific radionuclides. The assay of alpha emitters is discussed including background, sensitivity, classification of alpha particle detection techniques, measurement of total emission rate of α particles in solid sources, measurement of α particle emitters in the gaseous phase, spectrometry of alpha particles, and the nuclear-track technique. Also, standardized sources and reference sources for low-level procedures are briefly discussed. An extensive list of references is included. (W.H.K.)

45262 (NIM-1427) PRESSED FUSED DISC: A TECHNIQUE FOR THE PREPARATION OF ORE SAMPLES TO BE ANALYSED FOR URANIUM BY X-RAY FLUORESCENCE. Austen, C. E.; Russell, B. G. (National Inst. for Metallurgy, Johannesburg (South Africa)). 28 Mar 1972. 8p. Dep. NTIS (U. S. Sales Only).

Because of the structure of some uranium minerals, erroneous results are sometimes obtained when x-ray-fluorescence measurements are made on the powdered rock. Samples are therefore fused, prior to analysis, with sodium tetraborate or lithium tetraborate. A modified fusion technique is described in which the rock sample is fused with fluorspar and lithium tetraborate, and the melt is pressed into a disc. The precision of x-ray-fluorescence analyses for uranium on the pressed fused discs is shown to be similar to that on discs prepared in the old way, but the preparation of the modified discs is much simpler and quicker. (auth)

45263 (RT/PROT-72-11) MEASUREMENT OF ^{137}Cs AND ^{90}Sr IN SOIL. Baglino, G.; Fratarcangeli, S.; Lalli, M. G.; Toerok, I. (Comitato Nazionale per l'Energia Nucleare, Rome (Italy)). 30 Mar 1972. 27p. (In Italian). Dep. NTIS (U. S. Sales Only).

An analytical method is described for determination of ^{90}Sr , Ca, ^{137}Cs , and K in soil samples of different chemical composition. Also reported are the results of analysis performed on 30 soil samples taken at several different sampling points. (auth)

45264 (SRARI-P-123) DETERMINATION OF THE CONCENTRATION OF GAMMA-EMITTING ISOTOPES IN EXTENDED SOURCES WITHOUT SAMPLING. Polyakov, V. I.; Chechetkin, Yu. V. (Scientific Research Atomic Reactor Inst., Melekes (USSR)). 1971. 28p. (In Russian). Dep. NTIS (U. S. Sales Only).

A scintillation γ spectrometer with a collimator is used to measure bulk and surface concentrations of isotopes in different source configurations including cylindrical ones. Analytical expressions are given enabling, as it is shown experimentally, specific activity determination with an error of 10 to 15%. The methods of calibrating a scintillation spectrometer without a collimator are described. (auth)

45265 (UCRL-51061(Vol.1)) COMPUTERIZED QUANTITATIVE ANALYSIS BY GAMMA-RAY SPECTROMETRY. VOLUME I. DESCRIPTION OF THE GAMANAL PROGRAM. Gunnink, R.; Niday, J. B. (California Univ., Livermore, Lawrence Livermore Lab.). 1 Mar 1972. Contract W-7405-eng-48. 78p. Dep. NTIS.

GAMANAL provides a complete qualitative and quantitative analysis of mixtures of radioactive species by computer interpretation of high-resolution gamma spectra. This volume describes the methods used for determining background, resolving and fitting complex peak groupings, determining the energies and absolute intensities of the gamma rays, and determining the amounts of the source nuclides. The latter are identified with the aid of a preprocessed decay-scheme library, and all inter-

ferences are resolved by a least-squares solution of the matrix of equations for the gamma intensities. (auth)

45266 IMPROVED METHOD FOR DETERMINATION OF ^{35}S IN PLANT MATERIAL USING OXYGEN FLASK COMBUSTION AND LIQUID SCINTILLATION COUNTING. Grundon, Noel J. (Univ. of Queensland, St. Lucia, Australia); Asher, Colin J. J. Agr. Food Chem.; 20: No. 4, 794-8(1972).

A safe oxygen flask combustion procedure is described for preparation of ^{35}S -labeled plant material for liquid scintillation counting. Hazards associated with flammable absorbing solutions are eliminated by using deionized water to trap the sample combustion products from the oxygen flask. Sulfur-35 activity in the resulting aqueous solution is then determined in a 1:1:2 mixture of toluene, Triton X-100, and water with an efficiency of 73%. With a 3-liter oxygen flask and a 20-min absorption period, the relationship between sample size and activity recovered was linear up to a sample weight of 1.2 g. Recoveries of ^{35}S added to plant material as sodium sulfate- ^{35}S or cysteine hydrochloride- ^{35}S were $100.0 \pm 2.3\%$ and $99.1 \pm 1.2\%$, respectively. Using five combustion flasks and an automatic liquid scintillation counter, a single operator can process about 50 samples per day. (auth)

45267 APPARATUS FOR MEASURING THE D_2O CONTENT IN A FLOW OF WATER. Strindehag, O. M. (to Aktiebolaget Atomenergi). Canadian Patent 893,931. 22 Feb 1972. Priority date 30 Oct 1969, Sweden.

The D_2O content of a flow of water is measured by passing the flow through a measuring chamber containing a gamma-radiating body consisting of a long-lived radioactive isotope in radioactive equilibrium with a short-lived radioactive gamma-radiating affiliated isotope having a gamma energy of more than 2.23 MeV. Photoneutrons are formed as a result of the heavy hydrogen's being bombarded with high-energy gamma photons. The photoneutrons are detected and counted by means of a neutron detector. (Canada)

45268 DETERMINATION OF Pu IN THE ENVIRONMENT BY EXTRACTION AND LIQUID SCINTILLATION COUNTING. Ghysels, J. P. Ind. At. Spatiales; 26: No. 1, 33-6(Jan 1972). (In French and English).

The extraction of Pu was effected with liquid ion exchangers (e.g., tertiary amines such as triisooctylamine). The Pu was put back into organic solution by extraction with DEHPA. After mixing with a liquid scintillator, the counting yield was determined (close to 100%). This method of analysis is intended for determination of Pu in biological substances. It was applied to the analysis of radioactive dusts of different origin. (France)

45269 NONDESTRUCTIVE ASSAY OF POWER REACTOR FUEL ASSEMBLIES. Weisbin, C. R. (Los Alamos Scientific Lab., N. Mex.); Augustson, R. H.; Hendricks, J. S.; (and others). Nucl. Technol.; 15: No. 3, 455-61(Sep 1972).

The total ^{235}U content of low enrichment power reactor fuel assemblies was determined by interrogation with Li(p,n) neutrons (200 to 500 keV) produced by a Van de Graaff proton accelerator and by measurement of fission induced delayed neutrons. To extend the application of this assay technique and to optimize its implementation, numerical calculations were performed using a modified Monte Carlo approach. Experimental measurements from a BWR-type fuel assembly verified the mathematical model employed. By incorporating results from both calculation and measurement, a system is being designed to assay the total ^{235}U content of power reactor fuel elements with an overall uncertainty within 2%. Using commercially available fuel handling equipment, the analysis time for the entire element is estimated to be approximately 30 minutes. (auth)

45270 X-RAY FLUORESCENCE ANALYSIS OF HEAVY ELEMENTS BY AN α -X COINCIDENCE METHOD. Aleksandrov, B. M.; Solov'ev, S. M.; Tyvin, L. I.; Eismont, V. P. At. Energ. (USSR); 32: No. 6, 491-3(Jun 1972). (In Russian).

The method of α -x coincidence was used to determine small quantities of ^{241}Am , $^{242-244}\text{Cm}$, and ^{252}Cf by using the characteristic L radiation that accompanies α decay. Typical spectra of α particles from the targets mentioned and of L radiation are plotted. (3 figures) (K.S.W.)

45271 ELASTIC NUCLEAR SCATTERING TECHNIQUES. Rubin, Sylvan (Stanford Research Inst., Menlo Park, Calif.). pp 657-85 of Modern Analytical Techniques for Metals and Alloys. Pt. 2. /Bunshah, R. F. (ed.). New York; John Wiley and Sons, Inc. (1970).

The theory of the method, including velocity change on scattering, scattering cross section, stopping power, and scattering yield, is discussed. Equipment and measurement problems and limitations are described. (20 references) (M.C.G.)

45272 RADIOMETRIC DETERMINATION OF TRACE AMOUNTS OF ANIONIC SURFACTANTS IN GROUND WATER AND POTABLE WATER. Taylor, C. G.; Waters, J. (Liverpool Polytechnic). Analyst; 97: No. 1156, 533-41 (Jul 1972).

The water sample is treated with an excess of ferroin solution labelled with iron-59. The solution is extracted with chloroform and the activity of a portion of the extract is measured. The anionic surfactant content of the sample is determined by the standard addition of sodium lauryl sulphate. The application of the method to river-water and tap-water samples gives results that agree with those obtained by using Abbott's methylene blue method. The present method requires 30 minutes for each reading, i.e., blank, sample, or sample plus standard. The over-all standard deviation for river-water samples was calculated to be $\pm 4.3\%$ at the level of 0.1 ppM of sodium lauryl sulphate. The limit of detection is 0.1 μg of sodium lauryl sulphate (0.005 ppM in a 20-ml sample). This limit is the equivalent of the average blank. Blanks are minimized by extracting at pH 4 with chloroform equilibrated with water, by performing only one extraction, by centrifuging for a fixed period and by removing the extract for counting with a capillary device. No species have been found to interfere when present at the levels that are normally encountered in river waters. Potential interferences by fluoride or iron are compensated for by the internal standard. (auth)

Separation Procedures

Refer also to abstracts: 45302 and 47014.

45273 (UJV-2820-Ch) STRONTIUM SORPTION ON THE SORBENTS OF BaSO_4 - CaSO_4 SYSTEM. Vojtech, O.; Selucka, H. (Ceskoslovenska Akademie Ved, Rez, Ustav Jaderneho Vyzkumu). 1972. 29p. Dep. NTIS (U. S. Sales Only).

The sorption of strontium on sorbents of the BaSO_4 - CaSO_4 system was studied in detail over the whole concentration range of this binary system. The sorbents were prepared by two different methods of thermal treatment which led to a formation of stable and metastable polycrystalline powders. Besides the kinetics of sorption, also course of dissolution of the CaSO_4 component was studied and specific surface areas of metastable sorbents were measured. Results led to conclusions which explain some specific sorption properties of the studied sorbents. Metastable mixed crystals BaSO_4 - CaSO_4 with a higher content of CaSO_4 were found to be good sorbents of strontium, regardless of the type of their crystal lattices. (auth)

45274 ZIRCONIUM TETRAFLUORIDE PROCESS. Craigen, W. J. S.; Joe, E. G.; Ritcey, G. M.; Melvanin, F. W.; Smart, B. C. (to Eldorado Nuclear Ltd.). Canadian Patent 899,029. 2 May 1972.

Anhydrous zirconium tetrafluoride can be prepared by preparing a clear solution of zirconyl nitrate in a nitric acid solution (at least 8N, with zirconium content between 50 and 150 g/l), precipitating zirconium tetrafluoride monohydrate by adding hydrofluoric acid at a temperature of at least 30°C, isolating and drying the precipitate, and using acid solution remaining after precipitation for the preparation of the zirconyl nitrate solution. Recycling of the acid solution must be accompanied by a bleed to prevent too much build-up of metallic impurities such as titanium or uranium salts of higher solubility than zirconium tetrafluoride. This removal of a portion of the reused acid also, when the feed material is zirconyl nitrate, prevents the nitrate ion concentration from becoming excessively high. Greater than 90% recovery of the metal ion from solution may be obtained from this process. (E.M.T.) (Canada)

45275 PAPER CHROMATOGRAPHY OF THE RARE EARTH ELEMENTS WITH THE DEVELOPER CONTAINING OF ETHYLENE GLYCOL MONETHERS, HYDROPHOBIC SOLVENTS, AND OXINE. Nagai, Hideo; Deguchi, Toshio; Nakai, Kiyoto (Kumamoto Univ., Japan). Jap. Anal.; 21: No. 6, 788-93 (Jun 1972). (In Japanese).

Separation of lanthanides (from La to Tb) was carried out on filter paper (Toyo Roshi No. 5B, 1.5×30 cm) impregnated with potassium chloride. Developing solvents used consisted of ethylene glycol monoether, oxine, acetate buffer and hydrophobic solvents such as chloroform. Test solutions of lanthanides were prepared by dissolving their nitrates in 1M nitric acid containing 2.0 to 2.5 mg of each ion per ml of the solution. About 0.2 μl of test solution was applied at the starting position of the filter paper strip in a narrow straight line and then developed as the usual ascending paper chromatography. The following five kinds of developers were used: (A) 55 ml of carbitol, 15 ml of 1M acetic acid-1M sodium acetate buffer (pH ca. 5), 30 ml of chloroform and 4.0 g of oxine, (B) 55 ml of carbitol, 15 ml of the acetate buffer (pH ca. 5), 30 ml of chloroform, and 3.5 g of oxine, (C) 45 ml of methyl cellosolve, 10 ml of the acetate buffer (pH ca. 4.5), 45 ml of chloroform and

2.0 g of oxine, (D) 40.5 ml of methyl cellosolve, 7.5 ml of the acetate buffer (pH ca. 4.5), 52 ml of chloroform and 2.5 g of oxine, and (E) 50 ml of cellosolve, 15 ml of the acetate buffer (pH ca. 4), 35 ml of chloroform and 2.0 g of oxine. Using the developer A or D, quaternary mixtures such as La-Ce-Pr-Nd were sufficiently separated. With the developer B the separation of Sm-Eu-Tb or Nd-Gd-Tb was achieved. Separation of La-Ce-Pr-Sm was possible by using the developer C. Separation of La-Ce-Pr, Ce-Pr-Sm, Nd-Sm-Gd or Eu-Tb was also possible with the developer E. In these five developers, A and D showed the most widely applicable properties. When a hydrophobic solvent such as chloroform was increased in a developer, the R_f value of the ionic band was depressed-reducing the tailing of the band. Accordingly, the optimum addition of chloroform was investigated to obtain the maximum difference of the R_f values between the two adjacent bands maintaining the minimum tailing of their bands. These optima were usually obtained at a slight excess addition of chloroform beyond the saturation of the developers. The hydrophobic solvents investigated other than chloroform were carbon tetrachloride, benzene, 1-butanol, 1-butyl acetate, diethyl ether, and ethyl methyl ketone, but chloroform showed the best result of all the developers used in this experiment. (auth)

45276 SUBSTANCE PURIFICATION BY ZONE MELTING IN A ROTATING CONTAINER. Kirgintsev, A. N.; Lastushkin, N. I.; Rybin, Yu. A.; Avvakumov, E. G. (Inst. of Inorganic Chemistry, Novosibirsk, USSR). Izv. Sib. Otd. Akad. Nauk SSSR, No. 9, Ser. Khim. Nauk; No. 4, 54-60 (Jul 1971). (In Russian).

Zone-purification plants designed as rotating containers around a horizontal axis or an axis located at a certain angle to the main horizon are described. The features of containers are described and the representative examples of substance (germanium, bismuth, antimony, tellurium) purification are given. High-speed (over 140 revs/min) and not low-speed (1-5 revs/min) is believed to be in order for zone purification. (auth)

45277 SOLVENT EXTRACTION OF EUROPIUM BY VERSATIC ACID 911. Shigematsu, Tatsuhiko; Nishimura, Sanji; Tanabe, Teruo; Kondo, Yoshio (Kyoto Univ.). J. Jap. Inst. Metals; 36: No. 5, 445-50 (May 1972). (In Japanese).

Equilibrium in the solvent extraction of europium by Versatic Acid 911 was studied. The organic phase was composed of Versatic Acid and benzene, and the aqueous solution of $\text{Eu}(\text{NO}_3)_3$ and HNO_3 was used. The ionic strength of the latter solution was maintained at unity by adding NaNO_3 . Radioactive ^{152}Eu was used as a tracer. From the measurements carried out at 25°C, the equation of extraction and the apparent equilibrium constant was determined as follows: $\text{Eu}^{3+}(\text{aq}) + 3.5 \text{R}_2\text{H}_2(\text{org}) \rightarrow \text{EuR}_3 \cdot 4 \text{RH}(\text{org}) + 3 \text{H}^+(\text{aq})$ and $K = [\text{EuR}_3 \cdot 4 \text{RH}]_0 [\text{H}^+]_{\text{aq}}^3 / [\text{Eu}^{3+}]_{\text{aq}} [\text{R}_2\text{H}_2]_0^{3.5} = 1.12 \times 10^{-3}$ where R_2H_2 refers the dimeric form of Versatic Acid in benzene solution. (auth)

45278 THIN-LAYER CHROMATOGRAPHY OF METAL IONS ON MICROCRYSTALLINE CELLULOSE IMPREGNATED WITH POTASSIUM CHLORIDE. Ishida, Koji (Kyorin Univ., Hachioji, Japan); Nakagomi, Yoshihito. Jap. Anal.; 21: No. 6, 810-12 (Jun 1972). (In Japanese).

Thin-layer chromatographic behavior of 32 metal ions on microcrystalline cellulose, Avicel SF, as well as that impregnated with potassium chloride was investigated in ethanol-6M hydrochloric acid media. R_f values of most ions on the latter thin-layer decreased with increasing ethanol concentration. This chromatographic behavior suggests that the formation of anionic chloro complexes and of slightly soluble potassium double salts may be responsible for the distribution of the metal ions on the microcrystalline cellulose impregnated with potassium chloride. The chromatographic system involving the chloride impregnated layer makes it possible to accomplish many separations of analytical interest which are difficult on the pure cellulose. (auth)

45279 RADIOMETRIC INVESTIGATION OF THE ABSORPTION OF POTENTIAL-DETERMINING IONS FROM AQUEOUS SOLUTIONS OF NON-IONOGENIC SURFACE-ACTIVE SUBSTANCES. Kabysch, G. M.; Aleinikov, V. G.; Glazman, Yu. M.; Strazhesko, D. N. (Kiev Technological Inst. of Light Industry. Inst. of Physical Chemistry, Kiev). Ukr. Khim. Zh.; 38: No. 2, 163-4 (Feb 1972). (In Russian).

The absorption of I^- ions by AgI powder in the presence of a series of alkyl ethers of polyethylene glycol with different lengths of hydrocarbon radicals and oxyethylene chains was studied. The labeled KI solution contained about 75% of ^{131}I , and as a result of isotopic exchange the activity of the KI decreased by a factor of ten. On addition to the solution of different nonionogenic surface active substances, the potential-determining I^- ions were not desorbed from the surface of the AgI. (tr-auth)

45280 SEPARATION OF THE RARE EARTH ELEMENTS BY ANION EXCHANGE. I. ANION EXCHANGE RESIN SEPA-

RATION OF Gd, Eu, Sm, Nd, AND Pr IN METHANOL-ACETIC ACID-NITRIC ACID MIXTURES. Kurokawa, Kazuo (Tokyo Metropolitan Hakuo Senior High School); Kakihana, Hidetake. Nippon Kagaku Kaishi; No. 5, 925-9 (May 1972). (In Japanese).

Ion exchange separation of gadolinium, europium, samarium, neodymium, and praseodymium was studied using mixed solutions of methanol-acetic acid-nitric acid. Satisfactory separation was obtained with the anion exchange resin column of Amberlyst-A 29 (nitrate form). The rare earths were strongly adsorbed on the top of the column from solutions consisting of 90% methanol, 5% glacial acetic acid, and 5% 7 mol/l nitric acid. Then, Gd, Eu, Sm, Nd, and Pr were eluted in this order by passing successively the four kinds of eluents: (flow rate: 0.7 to 0.8 ml/min). (1) MeOH-17 mol/l HOAc-7 mol/l HNO₃ (volume ratio: 90:5:5). (2) MeOH-10 mol/l HOAc-7 mol/l HNO₃ (80:15:5). (3) MeOH-8 mol/l HOAc-7 mol/l HNO₃-H₂O (75:15:5:5) and (4) H₂O. It is confirmed that up to 0.007 mmol of Gd, Eu, Sm, Nd, and Pr are satisfactorily separated with a column of 1.1 cm diameter and 21 cm length. (auth)

45281 RAPID SEPARATION OF METAL CHELATES BY COLUMN LIQUID-LIQUID CHROMATOGRAPHY USING ULTRA-VIOLET DETECTION. Huber, J. F. K. (Univ., Amsterdam); Kraak, J. C.; Veening, Hans. Anal. Chem.; 44: No. 9, 1554-9 (Aug 1972).

The separation of metal- β -diketonates was accomplished by column liquid-liquid chromatography. Included in the study are the acetylacetonates and the trifluoroacetylacetonates of Be(II), Al(III), Cr(III), Fe(III), Co(II), Co(III), Ni(II), Cu(II), Zn(II), Zr(IV), and Ru(III). A ternary liquid-liquid system composed of water, 2,2,4-trimethylpentane, and ethanol was employed. The water-rich phase was used as the stationary medium while the water-poor phase served as eluent. Several of these two phase systems, varying in quantitative composition, were evaluated for chromatographic selectivity. The addition of a trace of chelating ligand to the phase system suppressed undesirable hydrolysis reactions of chelates. A six-component mixture of metal acetylacetonates can be separated in less than 25 minutes. (auth)

45282 COMBINED ION EXCHANGE-SOLVENT EXTRACTION (CIESE) STUDIES OF METAL IONS ON ION EXCHANGE PAPERS. SYNERGISTIC EFFECTS AND CHROMATOGRAPHIC SEPARATIONS. De, Anil K.; Bhattacharyya, Chitta R. (Dept. of Chemistry, Visva-Bharati, India). Anal. Chem.; 44: No. 9, 1686-8 (Aug 1972).

SA-2 and SB-2 ion exchange papers were used to describe several selected chromatographic systems. Cation exchange systems, anion exchange systems, and synergistic effects are discussed. Separations achieved are given for Nb-Ce-La, Zr-Ce-Th, and V-Nb-Ta systems. (M.C.G.)

45283 SELECTIVE ION EXCHANGERS WITH ISONITROSO-ACETAMIDE FUNCTIONAL GROUPS. II. PROPERTIES AND APPLICATION. Sykora, V.; Dubsky, F. (Inst. of Chemical Tech., Prague). Collect. Czech. Chem. Commun.; 37: No. 5, 1504-13 (May 1972).

Selective ion exchangers with isonitrosoacetamide functional groups ($-\text{NH} \cdot \text{CO} \cdot \text{CH}=\text{NOH}$) attached to a polycondensate of m-aminophenol with formaldehyde (I) or to a styrene-divinylbenzene copolymer (II) were evaluated from a static and a dynamic point of view. The two types of ion exchanger have proved to have practically identical properties, given by the reactivity of isonitrosoacetanilide. The high selectivity of the ion exchangers for Pd²⁺ was utilized to separate palladium in its dilute solutions from an excess of ordinary metals and silver. The exchange capacities for both types of ion exchanger were determined for several cations, including UO₂²⁺. (auth)

45284 SELECTIVE EXTRACTION OF GALLIUM WITH N-BENZYLANILINE IN CHLOROFORM FROM HYDROCHLORIC ACID MEDIA. Khosla, M. M. L. (Defence Lab., Jodhpur, India); Rao, S. P. Anal. Chim. Acta; 61: No. 1, 156-8 (Aug 1972).

N-benzylaniline in chloroform is reported as a new extractant for gallium from hydrochloric acid media. Gallium(III) is readily and quantitatively extracted from 6.5 to 7.5M hydrochloric acid; the phase separation is rapid, and gallium can be easily stripped from the organic phase with water and determined by EDTA titration. The extraction is highly selective if thioglycolic acid is added as masking agent. (D.H.M.)

45285 CHEMISTRY OF GALLIUM [1-5]. PART VI. SEPARATION OF GALLIUM FROM FOREIGN ELEMENTS. Rafaeloff, R. (Soreq Nuclear Research Centre, Yavne, Israel). Radiochim. Acta; 17: No. 2, 113-14 (Jul 1972).

Methyl isobutyl ketone was used for the separation of gallium(III) from Al, Sc, Y, La, Eu(III), Tm, Dy, Ge(IV), Cu(II) and Fe(III). The effect of acid concentration, gallium concentration and foreign elements was studied. The gallium can be quantitatively stripped from the organic phase by washing with small quantities of water. (auth)

Inorganic, Organic, and Physical Chemistry

Refer also to abstracts: 54310 and 45311.

Chemical and Physico-chemical Properties

Refer also to abstracts: 45259, 45353, 45388, 45417, 45606, and 45978.

45286 (MLM-1913) CARBON-13: A SELECTED ABSTRACT BIBLIOGRAPHY. Stringham, W. S. (comp.) (Mound Lab., Miamisburg, Ohio). 15 May 1972. Contract AT(33-1)-Gen-53. 205p. Dep. NTIS.

A listing of 627 selected references and abstracts is presented of chemical and physical properties, analytical and production methods, and uses of carbon-13. The abstracts are listed chronologically and numbered serially for each year from 1962 through 1971. An authors index and key word index are included. (auth)

45287 (SC-R-722769) FUSED SALTS. A Bibliography. VOLUME I. CITATIONS LISTING. VOLUME II. SUBJECT/TITLE INDEX. VOLUME III. AUTHORS INDEX. Clark, Paul V. (comp.) (Sandia Labs., Albuquerque, N. Mex.). Jun 1972. 1149p. Dep. NTIS \$12.00.

These Volumes are issued separately, but cataloged as a unit.

This publication incorporates the bibliographic data published in SC-R-68-1680 (Fused Salt Mixtures: Eutectic Compositions and Melting Points); SC-R-69-1386 (Fused Salt Mixtures: Specific Conductivity Tables); and, through reference number 6373 of SC-R-71-3315 (Fused Salts, A Bibliography). It is provided to furnish supplementary access points (subject terms and title words) not available in the original publications. (auth)

45288 (TID-26106) CHEMISTRY OF HTGR. de Nordwall, H. J. (Oak Ridge National Lab., Tenn.). 1972. 44p. Dep. NTIS.

Chemical problems associated with the HTGR include: control of helium purity, boiler deposits, evaluation, fuel studies, fission product studies, and coolant studies. These problems are discussed using tables and graphs. (M.C.G.)

45289 METHOD OF PREPARING SECONDARY ALKALINE AMINE DERIVATIVES. Lambert, Irma; Ravoire, Jean (to Commissariat a l'Energie Atomique). French Patent 2,088,610. 13 Dec 1971. Filed 17 Apr 1970. (In French).

A method for preparing alkaline derivatives of secondary amines having an RR'NM formula (R and R' are aliphatic radicals with between 1 and 6 carbon atoms and M is an alkali metal, i.e., Li, Na, K, Rb, or Cs) is described. The method consists of reacting an amine having an RR'NH formula with an alkaline amide, MNH₂, in suspension in the amine solution. The suspension is obtained by adding the amine to the alkaline amide in ammonia. These secondary alkaline amine derivatives are used as catalysts in isotopic exchange reactions between the corresponding amines and hydrogen. (France)

45290 ORGANIC DISULFIDES AND RELATED SUBSTANCES. 33. SODIUM 4-(2-ACETAMIDOETHYL)DITHIOBUTANESULFINATE AND RELATED COMPOUNDS AS ANTIRADIATION DRUGS. Field, Lamar; Khim, Yong H. (Vanderbilt Univ., Nashville). J. Med. Chem.; 15: No. 3, 312-15 (Mar 1972).

The compound AcNH(CH₂)₂SS(CH₂)₄SO₂Na (1), a promising anti-radiation drug, disproportionates far more rapidly in solution to the symmetrical disulfides 2 (the amide) and 3 (the sulfinate) than would be expected from behavior of analogous compounds having no SO₂Na moiety; solid 1 is stable. The disproportionation is reversible. An improved preparation of 1 minimizes disproportionation by use of lower temperatures, precipitation instead of concentration, and minimum contact with polar solvents. Radioprotective activity of an improved 1 confirms earlier results; 3 is active as well, but 2 is inactive. The trisulfide [NaO₂S(CH₂)₄S]₂S (9) was prepared in 80% yield from 1,2-dithiane 1,1-dioxide (6) using Na₂S. Thus far, 9 has shown ALD₅₀ > 900 mg/kg (ip or po) and 73 to 100% protection of mice at doses of 38 (ip)-300 (po) mg/kg in mice. The promising activity of 3 and (especially) 9 is noteworthy since neither contains a nitrogen function. (auth)

Isotope Effects

Refer also to abstracts: 45342 and 45343.

45291 MICROWAVE SPECTRUM, STRUCTURE, AND DIPOLE MOMENT OF ETHYLENE SULFONE. Kim, Hyunyoung (Univ. of Missouri, Columbia). J. Chem. Phys.; 57: No. 3, 1075-7 (1 Aug 1972).

The microwave rotational spectra of the ^{32}S and ^{34}S isotopic species of ethylene sulfone were observed. The rotational constants A, B, and C in megahertz are: 6676.06, 4924.01 and 3759.97 for the ^{32}S species; and 6679.20, 4917.07, and 3757.64 for the ^{34}S species. The structural parameters obtained are $r(\text{S}-\text{O}) = 1.42 \pm 0.02 \text{ \AA}$, $r(\text{C}-\text{C}) = 1.586 \pm 0.01 \text{ \AA}$, $r(\text{C}-\text{S}) = 1.76 \pm 0.02 \text{ \AA}$, and $\angle\text{OSO} = 124 \pm 2^\circ$, with the assumption that the C-H distance and $\angle\text{HCH}$ and $\angle\text{H}_2\text{CC}$ angles are the same as those of ethylene sulfide. From the measurements of Stark shifts, the dipole moment was determined to be $4.47 \pm 0.02 \text{ D}$. (auth)

45292 SOLVENT DEUTERIUM ISOTOPE EFFECTS IN ACID-BASE REACTIONS. VII. SEPARATION OF THE OVERALL ISOTOPE EFFECT IN THE DISSOCIATION OF PICRIC ACID INTO TRANSFER AND ISOTOPE EXCHANGE EFFECTS. Salomaa, Pentti (Univ. of Turku, Finland). Suomen. Kem.; 45: No. 5-6, 149-54(1972).

Changes in standard free energies were measured for the transfer of protopic acid from benzene to light water, the transfer of deuteriopic acid from benzene to heavy water, and the transfer of 1, 3, 5-trinitrobenzene from light to heavy water. On the basis of two assumptions, the validities of which were tested by comparison with experimental data, the results allow calculation of the relative acid strengths of picric acid in light and heavy water. When the present results were combined with other thermodynamic data, the overall isotope effect on the pK value of picric acid can be divided into its constituent parts, namely, effect involved in the transfer of the light isotope species from light to heavy water and isotope exchange effects associated with replacements of protium by deuterium in these species. The contribution of the former effects is about 20% of the overall effect. (auth) (Finland)

45293 ISOTOPE EFFECT ON FRANCK-CONDON FACTORS AND r-CENTROIDS FOR THE NEW ($^1\Sigma_u^+ - ^1\Sigma_g^+$) SYSTEM OF ISOTOPIC C_2 MOLECULE. Singh, I. D.; Maheshwari, R. C. (Univ. of Gorakhpur, India). Indian J. Pure Appl. Phys.; 9: No. 5, 296-7(May 1971).

A partial array of Franck-Condon factors and r-centroids was computed based on a Morse model for the bands of $^{12}\text{C}_2$ and $^{13}\text{C}_2$ molecules of astrophysical importance, observed and analyzed by G. Herzberg and A. Lagerqvist [Can. J. Phys., 46 (1968), 2363]. The ratio of Franck-Condon factors for $^{12}\text{C}_2$ and $^{13}\text{C}_2$ molecules is greater than unity for the $\Delta\nu = 0$ sequence and decreases with transitions to higher levels in both the ν' and ν'' progressions. The isotopic effect on r-centroids is small but significant. (auth)

45294 TEMPERATURE DEPENDENCE OF THE ISOTOPE EFFECT FOR ELECTROMIGRATION OF RUBIDIUM IONS IN MOLTEN RUBIDIUM NITRATE. Lunden, Arnold; Floberg, Allan; Mattsson, Ronny (Chalmers Inst. of Tech., Goteborg). Z. Naturforsch.; 27a: No. 7, 1135-8(Jul 1972).

The relative difference ($\Delta b/b$) between the internal electromigration mobilities of ^{85}Rb and ^{87}Rb in molten RbNO_3 was measured over the range 355 to 500°C. The mass effect $\mu = (\Delta b/b)/(\Delta m/m)$ has a complicated temperature dependence. Thus, the largest mass effect, $-\mu = 0.061$, was obtained at 445°C, while it is about 0.033 at 350°C and 0.041 at 500°C. A similar temperature dependence was found by Saito et al. for μ_{Na} in pure NaNO_3 , and for both μ_{Rb} and μ_{Na} maxima were found also in nitrate mixtures (in $\text{KNO}_3\text{-RbNO}_3$ and $\text{NaNO}_3\text{-KNO}_3$). (auth)

45295 DECAY TIMES, INTENSITIES, AND THEIR TEMPERATURE VARIATION OF LUMINESCENCE OF AQUEOUS AND DEUTERATED SOLUTIONS OF EUROPIUM ION. Pant, D. D.; Bhatt, B. C.; Sanwal, D. N. (Th. D. S. B. Government Coll., Naini Tal, India). Indian J. Pure Appl. Phys.; 9: No. 5, 324-6 (May 1971).

Because of the contradicting nature of the results reported by J. L. Kropp and M. W. Windsor [J. chem. Phys., 39 (1963), 2769; 42 (1965), 1599] and P. K. Gallagher [J. chem. Phys., 43 (1965), 1742] in their studies on luminescence of Eu^{3+} ions in water and heavy water solutions, a careful study of the decay times and relative intensities was made. The results of the present study are in disagreement with the results reported by Gallagher. The temperature variation of decay times and relative intensities in the range 80 to 300°K shows a λ -type behavior for the solutions in $\text{D}_2\text{O-H}_2\text{O}$ mixtures. The temperature variation characteristics are explained on the basis of the diffusion of H^+ and OH^- ions. The dependence of decay times and intensities on H_2O concentration indicates the Stern-Volmer mechanism of quenching. The various processes involved in the mechanism of quenching are discussed. (auth)

45296 ESR EVIDENCE FOR THE INFLUENCE OF ETHANOL ON THE HYDRATED COMPLEXES OF COPPER IONS. Znamirovski, V.; Cozar, O. (Univ., Cluj). Acta Phys. Pol.; A42: No. 1, 3-7(Jul 1972).

An ESR study was performed on the steric effect of ethanol

molecules on the first hydration shell of copper ions. The resolution of the hyperfine structure increases with the ethanol concentration. An H-D isotope effect, which appears in the hyperfine structure, is mentioned and is explained by the preferential association of the ethanol with H_2O . (auth)

45297 NUCLEAR MAGNETIC RELAXATION STUDY OF FORMIC ACID. ISOTOPE EFFECT AND ^{13}C RELAXATION. Ozawa, Hiroshi; Arata, Yoji; Fujiwara, Shizuo (Tokyo Univ.). J. Chem. Phys.; 57: No. 4, 1613-15(15 Aug 1972).

The transient recovery after adiabatic inversion was observed separately for the magnetizations of the $^{12}\text{C-H}$, $^{13}\text{C-H}$, and O-H protons of formic acid. A large difference was observed between the decay constants for the $^{12}\text{C-H}$ and $^{13}\text{C-H}$ protons. It was concluded that this difference is due to (1) flip of the ^{13}C spin resulting in an exchange of the two $^{13}\text{C-H}$ proton signals, and (2) intramolecular dipolar coupling between the ^1H and ^{13}C spins. The above results were used along with the nuclear Overhauser effect data to estimate the ^{13}C spin-lattice relaxation time. It was also shown that the difference in relaxation times between the $^{12}\text{C-H}$ and $^{13}\text{C-H}$ protons is primarily due to the intramolecular dipolar coupling between the ^1H and ^{13}C spins. (auth)

45298 QUENCHING OF INFRARED CHEMILUMINESCENCE: RATES OF ENERGY TRANSFER FROM HF ($\nu \leq 5$) TO CO_2 AND HF, AND FROM DF ($\nu \leq 3$) TO CO_2 AND HF. Airey, J. R. (Naval Research Lab., Washington, D. C.); Smith, I. W. M. J. Chem. Phys.; 57: No. 4, 1669-76(15 Aug 1972).

The intensity of HF (and DF) infrared chemiluminescence from vibrationally excited molecules produced in the chain reaction $\text{H} + \text{F}_2 \rightarrow \text{HF} + \text{F}$; $\text{F} + \text{H}_2 \rightarrow \text{HF} + \text{H}$ (and its deuterium analogue) was studied as a function of the concentration of added CO_2 and HF. Using a steady-state analysis, rate coefficients were determined for transfer of vibrational energy from HF ($\nu \leq 5$) and DF ($\nu \leq 3$) to (vibrationally cold) CO_2 and HF. The results are discussed in terms of possible mechanisms for energy transfer and the importance of the processes in chemical lasers. (auth)

45299 APPLICATION OF PERTURBATION THEORY FOR THE CALCULATION OF MOLECULAR CONSTANTS DUE TO SMALL MASS CHANGES AND ON THE DETERMINATION OF FORCE CONSTANTS FROM VERY HEAVY ISOTOPE SUBSTITUTION. Mueller, A.; Schmidt, K. H.; Mohan, N. (Univ., Dortmund, Ger.). J. Chem. Phys.; 57: No. 4, 1752-7(15 Aug 1972).

The first-order perturbation theory was used to develop the expressions for studying the mass effect on Coriolis coupling constants in isotopically substituted molecules. The corresponding treatment reported by Wilson, Decius, and Cross for the calculation of isotopic frequency shifts has been applied to a large number of molecules with heavy and very heavy atom isotopic substitution, for which the experimental data were published in recent years. The use of a simple method (L matrix approximation) for the calculation of isotope shifts in molecules with small mass coupling, from the knowledge of the geometrical parameters only, is suggested. The results obtained using the perturbation theory for the calculation of $\Delta\nu$ and $\Delta\zeta$ are found to be quite encouraging. An outline of the utility of data on very heavy isotopic substitution for the determination of force constants is given with illustrative examples. (auth)

45300 ISOTOPE EFFECTS IN THE BARTON REACTION. Ferguson, Robert Nicholas. Baltimore; Johns Hopkins Univ. (1971). 346p. University Microfilms Order No. 72-16,912. Thesis.

The primary deuterium isotope effect for hydrogen abstraction in the photolysis of the nitrites of the epimeric alcohols at carbon-11 in apollane (1,4,4,7-tetramethyl tricyclo[5.3.1.0^{2,6}]undecane) was determined. Both of these nitrites give the same oximino alcohol on photolysis. The isotope effect served both as a probe of the transition state for hydrogen abstraction in this photolysis and as a method of comparison of the alkoxy radicals derived from each of the epimeric nitrites. A nondeuterated olefin alcohol was also prepared and reduced with deuterio dimide to give a 2,3 α -dideuterated alcohol. Photolysis of this nitrite gave a surprising secondary isotope effect of 1.7. The data indicate that most of this effect is due to only one (3 α) of the two deuteriums. The corresponding epimeric tertiary alcohols structures 10 and 11 were prepared and these two nitrites upon photolysis gave the same photo oximino alcohol. The corresponding β -deuterated tertiary nitrites of these alcohols were prepared and photolyzed and the isotope effect was found to be ca. 1.1 for each epimer. This value is much lower than that found for the secondary apollanols. The results with the secondary and tertiary systems indicate that the epimerized alkoxy radical has a very similar selectivity to the alkoxy radical generated immediately upon photolysis, and that this is not an "activated" alkoxy radical. The marked difference in primary isotope effects between the secondary and tertiary systems leads to the conclusion that considerable variation is possible in the

exact geometry and extent of hydrogen transfer in the transition state for intramolecular alkoxy radical hydrogen abstraction. (Diss. Abstr. Int., B)

45301 ISOTOPE EFFECTS UPON KINETIC ENERGY RELEASE IN METASTABLE ION FRAGMENTATIONS. Bertrand, M.; Beynon, J. H.; Cooks, R. G. (Purdue Univ., Lafayette, Ind.), *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 346-50 (Aug 1972).

Values of T_H/T_D , where T is the kinetic energy released in unimolecular ionic reactions, were measured for H^+ loss from a number of metastable molecular ions. It is suggested that H^+ loss should give $T_H/T_D \approx 1$ and only if special factors intervene does this value deviate appreciably from unity. Experimental data are in accord with this postulate. (M.C.G.)

Isotope Exchange and Isotope Separation

45302 (COO-1222-55) TECHNICAL PROGRESS REPORT ON FUNDAMENTAL STUDIES OF SEPARATION PROCESSES. Rogers, L. B. (Purdue Univ., Lafayette, Ind. Dept. of Chemistry), *Jul 1972, Contract AT(11-1)-1222, 19p. Dep. NTIS.*

Progress was made in three different major areas. Using high-precision gas chromatography, a mathematical technique, principal component analysis, was shown to be a promising means for analyzing chromatographic peak shapes. A study of electrostatic effects on gas chromatograms obtained using liquid-crystal stationary phases appears to have uncovered a new type of diffusion-limited phenomenon. Studies of micellar and ion-exchange equilibria were shown to be feasible, using gels as column packings, at pressures up to 50,000 psi. Using a relatively inexpensive quadrupole mass spectrometer as a detector for a gas chromatograph, enrichment of the $^{13}C/^{12}C$ ratio in carbon dioxide, after passage through a short column of silica, was confirmed. Currently, the enrichment is being examined under different conditions in the hope of achieving a sufficient separation to allow analyses to be made without using a mass spectrometer. (auth)

45303 IDENTIFICATION OF REACTION INTERMEDIATES BY MICROWAVE SPECTROSCOPY. THE CATALYTIC REACTIONS BETWEEN PROPYLENE AND DEUTERIUM OVER ZINC OXIDE. Naito, Shuichi (Sagami Chemical Research Center, Sagami-hara, Japan); Kondo, Toshikiko; Ichikawa, Masaru; Tamaru, Kenzi. *J. Phys. Chem.*; 76: No. 15, 2184-5 (20 Jul 1972).

The identification of the reaction intermediate in the propylene-deuterium exchange reaction over zinc oxide is reported and the mechanism of the deuterium addition to propylene is also discussed. The deuterium distribution of the product propylene- d_1 suggested that the reaction intermediate of the propylene-deuterium exchange reaction was a π -allyl species. However since propane- d_2 was formed predominantly from the beginning of the reaction, it was concluded that propane- d_2 is formed by the simple addition of deuterium to the double bond of normal propylene or to the propyl group in the surface. It was concluded that hydrogenation and isomerization proceed through different reaction intermediates over zinc oxide. (D.H.M.)

45304 IRON(II) AND IRON(III) ISOTOPE EXCHANGE IN WATER-DIMETHYL SULFOXIDE MIXTURES. Henry, Marian S.; Reynolds, Warren L. (Univ. of Minnesota, Minneapolis). *Int. J. Chem. Kinet.*; 4: No. 1, 37-48 (Jan 1972).

The rate of the isotope exchange reaction between iron(II) and iron(III) perchlorates was measured in a solvent mixture containing a 3:2 mole ratio of water to dimethyl sulfoxide over the temperature range from 25 to $-98^\circ C$. In this temperature range, the reactants can diffuse together faster than they can undergo isotope exchange. The activation enthalpy and entropy for the acid-independent reaction were 6.0 ± 1.2 kcal/mole and -38 ± 17 cal/deg mole, respectively. Below $-22^\circ C$, the acid-dependent exchange reaction did not contribute significantly to the exchange. In liquid media at -112 and $-117^\circ C$ and in a solid glass at $-136^\circ C$, no isotope exchange was observed over the period of a calculated half-life for the reaction. At these temperatures, the rate at which the reactants diffuse together is slower than the calculated rate of isotope exchange. In a solid glass at $-196^\circ C$, no isotope exchange was observed over the period of one week. (auth)

45305 HYDROGEN-DEUTERIUM EXCHANGE KINETICS OF CYTOCHROME C BETWEEN 20 AND $38^\circ C$. Nabedryk-Viala, Eliane; Mestelan-Nonnenmacher, Genevieve; Calvet, Paul (CEN de Saclay, Gif-sur-Yvette, France). *Compt. Rend., Ser. D*; 274: No. 9, 1414-17 (28 Feb 1972). (In French).

The hydrogen-deuterium exchange kinetics of ferricytochrome c (taken from the heart of a horse) were studied at a pH of 7 and at temperatures between 20 and $38^\circ C$. Increasing the temperature increased the rate of exchange between the hydrogen atoms and diminished the number of nonexchanged hydrogen atoms in a 24-hr period from 40 to 28. (France)

45306 STUDY OF EXCHANGE REACTIONS EMPLOYING A COMPLEMENTARY SHOCK TUBE FACILITY. Nika, G. Gerald. New Orleans; Louisiana State Univ. (1971). 210p. University Microfilms Order No. 72-20,803.

Thesis.

A complementary shock tube facility was constructed, employing time-of-flight mass spectrometry and infrared emission spectroscopy as analytical tools. The purpose of this facility is to study homogeneous, high-temperature gas phase exchange reactions, principally the isotopic exchange reaction $H_2 + D_2 = 2 HD$. This system is the prototype for other isotopic exchanges and has received considerable theoretical and experimental attention. Bauer's single-pulse shock tube study of this reaction led to the proposal of a vibrational energy mechanism since molecular or atomic mechanisms cannot explain the experimental data. It is demonstrated that molecular or atomic mechanisms are at variance with experimental observations and the acceptability of the energy mechanism is argued against. Three exchange reactions were studied; $HCl + D_2$, $H_2 + D_2$, and the self-exchange of HD . The first system was investigated using the complementary facility to demonstrate the validity of the time-of-flight-shock tube apparatus. This study covered the temperature range 1700 to $2800^\circ K$ and the results from the time-of-flight and infrared emission shock tubes agree within one standard deviation for the Arrhenius parameters. A non-linear growth of product mole fraction was determined by the two sets of experiments. The combined data is given by the equation $(1 - f_{DCI})[1 + (f_{HCl}/f_{DCI})_{eq}] = \exp(-k[M]^2)$ where, $k = 10^{16.12 \pm 0.30} \exp(-34,340 \pm 3,130/RT)$, $cm^3/mol \text{ sec}^2$. The $H_2 + D_2$ exchange reaction and the self-exchange of HD were studied only on the time-of-flight-shock tube. For the forward reaction the temperature range was 1800 to $3000^\circ K$, for the reverse reaction 1800 to $2800^\circ K$. Deuterium hydride mole fraction exhibited a nonlinear time dependent growth and is given by the rate expression $[1 - 2f_{HD}] = \exp(-k[M]^2)$ where, $k = 10^{16.93 \pm 0.24} \exp(-44,370 \pm 2,510/RT)$, $cm^3/mol \text{ sec}^2$. The data obtained from the HD self-exchange supports a nonlinear time dependent growth of H_2 and D_2 mole fractions. Using an average mole fraction basis the reaction profile for the back reaction is $[1 - 4f_{av}] = \exp(-k^1[M]^2)$ where, $k^1 = 10^{16.67 \pm 0.21} \exp(-40,980 \pm 2,250/RT)$, $cm^3/mol \text{ sec}^2$. All reaction systems are sufficiently described by a quadratic time dependence for product formation. Experimental activation energies are low with respect to values predicted by molecular or atomic mechanisms. The quadratic time dependence and inert gas dependence rule out a four-center mechanism, while a vibrational energy sequence is ruled out by the non-linear time dependence of product formation. An atomic mechanism supports the quadratic time dependence but fails to predict the observed activation energy or total reaction order. (Diss. Abstr. Int., B)

45307 ISOTOPIC SEPARATION BY PHOTOPREDISSOCIATION. Yeung, Edward S.; Moore, C. Bradley (Univ. of California, Berkeley). *Appl. Phys. Lett.*; 21: No. 3, 109-10 (1 Aug 1972).

A monochromatic source may be used to excite individual isotopic lines in a well-resolved absorption spectrum for which the upper state is predissociated. Isotopic enrichment is thus achieved in the dissociation products, which may be stable molecules or may be chemically trapped if they are reactive. As an example, isotopic separation of hydrogen and deuterium is performed by exciting formaldehyde with a frequency-doubled ruby laser. (auth)

Feed Materials Processing

Refer also to abstracts: 45389, 45390, 45405, and 46080.

45308 (DOCKET-701327-1) URANIUM HEXAFLUORIDE FACILITY. Environmental Report. (Allied-Gulf Nuclear Services, Barnwell, S. C.). 30 Jun 1972. 538p. Dep. NTIS.

Information is submitted in support of an application for license to possess and use special nuclear material, source material, and by-products material associated with the conversion of uranyl nitrate to UF_6 . The effects on the environment of the construction and operation of the facility are described. A description is given of the natural resource usage and disposition of radioactive, chemical, and sanitary waste. The site, plant, environmental effects of site preparation and plant construction, environmental effects of plant operation, accidents, alternatives to plant construction and operation, and costs and benefits are discussed. The independently managed environmental monitoring program, the impacts of liquid discharges on Lower Three Runs Creek, transportation requirements associated with UF_6 facilities, and a literature survey concerning environmental effects of several fluoride-associated compounds that are used in the UF_6 facility are given in the Appendixes. (M.C.G.)

Radiation Chemistry

Refer also to abstracts: 45632, 45672, and 46084.

45309 (AE-453) NUCLEAR CHEMISTRY. Progress Report, 1971. (Aktiebolaget Atomenergi, Studsvik (Sweden)). 1972. 21p. Dep. NTIS (U. S. Sales Only).

ANILINE—radiolysis of aqueous solutions of pure and substituted, yield of cyclohexadienyl-type radical from pulse BENZYL (FREE RADICAL)—spectra of, produced by pulse radiolysis of aqueous acid toluene solutions
CARBON—determination in stainless steel and Zircaloy by deuteron and proton activation
ELECTRONS—effects of pulsed, on ice and ice-d₂, optical absorption spectra of transient electrons from
ICE—radiolysis of, optical absorption spectra of transient electrons from pulse
ICE-d₂—radiolysis of, optical absorption spectra of transient electrons from pulse
IODINE-125—purification of trace amounts of, by water vapor distillation in oxidizing atmosphere
OXYGEN—determination in stainless steel and Zircaloy by deuteron and proton activation
STAINLESS STEEL—analysis for carbon and oxygen by deuteron and proton activation
TOLUENE—radiolysis of aqueous acid solutions of, spectra of benzyl radicals from pulse
ZIRCALOY—analysis for carbon and oxygen by deuteron and proton activation (P.C.H.)

45310 (BARC-595) WATER CHEMISTRY STUDIES VII: A GENERAL REVIEW OF WATER CONDITIONS AND WATER TREATMENT IN NUCLEAR REACTORS. Mathur, P. K.; Venkateswarlu, K. S. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 36p. Dep. NTIS (U. S. Sales Only).

A general review is presented covering the problems of water conditions in water-moderated and -cooled nuclear reactors of different types, viz., pool and tank reactors, PWR's and BWR's. The different sources of impurities in the reactor water, its stringent specifications, and the degradation by corrosion, radiolysis, etc., are discussed. The methods and the equipment used in purification, e.g., ion exchange, filtration and degasification, are discussed in detail. Uses of additives to control pH in the coolant, as poisons for emergency shutdown and shim purposes, and for scavenging dissolved oxygen in the coolant are also discussed. (auth)

45311 (BARC-599) CHEMISTRY DIVISION ANNUAL REPORT, 1970-1971. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 163p. Dep. NTIS (U. S. Sales Only).

Research progress is reported in three areas: preparation of pure materials, radiation chemistry, and solid-state studies. References to published papers are given. (M.C.G.)

45312 (COO-2147-2) PREPARATION AND CHARACTERIZATION OF BIOCOMPATIBLE POLYMERS BY RADIATION GRAFTING AND RADIOISOTOPE TECHNIQUE. Technical Progress Report, June 15, 1971-May 15, 1972. Andrade, J. D. (Utah Univ., Salt Lake City. Div. of Materials Science and Engineering). May 1972. Contract AT(11-1)-2147. 25p. Dep. NTIS.

Grafted gel surfaces for medical applications were prepared by γ -induced grafting of hydroxyethyl methacrylate (HEMA) and other hydroxyalkyl methacrylates on inert polymer substrates. Data on wetting and wettability of the grafted gel surfaces were obtained. Protein adsorption by the gel interfaces was studied. (M.C.G.)

45313 (COO-3029-1) ANNUAL PROGRESS REPORT, JUNE 1, 1971-MAY 31, 1972. Turkevich, John (Princeton Univ., N. J.). 1972. Contract AT(11-1)-3029. 6p. Dep. NTIS.

Studies carried out included: radiocolloids, radiation effects on ferrocene adsorbed on glass, and production of hydrogen. (M.C.G.)

45314 (COO-3286-1) RADIATION CHEMISTRY OF NUCLEIC ACIDS. ISOLATION AND CHARACTERIZATION OF THYMINE GLYCOLS. Hahn, Bo-Sup; Wang, Shih Yi (Johns Hopkins Univ., Baltimore, Md. Dept. of Biochemistry). [1971]. Contract AT(11-1)-3286. 7p. Dep. NTIS.

Dihydroxydihydrothymine was identified in irradiated aqueous solutions of thymine and the chemistry of the cis and trans configurations studied. (M.C.G.)

45315 (COO-3286-3) CHEMICAL STUDY OF THE EFFECTS OF RADIATION ON NUCLEIC ACIDS AND RELATED COMPOUNDS. Three Year Progress Report, June 1969-May 1972. Wang, Shih Y. (Johns Hopkins Univ., Baltimore, Md. Dept. of Biochemistry). May 1972. Contract AT(11-1)-3286. 10p. Dep. NTIS.

The chemical natures of radioinduced damage to nucleic acids

and their components were studied. Isolation and characterization of thymine glycols and a thymine hydroperoxide, chemical synthesis of radiation products of thymine, and the effects of thymine hydroperoxide on chemical compounds, transforming DNA, and chromosomes are discussed. (M.C.G.)

45316 (ORO-3909-3) SYNTHESIS, IDENTIFICATION, AND PURIFICATION OF C₆ AND C₇ FLUOROCARBONS FROM RADIIOLYSIS. Final Report. Askew, W. C.; Ward, C. H.; Hardie, F. W.; Parker, B. R.; Maduskar, K. D. (Auburn Univ., Ala. Dept. of Chemical Engineering). Dec 1971. Contract AT(40-1)-3909. 84p. Dep. NTIS.

Synthesis of appreciable amounts of large-molecular-weight fluorocarbons from CF₄ and C₂F₆ by ⁶⁰Co γ rays using flow and batch reactors with and without granular carbon has not been feasible although small amounts of large molecular weight fluorocarbons were made. A new reaction mechanism involving abstraction reactions has been proposed for the radiolysis of saturated fluorocarbons. ¹⁹F NMR spectroscopy can be used for the identification of separated fractions of perfluoroalkanes. Branched chain perfluoroalkane isomers can be easily differentiated from the corresponding n-perfluoroalkane and differentiation among the various isomers is also possible. Semiquantitative analysis of mixtures is feasible if a given component is present in amounts greater than about 10%. Gas chromatographic methods are presented for qualitative and quantitative analysis of perfluoroalkanes. (auth)

45317 ESR STUDIES OF ELECTRON IRRADIATED K₃Ir^{III}(CN)₆ IN KCl SINGLE CRYSTALS. I. LIGAND HYPERFINE STRUCTURE IN Ir^{III}(CN)₆ SPECIES. Vugman, N. V.; Muniz, R. P. A.; Danon, J. (Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro). J. Chem. Phys.; 57: No. 3, 1297-1300(1 Aug 1972).

Electron irradiation of the diamagnetic [Ir(CN)₆]³⁻ ion in a KCl host lattice produces two paramagnetic species. ESR studies reveal that one of the species can be assigned to the pentacoordinated [Ir(CN)₅]³⁻, of approximate C_{4v} symmetry, with 5d⁷ low spin configuration, arising from the trapping of an electron in a d₂ orbital followed by the loss of an axial cyanide. Ligand hyperfine structure with four equivalent and one nonequivalent nitrogens is observed and confirmed by isotopic substitution. The g values and the ligand hyperfine tensors are presented and discussed. Ligand f_s and f_p spins densities are calculated and discussed. (auth)

45318 RADIATION-INDUCED COPOLYMERIZATION OF VINYLPHOSPHONIC DICHLORIDE WITH METHYL METHACRYLATE AND STYRENE. Fujii, Shigetada; Maeda, Katsuaki (Government Industrial Research Inst., Nagoya, Japan). J. Appl. Polym. Sci.; 16: No. 6, 1331-8(Jun 1972).

The γ -induced copolymerization of vinylphosphonic dichloride (VPDC) with methyl methacrylate (MMA) and styrene (St) was studied at 25°C (liquid phase) and -78°C (solid phase). The reaction mechanisms are discussed. The reactivity ratios for the copolymerization of the VPDC-MMA system were determined as follows: 25°C: r₁ = 0.084, r₂ = 6.6 and -78°C: r₁ = 0.081, r₂ = 2.1. The difference between the reactivity between the liquid-phase (25°C) and solid-phase (-78°C) copolymerization is mainly attributable to the r₂ value. The behavior of the liquid-phase copolymerization of the VPDC-St system was anomalous, the r₁ value being negative in the range from 0 to 80 mole-% of VPDC monomer. In the solid-phase (-78°C) copolymerization for the VPDC-St system, the reactivity ratios r₁ and r₂ were 0.097 and 1.6, respectively. The rate of copolymerization (R_p) at 25°C, for both the VPDC-MMA and VPDC-St systems, passes a maximum point at a certain monomer concentration, suggesting that the composition of copolymer is considerably affected by R_p. This phenomenon was interpreted by the assumption that an energy transfer reaction from VPDC monomer to the other vinyl compound can easily occur. (auth)

45319 TRAPPED ELECTRONS PRODUCED IN ETHANOL GLASS AT 4°K. Hase, Hirotomo; Warashina, Tetsuo; Noda, Masato; Namiki, Akira; Higashimura, Takenobu (Kyoto Univ.). J. Chem. Phys.; 57: No. 3, 1039-45(1 Aug 1972).

Optical absorption and ESR measurements were carried out on e⁻ produced in ethanol at 4°K. The same e⁻ yield for γ radiolysis at 4 and 77°K indicates that electrons are stabilized in pre-existing traps at 4°K where molecular dipoles remain unrelaxed. e⁻ prior to solvation can be classified into two groups. One with $\lambda_{\text{max}} = 1500$ nm, W_g = 4 × 10³ cm⁻¹ and $\Delta H_{\text{pp}} = 5.5 \pm 0.5$ G is easily photobleached by the infrared light. The other with broad absorption band in the visible and $\Delta H_{\text{pp}} = 13.5 \pm 1.5$ G is not photobleached by the infrared light. The successive shift of the absorption spectrum to the higher energy side on warming is interpreted by the molecular reorientation mechanism. e⁻ decay is observed during the solvation process, depending on time required for the solvation. The blue shift of the absorption spectrum on reducing the temperature is attributed to contraction of electron traps. (auth)

45320 RADIATION-INDUCED GRAFT COPOLYMERIZATION OF MIXTURES OF STYRENE AND n-BUTYL ACRYLATE ONTO CELLULOSE AND CELLULOSE TRIACETATE. Matsuzaki, Kei; Nakamura, Shigeo; Shindo, Shuichi (Tokyo Univ.). *J. Appl. Polym. Sci.*; 16: No. 6, 1339-55 (Jun 1972).

Mixtures of styrene and n-butyl acrylate of various compositions were grafted onto cellulose and cellulose triacetate fibers pre-irradiated with γ rays at 0°C in air. Monomer reactivity ratios of the grafted copolymers were found to be different from those of the nongrafted copolymers or those of AIBN-initiated copolymers. The active species initiating the graft copolymerization were trapped radicals for cellulose and peroxides for cellulose triacetate. Kinetic investigations of the graft copolymerization of styrene onto preirradiated cellulose triacetate fibers were also carried out, and it was found that the kinetic scheme for radical polymerization is also applicable to graft copolymerization in a heterogeneous system. (auth)

45321 RECOMBINATION PRODUCTS FROM THE RADIOLYSIS OF TRICAPROIN. Letellier, P. R.; Nawar, W. W. (Univ. of Massachusetts, Amherst). *J. Amer. Oil Chem. Soc.*; 49: No. 4, 259-63 (Apr 1972).

A low-molecular-weight triglyceride, tricaproin, was selected to study the higher molecular weight products produced by irradiation. In a previous publication identification of the primary radiolytic products in tricaproin was reported. In the present work, 28 compounds considered to have been produced by the combination of free radicals were identified. Among these are hydrocarbons, ketones, esters, alkanediol diesters, and glyceryl ether diesters. Reaction mechanisms for the production of these compounds are discussed. (auth)

45322 COMPARISON OF THE RADIOLYSIS OF LIQUID BROMOTRIFLUOROMETHANE WITH THE RADIOLYSIS OF LIQUID TRIFLUOROIODOMETHANE. McAlpine, I.; Sutcliffe, H. (Univ. of Salford, Eng.). *J. Phys. Chem.*; 76: No. 15, 2070-1 (20 Jul 1972).

The radiolysis of bromotrifluoromethane in the liquid phase at room temperature was examined. The products and 100 eV yields are: tetrafluoromethane, 0.62; hexafluoroethane, 0.25; dibromodifluoromethane 0.98; and bromine, 0.22. In the presence of iodine the products and some 100-eV yields are: tetrafluoromethane, 0.05; hexafluoroethane, 0.11; trifluoroiodomethane, 3.54; dibromodifluoromethane, and probably bromodifluoroiodomethane. These results are compared with those obtained previously for liquid trifluoroiodomethane. (auth)

45323 ELECTRON SPIN RESONANCE STUDY OF THE FAR-ULTRAVIOLET PHOTOLYSIS OF SOME INORGANIC AND ORGANIC COMPOUNDS. Kuwata, Keiji; Kotake, Yashige; Inada, Koji; Ono, Mitsuo (Osaka Univ.). *J. Phys. Chem.*; 76: No. 15, 2061-9 (20 Jul 1972).

Intermediates in the solid-phase photolysis of alkanes, alcohols, ethers, water, and hydrogen peroxide were investigated at 20 and 77°K. Free-radical intermediates in the vapor-phase photolysis were also examined by a quenching technique at low temperature. In alkanes, alcohols, and ethers formation of ionic species is found when the energy of the far-uv light is higher than the ionization thresholds of these compounds. Also, photolytic conversion of the ionic species into secondary free radicals was examined. Photolytic conversion of the primary free radicals into secondary radicals by successive irradiations with uv light was also studied. In ice and solid hydrogen peroxide photolyzed at low temperature, a singlet spectrum with a line width of ca 13 G was observed. In connection with this system, some other systems were also studied, leading to the assignment of this signal to the HO_2 radical. (auth)

45324 PROTON ENDOR OF γ -IRRADIATED Y-TYPE ZEOLITES. Vedrine, Jacques C.; Hyde, James S.; Leniart, Daniel S. (Varian Associates, Palo Alto, Calif.). *J. Phys. Chem.*; 76: No. 15, 2087-97 (20 Jul 1972).

The initial application of electron-nuclear double resonance (ENDOR) to zeolites is presented and is concerned to a large degree with methodology. Hydrogen atoms are trapped during γ irradiation and the interactions between the atoms and surrounding protons are measured by ENDOR. Motion of the hydrogen atom modulates the dipolar interaction, causing nuclear relaxation of the protons which determines the nature of the ENDOR response. Well-resolved lines have been observed in H-Y-type zeolites. In this system it was determined that the hydrogen atom is trapped in the center of the sodalite cage, and interactions with four different kinds of protons have been found. Sample treatment cycles found to be reversible using other methods were found to be irreversible using ENDOR spectroscopy. (auth)

45325 RADIATION-INDUCED POLYMERIZATION. I. A PYROTECHNIC BINDER. Biggs, William T. (Naval Ammunition

Depot, Crane, Ind.): Parrish, C. F. *J. Appl. Polym. Sci.*; 16: No. 7, 1779-89 (Jul 1972).

In the past, binder systems had to be limited to systems where the catalytically initiated polymerization exotherm was low. With radiation-initiated polymerization, this is not a problem, and the number of monomers which can now be investigated is unlimited. Based on the styrene-benzoyl peroxide system, addition of peroxide sensitizers significantly decrease the radiation dose required for 100% conversion. (auth)

45326 I. RADIATION CHEMISTRY OF n-BUTYROPHENONE IN BENZENE SOLUTION. II. RADIATION-INDUCED ALKYLATION OF CAFFEINE BY AMINES. Stankunas, Alexander Robert. (Riverside, Calif.; Univ. of California (1971)). 140p. University Microfilms Order No. 72-17,048.

Thesis.

The irradiation of benzene solutions of n-butyrophenone leads to the formation of acetophenone and ethylene via a type II process. The G value for this reaction (molecules of product per 100 eV) was found to vary inversely with the dose. It is proposed that the observed decrease in G value is primarily due to the presence of biphenyl, which is produced by irradiation of solvent. An analysis of the rates of various reactions in the system indicate that biphenyl is responsible for at least 80% of the quenching activity. The reactions leading to acetophenone are shown to be strongly dependent on the dose rate. The ethylene produced in the reaction is shown to be unimportant as a quencher of excited butyrophenone. A new reaction, leading to the alkylation of caffeine by amines is reported. The relative reactivity of various alkyl amines under the influence of uv or gamma irradiation is given. Spectral techniques were employed to investigate the possibility of complex formation between the purine and the amines, with generally negative results. Based on data from quenching studies and product structure, a mechanism for the reaction is proposed. Abstraction of a hydrogen atom from the amine by a triplet excited caffeine is followed by radical addition at the 8-position of the caffeine. Loss of ammonia, and final rearrangement to product, follow. In addition, a new radiation-induced reaction between caffeine and acetonitrile to produce a methyl-cyano caffeine is reported. (Diss. Abstr. Int., B)

45327 INVESTIGATION OF THE RADIATION POLYMERIZATION OF METHYL METHACRYLATE-KAOLIN CLAY COMPOSITES. Beeson, James Jackson. Rolla, Mo.; Univ. of Missouri (1972). 156p. University Microfilms Order No. 72-18,182.

Thesis.

A kinetic and characterization study was made on the effects of gamma rays (^{60}Co) on the polymerization of a 50-50 weight mixture methyl methacrylate (MMA)-kaolin clay system. The effect of dose rate (7.35 to 24.9 rads per second), temperature (25 to 75°C), and total dose on the percentage conversion of monomer to polymer was studied. The rate of formation of polymer at 25°C in the composite system was found to be faster when compared to a bulk MMA system at the same experimental conditions. This acceleration showed that the clay had a catalytic effect on the formation of polymer. The effect decreased as temperature increased. Rate data and apparent activation energies are reported. Two types of polymethylmethacrylate (PMMA) were formed in the composite. One type was called homopolymer and could be removed from the composite by extraction with organic solvents. The other type was called inserted polymer and could only be removed by dissolving the clay matrix with hydrofluoric acid. Characterization studies showed that the homo and inserted polymer exhibited the following characteristics: 1. NMR spectra showed that both polymers had increased isotacticity as compared to bulk polymer; 2. the viscosity average molecular weights of the homopolymer ranged from (0.4×10^4) to (5×10^4) depending upon the dose rate, temperature and total dose received; 3. gel permeation chromatography studies confirmed the increase of molecular weight for the homopolymer and showed the same peak molecular weights and distribution for the inserted polymer; 4. infrared spectroscopy showed no differences in the structure of either type of polymer. (Diss. Abstr. Int., B)

45328 RADIATION-INDUCED FORMATION OF EXCITED STATES OF AROMATIC HYDROCARBONS IN BENZENE AND CYCLOHEXANE. II. YIELDS OF EXCITED SINGLET AND TRIPLET STATE SOLUTE MOLECULES. Dainton, Frederick; Morris, T.; Salmon, G. A.; Thompson, G. F. (Univ. of Leeds, Eng. Cookridge Hospital, Leeds, Eng.). *Proc. Roy. Soc. (London)*, Ser. A; 328: No. 1575, 457-79 (27 Jun 1972).

The concentration dependences of the yield of excited triplet states, $G(^3S^*)$, and the concentration dependences of the relative yield of radiation-induced fluorescence obtained on pulse radiolysis of solutions of naphthalene, anthracene, pyrene and 2,5-diphenyloxazole (PPO) are reported. The yields of excited singlet state solute molecules, $G(^1S^*)$, formed on pulse radiolysis of

naphthalene-cyclohexane and naphthalene-benzene solutions was determined by comparing the intensity of the radiation-induced fluorescence with that obtained on photo-excitation. It is concluded that intersystem crossing from the excited singlet state is an important process in the formation of the high yield of triplet excited states of the solute. Under certain conditions this process accounts for up to 50% of $G(^3S^*)$ in cyclohexane and the entire $G(^3S^*)$ in benzene solutions. (auth)

45329 RADIATION-INDUCED FORMATION OF EXCITED STATES OF AROMATIC HYDROCARBONS IN BENZENE AND CYCLOHEXANE. III. EFFECT OF SINGLET STATE QUENCHERS. Dainton, Frederick; Morrow, T.; Salmon, G. A.; Thompson, G. F. (Univ. of Leeds, Eng. Cookridge Hospital, Leeds, Eng.). *Proc. Roy. Soc. (London), Ser. A*; 328: No. 1575, 481-96 (27 Jun 1972).

The contribution to naphthalene and pyrene triplet state formation of intersystem crossing from the excited singlet state was determined for both cyclohexane and benzene solutions using the singlet state quenchers nitromethane and xenon. In agreement with conclusions reported previously, intersystem crossing plays an important role; under certain conditions up to 50% of the total triplet yield in naphthalene-cyclohexane solutions has this origin. The extinction coefficient for naphthalene triplet absorption in cyclohexane at $\lambda_{\max} = 412.5$ nm is $20,000 \pm 5,000$ l mol⁻¹ cm⁻¹. (auth)

45330 RADIATION-INDUCED FORMATION OF EXCITED STATES OF AROMATIC HYDROCARBONS IN CYCLOHEXANE. IV. EFFECT OF ELECTRON SCAVENGERS. Dainton, Frederick; May, R.; Morrow, T.; Salmon, G. A.; Thompson, G. F. (Univ. of Leeds, Eng. Cookridge Hospital, Leeds, Eng.). *Proc. Roy. Soc. (London), Ser. A*; 328: No. 1575, 497-513 (27 Jun 1972).

The effects of the electron scavengers nitrous oxide, nitromethane, carbon dioxide, sulphur hexafluoride, and chloroacetic acid on the yields of naphthalene excited states observed in pulse radiolysis of deaerated naphthalene-cyclohexane solutions were investigated. The excited-state yields were reduced by all the solutes examined although they showed marked differences in their effects on the formation of singlet and triplet excited naphthalene. Of the various competing electron scavengers investigated only the results of the nitromethane system at $[CH_3NO_2] > 10^{-2}$ mol/l could be explained solely in terms of a simple competition for free and geminate ions by the two solutes. However, it was found that the nitrous oxide and carbon dioxide systems could be rationalized on a similar model by taking account of possible secondary ion reactions and/or excited state formation as a result of ion recombination between $Naph^+$ and the negative ion of the competing scavenger. The secondary process occurring in the sulphur hexafluoride and chloroacetic systems are more complex and an empirical fit of the data was not possible. (auth)

45331 RADIATION-INDUCED FORMATION OF EXCITED STATES OF AROMATIC HYDROCARBONS IN BENZENE AND CYCLOHEXANE. I. RADIATION INDUCED FLUORESCENCE FROM SOLUTIONS OF AROMATIC HYDROCARBONS IN CYCLOHEXANE. Frankevich, E. L.; Morrow, T.; Salmon, G. A. (Univ. of Leeds, Eng. Cookridge Hospital, Leeds, Eng.). *Proc. Roy. Soc. (London), Ser. A*; 328: No. 1575, 445-56 (27 Jun 1972).

The nature of radiation-induced fluorescence from solutions of anthracene, naphthalene, and pyrene in cyclohexane was investigated. Evidence is presented for the formation of excited singlet solute molecules via charge recombination processes. Rate constants, measured by the pulse radiolysis technique, are reported for the quenching of excited singlet state molecules of naphthalene and pyrene by oxygen, xenon, iodine, 1,3-cyclohexadiene, nitrous oxide, sulphur hexafluoride and nitromethane. (auth)

45332 RADIOLYSES OF MIXTURES OF AMMONIA AND HYDROCARBONS. II. FORMATION OF HYDROGEN FROM MIXTURES OF AMMONIA AND OLEFINS (ETHYLENE, PROPYLENE, AND BUTADIENE). Hamanoue, Kumao; Okamoto, Jiro; Tokunaga, Okihiro; Danno, Akibumi (Japan Atomic Energy Research Inst., Tokai). *Bull. Chem. Soc. Jap.*; 45: No. 5, 1306-10 (May 1971).

Gamma radiolyses of gaseous mixtures of ammonia and olefins (ethylene, propylene, and butadiene) were investigated over the whole composition range of mixtures, and consideration was given on the primary processes of hydrogen formation. From the extrapolated values of $G(H_2)$ obtained in the γ -radiolyses of the mixture of ammonia and olefins, it was found that the non-scavengeable $G(H_2)$ of ammonia, ethylene, propylene and butadiene were 0.68-0.72, 1.26, 1.06, and 0.33, respectively. Gas-phase radiolyses of ammonia-perdeuterated olefin mixtures were also investigated. From kinetic calculation on the data of isotopic mixtures of hydrogen, the following results were obtained; (1) both thermal and non-scavengeable hydrogen molecules are produced by the γ -radiolysis of ammonia, and the non-scavengeable hydrogen mole-

cules are produced for the most part by a unimolecular process, (2) almost all the hydrogen molecules produced by the γ -radiolyses of olefins (ethylene, propylene, and butadiene) are non-scavengeable; a unimolecular process plays an important role in the case of ethylene, and in other cases a bimolecular process. $G(H_2)$ of the butadiene radiolysis is very small compared with that of other olefins and with the calculated value of butadiene, i.e., $G(H_2) < 1$. The discrepancies can be interpreted in terms of the addition reaction of the "hot" hydrogen radicals to butadiene to form t-2-butene. (auth)

45333 THEORETICAL CONSIDERATION OF THE ELECTRON-SCAVENGING PROCESS IN LIQUID HYDROCARBONS. III. INTEGRAL TRANSFORMS OF THE DISTRIBUTION FUNCTIONS. Yamazaki, Hideo; Shinsaka, Kyoji (Tokyo Inst. of Tech.). *Bull. Chem. Soc. Jap.*; 45: No. 5, 1335-8 (May 1972).

The relations of integral transforms between various distribution functions are discussed using the potential control approximation ignoring the diffusion term in the Smoluchowski equation; the initial spatial distribution of the electron-ion pair or the initial distribution of electrons from the central ion core in a spur, $\chi(r,0)$; the relative G-value of electron recombination, $G(k_e C_e)$, as a function of the concentration of the scavenger and the rate constant; the time dependence of the electron concentration, $N(t)$, after pulse irradiation; and the relaxation time spectrum, $K(n)$, of the decay of the electron concentration after pulse irradiation or of the weight function, $K(n)$, in the initial exponential distribution of thermalized electrons in a spur are shown to be connected by the square diagram of Laplace transforms:

$$\begin{array}{ccc} & K(n) & \xrightarrow{\text{Laplace Transform}} \chi(r,0) \\ \text{Laplace Transform} \downarrow & & \downarrow \text{Laplace Transform} \\ N(t) & \xrightarrow{\text{Laplace Transform}} & G(k_e C_e) \end{array}$$

By the use of this relation, the various distribution functions are evaluated from the empirical formulas of $G(k_e C_e)$ proposed by Schuler et al. and by Hummel. (auth)

45334 ACTIVE SPECIES OF γ -IRRADIATED VINYL ACETATE AND THEIR CHANGES ON IRRADIATION OF VISIBLE LIGHT. Hirai, Hidefumi; Fujiwara, Masato (Tokyo Univ.). *Nippon Kagaku Kaishi*; No. 5, 968-74 (May 1972). (In Japanese).

Free radicals produced by the γ -irradiation of vinyl acetate in the solid state were studied with electron spin resonance (ESR) and the changes of these radicals on the visible light irradiation were also examined. An oriented solid of vinyl acetate was prepared for ESR measurement by the solidification of vinyl acetate around an aluminum rod at -196°C or -117°C and with subsequent γ -irradiation (dose; 8×10^6 rad) at -196°C . The oriented crystalline solid prepared at -117°C gives an ESR spectrum consisting of a singlet and additional absorptions in the dark at -196°C . After irradiation by visible light for 10 minutes, ESR spectra of the oriented crystalline solid exhibit a distinct hyperfine structure and a definite anisotropy on the rotation of the solid in the dc magnetic field. These spectra consist of a quartet ($A_1 = 12.5$ G), an anisotropic triplet and an equivalent four lines ($A_1 = 103$ G, $A_2 = 16$ G), corresponding to the radicals, $\cdot CH_3COO\cdot$, $\cdot CH_2COOCH = CH_2$, and $CH_2 = CH$, respectively. Prolonged irradiation with visible light for 300 minutes yields the apparent five lines, consisting of a pair of triplet ($A_1 = 10$ G, $A_2 = 20$ G) and the above-mentioned triplet corresponding to the radicals $\cdot CH_2CHO$ and $\cdot CH_2COOCH = CH_2$, respectively. The spectrum of the γ -irradiated glassy solid prepared at -196°C consists of strong four lines, which is confirmed to be isotropy, and of a weak doublet split by 120 G, corresponding to the vinyl radical. Irradiation of the glassy solid with visible light for 30 minutes gives a sharp quartet ($A_1 = 23$ G) assigned to $CH_3\cdot$ produced by the photolysis of $CH_3COO\cdot$. A broad triplet ($A_1 = 22$ G) becomes observable on elevation of the temperature to -160°C and remains up to -100°C . This triplet is assigned to $\cdot CH_2COOCH = CH_2$. In conclusion, both the crystalline and the glassy solids produce the same radicals on γ -irradiation. (auth)

45335 SYNTHESIS OF OXYGEN-CONTAINING HALOGENATED COMPOUNDS. II. γ -RAY INDUCED REACTION OF CYCLOHEXANONE IN THE PRESENCE OF CHLORINATED HYDROCARBONS. Yumoto, Takaari (Government Industrial Research Inst., Nagoya, Japan); Matsuda, Tatsuo. *Nippon Kagaku Kaishi*; No. 5, 940-4 (May 1972). (In Japanese).

Cyclohexanone was irradiated with γ rays in the presence of some chlorinated hydrocarbons. The main products were confirmed to be 2-(1-cyclohexenyl)cyclohexanone [1] and 2,6-di(1-cyclohexenyl)cyclohexanone [2]. Any expected condensation product between cyclohexanone and chlorinated hydrocarbons was not detected. The rates of formation of [1] and [2] were found to be nearly proportional to the dose rate. A maximum yield of 47.5% of

[1] was obtained in the presence of 20% moles of tetrachloroethylene. The effect of the coexistence of chlorinated hydrocarbons on the yields of [1] and [2] decreased in the order: tetrachloroethylene > trichloroethylene > cis-1,2-dichloroethylene, carbon tetrachloride > chloroform > methylene chloride > 1,1-dichloroethane. The reaction mechanism for the formation of [1] and [2] was discussed. (auth)

45336 INDUCING DYEABILITY IN POLYPROPYLENE BY RADIATION INDUCED GRAFTING. DasGupta, S. (to Atomic Energy of Canada Ltd.). Canadian Patent 892,730. 8 Feb 1972.

A process for improving the dyeability of polypropylene, particularly polypropylene fibers and fabrics, is described. The polypropylene is impregnated with polymerizable unsaturated organic monomer containing hydrophilic groups, either individually or in admixture, or as mixtures with styrene. The monomer(s) are in solution in a substantially anhydrous primary alcohol, preferably methanol or ethanol. The monomer-impregnated polypropylene is subjected to high energy ionizing radiation, such as ^{60}Co gamma radiation, and the monomers are thereby graft copolymerized with the polypropylene. Such grafted polypropylene products can be dyed very easily. The dyed fabrics are soft to handle, have dye stability, light- and washfastness, moisture retention, resistance to static charge, and retain all of the desirable properties of ungrafted polypropylene. (Canada)

45337 1,6-GLUCODIALDOSE AS ONE OF THE RADIOLYSIS PRODUCTS OF D-GLUCOSE-6-PHOSPHATE. Kochetkov, N. K.; Kudryashov, L. I.; Chlenov, M. A.; Grineva, L. P. (Inst. of Organic Chemistry, Moscow). Dokl. Akad. Nauk SSSR; 202: No. 4, 847-9 (1 Feb 1972). (In Russian).

The radiolysis of D-glucose-6-phosphate was studied by subjecting a 10^{-2} M aqueous solution to ^{60}Co gamma radiation at a rate of 6.5×10^{16} eV/ml · sec. It was postulated that the radiation caused a rupture of the phosphate bond, and to a significant degree carbon-6 is oxidized. The radiolysis was conducted under different gaseous atmospheres (N_2 , N_2O , and O_2). In all cases one of the chief radiolysis products was 1,6-glucodialdose. (K.S.W.)

45338 RADIATION DAMAGE TO POLYMER FIBER STRUCTURE AS INVESTIGATED BY THE METHOD OF SMALL-ANGLE X-RAY SCATTERING. Plavnik, G. M.; Goryaev, V. M.; Spitsyn, Vikt. I.; Ryabchikova, G. G.; Berestnev, V. A. (Inst. of Physical Chemistry, Moscow). Dokl. Akad. Nauk SSSR; 202: No. 4, 878-81 (1 Feb 1972). (In Russian).

Experiments were conducted on fibers of Kapron cord in air and in vacuum by subjecting them to various forms of ionizing radiation: accelerated electrons at energies of 1 to 5 MeV or gamma radiation from a ^{60}Co source, varying the dose from 0.5 to 3,000 MR. X-ray diffraction patterns were taken with a four-slit small-angle camera, using $\text{CuK}\alpha$ radiation. Significant differences were apparent at large radiation doses: for irradiation in air the intensity of reflection dropped by a factor of 4 for a 500 MR dose relative to the original sample; irradiation in a vacuum yielded an analogous decrease in the intensity only at a dose of 1,500 MR. It was concluded that the method of small-angle scattering of x rays is extremely sensitive to processes evolving in polymeric fibers subjected to radiation, and allows quantitative evaluation of the extent of these processes. (K.S.W.)

45339 METHOD FOR THE POLYMERIZATION AND/OR THE CROSS-LINKING OF ACRYLIC POLYMERS OR COPOLYMERS. Blin, Marie-Francoise; Gaussens, Gilbert (to Commissariat a l'Energie Atomique). French Patent 2,076,863. 20 Sep 1971. Filed 30 Jan 1970. (In French).

A method for obtaining radiation-induced polymerization and/or cross-linking of acrylic polymers in the presence of air was developed, for all thicknesses of material. The process consists of preparing an acrylic saturated resin dissolved in vinyl monomers, and the subsequent radiation-induced shaping and cross-linking of the resin in the presence of air. (France)

45340 ABSORPTION SPECTROSCOPY OF TRANSIENT SPECIES. Herkstroeter, William G. (Eastman Kodak Co., Rochester, N. Y.). pp 521-76 of Physical Methods of Chemistry. Pt. IIIB. /Weissberger, Arnold (ed.). New York; John Wiley and Sons, Inc. (1972).

Absorption spectroscopy of transient species is reviewed including absorption of stabilized transients utilizing conventional spectroscopic techniques, flash photolysis, pulse radiolysis, optimum solute concentrations with lateral excitation, determination of transient extinction coefficients, and transient kinetics. (D.H.M.)

45341 ELECTRON SPIN RESONANCE STUDY OF RADICALS FORMED BY HIGH-ENERGY IRRADIATION OF K_2PdCl_4 AND K_2PtCl_4 . Krigas, Thomas Michael. East Lansing, Mich;

Michigan State Univ. (1971). 138p. University Microfilms Order No. 72-16,460.

Thesis.

All known square-planar complexes of Pt(II) and Pd(II) are diamagnetic with d^8 electron configurations and therefore cannot be studied by electron spin resonance spectroscopy. Several new paramagnetic species were produced by high-energy irradiation of solid diamagnetic compounds to give Pt and Pd containing radicals in which one-electron oxidation or reduction has occurred. The structures of these species were obtained from electron spin resonance studies. The X-band electron spin resonance spectra of single crystals of K_2PdCl_4 and K_2PtCl_4 that were irradiated by γ rays from a ^{60}Co source or by 1-MeV electrons were studied at temperatures between 77 and 296°K with the primary purpose of elucidating the structure and bonding in species showing unusual oxidation states of palladium and platinum. Irradiation of $\text{K}_2\text{Pd(II)Cl}_4$ produced two identifiable paramagnetic radicals: one is shown to be $(\text{Pd(I)Cl}_4)^{3-}$ with a $4d^9$ electron configuration and the unpaired electron in a $d_{x^2-y^2}$ orbital; the second radical is believed to be $(\text{Pd(III)Cl}_4)^{2-}$ with the unpaired electron in a $4d_{z^2}$ platinum atomic orbital. Crystals of $\text{K}_2\text{Pt(II)Cl}_4$ irradiated at 77°K show two groups of electron spin resonance lines: one belongs to a radical that shows hyperfine interaction with two equivalent platinum nuclei; the second arises from a radical showing hyperfine interaction with one platinum and three chlorine nuclei. It is suggested that these species are $(\text{Pt(II)Cl}_4)^-$, $(\text{Pt(III)Cl}_4)^{3-}$ and $(\text{Pt(I)Cl}_4)^{2-}$, respectively, where the metal electron configurations are $5d^{10}$ and $5d^9$. On warming to 125°K some of the dimeric radical is converted to a new species whose spectra are consistent with those predicted for $(\text{Pt(III)Cl}_4)^{2-}$. Thus, each identifiable radical contains platinum or palladium in which the original diamagnetic, low spin, d^8 configuration of the metal ion M(II) ($\text{M} = \text{Pt or Pd}$) was oxidized or reduced to form paramagnetic species with the metal in the unusual oxidation states M(III) or M(I) . Energy level schemes for each radical are proposed based upon the electron spin resonance spectra. The nature of the metal-chlorine bonds in the new species is discussed and the extent of covalency is estimated by using a simplified molecular orbital picture. (Diss. Abstr. Int., B)

45342 KINETIC ISOTOPE EFFECTS IN FREE RADICAL CHEMISTRY. Kneipp, Kenneth George. Baton Rouge, La.; Louisiana State Univ. (1971). 226p. University Microfilms Order No. 72-17,778.

Thesis.

Kinetic isotope effects for hydrogen atom abstraction from mercaptans by free radical species were investigated. A variety of structurally different radicals were generated by thermolytic, photochemical, and radiolytic techniques and allowed to abstract hydrogen, deuterium, or tritium atoms from the S-H position of a mercaptan molecule. The rate ratio, or isotope effect, k_H/k_D or k_H/k_T , was measured in all cases as a function of temperature. A useful relationship was developed whereby the magnitude of the observed isotope effect, or isotopic selectivity, for a given radical may be correlated to its stability or reactivity. Such a pattern of isotope effects for abstraction should prove useful as a diagnostic test for radical species whose reactivity may otherwise be difficult to assess. Secondary deuterium isotope effects on thermolysis of perester initiators were measured in order to elucidate the mechanism of decomposition of these free radical precursors. The results are shown to be in good agreement with similar secondary isotope effect measurements reported in the literature and the mechanistic implications are consistent with results obtained using a variety of investigation techniques. (Diss. Abstr. Int., B)

45343 I. RADIATION DAMAGE IN MALONIC ACID; AN ACYL RADICAL. II. DEUTERON DISTANT ENDOR STUDIES. McCalley, Roderick Canfield. Cambridge, Mass.; Harvard Univ. (1971). 250p. University Microfilms Order No. 72-18,635.

Thesis.

The radical $\text{RCH}_2\dot{\text{C}}=\text{O}$ (where R- is HOOC-) was unexpectedly discovered in x-irradiated malonic acid single crystals. Its most obvious ESR parameters, g tensor (1.9964, 2.0018, 2.0039) and "outer" ^{13}C hyperfine tensor ($498\frac{1}{2}$, 383, $372\frac{1}{2}$ MHz), indicate a $\dot{\text{C}}=\text{O}$ type "g" radical, but are not sufficient to unambiguously distinguish the proposed species from $\cdot\text{COOH}$. An "inner" ^{13}C hyperfine splitting (172, 133, 132 MHz) and the two proton hyperfine tensors (+6, -13, -16 MHz), as well as the geometry of all the tensors, do confirm the $\text{RCH}_2\dot{\text{C}}=\text{O}$ assignment. An analogous species, located at the opposite end of the malonic acid molecule, exhibits a hindered rotation of its $\dot{\text{C}}=\text{O}$ group at room temperature, averaging out 75% of the magnetic anisotropy in the plane $\perp$ to its C-C axis. For these substituted acetyl radicals, the spin distribution and properties (especially the large second ^{13}C interaction and the anomalously low and negative β -proton splittings) are discussed and explained in terms of general "rules" for

spin-delocalization and dipolar coupling. In deuterated malonic acid (d-Mal) about 45 to 50% of the radicals created by a short X-irradiation at 300°K are $\text{RCH}_2\dot{\text{C}}=\text{O}$, compared to only 25 to 30% in h-Mal. For the aging of $\text{RCH}_2\dot{\text{C}}=\text{O}$ at room temperature, the half-time is less than 2 hr in h-Mal and about 1 day in d-Mal. The ENDOR spectra of d-Mal at 4°K include very strong and narrow signals from the "distant" deuterons and naturally abundant ^{13}C (up to 80% and 25%, respectively, of the ESR baseline). Enough geometrical information is available from the splitting tensors to draw a fairly comprehensive picture of the 4°K crystal structure, particularly with respect to the manner in which the room temperature center of symmetry (between the two molecules in the unit cell) seems to be lost in cooling to 4°K. (Diss. Abstr. Int., B)

45344 STUDIES OF CONTINUOUS DIELECTRIC MODELS FOR SOLVATED ELECTRONS. Meglen, Robert Rudolf. Boulder, Colo.; Univ. of Colorado (1971). 200p. University Microfilms Order No. 72-17,286.

Thesis.

The evolution of dielectric models for the solvated electron is described. Early models greatly influenced the basic assumptions of the more recent models. The early particle-in-a-spherical box model was modified by J. Jortner. The Jortner cavity model is discussed. The basic assumptions made in the Jortner model were evaluated. A Self Consistent Field-Continuous Dielectric (SCF-CD) model for the solvated electron is presented. This model does not require the assumption of a cavity. The wave functions and energies are determined self-consistently by a numerical integration of the Schrödinger Equation. This model gives results that are comparable to the cavity model. Three modifications of the SCF-CD model are given. The effect of introducing a cavity into the model was evaluated by performing a set of SCF calculations with a variety of assumed cavity radii. The results of these calculations are compared to the results of the Jortner cavity calculations. A second modification of the SCF-CD model allows a continuously variable solvent density in the neighborhood of the additive electron. The reduction in solvent density is assumed to be proportional to the additive electron density, and it is treated self-consistently. The third modification makes use of the pseudopotential method. A repulsive term is added to the potential energy function to account for the additive solvated electron. The transition energy of the solvated electron in various solvents, the width of the absorption band, the temperature, and pressure dependence of the absorption maximum are discussed. None of the SCF-CD modifications provides a satisfactory explanation of the observed properties of the solvated electron. The variation in transition energy with solvent is difficult to interpret with present models. The width of the absorption band can not be explained by a Rydberg type series of electronic transitions. The most serious shortcoming of these modifications is that they predict the wrong sign for the temperature and pressure dependence of the absorption maximum. It was concluded that these difficulties are not caused by the assumption of a continuous dielectric medium. The difficulty is believed to be due to an incorrect form of the polarization potential. A brief description of some calculations of the onset of metallic conductivity is included in an appendix. This calculation treats the onset of metallic conductivity as a case of the Mott transition. (Diss. Abstr. Int., B)

45345 ROLE OF FLUORIDE ION IN IRRADIATED AQUEOUS SYSTEMS. Bales, Barney L. (California State Univ., Northridge); Kevan, Larry. J. Chem. Phys.; 57: No. 4, 1813-14 (15 Aug 1972).

A mechanism for explaining the role of fluoride ions in irradiated aqueous systems is proposed that led to the conclusion that more e^- is produced in F^- -doped ice because of chemical means rather than physical means; the same number and type of traps are available in ice and F^- -doped ice and are unrelated to the presence of that ion. (D.H.M.)

45346 γ RADIOLYSIS OF GASEOUS CARBON DIOXIDE AT HIGH DENSITIES. Yoshimura, M.; Chosa, M.; Soma, Y.; Nishikawa, M. (Tokyo Univ.). J. Chem. Phys.; 57: No. 4, 1626-31 (15 Aug 1972).

The γ radiolysis of gaseous carbon dioxide was investigated at $50 \pm 1^\circ\text{C}$ over the density range of 0.2 to 0.6 g/ml. At density of around 0.5 g/ml, $\text{G}(\text{CO})$ from pure carbon dioxide showed a very marked increase. Addition of SF_6 at these densities efficiently reduced $\text{G}(\text{CO})$. From this it appears that formation of carbon monoxide results mostly from charge neutralization. Examination of the mechanisms in the radiolysis of carbon dioxide in gas and liquid phase, and a calculation on a simple model indicate that this neutralization is of geminate character. Upon addition of hydrocarbons which react with oxygen atoms as well as the addition of both hydrocarbons and SF_6 , $\text{G}(\text{CO})$ passed

through a maximum and reached the value of 2.2 at higher concentrations of hydrocarbons, independent of density. It was interpreted that (1) the positive ions involved in neutralization are probably clustered CO_2^+ in agreement with the above inference, and that (2) there are at least two species responsible for the rapid re-oxidation of CO. (auth)

45347 ESR STUDY OF THE VINYL-TYPE RADICAL TRAPPED IN A SINGLE CRYSTAL OF MALEIC ACID IRRADIATED AT 77°K. Eda, Bunzo; Iwasaki, Machio (Government Industrial Research Inst., Nagoya, Japan). J. Chem. Phys.; 57: No. 4, 1653-60 (15 Aug 1972).

The substituted vinyl radical, $\text{DOOC}-\dot{\text{C}}\text{H}=\text{CH}$, trapped in an irradiated single crystal of maleic acid- d_2 was studied by ESR to elucidate the geometrical and electronic structure of the vinyl-type radical. The hyperfine tensors for the ^{13}C and β proton couplings were determined to be (+133.8, +75.3, +62.2)G and (+59.5, +57.5, +56.8)G, respectively, in addition to the α proton coupling and g tensors preliminarily reported. All the magnetic tensors determined are consistent with the planar structure of the vinyl-type radical in which the unpaired electron occupies the in-plane sp hybrid orbital. The ^{13}C hyperfine tensor indicates the relatively large spin density in the $\text{C}_{2\text{PO}}$ orbital which is consistent with the small s character of the vinyl-type radical. The relative orientation of the hyperfine tensor of the α proton to that of ^{13}C gives the clear evidence for the bent structure of the radical carbon atom. The dipolar tensor of the α proton coupling has been calculated using the spin distribution and the geometry obtained in our experiments, and the results are in good agreement with the observation in its magnitude as well as its directions. The apparent difference in the α proton coupling between $\text{C}-\text{H}$ π and σ radicals has been discussed in relation to the sign of the isotropic coupling. (auth)

45348 RADICAL ION SATURATION IN SOME SULFUR COMPOUNDS X-IRRADIATED AT 4.2°K. Box, Harold C.; Potter, William R.; Budzinski, Edwin E. (Roswell Park Memorial Inst., Buffalo). Contract AT(30-1)-3558. Radiat. Res.; 51: No. 1, 10-14 (Jul 1972).

Single crystals of dimethylthiourea and dibenzoyl disulfide were exposed to ionizing radiation at 4.2°K. At this temperature positive and negative free radical ions are stable and can be identified from their electron spin resonance (ESR) absorption spectra. The concentration of radical ions was measured as a function of dose and it was found that the concentrations approach limiting values of the order of 10^{18} radicals per gram. This "saturation" effect is attributed to nongeminate recombination of electrons with cations. (auth)

45349 ELECTRON SPIN RESONANCE STUDIES AND INDO MO-CALCULATIONS OF ELECTRON IRRADIATED PEPTIDES. A SINGLE CRYSTAL OF N-ACETYLGLYCINE. Saxeboel, Gunnar; Meloe, Thor B.; Henriksen, Thormod (Univ. of Oslo). Radiat. Res.; 51: No. 1, 31-44 (Jul 1972).

The radicals induced by radiation in single crystals of N-acetyl glycine, grown from both H_2O and D_2O solutions, were studied using both ESR spectroscopy and INDO MO-calculations. At room temperature the radical $\text{H}_3\text{C} \cdot \text{CO} \cdot \text{NH} \cdot \dot{\text{C}}\text{H} \cdot \text{COOH}$ was formed as previously demonstrated. In these experiments the detailed distribution of the unpaired spin density was studied both through the ESR observations and by MO-calculations. The two methods were in satisfactory agreement. It was found that the unpaired electron was largely delocalized and even δ -proton interaction was found. Furthermore, the observation that unpaired spin density can migrate through the peptide group may have some bearing with regard to the processes taking place in irradiated proteins. (auth)

45350 RADIOLYSIS OF AQUEOUS SOLUTIONS OF NICKEL(II) ETHYLENEDIAMINE TETRAACETATE. Bhattacharyya, S. N.; Kundu, K. P. (Saha Inst. of Nuclear Physics, Calcutta). Radiat. Res.; 51: No. 1, 45-55 (Jul 1972).

Neutral aqueous solutions of nickel(II) ethylenediaminetetraacetate (N) were irradiated with x rays both in the presence and in the absence of dissolved oxygen, and the radiolytic degradation of the metal chelate was determined. Hydrogen peroxide and formaldehyde that were produced in the system were also determined. A radiolytic mechanism was proposed and discussed where it was argued that the radiolytic degradation at the ligand is due to OH only. Effect of methanol and formate was also studied and the ratios of the rate constants, $\text{K}(\text{OH} + \text{CH}_3\text{OH})/\text{K}(\text{OH} + \text{N})$ and $\text{K}(\text{OH} + \text{HCOO}^-)/\text{K}(\text{OH} + \text{N})$ were found to be 0.32 and 1.64, respectively. (auth)

45351 EFFECT OF PRESSURE ON THE RATES OF REACTION OF SOLVATED ELECTRONS IN LIQUID ETHANOL. Jha, Kamal N.; Freeman, Gordon R. (Univ. of Alberta, Edmonton). J. Chem. Phys.; 57: No. 4, 1408-14 (15 Aug 1972).

At a γ radiolysis dose of 1×10^{18} eV/g in ethanol at 296°K the hydrogen yields in neutral ethanol were $G(H_2) = 5.0$ at 1 bar and 5.5 at 5.3 kbar; in 1 mM and 1 M sulfuric acid $G(H_2) = 5.7$, independent of pressure. The electron scavengers nitrobenzene, acetone, and naphthalene reduce the hydrogen yields in neutral and acidic ethanol at 1 bar and 5.3 kbar; increasing the pressure greatly reduces the ease of scavenging the free ions but has only a small effect on scavenging efficiency in the spurs. Using as a standard the reaction $e_{sol}^- \rightarrow C_2H_5O_{sol}^- + H$ with $k = 1.2 \times 10^5 \text{ sec}^{-1}$ at 1 bar and $\Delta V^\ddagger = -14.4 \text{ cm}^3/\text{mole}$ averaged between 0 and 5.3 kbar, one obtains the following values for $k(M^{-1} \text{ sec}^{-1})$ and $\Delta V^\ddagger (\text{cm}^3/\text{mole})$: $e_{sol}^- + H^+$, 2.8×10^{10} at zero ionic strength, +4.5; $e_{sol}^- + C_6H_5NO_2$, 1.5×10^{10} , +7.5; $e_{sol}^- + \text{naphthalene}$, 5.0×10^9 , +5.6; $e_{sol}^- + CH_3COCH_3$, 4.2×10^9 , +5.1. The diffusion coefficient of solvated electrons in ethanol appears to correlate with the liquid viscosity; this is different from the behavior of electrons in alkanes. In ethanol $\Delta V^\ddagger_{\text{diffusion}} \approx +7 \text{ cm}^3/\text{mole}$ for e_{sol}^- and $+4 \text{ cm}^3/\text{mole}$ for H_{sol}^+ . The free ion yield in ethanol is $G(e_{sol}^-)_{fi} = 1.7$, independent of pressure from 1 bar to 5.3 kbar; the model of Freeman and Fayadh [J. Chem. Phys. 43: 86 (1965)] is consistent with this behavior and it predicts that G_{fi} in methanol and water is also independent of pressure. (auth)

45352 CROSSLINKED POLYVINYL CHLORIDE OR POLY-VINYLDENE FLUORIDE FILM. Hahn, Ernest A. (to PPG Industries, Inc.). U. S. Patent 3,676,192. 11 Jul 1972. Filed 6 Apr 1970.

Coating compositions comprising poly(vinyl chloride) or poly(vinylidene fluoride) are prepared by mixing a solution of the poly(vinyl chloride) or poly(vinylidene fluoride) in an active solvent with an acrylic monomer and subjecting the mixture to ionizing irradiation. The cured films are hard, stain-resistant, heat-resistant, and mar-resistant. (Official Gazette)

45353 ION-MOLECULE REACTIONS IN AMMONIA AT HIGH PRESSURES. Wincel, Henryk (Inst. of Nuclear Research, Warsaw). J. Mass Spectrom. Ion Phys., 9: No. 3, 267-85 (Aug 1972).

Ion-molecule reactions in gaseous ammonia were examined at pressures up to 2 torr in a high-pressure mass spectrometer. Solvation of ammonium ions as well as other ions in both NH_3 and ND_3 studies was observed. The stability of the solvates $NH_4^+(NH_3)_n$ diminishes with increasing degree of solvation. The effect of the energy of the ionizing electrons on the yields of solvated and other ions was studied. Evidence for the formation of excited states of NH_4^+ , $NH_4^+NH_3$, $NH_4^+(NH_3)_2$, etc., and collisional de-excitation processes was obtained. Only NH_4^+ ions reach thermodynamic equilibrium at the ion-source pressures used in these experiments. (auth)

45354 ELECTRON AND HYDROGEN ATOM ATTACHMENT TO AROMATIC CARBONYL COMPOUNDS IN AQUEOUS SOLUTION. ABSORPTION SPECTRA AND DISSOCIATION CONSTANTS OF KETYL RADICALS. Hayon, E.; Ibatata, T.; Lichtin, N. N.; Simic, M. (Army Natick Labs., Mass. Boston Univ.). J. Phys. Chem., 76: No 15, 2072-8 (20 Jul 1972).

The technique of pulse radiolysis-absorption spectroscopy was used to study the addition of hydrated electrons and H atoms to a number of compounds incorporating both carbonyl and aromatic groups. The absorption spectra of the electron adducts and their oxygen-protonated conjugate acids were determined for acetophenone, 9,10-antraquinone-2-sulfonate, benzamide, benzil, benzoin, benzophenone, and fluorenone. The pK_a values for all of these radicals except those derived from benzoin were evaluated as were the bimolecular decay rates of the anion radicals. Bimolecular decay rates of the protonated electron adducts of acetophenone, 9,10-antraquinone-2-sulfonate, and benzamide were also measured. Specific rates of addition of the hydrated electron to all the substrates for which resultant spectra were determined as well as anthrone, dibenzylketone, 1,4-naphthoquinone-2-sulfonate, 1,2-naphthoquinone-4-sulfonate, and phenylacetone have been found to fall in the range $(1.1-3.6) \times 10^{10} M^{-1} \text{ sec}^{-1}$. Spectra of the H-atom adducts and bimolecular specific rates of decay of these adducts have been determined for acetophenone, 9,10-antraquinone-2-sulfonate, benzamide, benzil, benzophenone, fluorenone, and 1,4-naphthoquinone-2-sulfonate. Systematic aspects of the dependence on structure of the reactivity toward addition of e_{aq}^- and H atoms, of the pK_a 's of protonated electron adducts, and of the various spectra are discussed. (auth)

45355 METHOD OF MODIFYING POLYMERIC MATERIALS, WOOD, AND ARTICLES MADE THEREFROM BY RADIATION-INITIATED GRAFT COPOLYMERIZATION THEREOF. Kabanov, Vitaly Yakovlevich; Voronkov, Nikolai Andreevich; Kochkin, Dmitry Alexandrovich; Spitsyn, Viktor Ivanovich; Kazamanova, Irada Vasilievna (to Institut Fizicheskoi Khimii Akademii

Nauk SSSR). British Patent 1,269,297. 6 Apr 1972. Filed 17 Jul 1970.

A method is presented for the modification of synthetic polymers and wood in order to render them fungus resistant. The method described comprises contacting the material with a trialkylacryloxystannane, and in particular triethylmethacryloxystannane, and solution graft polymerising this onto the material under the action of γ -radiation. Irradiation is carried out until not less than 0.5% by weight of the oxystannane, based on the weight of polymer or wood, has been grafted. The materials so produced are fungus resistant and have long service life, since the grafted material cannot be removed under practically all service conditions. The preferred modes of conducting the process are described. The dosage of γ radiation for polymers is suitably 0.1 to 0.5 MR and the absorbed dose rate 0.3 to 5 R/sec, and for wood 80 to 100 R/sec with an exposure time of 30 to 50 hr. Irradiation may be before or after contact with the oxystannane. A number of examples are given. (UK)

Radiochemistry and Nuclear Chemistry

Refer also to abstracts: 45288, 45311, and 45927.

Hot-Atom Chemistry

45356 EFFECTS OF PRE-IRRADIATION TREATMENTS ON THE CHEMICAL BEHAVIOR OF ^{64}Cu RECOIL ATOMS IN NEUTRON-IRRADIATED COPPER PHTHALOCYANINE. Kudo, Hiroshi (Japan Atomic Energy Research Inst., Tokai). Bull. Chem. Soc. Jap.; 45: No. 5, 1311-13 (May 1972).

The chemical behavior of ^{64}Cu recoil atoms was studied in copper phthalocyanine that had been quenched from an elevated temperature or exposed to ^{60}Co γ -rays before the neutron irradiation. In the quenched β -crystal, the initial retention was found to increase with an increase in the temperature or pre-irradiation heating, while the plateau retention of the post-neutron annealing reaction decreased. These results were compared with those for cadmium phthalocyanine and were understood in terms of "dislocation" in the crystal, the recoil atoms being assumed to be trapped and stabilized in the "dislocation." Although the variations were small, the irradiation with ^{60}Co γ -rays caused a change in the chemical behavior of the recoil atom. The role of some radiation-induced species was discussed. (auth)

45357 NUCLEAR RECOIL REACTIONS IN ORGANOMANGANESE COMPOUNDS. VI. $\text{IMn}(\text{CO})_5$ AND $\text{BrMn}(\text{CO})_5$. De Jong, I. G.; Wiles, D. R. (Carleton Univ., Ottawa). Can. J. Chem.; 50: No. 7, 961-5 (1 Apr 1972).

Radiochemical reactions of ^{82}Br , ^{128}I , and ^{56}Mn were studied in neutron-irradiated $\text{BrMn}(\text{CO})_5$ and $\text{IMn}(\text{CO})_5$. The parent compounds were found to contain 49 and 28% of the ^{82}Br and ^{128}I , but only 6 and 10% of the ^{56}Mn , respectively. The dimeric $[\text{XMn}(\text{CO})_4]_2$ accounted for 28% of the ^{82}Br , but only about 2% of the ^{56}Mn . Some $^{56}\text{Mn}_2(\text{CO})_{10}$ was also found. The radiochemical yields of several products were found to increase substantially on thermal treatment of the target material. Reaction pathways are suggested which can account for the thermal effects, which involve halogen atom transfer and carbonyl transfer in the solid. (auth)

45358 CHEMICAL EFFECTS ASSOCIATED WITH THE THERMAL NEUTRON IRRADIATION OF SOLID ACETANILIDE (ANTIFIBRIN). Barakat, M. F.; Zahran, A. H. (Atomic Energy Authority, Cairo). Radiochim. Acta; 17: No. 2, 65-8 (Jul 1972).

Reactions of recoil ^{14}C atoms were studied in a neutron-irradiated acetanilide system. The reaction mechanisms in solid systems are discussed with reference to those occurring in simple gaseous and liquid systems. The "random fragmentation" hypothesis seems to offer an adequate explanation. Most of the products identified were the result of different recombination processes between fragments formed by the severe fragmentation of the irradiated material. ^{14}C -labeled acetanilide was obtained with the highest radiochemical yield. The radiochemical yields of the simple synthetic products were higher than those of more complicated products. Labeled methyl derivatives were not detected. The infrared spectrum of the polymer separated showed a marked similarity, in absorption bands, to the spectrum of pure acetanilide except for the possible appearance of $-\text{CH}_2-$ groups in the polymer spectrum. (auth)

45359 CHARGE STATES OF ^{208}Pb AND ^{208}Tl FORMED BY α DECAY OF ^{212}Po AND ^{212}Bi . Meyer, J.; Paulus, J. M.; Abbe, J. Ch. (Centre de Recherches Nucleaires, Strasbourg). Radiochim. Acta; 17: No. 2, 76-81 (Jul 1972). (In French).

Charge states of ^{208}Pb and ^{208}Tl formed from the α decay of a

solid deposit of ^{212}Po and ^{212}Bi were analyzed by electrostatic deflection. In the absence of an internal conversion, the charge of recoils varies from -1 to +6. The yields of $^{208}\text{Pb}^+$ and $^{208}\text{Pb}^0$, $^{208}\text{Tl}^+$ and $^{208}\text{Tl}^0$ are of the same order of magnitude, but the intensities of $^{208}\text{Pb}^{n+}$ and $^{208}\text{Tl}^{n+}$ decrease strongly when the charge increases. The mean charge is less than one. The backing influences the charge carried by the recoil as it leaves the surface. The diffusion of the parent into the backing diminishes the yields of the multicharged daughters. When α decay is followed by internal conversion, the charge spectrum of $^{208}\text{Tl}^{n+}$ lies between +5 and +20 with a mean charge of +12. This charge distribution is the result of vacancy cascades initiated by internal conversion in the L or M shell. In this case, no effect of the backing has been observed. (auth)

45360 CHEMICAL EFFECTS OF ISOMERIC TRANSITION IN BROMOETHANE MIXTURES. da Fonseca, A. J. R.; Shaw, D.; Shaw, P. F. D. (Oxford Univ.). *Radiochim. Acta*; 17: No. 2, 81-95 (Jul 1972).

The distribution of products from the decay of $^{80\text{m}}\text{Br}$, combined initially as bromine or bromoethane, was determined in bromoethane-bromine mixtures at room temperature. Comparative results at -80°C were also obtained with $^{80\text{m}}\text{Br}$ present initially as hydrogen bromide. Analysis was made using novel G.L.C. separation involving two columns in series. While the overall magnitude of the yields of ^{80}Br labeled products is roughly independent of the initial form of $^{80\text{m}}\text{Br}$, the intensity of the chemical damage, judged by the degree of bromination or the extent of C-C bond schism, appears to be correlated with the magnitude of the charge transferred from the freshly formed ^{80}Br atoms to bromoethane molecules. While radiolysis by Auger electrons may be partly responsible for part of the diffusive yields, it cannot account for the overall distribution of atoms which have undergone internal conversion, or, indeed, any other form of nuclear activation studied in the bromoethane-bromine system. It was also possible to show that the kinetic energy gained by mutual repulsion of charged atoms is insufficient to account for the observed product distribution. Several of the most important features of the distribution are satisfied by a model which assumes that dissociation (by mutual electrostatic repulsion) and neutralization occur at similar rates so that the faster dissociative processes, and the free radicals produced by them, determine the nature of the final products. (auth)

45361 CHEMICAL CONSEQUENCES OF THE REACTION $^{127}\text{I}(n,\gamma)^{128}\text{I}$ IN MIXTURES OF IODINE AND n-BUTANE IN THE GAS PHASE. Lagarde, G.; Abbe, J. Ch.; Paulus, J. M. (Centre de Recherches Nucleaires, Strasbourg). *Radiochim. Acta*; 17: No. 2, 96-101 (Jul 1972). (In French).

Gaseous mixtures of iodine and n-butane were submitted to neutron irradiation. Hot atom reactions yield to the following products: CH_2I_2 (31%), CH_3I (22%), $\text{C}_2\text{H}_5\text{I}$ (21%), $\text{C}_2\text{H}_5\text{I}$ (13%), $n\text{-C}_3\text{H}_7\text{I}$ (9%) and $s\text{-C}_3\text{H}_7\text{I}$ (4%) with a total organic yield of 1.10%. The results are in good agreement with the kinetic theory of Estrup-Wolfgang and give a calculated reactivity integral $I = 0.0071 \alpha_{\text{Ar}}$. The other reactions arise from excited iodine atoms or ions resulting in the formation of CH_3I (80%) and $\text{C}_2\text{H}_5\text{I}$ (20%) with a total organic yield of 0.94%. This study reveals the importance of decomposition reactions of the primary products. (auth)

45362 CHEMICAL CONSEQUENCES OF THE REACTION $^{127}\text{I}(n,\gamma)^{128}\text{I}$ IN MIXTURES OF IODINE AND n-BUTANE IN THE LIQUID PHASE. Lagarde, G.; Abbe, J. Ch.; Paulus, J. M. (Centre de Recherches Nucleaires, Strasbourg). *Radiochim. Acta*; 17: No. 2, 102-4 (Jul 1972). (In French).

The total organic yield of ^{128}I produced by radiative neutron capture in solutions of iodine and n-butane is 26% (for an iodine mole fraction of 0.003). The relative abundance of $\text{C}_4\text{H}_9\text{I}$ is 51%. The comparison of these results with those obtained in the gaseous phase reveals an important phase effect. The observed phenomena are interpreted as resulting from the direct interaction between ^{128}I and butane and from the biradical reactions arising from the autoradiolysis of butane. (auth)

45363 ESTRUP-WOLFGANG THEORY OF HOT REACTIONS. PART II. α , THE ENERGY LOSS PARAMETER. Urch, D. S. (Queen Mary Coll., London). *Radiochim. Acta*; 17: No. 2, 105-10 (Jul 1972).

The possibility is considered that α , the energy loss parameter used in the Estrup-Wolfgang theory of hot reactions, might vary with energy. Calculations by Estrup have shown that α probably is a function of energy. It is shown that only slight variations in α could vitiate the conclusions drawn from the theory as to the relative energies of "hot" atoms that produce particular products. Attempts are made to provide criteria for judging the relative energies of atoms that initiate different reactions. Allowing α to be

a function of energy provides an explanation for all the deviations from "classical" kinetic theory behavior that have so far been observed. (auth)

45364 ANNEALING REACTIONS IN POTASSIUM CHROMATE. PART 3. DOPANT-ATOM AND RECOIL-ATOM ANNEALING IN NEARLY IDENTICAL CRYSTALS OF POTASSIUM CHROMATE. Collins, Carol H.; Ackerhalt, Robert E.; Collins, Kenneth E. (State Univ. of New York, Buffalo. Western New York Nuclear Research Center, Buffalo). *Radiochim. Acta*; 17: No. 2, 73-5 (Jul 1972).

Thermal annealing of ^{51}Cr atoms, introduced into nearly identical K_2CrO_4 crystals as $^{51}\text{Cr}(\text{III})$ dopant and as ^{51}Cr recoil atoms, indicates several different categories of annealing reactions. The $^{51}\text{Cr}(\text{III})$ -dopant atoms and ^{51}Cr recoil atoms anneal at the same rates to identical quantities of $^{51}\text{Cr}(\text{VI})$ at temperatures above 150° , indicating that thermal annealing processes in K_2CrO_4 which occur above this temperature are insensitive to the exact structure of the annealable ^{51}Cr sites and reflect rather the properties of the bulk crystals. At intermediate temperatures (0 to 150°), the recoil atoms anneal at faster rates than the dopant atoms while at temperatures below 0° , comparable fractions of dopant atoms and recoil atoms anneal by processes too fast to follow by analysis at 0° . (auth)

Properties of Radioactive Materials

Refer also to abstract: 45313.

45365 (IEA-183) OPTIMIZATION STUDY OF THE PRECIPITATION OF THORIUM OXALATE. Ikuta, A.; Costa, E. C. (Instituto de Energia Atomica, Sao Paulo (Brazil)). Aug 1969. 20p. Dep. NTIS (U. S. Sales Only).

The maximization of the effects of 4 variables (time, temperature, initial free HNO_3 concentration, and excess oxalic acid) on the filtration rate of $\text{Th}(\text{C}_2\text{O}_4)_2 \cdot 6\text{H}_2\text{O}$ precipitated from a 60g $\text{ThO}_2/1$ thorium nitrate solution was studied. The precipitations were carried out in a batch agitated lab-scale reactor at constant weight of $\text{Th}(\text{C}_2\text{O}_4)_2 \cdot 6\text{H}_2\text{O}$, while in the filtration step a constant volume was filtered under constant vacuum through filter papers of the same lot. The optimization was effected by the Steepest Ascent Method employing data from a half-replicate of a 2^4 factorial design. A total of 25 tests were required. The analysis of the data indicated that maximum filtration rate for a 60g $\text{ThO}_2/1$ solution is obtained by precipitation at $60 \pm 4^\circ\text{C}$ in a $3.1 \pm 0.3\text{N}$ HNO_3 solution, using $20 \pm 3\%$ excess oxalic acid with a precipitation-digestion period of 28 ± 2 min. A check on these data was obtained by individual parametric variation around the maximum. Solutions containing 40 and 80 g $\text{ThO}_2/1$ were also precipitated at these conditions, however, results indicated the need for optimization at each particular solute concentration level. (auth)

45366 (JNRR-P6-6402) STUDY OF TRANSURANIUM ELEMENT HYDROLYSIS. I. STUDY OF HYDROLYSIS BY PARTITION CHROMATOGRAPHY (UO_2^{2+} , PuO_2^{2+}). Korotkin, Yu. S. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Nuclear Reactions). 1972. 15p. (In Russian). Dep. NTIS (U. S. Sales Only).

A possibility is considered of application of reverse phase partition chromatography to hydrolysis study. The scheme suggested was tested by studying the hydrolysis of uranyl and plutonyl ions. (auth)

45367 X-RAY PHOTOELECTRON SPECTRA OF URANIUM OXIDES. Chadwick, D.; Graham, J. (Central Electricity Research Labs., Leatherhead, Eng.). *Nature (London) Phys. Sci.*; 237: No. 77, 127-8 (19 Jun 1972).

X-ray photoelectron spectroscopy is an analytical technique that may prove useful in ascertaining the oxidation state of U in particular environments, and an examination was made of the spectra of UO_2 , U_3O_8 , and UO_3 , as well as of U metal. Details are given of the experimental technique, and the basis of the method, and results are shown tabulated. The measurements indicated a chemical shift of 1 to 1.2 eV between the 4f core levels in UO_3 and UO_2 . The observed shift between UO_2 and U_3O_8 was probably just significant (0.4 eV), although the 4f binding energies for U_3O_8 are distinctly different from those for UO_3 . The binding energies observed to increase from UO_2 through U_3O_8 were probably just as significant (0.4 eV), although the 4f binding energies for U_3O_8 are distinctly different from those for UO_3 . The binding energies were observed to increase from UO_2 through U_3O_8 to UO_3 , which is to be expected on the basis of the increase in the formal U oxidation state from $\text{UO}_2[\text{U}(\text{IV})]$ to $\text{UO}_3[\text{U}(\text{VI})]$. At room temperature U_3O_8 , supposedly a mixed valence state compound, gave only single peaks with a line width comparable with the other oxides. Results for oxide-coated U metal indicated that the crating has a stoichiometry close to UO_2 . Some interesting observations on abrasively cleaned metal are mentioned. (UK)

45368 DETERMINATION OF THE STRUCTURE OF MONOHYDRATED URANYL DIFORMATE USING X-RAY DIFFRACTION. Claudel, Bernard; Mentzen, Bernard; Navarro, Alain; Puaux, J. P. (Institut National des Sciences Appliquées, Villeurbanne, France). *Compt. Rend., Ser. C*; 274: No. 16, 1417-19 (17 Apr 1972). (In French).

Monohydrated uranyl diformate crystallizes in a formic acid medium as crystals having an orthorhombic symmetry. The x-ray diffraction patterns of monocrystals and powders were studied. The lattice was confirmed to be orthorhombic with the following lattice parameters: $a = 5.84 \pm 0.01\text{Å}$; $b = 11.55 \pm 0.02\text{Å}$; $c = 19.40 \pm 0.04\text{Å}$; the space group being $Fddd$. (France)

45369 IONIC SOLVATION NUMBERS FROM COMPRESSIBILITIES AND IONIC VIBRATION POTENTIALS MEASUREMENTS. Bockris, J. O'M.; Saluja, P. P. S. (Univ. of Pennsylvania, Philadelphia). *J. Phys. Chem.*; 76: No. 15, 2140-51 (20 Jul 1972).

Results are presented of work the general aim of which is to extend and deepen the model of the solvent sheath near ions in aqueous solutions. The experimental measurements reported are those of the compressibility of the solution. A modification of the method reported previously was used to give sound velocity at frequency of 5 MHz to an accuracy of $\pm 0.4 \text{ m sec}^{-1}$. The velocity measurements are converted into values of the compressibility, β , by the use of a well-known equation. Such measurements are then converted by the equation of Passynski into solvation numbers. It is desired to obtain solvation numbers. Ionic vibration potentials due to Madan and to Zana and Yeager were utilized. These give the difference of the masses of the ions of a salt, together with the water molecules which move with the ions when they receive an ultrasonic pulse. The individual solvation numbers were found to be $\text{Li}^+ = 4.0$, $\text{Na}^+ = 4.5$, $\text{K}^+ = 3.5$, $\text{Rb}^+ = 3.0$, $\text{Cs}^+ = 2.5$; $\text{Ba}^{2+} = 11.0$, $\text{Ca}^{2+} = 9.0$, $\text{Sr}^{2+} = 9.0$, $\text{Mg}^{2+} = 10.0$; $\text{La}^{3+} = 15.0$, $\text{Ce}^{3+} = 18.0$, $\text{Th}^{4+} = 22.0$, $\text{F}^- = 4.0$, $\text{Cl}^- = 2.0$, $\text{Br}^- = 1.8$, and $\text{I}^- = 1.5$. A theoretical discussion of the meaning of the compressibility and ionic vibration potential measurements is given. It is stressed that the discussion is made on the basis of a definition of solvation number (SN). It will be noted that this solvation number definition is time dependent. Correspondingly, a relationship between the reorientation time of the solvent and the time during which the ion waits in its movement in solution is shown and supports the definition given. An argument is given, which suggests that the compressibility and the ionic vibration potential measurements measure the same quantity. (auth)

45370 PRECIPITATION OF SOME ACTINIDE ELEMENT COMPLEX IONS BY USING HEXAMMINE COBALT(III) CATION. IV. THE PRECIPITATION OF URANIUM(VI) PEROXO COMPLEX ION WITH HEXAMMINE COBALT(III) ION. Hoshi, Michio; Ueno, Kaoru (Japan Atomic Energy Research Inst., Tokai). *J. Inorg. Nucl. Chem.*; 34: No. 7, 2277-82 (Jul 1972).

The precipitation behaviors of uranium(VI) peroxo complex ion were investigated with hexamminecobalt(III) cation. This precipitation reaction produces $[\text{Co}(\text{NH}_3)_6][(\text{UO}_2)_2(\text{O}_2)_4]_3 \cdot n\text{H}_2\text{O}$, which changes to monoperoxo species by hydrolysis in water. Monoperoxo species i.e. $[\text{Co}(\text{NH}_3)_6][(\text{UO}_2)_2(\text{O}_2)_2(\text{OH})_4]_3 \cdot 12\text{H}_2\text{O}$ belongs to a tetragonal system with $a = 7.18 \text{ Å}$ and $c = 13.02 \text{ Å}$. (auth)

45371 PROCESS FOR RECOVERING BY-PRODUCTS FROM URANIUM DIOXIDE PRODUCTION. Pirk, H.; Muller, H.; Hackstein, K. (to NUKEM-Nuklear-Chemie und Metallurgie Gesellschaft mbH). Canadian Patent 891,958. 1 Feb 1972. Priority date 23 Apr 1969, Germany.

A process is described for the recovery of uranium, ammonium hydrogen fluoride, ammonia, carbon dioxide, and water from the filtrates and waste gases resulting from the conversion of uranium hexafluoride to uranium dioxide by the precipitation of the uranium as tetraammonium uranyl tricarbonate (AUC) and the associated reductive pyrohydrolysis of this salt. (Canada)

45372 EPR STUDY OF VO^{2+} IN $\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ SINGLE CRYSTAL. Krishnamurthy, M. V. (Bhabha Atomic Research Centre, Trombay, India). *J. Phys. Chem. Solids*; 33: No. 8, 1645-9 (Aug 1972).

The electron paramagnetic resonance of VO^{2+} ion doped in uranyl nitrate hexahydrate single crystals was studied. The room temperature studies show that the VO^{2+} ion undergoes rapid reorientation or hindered motion in the crystal lattice. These motions are effectively quenched at liquid nitrogen temperature. The low temperature studies reveal that the V-O axis of the vanadyl ion is randomly oriented in the lattice and this randomness gives rise to the observed powder spectrum. The line-widths of the hyperfine lines at room temperature are found to depend on the nuclear quantum number, m_I . This line-width variation is satisfactorily explained employing McConnell-Kivelson theory based on the model of a rapid tumbling paramagnetic complex. (auth)

45373 THORIUM-ACETATE AND THORIUM-PROPIONATE COMPLEXES. Tedesco, P. H.; Anon, M. C. (Univ. Nacional, La Plata). *J. Inorg. Nucl. Chem.*; 34: No. 7, 2271-6 (Jul 1972).

Equilibrium of complexes formation was studied for the systems thorium-acetate and thorium-propionate at 25°C and ionic strength 1.0. At the experimental conditions only the first stability constant was possible to determine, using the potentiometric and spectrophotometric methods, which gave results in good agreement. (auth)

45374 STUDY ON CATALYSIS OF URANIUM OXIDE AND ITS MIXED CATALYSTS. II. STRUCTURE OF URANIUM-ANTIMONY OXIDE CATALYST AND CATALYTIC ACTIVITY FOR OXIDATION OF PROPYLENE TO ACROLEIN. Nozaki, Fumio; Okada, Hiroshi (Chiba Univ., Japan). *Nippon Kagaku Kaishi*; No. 5, 842-9 (May 1972). (In Japanese).

A uranium antimony oxide catalyst, claimed in the patents for the oxidation of propylene to acrolein, was investigated from a standpoint of structure and activity relationships. Catalytic activity tests were carried out by a flow system and a pulse reaction method. X-ray pattern, D.T.A.-T.G.A. curve and low frequency infrared spectrum of the U-Sb binary oxide catalyst system were measured for characterizing the catalyst structure. The following conclusions were obtained. (1) High selectivity of propylene to acrolein in excess of 70% was achieved over the catalysts with a lower atomic U/Sb ratio than 4/6 and those calcined at high temperatures above 700°C. (2) The formation of the double oxide ($\text{USb}_3\text{O}_{10}$) was observed by calcination at about 620°C. On the basis of direct relationship between a calcination temperature above 700°C and appearance of catalytic activity for selective acrolein formation, $\text{USb}_3\text{O}_{10}$ was recognized as the active component of the catalyst. (3) From the experiments on reducing the catalyst with propylene pulses and reoxidizing the catalyst with oxygen pulses, interesting information on the catalysis was obtained. (auth)

45375 EXISTENCE OF REGULARITIES, TETRAD, OR DOUBLE-DOUBLE EFFECT IN THE CHANGES IN UNIT CELL VOLUMES OF THE COMPOUNDS OF THE ACTINIDES IN DIFFERENT OXIDATION STATES. Siekierski, S.; Fidelis, I. (Inst. of Nuclear Research, Warsaw). *J. Inorg. Nucl. Chem.*; 34: No. 7, 2225-31 (Jul 1972).

It is shown that the contraction of the actinides as measured by the changes with atomic number, Z, of the unit cell volumes of their compounds in the +3, +4, and +6 oxidation states, shows the same double-double effect (regularities, or tetrad effect) as that observed in the case of the lanthanides. (auth)

45376 IDENTIFICATION OF AN AmCl_6^{3-} OCTAHEDRAL COMPLEX FROM THE Am(III) ABSORPTION SPECTRUM IN ETHANOL SOLUTIONS OF HCl. Barbanel, Yu. A.; Kotlin, V. P.; Gorski, A. G. *Dokl. Akad. Nauk SSSR*; 202: No. 4, 830-3 (1 Feb 1972). (In Russian).

At certain concentrations of HCl in ethanol solutions of Am(III) a spectrum was obtained, which was used to assign an octahedral configuration to the americium present. Additional work with Pr(III) and Nd(III) aided in the substantiation of the spectroscopic identification of a AmCl_6^{3-} complex, and in confirming the existence of the set of complexes LnCl_6^{3-} . (K.S.W.)

45377 CHEMICAL SHIFTS IN THE $L_{\alpha 1}$ X-RAY LINE OF SOME URANIUM COMPOUNDS. Vdovenko, V. M.; Makarov, L. L.; Suglobova, I. G.; Volkov, V. A.; Chibisov, N. P. (Leningrad State Univ.). *Dokl. Akad. Nauk SSSR*; 202: No. 4, 868-70 (1 Feb 1972). (In Russian)

Chemical shifts in the $L_{\alpha 1}$ line of U for a series of chlorine- and oxygen-containing compounds were obtained: for example in the pairs $\text{CsUCl}_5 - \text{UCl}_4$, $\text{Cs}_2\text{UCl}_6 - \text{UCl}_4$, $\text{UCl}_3 - \text{UO}_3$, $\text{K}_2\text{UCl}_6 - \text{Rb}_2\text{UCl}_6$, and $\text{UO}_2 - \text{U}$, shifts accompanying the change in chemical state of the U amounted to as much as $\sim 1 \text{ eV}$. These shifts are ascribed to a participation of the 5f electrons in the electron structure of U. (K.S.W.)

45378 U_3X_5 TYPE SULFIDES AND SELENIDES OF U AND Ca, Sr, Ba, Pb, AND Eu (X = S, Se). Brochu, Robert; Padiou, Jean; Prigent, Jacques (Univ., Rennes, France). *Compt. Rend., Ser. C*; 274: No. 10, 959-61 (6 Mar 1972). (In French).

MU_2S_5 (M = Pb, Eu) and MU_2Se_5 (M = Ca, Sr, Ba, Pb, Eu) were synthesized. Studies on monocrystals of PbU_2X_5 (X = S, Se) confirm the orthorhombic cell with the probable $\text{P2}_12_12_1$ space group. The structural similarities between U_3S_5 and MU_2Se_5 are confirmed by the existence of continuous solid solutions: $\text{M}_x\text{U}_{(3-x)}\text{Se}_5$ ($0 \leq x \leq 1$). (France)

45379 FLUORINATION OF UO_2 AND U_3O_8 BY GASEOUS BROMINE TRIFLUORIDE. Iwasaki, M.; Sakurai, T. (Japan Atomic Energy Research Inst., Tokai). *J. Inorg. Nucl. Chem.*; 34: No. 7, 2189-2200 (Jul 1972).

The reactions of UO_2 and U_3O_8 with gaseous BrF_3 were studied gravimetrically. The specimens of uranium oxides were in pelletized form. The kinetic equations were determined as follows: $k[\text{g}/\text{cm}^2 \cdot \text{h}] = 0.346 \exp(-3900/\text{RT})$ for UO_2 , $k[\text{g}/\text{cm}^2 \cdot \text{h}] = 0.024 \exp(-950/\text{RT})$ for U_3O_8 . The activation energies were 3.9 and 0.95 kcal/mole for UO_2 and U_3O_8 , respectively. A little amount of solid intermediate was formed for both oxides. Their chemical form, however, was not identified. The reaction process was discussed by considering the chemical behavior of BrF_3 . (auth)

45380 COUPLING SCHEME IN URANYL COMPLEXES. Goerllier-Walrand, C.; Vanquickenborne, L. G. (Univ. Leuven, Heverlee, Belg.). *J. Chem. Phys.*; 57: No. 4, 1436-40 (15 Aug 1972).

Simple theoretical considerations of the ligand field type were applied to the electronic structure of insular uranyl complexes. The study of the consequences of equatorial ligation is used as a means to test the applicability of different coupling schemes for the isolated uranyl ion. From the analysis of the spectra of a large number of different uranyl complexes of D_{nh} symmetry, it was previously shown how the effect of equatorial ligation consists in lifting certain degeneracies of the free uranyl electronic energy levels and introducing variations in the corresponding transition probabilities. In this note the equatorial field operator is successively applied to uranyl wavefunctions derived in a Russell-Saunders approximation, in an ω - ω coupling approximation, and in an intermediate scheme. Only Russell-Saunders coupling allows a satisfactory explanation of the observed splitting and intensity pattern. (auth)

Radioactive Waste Treatment

Refer also to abstracts: 45190 and 46098.

45381 (ARH-SA-120) PHOSPHATE SOLIDIFICATION OF CAUSTIC PROCESS SOLUTION WASTE. Puryear, Dean A.; Buckingham, J. Steve; Campbell, Milton H. (Atlantic Richfield Hanford Co., Richland, Wash.). May 1972. Contract AT(45-1)-2130. 14p. (CONF-720628-2). Dep. NTIS.

From twenty-seventh ACS northwest regional meeting; Corvallis, OR. (15 Jun 1972).

A process for solidifying concentrated high-salt process solution waste with phosphoric acid is investigated on a laboratory-scale. Discussion of the proposed application of this process is presented, as well as the tests and results used to establish the parameters for solidification candidates. Synthetic high-salt caustic wastes are used to accumulate data on the kinetics of the reaction. The product is postulated to be the chemical complex $7(\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O})\text{NaOH}$ or a derivative thereof. Quality of the product is directly related to wt % water, total sodium molarity, and the amount of monovalent sodium salt present in the waste candidate. (auth)

45382 (ARH-SA-130) PLUTONIUM-URANIUM PARTITIONING BY A REFLUX SOLVENT EXTRACTION FLOWSHEET. Bruns, L. E. (Atlantic Richfield Hanford Co., Richland, Wash.). 4 May 1972. 22p. (CONF-720607-19). Dep. NTIS.

From eighteenth annual American Nuclear Society conference; Las Vegas, NV. (18 Jun 1972).

A solvent extraction flowsheet was developed for the processing of unirradiated plutonium-uranium scrap. By refluxing plutonium product as plutonium(III) in a multistage plutonium stripping-partitioning system, uranium is salted into the organic and pure plutonium is quantitatively separated from uranium. A hydrofluoric acid strip removes the final traces of plutonium from the organic phase containing the uranium. The flowsheet was plant-tested; and problems evident in the plant test were solved by comprehensive laboratory tests. Plant flowsheets were successfully tried; further laboratory tests were made to enhance column control. (auth)

45383 (ICP-1008) DTA-X-RAY DIFFRACTION EXAMINATION OF CHEMICAL REPROCESSING WASTES CONTAINING SODIUM AND POTASSIUM NITRATES. Newby, B. J.; Dickerson, Edwarda S. (Allied Chemical Corp., Idaho Falls, Idaho). Apr 1972. Contract AT(10-1)-1375. 32p. Dep. NTIS.

Sodium and potassium nitrates pose problems in fluidized-bed calcination; the large temperature range over which the nitrates exist in the molten-undecomposed state makes the fluidized-bed prone to agglomeration. Differential thermal analyses combined with x-ray diffraction were used to scope methods for lowering the decomposition temperature of sodium and potassium nitrates in wastes. The decomposition temperature of sodium nitrate in wastes can be lowered effectively by adding alumina, aluminum nitrate, sulfuric acid, ammonium sulfate, or magnesium sulfate. The same reagents are potential reagents for lowering the decomposition temperature of potassium nitrate in wastes. (auth)

45384 METHOD OF DECONTAMINATING CESIUM. Berton, Yves; Chauvet, Paul; des Ligneris, Jean (to Commissariat a l'Energies Atomique). French Patent 2,067,743. 26 Jul 1971. Filed 14 Nov 1969. (In French).

A method for decontaminating Cs, applicable to the decontamination of effluents arising from the processing of irradiated nuclear fuels, is described. The method involves the preparation of a colloidal nickel ferrocyanide suspension and the addition of this suspension to the effluent (at a pH not exceeding 11). This ratio of the Ni and ferrocyanide ion concentrations in the colloidal suspension is less than 1.5. (France)

45385 ESTER PROGRAM: HIGH-LEVEL RADIOACTIVE WASTE SOLIDIFICATION. BENCH-SCALE PHOSPHATE GLASS VITRIFICATION TESTS WITH SIMULATED FISSION PRODUCT SOLUTIONS. Bocola, W.; Donato, A. (CNEN, Rome). *Energ. Nucl. (Milan)*; 19: No. 6, 390-401 (Jun 1972).

The ESTER Program, carried out by the Casaccia Center of CNEN, is aimed at contributing to the solution of problems of treatment and disposal of high-level radioactive wastes resulting from nuclear fuel reprocessing. The process studied in the frame of this program is the fixation of fission product solutions in stable and insoluble glasses. Results and the practical experience achieved from twenty-seven phosphate glass vitrification runs, carried out in a bench-scale plant with six different types of simulated fission product solutions are described. The solids produced, containing 2 to 16% in moles and 4 to 22 wt % of fission product oxides, are of the glassy type and in some cases of ceramic type. Their leaching rates, determined at 25°C in distilled water, are 10^{-6} g $\text{cm}^{-2}\text{d}^{-1}$. Ruthenium volatilization is lower than 0.75% when red phosphorus is used as a glass former additive, but higher in the case of phosphoric acid. (auth)

45386 METHOD OF DISPOSING OF RADIOACTIVE ORGANIC WASTE SOLUTIONS. Knoll, Kenneth C. (to United States Atomic Energy Commission). U. S. Patent 3,666,673. 30 May 1972. Filed 24 Dec 1969.

Radioactive organic waste solutions may be disposed of and the radioactive material contained therein may be concentrated by passing the wastes as a liquid or vapor into a suitable preheated oxidation catalyst in the presence of air and collecting the radioactive residue from the oxidized organic wastes. (Official Gazette)

45387 SYSTEM FOR TOTAL IODINE RETENTION. Maeck, William J. (to United States Atomic Energy Commission). U. S. Patent 3,658,467. 25 Apr 1972. Filed 28 Jul 1969.

A method of absorbing and retaining air-borne inorganic iodine and organic iodine species by passing a gaseous stream containing these iodines through a filter bed of synthetic zeolite in a metal ion-exchanged form, which metal is reactive with iodine, is described. (Official Gazette)

45388 MODIFIED MACROPOROUS STYRENE-DIVINYLBENZENE COPOLYMERS AS THE SORBENTS WITH SELF-FIXATION ABILITY FOR THE REMOVAL OF VAPORS OF RADIOIODINE AND ITS COMPOUNDS. PART I. EQUILIBRIUM SORPTION OF METHYL IODIDE VAPORS ON NONMODIFIED MACROPOROUS STYRENE-DIVINYLBENZENE COPOLYMERS. Cejnar, Frantisek (Ustav Jaderneho Vyzkumu, Rez, Czech.). *Jad. Energ.*; 18: No. 6, 199-200 (Jun 1972). (In Czech).

Sorption and desorption isotherms were obtained for fission product iodine as methyl iodide on nonmodified macroporous styrene-divinylbenzene copolymers. The effects of various percentages of divinylbenzene on the sorption were studied. (M.C.G.)

45389 SORPTION OF POLYVALENT ELEMENTS ON SILICA GEL. II. SORPTION OF PROTACTINIUM ON SILICA GEL FROM MIXED AQUEOUS-ORGANIC SOLUTIONS OF HCl AND HNO_3 . Caletka, R. (Nuclear Research Inst., Rez, Czech.). *Collect. Czech. Chem. Commun.*; 37: No. 5, 1494-1503 (May 1972).

The effect of methanol, ethanol, 1-propanol, 2-propanol, acetone, and dioxane on the sorption of protactinium on silica gel was studied. It was found that in the presence of these solvents the sorption of protactinium decreases and the sorption kinetics are slower. The values of the distribution coefficients depend not only on the composition of the solution but, in several cases, also on its age. The sorption kinetics are mainly governed by the structure of silica gel. The effect of alcohols on the sorption of protactinium increases with the increasing number of carbon atoms in the alcohol molecule. The results are explained by the formation of ionic associates. The associates in solutions with the concentrated hydrochloric acid have the probable formula $(\text{SH}(\text{H}_2\text{O})_n \cdot \text{S})_{x-5} \cdot \text{PaCl}_x$ (where S denotes the organic solvent molecule and $x = 6$ to 8). In the case of acetone and dioxane, the formation of the well-sorbed associates $(\text{SH} \cdot \text{H}_2\text{O})_{x-5} \cdot \text{PaCl}_x$ is assumed. The sorption of protactinium from the hydrochloric and nitric acid

solutions in the presence of methanol and ethanol is compared. (auth)

45390 . SORPTION OF POLYVALENT ELEMENTS ON SILICA GEL. III. ISOLATION OF PROTACTIONIUM BY SORPTION ON SILICA GEL FROM THE SOLUTIONS OF HCl and H_2SO_4 . Caletka, R. (Nuclear Research Inst., Rez, Czech.). Collect. Czech. Chem. Commun.; 37: No. 5, 1684-9(May 1972).

The sorption of trace amounts of protactinium on silica gel was studied under dynamic conditions together with the desorption effectiveness of various aqueous and aqueous-organic solutions. The possibility to separate protactinium, zirconium, hafnium, and niobium in the hydrochloric or sulfuric acid solutions was proved experimentally. Protactinium was also separated from the activated thorium salts and from the uranium ore. The mixture of 30% of 11.5M-HCl and 70% of 2-propanol or 10M- H_2SO_4 can be used for the elution of protactinium adsorbed on the column of silica gel. (auth)

45391 SORPTION OF POLYVALENT ELEMENTS ON SILICA GEL. IV. SEPARATION OF THE TETRAVALENT AND PENTAVALENT PROTACTIONIUM. Caletka, R.; Knobloch, V. (Nuclear Research Inst., Rez, Czech.). Collect. Czech. Chem. Commun.; 37: No. 5, 1690-2(May 1972).

The possibility of the tetravalent and pentavalent protactinium separation on silica gel was studied and it was found that in the presence of hydrochloric and sulfuric acids the protactinium adsorbed on the silica gel column is reduced and washed out by chromium(II) solutions whereas the nonreduced fraction of protactinium remains adsorbed on the column. The method was tested with trace amounts of the ^{233}Pa nuclide. It was found that under similar conditions the elution of niobium from silica gel proceeds only partly. (auth)

Radioisotope Production

Refer also to abstracts: 45223 and 45260.

45392 (N-72-15068) CYCLOTRON PRODUCTION OF ^{123}I : AN EVALUATION OF THE NUCLEAR REACTIONS WHICH PRODUCE THIS ISOTOPE. Sodd, Vincent J.; Scholz, Kenneth L.; Blue, James W.; Wellman, Henry N. (National Aeronautics and Space Administration, Cleveland, Ohio. Lewis Research Center). Oct 1970. 45p. (NASA-TM-X-67594; BRH/DMRE-70-4). NTIS.

Various nuclear reactions are described by which ^{123}I , a low radiation dose radiopharmaceutical, can be cyclotron-produced. Methods of directly producing ^{123}I and those that indirectly produce the radionuclide through the beta (+) decay of its natural precursor, ^{123}Xe . It is impossible to separate the radioiodine contaminants that occur in the direct method. Thus, it is preferable to produce pure ^{123}I from ^{123}Xe , which is easily separated from the radioiodines. Among the characteristics of ^{123}I is the capability of reducing the patient dose in a thyroid uptake measurement to a very small percentage of that delivered by the more commonly used ^{131}I . (auth) (STAR)

45393 CONVERSION OF CALIFORNIUM α -HYDROXYISOBUTYRATE TO OXIDE WITH CARBOXYLIC ACID RESIN. Harbour, R. M.; Hale, W. H. Jr. (E. I. duPont de Nemours and Co., Aiken, S. C.). Radiochem. Radioanal. Lett.; 10: No. 6, 339-44 (15 Jul 1972).

A method was developed to recover ^{252}Cf from ion exchange product solutions and prepare $^{252}Cf_2O_3$. The process was demonstrated with ^{252}Cf tracer and bulk neodymium stand-in; an ammonium α -hydroxyisobutyrate-Nd-Cf solution was adjusted to pH 9, Nd-Cf was sorbed on carboxylic acid resin, and the oxide was prepared by calcination at 600°C. (auth)

45394 PRODUCTION OF CARRIER-FREE ^{54}Mn BY THE α -PARTICLE BOMBARDMENT OF NATURAL CHROMIUM. Akiha, Fumimasa (Hiroshima Univ., Japan); Yuita, Koichi; Aburai, Tamaru; Nozaki, Tadashi; Murakami, Yukio. Nippon Kagaku Kaishi; No. 5, 908-13(May 1972). (In Japanese).

For the production of ^{54}Mn by the α - and ^3He -particle reactions on natural chromium, their excitation curves and thick-target yields were measured. Various conditions in the production procedure, especially the chemical separation of manganese in a hydrochloric acid solution of the bombarded target after the removal of ^{52}Fe were examined. A stack of thin chromium plates was bombarded by a beam of α - or ^3He -particles of 44-MeV maximum energy, and the manganese activities were measured non-destructively or after the chemical separation. The excitation curve of the α -reaction shows a maximum of 490 mb near 30-MeV energy, and that of the ^3He -reaction, a maximum of 80 mb near 18-MeV. The thick-target yields for 40-MeV incident energy are 6 $\mu Ci/\mu Ah$ and 1.1 $\mu Ci/\mu Ah$ for the α reaction and the ^3He reaction, respectively. The α reaction thus proved to give a higher

yield of ^{54}Mn than the other possible reactions. The bombarded chromium target was dissolved in a hydrochloric acid solution (8N) from which the ^{52}Fe was removed by an extraction method. The remaining solution was saturated with pure HCl gas by bubbling it through the solution, from which almost all the nonradioactive chromium and other active contaminants were removed by an anion exchange method. The manganese solution thus obtained was evaporated to dryness, and the Cl^- -free residue was dissolved in 50 ml of 10 mol/l H_2SO_4 together with 2g of KIO_3 . Then the Mn was distilled out as Mn_2O_7 from the resulting solution into a trap containing a small amount of dilute HCl solution plus H_2O_2 . To maintain the solution at 10 mol/l of sulfuric acid concentration and 166°C, which is the optimum condition for distillation, azeotropic nitric acid vapor (120.5°C) was added gradually to the boiling solution. The ^{54}Mn was obtained in a carrier-free state. The chemical yields over all the chemical procedure proved to be >90% with a sufficient radiochemical purity. (auth)

45395 PRODUCTION OF CARRIER-FREE $^{197m}, ^{197}Hg$ WITH A CYCLOTRON. Wilkniss, P. E.; Beach, L. A.; Marlow, K. W. (Naval Research Lab., Washington, D. C.). Radiochim. Acta; 17: No. 2, 110-13(Jul 1972).

The practical aspects of the production of carrier-free mercury in a cyclotron by proton bombardment of a gold target are discussed. A liquid nitrogen cooled target assembly and the radiochemical procedure for the recovery of carrier-free mercury from the gold target are described. Gamma spectra obtained with a Ge(Li) detector and a Si(Li) detector are shown. Experimental yield determinations for $^{197m}, ^{197}Hg$ are compared with theoretical calculations. (auth)

45396 UTILIZATION OF RESEARCH REACTORS FOR RADIO-ISOTOPE PRODUCTION. Rahman, M. Matiur (Pakistan Inst. of Nuclear Science and Tech., Rawalpindi). Nucleus (Karachi); 8: No. 4, 29-32(1971).

The radioisotope production facility at the Pakistan Institute of Nuclear Science and Technology (PINSTECH) is described. It is suggested that efforts should be directed towards the establishment of methods for the selection of target materials and the determination of chemical processes for production. Besides, some basic research on nuclear reactions and the chemical effects of nuclear transformation is also vital for radioisotope production, to say nothing of the guaranteeing of high radiochemical purity for the products, procedures of activation analysis of target materials, etc. (auth)

Reactor Fuel Reprocessing

Refer also to abstracts: 45308 and 47300.

45397 (CONF-720903-9) CONTINUOUS OR SEMICONTINUOUS LEACHER FOR LEACHING SOLUBLE CORE MATERIAL FROM SHEARED SPENT NUCLEAR FUEL TUBES. Odom, C. H. (Oak Ridge National Lab., Tenn.). 1972. 32p. Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

The effective dissolution of sheared liquid-metal fast breeder reactor fuels is considered. Concepts and models representative of current approaches to the continuous or semicontinuous dissolution of these fuels were developed to assess the magnitude of the difficulties to be encountered in controlling the flow, residence time, agitation, and transfer of the sheared material through such a device. The performance of many of the models developed was successfully demonstrated, but the model of the compartmented rotary drum continuous leacher with sloped transfer ducts is representative of the most effective and straightforward design required to meet all of the performance requirements. (auth)

45398 (DOCKET-50201-125) WEST VALLEY REPROCESSING PLANT. Quarterly Report for April 1, 1972-June 30, 1972, Part I. (Nuclear Fuel Services, Inc., West Valley, N. Y.). 14 Jul 1972. 8p. Dep. NTIS.

Environmental samples were analyzed for ^{131}I , alpha activity, and beta activity. Total discharges in liquid and gaseous effluents are given for gross α , gross β , tritium, ^{90}Sr , ^{129}I , ^{85}Kr , and ^{131}I . Surveillance tests and filter replacements are listed. (M.C.G.)

45399 (DOCKET-50201-126) WEST VALLEY REPROCESSING PLANT. Technical Specifications, Request for Approval of Revisions. (Nuclear Fuel Services, Inc., Rockville, Md.). 19 Jul 1972. 48p. Dep. NTIS.

Revisions are requested for technical specifications for: form of materials, extractant concentration, plutonium ion exchange operation, caustic concentration in carbon steel waste storage tanks, solid radioactive waste burial, evaporator steam pressure, respiratory protection equipment, process instrumentation, high

radiation area access, contamination and radiation control, slump alarms and eductors, dissolver dilution air, boric acid, and solvent analysis. (M.C.G.)

45400 (DOCKET-50201-127) WEST VALLEY REPROCESSING PLANT. Low Level Waste Disposal. (Nuclear Fuel Services, Inc., Rockville, Md.). 21 Jul 1972. 2p. Dep. NTIS.

Lagoon No. 3 was scraped clean of settled contamination and transfers to it will be from Lagoons 4 and 5, which receive discharge from the Low Level Waste Treatment facility. Piping was installed between holding lagoons at the solid waste burial site and the feed lagoon for the Low Level Liquid Waste Treatment plant. (M.C.G.)

45401 (DOCKET-50201-128) WEST VALLEY REPROCESSING PLANT. Comments Concerning Nuclear Criticality. (Directorate of Licensing (AEC), Washington, D. C.). 8 Aug 1972. 3p. Dep. NTIS.

Criticality in the proposed intercycle evaporators and plutonium solvent extraction system is considered. (M.C.G.)

45402 (DOCKET-50268-31) MIDWEST FUEL RECOVERY PLANT. Notice of Consideration of Issuance of a Facility Operating License and Notice of Opportunity for Hearing. (Directorate of Licensing (AEC), Washington, D. C.). 1 Aug 1972. 5p. Dep. NTIS.

45403 (DOCKET-50268-32) MIDWEST FUEL RECOVERY PLANT. Final Safety Analysis Report, Amendment 27. (General Electric Co., San Jose, Calif. Nuclear Fuel Dept.). 9 Aug 1972. 5p. Dep. NTIS.

An amendment is submitted for requesting temporary use of the service steam system to supply process steam during uranium cold run testing. The change is required because of the failure of the original process supply steam generator. Safety precautions that will be taken are listed. (M.C.G.)

45404 (DOCKET-50268-33) MIDWEST FUEL RECOVERY PLANT. Final Safety Analysis Report, Amendment 28. (General Electric Co., San Jose, Calif. Nuclear Fuel Dept.). 8 Aug 1972. 6p. Dep. NTIS.

A proposal is submitted for operation of the Wiped Film Evaporation System prior to commencement of normal plant operations. It is designed to concentrate wastes in the low-activity waste vault by evaporating their water content. (M.C.G.)

45405 (ICP-1006) IDAHO CHEMICAL PROGRAMS ANNUAL TECHNICAL REPORT, FISCAL YEAR 1971. Bower, J. R. (ed.) (Allied Chemical Corp., Idaho Falls, Idaho. Idaho Chemical Programs Operations Office). May 1972. Contract AT(10-1)-1375. 82p. Dep. NTIS.

The operating experience in fuel processing, waste calcining, and other waste management activities at the ICPP for FY 1971 is summarized. Processing of zirconium fuels was highly satisfactory, and successful startup and operation of a fluidized-bed denitration process for conversion of product uranyl nitrate to UO_3 was achieved. The first campaign for fluid-bed calcination of reactor fuel wastes using in-bed combustion of fuel oil to supply heat was completed with 100% on-stream time and at an increased processing rate over the original WCF (Waste Calcining Facility) system. Technical support activities in areas of current and proposed operations at the ICPP are also presented. These include: corrosion studies on processing equipment and fuel storage facilities; modifications to ICPP processes; developments in the electrolytic-dissolution process; developments in the combustion-dissolution process for graphite fuels; plans for future fuel storage; pilot-plant calcination of varied waste solutions; plans for improved environmental protection at the ICPP. Research programs in other areas include: safeguards in uranium accountability; burnup determination for fast breeder reactor fuels; iodine behavior and iodine adsorbents; ion-control membranes for separations processes. (auth)

45406 (ORNL-TM-2871) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR JANUARY 1970, NO. 11. (Oak Ridge National Lab., Tenn.). Feb 1970. Contract W-7405-eng-26. 102p. Dep. NTIS.

This report continues a series outlining progress in the development of methods for the reprocessing and fabrication of LMFBR fuels. Development work is reported on problems of irradiated fuel transport to the processing facility, the dissolution of the fuel and the chemical recovery of the PuO_2-UO_2 values, the containment of volatile fission products, product purification, preparation of fuel material by the sol-gel process, conversion of fuel processing plant product nitrate solutions to solids suitable for shipping and for fuel fabrication, fuel fabrication of sol-gel materials, and fuel evaluation studies, both in-pile and out-of-pile. Pertinent experimental results are presented for the information

of those immediately concerned with the field. Detailed description of experimental work and data are included in topical reports and in the Chemical Technology Division and Metals and Ceramics Division Annual Reports. (auth)

45407 (ORNL-TM-2918) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR FEBRUARY 1970, NO. 12. (Oak Ridge National Lab., Tenn.). Mar 1970. Contract W-7405-eng-26. 89p. Dep. NTIS.

FISSION PRODUCTS—chemistry in oxide fuel dissolution

IODINE—chemistry in oxide fuel dissolution

—volatilization from nitric acid solutions

—separation from fuel reprocessing off-gases by sorption on zeolites or mercury nitrates

LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (Feb. 1970) on

METHANE, IODO—separation from fuel reprocessing off-gases by sorption on zeolites or mercury nitrates

PHOSPHORIC ACID, TRIBUTYL ESTER—stability in Purex Process

PLUTONIUM—purification in Purex Process

—shipping costs for

PLUTONIUM OXIDES PuO_2 —production of sinterable powders of, from fuel reprocessing solutions

— PuO_2-UO_2 , preparation of sols of

— PuO_2-UO_2 , preparation by sol-gel process

— PuO_2-UO_2 , fabrication of sol-gel-derived

PUREX PROCESS—solvent stability in

REACTOR FUEL ELEMENTS—shipment of spent LMFBR, criticality and heat dissipation studies on

—reprocessing of stainless steel-clad oxide, shearing studies in

REACTOR FUELS—reprocessing of LMFBR, research progress (Feb. 1970) on aqueous

—performance studies on oxide

TRANSURANIUM PROCESSING PLANT—iodine evolution from dissolver solutions in

URANIUM OXIDES UO_2 —production of sinterable powders of, from fuel reprocessing solutions

—preparation of sols of, CUSP process for

— PuO_2-UO_2 , preparation of sols of

— PuO_2-UO_2 , preparation by sol-gel process

— PuO_2-UO_2 , fabrication of sol-gel-derived

WEST VALLEY PROCESSING PLANT—evaluation for handling LMFBR fuels (M.C.G.)

45408 (ORNL-TM-2949) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR MARCH 1970, NO. 13. (Oak Ridge National Lab., Tenn.). Apr 1970. Contract W-7405-eng-26. 97p. Dep. NTIS.

CARBON—reactions with nitrogen and uranium dioxide

FISSION PRODUCTS—solubility in nitric acid

FORMIC ACID—reactions with plutonium dioxide sols

IODINE—volatilization from spent oxide fuels in nitric acid dissolution

—separation from off-gases in LMFBR fuel reprocessing

—chemistry in acidic and alkaline solutions

LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (March 1970) on

METHANE, IODO—separation from off-gases in LMFBR fuel reprocessing

NEUTRONS—effects on plutonium dioxide-uranium dioxide fuels

NITROGEN—reactions with carbon and uranium dioxide

PLUTONIUM OXIDES PuO_2 — PuO_2-UO_2 , dissolution of irradiated, in nitric acid

—preparation from fuel reprocessing solutions, evaluation of processes for

— PuO_2-UO_2 , preparation by sol-gel process

—reactions of sols of, with formic acid

— PuO_2-UO_2 , preparation of microspheres of

— PuO_2-UO_2 , fabrication of sol-gel-derived

— PuO_2-UO_2 , fabrication of fuel rods of

— PuO_2-UO_2 , radiation effects on, neutron

PUREX PROCESS—flowsheet development for LMFBR fuels

RADIOCHEMICAL PROCESSING PLANTS—evaluation for reprocessing LMFBR spent fuels

REACTOR FUEL ELEMENTS—shipment of spent LMFBR, criticality and heat dissipation studies for

—reprocessing of stainless steel-clad oxide, shearing studies for

—swelling and gas release in stainless steel-clad oxide

REACTOR FUELS—performance studies on oxide

—reprocessing of LMFBR, research progress (March 1970) on aqueous

—photon spectra of spent LMFBR

—neutron source strength of spent LMFBR

URANIUM OXIDES UO_2 — PuO_2 — UO_2 , dissolution of irradiated, in nitric acid
 —preparation from fuel reprocessing solutions, evaluation of processes for
 —reactions with carbon and nitrogen
 —preparation by sol-gel process
 — PuO_2 — UO_2 , preparation by sol-gel process
 — PuO_2 — UO_2 , preparation of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , fabrication of fuel rods of
 — PuO_2 — UO_2 , radiation effects on, neutron (M.C.G.)

45409 (ORNL-TM-2996) LMFBR FUEL CYCLE STUDIES
 PROGRESS REPORT FOR APRIL 1970, NO. 14. (Oak Ridge National Lab., Tenn.). May 1970. Contract W-7405-eng-26. 82p. Dep. NTIS.

FISSION PRODUCTS—volatilization from oxide fuels
 —solubility in nitric acid
 IODINE—separation from fuel reprocessing off-gases
 —sorption on zeolites
 —chemistry of uncommon volatile species
 LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (April 1970) on
 NEUTRONS—effects on oxide fuels
 ORGANIC COMPOUNDS—oxidation in fuel reprocessing off-gas
 PLUTONIUM CARBONITRIDES— $PuCN$ — UCN , preparation of
 PLUTONIUM OXIDES PuO_2 —preparation from fuel reprocessing solutions, evaluation of methods for
 — PuO_2 — UO_2 , dissolution in nitric acid
 — PuO_2 — UO_2 , preparation of sols of
 — PuO_2 — UO_2 , preparation of microspheres from sols of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , analysis by γ spectroscopy
 — PuO_2 — UO_2 , radiation effects on, neutron
 PUREX PROCESS—flowsheet development for LMFBR fuels
 REACTOR FUEL ELEMENTS—shipment of spent LMFBR, criticality and heat dissipation studies on
 —cladding reaction with oxide fuels in
 REACTOR FUELS—reprocessing of LMFBR, research progress (April 1970) on aqueous
 —dissolution of spent oxide, in nitric acid
 —performance evaluation of oxide
 TRANSURANIUM PROCESSING PLANT—iodine evolution and trapping studies at
 TRITIUM—volatilization from oxide fuels
 URANIUM CARBONITRIDES— $PuCN$ — UCN , preparation of
 URANIUM OXIDES UO_2 —preparation from fuel reprocessing solutions, evaluation of methods for
 —preparation of sols of, CUSP process for
 — PuO_2 — UO_2 , dissolution in nitric acid
 — PuO_2 — UO_2 , preparation of sols of
 — PuO_2 — UO_2 , preparation of microspheres from sols of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , analysis by γ spectroscopy
 — PuO_2 — UO_2 , radiation effects on, neutron (M.C.G.)

45410 (ORNL-TM-3018) LMFBR FUEL CYCLE STUDIES
 PROGRESS REPORT FOR MAY 1970, NO. 15. (Oak Ridge National Lab., Tenn.). Jun 1970. Contract W-7405-eng-26. 94p. Dep. NTIS.

FISSION PRODUCTS—volatilization from oxide fuels
 IODINE—chemistry in nitric acid solutions in fuel reprocessing
 —sorption on zeolites
 LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (May 1970) on
 METHANE, IODO—sorption on zeolites
 ORGANIC COMPOUNDS—oxidation in fuel reprocessing cycle
 PLUTONIUM—purification in Purex Process
 PLUTONIUM OXIDES PuO_2 — PuO_2 — UO_2 , preparation by sol-gel process
 — PuO_2 — UO_2 , preparation of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , radiation effects on, neutron
 RADIOCHEMICAL PROCESSING PLANTS—evaluation for LMFBR spent fuel reprocessing
 REACTOR FUEL ELEMENTS—shipment of spent LMFBR, cask evaluation studies for
 —reprocessing of, volatile fission product removal in head-end
 —cladding reactions with oxide fuel in
 —cladding of oxide, economic and performance evaluation of materials for
 REACTOR FUELS—reprocessing of LMFBR, research progress (May 1970) on
 —radiation properties and shielding of LMFBR

—performance of oxide
 URANIUM OXIDES UO_2 —preparation by sol-gel process
 — PuO_2 — UO_2 , preparation by sol-gel process
 — PuO_2 — UO_2 , preparation of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , radiation effects on, neutron

45411 (ORNL-TM-3072) LMFBR FUEL CYCLE STUDIES
 PROGRESS REPORT FOR JUNE 1970, NO. 16. (Oak Ridge National Lab., Tenn.). Jul 1970. Contract W-7405-eng-26. 94p. Dep. NTIS.

FISSION PRODUCTS—volatilization from oxide fuels
 —diffusion of gaseous, from oxide fuels
 IODINE—volatilization from nitric acid
 —separation from fuel reprocess off-gas
 —sorption on zeolites
 LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (June 1970) on
 METHANE, IODO—separation from fuel reprocess off-gas
 —sorption on zeolites
 NEUTRONS—effects on oxide fuels
 NITRIC ACID—reactions with sodium
 PLUTONIUM—purification in Purex Process
 PLUTONIUM CARBONITRIDES— $PuCN$ — UCN , preparation of
 PLUTONIUM OXIDES PuO_2 — PuO_2 — UO_2 , dissolution in nitric acid
 —preparation from fuel reprocessing solutions, evaluation of methods for
 — PuO_2 — UO_2 , preparation by sol-gel process
 — PuO_2 — UO_2 , preparation and properties of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived, sphere-gas process for
 — PuO_2 — UO_2 , radiation effects on, neutron
 RADIOCHEMICAL PROCESSING PLANTS—evaluation for LMFBR fuel reprocessing
 REACTOR FUEL ELEMENTS—shipment of spent LMFBR, cask design studies for
 —performance studies on stainless steel-clad oxide
 —fuel pellet densification in
 REACTOR FUELS—reprocessing of LMFBR research progress (June 1970) on
 —criticality studies on LMFBR
 —fission product release from oxide
 SODIUM—reactions with nitric acid
 URANIUM CARBONITRIDES— $PuCN$ — UCN , preparation of
 URANIUM OXIDES UO_2 — PuO_2 — UO_2 , dissolution in nitric acid
 —preparation from fuel reprocessing solutions, evaluation of methods for
 —preparation of sols of, CUSP process for
 — PuO_2 — UO_2 , preparation by sol-gel process
 — PuO_2 — UO_2 , preparation and properties of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived, sphere-gas process for
 — PuO_2 — UO_2 , radiation effects on, neutron (M.C.G.)

45412 (ORNL-TM-3095) LMFBR FUEL CYCLE STUDIES
 PROGRESS REPORT FOR JULY 1970, NO. 17. (Oak Ridge National Lab., Tenn.). Aug 1970. Contract W-7405-eng-26. 76p. Dep. NTIS.

FISSION PRODUCTS—volatilization from oxide fuels
 —diffusion from oxide fuel
 IODINE—release in fuel reprocessing cycle
 —spectra of species of, in nitric acid solution
 —separation from gas streams by sorption on porous plastics or zeolites
 LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (July 1970) on
 NEUTRONS—effects on oxide fuels
 PLUTONIUM—purification in Purex Process
 PLUTONIUM OXIDES PuO_2 — PuO_2 — UO_2 , dissolution in nitric acid
 —preparation from solutions from fuel reprocessing cycle
 — PuO_2 — UO_2 , preparation of microspheres of
 — PuO_2 — UO_2 , fabrication of sol-gel-derived
 — PuO_2 — UO_2 , radiation effects on, neutron
 REACTOR FUEL ELEMENTS—shipment of spent LMFBR, cask evaluation studies for
 —thermal conductivity of sheared stainless steel-clad oxide
 —reprocessing of, volatilization of fission products in head-end
 —mechanical interaction at fuel-cladding interface in
 —fission product release in oxide
 REACTOR FUELS—reprocessing of LMFBR research progress (July 1970) on
 —performance of oxide
 URANIUM OXIDES UO_2 — PuO_2 — UO_2 , dissolution in nitric acid

- preparation from solutions from fuel reprocessing cycle
- preparation of sols of, CUSP process for
- preparation of microspheres of
- $\text{PuO}_2\text{--UO}_2$, preparation of microspheres of
- $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
- $\text{PuO}_2\text{--UO}_2$, radiation effects on, neutron (M.C.G.)

45413 (ORNL-TM-3127) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR AUGUST 1970, NO. 18. (Oak Ridge National Lab., Tenn.). Sep 1970. Contract W-7405-eng-26. 78p. Dep. NTIS.

- FISSION PRODUCTS—reactions in oxide fuels
- IODINE—spectra in nitric acid solutions
 - volatilization from nitric acid solutions
 - separation from gas streams in fuel reprocessing
 - distribution between aqueous and gaseous phases
- LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (August 1970) on
- METHANE, IODO—sorption on zeolites
- PLUTONIUM CARBONITRIDES— PuCN--UCN , preparation studies on
- PLUTONIUM OXIDES $\text{PuO}_2\text{--PuO}_2\text{--UO}_2$, dissolution in nitric acid
 - preparation from solutions from fuel reprocessing
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, neutron
- RADIOCHEMICAL PROCESSING PLANTS—evaluation for LMFBR spent fuel reprocessing
- REACTOR FUEL ELEMENTS—shipment of spent LMFBR, cask evaluation studies on
 - reprocessing of oxide, voloxidation fission product removal in head-end
- REACTOR FUELS—reprocessing of LMFBR research progress (August 1970) on aqueous
 - performance of oxide
- URANIUM CARBONITRIDES— PuCN--UCN , preparation studies on
- URANIUM OXIDES $\text{UO}_2\text{--PuO}_2\text{--UO}_2$, dissolution in nitric acid
 - preparation from solutions from fuel reprocessing
 - preparation of sols of, CUSP process for
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, neutron (M.C.G.)

45414 (ORNL-TM-3180) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR SEPTEMBER 1970, NO. 19. (Oak Ridge National Lab., Tenn.). Oct 1970. Contract W-7405-eng-26. 92p. Dep. NTIS.

- DISSOLVERS—forward mixing in continuous
- IODINE—properties in nitric acid solutions
 - sorption on porous plastic
 - chemistry of uncommon volatile species of
- KRYPTON—volatilization from oxide fuels
- KRYPTON-85—release from irradiated oxide fuels
- LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (Sept 1970) on
- METHANE, IODO—reactions with nitric acid
 - sorption on zeolites
 - sorption on porous plastic
- NITRIC ACID—reactions with methyl iodide
- PLUTONIUM OXIDES PuO_2 —preparation from solutions from fuel reprocessing
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process
 - $\text{PuO}_2\text{--UO}_2$, preparation of microspheres
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived
- RARE EARTH OXIDES—rare earth oxides- UO_2 , crystal structure of solid solutions in
- REACTOR FUEL ELEMENTS—shipment of spent LMFBR, cask evaluations for
 - reprocessing of, fission product removed by voloxidation in head-end
 - performance modeling of clad oxide
- REACTOR FUELS—reprocessing of LMFBR, research progress (Sept 1970) on aqueous
- SOLVENT EXTRACTION PROCESSES—evaluation for LMFBR fuel reprocessing
- TRITIUM—distribution in Zircaloy-clad uranium dioxide fuel rods
- URANIUM OXIDES UO_2 —tritium distribution in Zircaloy-clad fuel rods of
 - preparation from solutions from fuel reprocessing
 - preparation of sols of, CUSP process for
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process

- $\text{PuO}_2\text{--UO}_2$, preparation of microspheres of
- $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
- $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived
- rare earth oxides- UO_2 , crystal structure of solid solutions in (M.C.G.)

45415 (ORNL-TM-3217) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR OCTOBER 1970, NO. 20. (Oak Ridge National Lab., Tenn.). Nov 1970. Contract W-7405-eng-26. 88p. Dep. NTIS.

- FISSION PRODUCTS—volatilization from oxide fuels
- IODINE—volatilization from nitric acid
 - spectra in nitric acid solutions
 - partition between aqueous and gaseous systems
 - chemistry of uncommon volatile species of
- KRYPTON—volatilization from oxide fuels
- LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (Oct 1970) on
- METHANE, IODO—sorption from gas streams on solids
 - reactions with nitric acid
- PLUTONIUM CARBONITRIDES— PuCN--UCN , preparation of
- PLUTONIUM OXIDES $\text{PuO}_2\text{--PuO}_2\text{--UO}_2$, dissolution in nitric acid
 - preparation from solutions from fuel reprocessing
 - $\text{PuO}_2\text{--UO}_2$, preparation of microspheres of
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived
- REACTOR FUEL ELEMENTS—cladding mechanical interactions with oxide fuel in, FMØDEL code prediction of
 - shipment of spent LMFBR, heat dissipation and shielding studies for
- REACTOR FUELS—reprocessing of LMFBR, research progress (Oct 1970) on aqueous
 - performance evaluation of oxide
- SOLVENT EXTRACTION PROCESSES—evaluation for reprocessing LMFBR fuels
- URANIUM CARBONITRIDES— PuCN--UCN , preparation of
- URANIUM OXIDES $\text{UO}_2\text{--PuO}_2\text{--UO}_2$, dissolution in nitric acid
 - preparation from solutions from fuel reprocessing
 - preparation of sols of, CUSP process for
 - $\text{PuO}_2\text{--UO}_2$, preparation of microspheres of
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived (M.C.G.)

45416 (ORNL-TM-3250) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR NOVEMBER 1970, NO. 21. (Oak Ridge National Lab., Tenn.). Jan 1971. Contract W-7405-eng-26. 93p. Dep. NTIS.

- AIR—separation of fission product iodine and tritium from
- FISSION PRODUCTS—volatilization from oxide fuels
- IODINE—volatilization from nitric acid solutions
 - spectra in nitric acid solutions
 - separation from air by cold trapping
 - chemistry of uncommon volatile species of
- LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (Nov 1970) on
- METHANE, IODO—separation from air by cold trapping
 - sorption on solids, research progress on
- PLUTONIUM—purification in Purex Process
- PLUTONIUM OXIDES $\text{PuO}_2\text{--PuO}_2\text{--UO}_2$, oxidation of unirradiated pellets of
 - $\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid
 - preparation from solutions from fuel reprocessing
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process
 - $\text{PuO}_2\text{--UO}_2$, preparation of microspheres
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived
 - $\text{PuO}_2\text{--UO}_2$, analysis of pellets of, for moisture
 - $\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived
- REACTOR FUEL ELEMENTS—shipment of spent LMFBR, heat dissipation studies for
 - reprocessing of clad oxide, head-end steps in
 - criticality studies on spent LMFBR
 - cladding mechanical interaction with fuel in
 - performance evaluation of clad oxide
- REACTOR FUELS—reprocessing of LMFBR, research progress (Nov 1970) on aqueous
- SODIUM—reactions of, survey on
- TRITIUM—release in reprocessing of oxide fuels
 - separation from air by cold trapping
- URANIUM CARBONITRIDES—preparation of
- URANIUM OXIDES UO_2 —preparation from solutions from fuel reprocessing
 - preparation of sols of, CUSP process for
 - $\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process
 - $\text{PuO}_2\text{--UO}_2$, preparation of microspheres
 - $\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived

- PuO₂—UO₂, analysis of pellets of, for moisture
- PuO₂—UO₂, radiation effects on sol-gel-derived
- PuO₂—UO₂, oxidation of unirradiated pellets of
- PuO₂—UO₂, dissolution in nitric acid (M.C.G.)

45417 (ORNL-TM-3281) LMFBR FUEL CYCLE STUDIES PROGRESS REPORT FOR DECEMBER 1970, NO. 22. (Oak Ridge National Lab., Tenn.). Jan 1971. Contract W-7405-eng-26. 85p. Dep. NTIS.

CARBON—reactions with nitrogen and uranium dioxide

FISSION PRODUCTS—diffusion in oxide fuels

IODINE—volatilization from nitric acid solutions

—separation from fuel reprocessing off-gases

—separation from aqueous solutions

—spectra in alkaline and nitrate solutions

LIQUID METAL FAST BREEDER REACTOR—fuel cycle development studies (Dec 1970) on

METHANE, IODO—reactions with nitric acid

NITROGEN—reactions with carbon and uranium dioxide

PLUTONIUM OXIDES PuO₂—PuO₂—UO₂, solubility in nitric acid

—preparation from solutions from fuel reprocessing

—PuO₂—UO₂, preparation by sol-gel process

—PuO₂—UO₂, preparation of microspheres of

—PuO₂—UO₂, fabrication of sol-gel-derived

—PuO₂—UO₂, radiation effects on sol-gel-derived

—PuO₂—UO₂, diffusion of fission products in

REACTOR FUEL ELEMENTS—performance of clad oxide

—shipment of spent LMFBR, cask evaluation for

REACTOR FUELS—reprocessing of LMFBR, research progress (Dec 1970) on

—radiation effects on restructuring of ceramic, model for

SODIUM—reactions of, review on

STAINLESS STEEL—creep of tubing of

URANIUM OXIDES UO₂—PuO₂—UO₂, solubility in nitric acid

—preparation from solutions from fuel reprocessing

—reactions with carbon and nitrogen

—preparation of sols of, CUSP process for

—PuO₂—UO₂, preparation by sol-gel process

—PuO₂—UO₂, preparation of microspheres of

—PuO₂—UO₂, fabrication of sol-gel-derived

—PuO₂—UO₂, radiation effects on sol-gel-derived

—PuO₂—UO₂, diffusion of fission products in (M.C.G.)

45418 PROCESSING OF SOLIDS ON FLUIDIZED BEDS.

Bergougnou, M. A. (Univ. of Western Ontario, London). *Chim. Ind., Genie Chem.*; 105: No. 9, 559-71 (Apr 1972). (In French).

Fluidization has extensive applications in the processing of solids: coal distillation, coal combustion, incineration of organic residues, elimination of radioactive residuary slurries, ore reduction (iron ore in particular), ore enrichment, and high temperature electrothermal processes. The agricultural, medical and nuclear applications (fuel processing and reprocessing, coating of nuclear fuel particles with pyrolytic carbon) are described. General considerations on the calculation of fluidized units for the processing of solids are given. (France)

45419 NEW METHOD OF NUCLEAR FUEL REPROCESSING.

Perrot, Michel (Ugine-Kuhlmann, Paris). *Ind. At. Spatiales*; 26: No. 1, 15-24 (Jan 1972). (In French and English).

The reprocessing of irradiated fuels by the volatilization of

fluorides is reviewed after a brief description of the different types of fuel and possibilities in Europe. The advantages of this technique are described on the basis of experimental results and comparative economic studies. (France)

45420 PROCESSING OF IRRADIATED NUCLEAR FUELS.

Mills, Alfred Leonard; Vivers, Robert William Jackson (to United Kingdom Atomic Energy Authority). U. S. Patent 3,672,848. 27 Jun 1972. Filed 31 Dec 1969.

Irradiated monocarbide nuclear fuel is dissolved in nitric acid and the solution is subjected to the oxidizing effect of ozone or electrolytic oxidation to effect oxidation of organic compounds in the solution. Oxidation is terminated at the onset of the conversion of plutonium in the solution to the hexavalent state. (Official Gazette)

45421 CONTAINMENT OF FISSION PRODUCT IODINE IN THE REPROCESSING OF LMFBR FUELS BY PYROCHEMICAL REACTIONS. Krumpelt, M.; Heiberger, J. J.; Maroni, V. A.; Steindler, M. J. (Argonne National Lab., Ill.). *Nucl. Technol.*; 15: No. 3, 391-5 (Sep 1972).

The reaction of elemental iodine with liquid zinc and the reaction of ZnI₂ with molten chloride salt were studied. These reactions, which occur when fission product iodine is released into a zinc-salt melt during the pyrochemical decladding of fuel elements, appear to be very rapid. The mechanism is postulated to include conversion of zinc iodide to a zinc chloride complex. The presence of zinc chloride in the melt was verified by Raman spectroscopy. The results of laboratory-scale experiments suggest that the process for decladding LMFBR fuels in a zinc-salt melt may be capable of effectively retaining fission-product iodine in an easily disposable form. (auth)

45422 METHOD OF EXTRACTING FUEL FROM NUCLEAR FUEL ELEMENTS. Ballet, Paul; Faugeras, Pierre; Lefort, Guy (to Commissariat a l'Energie Atomique). French Patent 2,081,176. 3 Dec 1971. Filed 13 Mar 1970. (In French).

A method of extracting the fuel core from a nuclear fuel element with a metallic cladding is described. An electrochemical method is used to cut the cladding. HNO₃ is introduced under pressure into the cladding in order to leach the fuel. The fuel material is dissolved in the electrolyte (HNO₃) arising from the leaching by concentration in a dissolver-evaporator; the distilled electrolyte is then recycled in the evaporator. The extraction is effected by nitric acid leaching of the UO₂—PuO₂ fuel from the stainless steel-clad fast neutron reactor elements. (France)

45423 EXTRACTION OF FISSION PRODUCT NOBLE METALS FROM SPENT NUCLEAR FUELS. MacCragh, Adolfo (to W. R. Grace and Co.). U. S. Patent 3,672,875. 27 Jun 1972. Filed 5 Sep 1969.

A process for separating fission product metals from spent nuclear fuels, in which the insolubles from fuel element dissolution are heated in a stream of halide gas to convert the residues to the halides, is described. Ruthenium and rhodium fractions are converted to the complex haloruthenite and halorhodite, followed by dissolution and distillation to remove ruthenium. The rhodium is recovered by neutralizing the undistilled portion and reducing to the metal. Palladium and technetium are recovered from the halides volatilized in the first step of the process. (Official Gazette)

ENGINEERING

Energy Conversion

Refer also to abstracts: 45432, 45487, 45704, and 47170.

45424 (N-72-15607) CONCEPTUAL DESIGN STUDY OF A NUCLEAR BRAYTON TURBOALTERNATOR-COMPRESSOR. (General Electric Co., Cincinnati, Ohio). Nov 1971. Contract NAS3-13449. 526p. (NASA-CR-113925; GESP-493). NTIS.

A comprehensive analysis and conceptual design study of the turboalternator-compressor components using He-Xe as the working fluid was performed. The study was conducted in three phases: general configuration analysis (Phase 1), design variations (Phase 2), and conceptual design study (Phase 3). During the Phase 1 analysis, individual turbine, alternator, compressor, and bearing and seal designs were evaluated. Six turboalternator-compressor (TAC) configurations were completed. Phase 2 consisted of evaluating one selected Phase 1 TAC configuration to calculate its performance when operating under new cycle conditions, namely, one higher and one lower turbine inlet temperature and one case with krypton as the working fluid. Based on the Phase 1 and 2 results, a TAC configuration that incorporated a radial compressor, a radial turbine, a Lundell alternator, and gas bearings was selected. During Phase 3 a new layout of the TAC was prepared that reflects the cycle state points necessary to accommodate a zirconium hydride moderated reactor and a 400 Hz alternator. The final TAC design rotates at 24,000 rpm and produces 160 kW(e), 480 V, 3-phase, 400 hertz power. (auth) (STAR)

45425 (N-72-16724) EFFECT OF TWO TYPES OF HELIUM CIRCULATORS ON THE PERFORMANCE OF A SUBSONIC NUCLEAR-POWERED AIRPLANE. Strack, William C. (National Aeronautics and Space Administration, Cleveland, Ohio. Lewis Research Center). Mar 1971. 29p. (NASA-TM-X-2237; E-5811). NTIS.

Two types of helium circulators are analytically compared on the bases of their influence on airplane payload and on propulsion system variables. One type of circulator is driven by the turbofan engines with power takeoff shafting while the other, a turbocirculator, is powered by a turbine placed in the helium loop between the nuclear reactor and the helium-to-air heat exchangers inside the engines. Typical results show that the turbocirculator yields more payload for circulator efficiencies greater than 0.82. Optimum engine and heat exchanger temperatures and pressures are significantly lower in the turbocirculator case compared with the engine-driven circulator scheme. (auth) (STAR)

45426 COMPOSITE WALLS FOR MAGNETOHYDRODYNAMIC APPARATUS. Blattmann, Henri; Javellaud, Jean; Yerouchalmi, David (to Commissariat a l'Energie Atomique). French Patent 2,082,735. 15 Nov 1971. Filed 25 Mar 1970. (In French).

Improvements to the composite walls of MHD apparatus are presented. They are applicable to both the electrodes and the insulating walls. The composite wall of a MHD generator is constructed from an assembly of principal elements, each element being composed of several types of subelement. The principal elements are assembled using trapezoidal prismatic spacers made of insulating refractory ceramic material contained between two sheets of corrugated steel. The spacers are made of magnesite or strontium zirconate. The principal elements of these walls can be divided into several different parts by transversely positioned corrugated steel sheets. (France)

45427 THERMAL CONTROL AND POWER FLATTENING FOR RADIOISOTOPIC THERMODYNAMIC POWER SYSTEM. Bienert, Walter B.; Levedahl, William J.; Streb, Alan J. (to Teledyne, Inc.). U. S. Patent 3,672,443. 27 Jun 1972. Filed 28 Jan 1969.

A thermal control device utilizing the heat pipe principle is described. Heat receiving and heat rejecting surfaces are spaced from each other along the length of the heat pipe. In a preferred form, a radioisotope fuel capsule is in thermal conducting relationship with the heat receiving surface and a noncondensable gas is mixed with the working fluid of the heat pipe. A noncondensable gas automatically expands as the threshold temperature of the system drops to block off more of the heat rejecting surface. In other alternate preferred embodiments, a nonvaporizable liquid is utilized in lieu of a noncondensable gas to reduce the area of the heat receiving surface as the threshold temperature of the system drops or a second material is mixed with the working fluid which changes to a solid phase to block the flow of the working fluid between the heat rejecting and heat receiving surfaces

as the threshold temperature of the system drops. (Official Gazette)

45428 LASER POWER STATIONS IN ORBIT. Hansen, C. Frederick; Lee, George (Ames Research Center, Moffett Field, Calif.). Astronaut. Aeronaut.; 10: No. 7, 42-55 (Jul 1972).

Application of nuclear-fueled laser satellites to beam power to lunar bases and other spacecraft are proposed. Cycles for converting laser radiation to electric energy are considered along with electrostatic ion propulsion. It is noted that such vehicles would carry only power conversion systems, appropriate thrusters, and fuels. (J.R.D.)

45429 NEW CONCEPT IN ENERGY TRANSMISSION. Gregory, Derek P. (Inst. of Gas Tech., Chicago). Pub. Util. Fortn.; 89: No. 3, 21-9 (3 Feb 1972).

As the U. S. moves from a fossil fuel economy toward a nuclear power age to meet burgeoning energy demands, the problems of electricity transmission, distribution, and storage become accentuated. An alternative exists to the growing number of overhead power lines appearing around cities. Hydrogen is a synthetic fuel that could be made from water and electricity at the power stations and used as a means of underground energy transmission. It is an ideal fuel in many respects for combustion to obtain heat or for reconversion to electricity near the user. Economically, transmission of a gas is far cheaper than transmission of electric power, especially underground. This concept is not without its problems; nevertheless, it surely deserves detailed consideration in the future. (auth)

45430 INVESTIGATION OF OPTIMUM CONDITIONS FOR ENERGY CONVERSION IN MHD GENERATOR. Zaporowski, B. (Technical Univ., Poznan, Poland). Pr. Inst. Masz. Przemy.; No. 54-55, 77-84 (1971). (In Russian).

Investigation of the efficiency of heat conversion into electric energy in the MHD generator as a function of initial energy parameters of the plasma is described. There are given values of energy parameters and physical properties of the plasma obtained by combustion of conventional fuels in the oxidizer of variable O₂-content, and its variable physical enthalpy, account taken of the losses of energy for cooling the combustion chamber and duct as well as for introducing the ionizing seed and the like. An analysis of the influence of energy parameters of the plasma on the output obtained from the unit volume of the duct, varying along its axis is presented. A short description of a MHD generator having a thermal power of 4 MW, is included. (NSA of Poland)

45431 ELECTRICAL PARAMETERS IN THE HALL-TYPE MAGNETOHYDRODYNAMIC GENERATOR WITH GAS PROPERTY NONUNIFORMITIES IN THE DIRECTION OF THE MAGNETIC FIELD. Celinski, Z. N. (Inst. of Nuclear Research, Warsaw). Rozpr. Elektrotech.; 17: 109-23 (1971).

The effects of gas velocity and temperature distributions in the direction of a magnetic field on the electrical parameters of the Hall-type MHD generator were considered. The nonuniform distribution of gas velocity leads to the formation of circulating currents in the x and y directions. The Ohmic losses increase slightly. The terminal values of the generator do not change sufficiently in comparison to the constant gas property generator. The internal resistance is practically unchanged; the terminal voltage and the generated power increase slightly. The expressions for the electrical values were derived and a numerical example was calculated. (NSA of Poland)

45432 RADIATIONNAYA TEKHNIKA. VYPUSK 6. (Radiation Technology. Number 6). Shtan, A. S. (ed.). Moscow; Atomizdat (1971). 256p.

This collection of 43 articles by various authors presents theoretical and experimental work on isotope thermoelectric generators, irradiation equipment, pulsed proton generators, nuclear methods of analysis, and gamma defectoscopy. A few papers deal with radiotherapy and x-ray diagnostic apparatus. The production of isotopes using neutrons, neutron activation and neutron resonance analysis is discussed. (K.S.W.)

45433 CONVERSION OF HIGH-TEMPERATURE PLASMA ENERGY INTO ELECTRICAL ENERGY. Post, Richard F. (to United States Atomic Energy Commission). U. S. Patent 3,663,360. 16 May 1972. Filed 13 Aug 1970.

High temperature plasma is produced in a containment zone of a controlled fusion reactor defined by a magnetic containment field from which field plasma emerges as a beam. The plasma beam is directed through the magnetic field of an expander which

expander field is coupled to the containment field and diminishes in intensity therethrough so that rotational energy of the plasma ions is converted to translational kinetic energy. An arrangement is provided at the outlet of the expander to separate electrons from the high energy ions which ions are then directed into a collector structure. The collector structure comprises a plurality of stages maintained at progressively higher positive potentials wherein the ions are progressively decelerated so that successive fractions of the ions pass residual kinetic energies less than the differential potential between successive collector stages. The collector stages are provided with a deflector means which selectively deflects the ions with low residual kinetic energies to impinge upon collector plates to create electrical current at the stage potential. In one disclosed embodiment, the selective deflector means utilizes a vapor jet charge exchange arrangement, in a second, the selective deflector means is an arrangement using crossed electric and magnetic fields, and, in a third, the selective deflector means utilizes particle focusing lenses. Electrical circuitry means are provided for combining and/or converting the individual collector currents for delivery to an electrical distribution system with means to establish the initial decelerating or retarding potentials. (auth)

45434 ELECTROCHEMICAL POWER-PRODUCING CELL. Cairns, Elton J.; Chilenskas, Albert A.; Steunenberg, Robert K.; Shimotake, Hiroshi (to United States Atomic Energy Commission). U. S. Patent 3,666,560. 30 May 1972. Filed 21 Sep 1970.

An electrochemical power-producing cell is described including a molten lithium metal anode, a molten selenium metal cathode, a paste electrolyte separating the anode from the cathode, an anode current collector, and a single layer of niobium expanded metal formed in corrugated shape as cathode current collector. In addition, means are provided for sealing the anode and the cathode from loss of lithium and selenium respectively. An insulator is provided between the anode housing and the paste electrolyte disk. (Official Gazette)

Facilities and Equipment

Refer also to abstract: 47300.

45435 (CEA-CONF-2008) HEAT PIPES. DESIGN AND INDUSTRIAL APPLICATIONS. Semeria, R. (Commissariat a l'Energie Atomique, Grenoble (France). Centre d'Etudes Nucleaires). 1971. 16p. (In French). (CONF-710651-3). Dep. NTIS (U. S. Sales Only).

From International conference on heat exchangers; Paris, France (16 Jun 1971).

Heat pipes are thermosiphons with vaporization where we can distinguish a boiler, a condenser, and eventually an adiabatic zone. To insure the returning liquid flow from the condenser to the boiler, surface tension forces, associated with the gravity forces, if need be, are used. For this, the condensing liquid is sucked by a capillary structure, generally situated against the inner wall. The review of the design methods, and particularly the prediction of the maximal performances shows the advantages and limitations of such devices. The main difficulties are technological for the heat pipes with high temperature liquid metals. The thermohydrodynamic limitations are: the maximum power (which can be calculated by a balance between friction forces and active forces), the maximum heat flux leading to the dry-out of the evaporator, and the critical conditions for the start up associated with the sonic conditions in the vapor phase. Heat pipes designed for industrial applications (mainly for space) are described. (auth)

45436 (CEA-CONF-2010) DESIGN CONSIDERATIONS OF HEAT EXCHANGERS FOR VAPORIZING FLUIDS. Fabrega, S.; Huyghe, J.; Ricque, R. (Commissariat a l'Energie Atomique, Grenoble (France). Centre d'Etudes Nucleaires). 1971. 13p. (In French). (CONF-710651-2). Dep. NTIS (U. S. Sales Only).

From International conference on heat exchangers; Paris, France (16 Jun 1971).

Aspects of determining the steady-state operating characteristics of certain types of vapor generating equipment are examined. A type of equipment considered is the vertical tube reboiler with upward flow. A calculation method applicable to industrial mixtures is described, wherein the evolution of mixture composition along the tube is accounted for. Based on results obtained with an experimental unit of this type, an attempt was made to develop correlations for the pressure drop and the heat transfer rate that would apply at relatively low pressures (a few bars). Also discussed are falling film, long tube vertical evaporators, and a boiling system often found in industrial situations. Methods and results are presented that permit predictions of the heat transfer coefficient and the pressure drop. For the case of horizontal, once-through vapor generators at high pressures, the influence of gravity

leads to non-symmetrical flow and heat transfer patterns inside the tubes. First results with Freon are presented to permit an analysis of this asymmetry, and in addition, to permit examination of the effect of elbows. Aspects of boiling outside horizontal or vertical tubes are considered in connection with the operation of low-pressure boilers often encountered in seawater desalting units (sub-atmospheric pressures) and in the chemical industry. (auth)

45437 (CONF-720904-1) DESIGN OF HIGH-TEMPERATURE LIQUID METAL SYSTEMS. Fraas, A. P. (Oak Ridge National Lab., Tenn.). 13 Sep 1972. 27p. Dep. NTIS.

From Conference on the impact of aerospace technology on society in the 70's; Los Angeles, CA. (11 Sep 1972).

A review of over two million hours of high-temperature liquid metal system operating experience gained under the Aircraft Nuclear Propulsion and Nuclear Electric Space Power Plant programs at ORNL during the past 22 years may prove helpful to those who have occasion to design high-temperature liquid metal systems for other applications. The nature of the problems involved in material selection, mass transfer and corrosion, heat transfer and heat transport, limitations imposed by oxidation of external surfaces of a system, the performance characteristics of various liquid metals relative to other heat transfer media such as high-pressure helium and various molten salts, thermal stress and thermal strain cycling, and problems posed by the fire hazard potential all play obviously important roles in the design of high-temperature liquid metal systems. In addition, experience has shown that liquid metal systems are singularly unforgiving of small errors in design, shakedown, and operation. This in turn means that much attention must be devoted to the detail design of components to provide a high degree of leak tightness, reasonable freedom from thermal strain cycling, and a host of special problems such as self welding. A set of design precepts for liquid metal systems is presented along with selected references as an aid to those interested in this field. (auth)

45438 (KFK-1556) CONCEPT OF AN IRRADIATION FACILITY FOR INVESTIGATION OF HIGH-TEMPERATURE THERMOELEMENTS AT TEMPERATURES TO 2000°C. Piel, D. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Material und Festkoerperforschung). Mar 1972. 29p. (In German). Dep. NTIS (U. S. Sales Only).

In addition to general indications with respect to W/Re thermocouples and the aims pursued in a high-temperature irradiation program an irradiation rig is described. This rig may be used to investigate the behavior of high-temperature thermocouples in the radiation field of a reactor, using a gas thermometer. The most important instability effects of thermocouples are discussed and given as experimental parameters. The irradiation capsule has been designed for insertion into the Br2 Reactor at Mol and for temperatures of about 2000°C. The individual heat transport mechanisms are described which underly the calculation of temperature distribution. (auth)

45439 (KFK-1566) RECUPERATIVE HEAT EXCHANGER FOR A FAST HELIUM-COOLED BREEDER REACTOR WITH HELIUM TURBINE. Savatteri, C. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Neutronenphysik und Reaktor-technik). Feb 1972. 32p. (In German). (EUR-4725). Dep. NTIS (U. S. Sales Only).

In a plant operated by a gas-cooled fast breeder reactor and a gas turbine the amount of heat transferred in the recuperative heat exchanger is almost identical to that in the reactor. Since the inlet temperatures of the turbine and the condenser are also fixed in advance, the efficiency of the process is decisively dependent on the heat exchanger. In order to obtain guide lines for the final design of the recuperator of a 1000 MWe plant, preliminary calculations of variations are carried out. This work investigates the influence of various parameters (tube diameter, pitch ratio, loss of pressure distribution) on the size of the heat exchanger. For better understanding of the problems the plant is briefly described. (auth)

45440 (MSAR-71-45) ENTRAINMENT MOISTURE SEPARATORS FOR FINE (1-10 μ) WATER-AIR-STEAM SERVICE: THEIR PERFORMANCE, DEVELOPMENT, AND STATUS. Final Report. Griwatz, Gunther H.; Friel, Joseph V.; Bicehouse, Jack L. (MSA Research Corp., Evans City, Pa.). Mar 1971. Contract AT(45-1)-2145. 163p. Dep. NTIS.

There is a need for moisture separators that are at least 99% efficient in removing entrained water particles in the 1 to 10 micron (μ) range from air streams. Five commercially available entrained moisture separators were tested for performance in the 1 to 10 micron particle-size range. The MSA Type G and the AAF Type T separators were greater than 99% efficient in removing particles in this range from atmospheric to PWR post-accident

(incident) conditions. The York Type 321 SR separator failed to contain the separated water within the drains of the unit and re-entrainment occurred. The Farr Type 68-44ZH and the Monsanto baffle-type separators were inferior because of visible and measured penetration. MSA test details and results are summarized. Survey results, presenting the status of entrainment separators and media for 1 to 10 micron service, together with methods of measuring and generating particles in this range, have been previously reported. (auth)

45441 (SRARI-P-120) REALIZATION OF SYNCHRONOUS ROTATION OF SOME ROTORS SUSPENDED IN A MAGNETIC FIELD. Rukolaine, G. V.; Kalebin, S. M.; Pelevin, G. A.; Nikol'skii, S. N. (Scientific Research Atomic Reactor Inst., Melekes (USSR)). 1971. 24p. (In Russian). Dep. NTIS (U. S. Sales Only).

A method for the synchronous rotation of any number of rotors suspended in a magnetic field is considered. The advantage is taken of such distinctive features of suspension in a magnetic field as low friction during rotation of the rotors, and the absence of mechanical effects on the angular rate of rotation. The operational principle of electronic tracing circuits made for specified synchronization is described. The following technical characteristics are available in the arrangement developed: the rotation period of the rotors suspended in a magnetic field varies no more than 0.25 μ s, and their synchronous rotation measured at the rate of 6000 rpm is performed accurately to $\pm 0.04^\circ$. (auth)

45442 DETERMINATION OF THE PERFORMANCE OF STEAM SEPARATORS WITH THE AID OF THE LAWS OF SIMILITUDE. Brandes, H. (Technische Univ., Hanover). Kerntechnik; 14: No. 5, 210-19 (May 1972) (In German and English).

In modern high-performance steam generators, especially those of nuclear design, a sufficient separation of the generated steam/water mixture cannot be obtained by gravitational means alone. Therefore, not only steam dryers but also so-called steam separators, usually of the centrifugal type, are installed to ensure a low steam content (carryunder) in the water returning to the heating section of the steam generator. A low carryunder is necessary to stabilize the flow and to ensure a good heat transfer in the heating section. An investigation of separators of a centrifugal type was performed to obtain empirical relations for the calculation of pressure drop and carryunder in future designs of similar separators. The laws of similitude were applied to analyze the test data. It is shown that, by a dimensional analysis, the pressure drop of the tested separator type can be satisfactorily predetermined whereas calculations of carryunder will yield only approximate results. (auth)

45443 DYNAMIC RESPONSE OF A PLATE TO TIME-DEPENDENT BOUNDARY CONDITIONS. Ramachandran, J. (Indian Inst. of Tech., Madras). Nucl. Eng. Des.; 21: No. 3, 339-49 (1972).

The dynamic response of a simply supported rectangular plate to time-dependent boundary conditions is considered. Results are presented for the cases when (1) any one of the boundaries and (2) any two opposite boundaries are subject to a certain prescribed time-dependent displacement. The analysis is extended to a case where the plate carries a concentrated mass situated at some point within the domain. Numerical examples are given for all the cases considered. (auth)

45444 DISTRIBUTION OF A RADIOACTIVE AEROSOL IN A MODEL CHAMBER. Zalmanzon, Yu. E.; Nikitin, V. I.; Chutkin, O. A. (Scientific Research Inst. for Equipment Construction, Moscow). Gig. Sanit.; No. 3, 47-51 (Mar 1972). (In Russian).

A chamber is described which can be filled with a radioactive aerosol which is generated by bubbling pure air through a solution of a salt of a radioisotope. Working with NH_4Br -labeled the beta-emitter ^{82}Br , the aerosol generated had an average activity of 2×10^{-12} to 5×10^{-14} Ci/liter. Samples were taken from the corners and other parts of the chamber in order to determine the effectiveness of mixing. The isotope concentration showed a vertical variation of from 20 to 90%. (K.S.W.)

45445 STRESS DISTRIBUTION IN NOZZLE-TO-CYLINDER CONNECTIONS FOR SMALL DIAMETER RATIOS. Van Campen, D. H.; Spaas, H. A. C. M. (Delft Univ. of Tech., Netherlands). Nucl. Eng. Des.; 21: No. 3, 368-95 (1972).

Development of a finite element computational method is described, which is intended for determining the detailed distribution of strain and stress (inclusive the peak stresses) in nozzle-to-cylinder intersections for sufficiently small d/D ratios (typically $d/D \leq 1/4$). The proposed method can be considered as an extension of the computational method for the biaxially loaded nozzle-to-flat plate connection presented previously, which is based upon the finite element method for solids of revolution under non-axisym-

metric loadings. The two computational methods (nozzle-to-flat plate and nozzle-to-cylindrical vessel) are compared with each other and the reliability of both is evaluated by comparison with experimental evidence. (auth)

45446 POOL IRRADIATOR WITH RADIATION SOURCE IN A ROTATING CONTAINER. Ransohoff, Jackson A. (to Neutron Products Inc.). U. S. Patent 3,665,183. 23 May 1972. Filed 28 May 1969.

An irradiator operable with a radiation source under the surface of a pool of water includes a gas filled space within the irradiator and between the irradiation source and various positions in which containers of material are irradiated. The various positions are at preselected distances from the source to allow simultaneous experiments and the gas filled space conserves radiation. The radiation source may be a plurality of rods within tubes arranged circumferentially to one of the containers of material to be irradiated and the containers of material to be irradiated may be rotated during irradiation. (Official Gazette)

45447 PRODUCTION IRRADIATOR. Ransohoff, Jackson A.; Sarkozi, Lewis P. (to Neutron Products, Inc.). U. S. Patent 3,676,675. 11 Jul 1972. Filed 29 May 1969.

Irradiation of material in packages is carried out continuously by a unique conveyor arrangement which transports closely spaced carriers containing the packages in vertical runs past a radiation source. The radiation conveyor utilizes extension arms to increase the radius of outside turns and allow the carriers to be spaced closer to the source and to each other. A transfer conveyor is provided to obviate the need for inside turns so that individual carriers and adjacent vertical runs may be closely spaced in the radiation zone. A loading station provides means for the horizontal loading and unloading of the packages in increments to provide for the integration of the radiation dose horizontally across the carrier length. An effective labyrinth radiation shield above the source shields the loading station. The radiation dosage per pass through the system may be varied by varying the speed of the conveyor or the strength of the radiation source plaque. (Official Gazette)

45448 APPARATUS FOR IRRADIATION IN A CONTROLLED ATMOSPHERE. Coleman, George E. (to PPG Industries, Inc.). U. S. Patent 3,676,673. 11 Jul 1972. Filed 18 Aug 1969.

Apparatus which permits treatment of products with ionizing radiation in a controlled atmosphere comprises an inner chamber positioned within an outer chamber. The workpiece to be treated passes through the inner chamber. The inner chamber has a radiation-permeable window in one surface and the second chamber has means for the passage of radiation in juxtaposition with the window in the outer chamber. The apparatus may also contain means for carrying the product to be treated, as well as cooling means at the end of the radiation path. (Official Gazette)

45449 METHOD OF FABRICATING A HEAT PIPE. Werner, Richard W.; Alexander, Everette E.; Comstock, Irving J. (to United States Atomic Energy Commission). U. S. Patent 3,665,573. 30 May 1972. Filed 18 May 1970.

A tube is provided having a plurality of closely spaced very thin longitudinal fins projecting into its interior. The tube is particularly useful as a heat pipe, the capillaries defined between adjacent fins serving as a wicking structure for conveying a heat transfer working fluid from one region of the pipe to another by capillary pumping. Fabrication of the internally finned tube is accomplished by a sacrificial mandrel method. (Official Gazette)

45450 SODIUM HEATED STEAM GENERATOR. Creek, Ronald B.; Gardner, Karl A. (to United States Atomic Energy Commission). U. S. Patent 3,662,718. 16 May 1972. Filed 2 Mar 1971.

A sodium-heated steam generator is described having a tube sheet which is subject only to saturated or near saturated steam temperatures. Double-walled evaporative and superheated steam tubes extend through said tube sheet into the liquid metal. In both the evaporative and superheated steam tubes the saturated steam passes through the annular passageways which contact the tube sheet. (Official Gazette)

45451 MIXER-SETTLER APPARATUS. Miller, William E.; Knighton, James B.; Bernstein, George J. (to United States Atomic Energy Commission). U. S. Patent 3,663,178. 16 May 1972. Filed 3 Jun 1969.

An apparatus is described for simultaneously mixing and separating multiple phase fluids in which one phase is retained as a batch phase and the other phases are continuously flowing. A pump-mixer continuously mixes the fluids and pumps them into a settling chamber where they settle out and means are provided in the settling chamber for returning one phase to the mixer from which it came and transferring the other phases to another pump-mixer. Apparatus is provided for retaining as a batch or captive phase either heavy or light phases. (Official Gazette)

45452 DENITRATOR 5 IN (29 Nov 1971) (Engineering Materials). (Union Carbide Corp., Oak Ridge, Tenn. Y-12 Plant). (CAPE-724).

7 drawings.

This denitrator is a scaled-down and modified version of a denitrator developed at Hanford. It is used to convert uranyl nitrate ($\text{UO}_2(\text{NO}_3)_2$) to uranium trioxide (UO_3). (For details see TID-4100 Suppl. 72) (F.S.)

45453 CONSISTENT THEORY FOR SPHERICAL SHELLS IN EQUILIBRIUM. Palusamy, S. (Sir George Williams Univ., Montreal); Lind, N. C. Nucl. Eng. Des.; 21: No. 3, 350-7 (1972).

A consistent theory in the sense that the displacement and stress fields not only satisfy respectively the kinematic and static requirements, but also the requirements of the principle of virtual work, is presented for a thin spherical shell in equilibrium; the results are useful in plastic limit analysis where this consistency is a prerequisite for the application of the theorems of limit analysis. (auth)

45454 DEVELOPMENT OF 50Hz 44 INCH TURBINE BLADE. Okubo, Atsuo (Takasago Machinery Works, Japan); Kadoya, Yoshiki; Sato, Tomohiko; Umemura, Sunao; Kuramoto, Yasuo. Tech. Rev., Mitsubishi Heavy Ind.; 9: No. 1, 21-30 (Feb 1972).

A 50-Hz, 44-inch low pressure end turbine blade was developed to build 1,000-MW class fossil- and nuclear-fueled power plants for reliability and high efficiency. Optical and traverse tests in high-speed cascade wind tunnel were carried out to determine the cascade performance of the blade, resulting in very low cascade losses. Rotating vibration tests using actual blades and disk revealed the blade had excellent vibratory characteristics. The design features and results of high-speed cascade test and rotating vibration test for the blade are described. (auth)

45455 MANUFACTURING DETAILS AND MATERIAL USED FOR MANUFACTURE OF THROTTLE, CONTROL, INTERCEPTOR, AND STOP VALVES FOR HIGH RATING STEAM TURBINES OF CONVENTIONAL AND NUCLEAR TYPE. Celaschi, Emilio (Franco Tosi SpA, Legnano, Italy). Termotecnica (Milan); 26: No. 5, 243-8 (May 1972). (In Italian).

The evolution of throttle, control, interceptor, and stop valves for 160, 320, and 600 MW conventional turbogenerator units is described. Descriptions of nuclear stations with outputs at 300 to 1200 MW are also included. Improved turbogenerator reliability is noted as a result of recent developments. (J.R.D.)

45456 IRRADIATOR APPARATUS WITH A FLUID FLOW-PATH DETERMINED BY COMMUNICATING CORE AND CASTING BAFFLES. Woodbridge, David D.; Mann, Leland A. (to Energy Systems, Inc.). U. S. Patent 3,671,741. 20 Jun 1972. Filed 12 Jan 1970.

An irradiator apparatus is described which is designed to perform a sterilization treatment or like process on waste material especially in the liquid state such as sewage or general industrial waste wherein a radiation source in the form of radioactive material is utilized to eliminate various types of micro-organisms or other undesirable elements which are carried by the liquid waste material. The apparatus includes a structure comprising a plurality of parallel baffles arranged about a substantially centrally located core mounted within the casing and structurally designed to support a plurality of radioactive material elements serving as the radiation source. The core also includes a plurality of parallel baffles which communicate with the baffles of the casing to define a somewhat serpentine flowpath along which the liquid travels as it passes through the irradiator. (Official Gazette)

45457 IRRADIATION APPARATUS FOR PROVIDING A HIGH INITIAL IRRADIATION OF THE PRODUCT. Green, William R. (to Atomic Energy of Canada Ltd.). U. S. Patent 3,673,409. 27 Jun 1972. Priority date 6 Feb 1970, Canada.

Apparatus is described for the radioactive irradiation of products whose composition requires a relatively high intensity of irradiation for a short period followed by a relatively low intensity of irradiation for a longer period. In order to efficiently utilize the source, the product is passed into the center of a rack of source material and thereafter passed around the outside of the rack. The product is cooled by water during irradiation and the same water is used to effect rotation of the product. (Official Gazette)

45458 (CISE-R-245) PDE: DIGITAL PROGRAM FOR THE CALCULATION OF THE THERMAL-HYDRODYNAMIC CONDITION IN A CIRCUIT OF A TWO PHASE MIXTURE WATER AND STEAM. Muscettola, M.; Barbieri, G. (Centro Informazioni Studi Esperienze, Milan (Italy)). Oct 1967. 175p. (In Italian). Dep. NTIS (U. S. Sales Only).

The digital program PDE (Pressure, Density, Energy) is described. It has been set up to compute the pressure conditions, the densities and the energy and mass content in a boiler. The hydraulic schemes which can be considered by the program are indicated. In input the program requires groups of data such as the constants of experimental correlations for pressure drops and steam volume fraction: the geometric characteristics and dimensions of the circuit through the number of zones, risers per zone, channels per riser, the number of cells, and for each cell its length, flow area, hydraulic diameter, direction of the flow, type of geometry, and type of discontinuity at the outlet extreme; and the thermal and hydraulic conditions imposed, i.e., the drum pressure and for each channel the mass flow rate, the inlet enthalpy (below as well as above saturation), and the thermal power transferred to the fluid specifying both the magnitude and the shape of the distribution along the channel (constant, sinusoidal and defined in a discrete number of points). Program output consists of data on the pressure, the steam quality, volume fraction, and the density and enthalpy of the mixture at the inlet and at the outlet of each cell; for the heated channels, the mass flow rates which would determine the thermal-crisis at the outlet of each cell; for each section of the circuit (i.e., risers, unheated and heated channel) and for each zone, the mean density, the mass, mean energy, and total energy of the mixture and the internal volume of the circuit; for the heated section of the channel of each zone, the mean density in every cell and the mean density weighted on the square of the heat flux, the mean being taken over the whole heated section of the channel; for the whole circuit the mean density weighted on the square of the heat flux, the total mass and energy of the fluid, and the total volume of the circuit. The program is written in FORTRAN IV for IBM 7040. (auth)

45459 (CISE-R-275) THRESHOLD OF PARALLEL CHANNEL FLOW INSTABILITY. Masini, G.; Possa, G.; Tacconi, F. A. (Centro Informazioni Studi Esperienze, Milan (Italy)). Sep 1968. 57p. (In Italian). Dep. NTIS (U. S. Sales Only).

In the framework of the CIRENE Project, an experimental program was conducted by CISE on parallel channel flow instability. A forced convection open loop circuit was built. The circuit is composed of two parallel channels, simulating at a reduced scale the reactor power channels. Flow instability threshold data are presented and briefly analyzed. The mechanism of flow instability is clearly of a dynamic nature. (auth)

45460 VAPOR TRANSPORT LIMITS OF LIQUID METAL HEAT PIPES. DeMichele, Don W. (Texas A and M Univ., College Station); Davis, Monte V. Nucl. Technol.; 15: No. 3, 366-83 (Sep 1972).

The mass flow choking limits have been calculated for fourteen potential working fluids in cylindrical heat pipes. The problem is analyzed with the vapor assumed to be a perfect gas and then as a compressible gas with viscous flow and both results compared to measured heat pipe performance. Based on the measured data the analysis has been extended to other potential working fluids and the results are tabulated. (auth)

45461 EXPLOSIVE INTERACTION OF MOLTEN METALS INJECTED INTO WATER. Bradley, R. H.; Witte, L. C. (Univ. of Houston, Tex.). Nucl. Sci. Eng.; 48: No. 4, 387-96 (Aug 1972).

A high-speed photographic investigation established the general nature of the explosive interaction between hot molten metal jets and subcooled (75°F) distilled water. The photographic sequences yielded basic information concerning the relationship between extent of material fragmentation and the intensity of explosive action. Prior theories for the explosive causative mechanism were shown to be inadequate to explain the observed explosive action. A thermally controlled initiating mechanism is hypothesized and the experimental evidence supports the hypothesis. Dimensional analysis yields a parameter dominated by the thermal characteristics of the jet system. (auth)

45462 EFFECT OF LATERAL ROD DISPLACEMENT ON LAMINAR-FLOW TRANSFER. Hsu, Chia-jung (Brookhaven National Lab., Upton, N. Y.). J. Heat Transfer; 94: No. 2, 169-73 (May 1972).

Heat transfer in longitudinal laminar flow through tightly packed rod bundles is analyzed for a rod displaced from its symmetrical position. The influences of the pitch/diameter (P/D) ratio, the magnitude of displacement/ $d_{\max}$ (σ), and the type of thermal boundary condition on the variation of the coolant temperature field and the rod-average Nusselt number

Heat Transfer and Fluid Flow

Refer also to abstracts: 45427, 45435-45437, 45439, 45442, 45449, 45450, and 45721.

are investigated. Average Nusselt numbers for the displaced rod are calculated for rod bundles with $1.00 \leq P/D \leq 2.0$ and $0 \leq \sigma \leq 0.8$. The results for the special case of $\sigma = 0$ (i.e., for symmetrical rod bundle) show excellent agreement with those reported by Dwyer and Berry, who analyzed numerically the corresponding problem for symmetrical rod bundles. (auth)

45463 RECENT DEVELOPMENTS IN THERMOPHYSICAL PROPERTIES RESEARCH. Winter Annual Meeting of the American Society of Mechanical Engineers, Washington, D. C., December 1, 1971. Cezairliyan, A. (ed.). HTD Vol. 3. New York; American Society of Mechanical Engineers (1971). 18p. \$3.50. (CONF-711218).

Papers are included in which discussions are presented on thermal conductivity, thermal diffusivity, thermal expansion, and thermal radiation properties. Other thermophysical and related properties, such as specific heat, melting point, and electrical resistivity, are also reviewed in connection with discussions of technique. (J.R.D.)

45464 EXPERIMENTAL INVESTIGATION OF HEAT TRANSFER FROM PARTIALLY IONIZED ARGON WITH AN APPLIED TRANSVERSE MAGNETIC FIELD. Roschke, E. J. (California Inst. of Tech., Pasadena). J. Heat Transfer; 94: No. 2, 174-80 (May 1972).

Wall heat transfer measurements were obtained for laminar flow of partially ionized argon flowing within the conducting walls of a square channel, with and without an applied transverse magnetic field. Tests were conducted for subsonic flows and for flows which were supersonic before a magnetic field was applied. Increases in Stanton number by a factor of as much as six were observed at field strengths approaching 10 kG as compared with values at zero magnetic field. These large increases in heat transfer are believed to have been due to (1) a small amount of joule heating augmented or accompanied by (2) magnetically induced ionization. Heat transfer and flow data were used to estimate effective values of the joule heating parameter, Hall coefficient, and current density. The experimental data have been compared with theoretical predictions for several limiting cases. (auth)

45465 IMPROVED POINT-MATCHING TECHNIQUES APPLIED TO MULTI-REGION HEAT TRANSFER PROBLEMS. France, D. M.; Ginsberg, T. (Argonne National Lab., Ill.). J. Heat Transfer; 94: No. 2, 203-10 (May 1972).

An analytical method is presented which extends the series solution of the Laplace and Poisson equations with irregular boundary conditions to multi-cell problems. The method employs a least-squares technique of satisfying the boundary conditions on the irregular boundaries and eliminates the use of a finite number of boundary points to satisfy these conditions. The technique is applied to the calculation of the fully developed temperature distribution of a constant-velocity fluid flowing parallel to a semi-infinite square array of circular nuclear fuel rods. The bounding wall of the array is located such that the flow area of the cell associated with the rod adjacent to the wall is different from the (equal) areas of all the other cells. The series solution is compared with a finite-difference solution for a sample case of two cells. The results for the semi-infinite array indicate that while the array temperature distribution is markedly affected by the difference in flow areas, the Nusselt numbers of the rods are relatively unaffected. Typical results are presented for a pitch-to-diameter of 1.2; the flow area of the first cell is 3.67 percent greater than the area of the other cells. (auth)

Materials Testing

Refer also to abstract: 45432.

45466 (AE-451) DEPTH DISTRIBUTION STUDIES OF CARBON IN STEEL SURFACES BY MEANS OF CHARGED PARTICLE ACTIVATION ANALYSIS WITH AN ACCOUNT OF HEAT AND DIFFUSION EFFECTS IN THE SAMPLE. Brune, D.; Lorenzen, J.; Witalis, E. (Aktiebolaget Atomenergi, Studsvik (Sweden)). 1972. 47p. Dep. NTIS (U. S. Sales Only).

Depth distribution studies of carbon in steel and iron were carried out in the concentration range 0.05 to 1%, using proton activation analysis. Surface content studies were performed in the concentration range 0.01 to 1% using deuteron activation analysis. The following reactions were utilized: $^{12}\text{C}(p,\gamma)^{13}\text{N}$ and $^{12}\text{C}(d,n)^{13}\text{N}$. Evaluations of depth distribution were based on resonances in the excitation function. The carbon content was determined with the aid of the positron emitter, ^{13}N , using either single-peak or coincidence measurements. The heat dissipation in the irradiated region of the samples was calculated, and the temperature rise was measured using thermocouples. The tem-

perature distribution within the hot zone subjected to irradiation by charged particles, together with the temperature distribution around this zone, was studied in order to estimate any effect this might have on the carbon diffusion. A device for automatic sample exchange which is remotely controlled is described. (auth)

45467 (BNWL-SA-4320) ^{252}Cf NEUTRON RADIOGRAPHY AT BATTELLE-NORTHWEST. Swinth, K. L. (Battelle-Northwest, Richland, Wash. Pacific Northwest Lab.). [nd]. 4p. (CONF-720902-1). Dep. NTIS.

From National topical meeting on the applications of californium-252; Austin, TX. (11 Sep 1972).

Work devoted to development and use of ^{252}Cf in neutron radiography is described. Discussions are included on exposure facilities, imaging techniques, and applications. (J.R.D.)

45468 SENSITOMETRIC MEASUREMENTS OF NON-SCREEN INDUSTRIAL FILMS EXPOSED BY X OR GAMMA RADIATION. Smoniewski, J. Pomiar, Automat., Kontr.; 17: 202-3 (1971). (In Polish).

The quality of a photographic image obtained by exposure of radiographic films with the aid of X or gamma radiation in industrial flaw detection technique has an important effect on the information about a part under test. This quality being dependent upon the quality of radiation, the effective energy of 85, 125 and 660 keV representing the range of practically applied exposures is selected for testing. A method of sensitometric measurement is described and numerical values of radiosensitivity and gamma indices are given. These data should facilitate the correct choice of the films and also proper exposure thereof. (NSA of Poland) (auth)

45469 ULTRASONIC FLAW DETERMINATION BY SPECTRAL ANALYSIS. Adler, Laszlo; Whaley, Hubert L. (to United States Atomic Energy Commission). U. S. Patent 3,662,589. 16 May 1972. Filed 2 Feb 1971.

A method of quantitatively determining the size and orientation of flaws within a material is given. A broadband pulse having a frequency width of at least several MHz is used for an input pulse from a transducer which is directed toward the test material which may contain a flaw, and the average frequency interval between interference maxima in the reflected spectrum is related to both the size and orientation of the flaw. An experimental method using two known angle changes allows determination of the size and angle of randomly oriented discontinuities within metal specimens. (Official Gazette)

45470 NONDESTRUCTIVE TEST FOR MEASURING THE STATE OF HEAT TREATMENT IN CLOSURE WELDS. Dodd, C. V.; Lu, C. C. (Oak Ridge National Lab., Tenn.). Mater. Eval.; 30: No. 7, 142-8 (Jul 1972).

The first-stage development on a nondestructive testing technique to measure the properties of resistance-welded closures on chemical munition containers has been completed. A preliminary technique using a magnetic circuit has been developed that measures temper of the weld, cross-sectional area of the weld, and variation of cap height from nominal. Measurements of magnetic properties as a function of temper and area were made of samples of AISI Type 1040 steel. A prototype probe to induce the magnetic circuit, link the magnetic circuit through the weld and measure the magnetic flux through the circuit was designed and constructed. Tests performed on six sample welds demonstrated the capability for measuring the temper, area, and variations in cap height. Recommendations for improving the accuracy, speed, and reliability of the measurements are given. (auth)

45471 FACTORS GOVERNING RADIOGRAPHIC CRACK DETECTABILITY IN STEEL WELD SPECIMENS. Becker, George L. (E. I. Du Pont de Nemours and Co., Inc., Wilmington, Del.). Mater. Eval.; 30: No. 7, 149-52 (Jul 1972).

The effects of energy level and radiation source position changes on crack detectability were evaluated on a crack in a steel weld. The weld was step-machined to present various crack depths on the radiographs. Steel plates were added progressively beneath this cracked wedge until the crack image was no longer discernible on the film. Both x and gamma radiation were used, and standard image quality indicators were added to correlate crack detectability with indicator sensitivity. Detectability parameters were established for the various energies and positions used. (auth)

45472 ULTRASOUND QUALITY CONTROL INSTALLATION. Boulanger, Gilbert; Samoeil, Alain (to Commissariat a l'Energie Atomique). French Patent 2,088,972. 13 Dec 1971. Filed 30 Apr 1970. (In French).

An apparatus is presented for the quality control of metallic elements (in particular the fuel cans of nuclear fuels) by ultrasound. It consists of a non-sealed vessel in which the water level is maintained constant with the aid of a feed device de-

signed to avoid bubbles of air and turbulence in the vessel. The measurement vessel is of parallelepiped form and separated into three adjacent compartments by two parallel partitions. The two partitions and the sides of the vessel are provided with orifices which face one another and allow the passage of the fuel cans. The installation also possesses a means of moving the fuel cans, measurement transducers which are attached to the opposite faces of the vessel in the central compartment, and a system for feeding water into the vessel. (France)

45473 POSSIBILITIES OF USING RADIOISOTOPE INSTRUMENTS IN FABRICATION OF HIGH POWER PLANTS OF SUPER-CRITICAL STEAM PARAMETERS. Lonski, E. D.; Zorchev, S. N.; Volkov, O. S.; (and others). *Izotopy SSSR*; No. 22, 14-17 (1971). (In Russian).

The possibility of employing radioisotope methods for the inspection of welded joints of boilers with capacities of 300 000, 500 000, and more kW is considered. (tr-auth)

Nuclear Explosions

Refer also to abstracts: 45542, 45571, 45628, and 45674.

45474 (JAB-99-88) OBSERVATIONS OF CHIMNEY DAMAGE TO RESIDENTIAL BUILDINGS IN GLENDALE, CALIFORNIA; SAN FERNANDO EARTHQUAKE, FEBRUARY 1971. Farhoomand, I.; Scholl, R. E. (Blume (John A.) and Associates Research Div., San Francisco, Calif.). May 1972. Contract AT(26-1)-99. 36p. Dep. NTIS.

Statistical observations of residential chimney damage, resulting from the San Fernando earthquake, for a selected control area within Glendale, California, are summarized. The procedures used in performing a structure inventory and damage survey are described. General statistical observations are presented, the variation of damage for different chimney types is examined, and suggestions are made for increasing chimney resistance to ground motion disturbances. (auth)

45475 (NP-19361) DESIGN OF A WATERWAY CONNECTING THE ORINOCO AND RIO NEGRO RIVERS IN THE FEDERAL TERRITORY OF AMAZONAS, VENEZUELA. Technical Report No. 153. Paz-Castillo, Fernando A.; Kruger, Paul (Stanford Univ., Calif. Dept. of Civil Engineering). Jun 1972. 251p.

Development of the Federal Territory of Amazonas in Venezuela depends upon the construction of adequate facilities for inland bulk-cargo transportation. A major facility would be the extension of navigability for freight cargo along the Orinoco River deep into the Federal Territory. The extension of navigability can be further developed by an excavation of the divide between the Orinoco River and the Rio Negro River, a major tributary of the Amazon River in Brazil. A navigation waterway would greatly assist the economic development of the Federal Territory and provide a means for extended commerce between Venezuela and its neighboring countries Brazil and Colombia. The geographical properties of the divide region were examined and a suitable waterway was designed which can be constructed by modern explosive excavation methods. An alignment of about 62 km connecting the Atabapo River tributary of the Orinoco, with the Guainia River tributary of the Rio Negro appears to be economically feasible. Estimates from sparse hydrometeorologic data indicate that sufficient flow exists to support a navigation channel with a minimum width of 77 m for barges of 3 m draught. It is estimated that the waterway could maintain an annual freight of 80 Mt. The waterway is designed for construction by a combination of high-energy chemical and nuclear explosions and appropriate blasting and dredging operations. The requirements for the selected alignment include a total of 92 kt high-energy chemical explosives for excavations less than 17 m deep, and 157 10-kt nuclear explosives for a 15.5 km excavation through the divide. The total cost of the waterway excavation, with present costs of chemical explosives and nuclear explosion service prices announced by the USAEC, is about \$195,000,000. The mechanical and radiological safety of the excavation has been examined and no undue restrictions are apparent. The Federal Territory of Amazonas in its still virtually undeveloped state presents little population relocation problems. A schedule for detonations and reentry was prepared which can achieve the waterway construction over a period of four years. Construction of the waterway could be currently achieved under provisions of all existing international treaties; especially under Article V of the Non-Proliferation of Nuclear Weapons Treaty. The further continuation of this study could serve as a model project by the International Atomic Energy Agency to achieve the beneficial aspects of nuclear explosives sought by its member nations. (auth)

45476 (NVO-38-33) PROJECTED RESERVOIR PERFORMANCE. RIO BLANCO PROJECT. van Poolen, H. K. (van

Poolen (H. K.) and Associates, Littleton, Colo. (USA)). Mar 1972. 88p. (PNE-RB-26). Dep. NTIS.

Data from three wells were analyzed to determine basic reservoir parameters and to predict a future reservoir performance assuming nuclear stimulation is carried out. Permeability, as calculated by pressure transient analysis, ranged from 0.0064 to 0.087 md in offset well Fawn Creek Government No. 1. Log analyses of RB-E-01 indicate an average porosity of 9.4%, average gas saturation of 57% and a net pay of 421 ft. The permeability from pressure transient analysis averages 0.025 md for the Fort Union and 0.011 md for the Mesaverde. This gives 7.75 md-ft for the Fort Union and 1.23 md-ft for the Mesaverde or a total of 8.98 md-ft. Reservoir performance was predicted using a single-layer model with an average permeability of 0.0212 md. Cumulative production after 25 years was 19.9 BCF for 640-acre spacing. A two-layer model using the calculated values for the Fort Union and the Mesaverde gave a 25 year cumulative production of 17.9 BCF. Additional predictions made using a range of permeabilities are also included. One value of pressure dependent permeability was run using 640- and 320-acre spacings. However, it is felt that the information used to determine the pressure dependent permeability was not necessarily valid and should not be given much consideration. (auth)

45477 (NVO-86(Rev.1)) REPORTS AVAILABLE IN PLOW-SHARE OPEN FILE. (Nevada Operations Office (AEC), Las Vegas). May 1972. 33p. Dep. NTIS.

A list is provided of those publications concerning Gasbuggy, Rulison, Rio Blanco, and Wagon Wheel that have been placed in the Plowshare Open Files by the Nevada Operations Office of the USAEC. (L.C.L.)

45478 (NVO-409-1) ECOLOGY OF HOT CREEK VALLEY: VEGETATION AND SOIL RESPONSE TO UNDERGROUND DETONATION. Annual Progress Report FY 1971. Tueller, Paul T.; Bruner, Allen D.; Davis, J. Barry (Nevada Univ., Reno (USA). Max C. Fleischmann Coll. of Agriculture). 1971. Contract AT-(26-1)-409. 135p. Dep. NTIS.

An investigation is in progress on the influence of ground shock as a result of underground nuclear detonations on the soils and vegetation in Hot Creek Valley, Nevada. An ecological resource inventory of soils and vegetation has been completed to serve as a base for the determination of possible long-term vegetation changes. Seven plant associations and 30 soil families were mapped from ninety-four vegetation macroplots and soil descriptions. An estimation equation for grass, forb, and shrub production from shrub ring widths was developed to evaluate changes in vegetation production. By comparing growth estimations from ring width analyses with actual yields, the effect of a treatment (underground detonation) can be evaluated independent of yield differences caused by normal environmental fluctuations. Other studies are in progress on both treatment and control sites to help determine possible post-detonation changes. These include sap flow velocity measurements to compare with changes in soil moisture uptake rates in woody plants; plant phenology and productivity determinations to evaluate growth and developmental changes in the life cycle of plants; and photographic remote sensing with color infrared film for the analysis of plant vigor. Revegetation attempts using several species and methods of establishment and the influence of ground shock on other ongoing ecological studies are reported on. (auth)

45479 (NVO-746-122) MILROW/CANNIKIN SEISMIC EFFECTS. Engdahl, E. R. (National Oceanic and Atmospheric Administration, Rockville, Md. Environmental Research Labs.). May 1972. Contract AT(29-2)-746. 36p. Dep. NTIS.

Seismic effects of the underground nuclear explosions Milrow (October 1969, about 1 megaton) and Cannikin (November 1971, under 5 megatons) were monitored by a network of continuously-recording, high-frequency, high-gain seismographs located on Amchitka and nearby islands. Each explosion was immediately followed by hundreds of small discrete events ($m_B < 4$), of a similar focal mechanism and with a characteristic low-frequency signature, which apparently occurred as a result of the deterioration of the explosion cavity. This activity intensified, then terminated, within minutes of a large complex multiple event and concurrent formation of a surface subsidence crater, signaling complete collapse of the explosion cavity (Milrow, 37 hours; Cannikin, 38 hours). On a smaller scale, apparently unrelated to the collapse phenomenon, were a number of explosion-stimulated tectonic events which occurred intermittently for several weeks following each explosion near the explosion cavity and up to 13 km southeast of Cannikin ground zero along the Island. These events were confined to the upper crust of the Island, had characteristically high-frequency signatures, and, near the Rifle Range Fault, had focal mechanisms which correlated with pre-existing faulting. The evidence points to a short-term interaction of the explosions

with the local ambient tectonic stresses of the Island through explosion-generated elastic body waves or changes in fluid pore pressure. Continuous monitoring of the natural seismicity of the region since 1969 reveals no other apparent evidence for an interaction between either Milrow or Cannikin and natural tectonic processes. The high level of natural seismicity in the subduction zone from 20 to 60 km beneath the Island and the apparent low level of stress in the upper crust of Amchitka suggest that these zones are effectively seismically decoupled. (auth)

45480 (NVO-1163-TM-32) COMPARISON OF RESPONSE SPECTRA FROM YUCCA FLAT AND PAHUTE MESA UNDERGROUND NUCLEAR DETONATIONS. Lynch, Robert D.; Williams, Spencer L. (Environmental Research Corp., Las Vegas, Nev.). Jun 1972. Contract AT(29-2)-1163. 78p. Dep. NTIS.

The statistical techniques of regression analysis and covariance analysis are applied to response spectrum data from underground nuclear events detonated at the Yucca Flat area of the Nevada Test Site. In the case of regression analysis, estimates of yield and distance scaling exponents are determined. In the case of covariance analysis, only yield scaling exponents are determined. In both analyses, the deviation of observed response data from the assumed relationships among the variables is measured by the standard error of estimate. The results of the Yucca Flat analysis are compared with previous results of similar analyses of response spectrum data from Pahute Mesa detonations at the Nevada Test Site. Yield scaling exponents differ significantly for events in the two detonation areas. An analysis determines that the relatively heterogeneous emplacement environment at Yucca Flat compared to that of Pahute Mesa is responsible for the yield scaling discrepancy. (auth)

45481 (NVO-1163-TM-33) ANALYSIS OF SEISMIC WAVES RECORDED IN LAS VEGAS, NEVADA FROM UNDERGROUND NUCLEAR DETONATIONS. Davis, Lawrence L. (Environmental Research Corp., Las Vegas, Nev.). Jun 1972. Contract AT(29-2)-1163. 72p. Dep. NTIS.

The seismic data from Nevada Test Site detonations recorded on a comprehensive instrumentation array in Las Vegas, Nevada, were analyzed to study the propagation of the surface wave trains which contain significant amplitudes, to define the variations in response throughout the city and to relate these variations to known geologic and geophysical conditions. The propagation of the Rayleigh wave was examined and a simplified crustal model between NTS and Las Vegas was derived from the dispersion characteristics of the Rayleigh wave. A late arriving, large amplitude surface wave appears on the recordings in the southeast portion of the array. This wave apparently arrives from a direction other than radial and is possibly a Love wave because of the extremely small vertical motion. Peak amplitudes and spectral amplitudes are largest in the center of Las Vegas Valley, presumably because the alluvial fill is thickest in the center of the valley. Correlation of amplitudes and spectra with surficial geology and soil type was inconclusive. (auth)

45482 (NVO-1229-177) LONG TERM RELEASE OF RADIOACTIVITY FROM RAINIER MELT-GLASS. Essington, E. H. (Teledyne Isotopes, Las Vegas, Nev.). Oct 1971. Contract AT(29-2)-1229. 21p. Dep. NTIS.

As a result of underground nuclear explosions much of the radioactivity produced is associated with melt-glass, formed partly as the condensation product of vaporized rock and device materials. Movement of these radionuclides away from the explosion zone by flowing ground water requires dissolution of the melt-glass and solubilization of the component radionuclides. Samples of melt-glass collected from the Rainier explosion zone at the Nevada Test Site were suspended in water of a chemical nature native to the area for periods of time ranging from a few minutes to nearly 4 years. Results of the study indicate that the major beta contributing radionuclides solubilized include ^{90}Sr , ^{137}Cs , and ^{125}Sb . At no time during the 4 year period were levels of these radionuclides found above reference Concentration Guide values specified for release of radioactivity to individuals or population groups in uncontrolled areas. (auth)

45483 (NVO-1229-180) DISTRIBUTION OF RADIOACTIVITY IN AND NEAR THE RAINIER RUBBLE CHIMNEY. Essington, E. H.; Forslow, E. J. (Teledyne Isotopes, Las Vegas, Nev.). Sep 1971. Contract AT(29-2)-1229. 91p. Dep. NTIS.

Underground nuclear explosions produce radioactive contaminants which can be transported with ground water. In order to accurately determine the variation in radioactive contaminant concentration in the hydrologic system in the vicinity of an underground nuclear explosion site, the initial spatial distribution and concentration of radionuclides in the cavity/rubble chimney complex must be known. An extensive analytical program was conducted to determine the spatial distribution of major radioactive contaminants in the one accessible rubble chimney, Rainier.

Concentrations of the fission products ^{137}Cs , ^{90}Sr , ^{106}Ru , ^{144}Ce , and ^{125}Sb ; and the activation products ^{134}Cs , ^{60}Co , $^{152-154}\text{Eu}$, and ^{55}Fe were measured on 5 identifiable rock types: melt-glass, contact melt, altered rubble, rubble, country rock, and an inseparable mixture of melt/rubble material. No data interpretation is given. (auth)

45484 (SC-DC-722296) LARGE SCALE CHARACTERIZATION TESTS OF GELLED SLURRY BLASTING AGENTS. Fastle, D. L.; Laursen, H. G.; Marker, T. F. (Sandia Labs., Albuquerque, N. Mex.). [nd]. 28p. Dep. NTIS.

A series of test events to characterize the large scale detonation properties of three gelled slurry blasting agents is described. The slurries chosen were DBA-22M, DBA-65T2, and DBA-105T2, all the claimed proprietary products of Intermountain Research and Engineering Co. of Salt Lake City, Utah. It was also decided that the test series would include reference TNT devices in the same geometry, same charge weight, and as near the same density as is practical. Results of the tests indicate that DBA-22M and DBA-65T2 have considerable military potential as blasting agents. It is further concluded from other data available on DBA-22M that overpressure and impulse data, while very valuable, do not wholly account for the superior performance of this slurry agent over TNT in a more realistic military situation. It is also concluded that 65T2 has possibilities as a cratering or earth moving agent. (J.R.D.)

45485 (SC-RR-720359) CANNIKIN: EFFECTS ON ARCHAEOLOGICAL SITES. Merritt, M. L. (Sandia Labs., Albuquerque, N. Mex.). Jul 1972. 19p. Dep. NTIS.

Examination before and after Cannikin reveals that archaeological Site 32 suffered loss by cliff fall of about 45 m² of previously undisturbed midden area, as well as cave-ins of the walls of excavations made during the previous summer. Turf slides occurred along the outer slopes of the bluff on which Site 32 was located; a crack or incipient slide opened at Site 67; and cracks parallel to and just inward from the edge occurred in Sites 31 and 32 and probably elsewhere, but these slides and cracks only affected material that had already moved from its original position and thus had lost its archaeological context. AEC construction caused no damage to archaeological sites. (auth)

45486 (UCRL-51235) METHOD FOR ESTIMATING HEIGHTS OF WATER WAVES GENERATED BY UNDERWATER CRATERING EXPLOSIONS. Garcia, W. J. Jr. (California Univ., Livermore. Lawrence Livermore Lab.). 7 Jun 1972. Contract W-7405-eng-48. 10p. Dep. NTIS.

A method is described for predicting the wave heights of water waves generated by underwater cratering explosions. The method of predicting wave heights is based upon underwater explosion data, the results of numerical model simulations, and laboratory experiments. It was found that the ratio of wave height to source displacement is independent of the details of the displacement-time history of the source motion. A method for simulating water wave generation by underwater cratering explosions with a numerical model is also discussed. (auth)

45487 (UCRL-73939) CORROSION AND SCALING BY STEAM IN NUCLEAR GEOTHERMAL POWER PLANTS. Krikorian, Oscar H. (California Univ., Livermore. Lawrence Livermore Lab.). 2 Jun 1972. 36p. (CONF-720607-24). Dep. NTIS.

From eighteenth annual American Nuclear Society conference; Las Vegas, NV. (18 Jun 1972).

The quality of steam produced in a nuclear geothermal well will significantly affect the design, operation, efficiency, and expense of a steam-turbine powerplant. Problem areas related to steam quality are: the carrying and deposition of minerals (scaling) by steam, the carrying of radioactive contaminants by steam, the amount and types of noncondensable gases, and the corrosion of alloys. Minerals carried by steam are usually salts (such as NaCl, NaOH, or sodium silicates) and silica. Deposition of minerals in the piping or turbine can reduce flow, give inefficient turbine operation, or cause material damage to the turbine. The salt deposits are water soluble and can be washed away as an operational procedure. Silica, however, poses serious problems because of its insolubility. The maximum permissible concentration of silica in steam is about 0.05 ppm, in order to reduce silica deposition to a negligible level. If temperatures of 620°K and steam pressures of 80 atm were to be used, the silica content of steam would be about 9 ppm, i.e., more than 2 orders of magnitude higher than the maximum permitted concentration. It is clear that such impure steam would need to be purified, either by scrubbing with water or by some other means, before introducing it into the turbine. Radioactive contaminants from fusion or fission explosives that may be volatile after 180 days are given as the fission products: ^{85}Kr , ^{103}Ru , ^{106}Ru , ^{106}Rh , ^{125}Sb , ^{127m}Te , and ^{137}Cs ; and induced activities: ^3H , ^{22}Na , ^{32}P , ^{35}S , ^{37}Ar , and ^{134}Cs .

For 1 Mt of yield, the amount of gamma radiation for the above species is 3.0×10^{16} MeV/sec for all-fission explosives, and 1.1×10^{15} MeV/sec for fusion explosives with 3 kt of fission. Fusion explosives also produce a high level of beta activity in the form of tritium. A thermodynamic model is used to roughly assess the volatilities of the main gamma emitters in steam. The approach used is to estimate the composition and amounts of volatile rock components and radioactivity retained in the steam when the major portion of the melted rock has cooled and separated from the steam phase. The radionuclide ^{85}Kr is, of course, a permanent gas while ^{125}Sb and $^{127\text{m}}\text{Te}$ are also expected to have high volatilities. For ^{103}Ru and ^{106}Ru , high volatilities are expected under oxidizing conditions with low volatilities expected under reducing conditions. The radionuclides ^{134}Cs and ^{137}Cs are believed to be significantly volatile, with their actual volatilities affected by the chloride and carbonate content of the rock formation. Radioactive impurities carried by steam are likely to coprecipitate with salts and silica in the turbine system, if the steam quality is poor. Noncondensable gases present in geothermal steam lead to three problems: enhanced corrosion to alloys; increased pumping costs to maintain turbine efficiency; and possible enhancement of carrying of minerals in steam. The noncondensable gas content of geothermal steam ranges from 0.01 to 5 wt %, with more typical values of 0.1 to 1 wt %. The gas CO_2 represents from 70 to 90% of the noncondensable gas. Next in importance are H_2S and H_2 , with lesser amounts of CH_4 , NH_3 , and N_2 . Limited information is available on the corrosion resistance of common alloys and other materials to geothermal water and steam. Tests have been performed on alloy coupons with water at 320 to 510°K and steam at 370 to 470°K for periods up to 150 days. The alloys most resistant to corrosion are titanium, austenitic stainless steels, and chromium-plated steels. Corrosion of alloys can occur from the presence of NH_3 , H_2S , CO_2 , and especially chlorides in steam. Also, corrosion is generally enhanced by the presence of oxygen. (auth)

45488 (USGS-474-148) GEOLOGIC AND HYDROLOGIC EFFECTS OF THE CANNIKIN UNDERGROUND NUCLEAR EXPLOSION, AMCHITKA ISLAND, ALEUTIAN ISLANDS, ALASKA. Amchitka-33. (Geological Survey, Denver, Colo.). Jun 1972. Contract AT(29-2)-474. 87p. Dep. NTIS.

The USGS studies related to the Cannikin Event included visible geologic effects, ground deformation measurement, analysis of high-speed motion photography, hydrologic effects, and a long-term program to analyze water on Amchitka Island for radionuclide content. Most visible effects were limited to an area within 2 km of ground zero. These included minor movement on faults, rockfalls and slumping along the sea cliffs, and about 1 m uplift along about 2 km of the Bering coastline. Displacement of less than 1 m occurred along a distance of about 600 m on two northeast-trending faults which lie 760 m southeast and 1,070 m northwest of GZ. The explosion cavity collapsed to the surface 38 hours after detonation forming an asymmetrical oval closed depression 915×1270 m in diameter and more than 10 m deep. The deepest point of the sink is 366 m southeast of GZ, which itself subsided only 5 m. Precise measuring of strain nets shows that a line 2 km long that trends northeast across GZ extended 1.2 m. Calculated strains decrease with slant distance from GZ out to a distance of 6 km. At that distance the strain was 2×10^{-5} . Principal strains show northeast-southwest extension. Flow of White Alice Creek, which drains the Cannikin site, has been reduced to about 4% of normal because of capture of most of the creek flow by the collapse sink. It is estimated that the chimney will fill in 1.5 years and that a lake 6 m deep will form in the sink before the normal flow of White Alice Creek can resume. The response of fluid pressures to the Cannikin Event was recorded in a number of boreholes, and a relation between fluid pressures (P) and slant ranges (R_s) has been established as $P = 963 R_s^{-2.14}$. Chemical analyses of water collected on and around Amchitka after the event were compared with preevent analyses as part of a long-term program of radiochemical monitoring. No measurable increase in radioactivity has been detected. (auth)

45489 LONG-RANGE DETECTION AND LOCATION OF SOME SHALLOW-DEPTH EXPLOSIONS IN THE NORTHWEST PACIFIC. Northrop, John (Naval Undersea Research and Development Center, San Diego, Calif.). Bull. Seismol. Soc. Amer.; 62: No. 3, 793-803 (Jun 1972).

During a survey of underwater explosions, a set of 382 closely-spaced events was located in the Northwest Pacific near the Emperor Seamounts. Later, it was learned that the shots were from a joint U. S.-Japan science cooperation seismic-refraction program. After-the-fact correlations were then made between the Soud Fixing and Ranging (SOFAR) fixes and shot-log data. Shots were composed of 3 to 288-lb Tetrytol or Seismogel demolition charges. Detonation depths varied from 60 to 420 ft. Results are that: (1) if only shots of 8.6 lb or greater are considered, a larger percentage (87.7) of shots composed of Seismogel

was successfully located than of those composed of Tetrytol (59.5); (2) the threshold of detection was lower (9 lb at 90 ft as compared to 12 lb at 110 ft) in the northern part of the profile (37°N) than in the southern part (32°N); and (3) signals from shots detonated over the southern edge of the Shatsky Rise were received at a higher level than those detonated off the Rise. The reason for (1) is the higher detonating velocity of Seismogel; for (2) it is the shoaling of the SOFAR channel with increasing latitude, and for (3) it is the effect of a highly reflective bottom which sloped toward the hydrophones. (auth)

45490 SEISMIC-SOURCE ENERGIES OF UNDERGROUND NUCLEAR EXPLOSIONS. Perret, William R. (Sandia Labs., Albuquerque, N. Mex.). Bull. Seismol. Soc. Amer.; 62: No. 3, 763-74 (Jun 1972).

Records of particle velocity in the free-field geological environment near underground nuclear explosions may be interpreted to yield the energy flux at a measuring station near or within the region of elastic response of the rock. Such data from 21 events in several types of rock have yielded energy fluxes. Total energies associated with the motion propagated into the seismic region have been derived from all but seven of these. The total energies are a measure of the seismic-source strength, and the ratios of these energies to those released by the explosion are measures of the energy coupled into the Earth as seismic motion. Explosions in granite, dolomite, or wet tuff produce seismic sources of the order of 2% of the explosively released energy. Similar explosions in porous rocks such as dry tuff and desert alluvium provide seismic sources an order of magnitude weaker than those in hard or wet rock. (auth)

45491 ELASTODYNAMIC NEAR-FIELD OF A FINITE, PROPAGATING TENSILE FAULT. Haskell, N. A.; Thomson, K. C. (Air Force Cambridge Research Labs., Bedford, Mass.). Bull. Seismol. Soc. Amer.; 62: No. 3, 675-97 (Jun 1972).

Displacement, particle velocity, and acceleration wave forms in the near-field of a finite, propagating tensile fault were computed by numerical integration of the Green's function integrals for an infinite medium. The displacement discontinuity (dislocation) on the fault plane is assumed to have the form of a unilaterally propagating, finite ramp function in time. Computer generated wave-form plots are presented for a variety of close-in locations with respect to the fault plane and for two different fault lengths. (auth)

45492 EXISTENCE OF NUCLEAR WEAPONS: A PRIME ENVIRONMENTAL THREAT. Gofman, John W. (Univ. of California, Berkeley). Environ. Aff.; 1: No. 4, 782-91 (Mar 1972).

The environmental threat of the existence, not just the testing, of nuclear weapons is argued. The subjects discussed include the reasons why nuclear weapons exist, the political aspects of the nuclear arms race, the prospects for limiting or eliminating nuclear weapons, and the significance to the nuclear arms race and the environmental effects of the Cannikin test. (L.C.L.)

45493 NUCLEAR EXPLOSIVES AND HYDROCARBONS. Cohen, P. (CEA, Paris). Rev. Inst. Fr. Petrole Ann. Combust. Liquides; 26: No. 10, 927-34 (Oct 1971). (In French).

The construction of a nuclear explosive device 12 in. in diameter which produces very little tritium is feasible in France. Such a device would be well adapted for contained nuclear explosions set off for the purpose of hydrocarbon storage or stimulation. The different aspects of setting off the explosive are reviewed. In the particular case of gas storage in a nuclear cavity in granite, it is demonstrated that the radiation dose received is small. (France)

45494 PREDICTION OF UNDERGROUND NUCLEAR EXPLOSION EFFECTS IN WAGON WHEEL SANDSTONE. Terhune, Robert W. (Univ. of California, Livermore). Nucl. Technol.; 15: No. 3, 431-46 (Sep 1972).

Project Wagon Wheel is a study to investigate the technical concept of nuclear stimulation of a natural gas reservoir located near Pinedale, Wyoming. The results are presented of computer calculations, simulating the explosion effects, based on equation-of-state measurements of core samples taken from El Paso Natural Gas (EPNG) Wagon Wheel No. 1 hole at a depth between 8,000 to 12,000 ft. The shear failure envelope for Wagon Wheel sandstone is almost identical to that for Hoggar granite. A calculation for Wagon Wheel sandstone for one kt energy at a depth of 300 m duplicated cavity radius, shear fracture radius, stressed region, peak particle velocity, and peak acceleration data measured from the nuclear experiments conducted in the Hoggar granite by the French. The conclusion drawn from this comparison is that the Hoggar granite chimneys and regions of increased permeability provide a reasonable model to assume for Wagon Wheel. Cavity and chimney dimensions are predicted for a single explosive detonated at a depth of 10,000 ft. (auth)

45495 STUDY OF TRAVEL-TIME RESIDUALS WITH RESPECT TO THE LOCATION OF TELESEISMIC EVENTS FROM BODY-WAVE RECORDS AT MOXA STATION. Bormann, P. (Zentral-Institut Physik der Erde, Potsdam, Ger.). Gerlands Beitr. Geophys.; 81: No. 1-2, 117-24(1972).

From frequency distributions of travel-time errors based on focal data of the world data centers no exact quantitative parameters on the total travel-time deviation from the Jeffreys-Bullen (J-B) travel-time model can be obtained. The same holds for speed anomalies in the focal region. However, the difference of the systematic travel-time errors of various waves indicates travel-time differences of the waves that are not included in the J-B model. Interpretation of the causes is hereby often facilitated. Among other things, inferences can be drawn to anomalous travel times in the region of the station or the reflecting region. When considering the systematic travel-time errors with inverse sign as a correction for the J-B travel times, then the accuracy of distance determinations and, hence, the accuracy of the location of seismic events by a single station can often be considerably increased. (auth)

45496 IDENTIFICATION OF TELESEISMIC EVENTS IN THE RECORDS OF MOXA STATION. Bormann, P. (Zentral-Institut Physik der Erde, Potsdam, Ger.). Gerlands Beitr. Geophys.; 81: No. 1-2, 105-16(1972).

Simple, efficient criteria for a routine identification of seismic events are given. With a minimum of technical and financial expenditure they enable earthquakes and underground nuclear explosions to be distinguished in the records of Moxa Station. In complex application, related to the signal-to-noise ratio (SNR), a very high identification rate is obtained. It corresponds about to the rate given in the literature for large station arrays. (auth)

45497 ELECTROMAGNETIC PULSE AND TIME-DEPENDENT ESCAPE OF NEUTRONS AND GAMMA RAYS FROM A NUCLEAR EXPLOSION. Sandmeier, H. A. (Los Alamos Scientific Lab., N. Mex.); Dupree, S. A.; Hansen, G. E. Nucl. Sci. Eng.; 48: No. 3, 343-52(Jul 1972).

The time-dependent escape of neutrons and gamma rays from a nuclear explosion is of interest in Plowshare and nuclear weapons effects investigations, particularly with regard to generation of the electromagnetic pulse (EMP). The neutron and gamma-ray leakage rates, spectra, and useful yield-related integral quantities are calculated for the 24 kt S-device. EMP parameters and fields at 500 m from the explosion of the S-device on the ground are presented and discussed. (auth)

45498 SEISMIC ARRAYS AND SEISMOLOGY. Husebye, Eystein S. Abstracts of Uppsala Dissertations from the Faculty of Science, No. 223. Uppsala; Univ. of Uppsala (1972). 13p. Thesis.

The concept of seismic arrays, seismic data processing methods, signal parameter extraction, and applications of array data in seismological research are discussed. Information is included on both continental arrays and large aperture seismic arrays. (auth)

Protective Structures and Equipment

Refer also to abstracts: 46104, 46107, and 46108.

45499 RADIATION SAFETY AROUND THE HOT FACILITIES AT THE PINSTECH. Shaikh, Unis; Mubarak, M. A. (Pakistan Inst. of Nuclear Science and Tech., Rawalpindi). Nucleus (Karachi); 8: No. 4, 13-27(1971).

The hot facilities at the Pakistan Institute of Nuclear Science and Technology (PINSTECH) are described. These facilities include a 5-MW swimming pool-type reactor, nuclear chemistry, and nuclear materials laboratories. (auth)

45500 SOME OBSERVATIONS ON RADIATION SAFETY IN VIRGINIA DENTAL OFFICES. Mizeras, A. V. (Bureau of Industrial Hygiene, Richmond, Va.). Va. Dent. J.; 48: 26-33(Apr 1971).

The factors with which the Virginia Health Dept. is concerned in its radiation safety survey program are discussed. These include collimation, filtration, film badges, lead shielding, portable lead shields, lead aprons, cones, tube leakage, workload, timers, film speed and processing, positions of personnel during x rays, and the use of specialized x-ray equipment. (P.C.H.)

45501 PROCEEDINGS OF THE 1972 CLEAN AIR CONFERENCE, MELBOURNE, 15th-18th MAY 1972. Parkville, Australia; Clean Air Society of Australia and New Zealand (1972). 308p. (CONF-720541).

Forty-six papers on clean air were presented at the conference. A separate abstract was prepared for each of two. Forty-four papers are not within the scope of NSA. (L.C.L.)

For abstracts of individual papers see: 45502 and 45521.

45502 REMOVAL OF RADIOACTIVE IODINE FROM VENTILATION EXHAUST STREAMS AT THE AAEC RESEARCH ESTABLISHMENT. Bonhote, P. A.; May, G. G.; Polson, H. J.; Weir, R. (Australian Atomic Energy Commission Research Establishment, Lucas Heights). pp 114-19 of Proceedings of the 1972 Clean Air Conference. Parkville, Australia; Clean Air Society of Australia and New Zealand (1972).

From International clean air conference; Melbourne, Australia (15 May 1972). See CONF-720541.

Since radioactive iodine is volatile, abundant in fission products, and selectively concentrated in the thyroid gland, it is usually the most important nuclide to be considered in assessing the risk of possible releases of fission products. At low mass concentrations in air, it forms many volatile organic compounds, the most abundant being methyl iodide. The elemental form is easily absorbed in activated charcoal, but the removal of methyl iodide at high humidities requires the use of special impregnated charcoals. The design criteria adopted at the Research Establishment are discussed and the absorption beds, the bed loader, bed housing, and handling procedures are described. The beds are tested at regular intervals, and results indicate a useful working life of one year. (auth)

ENVIRONMENTAL AND EARTH SCIENCES

45503 (COO-2195-6) PLANETARY TRANSPORT PROCESSES IN THE UPPER ATMOSPHERE. Annual Progress Report. Newell, Reginald E. (Massachusetts Inst. of Tech., Cambridge, Dept. of Meteorology). 31 May 1972. Contract AT(11-1)-2195. 16p. Dep. NTIS.

Results of research on fundamental processes which govern the structure, mass, and momentum of the upper atmosphere are reported. Lists of publications, theses, reports, and papers associated with the research are included. (J.R.D.)

Minerals and Ores

Refer also to abstracts: 45566, 45569, 45572, 46156, and 47300.

45504 PECULIARITIES OF FORMATION OF AN INCREASED URANIUM CONCENTRATION IN OXIDIZING CONDITIONS. Borkov, F. P.; Egorov, N. I.; Zaitsev, E. V. (Moscow Geological Inst.). *Izv. Vyssh. Ucheb. Zaved., Geol. Razved.*; No. 5, 51-7(1971). (In Russian).

The peculiarities of the formation of an enhanced U concentration under oxidizing conditions were examined. Similar concentrations are generally adapted to an oxidation zone of U deposits and are scarcely ever encountered in rocks with a Clark content of uranium. The U concentrations in oxidation zones of rocks with normal or somewhat increased Clark radioactive elements are usually adapted to regions of arid climate, and their formation is considered to be a result of an evaporative concentration of elements in ground waters (insolation). The U concentrations in the ore phenomena under consideration did not agree with these notions, but instead are examples of favorable geological conditions and geochemical circumstances. An epigenetic accumulation of U might occur under conditions of an oxidizing medium through its fixation in the hexavalent state. (tr-auth)

45505 DISTRIBUTION AND GENESIS OF RADIOACTIVITY IN COALS OF THE PECHORA RIVER BASIN. Gipsh, A. A.; Kapaturin, G. G.; Yudovich, Ya. E. *Izv. Vyssh. Ucheb. Zaved., Geol. Razved.*; No. 6, 61-70(Jun 1971). (In Russian).

The radioactivity of coals of the Pechora basin was studied in 77 samples chosen from mines of the Vorkutsk, Yun'-Yayinck, and Khal'mer'yu deposits. The content of the radioactive elements ranged between 0.4 and 10 g/T, was typically 2 g/T, and was related to the ash content. The essential carriers of radioactive elements in coal are the terrigenous components of ash—clay minerals. For low-ash strata, the increase of activity with increased ash content occurred more rapidly than high-ash strata. In Vorkutsk coals the fraction of activity due to uranium decreased with a rise in the ash content; the fraction due to potassium increases. The content of the potassium carrier (hydromica), with increase of ash content rose more rapidly than the U carrier (kaolinite). The Th content in these coals was extremely small. (tr-auth)

45506 PROBLEM OF EPIGENETIC ZONALITY IN URANIUM ORE DEPOSITS OF CARBONATE ROCKS. Shmariovich, E. M. *Litol. Polez. Iskop.*; No. 4, 105-16(Jul 1971). (In Russian).

The general characteristics of morphological features of uranium mineralization are provided by deposits in carbonate rocks and confirm their epigenetic character. In an instance of one of the deposits, an adaptation of the ore to an outcrop of an ancient reddish zone of stratified oxidation was observed. Study has substantiated the incompetence of the scheme of "oxygenless zonality" of uranium ore in carbonate oil-bearing rocks. Lime-stone recrystallization near ore zones is a result of secondary reestablishment of stratified-oxidized rocks in conjunction with the onset of gleyey (sic) water and petroleum. It is proposed that the intrusion of restoration fluids into carbonate rocks is accomplished along fracture zones. (tr-auth)

45507 SPECTROMETRY AND COMPUTERS USED FOR MAPPING DEPOSITS. Bland, Frederick. *Aust. Mining*; 64: No. 5, 56-7(May 1972).

A description is given of the airborne gamma spectrometric process as used for mapping uranium deposits in Australia. The theory behind the process is outlined and the installation of the 10 crystal spectrometer and computer processing of the data accumulated on tape are briefly described. (Australia)

45508 PRESENT STATUS AND PROSPECTS OF RADIOACTIVE ORE EXPLORATION IN YUGOSLAVIA. Com. Naz. *Energ. Nucl., Notiz.*; 18: No. 5, 38-40(May 1972). (In Italian).

A review of the present situation of uranium exploration and a description of ore deposits in Yugoslavia are presented. (auth)

Radioactive Effluents

Refer also to abstracts: 45398, 46102, 47401, 47404, 47413, 47424, 47435, 47446, 47465, 47471, 47473, 47479, 47486, 47487, 47489, 47497, 47498, 47502, 47510, 47516, 47518, 47532, 47534, 47535, 47543, 47545, and 47564.

45509 (KFK-1560) ENVIRONMENTAL BURDEN OF TRITIUM. Koenig, L. A. (Kernforschungszentrum, Karlsruhe (West Germany)). Feb 1972. 26p. (In German). Dep. NTIS (U. S. Sales Only).

A review is presented on the present state of knowledge with respect to the tritium burden on the environment. Following a compilation of the most important characteristics, the theoretical relationship between tritium incorporation and radiation burden is discussed. Sources of environmental tritium contamination are the tritium produced by nature and in nuclear fission reactors, the use of tritium-containing luminous colors, and, above all, the nuclear weapons tests. The tritium released undergoes a worldwide spread by the circulating water. The radiation burden due to this tritium contamination and evaluated by concentration measurements amounts presently to 0.1 mrem/g. An increase of this radiation burden is not anticipated until the year 2000. In case of realization of fusion reactors considerable attention will have to be paid to the tritium problem, since the tritium requirements of a fusion reactor amount to 4×10^5 Ci/MW, while in the heavy-water reactor only about 10^3 Ci ^3H /MW and in the light-water reactor about 10^1 Ci ^3H /MW are produced. (auth)

Radioactivity Transport and Monitoring

Refer also to abstracts: 45205 and 47623.

45510 (AAEC/TM-627) ENVIRONMENTAL HAZARDS OF FOSSIL AND NUCLEAR POWER PRODUCTION. Watson, G. M. (Australian Atomic Energy Commission Research Establishment, Lucas Heights). 1972. 20p. (CONF-720627-2). Dep. NTIS (U. S. Sales Only).

From Symposium on the role of nuclear energy in Australia's development; Canberra, Australia (1 Jun 1972).

It is now essential for engineers to recognize the environmental hazards which may be associated with their constructions and to make proper allowance for them when costing different solutions to a problem. Both nuclear and fossil-fueled power stations present environmental hazards, and these can be traced from mines through processing plants to power station wastes. It is necessary that engineers develop an appropriate basis of comparison if they are to assess the relative environmental costs of nuclear and fossil-fueled power stations; this must allow for effects on health, ecological effects, and aesthetic effects. The biological effects of radiation have been fairly well quantified with respect to major exposures, but there is still uncertainty on the precise form of dose-response relationships at low doses, such as might be relevant to industrial and public health. There are also uncertainties for non-radioactive pollutants, and it is probably not very logical to treat the dose-effect relationships of radioactive and chemical pollutants as being quite different in nature, as is commonly done at present. There are internationally accepted standards for allowable radiation exposure. It is possible to base allowable discharges of radioactive contaminants to the environment on these standards, provided the modes of dispersion are understood and possible pathways to the exposure of man are recognized. The philosophy of radiation protection recognizes that although radiation risks can be made very small, they cannot be abolished entirely, and that these risks must be balanced against the benefits of using radiation. These risks can be quantified and compared with other hazards of work and life which people accept, voluntarily or involuntarily. There are insufficient data available on the effects on health of various pollutants, and there is too little public agreement on the relative worth of mined and unmined natural resources, to allow a detailed comparison of the environmental hazards of nuclear and fossil-fueled power stations, but it is possible to begin this comparison in limited areas. (auth)

45511 (HASL-257) FALLOUT PROGRAM. Quarterly Summary Report, March 1, 1972-June 1, 1972. Hardy, Edward P. Jr. (New York Operations Office (AEC), N. Y. Health and Safety Lab.). 1 Jul 1972. 201p. Dep. NTIS.

Activities related to the HASL fallout program conducted during

the period from March 1, 1972 to June 1, 1972 are presented in nine sections. A separate abstract was prepared for each of the sections. (R.C.K.)

For abstracts of individual sections see: 45516-45519, 45525, 45526, and 45538-45540.

45512 (HASL-257(App.)) FALLOUT PROGRAM. Quarterly Summary Report, March 1-June 1, 1972. Appendix. (New York Operations Office (AEC), N. Y. Health and Safety Lab.). 1 Jul 1972. 437p. Dep. NTIS.

Data are given on ^{90}Sr and ^{89}Sr in monthly deposition at world land sites, radiostrontium deposition at Atlantic Ocean Weather stations, radionuclides and lead in surface air, and radiostrontium in milk and tapwater. A table of conversion factors and radionuclides of interest in the HASL fallout program are also listed. (P.C.H.)

45513 (UCRL-51242) ENVIRONMENTAL LEVELS OF RADIOACTIVITY IN THE VICINITY OF THE LAWRENCE LIVERMORE LABORATORY JANUARY THROUGH DECEMBER 1971. Gudiksen, P. H.; Lindeken, C. L.; Gatrousis, C.; Anspaugh, L. R. (California Univ., Livermore. Lawrence Livermore Lab.). 14 Jun 1972. Contract W-7405-eng-48. 47p. Dep. NTIS.

An environmental surveillance program is continuing in the Livermore Valley and Site 300 areas to determine to what extent, if any, the Lawrence Livermore Laboratory has contributed to the levels of radioactivity in the surrounding environment. The results of analyses for gross radioactivity and for specific radionuclides of interest in a variety of environmental samples are presented. Environmental levels of radioactivity were found to be below the concentration guide levels in AEC Manual Chapter 0524. Airborne particulate samples showed gross beta activities similar to those measured during previous years. Gamma spectral measurements of these samples revealed detectable activities of typical gamma emitters observed in global fallout, including fresh fission product debris from the 1970 Chinese and French nuclear weapons tests. Isotopic uranium analyses of airborne particulates collected at Site 300 indicated that the uranium was depleted in ^{235}U with respect to natural uranium. Soil samples collected within the Livermore Valley and Site 300 showed that the only activity contributed by the Laboratory was some low level ^{239}Pu contamination in one small on-site area and in one of the holding ponds at the Livermore Sewage Treatment Plant where the Laboratory's liquid effluent is processed. Samples of this effluent were also collected at various stages throughout the treatment plant to detect any possible build-up of activity within the facility. Gamma spectral analyses of milk and vegetation samples revealed no gamma-emitting radionuclides other than those present naturally or in global fallout. The results of the tritium analyses of vegetation samples revealed significant variation from one location to another which may be due to Laboratory operations. The activities of samples collected on-site and in off-site areas generally downwind from the Laboratory are up to 2 orders of magnitude higher than those observed in areas where the Laboratory's contribution should be minimal. Water samples collected in the Livermore Valley exhibited normal gross beta and tritium activities, as did those from Site 300, except from one spring with water containing tritium activities that are about 4 orders of magnitude (but less than 1% of standard) greater than the others. Assessment of the doses to an individual from the observed environmental activities indicated the contribution from artificially produced radionuclides was small in comparison with that from natural processes. An adult in this area will receive less than 1 mrem/yr from the artificially produced radionuclides, with the exception of ^{144}Ce , which may deliver several mrem/yr to the organs of interest. (P.C.H.)

45514 GASEOUS PLUME DIFFUSION ABOUT ISOLATED STRUCTURES OF SIMPLE GEOMETRY. Meroney, Robert N.; Yang, Bing-tsuen (Colorado State Univ., Fort Collins). pp 1022-9 of Proceedings of the Second International Clean Air Congress. /Englund, H. M. (ed.). New York: Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

The state of gaseous dispersion prediction in the vicinity of buildings and results on the effects of building orientation, leakage location, and thermal stratification are reported. The study led to the following conclusions: dispersion patterns differ in regions with and without the presence of a building structure. The ground-level concentration variation with longitudinal distance in the wake region shows much flatter slopes than those in open fields; for a specific building orientation, dispersions are similar for different release ports (from the top, middle, and bottom of the building height). Strong turbulent mixing motions are believed to smooth out any effects from the origin of release; aerodynamic effects due to building orientations (0° , 45° , 90° , 135° , 180°) cause

a slightly different concentration distribution in the cavity and near wake region. This difference depends on the portion of effluent which is initially carried downwind by convective motions; further downwind the dispersion will be independent of the original building shapes; mechanical turbulence dominates the dispersion behavior. The stratification becomes more important further downwind; inversion stratification causes about 8% higher ground concentration than obtained for neutral stratification; strong inversion stratification causes smaller transverse dispersion and "freezes" the plume growth in the vertical direction; and the plume growth in the transverse direction is about 3 to 5 times greater than that in the vertical direction for both neutral and stabilized stratified shear flows. (auth)

45515 DATA. SECTION IV. OTHER DATA. Radiat. Data Rep.; 13: No. 6, 377-84 (Jun 1972).

Data are reported on off-site surveillance activities around the Nevada Test Site during July through Dec. 1971 including measurements of the gamma dose rates at 13 locations and the maximum ^{131}I content measured in off-site milk samples; gamma radioactivity in the environs of the Brookhaven National Laboratory during Jan. to July 1968; gross beta radioactivity in liquid waste effluents at the BNL site boundaries during Jan. to July 1968; and the location of existing and proposed U.S. nuclear power plants. (L.C.L.)

Atmosphere

Refer also to abstracts: 45501 and 45861.

45516 (HASL-257, pp 1,2-17) MERIDIONAL TRANSPORT OF ^{90}Sr IN THE TROPOSPHERE. Servant, J. (Centre de Physique Atomique, Toulouse); Hoang, C. T. 1 Jul 1972.

In Fallout program.

Data on ^{90}Sr concentrations in surface air and in fallout, put in U.S.A.E.C. Health and Safety Laboratory, have been analyzed to interpret the dispersion of ^{90}Sr in the troposphere during the moratorium 1963 to 1965. Different processes, such as eddy diffusion, mean meridional motion, and stratospheric-tropospheric exchange were examined in relation with this analysis. At low latitudes, of both hemispheres, the distribution of ^{90}Sr , for the most part, can be accounted for by a meridional circulation. In the midlatitude (30°N - 20°S), the direct injection from the stratosphere seems to play a major role in the distribution observed. Some difference in the behavior of ^{90}Sr in the southern hemisphere appears to result from an additional source in the upper troposphere of the southern equatorial zone, due to the interhemispheric exchange. At 70°N , in winter, a transfer of ^{90}Sr at the ground level from the midlatitudes is suggested. (auth)

45517 (HASL-257, pp 1,18-32) HIGH FALLOUT IN THE WESTERN UNITED STATES. An Explanation. Volchok, H. L. 1 Jul 1972.

In Fallout program.

Through analysis of the ^{90}Sr concentrations in precipitation at conterminous U. S. sites, plus ^{90}Sr , ^{137}Cs , and ^{239}Pu data in surface air, several hypotheses for systematically high fallout in the western U. S. can be rejected. The most plausible explanation, consistent with all of the data, requires a preferential longitude for stratosphere-troposphere exchange at about 110° (west). This is close to the longitude of Salt Lake City, Utah, and with this mechanism, essentially all of the high ^{90}Sr fallout at that location is reasonably understood. (auth)

45518 (HASL-257, pp 1,33-50) UPDATING STRATOSPHERIC INVENTORIES TO JULY 1971. Krey, P. W.; Krajewski, B. T. 1 Jul 1972.

In Fallout program.

The stratospheric inventories of ^{90}Sr , ^{95}Zr , and ^{137}Cs are reported to July 1971. The average $^{137}\text{Cs}/^{90}\text{Sr}$ ratio calculated from the inventories in each hemisphere during 1970 and 1971 (1.45) agreed with the theoretical production ratio of 1.44. Revised calculations indicated that the Chinese test of 10/14/70 and the large French tests of 5/31 and 7/5/71 injected into the stratosphere about 200 and 240 kCi of ^{90}Sr , respectively. Approximately 40 kCi of ^{90}Sr was injected into the Southern Hemisphere stratosphere probably by the intermediate yield French test of 6/12/71. (auth)

45519 (HASL-257, pp III,53-8) PROTECTION SERVICE, SITE SURVEY AND METEOROLOGY SECTION. Quarterly Report. (EURATOM Joint Nuclear Research Centre, Ispra, Italy). 1 Jul 1972.

In Fallout program.

Data are given on ^{90}Sr , ^{89}Sr , ^{137}Cs , ^{239}Pu , and ^{238}Pu contents in air and wet and dry deposition at Ispra for Sept. 1971 through

March 1972 and ^{90}Sr content in milk at Ispra for 1960 through 1971. The chemical procedures and counting techniques used are described. (P.C.H.)

45520 (NP-19357) COMPILATION OF THE RESULTS OF MONITORING ^{90}Sr AND ^{137}Cs DUE TO FALLOUT IN THE ARGENTINE REPUBLIC. (Comision Nacional de Energia Atomica, Buenos Aires (Argentina), Division Estudios Ambientales, 1972, 11p. (In Spanish). (RS-34/66). Dep. NTIS (U. S. Sales Only).

For more than ten years systematic measurements on the levels of contamination of the Argentina biosphere due to radioactive fallout have been carried out. The results of these measurements with reference to strontium-90 and cesium-137 are given. (tr-auth)

45521 DEPENDENCE BETWEEN RADIOACTIVITY AND CONCENTRATION OF ATMOSPHERIC AEROSOLS. Glowiak, Bohdan J.; Palczynski, Ryszard J. (Technical Univ., Wroclaw), pp 155-61 of Proceedings of the 1972 Clean Air Conference. Parkville, Australia; Clean Air Society of Australia and New Zealand (1972).

From International clean air conference; Melbourne, Australia (15 May 1972). See CONF-720541.

An investigation of the natural radioactivity of atmospheric aerosols and the degree of dust concentration in the air in the Wroclaw area was carried out over a two-year period. Air was pumped through a blotting paper band filter and the radioactivity of precipitated aerosols was measured by means of a Geiger-Mueller counter. It was found that the solid air pollutants showed radioactive properties, caused, first of all, by the presence of natural isotope products of radon and thoron decay. Radioactive concentrations within the limits from 0.05×10^{-10} to 12.18×10^{-10} $\mu\text{Ci}/\text{cm}^3$ with a mean value of 1.7×10^{-10} $\mu\text{Ci}/\text{cm}^3$ were observed. The concentration of aerosols in the air investigated was measured by means of a photometric method which determined the degree of filter blackening. Measurements were expressed in dimensionless units based on a pattern scale. The summer and winter measurement results are discussed separately. Correlation coefficients, standard deviations and the parameters of the regression equations of each measurement series were calculated. The investigation showed that the dependence between the concentration of atmospheric aerosols and their radioactivity could be described by exponential equations with correlation coefficients ranging from 0.708 to 0.853. (auth)

45522 ESTIMATION OF THE TURBULENT DIFFUSION COEFFICIENT AND VERTICAL COMPONENT OF WIND VELOCITY ACCORDING TO NATURAL RADIOACTIVITY DISTRIBUTION. Amiranashvili, A. G.; Okroashvili, G. G.; Khundzhua, T. G. (Inst. of Geophysics, Tbilisi, USSR). Soobshch. Akad. Nauk Gruz. SSR; 66: No. 3, 585-7 (Jun 1972). (In Russian).

A formula was obtained for determining the turbulent diffusion coefficient and vertical component of wind velocity. The calculations were made under the premise that the Rn present was in equilibrium with its decay products. Values were estimated for territory of Georgia, USSR. (auth)

45523 DATA. SECTION III. AIR AND DEPOSITION. Radiat. Data Rep.; 13: No. 6, 367-76 (Jun 1972).

Data for Feb. 1972 on the gross beta radioactivity in surface air and atmospheric precipitation sampled at 70 U.S., 24 Canadian, and 12 Pan American sampling stations and at 14 air sampling stations in California are presented. (L.C.L.)

45524 STUDIES OF THE METEOROLOGICAL INFLUENCES ON ATMOSPHERIC RADIOACTIVITY AND ELECTRICITY AT GROUND LEVEL. Israelsson, Sven. Abstracts of Uppsala Disertations from the Faculty of Science, No. 200. Uppsala; Univ. of Uppsala (1972). 36p.

Thesis.

The influences of meteorological conditions upon the atmospheric radioactivity and its connection with the atmospheric-electric phenomena near the ground were studied. The subjects discussed are divided into four categories; exhalation of radon from the ground; natural radioactivity in the air near the ground; artificial radioactivity in the air near the ground, and the relation between atmospheric electric conductivity and radioactivity in the air near the ground. Summaries are presented of ten papers entitled: natural radioactivity in the soil gas and exhalation of natural radioactivity from the ground surface; radon and thoron in the air near the ground; natural α -activity in the air near the ground; measurements of the β -activity in the air near the ground; natural β -activity in the air near the ground; meteorological influences upon the artificial β -activity in the air near the ground; measurements of the natural β -activity and the atmospheric polar conductivity at ground level; variations of electric conductivity and small-ion density in the air near the ground; and conductivity and concentration of small-ions in the air near the ground. (L.C.L.)

Soil

45525 (HASL-257, pp 1,71-94) ^{90}Sr FALLOUT IN UTAH. Hardy, E. P.; Volchok, H. L.; Krey, P. W. 1 Jul 1972.

In Fallout program.

The annual deposition rate of ^{90}Sr (mCi per km^2 per year) derived from monthly precipitation samples was the same at Salt Lake City and New York City from 1957 to 1971. ^{90}Sr deposition concentrations (mCi per km^2 per cm of rain) based on soil analyses up to 1971 were higher in Salt Lake City and other regions of Utah than in other areas of the United States. These data along with ^{90}Sr to ^{90}Sr ratios at appropriate sampling locations indicated that the stratosphere is the major source of ^{90}Sr fallout in Salt Lake City, although about 16 mCi per km^2 or 10 percent of the ^{90}Sr debris probably came from activities at the Nevada Test Site prior to 1957. A limited topographic study in Utah in 1971 indicated that the highest ^{90}Sr deposition concentrations were found at sites closest to the windward side of a mountain front whereas the lowest concentrations were found at high elevation sites within the mountain range. However, it is believed that topography plays a minor role and the preferential stratospheric intrusions are more likely the cause of the high deposition concentrations in Utah. (auth)

45526 (HASL-257, pp 1,95-118) PLUTONIUM FALLOUT IN UTAH. Hardy, E. P.; Krey, P. W.; Volchok, H. L. 1 Jul 1972.

In Fallout program.

Accumulated plutonium fallout in the north central and southeastern sections of Utah was measured in soil samples collected in June 1971. The maximum values in Utah of the deposition (mCi ^{239}Pu and ^{240}Pu per km^2) and deposition concentration (mCi per km^2 per cm of precipitation) were 2.2 and 3.8 times the maximum values found anywhere else in the United States. By comparing the $^{239,240}\text{Pu}$ to ^{90}Sr activity ratios of the Utah soils with the average activity ratio of 32 Northern Hemisphere soils collected in 1970-71, it was estimated that up to 60 percent of the total $^{239,240}\text{Pu}$ activity deposited at some Utah sites was from a source other than the stratospheric reservoir. Soil samples taken at a site near the University of Utah from 1959 through 1971 revealed that the excess plutonium was delivered prior to 1959. Mass isotopic analyses indicate that the Nevada Test Site is the probable second source and that about 3.5 mCi $^{239,240}\text{Pu}$ per km^2 or twice the level expected from global fallout, deposited in the Salt Lake City area from NTS. Excess plutonium was not evident at sites north of Salt Lake City and due east in the valley regions beyond the Wasatch range. (auth)

45527 (TID-26103) MEASUREMENT OF RADIOACTIVE FALLOUT BY FIELD TOTAL ABSORPTION GAMMA RAY SPECTROSCOPY. Report No. 3. Foote, R. S. (Texas Instruments Inc., Dallas, Science Services Div.). 30 Sep 1964. Contract AT(40-1)-3147, 75p. Dep. NTIS.

One and three-fourths years of continuous operation of the field gamma ray spectrometer has demonstrated: gamma ray emitting fission product fallout may be accurately measured and results reported within approximately 24 hr; near-surface amounts of the natural gamma emitters ^{214}Bi , ^{208}Tl , and ^{40}K can be measured with sufficient precision to study quantitatively the effects of surface water absorption, diurnal variations caused by meteorological effects on radon gas diffusion up to the ground surface and the fallout of radon daughter elements in rain showers; instrumental errors can be minimized by computer correction of the data by internal calibration using photopeaks which appear in all spectra; the ^{214}Bi can show daily variations as great as 100% caused by the diurnal radon "breathing" of the soil; absorption of natural gamma radiation from soils by increased soil moisture content is much greater than absorption of artificial fallout radiation which originates within a thin surface layer; the components of the field spectra have been adequately identified and accounted for down to a photopeak energy of 0.5 MeV, as shown by the residue spectra; ^{54}Mn and ^{125}Sb have been determined to be (or to have been) present in the gamma ray environment; and the region in energy about 0.5 MeV is believed to be a mixture of emitters which provide 0.5 MeV plus amounts of other gamma energies. Basically this is believed to be annihilation radiation from near-surface sources and from cosmic radiation as well as from ^{106}Rh and ^{103}Ru . The majority of this radiation has been considered to be annihilation radiation for the data reduction. Fallout measurements during the first half of the calendar year 1964 indicate that: no fallout of newly created isotopes occurred; ^{144}Ce - ^{144}Pr and ^{95}Zr - ^{95}Nb continued to decay according to their respective half lives during periods of no rain; ^{137}Cs continued to increase; ^{54}Mn has increased very slightly over the last one and one-half years; and ^{125}Sb has increased 41% since 1 Jan. 1964. Comparison of the matrix and least squares data reduction methods indicated that: the matrix technique is the most efficient in analyzing fallout data

where the components have half-lives greater than several days and may be identified by study of successive residue spectra; and the more time consuming and complex least squares method is best applied with less well known mixtures involving isotopes with shorter half lives. Analysis of data based on the 11×11 matrix reduction and upon a single record per day shows the pattern of buildup and decay of fallout radioisotopes detected over the one and one-half year period. (auth)

45528 DEPENDENCE OF IODIDE SORPTION ON VARIOUS SOIL COMPONENTS. Knaelmann, M. (Technische Univ., Hanover). Atomkernenergie; 19: No. 4, 329-32(1972). (In German).

The sorption of radioiodine from aqueous sodium iodide solutions by soils, which is important from the view of radiation protection, depends on the iodide concentration and the kind of soil. Data of previous experiments are used to test the relationship between several soil characteristics and the relative sorption. The method for determining the anion exclusion volume, which was used to calculate the sorption data, is compared with another method. The partial correlation coefficients calculated by these sorption data and the significant correlations are shown. (auth)

45529 GAMMA RADIATION OF SOME PLANTS AND SOILS FROM A URANIFEROUS AREA IN NEW ZEALAND. Whitehead, Neil E.; Brooks, Robert Richard; Coote, G. E. (Massey Univ., Palmerston North, N. Z.). N. Z. J. Sci.; 14: No. 1, 66-76(1971).

The amounts of several γ -emitters in plants and associated soils from a uraniferous area were determined by γ spectrometry with the use of a germanium detector operating in the low energy region. The following isotopes were absorbed by plants: ^{210}Pb , ^{227}Ac , ^{238}U , and ^{226}Ra . The average plant soil ratios (dry wt) were 0.092, 0.170, 0.197, and 0.815, respectively. The contribution of each radionuclide and its daughters to the radioactivity of the plant samples was 2.2, 3.3, 81.5, 1.9, and 10.4%, respectively. The remainder, 0.7%, was due to ^{228}Th and its daughters. (Euratom)

Water

Refer also to abstract: 45924.

45530 (BNWL-SA-4205(Rev.)) OCEANIC DISTRIBUTIONS AND RELATIONSHIPS OF ^7Be AND FISSION PRODUCTS. Silker, W. B.; Young, J. A.; Petersen, M. R. (Battelle Pacific Northwest Labs., Richland, Wash.). 1972. 18p. (CONF-720708-6). Dep. NTIS.

From Symposium on the Interaction of radioactive contaminants with the constituents of the marine environment; Seattle, WA. (10 Jul 1972).

The vertical and horizontal distribution of ^7Be in the world's oceans provides a potential basis for estimation of the distributions of other conservative fallout radionuclides as a function of time and space. Such information is essential to any study of biogeochemical processes occurring within the marine biosphere. Surface air concentrations of ^7Be , a cosmogenic radionuclide, vary seasonally with the northern hemisphere maximum occurring during June and the minimum in December, however the concentrations are uniform from year to year. If the transfer of ^7Be across the air-sea interface is proportional to its air concentration, then ^7Be in surface water should parallel the seasonal atmospheric fluctuation, and the water column would reflect the transfer during the preceding months. The ratio of the surface concentration of ^7Be to its water column inventory, for twenty-one northern hemispheric profiles, did indeed exhibit a strong seasonal change which was independent of latitude, longitude, and year of sampling. Highest ratios were observed during July and August and the lowest during the early spring months, in accordance with the pattern that would be predicted from the variation in air concentration. It is estimated that the ^7Be water column inventory can be determined within $\pm 10\%$ at the 95% confidence limits from a single measurement of the surface water concentration, and the annual deposition of ^7Be be approximated within $\pm 20\%$. Oceanic deposition of a given fission product radionuclide should be directly related to the proportionality between its measured air concentration and that of ^7Be . Thus, the cumulative input of fission products with half-lives comparable to ^7Be could be determined by applying the time integrated ratios of fission product to ^7Be air concentration to the measured concentration of ^7Be in surface seawater; this would require appropriate corrections for relative decay rates and vertical mixing within the water column. A more complicated system occurs in the case of longer lived fission products because advective water mass transport becomes an important factor in the oceanic distributions. Eventually, air concentration ratios of fission products and ^7Be , together with surface water concentrations of ^7Be would be employed in a dynamic model of ocean mixing and circulation

to predict the temporal and spatial distributions of long-lived radionuclides in the north Pacific Ocean. Once completed and verified, this model should provide a useful tool for predicting future oceanic distributions of radioactivity from nuclear weapons testing. (auth)

45531 (COO-3563-3) CONCENTRATIONS AND DISTRIBUTIONS OF LONG-LIVED FALLOUT RADIONUCLIDES IN OPEN OCEAN SEDIMENTS. Noshkin, V. E.; Bowen, V. T. (Woods Hole Oceanographic Institution, Mass.). 1972. Contract AT-(11-1)-3563. 22p. (CONF-720708-4). Dep. NTIS.

From Symposium on the interaction of radioactive contaminants with the constituents of the marine environment; Seattle, WA. (10 Jul 1972).

Analyses of deep open-ocean sediment cores from the north and south Atlantic and the Mediterranean Sea showed the presence of ^{90}Sr , ^{137}Cs and $^{239,238}\text{Pu}$ in measurable amounts distributed to depth within the sediment columns. Comparison of the quantities of these radionuclides accumulated in the sediments with either estimated integrated deliveries to the sea surface or the measured inventories in the overlying water masses has been used to estimate the residence times of the radionuclides in the water column. For ^{90}Sr and ^{137}Cs these computed residence times were orders of magnitude shorter than those estimated for the stable elements. Plutonium was removed from the water column significantly more rapidly than either ^{90}Sr or ^{137}Cs . The ratio of ^{137}Cs to ^{90}Sr in the sediments was higher than that found in precipitation, suggesting that a larger fraction of ocean water ^{137}Cs is associated with sedimenting particles than has been previously assumed. A simple model assuming plutonium to associate with a mixed population of particles, sinking at rates of 392 to 70 m per year, predicts very well both the relation of sediment inventory of plutonium to depth of sediment, and the plutonium distribution vertically in the water column. (auth)

45532 ^{90}Sr IN THE GREAT LAKES: CONCENTRATION-TIME MODEL. Lerman, Abraham (Northwestern Univ., Evanston, Ill.). J. Geophys. Res.; 77: No. 18, 3256-64(20 Jun 1972).

Concentration-time models for ^{90}Sr in the Great Lakes deal with transient concentrations of a radionuclide in a chain of mixed lakes and in a chain of periodically stratified lakes. In the model, the lakes receive time-dependent input of ^{90}Sr from atmosphere and land (0 to 10% of annual deposit in the land drainage basins). Removal of ^{90}Sr from lake water is by outflow and decay. In the five Great Lakes computed concentrations of ^{90}Sr rise from nil (assumed) in 1953 to a maximum in 1964 to 1965, followed by a gradual decline until 1969. Computed concentrations are comparable to the range of reported values. However, a closer agreement between the model and reported concentrations requires the rates of input in 1962 to 1964 to be higher than the mean, as well as the rates of removal to be faster (by 6 to 8%) than the removal by decay and mean outflow. Seasonal stratification of the lakes has virtually no effect on mean annual concentrations of ^{90}Sr . (auth)

45533 SHALLOW-WATER ^{90}Sr ANOMALY ABOUT THE ANTILLES ARC, 1970. Bowen, Vaughan T.; Metcalf, W. G.; Burke, John C. (Woods Hole Oceanographic Institution, Mass.). J. Mar. Res.; 30: No. 1, 112-20(5 Jan 1972). (NYO-2174-124).

Vertical profiles about the southeastern approaches to the Caribbean in early 1970 have shown a consistent ^{90}Sr inversion, with the maximum concentrations at depths of about 100 m. It appears that four water masses may be involved in this area, in a very complicated mixing and overlayering phenomenon. (auth)

45534 RADIOLOGICAL SURVEYS OF PEARL HARBOR, HAWAII, AND ENVIRONS, 1966-1968. Cahill, Daniel F.; Harvey, H. D. Jr.; McCurry, D. C.; (and others) (Eastern Environmental Radiation Lab., Montgomery, Ala.). Radiat. Data Rep.; 13: No. 6, 323-34(Jun 1972).

From Apr 1966 through June 1968, the Eastern Environmental Radiation Laboratory of the Environmental Protection Agency (formerly the Southeastern Radiological Health Laboratory, Bureau of Radiological Health, U.S. Public Health Service) conducted radiological surveys of a number of U.S. ports which service nuclear-powered submarines. These surveys were undertaken to determine whether nuclear submarine operations, which include berthing, repair, and servicing, had resulted in radioactivity levels in the harbor which could contribute a detectable radiation exposure to the public. Six surveys were conducted at five ports with histories of extensive nuclear vessel activities, namely: Thames River, New London, Conn.; Hampton Roads, Va., including Portsmouth, Norfolk, and Newport News, Va.; San Diego Harbor, Calif.; Mare Island Naval Shipyard, Vallejo, Calif.; and Pearl Harbor, Hawaii. The conduct of the onsite surveys performed at Pearl Harbor in 1966 and 1968 is described and the results of the laboratory analyses of numerous environmental samples are pre-

sented. It is concluded that: small but detectable quantities of ^{60}Co were found exclusively in the sediment phases of the harbor; neither the harbor water nor any other environmental samples contained detectable ^{60}Co activity; the total amount of ^{60}Co in the harbor sediment is small compared to the natural radioactivity present in Pearl Harbor and decreased significantly from 1966 to 1968; the levels of ^{60}Co activity in the harbor were well below levels that could contribute a detectable radiation exposure to the public and do not constitute a hazard to public health or the environment; and investigation of the physical and chemical characteristics of ^{60}Co in marine sediments indicates that ^{60}Co and possibly other neutron-activated corrosion products are released in a form which is not biologically available to marine or other forms of life. (auth)

45535 RADIOACTIVITY IN LAKE MICHIGAN WATER AND BIOTA, VICINITY OF BIG ROCK POINT NUCLEAR REACTOR, 1960-69. Fetterolf, Carlos M., Jr.; Seeburger, D. James (Michigan Dept. of Natural Resources, Lansing). pp 211-18 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

Since two years prior to the startup of the Big Rock Nuclear Plant the aquatic biota has been monitored for gross β radioactivity at the discharge and at two remote control stations. Background levels, which were similar at all stations prior to criticality, fluctuated due to atmospheric fallout from nuclear weapon testing. Since criticality, radioactivity of aquatic biota at control stations has gradually decreased to near the minimum preoperational levels while radioactivity of biota in the discharge vicinity has gradually increased to near or slightly above the preoperational peaks caused by weapon testing. Surface water samples have been collected statewide and analyzed for gross β radioactivity during the past nine years. Radioactivity of surface water in the reactor vicinity is similar to that at other Michigan locations. (auth)

45536 RADIONUCLIDE CONCENTRATIONS NEAR THE BIG ROCK POINT NUCLEAR POWER STATION. Nelson, Donald M.; Romberg, Gerald P.; Prepejchal, William (Argonne National Lab., Ill.). pp 268-76 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

During the summer and fall of 1970, the concentrations of gamma-ray emitting nuclides were determined in samples of water, sediments, and organisms collected near the Big Rock Point reactor (Charlevoix, Michigan). Reactor-produced ^{65}Zn , ^{137}Cs , ^{134}Cs , and ^{60}Co were measurable at distances of up to a few kilometers along the shore in crayfish, water snails, Potamogeton sp., Ranunculus sp., Chara sp., Cladophora sp. and periphyton. In general, concentrations at 1 km are less than one-tenth those found in the same organism at the discharge point, with somewhat higher concentrations being found toward the east (into Little Traverse Bay). Water and bottom samples were found to be quite variable, with concentrations in bottom sands being somewhat lower than the concentrations in organisms living at the same place. Several samples of small fish were also taken from the discharge channel. One meal per day of the most radioactive of these (bluegill) would contribute about 1% of the permissible daily ingestion of these four nuclides. (auth)

45537 DATA, SECTION II. WATER. Radiat. Data Rep.; 13: No. 6, 360-6 (Jun 1972).

Data are presented on the gross radioactivity in surface waters sampled at 25 locations in the U.S. during Oct. 1971; ^{90}Sr content of surface waters sampled at 5 locations in the U.S. during Jan. to Apr. 1971; monthly gross beta radioactivity and radioisotope content of California water supplies sampled during 1970; and the gross radioactivity of California sewage samples measured during the same year. (L.C.L.)

Ecosystems and Food Cycles

Refer also to abstracts: 45478, 45762, 45808, 45903, 45909, 45914, 45919, and 45926.

45538 (HASL-257, pp I,51-70) ^{90}Sr IN HUMAN BONE: 1971 RESULTS FOR NEW YORK CITY AND SAN FRANCISCO. Bennett, B. G. 1 Jul 1972.

In Fallout program.

Results are presented of determinations of ^{90}Sr content of 254 specimens of human vertebrae obtained during 1971 in New York City and San Francisco. The average ^{90}Sr to Ca ratios for adult

vertebrae, 0.80 pCi/g Ca in San Francisco and 1.40 pCi/g Ca in New York represent 7 to 10% decreases from average values of the previous year. Average ^{90}Sr concentrations in children's bones were somewhat higher than the adult values. An improved bone model has been formulated to describe the variations in ^{90}Sr content of bone and provide correlation with dietary ^{90}Sr intake. The two compartment model accounts for both short and long term retention of ^{90}Sr in bone. Regression analysis of the 11 years of survey data provides values of the relative retention of dietary ^{90}Sr and the effective bone turnover rate. These parameters of the bone model are constant for adults but age dependent for children. The bone model gives satisfactory description of observed bone ^{90}Sr levels and allows reliable assessment of the long-term behavior of ^{90}Sr in man. (auth)

45539 (HASL-257, pp II,1-6) HASL FALLOUT PROGRAM DATA. 1 Jul 1972.

In Fallout program.

The results of measurements of the ^{90}Sr content of foods purchased in New York City and San Francisco in the first quarter of 1972 are given. Estimates of the daily intake of ^{90}Sr based on these measurements and on the revised consumption statistics given in most recently available U. S. Dept. of Agriculture Report (U.S.D.A. ARS62-16) are also listed. (P.C.H.)

45540 (HASL-257, pp III,45-52) ^{137}Cs IN VARIOUS CHICAGO FOODS. (Collection Months, January and April 1972). Brar, S. S. (Argonne National Lab., Ill.). 1 Jul 1972.

In Fallout program.

Since April 1961, the ^{137}Cs and potassium content of the Chicago portion of Tri-City Diet Sampling Program has been determined in bulk food samples by gamma spectrometry using a 4" x 4" NaI (Tl) crystal. The individual food components were counted for a minimum of 100 minutes, and from these measurements composite daily and yearly food intakes were obtained. Cesium-137 activity in food, now, is an order of magnitude lower than it was a few years ago; consequently, a new procedure for these measurements has been adopted in order to improve accuracy. The same variety of foods (all fresh vegetables, all fresh fruits, etc.) are composited before measurement, and the samples are counted a minimum of 400 minutes. The results for the January and April 1972 quarters are tabulated. (auth)

45541 (NYO-2174-130) PLUTONIUM IN NORTH ATLANTIC OCEAN ORGANISMS; ECOLOGICAL RELATIONSHIPS. Noshkin, Victor E.; Bowen, Vaughan T.; Wong, Kai M.; Burke, John C. (Woods Hole Oceanographic Institution, Mass.). 1971. 19p. (CONF-710501-55). Dep. NTIS.

From third national symposium on radioecology; Oak Ridge, Tenn. (10 May 1971).

A considerable series of north Atlantic Ocean organisms were analyzed, by radiochemistry and alpha spectrometry, for fallout plutonium-238, -239, and -240. Organisms from the near-shore environment were selected to show the effects, on plutonium uptake, of variations in feeding habits, association with sediment or with absorptive surfaces, and of trophic level. In general plutonium concentrations are higher in organisms feeding on sediment or on surfaces than in those drawing largely on the water itself. There is some evidence that plutonium concentrations are higher in organisms of higher trophic levels. (auth)

45542 (PNE-529) ECOLOGICAL AND ENVIRONMENTAL EFFECTS FROM LOCAL FALLOUT FROM SCHOONER. 2. THE BETA AND GAMMA RADIATION EFFECTS FROM CLOSE-IN FALLOUT. Rhoads, William A.; Kantz, Asher D.; Ragsdale, H. L. (EG and G, Inc., Goleta, Calif.). 30 Jun 1972. Contract AT(29-1)-1183. 67p. Dep. NTIS.

Project Schooner, a nuclear cratering event with approximately eight to ten times the yield of two previous Nevada Test Site cratering events, Palanquin and Cabriolet, offered the opportunity to further investigate radiation doses and short-term effects to vegetation and the environment, where a larger affected area was anticipated and where topography might be expected to further influence the results of close-in fallout. Specially constructed dosimeters were placed at 92 locations, forming an arc with a radius of approximately 1.7 to 2.0 km, beginning about 12° west of north with respect to GZ. This arc was about 3.35 km long with a mean distance between dosimeter stations of 36 m. Dosimeters were positioned in vertical arrays at 25 cm, 1 m, and 3 m above the surface away from shrubs, as well as on the soil surface, and on shrubs. To determine whether preventing direct fallout from reaching the shrubs would protect them, polyethylene sheets were placed over shrubs at alternate stations. Twelve days after detonation the dosimeters were removed from the field and the protective sheets were removed. No effects were observed on the vegetation except a dusty covering on the unprotected shrubs. In April (D plus 4 months) the first effects attributable to radiation were noted. During the following months at the center of the fallout

pattern all *Artemisia* shrubs lost their leaves and died except those which had been covered with plastic sheets. Elsewhere along the arc there was a "skirting effect" in which the lower parts of larger shrubs were defoliated while the tops remained near normal. Small shrubs were completely killed. Increasingly larger shrubs were killed with higher doses. Beyond the dosimetry arc, a helicopter survey allowed an assessment of the extent of vegetation damage. South of the crater, study sites were established in the most diverse environments resulting from the crater formation. Vegetation analysis at these sites showed successional changes which might be expected to return the area to conditions approximating undisturbed conditions. (auth)

45543 (RLO-2225-T-1-9) DECLINE OF ^{65}Zn IN MARINE MUSSELS FOLLOWING THE SHUTDOWN OF HANFORD REACTORS. Seymour, Allyn H.; Nelson, Victor A. (Washington Univ., Seattle, Lab. of Radiation Ecology). 1972. Contract AT(45-1)-2225. 15p. (SM-158/16; CONF-720708-7). Dep. NTIS.

From Symposium on the interaction of radioactive contaminants with the constituents of the marine environment; Seattle, WA. (10 Jul 1972).

The shutdown of the last of the eight original plutonium production reactors at the Hanford Atomic Plant in February 1971 eliminated the source of practically all of the radionuclides in the Columbia River and in the plume of Columbia River water in the Pacific Ocean. Zinc-65, perhaps the most biologically important of the Hanford-produced radionuclides in the marine environment, continues to be present in both the biota and the water fifteen months after shutdown. The amounts of ^{65}Zn in mussels at the mouth of the Columbia River and along the Washington Coast have been monitored since 1961, and data from these observations have been used as reference values to measure the loss of ^{65}Zn related to the shutdown of the reactors. The amount of ^{65}Zn in the soft tissues of *Mytilus edulis* from the North Jetty at the mouth of the Columbia River declined from 40 pCi/g of dry tissue in February 1971, to 6 pCi/g, dry, in May 1972. The effective half-life for ^{65}Zn was calculated to be 130 days and the loss of ^{65}Zn by processes other than physical decay, in terms of half-life, was calculated to be 277 days. The latter value is called ecological half-life rather than biological half-life, since there is some uptake of recycled ^{65}Zn but at a rate less than the rate of loss of ^{65}Zn by biological processes. At North Head, 5 km north of the Columbia River, the amounts of ^{65}Zn and Zn in *M. californianus* and the rate of decline was less than comparable values for North Jetty mussels, although the specific activity values, $\mu\text{Ci } ^{65}\text{Zn/g Zn}$, for mussels from both locations, collected on the same day, were similar. The ^{65}Zn effective half-life value for North Head mussels was 149 days. The amount of ^{65}Zn in mussels from seven locations on the Washington Coast and in the Strait of Juan de Fuca, to a distance of 340 km from the Columbia River, declined in a regular manner in respect both to distance from the Columbia River and to time after reactor shutdown. (auth)

45544 (RLO-2225-T-14-4) NATURAL ^{210}Pb AND ^{210}Po IN A MARINE ENVIRONMENT. Schell, W. R.; Jokela, T.; Eagle, R. (Washington Univ., Seattle, Lab. of Radiation Ecology). 1971. Contract AT(45-1)-2225. 29p. (SM-158/47; CONF-720708-2). Dep. NTIS.

From Symposium on the interaction of radioactive contaminants with the constituents of the marine environment; Seattle, WA. (10 Jul 1972).

A regional sampling program was initiated to determine the existing levels of ^{210}Pb and ^{210}Po in sediment, organisms, and water from different depths in a salt water sound and a fresh water lake. The collections were made with plankton nets, midwater trawls, sediment corers, and a large volume water sampler consisting of 0.3 μ Millipore filters and Al_2O_3 sorption beds (200 to 2,000 liters). The efficiency of collecting lead from fresh and salt water by the Al_2O_3 beds was determined. The sample analysis was made using ^{210}Po tracer for yield determination and low background alpha spectroscopy which separates the energies of ^{210}Po from ^{210}Pb ; the ingrowth and decay of ^{210}Po were used to determine ^{210}Pb . Computer programs were completed for data analysis, sorting, and retrieval. One interpretation of the early results of sample analysis from the fresh water and salt water environments was that the processes of erosion and leaching can contribute significantly to the concentration of ^{210}Pb and ^{210}Po in the aquatic environment. Regional differences were found which depend on the geological strata. Both the erosion process and the atmospheric fallout process must be considered as sources of these radionuclides in the aquatic environment. An unusually large concentration of ^{210}Po was found in the liver of a sablefish, *Anoplopoma fimbria*; the zooplankton in the same region also had a high ^{210}Po concentration. The nektonic crustacea have the highest $^{210}\text{Po}/^{210}\text{Pb}$ ratio of the organisms collected at ca 100. (auth)

45545 (UCRL-73744) ACCUMULATION AND LOSS OF COBALT AND CESIUM BY THE MARINE CLAM, *MYA ARENARIA*, UNDER LABORATORY AND FIELD CONDITIONS. Harrison, F. L. (California Univ., Livermore, Lawrence Livermore Lab.). 6 Jul 1972. 35p. (CONF-720708-7). Dep. NTIS.

From Symposium on the interaction of radioactive contaminants with the constituents of the marine environment; Seattle, WA. (10 Jul 1972).

The accumulation of ^{60}Co and ^{137}Cs by clams was followed in the laboratory in aquarium systems in which the levels of radioactive and stable cesium and cobalt were held constant. At 3- to 21-day intervals during the 6-month accumulation period, groups of 8 animals were removed from the water and their soft tissues were dissected into 7 different organs and tissues. The concentrations of both ^{60}Co and ^{137}Cs in the edible tissues increased rapidly and then leveled off at concentrations of ^{60}Co that were about 200 times that of the water and of ^{137}Cs that were about 5 times that of the water. Variations in concentration among the body parts were larger for ^{60}Co than for ^{137}Cs . The effects of stable element concentration in water on the accumulation rates were assessed in groups of animals maintained in seawater containing 0.5, 2.5, and 12.5 $\mu\text{g/litre}$ each of stable cobalt and cesium. Little regulation of cesium accumulation was observed at these concentrations; the rates of accumulation increased with the concentration in water. Some regulation of cobalt occurred at the higher concentrations. The accumulation of ^{60}Co and ^{137}Cs was followed also in clams introduced into a discharge canal receiving radioactive waste from a boiling-water reactor. Groups of 6 animals were sampled at 3-week intervals and concentrations were measured in the pooled, total soft tissues. Results obtained in the field will be compared to those obtained in the laboratory. Loss of ^{60}Co and ^{137}Cs was followed in animals that had accumulated a radioactive burden in the laboratory and then were transferred to unfiltered oceanic water at a marine station. Loss of ^{60}Co by the whole clam was monophasic, that of ^{137}Cs was biphasic. The kinetics of accumulation and loss were described by exponential equations. The rate constants of accumulation and loss and the steady-state concentration factors are being used in models to predict the dose to man from the consumption of animals exposed to radioactive releases. (auth)

45546 ARTIFICIAL RADIOACTIVITY OF FALLOUT AND FOOD IN THE NORTHERN DISTRICTS OF GDR IN THE YEAR 1969. Schulze, H.-A.; Randow, F. (Univ., Rostock, Ger.). Z. Gesam. Hyg. Ihre Grenzgeb.; 17: 255-7 (Apr 1971). (In German).

The radioactive nuclides ^{90}Sr and ^{137}Cs that occurred in 1969 in milk and cereals in the three northern counties of the GDR (Mecklenburg) and in oversea imports were measured. Apart from the nuclides, the total- β -activity was determined in the fallout. The annual amounts of fallout radioactivity were 26.12 nCi β/m^2 , 0.46 nCi $^{90}\text{Sr}/\text{m}^2$, and 1.18 nCi $^{137}\text{Cs}/\text{m}^2$ in Rostock and were, with the exception of the total- β -activity which remained unchanged, below the values of last year with 54 percent and 76 percent, respectively. The contamination of milk amounted to 5 pCi $^{90}\text{Sr}/\text{l}$ and 15 pCi $^{137}\text{Cs}/\text{l}$, and that of cereals to 22 pCi $^{90}\text{Sr}/\text{kg}$ and 38 pCi $^{137}\text{Cs}/\text{kg}$. Citrus and other tropical fruit contained 4 to 52 pCi $^{90}\text{Sr}/\text{kg}$ and 1 to 17 pCi $^{137}\text{Cs}/\text{kg}$. Tea contained 600 to 4,400 pCi $^{90}\text{Sr}/\text{kg}$ and 500 to 800 pCi $^{137}\text{Cs}/\text{kg}$, whereas the values for coffee were 32 pCi $^{90}\text{Sr}/\text{kg}$ and 105 pCi $^{137}\text{Cs}/\text{kg}$. (auth)

45547 METHOD OF EVALUATING CHEMICAL POLLUTION LEVELS IN AQUATIC MEDIA AND FOOD CHAINS. Bittel, R.; Lacourry, G. (CEN, Fontenay-aux-Roses, France). Rev. Int. Oceanogr. Med.; 22: 129-41 (1971). (In French).

The transfer of radioactive pollution in aquatic food chains can be evaluated using a simple mathematical model: $D = \Sigma D(Q, F, X)$ in which D = radiation dose delivered to the organism, Q = ingested quantity of each alimentary carrier, F = transfer factor from the environment to the organism, and X = environmental pollution level. The summation was carried out for the various foods and radionuclides considered. Compliance with the protection standards for man requires $D < D_1$ (D_1 = dose limit). This investigation demonstrates the feasibility of setting up a very simple model for the transfer of chemical pollution in marine food chains. This model simplifies the transfer parameter inventory, and enables the following parameters to be defined: acceptable daily dose; level limit for water; and critical food consumption. Indications are given of future research projects together with their priorities. (France)

45548 CONTENT OF CESIUM IN MUSCLES OF BROILERS, HENS, AND PARTRIDGES. Szulc, M.; Bruniak, A. (Warsaw Agricultural Univ.). Med. Wet. (Poland); 27: 297-9 (1971). (In Polish).

The activity of cesium-137 in muscles of broilers, hens from individual farms, and partridges has been measured. Materials

were gathered from the Warsaw area in the autumn and winter season 1968 to 69. The measurement of radioactivity of ^{137}Cs was performed with gamma spectrophotometry by means of scintillation probe equipped with a NaI crystal and a single channel analyzer of impulse amplitude. The findings were expressed in activity units of ^{137}Cs in ash and muscles of birds examined, i.e., pCi/kg of ash and pCi/kg of fresh muscle. They showed that the muscles of broilers contained the highest content of ^{137}Cs (157.7 pCi/kg of muscle and 13.478 pCi/kg of ash); lower contents were found in muscles of hens (155.3 pCi/kg of muscle and 12.626 pCi/kg of ash), and the lowest in muscles of partridges (143.0 pCi/kg of muscle and 10.362 pCi/kg of ash). In addition, it was revealed that the content of ^{137}Cs in muscles of birds was lower than those in mammals. (NSA of Poland)

45549 PASSAGE OF ^{90}Sr FROM BOVINE BONES TO RECEIVED FOOD PRODUCTS. Szulc, M.; Mierzevska, H. (Warsaw Agricultural Univ.). *Med. Wet.* (Poland); 27: 346-9 (1971). (In Polish).

The basic purpose of the investigations performed in 1967 to 69 was to establish if raw bone material contaminated with radioactive strontium may be used in good processing. The investigations were carried out with bovine cervical vertebra and shoulders derived from normal animal carcasses. Basic technological processes commonly used in the processing of alimentary bones, e.g., boiling in tap water-equivalent of broth production, boiling in acidified water-equivalent of acidified soup production, bone dressing in the course of gelatin proanalysis (method of F. Bryant, A. Morgan, and G. Spicer), and then on the basis of measurement of radioactivity by the use of anticoincidental system were examined. It was found that in the examined processes of bovine bone dressing, supravital contaminated with ^{90}Sr , an average of 11.9%, 14.3% of ^{90}Sr were found in brews, and about 0.0006% in gelatin. The results obtained indicated that bovine bones heavily contaminated with ^{90}Sr cannot be used either for broth or for production of acidified soups; but it seems that they can be used as a raw material for gelatine production. (NSA of Poland)

45550 ACCUMULATION OF ^{90}Sr IN WHEAT ON SOD-PODZOLIC SOILS. Gulyakin, I. V.; Yudinseva, E. V.; Makarov, K. I. *Agrokimiya*; No. 4, 84-92 (1971). (In Russian).

Thirty-two sod-podzolic soils were investigated. A close relation was observed between the ^{90}Sr accumulation in wheat and the quantity of exchangeable Ca in the soil, degree of the soil radionuclide stability, and the ^{90}Sr content in the plants at early developmental stages. The soil Ca/Sr relation was altered by the soil acidity and the amount of humus. (Euratom)

45551 ACCUMULATION AND DISTRIBUTION OF RADIOACTIVE ELEMENTS IN NATURAL PHYTOCOENOSSES OF THE CARPATHIANS. Shevchuk, A. I. (Inst. of Botany, Kiev). *Ukr. Bot. Zh.*; 25: No. 4, 18-24 (1968). (In Ukrainian).

A study was made during 1962 to 1965 on the accumulation and distribution of radioactive isotopes in the natural plant associations of the forest, subalpine, and alpine belts of the Carpathians. The accumulation of β -emitters per unit area was determined in associations dominated by the European spruce forest, Siberian juniper forest, *Sesleria bielzii* grass community, and *Festuca rubra* grass community. The data obtained were correlated with the amount of radioactive fallout during the years in question. The maximum amount of fallout was in 1963, followed by significant successive decreases in 1964 and 1965. The changes in the content of ^{90}Sr in herbaceous associations corresponded to variations in the ^{90}Sr fallout from year to year during the period in question. In arboreal and shrub associations there was a cumulative increase in the content of ^{90}Sr because of retention of this isotope from the fallout of earlier years by perennial organs of the plants. From Sept. 1962 to Sept. 1963 and Sept. 1963 to Sept. 1964 the spruce forest accumulated 55.8 and 60.8% of ^{90}Sr , respectively, that fell on its surface from the atmosphere and the Siberian juniper association (29.8 and 30.2%, respectively). Coniferous tree and shrub associations, by binding a considerable part of the ^{90}Sr and preventing it from subsequent migration, play an important sanitary role at present. In herbaceous associations, the principal amount of radioactive elements from aerial parts of plants is removed by animals during grazing. (Euratom)

45552 DATA. SECTION I. MILK AND FOOD. *Radiat. Data Rep.*; 13: No. 6, 347-59 (Jun 1972).

The monthly averages and 12-month averages of ^{137}Cs and ^{90}Sr content in milk samples collected during Feb. 1972 and during Mar. 1971 to Mar. 1972 at 61 U.S. sampling stations, at 16 Canadian sampling stations, and at 7 sampling stations in Central and South America are tabulated. ^{140}Ba and ^{89}Sr were detected in only 4 samples from U.S. stations and are noted. (L.C.L.)

45553 RADIOECOLOGY OF $\text{Ce-}^{144}\text{Pr}$ IN A LITTORAL FRESH-WATER MICROCOSM. Trabalka, John Richard. *Ann*

Arbor, Mich.; Univ. of Michigan (1971). 225p. University Microfilms Order No. 72-15,021.

Thesis.

Results are reported from a study conducted over a period of 760 days in which natural conditions were simulated in two 5000 liter tanks which contained replicate microecosystems. Starting on day 493, the water in one tank was maintained at a mean concentration of 1.2×10^{-4} microcuries/cc of cerium-144 + praseodymium-144 until day 557. Throughout the study period the biological communities, consisting of over 200 species, were studied comparatively in the two tanks. Populations of the fathead minnow (*Pimephales promelas*) completed two life-cycles before day 341. The fish spawned three times between days 341 and 598; a total of 6464 larvae was examined microscopically for abnormalities. The dose to the fish gonads was estimated to be 16.5 rads, and to the fish embryos, 78 millirads in the treated tank. No statistically significant differences between the treated and control tanks were observed either in the frequency of abnormalities in fish larvae, or in the biological communities. At the end of the experiment, over 99% of the added radioactivity was in the sediment and 0.1% was in the water. (Diss. Abstr. Int., B)

45554 ASSESSMENT AND CONTROL OF ENVIRONMENTAL CONTAMINATION: EXPERIENCE WITH ARTIFICIAL RADIOACTIVITY. Butler, G. C. (National Research Council, Ottawa); Karol, I. L.; Lindell, B. G.; Stevens, D. J.; Zeleny, V. *Biol. Conserv.*; 4: No. 3, 177-84 (Apr 1972).

The basic principles followed in assessing and controlling radioactive contamination of the environment are outlined. These principles largely stem from the nature of the relationship between dose and effects of radiation. Acute effects only occur at doses and dose-rates much higher than those likely to arise from environmental contamination, except in major accidents. For delayed effects, both genetic and somatic, it is assumed that the increment in biological effect per unit increment in dose is the same at all doses, i.e., there is a linear non-threshold dose relation, and the biological effect of a given dose is independent of the rate at which the dose is delivered. These assumptions are conservative since they imply that no doses are "safe" in absolute terms but that all doses involve a risk proportional to their magnitude, and that relying on them will, in most practical cases, result in overestimates of the risk of harm at low doses and low dose-rates. The assumptions justify the use of the concepts of "dose-commitment"—the total average dose to be received by the members of a population, living and to be born, as a result of a release of radioactivity in the environment—and of "harm commitment" (a measure of the total harm expected in that population as a consequence of the dose commitment received). The application of these concepts to the establishment of norms for the protection of the public and of surveillance programmes is discussed. (auth)

45555 CONVENTIONAL AND RADIOACTIVE POLLUTION IN THE ENVIRONMENT. Cigna, Arrigo A. (CNEN, Milan). *Com. Naz. Energ. Nucl., Notiz.*; 18: No. 5, 25-7 (May 1972). (In Italian).

A comparison is made between risks of various kinds in the development of nuclear energy, the levels allowed for peaceful uses, and the natural and artificial types of radiation. A typical exposure for the population of the U.S. is discussed, involving the natural background, diagnostic x rays, fallout from nuclear explosions, color TV sets, luminous dials, and employment in nuclear plants. Risks vs benefits are evaluated, and maximum permissible limits are defined. Causes and probability of death or serious injury resulting from non-radiation causes (auto accidents, drowning, cancer, etc.) are examined. (K.S.W.)

45556 REPORT OF COMMITTEE 4 ON PRINCIPLES OF ENVIRONMENTAL MONITORING RELATED TO THE HANDLING OF RADIOACTIVE MATERIALS. CIPR-7. Fontenay-aux-Roses, France; Centre d'Etudes Nucleaires ([1971]). 26p. (In French).

Translation of Report of Committee 4 on Principles of Environmental Monitoring Related to the Handling of Radioactive Materials, Oxford, Pergamon Press.

Environmental monitoring is the term applied to radiation and radioactivity measurements made outside the limits of installations using radioactive substances or sources. One of the principal objectives of the environmental monitoring program is the evaluation of the real or potential exposure of man to radioactive substance or radiations present in the environment, and the determination of the maximum probable limits of such an exposure. The objectives and the different factors intervening in the development of routine monitoring programs for the exterior of installations were studied. Preliminary studies concerned with the putting into service of these operations enabled information about nuclides, modes of transfer, critical groups reference man, and normal op-

eration to be obtained. Monitoring programs in the case of emergencies (accidental irradiation) are also studied. (France)

Radiometric Techniques

Refer also to abstracts: 45205, 45444, and 45658.

45557 REAPPRAISAL OF THE $^{40}\text{Ar}/^{39}\text{Ar}$ STEPWISE DEGASSING TECHNIQUE. Brereton, N. Robin (Cambridge Univ., Eng.). *Geophys. J. Roy. Astron. Soc.*; 27: No. 5, 449-78 (May 1972).

The principal advantage of the $^{40}\text{Ar}/^{39}\text{Ar}$ dating method lies in the stepwise degassing technique. Despite the widespread use of the technique, each worker seems to adopt an individual criterion of choice, as to how the data should be treated, so as to extract the maximum amount of information from it. A detailed analysis of the data taken during a number of stepwise degassing experiments was made to decide upon the most useful treatment of results. The basic principles of the technique are reviewed, and a method of data analysis is presented, which enables a detailed study of the argon liberation characteristics of the sample to be made. It also allows the age of the sample to be determined, such that no direct correction is made for atmospheric contaminant ^{40}Ar . This avoids the fundamental assumption that all ^{36}Ar , used to make this correction, is of atmospheric origin. (auth)

45558 PRELIMINARY DATA ON THE TRITIUM CONTENT OF AQUIFER SHEETS IN NORTHWEST AND CENTRAL (ALGERIA) SAHARA; HYDROGEOLOGICAL CONSEQUENCES. Conrad, Georges (Rouen Univ., Mont-Saint-Aignan, France); Olive, Philippe. *Compt. Rend., Ser. D*; 274: No. 14, 2019-22 (5 Apr 1972). (In French).

The interpretation of the tritium activities of the principal aquifer sheets was effected using isotopic data collected (since 1964) on precipitations and surface hydrology. The present day surface aquifer recharge linked to the hydrographic network appears at Saoura, the Chech erg and Ahnet-Mouydir. In contrast no rapid contributions of precipitations occurring after 1955 to the feeding of the economically important sheets of the septentrional Sahara has taken place. (France)

45559 WELL LOGGING METHOD FOR DETERMINING THE PRESENCE OF ALUMINUM AND MAGNESIUM. Youmans, Arthur H. (to Dresser Industries, Inc.). U. S. Patent 3,662,173. 9 May 1972. Filed 27 Oct 1969.

An activation well logging method for determining the presence and quantity of aluminum and magnesium in formations surrounding a borehole is described. The formations are irradiated with 14 MeV neutrons which produce aluminum and magnesium activity with half lives of 20 milliseconds and gamma ray energy of 473 keV. The neutron source is turned on for a period of time sufficient to activate aluminum and magnesium and then turned off for 20 milliseconds. During the first 4 milliseconds while the source is off, no measurement is made. During the remaining 16 milliseconds, pulses are counted which fall in the amplitude range representing gamma ray energies from 450 keV to 500 keV. The rate of occurrence of these pulses is an indication of the aluminum and magnesium content of the formation. To correct for undesired background, the rate of occurrence of pulses in the energy intervals immediately above and below the 450 to 500 keV interval is determined and subtracted from the 450 to 500 keV measurement. (Official Gazette)

45560 DEVELOPMENTS IN GEOPHYSICAL LOGGING INSTRUMENTS AND METHODS. Andrassy, L.; Barat, I.; Liszt, Ferenc; (and others). Magy. Allami Eotvos Lorand Geofiz. Int. Evi Jelentese; 1969: 89-93 (1970). (In Hungarian).

More than half the effort of the well logging group of the Geophysical Institute in Hungary in 1962 was devoted to instrumental development. Radioactive logging equipment with gas-discharge and semiconductor detectors was constructed. The surface pulse-processing and control unit, two linear ratemeters, a pulse-separating calibrating generator, a universal stabilized supply unit, and a mass voltage converter have been finished. Scintillation radioactive equipment was further improved. Precision circuits have been designed for better signal transmission in energy-selective borehole instruments. The circuit design of spectrum-frequency logging equipment was given its final shape. Acoustic logging equipment was used in practical measurements in velocity logging, cement-bond logging, and porosity determinations. Work continued on the development of calibrating equipment for radioactive logging apparatus. Activation analysis with a neutron generator is now routine in the laboratory, particularly with an automatic bauxite analyzer designed and built in the Geophysical Institute. Digital well logging apparatus is being developed in co-operation with other institutions. (Geophys. Abstr.)

45561 THERMAL NEUTRON ACTIVATION RADIOACTIVITY LOGGING METHOD. Youmans, Arthur H. (to Dresser Industries, Inc.). U. S. Patent 3,665,195. 23 May 1972. Filed 16 Oct 1969.

A method is described for determining the existence and quantity of an element in a formation. A portion of the formation is irradiated with neutrons and the gamma rays emanating from the formation due to induced radioactivity, or activation from thermal neutron capture by the element of interest are detected and recorded. The thermal neutron capture cross section of the formation is determined. The product of the gamma ray emission and the thermal neutron capture cross section is obtained as a quantitative indication of the element's abundance in the formation. (Official Gazette)

45562 RADIOACTIVITY WELL LOGGING SYSTEM HAVING A PAD MOUNTED NEUTRON SOURCE AND A CENTRALIZED RADIATION DETECTOR THAT PROVIDES COMPENSATION FOR BOREHOLE FLUID DENSITY AND BOREHOLE DIAMETER VARIATIONS. Youmans, Arthur H. (to Dresser Industries, Inc.). U. S. Patent 3,662,172. 9 May 1972. Filed 24 Oct 1969.

A well logging instrument adapted for traversing an earth borehole has a plurality of neutron sources mounted in pads which contact the wall of the earth borehole. The radiation detector is centralized in the borehole and the system provides compensation for natural gamma radiation and variations in borehole fluid density and borehole diameter. (Official Gazette)

45563 PULSE NEUTRON LOGGING TOOL CONTROL AND TRANSMISSION SYSTEM. Frentrop, Arthur H.; Wahl, John S. (to Schlumberger Tech. Corp.). U. S. Patent 3,662,179. 9 May 1972. Filed 25 Apr 1969.

An illustrative embodiment of the invention measures the thermal neutron decay time in earth formations traversed by a borehole. For example, with a latching relay downhole, only one electrical conductor and a ground return is needed to send power and neutron generator control signals to the logging tool. The repetition rate and the duration of the neutron bursts, as well as the decay time measurements, all are regulated automatically through downhole circuits in response to the neutron characteristics of the formation under study. The single conductor and ground return also transmit the decay time measurements to the earth's surface. A gate enabling relay downhole also responds to direct current signals and selects detector count information for transmission to the surface through a manipulation of the thermal neutron decay time information signal polarity. Other information signals, of which casing collar locator signals are typical, also can be transmitted through the conductor. (Official Gazette)

45564 APPLICATION OF WELL LOGGING METHODS IN HYDROCARBON PROSPECTING, DRILLING, AND PRODUCTION TO THE STUDY OF GEOSTATIC PRESSURE PROBLEMS. Fertl, W. H.; Timko, D. J. (Continental Oil Co., Ponca City, Okla.). *Rev. Inst. Fr. Petrole Ann. Combust. Liquides*; 26: No. 9, 687-714 (Sep 1971). (In French).

A number of well logging methods are being used by the oil industry to detect and evaluate overpressure formations. World-wide experience in exploring for oil and gas indicates a significant correlation between the presence and magnitude of geopressures and the log-derived shale-sand ratio of sedimentary sections penetrated by the drill bit. Knowledge of the expected pore pressure and fracture gradient is the basis for making the best of modern drilling techniques. This prevents a breakdown of exposed formations and contains the high-pressure fluids in deeper intervals, thereby reducing blowout hazards. In several sedimentary basins throughout the world, the magnitude of log-derived in situ rock parameters, such as pore-fluid pressures derived from conventional well logs, correlates very well with the distribution and type of hydrocarbon accumulations. Pulsed neutron logging devices can be applied to monitor pore pressure variations behind casing. (France)

45565 RADIOACTIVITY OF WASTE WATERS IN NUCLEAR RESEARCH INSTITUTE, REZ, AND ENVIRONMENT. Krepelka, Jiri (Nuclear Research Inst., Rez, Czech.); Chysky, Jaromir. *Jad. Energ.*; 18: No. 5, 146-50 (May 1972). (In Czech).

A system of drainage of liquid wastes and a method of liquidation of radioactive wastes in the Nuclear Research Institute are briefly described. The radioactivity of waste waters discharged into the river is controlled discontinuously. The gross beta-gamma activity is determined in samples. The control also includes a continuous measurement of the activity of chemical wastes discharged separately. Both methods are adequate in view of the isotopes that might be present in a given mixture. In the period of 1963 to 1971, more than 2000 samples were measured. The values obtained

were very low, which shows a good function of the liquidation station. No effect on the activity of the river Vltava, into which the wastes were discharged, was found. Therefore, a measurable increase in the exposition by ionizing radiation could not occur in the inhabitants using the Vltava water. The continuous measurement showed several times an accidental uncontrolled discharge of the activity into the chemical drainage and into the river. However, it was found that the maximum permissible concentrations for drinking water could not be exceeded in the river Vltava, which was also proved. The results showed that the amount of activity treated and the qualitative isotopic composition need be known for a complex hygienic evaluation of the system of liquidation of wastes in larger nuclear plants. (auth)

45566 DETERMINATION OF DENSITY AND EFFECTIVE ATOMIC NUMBER OF ROCKS AND ORES BY GAMMA-GAMMA METHOD IN THE PRESENCE OF AN INTERMEDIATE ZONE. Artsybashev, V. A.; Ivanyukovich, G. A. (Leningrad Mining Inst.). *Izv. Vyssh. Ucheb. Zaved., Geol. Razved.*; No. 9, 97-101 (Sep 1971). (In Russian).

The density and effective atomic number of rocks and ores were determined by the gamma-gamma method in the presence of the so-called "intermediate zone", formed because of irregularities in the aperture walls, of caverns, crusts, etc. Certain conditions were observed in the measurements: constant geometry and constancy in the source and the detector. However the gamma field depended on the gamma quantum properties of the medium and the intermediate zone: $N = F(\rho, Z_{eff}, h)$, where N is the measured gamma field characteristics, ρ is the density of the medium, Z_{eff} is the effective atomic number, and h is the level of the intermediate zone—the distance from the zone surface to the bore wall. Two variants of the solution to the problem are explained for simultaneous determination of ρ , Z_{eff} , and h . The results of experimental testing on a bore-hole model are interpreted. A 2π probe with a ^{60}Co source and scintillation detector were employed. The body of the probe was made of Al and had a thickness of 3 mm. No supplementary filtration of the gamma radiation was supplied. The initial threshold of the recording of the gamma quanta was 20 to 30 keV. The integral beam of scattered gamma radiation was recorded. (tr-auth)

45567 APPLICATION OF NEUTRON ACTIVATION TO THE DETERMINATION OF ELEMENTS IN BOREHOLES. Wichmann, P. A. *Rev. Inst. Fr. Petrole Ann. Combust. Liquides*; 26: No. 9, 715-29 (Sep 1971). (In French).

Some of the neutron activation logging methods which have been used over the past few years are reviewed. These methods depend almost entirely on the half-lives and radiation intensity of the activation products. The elements which are most conveniently identified in this manner include oxygen, silicon, and aluminum. Field examples of the logs and/or measurements involved in each of these determinations are given. The application and relative value of each of these determinations, the discussion of limitations of the presently available field techniques, and possible solutions of the limiting problems are discussed. The most likely solution of these problems is primarily related to the use of gamma spectral measurements to define the activation products. Field examples of test data supporting the validity of these solutions are also presented. The method is shown to be a viable approach for determining lithologic components (through elemental assay) in well bores containing a wide range of conditions. While having certain limitations, the techniques described have the almost singular advantage of being direct measurements of the elemental constituents under investigation. (France)

45568 ISOTOPE RATIO TRACER METHOD: APPLICATIONS IN ATMOSPHERIC SULFUR POLLUTION STUDIES. Manowitz, Bernard; Smith, Maynard E.; Steinberg, Meyer; Newman, Leonard; Tucker, Walter D. (Brookhaven National Lab., Upton, N. Y.). pp 993-9 of Proceedings of the Second International Clean Air Congress. /England, H. M. (ed.). New York; Academic Press, Inc. (1971).

From second international clean air congress conference; Washington, D. C. (6 Dec 1970). See CONF-701229.

The isotope ratio tracer method (IRT) is based on the natural variation in the ratios of the stable isotopes in sulfur coming from different sources. It is possible with the IRT method to measure changes in the ratio of ^{32}S to ^{34}S that may result from isotope exchange following chemical reaction with sufficient precision to indicate the degree of conversion that has occurred. The use of the sulfur hexafluoride tracer system in conjunction with IRT greatly increases its usefulness. (D.H.M.)

45569 AGE OF CUPROUS UDOKAN SERIES (OLEKMO-VITIM MOUNTAIN SYSTEM) BY URANIUM-LEAD METHOD. Bogdanov, Yu. V.; Iskanderova, A. D.; Levskii, L. K. *Izv. Akad. Nauk SSSR, Ser. Geol.*; No. 9, 29-34 (Sep 1971). (In Russian).

The age estimation of the Udokan series (Olekmo-Vitim mountains) was conducted by means of the radiogenic lead occluded in the rock samples. In order to establish the time of accumulation of the Udokan sediment, the lead content and isotopic composition of six types of sandstones were determined. Both U and Th data were obtained on three of the samples. The isotopic composition of Pb was measured on the MI-1309 mass spectrometer by thermionic emission using a dry aluminum silicate emitter. The U in the sandstones was determined by radiochemical and luminescence methods. The Th was determined by radiochemical and colorimetric (Arsenazo III) methods. The age, from the U-Pb method, of the deposits of the Udokan series was confirmed by argon dating of granitoids—faulted rock of the Udokan series. They verified the correctness of the connection of the copper-bearing carbonate-terigenous strata of the Udokan series with the lower Paleozoic. (tr-auth)

Site Surveys

Refer also to abstract: 45510.

45570 (EUR-4841) SITE CHARACTERISTICS. de Bortoli, M.; Gaglione, P.; Gandino, C. (European Atomic Energy Community, Ispra (Italy). Joint Nuclear Research Center). 1972. 98p. (In Italian). Dep. NTIS (U. S. Sales Only). EUR FB 125,-.

The main features of the Ispra site concerning the meteorological, geologic, hydrologic demographic, and ecological aspects are described. The information was gathered partly from reports available at the Ispra Euratom JRC, partly from the Ufficio Provinciale di Statistica di Varese, and partly from inquiries performed directly by the Site Survey Meteorology Section of the Division Protection. (auth)

45571 (NVO-55) PROJECT WAGON WHEEL: PRELIMINARY SITE EVALUATION REPORT. (Nevada Operations Office (AEC), Las Vegas). Mar 1970. 31p. (PNE-WW-2). Dep. NTIS.

After reviewing currently available data and conducting preliminary on-site investigations, the proposed Wagon Wheel site, at this time, is recommended for further study and investigation as a possible location for a nuclear explosive gas stimulation experiment. In preparing the evaluations of the Wagon Wheel area, it was assumed that the experiment would utilize an explosive having a yield of 100 kt buried at a depth of at least 10,000 feet. It should be understood that when a definitive technical program has been agreed to by industry and government, some revisions to the following statements may be required. Several areas where further detailed studies will be required include: nearby large reservoir sites; mining operations within a 184 km radius; and abandoned coal mines at Rock Springs, Wyoming, 121 km south of Wagon Wheel. A factor which should be given early consideration is the season of the year most suitable for detonation, since each season has some particular limitation. In summer, there is considerable tourist activity. During the fall, the hunting season provides considerable income for this area. The winter season is accompanied by heavy snow and severe cold which might present operational difficulties, particularly if any evacuation was necessary. The spring and early summer seasons, because of melting snow and heavy run-off, have full reservoirs and lakes. Also, heavy ground saturation could increase the potential for triggering earth movement and ground motion. These conditions should be examined closely to determine which would be the least restrictive season for detonation. These statements do not attempt to cover the total program of detailed effects evaluations to be conducted during the project definition phase of the project, but they do reflect the types of efforts which will be involved in the detailed studies of the Wagon Wheel site. (auth)

45572 (WASH-1523) LEASING OF AEC CONTROLLED URANIUM BEARING LANDS, COLORADO, UTAH, NEW MEXICO. Environmental Statement. (Atomic Energy Commission, Washington, D. C.). Mar 1972. 130p. Dep. NTIS.

The USAEC proposes to lease certain AEC-controlled lands in the Colorado Plateau region for the production of uranium-vanadium ores. About 25,000 acres of land are available for leasing in about two dozen tracts, in western Colorado, eastern Utah, and northwestern New Mexico. The lands are located in long established mining districts and most were mined previously under AEC leases until 1962. The known reserves of U-V ores on these lands are substantial and the potential for further ore discovery is also good. The known reserves alone when produced and delivered to a mill have a market value on the order of \$45 to 50 million. An appraisal of the AEC controlled lands was made to determine the potential adverse environmental effects of the proposed leasing program. A multi-disciplined team was used which included experts in various fields from the Bureau of Land Management of the Department of Interior, an AEC contractor,

Lucius Pitkin, Inc., the Forest Service, Department of Agriculture, and the AEC. Factors considered included: present multiple use practices including range and livestock, wildlife, and recreation; prehistoric and historic values; soil and watershed, forestry, roads, other mineral values; mining requirements for water and other resources; water, air, and other pollution possibilities from mining and milling operations; proposed land use projects; and potential effects on local communities, water, power supplies, schools and other facilities. As a result of this review, draft stipulations were developed, and will be included in lease agreements to minimize potential adverse effects of the conduct of exploration and mining activities on the AEC controlled lands. The alternatives to this plan are to not lease, and return lands to the public domain or to not lease, and maintain lands in withdrawn status. Considering the substantial anticipated benefits versus the rather minimal adverse environmental effects and dollar costs, and in view of the range of reasonable alternatives, it was determined that the proposed leasing program should be undertaken. (L.C.L.)

Thermal Effluents

Refer also to abstracts: 47397, 47401, 47404, 47413, 47424, 47435, 47437, 47446, 47448, 47465, 47471, 47473, 47479, 47486, 47487, 47489, 47497, 47498, 47502, 47510, 47516, 47532, 47534, 47535, 47539, 47543, 47545, and 47564.

45573 (AE-448) LOW-GRADE HEAT FROM THERMAL ELECTRICITY PRODUCTION: QUANTITY, WORTH, AND POSSIBLE UTILISATION IN SWEDEN. Christensen, Joergen (Aktiebolaget Atomenergi, Studsvik (Sweden)). 8 Oct 1971. 101p. Dep. NTIS (U. S. Sales Only).

In a thermal cycle working process, entropy is degraded from one temperature to a lower temperature, T_2 . The loss of heat at T_2 is an inescapable part of the process; if the entropy flow is $\dot{S}$, the flow of this degradation heat is $\dot{S} \times T_2$. The ratio of heat flow to mechanical power or, in an electric generating plant, say, the cooling/electricity power ratio may be influenced in several ways, but radical changes can hardly be expected, not even with fusion power. Swedish degradation heat flow by 2000 A.D. is estimated to be 70 GW $\pm$ 14 GW. The pricing of heat from a combined plant is and must be arbitrary. A pricing method based on energy coupled to the electricity market is chosen here. Low-grade heat at temperatures above that of the cooling outlet from electricity-only plants is not cost free. Among the ways of utilizing low-grade heat considered are distillation of sewage, heating of soil, heating of greenhouses, heating of water for aquaculture, and chemical processes. (auth)

45574 (CONF-720619-1) MEASUREMENT OF LIQUID DROPLET EMISSIONS FROM COOLING TOWERS AND PROCESS STACKS. Fish, Birney R.; Duncan, Joseph R. (Tennessee Univ., Knoxville). 1971. 21p. Dep. NTIS.

From sixty-fifth meeting of the air-pollution control association; Miami Beach, FL. (18 Jun 1972).

Many air pollution control regulations specifically exclude from their definition of an air contaminant that which is termed "uncombined" water droplets. However, there is an increasing awareness that water drops from some sources contain various salts, fungicides, corrosion inhibitors, and other dissolved or entrained material. Measurement of the emission rate of water droplets requires special techniques similar to those that have been used in studying the microphysics of clouds and fog. A new method of water droplet size analysis was developed, combining the techniques of the impaction stain methods with standard filter paper air sampling. The air filter is sensitized by soaking it in an aqueous solution of potassium ferricyanide, drying it, and then dusting it with finely ground ferrous ammonium sulfate. These two compounds dissolve in a captured water droplet and react to form an insoluble blue dye which precipitates near the evaporating periphery of the droplet leaving a ring-like blue stain. Droplet diameter can be related to stain diameter by calibrating the sensitized paper using known size droplets. Field tests confirm the validity of the air filter-strain method in measuring the emission of droplets from a water cooling tower. (auth)

45575 PREDICTION OF HEATED EFFLUENT DISPERSION IN A STREAM USING SIMILITUDE TECHNIQUES. DeLong, Max Myron. Ames, Iowa; Iowa State Univ. (1972). 121p. University Microfilms Order No. 72-19,981. Thesis.

An experimental study is done to develop prediction equations for heated effluent dispersion using open channel models designed by similitude techniques. In some models the flow depth is intentionally distorted compared to the horizontal dimension. The

principles of engineering similitude are used to analyze open channel flow, and sixteen dimensionless ratios, or Pi terms, are developed to describe the phenomenon of heated effluent dispersion. Distortion factors are defined to characterize the intentional and unavoidable deviations of the model design conditions from those of a true model. Seven open channel systems were paired in different combinations to achieve models with different length scales and different degrees of depth distortion. Temperature data were obtained by using thermistors. Prediction factors were developed from the experimental data. The relation of prediction factor with depth and slope distortion factors and length scale was determined. (Diss. Abstr. Int., B)

45576 LIVESTOCK SHELTERS AS HORIZONTAL COOLING TOWERS. Hirst, Eric (Oak Ridge National Lab., Tenn.). Agr. Eng.; 53: No. 5, 18-19 (May 1972).

A proposal is presented for using power plant waste heat for providing optimal thermal conditions for raising small farm animals such as chickens and pigs. The quantity of heat rejected from U. S. power plants, the heat requirements for raising broilers and pigs in various geographical locations, and shelter and ventilation requirements, and the advantages and problems associated with this use of power plant waste heat are discussed. (L.C.L.)

45577 PROCEEDINGS OF THE FOURTEENTH CONFERENCE ON GREAT LAKES RESEARCH, TORONTO, ONTARIO, APRIL 19-21, 1971. Ann Arbor, Mich.; International Association for Great Lakes Research (1971). 844p. (CONF-710431).

Seventy-five papers on Great Lakes research were presented at the conference. A separate abstract was prepared for each of 7 papers. One paper was previously abstracted for NSA. For an abstract of this see CONF-710431 in the Report Number Index. Sixty-seven papers were not within the scope of NSA. (L.C.L.)

For abstracts of individual papers see: 45535, 45536, and 45578-45582.

45578 INFLUENCES OF THERMAL EFFLUENTS UPON AQUATIC PRODUCTION IN LAKE ONTARIO. Fenlon, M. W.; McNaught, D. C.; Schroder, G. D. (State Univ. of New York, Albany). pp 21-6 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

Additions of heat to the Lake Ontario ecosystem at Nine Mile Point have increased the standing crop of *Bosmina* 25.0 times and *Daphnia retrocurva* 1.2 times in the overall study area. Adjacent to the outfall these same populations increased 123.8 and 2.4 times respectively. At the same time primary production was not significantly affected. It is suggested that turning-over these large populations of herbivores by fishing their predators may reduce the likelihood of algal blooms as thermal additions to the Great Lakes are increased in the 1980's. (auth)

45579 THERMAL PLUME DATA ACQUISITION, DOCUMENTATION, AND INITIAL ANALYSIS. Abu-Shumays, I. K.; Phillips, D. L.; Prastine, S. M. (Argonne National Lab., Ill.). pp 495-506 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

A brief description is presented of the temperature data acquisition and documentation procedure adopted by interdisciplinary group at Argonne National Laboratory during the summer of 1970. The 12 August, 1970 thermal plume at the Commonwealth Edison Waukegan Station is used to illustrate the procedure. The aim was to employ relatively simple and economic methods of acquiring and analyzing the data needed to explore thermal plume structure. The accuracy of the procedure is analyzed and an initial computer program for documenting the data and displaying plume patterns is described. The statistical nature of the thermal plume is stressed, and what might be called a statistical plume and an average or representative plume is discussed. (auth)

45580 NATURAL AND UNNATURAL WATER TEMPERATURES IN ZION-WAUKEGAN, ILLINOIS AREA OF SOUTHWEST LAKE MICHIGAN. Beer, Lawrence P. (Industrial Bio-Test Labs., Inc., Northbrook, Ill.). pp 507-21 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

A well-defined thermal bar observed offshore from the Zion-Waukegan, Illinois area of southwest Lake Michigan on Apr. 23, 1970 moved offshore a distance of at least 6,080 m by Apr. 30 a velocity of movement of at least 0.01 m/sec. A poorly defined thermal bar was observed on Dec. 18, 1970. Ambient water tem-

peratures during the period of summer stratification varied considerably between two locations 6 to 8 mi apart. During the fall stratification was reduced and winter water temperatures were nearly isothermal in the near-shore waters. Thermal plumes from the Waukegan Generating Station (Waukegan, Illinois) varied in surface area from approximately 50 acres to approximately 174 acres during the five days of observation (during Apr. 23, 1970 to Feb. 16, 1971). Plumes were directed generally downwind, except in the case of onshore winds. Then, the plume assumed a pear shape east of the outfall and did not move upshore or downshore. Water temperatures at the lake bottom were affected by the plume from as little as 153 m to as great as 1,980 m from the outfall. The distance affected appeared to be a function of the depth of the lake in the downwind direction. The thermal plumes tended to rise to the surface and exchange heat with the atmosphere except when ambient winter waters were cold enough ($<4^{\circ}\text{C}$) to allow the warmer, more dense plume water to sink to the bottom. In this case, however, mixing of plume water with ambient colder water apparently was rapid. (auth)

45581 THERMAL PLUME STUDY AT DOUGLAS POINT, LAKE HURON. Csanady, G. T.; Crawford, W. R.; Pade, B. (Univ. of Waterloo, Ont.), pp 522-34 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971). From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

The temperature structure was explored of the cooling water effluent plume of the Douglas Point nuclear generating station. Two thermistor probes were carried on a short sampling boom on a catamaran and the plume was criss-crossed many times to provide fairly detailed cross-sections of thermal structure. Current speeds were also measured, so that the heat flux at successive sections of the plume could be estimated, although only with moderate accuracy. The results may be presented in terms of a four component system: (1) the power station; (2) the "identifiable plume" defined as that body of water lying within the 0.5°C excess temperature isotherm; (3) the atmosphere; and (4) the remainder of Lake Huron. All the injected heat flows from (1) to (2), a fraction of it flows from (2) to (3), the remainder from (2) to (4) in the first instance, although it is ultimately transferred to (3), but over a rather larger area and at rather lower excess temperatures than the flux going directly from (2) to (3). The magnitude of this "direct" heat flux to the atmosphere (as a fraction of the total rejected heat) varied somewhat, and the estimate was also subject to considerable random scatter, but it seemed to be approximately $\frac{1}{3}$. The "identifiable plume" was shallow and broad, its typical depth being 1.2 to 1.5 m, width 600 m and length 2300 m. The shallowness was somewhat remarkable: the densimetric Froude number based on this depth and current speed appears to be around unity. (auth)

45582 EXPERIMENTAL AND NUMERICAL INVESTIGATIONS OF RECTANGULAR JET. Paul, John; Prahl, Joseph (Case Western Reserve Univ., Cleveland), pp 607-17 of Proceedings of the Fourteenth Conference on Great Lakes Research. Ann Arbor, Mich.; International Association for Great Lakes Research (1971).

From fourteenth conference on Great Lakes research; Toronto, Ontario, Canada (19 Apr 1971). See CONF-710431.

Experimental and numerical results are presented for a rectangular constant temperature jet, horizontally entering a basin

of constant depth equal to the vertical extent of the jet. The experimental investigation is carried out on a 6 ft $\times$ 20 ft (1.83 m $\times$ 6.10 m) water table. Velocity profiles were measured by a conventional hot film constant temperature anemometer. The numerical investigation is based upon a vertical integration of the three-dimensional, time-dependent Navier-Stokes equations for an incompressible fluid. The final equations solved are essentially two-dimensional volume flux equations with parameters in the inertial and bottom friction terms. Results, both experimental and numerical, indicate that the effect of bottom friction is to introduce a diffusive behavior to the jet. (auth)

45583 THERMAL POLLUTION: A POTENTIAL THREAT TO OUR AQUATIC ENVIRONMENT. John, James E. A. (Univ. of Maryland, College Park). Environ. Aff.; 1: No. 2, 287-98 (Jun 1971).

In the next decade and beyond, with the very large amount of thermal energy to be discharged into our rivers, lakes and streams, there is a very real danger of degrading our aquatic environment. Up to the present time, there have been some reported cases of fish kills due to heated water effluents, yet not a large number. Rather, what has been reported are subtle effects that have been observed by scientists. It is clear that, if the number of power plants on lakes and rivers is allowed to increase without control, these subtle effects can only multiply and increase in magnitude, producing irreversible effects on our environment. One of the key factors in the control of thermal pollution is the selection of a suitable site for a power plant. In the past, site selection has been based mainly on economic considerations. In the future site selection must also take into account the effect of the power plant on the ecology of the surrounding area. The engineer will have to work hand in hand with the biologist on both site selection and operation of the plant. In many locations, it may be necessary to incur the added cost of cooling towers or ponds to protect rivers, lakes and streams. It is clear that, in the long run, the ultimate goal, and one that seems within the reach of today's technology, should be to utilize the waste heat from a power plant for the benefit, rather than to the detriment, of man and his environment. (auth)

45584 ATMOSPHERIC EFFECTS OF THERMAL DISCHARGES INTO A LARGE LAKE. Carson, James E. (Argonne National Lab., Ill.). J. Air. Pollut. Contr. Ass.; 22: No. 7, 523-8 (Jul 1972).

Concern for the biological and ecological effects of heated water has resulted in legal actions that will prevent power companies from dumping the waste heat from the majority of their new generating units into rivers and lakes. Many nuclear- and fossil-fueled plants now under construction, and even some now online, are being required to change from once-through cooling systems to other methods, such as wet cooling towers, cooling ponds, and spray canals, despite higher costs and lower thermal efficiencies. Yet, these alternate cooling procedures are not without their own environmental problems. The primary weather change due to once-through cooling on a large water body is a small local increase in foginess at the plant outfall. But the relative probability of significant local meteorological effects is much higher with alternate cooling procedures, since these reduce the area of heat and moisture transfer. It is therefore concluded that, from a meteorological point of view, the least undesirable way to dispose of waste heat is by using once-through cooling on large water bodies. (auth)

INSTRUMENTATION

Radiation Detection Instruments

Refer also to abstracts: 46083 and 47203.

General Detectors and Monitors

Refer also to abstracts: 45261 and 45673.

45585 (AERE-HP/M-74) MONITOR FOR ESTIMATING TRITIUM IN URINE. Booker, D. V.; Megaw, W. J.; Pittendrigh, L. W. D. (Atomic Energy Research Establishment, Harwell (England)). 29 Apr 1954. 9p. Dep. NTIS (U. S. Sales Only).

The use of an ionization chamber to determine levels of tritium present in urine is described. Acetylene from the action of urine on calcium carbide is passed through the chamber and from the ionization current reading, levels of tritium from 0.1 to 12 maximum permissible levels can be determined. (auth)

45586 (ARH-2432) RELATIVE EFFICIENCY AS A FUNCTION OF TIME FOR ATLANTIC RICHFIELD HANFORD COMPANY COAXIAL Ge(Li) DETECTORS. Zimmer, William H. (Atlantic Richfield Hanford Co., Richland, Wash.). May 1972. Contract AT(45-1)-2130. 35p. Dep. NTIS.

The performance and recovery of four detector systems are reported as typical of Ge(Li) detector performance at Hanford. (auth)

45587 (BNL-50332) OPERATING CHARACTERISTICS OF NEUTRON WELL COINCIDENCE COUNTERS. Sher, R. (Brookhaven National Lab., Upton, N. Y.). 1 Jan 1972. 47p. Dep. NTIS.

The neutron well coincidence meter consists of a bank of thermal neutron detectors, either BF₃ or ³He counting tubes, embedded in a moderating medium (usually polyethylene) for detecting spontaneous fission neutrons from Pu sources in the presence of a large random background. The detector system has principally been used to determine ²⁴⁰Pu contents of Pu-bearing samples such as feed material and fabrication plant output. Measured properties including efficiencies, linearity, geometric effects, matrix effects, and self-multiplication effects are discussed. A detailed description of the electronics is given, and dead-time corrections are discussed. Uncertainty considerations including statistical errors and calibration errors and the dependence of uncertainty on count time and source strength are detailed. Representative data obtained under field conditions with the neutron well coincidence counters are given. (W.H.K.)

45588 (COO-2114-2) EFFICIENCY STUDIES OF PILOT Y, PILOT F, AND NE 110 SCINTILLATOR. Messner, Robert; Nauenberg, Uriel (Colorado Univ., Boulder (USA)). Duane Physical Lab.). 1972. 6p. Dep. NTIS.

The scintillators tested had dimensions of 24" x 2" x 1/8". The efficiency was measured as a function of position along the long dimension. It was found that only Pilot Y plastic scintillator had a constant counting efficiency over the range of 24". (3 figures) (R.W.R.)

45589 (JEN-247) NEW APPROACH TO β - γ COINCIDENCE COUNTING. ADVANCE REPORT ON THE "SAMAR" ELECTRONIC SYSTEM. De Carlos, J. E.; Granados, C. E. (Junta de Energia Nuclear, Madrid (Spain)). 1972. 47p. (In Spanish). Dep. NTIS (U. S. Sales Only).

In $4\pi\beta$ - γ coincidence measurements, precision on the evaluation of coincidence counting losses is made difficult because of complex overlapping effects between the β - and γ -side dead times due to precursive counted events. In this context the SAMAR electronic system is aimed to give a precise way of automatic counting and reduce the need for calculated corrections. The configuration and basic features of the SAMAR are described. The SAMAR has been conceived in such a manner that both beta and gamma chains are sharing a common and nonextending dead time which is simultaneously applied to both channels. The shared dead time is made to be the only one inserted throughout the chains. Overlapping effects vanish and the three counting channels have identical transmission ratios. A new dead time circuit based on fast linear gates as blocking elements has been developed. Application of the two-oscillator Mueller's method evidences a fully non-extending character. Automatism is implemented by using a live timer corrective channel controlling the counting scalars. (auth)

45590 (ORNL-tr-2532) COMPARISON OF THE BACKGROUND SPECTRA IN EIGHT GERMAN WHOLE-BODY COUNTERS

AND A MODEL STRUCTURE WITH LAYERED SHIELD. Futtermenger, W. (Technische Universitaet, Hanover (West Germany)). Translated for Oak Ridge National Lab., Tenn., from report ABS-TUH-1065. 14p. Dep. NTIS.

The sources of background radiation in whole-body counters are discussed including radiation from wall material, cosmic radiation, radon radiation from the interior of the whole-body counter, and contamination and radiation from built-in instruments. The background spectra in eight German whole-body counters and a model structure were measured and compared. The various shielding designs used in the facilities are described and the construction costs listed and compared. (W.H.K.)

45591 MEASUREMENTS OF THE ABSOLUTE RADIATIVE EFFICIENCY OF CATHODOLUMINESCENT PHOSPHORS. Meyer, V. D. (Bayside Research Center, N. Y.). J. Electrochem. Soc.; 119: No. 7, 920-6(Jul 1972).

The absolute efficiency of cathodoluminescent phosphors was measured as a function of voltage and current density. The results are in good agreement with the highest values previously reported. Changes of slope in the luminance vs voltage curves indicate that changes in the voltage dependence may occur at x-ray excitation potentials; these observations are related to results reported by others for x-ray excitation of scintillation crystals. The indications are that it may not be possible to assume that the intrinsic efficiency is independent of the kinetic energy of the incident electron beam unless the energy exceeds the K x-ray levels of the host. (auth)

45592 APPARATUS FOR CONTINUOUS MEASUREMENT OF ACTIVE UPTAKE OF RADIOACTIVE SUBSTANCES. Loos, Eckhard (Univ., Regensburg, Ger.). Anal. Biochem.; 47: No. 1, 90-101(May 1972).

An apparatus is described allowing the registration of uptake kinetics of radioactive compounds by cells. Cells are held in a thin layer close to a scintillation crystal so that uptake of a labeled compound is seen as an increase in measured radioactivity. As a demonstration of the method, accumulation and countertransport kinetics are shown with 3-O-[¹⁴C]-methyl-D-glucose for *Chlorella vulgaris* and for conidia of *Neurospora crassa*; for *Chlorella* a K_m of 6×10^{-4} M is found for the uptake. Parallel experiments with the membrane filter technique show good agreement and can be used to calibrate the apparatus. (6 figures, 1 table) (auth)

45593 SUBTRACTION AND POSITRON SCANNING WITH THE DUAL HEADED WHOLE BODY SCANNER. Saegusa, Kenji; Kakehi, Hirotake; Arima, Akira (Chiba Univ., Japan). Radioisotopes (Tokyo); 21: No. 6, 342-7(Jun 1972). (In Japanese).

The 5 in. dual headed whole body scanner, which is used in routine tests, has three operation modes of addition, subtraction, and positron scanning besides the ordinary one. The diagnostic usefulness of the subtraction and positron modes was studied. The word "subtraction scanning" in this study has different meaning from the usual subtraction technique. In this subtraction method, the image on the scintigram can be made by counting differences of output signals from upper and lower detectors. Therefore, two scintigrams, positive and negative, can be obtained simultaneously. Collimators of good resolution are not always necessary for this technique. Positron scanning is one of the established scanning techniques. However, unless the sensitivity of crystal is high enough, it is usually difficult to get a good result. In this study two detectors were used, one with a focusing collimator and the other with no collimator. Both scannings, subtraction and positron, have less counting dots on the scintigrams compared with the ordinary scanning. If larger doses of radioisotopes with shorter half-lives and with no beta particles are used, scintigrams of good contrast and sharp quality can be obtained by these techniques. (auth)

45594 EFFECT OF TEMPERATURE ON VARIOUS TYPES OF THE IONIZATION CHAMBER. Okabe, Shigeru; Tabata, Tatsuo; Tsumori, Kunihiro. Annu. Rep. Radiat. Center Osaka Prefect.; 12: 45-8(1971).

The dependence of ionization current on temperature was investigated for various types of the chamber by using pulsed X rays from a linear electron accelerator and γ rays from a ⁶⁰Co source. In the glass-wall and the metal-wall chambers equipped with parallel-plate electrodes of Au and filled with various kinds of gases, both average and peak ionization currents decreased rather remarkably with increasing temperature. In the cylindrical chamber having a cylindrical Au-electrode, the currents showed

a smaller decrease. Possible causes of the temperature effect are briefly discussed. (auth)

45595 TELEVISION TYPE NUCLEAR RADIATION CAMERA SYSTEM. Asars, Juris A.; Schneeberger, Robert J. (to CGR Medical Corp.). U. S. Patent 3,668,396. 6 Jun 1972. Filed 10 Nov 1969.

A television type camera system in which TV type scanning is used for readout of a spatially distributed nuclear radiation image is described. The scanning system provides means of time integration of the video pulses derived from a single radiation event and provides discrimination of the integrated pulses generated by a single event in the spatially distributed image. In one embodiment of the invention, conventional TV scanning is utilized and circuitry is associated with the system to permit time integration of the multiple video pulses derived from a single event for subsequent discrimination. A second embodiment utilizes a non-conventional scanning system to permit the electron scanning beam to stop and integrate a single event and subsequent discrimination of the signal. (auth)

45596 SCINTILLATION CAMERA HAVING IMPROVED PERIPHERAL RESPONSE. Walker, William G. (to Nuclear-Chicago Corp.). U. S. Patent 3,668,395. 6 Jun 1972. Filed 17 Dec 1969.

A substantial increase in the diameter of the useable light image generated by a scintillation camera equipped with a multiple collimator is achieved by using a collimator that does not include holes which allow radiation to strike the periphery of the scintillator. (auth)

45597 NUCLEAR RADIATION DETECTION DEVICE UTILIZING DIAMOND DETECTOR WITH INJECTING AND BLOCKING CONTACTS. Kozlov, Stanislav Fedorovich. U. S. Patent 3,668,400. 6 Jun 1972. Priority date 11 Sep 1968, Russia.

A device for detecting nuclear radiations, comprising a nuclear radiation detector based on a diamond crystal plate, having a blocking and an injecting contact, a detector power supply associated with the injecting contact, and an amplifier with recording means associated with the blocking contact, is described. (auth)

45598 NEW READING SYSTEM FOR AN IONOGRAPHIC DETECTOR. Chapuis, A. M. (CEN, Fontenay-aux-Roses, France); Soudain, G.; Gerard, N. Radioprotection; 7: No. 1, 13-20(1972). (In French).

The track density may be obtained from an ionographic detector from measurements of the light transmitted by the detector, which incorporates a thin film of colored cellulose nitrate in which holes correspond to the particle tracks. Results obtained with a U source and radioactive aerosols are presented. (France)

45599 SCATTERED NEUTRONS AND THEIR EFFECTS ON THE CALIBRATION OF BONNER SPHERE COUNTERS. Heinzelmann, M.; Rohloff, F.; Schueren, H. (Kernforschungsanlage, Juelich, Ger.). Atomkernenergie; 19: No. 4, 312-16(Jul 1972). (In German).

The efficiency of Bonner sphere counters has been measured by use of an Am-Be-neutron source. Special attention has been given to the variation of measured counting rate due to scattered neutrons at different scattering levels. Measurements have proven that the hypothesis of a quadratical dependence on distance and a scattering rate independent of distance gives as good results as an additional evaluation of an effective center of the Bonner sphere. Measurements at a very large distance from the source are prone to give too high values of efficiency, as the scattering rate is no longer constant for distances greater than one meter. (auth)

45600 FRONTIERS IN RADIATION DETECTION. Res./Develop.; 23: No. 5, 24-6(May 1972).

The development of cadmium telluride, gallium arsenide, and mercuric iodide semiconductor detectors is briefly discussed. (W.H.K.)

45601 QUANTITATIVE EVALUATION OF RADIOISOTOPE DISTRIBUTION IN VIVO BY ISOSENSITIVE SCANNER PLUS 4096-WORD MULTI-CHANNEL ANALYZER COUPLING. Hisada, Kin-ichi; Kojima, Kazuhiko; Matsudaira, Masamichi; Hiramatsu, Hiroshi (Kanazawa Univ., Japan). Radioisotopes (Tokyo); 21: No. 6, 348-52(Jun 1972).

In an isosensitive radioisotope scanning system the delineation of the radioisotope distribution is completely independent of depth. The coupling of a 4,096-word multi-channel analyzer to the isosensitive scanner makes it possible to evaluate radioisotope distribution in vivo quantitatively and provides various new diagnostic tools; for example, spleen/liver ratio, pancreas/liver ratio, profile histogram of lung perfusion scan, and tumor net counts in brain scan. (auth)

45602 USE OF A CONVERGENT HOLE COLLIMATOR FOR TRANSMISSION SCINTIPHOTO AND FOR THE COMBINED X-PHOTO AND SCINTILLATION IMAGE. Saito, Hiroshi; Tamiya, Tadashi (Nagoya Univ., Japan). Radioisotopes (Tokyo); 21: No. 6, 338-41(Jun 1972). (In Japanese).

Transmission scintiphotos were taken using a gamma-ray point source and a convergent 1261 hole focusing collimator. In addition, X-photo and scintillation images taken by keeping the X-ray tube-film distance equal to focus-scintillator distance had the same rate of magnification and both the images were superimposable. If an attachment is used to reproduce the same geometry in recording scintillation image and X-photo, these images can be double exposed on the same X-ray film. Thus the localization of the scintillation image can be easily determined especially in combination with X-photos. The present method would be useful not only for diagnosis, but also for therapy. (auth)

45603 RADIATION CONTROL AND MONITORING SYSTEM. Bean, Laurence R. (Univ. of Kentucky, Lexington). J. Amer. Dent. Ass.; 85: No. 1, 90-3(Jul 1972).

A device that can be installed in existing dental X-ray units to control and monitor radiation exposure to the patient is described. The radiation control unit emits a preset value of radiation and terminates the exposure. The attached printing device prints information set on the machine, such as milliroentgen values and kilovoltages, the tooth number, and data. The device utilizes an ionization chamber. (auth)

45604 PROLONGED ETCHING FACTOR IN SOLID STATE TRACK DETECTION AND ITS APPLICATIONS. Khan, H. A.; Durrani, S. A. (Univ. of Birmingham, Eng.). Radiat. Eff.; 13: No. 3-4, 257-66(Apr 1972).

It was observed that the number density of etch pits resulting from spontaneous or induced fission within a solid state track detector (SSTD) is a function of the duration of etching, other conditions being constant. As the etching proceeds, the net number of etch pits observed per unit area continues to increase. This necessitates a modification of the conventional equations governing the observed etch-pit densities in SSTD. A prolonged etching factor $f(t)$ is defined, and its value measured as a function of time under various specific etching conditions, in a number of materials of interest, including "reference glass". Furthermore, the theoretical relationship of this factor with the general velocity of etching in a given material, and with the average range R of fission fragments in it, is utilized in deriving self-consistent and very satisfactory values of R in the above-mentioned materials. (auth)

45605 DIRECTIONAL PROPERTIES OF CYLINDRICAL SCINTILLATORS SCREENED BY A SLEEVE-TYPE SCREENING. Petr, Ivo. Jad. Energ.; 18: No. 7, 233-6(Jul 1972). (In Czech).

The suitability of using a sleeve-type lead screening of maximum thickness 1 cm for obtaining directional detection properties for gamma-rays is discussed. The mathematical expressions for the detection cross section are derived and the value of the effective linear absorption coefficient of the shielding was determined from the comparison of theoretical and experimental results. Finally, the upper energy limit where the screening is suitable is determined. (auth)

45606 SPECTRAL AND SCINTILLATION PROPERTIES OF THE PYRIMIDINE DERIVATIVES. Visotskaite, S. B.; Zhuravskene, E. A.; Paeda, R. I.; Yasinskas, L. L.; Kundra, A. A. (Vilnyus State Univ., USSR). Liet. Fiz. Rinkiny; 11: No. 5, 871-80(1971). (In Russian).

The absorption and fluorescence spectra of 29 derivatives of pyrimidine were investigated. The scintillation properties of the pyrimidine derivatives were also examined. The relative scintillation efficiency of the above compounds used as primary soluble substances is 13 to 30% in comparison with PPO and p-terphenyl in dioxane. As secondary soluble substances some of them increase the counting efficiency in PPO and p-terphenyl scintillators in dioxane. (3 figures, 2 tables) (auth)

45607 NEW TWO-CHANNEL SCANNER: THE SCINTICART NUMERIC T. II. EXPERIENCES IN PRACTICE. Krasznai, I. (Belgyogyszati Klinika, Budapest); Foeldes, J.; Billing, A.; Ambro, P. Izotoptechnika; 15: No. 1, 37-45(Apr 1972). (In Hungarian).

The results of experiments using phantoms and their application to clinical situations are presented. (8 figures, 1 table) (auth)

45608 PREAMPLIFIER FOR CHERENKOV SPECTROMETER. Khurshudyan, L. S.; Muradyan, E. G. (Inst. of Physics, Erevan). Izv. Akad. Nauk Arm. SSR, Ser. Fiz.; 6: No. 3, 223-5(1971). (In Russian).

Some peculiarities in the linear transmission of nanosecond pulses from radiation detectors by emitter follower into a cable with small wave impedance are considered, the feeding of the preamplifier being by the same coaxial cable as that transmitting the signal. A practically useful circuit for a preamplifier providing a transmission (with nonlinearity not worse than 1%) of pulses with a magnitude from 0.008 to 8 volts and duration from 10 ns to 10 ps into a cable with an impedance of 50 ohms is described. (1 figure) (auth)

45609 DEVICE FOR CONTROLLING SLOW DRIFT IN IONIZATION CHAMBERS. Dalmazzone, Jean (to Commissariat a l'Energie Atomique). French Patent 2,083,752. 22 Nov 1971. Filed 26 Mar 1970. (In French).

A device for controlling slow drift in a dc ionization chamber is presented. A reference chamber containing a fixed weak radiation source of long half-life is placed in parallel with the measurement ionization chamber. A switching system permits the connection (separate or in parallel) of the reference chamber and the measurement chamber to an associated electronic unit. According to the construction the monitor may have its own atmosphere, or an atmosphere joined to the common atmosphere; it contains a ^{14}C or ^{241}Am source; the switching system may be a highly insulated single or double switch. (France)

45610 USE OF SOLID TRACK DETECTORS IN REACTOR EXPERIMENTS. Kuznetsov, V. A.; Mogil'ner, A. I.; Koroleva, V. P.; Samovarov, V. S.; Chernov, L. A. At. Energ. (USSR); 32: No. 6, 481-3 (Jun 1972). (In Russian).

Methods of utilizing nitrocellulose for recording α particles (from ^{10}B and ^6Li) and mica for recording ^{235}U fission fragments were investigated. The nitrocellulose was in the form of motion picture film or x-ray film, for which the following etching conditions are suggested: 30% NaOH, 60°C, for 15 min. Etching of the mica was conducted with 40% HF, 60°C, for 30 min. (7 figures) (K.S.W.)

45611 APPLICATION OF TRACK STRUCTURE THEORY TO DOSIMETRY AND RADIOBIOLOGY. Homayoonfar, Mehdi. Lincoln, Nebr.; Univ. of Nebraska (1971). 258p. University Microfilms Order No. 72-15,988.

Thesis.

By the application of δ -ray theory, a model for cell survival after energetic ion bombardment is proposed. The survival curves for some cells are both sigmoid and exponential types, and are accounted for on the conceptual basis of ion-kill and gamma-kill modes and multitarget single-hit survival hypothesis. Tracks formed in nuclear emulsion by energetic heavy ions are made up of grains sensitized by the passage of a single ion in a model called ion-kill. The response of cells is not the same as emulsion grains, because cells also can be inactivated by a gamma-kill mode, involving the cumulative effects of overlapping δ rays from several ions. Survival curves are calculated and compared with experimental survival data for T-1 human kidney, HeLa, Chinese hamster, haploid yeast, and bacterial spores. The proposed model is extended to those cells whose experimental survival curves have big shoulders to x and gamma rays and heavy ions. The cellular model is further extended to the problem of radiation therapy and mixed radiation environments like neutrons and stopped pions. The predicted survival curves for this mixed radiation are compared with existing experimental data for T-1 human kidney as well as HeLa cells irradiated with 14-MeV neutrons; agreement is good. The δ ray theory was extended to analyze the response of various detectors like the LiF thermoluminescent dosimeter, the Fricke dosimeter, the silver-activated phosphate glass dosimeter, the creation of free radicals in solid biological substances, solid and liquid organic scintillators and other systems. The results show that all are well described by the application of the δ -ray theory. Difficulties arising from the use of LET as a major and basic tool to describe radiation quality, cross section, radiosensitivity, OER, RBE, etc., are discussed. (TSS) (Diss. Abstr. Int., B)

45612 TRACK EFFECTS IN PHYSICAL, CHEMICAL, AND BIOLOGICAL SYSTEMS. Sharma, Subhash Chander. Lincoln, Nebr.; Univ. of Nebraska (1971). 231p. University Microfilms Order No. 72-16,014.

Thesis.

The universal applicability of the delta-ray theory of the track structure associated with heavy ions to several physical, chemical, and biological systems is shown. The theory is extended to various radiation detectors like the silver-activated phosphate glass dosimeter, the LiF thermoluminescent dosimeter, the creation of free radicals in solid biological substances (esr experiments), solid and liquid organic scintillators, and the chemical systems like the ferrous sulphate (Fricke) dosimeter. The theoretical yield curves are in good agreement with experi-

mental data. No saturation cross section is observed for these detectors. The nonlinearity and the multiple valuedness of the response curves for various radiation detectors are consequences of the characteristic dose and the size of the sensitive element. Variables describing the bombarding ion and those describing the medium are not separable. A cell model is proposed which accounts for the survival curves observed when elementary biological systems are bombarded with heavy ions on the basis of a 1-hit multitarget survival hypothesis. The dosage of ionization energy deposited by the delta rays from a single ion may itself inactivate a cell along the ion's path. This is most likely to occur at high specific energy loss. Since cells remember, cells not inactivated by a single ion may be inactivated by cooperative effects from several ions. Survival curves are calculated and compared with experimental data for spores, yeast, and various mammalian cells. The model proposed for cell radiation response is further applied to the prediction of cell survival curves for 14-MeV neutrons, stopped negative pions, and any radiation which gives rise to secondary charged particles whose energy spectrum is known. The models proposed were also investigated for predicting the radiation hazards and detection of heavy ions in cosmic rays. (TSS) (Diss. Abstr. Int., B)

45613 SEQUENTIAL SAMPLER. Teel, Dale M.; Putnam, John H. Jr. (to United States Atomic Energy Commission). U. S. Patent 3,657,920. 25 Apr 1972. Filed 6 May 1970.

A sampler for collecting a series of samples of aerosols from the atmosphere is described. The sampler utilizes a removable head having several sampling units, which are sequentially actuated to obtain samples at regular intervals over long periods of time. The head carrying the sampling units is readily replaceable, and the whole device is designed for remote operation. (2 figures) (auth)

45614 MINIATURE LOW-BACKGROUND COUNTERS FOR MEASUREMENT OF ^{37}Ar AND ^{39}Ar . Fisenko, A. V.; Kolesnikov, E. M. (Inst. of Pure Chemistry and Analytic Chemistry, Moscow). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1654-5 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 62-3 (1971).

The construction of miniature counters having an internal filler which is used to measure ^{37}Ar and ^{39}Ar , while operating in either the Geiger or proportional modes, is described. Information is presented on the magnitude of the background as a function of cathode material, cathode thickness, or the voltage on the guard electrodes. (auth)

45615 RECORDING OF ALPHA PARTICLES BY SOLID TRACK DETECTORS BASED ON CELLULOSE. Bondarev, V. N.; Korzin, E. G.; Parfanovich, D. M. Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1648-50 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 56-9 (1971).

A study was made of solid track detectors based on cellulose. For films made of nitro- and acetyl-cellulose the energy ranges of α particles which can be recorded with 100% efficiency are established, and the dependences of the track dimensions of α -particle energy are obtained. The results of measurements of the space distribution of the density of thermal-neutron capture in a boron absorptive rod having a small diameter measured by means of a nitro-cellulose detector are presented. (auth)

45616 TIME MEASUREMENTS WITH A GERMANIUM DETECTOR. Akimov, Yu. K.; Andert, K.; Kalinin, A. I.; Churin, I. N.; Shuravin, V. N. (Joint Inst. of Nuclear Research, Dubna, USSR). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1641-4 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 51-4 (1971).

A low-noise preamplifier with a rise time of 5 nsec and a fast shaper with compensation of the dependence of the time resolution on amplitude scatter and on changes in the leading edge of the pulses was developed for time measurements with a germanium detector. In recording γ quanta from ^{60}Co by a germanium detector having a volume of 3 cm^3 connected in coincidence with a scintillation counter, a time resolution of 2.2 nsec was obtained in the energy range from 0.07 to 1.33 MeV; the width of the coincidence curve at the 0.1 curve-height level was 12.5 nsec. In a narrow energy range a time resolution of 0.9 nsec was obtained. (auth)

Radiation Dosimeters

Refer also to abstracts: 45672 and 45808.

45617 (FEI-296) STUDY OF THE CHARACTERISTICS OF EMERGENCY TRACK DETECTORS. Obaturov, G. M.; Chumbarov, Yu. K. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoy Energii SSSR, Obninsk. Fiziko-Energeticheskii Institut). 1971. 30p. (In Russian). Dep. NTIS (U. S. Sales Only).

Mica track neutron detectors with layers of ^{237}Np , natural U,

and ^{238}U were investigated; and on the basis of the results a track dosimeter was designed with two layers of natural U and a layer of ^{239}U and ^{237}Np , making it possible to determine neutron doses for both the whole energy spectrum and for various energy intervals with acceptable accuracy. Results are given of comparative studies of track dosimeters, thermoluminescent dosimeters, and tissue-equivalent, low-efficiency Geiger counters for critical assemblies. (tr-auth)

45618 (ICRU-21) RADIATION DOSIMETRY: ELECTRONS WITH INITIAL ENERGIES BETWEEN 1 AND 50 MeV. (International Commission on Radiation Units and Measurements, Washington, D. C.). 15 May 1972. 64p. ICRU Publications, Washington, D. C. \$4.00.

The basic parameters of electron beams are discussed including the determination of electron energy, energy spectra of electron beams, measurement of energy fluence, and energy loss and multiple scattering. General concepts in determining electron absorbed dose are discussed; and calorimetric methods, ionization methods, chemical methods, and solid state methods are considered and compared. Electron beam monitors are discussed under the headings: transmission ionization chambers, secondary emission monitors, induction monitors, partial beam monitoring by ionization chambers, partial beam monitoring by collection of electrons, stability of monitors as a function of machine operating conditions, and choice of monitor. Measurement and control of beam characteristics including determination of the virtual source, beam direction, and beam profile, and measurement of fluctuations of beam direction and beam energy are discussed. Determination of absorbed dose at a reference point, calibration of dosimeters, determination of the absorbed dose in water at any point in the phantom, phantoms, factors influencing the absorbed dose distribution, and beam contamination by bremsstrahlung and neutrons are examined. The energy range (1 to 50 MeV) was chosen to include radiations produced by linear and circular accelerators and which are mainly used for medical, radiochemical, and radiobiological purposes. An extensive list of references is included. (W.H.K.)

45619 IMPORTANCE OF TISSUE-EQUIVALENT DETECTORS USED IN RADIOMETRY. Frolova, A. V.; Kolotilova, V. G.; Krongauz, A. N.; Shagimardanov, N. Sh. (Moscow Scientific-Research Roentgeno-Radiological Inst.). *Med. Radiol.*; 16: No. 12, 62-5 (Dec 1971). (In Russian).

Computer calculations are given for the elemental composition of plastic tissue-equivalent scintillators used for long-wave x radiation. The effective atomic number should be different for different radiation energies, and disregard of this fact results in error in the determination of the absorbed dose and especially its distribution throughout the irradiated volume. (tr-auth)

45620 DOSE-RATE MEASUREMENT BY SEVERAL RADOCON PROBES. II. Kanazawa, Tamotsu; Ohnishi, Tokuhiko; Furuta, Junichiro. *Annu. Rep. Radiat. Center Osaka Prefect.*; 12: 81-3 (1971).

In the dose-rate measurements of gamma-rays, the observed values by a Radocon probe are often found to change to lower values. The experiments were performed to check the causes for this change. The deterioration and the temperature rise of the probe by gamma-irradiation are considered to be the causes. (auth)

45621 PHANTOM-DOSIMETER WITH LIQUID-FILLED IONIZATION CHAMBER. Linchevskaya, G. A.; Luk'yanov, L. A. (Moscow Research X-Ray Radiological Inst.). *Med. Radiol.*; 17: No. 3, 87-9 (Mar 1972). (In Russian).

A measuring instrument consisting of an ionization chamber filled with a tissue-equivalent liquid, a special small-size source of stabilized high tension, and a Y-2 type electrometric amplifier is described. The liquid-filled ionization chamber measures directly the ionizing radiation energy absorbed in the tissue-equivalent medium (i.e. it plays the part of a radmeter). The liquid-filled chamber is encased in a housing made of plexiglas and there is also a set of plates of varying width which are placed on both sides of the liquid-filled detector and can be taken off and interchanged with various tissue- and bone-equivalent plates (phantoms). The main purpose of the proposed as a detector phantom-dosimeter is dosimetry of radiations within a broad range of energies (x-rays, gamma rays, high-energy bremsstrahlung, neutrons, and protons). (auth)

45622 THERMOLUMINESCENT DOSIMETRY SYSTEM FOR DETERMINING QUALITY. Szymczyk, W.; Byrski, E.; Polak, B. (Inst. of Oncology, Krakow). *Radiobiol.-Radiother.*; 12: No. 6, 797-801 (1971).

Two thermoluminescent phosphors LiF:Mg, Ti and $\text{CaF}_2\text{:Mn}$ made in Poland were used for effective energy determination of an X-ray beam penetrating a water medium. The results obtained are compared with the published data. (auth)

45623 COMMERCIAL THIMBLE CHAMBERS FOR ABSORBED DOSE MEASUREMENTS AT HIGH ENERGY ELECTRON RADIATION. Svensson, H.; Pettersson, C.; Hettlinger, G. (Umeaa Univ., Sweden). *Acta Radiol., Ther., Phys., Biol.*; 10: No. 5, 504-12 (Oct 1971).

The absorbed dose calibration factors for 6 types of commercial thimble ionization chambers at the point of maximum absorbed dose were determined with electron beam energies between 8 and 30 MeV. The influence of several correction factors dependent on the chamber construction are discussed. The properties of a thimble ionization chamber suitable for absorbed dose measurements in high energy electron beams are summarized. (auth) (Sweden)

45624 PRESENT STATUS OF PERSONNEL DOSIMETERS. Moriuchi, Yasuyuki. *Nippon Genshiryoku Gakkaishi*; 14: No. 6, 295-306 (Jun 1972). (In Japanese).

Basic considerations on the significance of personnel dosimeters are discussed; these are followed by an analytic examination of the factors that affect the precision and accuracy of the dose measurements. The factors are classified into the following four categories: theoretical characteristics based upon the dose measuring principles, quality control techniques applied to the detector elements and measuring instruments, techniques of pre- and/or post-irradiation treatments, and techniques of normalization against dose standards. Four kinds of personnel dosimeters currently used in Japan (film badge, pocket dosimeter, glass dosimeter, and thermoluminescent dosimeter) were inspected in accordance with the national exposure standards issued by ETL. The results obtained are presented with reference to systematically and objectively reconsidered theoretical characteristics such as the energy response and to data indicating the level of reliability of personnel dose monitoring systems available today in Japan. The results reveal that the reliability is largely dependent on the suitability of the monitoring system for securing good traceability against the dose standards and on proper post-irradiation treatment. It is concluded that among the four kinds of dosimeters, the film badge system sometimes appears to be most doubtful in accuracy. In view of the increasing importance of measurements of genetically significant dose, medically considered scattering radiation dose, and environmental radiation dose, the reliability of the available personnel dose measurements would appear to call for improvement the world over. (15 figures, 1 table) (auth)

45625 DOSIMETRY PROBLEMS WITH BREMSSTRAHLUNG IN THE GeV RANGE. Zywieltz, Friedrich (Univ., Hamburg). *Stud. Biophys.*; 23: No. 1, 61-8 (1970). (In German).

An experimental determination of the absorbed dose of high-energy bremsstrahlung ($E_{\text{max}} = 6200 \text{ MeV}$) was undertaken using the ferrous sulfate dosimeter. The experiments were carried out at the Deutsches Elektronen-Synchrotron (DESY). The irradiation technique, the beam properties, and the dosimetry are described. From the irradiation of the ferrous sulfate solution in a 6.2-GeV photon beam and the data of the time-integrated quantum flux by a quantimeter a dose-flux relation of $0.73 \times 10^{-7} \text{ rad/Q}_{\text{eff}} \text{ cm}^{-2}$ resulted. For the calculation of the absorbed dose, a G value of 15.6 was used. The development of the electromagnetic cascade in a water phantom leads to an increase in dose rate with depth, and this was measured with the ferrous sulfate dosimeter for various energies. (5 figures) (auth)

45626 MICRODOSIMETRY MEASUREMENTS OF FAST NEUTRONS WITH A TISSUE-EQUIVALENT PROPORTIONAL COUNTER OF 1 CM DIAMETER. Krueger, H.; Graul, E. H. (Philipps-Univ., Marburg, Ger.). *Stud. Biophys.*; 23: No. 1, 27-36 (1970). (In German).

For neutron-therapy investigations it is important to know not only depth-dose curves but also the changes of radiation quality in the human body. A ball-shaped tissue-equivalent proportional counter of 1 cm diameter, which was built for this purpose, is briefly described. LET spectra as well as the spectra of "local specific energy" and their depth-dependent changes were inferred from the microdosimetric measurements made with the proportional counter in a tissue-equivalent trunk phantom. Discussions of the spectra measured are included. (8 figures) (auth)

45627 DOSIMETRY OF PULSED X-RAYS AND ELECTRONS FROM A LINEAR ACCELERATOR. Antoku, Shigetoshi; Sunayashiki, Tadashi; Takeoka, Seiji; Takeshita, Kenji (Hiroshima Univ.). *Hiroshima Daigaku Genbaku Hoshano Igaku Kenkyusho Nenpo*; No. 12, 69-76 (1971). (In Japanese).

To use ionization chambers for dosimetry of pulsed radiations from the accelerator, the ion-recombination and energy dependence of the ionization chambers were studied by comparing them with chemical and thermoluminescence dosimeters. The ionization chambers indicated lower readings than the true values by recombination of ions. Readings of an Ionex 0.6-cc chamber were

about 20 percent lower than those of the chemical and thermoluminescent systems, at a dose rate of 1,500 rad/min (1.7 esu/pulse). The collection efficiencies of Ionex and Radocon dosimeters determined experimentally agreed with those calculated using $750 \text{ V} \cdot \text{cm} \cdot \text{esu}$ as a constant μ in a theoretical formula. The sensitivity changes according to radiation and its energy were established for both dosimeters. The sensitivity increased by 7–9 percent for 10- to 30-MeV electrons and decreased by 0–5 percent for 10- to 35-MV x-rays according to a ^{60}Co γ -ray standard. Field homogeneities in effective fields (80% of total area) were less than 10 percent for both electrons and x-rays in the therapy mode. They were less homogeneous for x-rays in the research mode. It is difficult to infer a specific dose to biological materials using ionization chambers because of variations in collection efficiency by output from the accelerator. Lack of homogeneity of fields was able to be reduced by rotating the irradiation cage. (auth) (Japan)

45628 UTILIZATION OF ALUMINA RADIOTHERMOLUMINESCENCE IN ZONE DOSIMETRY. EXPERIMENTAL APPLICATION IN FRENCH NUCLEAR TESTS IN THE PACIFIC OCEAN. Portal, G. (CEN, Fontenay-aux-Roses, France); Marchal, P.; Medioni, R.; (and others). Radioprotection; 7: No. 1, 21–43 (1972). (In French).

Roof tiles containing quartz and feldspars were used to estimate the radiation doses resulting from the atomic bomb explosions at Hiroshima and Nagasaki. This method might be used in other applications, for example, the dosimetry of criticality accidents. It was suggested that a large quantity of radiothermoluminescent material could be incorporated systematically in the bricks used in certain constructions. Investigations led to the selection of an industrial quality of aluminum oxide which could be marketed at a reasonable price. The crystallographic and chemical characteristics of this material were examined together with its radiothermoluminescent properties. The material presents a thermoluminescent curve having four energy levels. Only the deepest levels are utilized. With ordinary reading materials, measurements between 0.5 and 10^6 rads can be effected. This material is almost insensitive to neutrons, a property enabling it to be used in mixed fields. Dosimetry effected with alumina bricks in conjunction with French nuclear tests in the Pacific is presented. (France)

45629 PERSONAL PHOTODOSIMETRY IN RADIODIAGNOSIS AND RADIATION THERAPY ROOMS. Balon, Z. P.; Lyalina, L. I.; Kotov, G. V. (Central Scientific-Research Inst. of Roentgeno-Radiology, Leningrad). Med. Radiol.; 17: No. 5, 43–6 (May 1972). (In Russian).

Results obtained by investigating characteristics of personal photodosimeters in a radiation range from 10 to 500 keV are summarized, and data from studies of a new improved personal photodosimeter are presented. Photodosimeters have an energy dependence within a range of 10 to 200 keV, the character of the sensitivity variations being similar for different types of photodosimeters and films. This restricts the possibility of using dosimeters for x-ray diagnosis and therapy. Methods of recording energy dependence in the x-ray range of 40 to 200 kV suggested by some authors may find practical use only after these recommendations will have been taken into account in designing the device under consideration with strictly standardized methods of measurements. (3 figures, 2 tables) (auth)

45630 COMPARISON OF NEUTRON AND GAMMA-RAY PERSONNEL DOSIMETERS AT CRITICAL ASSEMBLIES. Keirim-Markus, I. B.; Kraiton, S. N.; Kuz'min, V. V.; (and others). At. Energ. (USSR); 33: No. 1, 563–5 (Jul 1972). (In Russian).

Comparative results of neutron and gamma-ray personnel dosimeters are presented. It is shown that data obtained by means of neutron-track dosimeters and thermoluminescent (CaS) dosimeters agree within 30% and, when measurements are conducted with sufficient statistical accuracy, 15%. There is agreement within the same limits for gamma-ray dosimeters: pocket film (IFK-2,3), glass (IKS), and LiF. The results obtained by means of neutron personnel dosimeters agree with LET spectra measured by a set of tissue-equivalent low-efficiency Geiger counters and neutron spectra measured by means of threshold detectors. (2 figures, 2 tables) (auth)

45631 FILM TECHNIQUE FOR DETERMINING THE ABSORBED DOSE AT EXIT FROM THE PATIENT DURING RADIATION EXPOSURE. Haus, A. G.; Strubler, K.; Marks, J. E. (Univ. of Chicago). Radiology; 104: No. 1, 197–200 (Jul 1972).

A film technique for measuring the absorbed dose at exit from a patient during radiation exposure was studied with the ^{60}Co and Van de Graaff teletherapy machines. Film response was investigated at several field sizes and depths. Results indicate that a standard film calibration curve can be obtained to calculate ab-

sorbed dose. The treatment verification film is obtained by inserting the film in a thin vinyl flexible cassette which remains beneath the patient throughout the complete radiation exposure. The absorbed exit dose can be obtained from the verification film with the calibration curve. (auth)

45632 RADIO-FREQUENCY (OSCILLOMETRIC) ALCOHOLIC CHLOBENZENE DOSIMETER. Horvath, Zsuzsanna; Banyai, Eva; Foldiak, Gabor (Magyar Tudományos Akademia Izotop Intezete, Budapest). Magy. Kem. Foly.; 76: No. 10, 504–5 (Oct 1970). (In Hungarian).

Oscillometric analysis as a new method in alcoholic chlorobenzene chemical dosimetry makes possible direct measurement in sealed ampoules, thus reducing the duration of a measurement from 8 to 10 minutes (in the case of conventional titration method) to 1 minute. By performing a sufficiently large number of measurements, the accuracy and reproducibility of the dosimetric system may further be improved. As shown by the results of thousands of measurements, the method can be used for routine work (for a point-to-point determination of radiation fields of irradiation facilities, regular dose monitoring, etc.). (auth)

45633 INFLUENCE OF A CONTAINER'S MATERIAL FOR INDICATIONS RENDERED BY THE CHEMICAL METHYL-ORANGE DOSIMETER. Kalmykov, L. Z. (Kharkov Scientific Research Inst. of Medical Radiology). Med. Radiol.; 17: No. 6, 90–3 (Jun 1972). (In Russian).

Evidence is offered proving that a methyl-orange solution in concentrations of $5 \cdot 10^{-6} \text{ M}$ in a 0.01 M nitric acid, employed as a chemical dosimeter, may be exposed to irradiation in vessels made of paraffin, polymethylacrylate (plexiglas), polyethylene, and plastic polyvinyl chloride in addition to glass vessels. (auth)

45634 MEASUREMENT OF ABSORBED DOSES WITH FLUID IONIZATION CHAMBERS. Linchevskaya, G. A.; Luk'yanov, L. A. (Moscow Scientific Research Roentgeno-Radiological Inst.). Med. Radiol.; 17: No. 6, 80–3 (Jun 1972). (In Russian).

The technique and results of direct measurements of absorbed radiation energy by means of ionization chambers using a tissue-equivalent fluid with high electric resistance (heptane + 3.5% HCl_4) are described. The operating principle of such fluid chambers is based on the fact that under the influence of ionizing radiation the conductivity of fluid dielectrics is increased. The dependence of the ionizing current on the electric field intensity in the chamber, dose, and radiation energy was investigated. Absorbed energy values in the operating volume of the fluid chamber were obtained. (auth)

45635 ABSORBED IONIZING RADIATION MEASURING DEVICE. Falk, Fred W. U. S. Patent 3,676,682. 11 Jul 1972. Filed 30 Oct 1968.

An ionization chamber for measuring absorbed radiation is disclosed. The chamber includes a housing which incorporates an anode, a cathode, and a space charge electrode. The anode comprises a conducting plate which has a plurality of protruding needle points with each of the needle points and the conducting plate being at the same electrical potential. The space charge electrode has a plurality of holes therein and is physically displaced from the anode with the holes of the space charge electrode being in registry with the needle points of the anode. The potential of the space charge electrode is less than that of the anode itself. (auth)

Nuclear Spectroscopic Instrumentation

Refer also to abstracts: 45264, 46144, 46893, and 46966.

45636 (BARC-591) SIMPLE TRANSISTORISED CONTROL UNIT FOR NEUTRON CRYSTAL SPECTROMETERS. Shah, V. M.; Singh, Virender; Rao, K. R. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 15p. Dep. NTIS (U. S. Sales Only).

A simple transistorized version of a control unit which can be used for auto-control of neutron spectrometers is described. Although the basic circuitry is discussed in terms of operation of a triple axis spectrometer (or a filter detector spectrometer), the same unit can be used for control of a diffractometer used for studying powder samples. The system is found to be reliable and extremely simple to operate. The design incorporates features of sequential and parallel drive of several motors in the forward and reverse directions. This helps in an optimum time scan of phonons, several of them one after another. (auth)

45637 (JINR-P13-6428) SIGNAL FILTRATION IN THE AMPLITUDE SPECTROMETRY SYSTEMS. Vakhtel, V. M.; Semenov, B. Yu. (Joint Inst. for Nuclear Research, Dubna (USSR), Lab. of Nuclear Problems). 1972. 23p. (In Russian). Dep. NTIS (U. S. Sales Only).

The following problems have been considered: (1) The production of the maximum signal-to-noise ratio in the standard systems. (When considering this problem the notion of the system weight (memory) function and that of the effective correlation coefficient are employed, which simplifies the procedure). (2) The achievement of maximally possible signal-to-noise ratio in the system of amplitude spectrometry. (3) The optimization of the parameter selection of some practically important types of RC-filters. (4) A proper choice of the time constant of the connecting RC-circuit between the amplifier and the analog-to-digital converter. The Appendix covers the system memory function and its relation to a widely-known notion of the system impulse response. Also, the material is systematized concerning the ways of describing the signal, the noise, and the system characteristics. This material is necessary in solving filtration problems. (auth)

45638 MAGNETIC FIELD INVERSELY PROPORTIONAL TO r WITH A PERIODIC AZIMUTHAL INHOMOGENEITY. Saulit, V. R. (Leningrad State Univ.). Sov. Phys.-Tech. Phys. (Engl. Transl.); 17: No. 1, 34-7 (Jul 1972).

Translated from Zh. Tekh. Fiz.; 42: No. 1, 48-53 (Jan 1972).

A magnetic field antisymmetric with respect to the plane $z = 0$ is specified by the expression $H(r, \varphi) = 1/r \sum_{k=0}^n (a_k \cos k\varphi + b_k \sin k\varphi)$, where a_k and b_k are arbitrary coefficients. The field in the space adjoining this plane and the shape of the appropriate pole piece are determined. The problem is solved in finite form without series expansions. The surface of each pole piece has a relatively simple geometric shape, which is easy to reproduce; it is formed by the geometric locus of points on straight lines intersecting in a single common point. This field can be used in the spectrometry of charged particles with moderate energies. (1 figure) (auth)

45639 FAST-SLOW COINCIDENCE CIRCUIT SYSTEMS FOR Ge(Li) DETECTOR. Sato, Yuji; Azuma, Toshio. Annu. Rep. Radiat. Center Osaka Prefect.; 12: 22-7 (1971).

To measure gamma-ray spectra from isotopes, a fast slow coincidence circuit system is tested. The resolving time of fast coincidence is 11 nsec FWHM for NaI-Ge. That of Ge-Ge is 21 nsec FWHM. A time-to-pulse height converter is used to measure short times of nsec order. The time delays between two inputs (NaI-Ge and Ge-Ge) are adjusted to make correct coincidence. (auth)

45640 SOURCE OF HIGH-INTENSITY NANOSECOND LIGHT PULSES. Elenski, Ya. S. (Inst. of Physics, Moscow). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 4, 1165-6 (Jul-Aug 1971).

Translated from Prib. Tekh. Eksp.; 14: No. 4, 183-4 (Jul-Aug 1971).

It is shown that a pulsed gas laser may be used to simulate the passage of showers of high-energy particles, as well as individual particles, through a scintillation counter. In either case the spectra and duration of the luminescence are close to the simulated values. (auth)

45641 PULSED ACCELERATOR TIME-OF-FLIGHT SPECTROMETERS. Rae, Ernest R. (Atomic Energy Research Establishment, Harwell, Eng.). pp 1-99 of Experimental Neutron Resonance Spectroscopy. /Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970).

The only technique that has proved successful for studying neutrons from 1 eV to 100 keV is that of neutron time of flight using a continuous-energy-spectrum neutron source. Although a number of accelerators can be used, the most versatile is the electron linac, and the bulk of this chapter is devoted to the present status of this device. The performances of a number of spectrometers are compared. The pulsed van de Graaff is also described, but its use is restricted to neutron energies above 2 keV. (52 figures, 3 tables, 166 references) (R.W.R.)

45642 MULTICHANNEL MODULAR SPECTROMETER FOR LOW-ENERGY ELECTRONS AND PROTONS. Vaisberg, O. L.; Zhurina, L. S.; Kovalenko, V. G.; (and others). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1631-3 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 42-4 (1971).

An electron and proton spectrometer is described which is designed for measurement of the characteristics of charged-particle fluxes in the range of energies from 0.03 to 10 keV. An energy analysis of the particles is carried out by electrostatic analyzers, while detection is performed by channel electron multipliers. (auth)

45643 LOW-ENERGY-ELECTRON SPECTROMETER HAVING A MULTIPLATE ANALYZER. Lein, E. L.; Ol'dekop, L. G.; Polenov, B. V.; (and others). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1634-7 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 44-8 (1971).

A spectrometer is described which was designed for studying

a flux of electrons having an energy of from 1 to 10 keV. The use of multiplate analyzers having a relatively high resolution allowed the sensitivity of the flux-density measurements to be increased in the presence of ultraviolet radiation. The fundamental electronic sections and the block diagram of the instrument are described. (auth)

45644 X-RAY SPECTROMETER HAVING A COOLED Si(Li)-DETECTOR. Arl't, R. D.; Banifatov, A. E.; Kalinin, A. I.; Petrov, M. M.; Tyvin, L. I. (Joint Inst. for Nuclear Research, Dubna, USSR). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1638-40 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 48-50 (1971).

An x-ray spectrometer with an Si(Li)-detector and a field-effect transistor mounted in a cryostat chamber is described. A resolution of 650 eV was obtained for the 6.4 keV line from ^{57}Co . (auth)

45645 CHARACTERISTICS OF CHARGED-PARTICLE DETECTION SECTIONS HAVING CHANNEL ELECTRON MULTIPLIERS. Kovalenko, V. G.; Polenov, B. V. Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1651-3 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 59-61 (1971).

The characteristics of the detection sections used in a low-energy charged-particle spectrometer are presented. The sections studied contain a cylindrical electrostatic analyzer and a channel electron multiplier having an input-window area $0.2 \times 0.6 \text{ cm}^2$. (auth)

High Energy Physics Instrumentation

45646 (RPP/H-89) PLASTIC THRESHOLD CERENKOV DETECTORS WITH WAVELENGTH SHIFTER AND QUENCHER. Sacharidis, E. J. (Rutherford High Energy Lab., Chilton (England)). [nd]. 27p. Dep. NTIS (U. S. Sales Only).

Tests on the threshold behavior of plastic Cherenkov radiators containing a wavelength shifter and quencher are described and discussed. It is shown that they lead to improved photoelectron yield combined with angular uniformity and low scintillation level above 0.500 GeV/c. (auth) (UK)

45647 (UCRL-51214) PSEUDO CLOUD-CHAMBER TRACKS AND THEIR USES. Ernst, Richard D. (California Univ., Livermore, Lawrence Livermore Lab.). 13 Apr 1972. Contract W-7405-eng-48. 27p. Dep. NTIS.

Pseudo cloud-chamber tracks are artificial tracks superimposed by double exposure on otherwise normal photographs of real cloud-chamber tracks. The pseudo tracks were originally developed as a means of training scanners to look for low-density (quark) tracks; since then, many other uses have arisen. Such uses include (1) checking the model used in generating the pseudo tracks, (2) investigating the effects of using different gases in the model, (3) training scanners to hand-count tracks under a microscope, (4) normalizing the hand-counted data, (5) keeping the scanners alert by using the pseudo tracks as "carrots," (6) determining how efficient the scanners are in finding quark tracks (and in the process proving that such tracks can be found), (7) analyzing the automatic scanning system, and (8) studying the feasibility of new approaches to the search for quarks. (auth)

45648 DETECTOR FOR HIGH ENERGY PARTICLES BASED ON THE PRINCIPLE OF USING PARTICLE RADIATION IN A LAMINATED MEDIUM. Arutyunyan, F. R.; Nazaryan, A. A.; Torgomyan, F. B.; Frangyan, A. A.; Ekimyan, G. A. (Inst. of Physical Research, Erevan). Dokl. Akad. Nauk Arm. SSR; 53: No. 1, 11-15 (1971). (In Russian).

The "resonance radiation detector" (RRD) is analogous to a Cherenkov counter, both in the concept of separating the direction of particle movement and in the threshold character of the radiation; but it differs from the Cherenkov counter in its ability to operate in the region of large values of E/mc^2 . The authors had previously used the RRD in the detection of muons in cosmic rays. Experiments are reported here on the use of the RRD for 2.8- and 3.7-GeV electrons from a synchrotron. The beam of π^- and e^- , after passage through a laminated medium, entered a gas threshold Cherenkov counter and a full absorption Cherenkov counter. The latter separated the electrons from the beam of π^- mesons. (3 figures) (K.S.W.)

45649 BUBBLE DENSITY DETERMINATION IN BUBBLE CHAMBERS. Ganovici, I.; Nichtin, Fl. An. Univ. Bucuresti, Ser. Stiint. Natur., Fiz.; 19: 9-17 (1970).

The gap distribution function was extended to negative gaps. The gap average length, size of the vacancies, and gap density were measured for two cases, viz. with and without consideration of the spots. Bubble density was measured from the average value of the bubble plus the spot size ($\bar{D}_{b+sp}$). (Rom. Sci. Abstr.)

45650 PERFORMANCE LOGIC OF SPARK CHAMBER INFORMATION READ-OUT BY TELEVISION AUTOMATION AND DATA LINK TO A COMPUTER. Klevalin, V. A.; Belyakov, E. S.; Bernshtein, L. I.; Lebedev, B. A.; Petrakov, A. V. *Izv. Akad. Nauk Arm. SSR, Ser. Fiz.*; 6: No. 3, 219-22(1971). (In Russian).

The on-line performance logic for spark chamber readout by television automation coupled to a small Razdan-2 computer is given. The information is transferred to computer during the backward motion of a vidicon beam. The information transference of 6 tracks in the spark chamber takes 100 μ s for each line for the direct motion of 300 μ s. (3 figures) (auth)

45651 SYSTEMATIC METHODS FOR THE ANALYSIS OF A SAMPLE OF NUCLEAR INTERACTIONS IN HYDROGEN BUBBLE CHAMBER. Huq, S. S. (Univ. of Rajshahi, Pakistan). *Nucleus* (Karachi); 8: No. 4, 63-76(1971).

Systematic methods of analysis of some 15,000 bubble chamber pictures of nuclear interactions involving inelastic collision of a 5 GeV/c K^+ beam on proton giving rise to four charged secondary tracks are presented. Final state reactions of the type $K^+p \rightarrow K^+\pi^+\pi^-p$, $K^+ + p \rightarrow K^+\pi^+\pi^-p$ (π^0 unseen), $K^+p \rightarrow p\pi^+\pi^-\pi^+$ (K^0 unseen) $K^+p \rightarrow K^+\pi^+\pi^-\pi^+$ (n unseen), and $K^+p \rightarrow pK^+K^+$ were uniquely identified from the analysis. (auth)

45652 COMPENSATION OF THE DRIFT OF CHARGED-PARTICLE TRACKS IN WIDE-GAP SPARK CHAMBERS. Avakyan, K. M.; Aivazyan, R. B.; Matevosyan, E. M. *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 6, 1645-7(1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 54-6(1971).

A method is described for supplying wide-gap spark chambers with high-voltage rf pulses. It is shown that for a sign-alternating field there is no displacement of the spark tracks relative to the true charged-particle trajectory due to drift of avalanches. A description is given of a generator which allows high-voltage rf pulses having a frequency of from 3 to 8 MHz to be obtained. (auth)

Radiometric Instruments

Refer also to abstracts: 45717 and 47267.

45653 (CSIRO-Trans-10705) CALIBRATION OF RADIOACTIVE LOGGING APPARATUS AND NEUTRON MODELING. Laszlo, Andrassy. Translated by G. Losonci for Commonwealth Scientific and Industrial Research Organization, Australia, from Magy. Allami Eotvos Lorand Geofiz. Int. Evi Jelentese; 137-40 (1968). 2p. Dep. NTIS (U. S. Sales Only).

A calibration technique for the two-detector gamma-gamma logging method using a borehole model was developed and gave the advantage of optimal pressing onto the wall. Calibration measurements for epithermal neutrons in water (100%), oily sand (45%), limestone (0%) and a glass tube model (6%) are shown. (Australia)

45654 SYSTEM AND METHOD FOR FACILITATING THE OPERATING SETUP OF A RADIATION GAUGE. Doering, George I. (to Industrial Nucleonics Corp.). U. S. Patent 3,668,398. 6 Jun 1972. Filed 26 Dec 1968.

Disclosed are a system and method for facilitating the setup of a nucleonic gage and automatic controller for measuring and controlling the thickness of a material. The gage includes conventional calibration circuits having variable voltage dividers set in accordance with the expected operating points for the material thickness center scale and gage sensitivity. In one embodiment, the sensitivity and operating point voltage dividers as well as the automatic controller set-point voltage divider are set with the aid of an auxiliary bridge including additional voltage dividers ganged with the voltage dividers of the gage calibration circuits. In another embodiment, one of the auxiliary bridge voltage dividers has a logarithmic response, enabling calibration for different materials to be established by changing the input voltage to the auxiliary bridge. (auth)

45655 APPARATUS FOR MEASURING HEIGHT OF A MOL-TEN METAL POOL. Blecherman, Sol S.; Ulion, Nicholas E.; Packer, Louis L. (to United Aircraft Corp.). U. S. Patent 3,668,386. 6 Jun 1972. Filed 13 Mar 1969.

An apparatus for detecting the liquid level in, or the liquid-solid interface at the bottom of, a pool of molten metal or alloy utilizing a radioactive isotope as a source of radiation and sensing the amount of radiation over or through the pool of metal and in which the melting is accomplished by an electron beam impinging on the pool surface is described. (auth)

45656 IMPROVEMENTS TO EQUIPMENTS FOR MEASURING THICKNESS BY BACKSCATTERING. Levenes, Guy (to Societe

Industrielle de Combustible Nucleaire). French Patent 2,070,925. 23 Aug 1971. Filed 11 Dec 1969. (In French).

A probe intended to be used in conjunction with apparatus for measuring the thickness of thin films by β particle backscattering is presented. The system utilizes a radioactive source and a Geiger-Muller counter. The axis of the source, the axis of the G M counter, and the normal to the surface examined lie in the same plane. The angles formed between the axis of the source and the axis of the G M counter and the normal are made as small as possible and arranged to be on either side of the normal. A description is given of probes primarily intended to measure the thickness of thin films deposited inside tubes, or cavities having cylindrical or conic surfaces, or on large surfaces. (France)

45657 RADIOMETRIC UNIT FOR MEASURING WEAR OF MATERIALS AND MACHINE PARTS. Ivanov, E. A.; Magarik, K. N. *Izotopy SSSR*; No. 22, 29-31(1971). (In Russian).

A two-channel automatic device for studying resistance to wear of machine parts which is based on the radioindicator method is described. A two-component isotopic mixture consisted of ^{60}Co and ^{75}Se in lubricating oil. The ^{60}Co was used to label a cast iron cylinder sleeve and ^{75}Se , a chrome-plated piston ring of a Diesel engine. The merit of the system was demonstrated by its sensitivity ($\sim 2 \times 10^{-8}$ Ci/l), its reliability, and its ease of operation. The possibility of automatic measurement and recording of the spectra of a multi-component isotopic mixture is explored. (tr-auth)

45658 SEMICONDUCTOR WELL LOGGING PROBE. Andrieux, Claude; Dumesnil, Patrice; Javellaud, Jean (to Commissariat a l'Energie Atomique). French Patent 2,070,479. 16 Aug 1971. Filed 5 Dec 1969. (In French).

A probe was developed for radioactive well logging using high-resolution γ spectrometry. The probe contains an intrinsic semiconductor nuclear particle detector device, a liquefied gas cryostat, a high voltage supply for the detector, and an appropriate preamplification device. The intrinsic semiconductor is made of Ge or Si in which the residual impurities are compensated by Li. This probe can be used to measure the ^{235}U content of ore. (France)

45659 RADIOACTIVE RADIATION-TYPE FLUID INTERFACE DETECTION SYSTEM. Bajek, Walter A.; Boyd, David M. Jr. (to Universal Oil Products Co.). U. S. Patent 3,668,392. 6 Jun 1972. Filed 20 May 1969.

Method and apparatus for locating one or more liquid-liquid interfaces within a closed vessel by utilizing a radiation-type interface detector are described. A traveling radiation source within the vessel and a traveling radiation detector outside the vessel, the paths of travel of both being vertical and the speed of both being synchronized, are used to provide constant path length between them. The instrument output is in the nature of a density profile. (5 figures) (auth)

Radiation Effects on Instrument Components, Instruments, and Electronic Systems

Refer also to abstracts: 45438, 45620, and 47230.

45660 (FRNC-CONF-29) NOISE IN Au-nSi DIODES OPERATING IN THE AVALANCHE REGION IN THE PRESENCE OF CHARGES INJECTED BY RADIATION. Peyre-Lavigne, A.; Cassanhol, B.; Peyre-Lavigne, M. (Toulouse Univ. (France). Centre de Physique Atomique et Nucleaire). 1971. 15p. (In French). (CONF-710958-1). Dep. NTIS (U. S. Sales Only).

From International conference on the noise in active semiconductor devices; Toulouse, France (21 Sep 1971).

Results are given of measurements made on Au-nSi and Al-pSi diodes (with leakage current collection rings made by metallization under vacuum) when the avalanche potential is reached. A temperature of 300°C is not exceeded during their manufacture, thus conserving the lifetime of the minority carriers. Surfaces as large as 12 mm² were prepared, which did not exhibit microplasmas at avalanche potentials reaching 1800 volts. This fact is in good agreement with theoretical values obtained by Van Overstraeten. The maximum coefficient of amplification did not exceed 20. The diodes (metal-semiconductor) were used to establish a model taking into account the current from the metal-semiconductor contact and the channel, the latter being linked to the surface potential and the density of surface states characteristic of a given passivation. This model enables good agreement to be obtained between theory and experiment in the analysis of shot noise. A

method is given for analyzing shot noise produced by radiation penetrating a diode operating in the multiplication region. (France)

45661 (N-72-17158) EFFECTS OF 22-MeV PROTONS ON SINGLE-JUNCTION AND SILICON-CONTROLLED RECTIFIERS. Beatty III, Marvin E. (National Aeronautics and Space Administration, Langley Station, Va. Langley Research Center). Feb 1972. 31p. (NASA-TN-D-6631). NTIS.

The effects of 22-MeV protons on various types of silicon single-junction and silicon-controlled rectifiers were investigated. The results show that low-leakage devices and silicon-controlled rectifiers are the most susceptible to radiation damage. There are also differences noted between single-junction rectifiers of the same type made by different manufacturers. This finding emphasizes the need for better selection of devices used in spacecraft. (auth) (STAR)

45662 (SC-DC-721232) DEVICE DEGRADATION FROM THE EFFECTS OF NUCLEAR RADIATION ON PASSIVATION MATERIALS. Coppage, F. N.; Graham, E. D. Jr. (Sandia Labs., Albuquerque, N. Mex.). Jul 1972. 19p. (CONF-720707-2). Dep. NTIS.

From Annual conference on nuclear and space radiation effects; Seattle, WA. (24 Jul 1972).

Bipolar transistors that utilized three different commercial passivation processes were studied in a nuclear radiation environment to determine if the passivation process influences the radiation tolerance of the device. Comparisons were made between devices obtained from the same manufacturer. The passivation processes studied featured silicon dioxide, silicon nitride, and aluminum oxide. Irradiations were performed with the devices in both a biased and unbiased mode. The results show that the device with silicon dioxide passivation degraded the most. The nitride devices having beam leads showed no saturation of the induced degradation while the other processes did for a dose of 1×10^7 (Si). All devices studied showed a small bias dependence. The addition of an aluminum oxide or silicon nitride passivation layer (over the silicon dioxide surface) for increased reliability definitely induces no increased radiation degradation. (auth)

45663 (SC-DC-721233) RADIATION EFFECTS IN MICROWAVE BIPOLAR TRANSISTORS. Graham, E. D. Jr.; Chaffin, R. J.; Gwyn, C. W. (Sandia Labs., Albuquerque, N. Mex.). [nd]. 11p. (CONF-720707-3). Dep. NTIS.

From Annual conference on nuclear and space radiation effects; Seattle, WA. (24 Jul 1972).

The effects of gamma and neutron radiation on the high frequency characteristics of microwave transistors (HP-35821E) are explored. From theoretical and experimental results, it is determined that the major effect is a decrease in the forward transfer coefficient (S_{21}) resulting from a neutron-enhanced decrease in cutoff frequency (f_c) (i.e., increase in emitter-to-collector transit time). For the recommended operating conditions ($V_{CE} = 15V$, $I_C = 15$ mA) no change in S-parameters was noted after 10^7 rads (Si) gamma. By increasing the collector voltage so that the epitaxial collector is entirely depleted, these devices can be operated at fluence levels of 8×10^{15} n/cm² ($E > 10$ keV, 14-Mev equivalent). (auth)

45664 (SC-DC-721237) CARRIER TRAPPING EFFECTS IN IRRADIATED PN JUNCTIONS. Gwyn, C. W. (Sandia Labs., Albuquerque, N. Mex.). [nd]. 25p. (CONF-720707-4). Dep. NTIS.

From Annual conference on nuclear and space radiation effects; Seattle, WA. (24 Jul 1972).

Numerical calculations have been used to study small signal trapping effects in neutron irradiated pn junctions as a function of frequency. Good agreement has been obtained between the calculations and experimental data in n-type silicon using two acceptor centers to model the trapping centers produced by neutron irradiation. A single level donor center has been used for p-type material. These calculations indicate that complex changes in the carrier distributions within the device can be responsible for a rather simple variation in the terminal characteristics of the device. Although good agreement has been obtained between the calculated and experimental results, the calculations suggest that another deep acceptor level (consisting of about 10% of the total trap density) may be present in n-type material which provides a more gradual change in the terminal capacitance with frequency than that noted using the two acceptor centers. (auth)

45665 (SC-DC-721240) RAPID ANNEALING OF FREQUENCY CHANGE IN CRYSTAL RESONATORS FOLLOWING PULSED X-IRRADIATION. King, J. C.; Sander, H. H. (Sandia Labs., Albuquerque, N. Mex.). Jul 1972. 34p. (CONF-720707-5). Dep. NTIS.

From Annual conference on nuclear and space radiation effects; Seattle, WA. (24 Jul 1972).

A transient negative frequency change in AT-cut 125 MHZ 5th overtone quartz resonators has been observed following exposure to an intense burst of x-rays. All natural, Z-growth synthetic and swept Z-growth synthetic resonators suffer a significant initial negative frequency offset. At room temperature, the transient frequency change in natural quartz anneals in approximately ten minutes to a relatively stable negative offset. For the synthetic quartz resonators, appreciable annealing of the initial negative offset continues for a longer period, finally arriving at an offset which is positive in the case of the Z-growth synthetic but negligibly small for swept Z-growth synthetic. The transient frequency change is attributed to a relaxation process which anneals above 165°K (Stage I). This annealable relaxation is superposed on a more stable part (Stage II) which is removed only by heating the crystal well above room temperature. The kinetics of Stage I annealing obeys a $t^{-1/2}$ relationship which is indicative of a three-dimensional diffusion limited annealing mechanism involving point defects. It is proposed that the transient change in frequency is due to the trapping of monovalent cations, most probably H⁺, at substitutional aluminum sites in the crystal lattice. The activation energy governing the diffusion of the defect to the trapping center is estimated to be 0.80 ± 0.5 eV. (auth)

45666 (SC-RR-720515) SEMICONDUCTOR DEVICE DEGRADATION BY HIGH AMPLITUDE CURRENT PULSES. Brown, W. D. (Sandia Labs., Albuquerque, N. Mex.). Jul 1972. 34p. Dep. NTIS.

The results of a lengthy and comprehensive investigation of semiconductor device degradation from gamma-induced nanosecond current pulses are given. Topics discussed include (1) previously published literature on pulse degradation and second breakdown, (2) experimental results obtained in several studies, (3) pulse damage recovery using several annealing techniques, (4) a model of the pulse damage, (5) possible methods of hardening against pulse degradation, and (6) a preventive measure that can be taken to eliminate pulse damage. (auth)

45667 (SC-T-722519) EFFECT OF GAMMA RADIATION ON THE OSCILLATION FREQUENCY OF AT-CUT PIEZOQUARTZ PLATES. Starodubtsev, S. V.; Chubarov, L. B. Translated by Patricia Newman (Sandia Labs., Albuquerque, N. Mex.), from Dokl. Akad. Nauk Uzb. SSR; 24: No. 1, 22-3(1967). 3p. Dep. NTIS.

The change in oscillation frequency of AT-cut native quartz plates subjected to cobalt-60 gamma radiation at 10^3 to 10^{10} rad was studied. Various types of self-excited oscillators employing either tubes or semiconductor triodes were used to generate oscillations in the plates. The oscillation frequency was 2 to 4 MHz. The plates were irradiated up to 2×10^{10} rad with ⁶⁰Co gamma radiation at a dose rate of 600 to 400 rad/sec at 42 to 36°C and 15,000 rad/sec at 90 to 100°; and they were irradiated up to 10^9 rad at a dose rate of 90 rad/sec at 29°C and 250 rad/sec at 32°C. No significant changes were observed in the oscillation frequency of the plates. The measurement accuracy was 0.01%. (W.H.K.)

45668 (UCRL-73974) X-RAY-INDUCED PHOTOCONDUCTIVITY IN DIELECTRIC FILMS. Weingart, R. C.; Barlett, R. H.; Lee, R. S.; Hofer, W. (California Univ., Livermore, Lawrence Livermore Lab.). 16 Jun 1972. 55p. (CONF-720707-10). Dep. NTIS.

From Annual conference on nuclear and space radiation effects; Seattle, WA. (24 Jul 1972).

The conductivity induced in films of polyethylene, epoxy, polytetrafluoroethylene, polyethylene terephthalate, polyimide, and glass by x rays at dose rates between 10^3 and 10^{10} rads/sec (dose in air) were measured. The films were 0.05 to 1.25 mm thick. The x-ray spectrum peaked in the vicinity of 10 keV, and the x-ray pulse width was about 40 nsec FWHM. X-ray-induced photo-currents were found to obey Ohm's law at low bias voltages (less than 1 kV). Above 1 kV, however, it was observed that the peak photoconductivity signals from some of the 0.05-mm-thick materials began to increase at a slightly faster than linear rate with bias voltage. The glass samples exhibited no apparent delayed conductivity, while the other sample materials showed various amounts. The magnitude of the delayed conductivity in polytetrafluoroethylene, polyethylene terephthalate, and polyimide depended on the electric field, an effect that is consistent with Poole-Frenkel field-assisted carrier generation. The magnitude and time dependence of the conductivity signals is quantitatively described by a simple trapping model using reasonable values for mobility, trap density, capture cross sections, and trap depths. (auth)

45669 EFFECT OF α PARTICLES ON THE OPERATING REGIME OF A CO₂ LASER. Popescu, Ion M. (Inst. of Atomic Physics, Bucharest); Preda, Alexandru M.; Enache, Ana. Rev. Roum. Phys.; 17: No. 2, 121-3(1972) (In French).

Experimental results from operating a CO₂ laser in the presence of an alpha source are presented. The laser supply voltage decreased, and the power output decreased due to a variation of the emission spectrum in the immediate region of the alpha particles emitted by the source. (tr-auth)

45670 EFFECTS OF X-RAY IRRADIATION ON MAS DIODES. Nakagiri, Masaru; Tsujide, Tohu (Nippon Electric Co., Ltd., Kawasaki, Japan). Jap. J. Appl. Phys.; 11: No. 7, 1054 (Jul 1972).

The shifting characteristics of the flat band voltage of metal-Al₂O₃-Si (MAS) diodes from x-ray irradiation under a certain bias voltage (B-X treatments) are described. B-X treatments shift the flat band voltage of MAS diodes with irradiation time until it reaches a certain saturated value, depending on the bias voltage, after irradiation of 20 minutes. When a sequence of B-X treatments at various bias voltages is carried out, the magnitude of flat band voltage is determined by the bias voltage of the last B-X treatment. It is concluded that x-ray irradiation creates hot electrons in the Al₂O₃ film and metal electrode, those excited electrons are transported by the electric field, and the charge transport forms a space charge in the Al₂O₃ film and it continues until the space charge compensates the external electric field due to the bias voltage. (2 figures) (R.W.R.)

45671 EFFECTS OF IMPURITIES ON GAMMA-IRRADIATED SILICON CRYSTAL EXAMINED BY PHOTOVOLTAIC EFFECT OF P-N JUNCTION DIODE. Yamaguchi, Masafumi; Nagai, Osamu (Nippon Telegraph and Telephone Public Corp., Tokai, Japan). Jap. J. Appl. Phys.; 11: No. 7, 1016-23 (Jul 1972).

Minority-carrier lifetime in ⁶⁰Co-gamma-irradiated silicon bulk crystals was examined by the photovoltaic effect of silicon p-n junctions, and following results were obtained. The damage constant, K, increases with impurity concentration of bulk crystals; K is smaller in p-type crystals than in n-type. For isochronal annealing, recovery stages of n-type bulk crystals at 100 to 150°C and 300°C are found; in p-type crystals, reverse annealing is observed at 200 to 300°C. From the temperature dependence of carrier lifetime, energy levels of the recombination centers at E_c - 0.42 eV for n-type and E_v + 0.32 and E_v + 0.25 eV for p-type crystals are found. Recovery stage of p-type bulk crystals due to gamma irradiation at 81 K is observed at about 150 K, but that of n-type is not observed between 80 and 300 K. The lifetime degradation in p-type crystals due to gamma irradiation at 81 K is reduced when the minority carrier is injected after the irradiation. (13 figures, 2 tables) (auth)

45672 RADIATION-RESISTANT PLASTIC INSULATORS. Sturm, Bernard J.; Parkinson, William W. (to United States Atomic Energy Commission). U. S. Patent 3,663,504. 16 May 1972. Filed 10 Nov 1970.

A high molecular weight organic composition useful as an electric insulator in radiation fields is provided and comprises normally a solid polymer of an organic compound having a specific resistance greater than 10¹³ ohm cm and containing phenyl groups and 1 to 7.5 weight percent of a high molecular weight organic phosphite. In one embodiment the composition comprises normally solid polystyrene having 7.5 weight percent tris-β-chloroethyl phosphite as an additive; the composition exhibited an increase in the post-irradiation resistivity of over an order of magnitude over the post-irradiation resistivity of pure polystyrene. The plastic was developed for use in pocket ionization monitors. (auth)

Miscellaneous Instruments and Components

Refer also to abstracts: 45243, 46093, and 47317.

45673 (UCRL-50007-71-3) HAZARDS CONTROL. Progress Report No. 41, September-December 1971. (California Univ., Livermore. Lawrence Livermore Lab.). 22 Mar 1972. Contract W-7405-eng-48. 27p. Dep. NTIS.

ATMOSPHERE—particle diffusion in, zinc sulfide tracer detector for measuring, pp. 1,2

FILTERS—performance testing of high efficiency air, pp. 11-13
LIVERMORE POOL TYPE REACTOR—operation and repair of, Sept.-Dec., 1971, pp. 14-16

PLUTONIUM-239—determination in waste drums, development of NaI(Tl) scanning system for, pp. 4-6

RAD DOS, THERMOLUMINESCENT—sensitivity to fluorescent and ultraviolet light, pp. 13,14

RESPIRATION—simulation of human, calibration and performance of mechanical apparatus for, pp. 7-11

RESPIRATORS—efficiency of cartridges for, effects of steady state and pulsating flow on, pp. 7

SOILS—radioactivity in, coring sampler for depth profile measurements of, pp. 18-19

VENTILATION—system at LLL, fire protection for, pp.16,17

X-RAY SOURCES—development of tube, for fluorescence analysis of trace elements, pp. 2,3 (C.E.S.)

45674 (UCRL-73665) REVIEW OF PROBLEMS ASSOCIATED WITH ACCELERATION MEASUREMENTS. MacLean, Lee M. (California Univ., Livermore. Lawrence Livermore Lab.). 15 May 1972. 12p. (CONF-720312-11). Dep. NTIS. From Underground nuclear test measurement symposium II; Menlo Park, CA. (21 Mar 1972).

A review of the past four years of acceleration measurements in underground nuclear test environments points out the excessively high percentage of "lost" data channels. Typical data traces from underground nuclear tests and high-explosive simulation tests illustrate a chronological review of problems associated with acceleration measurements. Various design philosophies have evolved to circumvent certain problems. Areas of concern include instrument-cable EMP, poor mount design, sensitive components of accelerometers, low resonant frequencies, and undamped response of accelerometers. As a result of this evaluation of typical recorded data, the validity of most underground accelerometer data is challenged. (14 figures, 1 table) (auth)

45675 CIRCUIT FOR CONTINUOUS ACQUISITION OF RATE RATIO OF TWO PULSE TRAINS. Seda, Josef; Sabol, Jozef (Technical Univ., Prague). Jad. Energ.; 18: No. 5, 163-6 (May 1972). (In Czech).

The problem of continuously dividing two signals, represented by two pulse trains, is considered. Some known ratio circuits are briefly discussed, and their advantages and disadvantages are pointed out. A new improved (especially as to the reliability and linearity of result with respect to the ratio of input signals) ratio circuit is designed. At its output there is available an analog or digital signal which is proportional to the ratio of the signals applied to the input. In addition to the block diagram of the ratio circuit, practical design using integrated circuits is also presented. Moreover, the possible application of the proposed circuit for the processing of signals from an ionizing radiation detector is examined. (7 figures) (auth)

45676 SHAPE OF THE MASS PEAK IN A THREE-DIMENSIONAL QUADRUPOLE MASS SPECTROMETER. Sheretov, E. P.; Zenkin, V. A. (Ryazan Radio Engineering Inst., USSR). Sov. Phys.-Tech. Phys. (Engl. Transl.); 17: No. 1, 160-2 (Jul 1972).

Translated from Zh. Tekh. Fiz.; 42: No. 1, 199-202 (Jan 1972).

Calculations of the shape of the mass peak in a three-dimensional quadrupole mass spectrometer are reported for various conditions for the introduction of the ionizing electron beam. It is shown that, as these conditions change, there is a change in the shape of the mass peak and in the position of the maximum-sensitivity line on the stability diagram. The shape is calculated for the phase method for introducing the electron beam. The calculated results are confirmed by experiment. (3 figures) (auth)

45677 NUCLEAR CHARGED SELF-SUSTAINING LASER. Morrison, Clyde A.; Wortman, Donald E.; Farrar, Ruben T. (to Secretary of the Army). U. S. Patent 3,667,068. 30 May 1972. Filed 17 Aug 1970.

A nuclear-charged solid-state laser which requires no external power supply is described. A large class of single crystals which contain a stable isotope that is potentially radioactive is disclosed. The crystal is doped with a laser impurity ion and is irradiated with nuclear energy such that the stable isotope becomes radioactive to provide a pumping source for the laser impurity ions therein. Optical resonator means can be provided by coating both ends of the crystal with highly reflective surfaces. The radioactive isotope provides a continuous internal power supply over the lifetime of its decay, and thus provides a potentially compact, portable, and inexpensive solid-state laser device. (2 figures) (auth)

45678 INVESTIGATION OF THE INJECTION CHARACTERISTICS OF SYMMETRICAL PIN-DIODES PREPARED BY THE METHOD OF ION IMPLANTATION. Akimov, Yu. S.; Gusev, V. M.; Kurinnyi, V. I.; Ryzhikov, I. V. Radio Eng. Electron. Phys. (USSR) (Engl. Transl.); 16: No. 10, 1714-22 (Oct 1971). Translated from Radiotekh. Elektron.; 16: No. 10, 1917-26 (Oct 1971).

A survey of the basic drift and diffusion theories of double injection relative to symmetrical PIN structures with different width of the I region (W/L_p = 0.3 - 15) is presented. Precise knowledge of the parameters of the individual regions of the P⁺PN⁺ structures, prepared by ion alloying of plates of high-ohmic p-silicon, permitted carrying out both a detailed qualitative and a detailed quantitative comparison of the experimental characteristics with

the theories predicted by drift and, in essence by diffusion double injection. The insufficiency of the diffusion model of Hall in analyzing the long PIN structures is shown and the Kleinman segment of the volt-ampere characteristics was investigated in detail. By quantitative comparison of experiment and theory, the influence of the nonuniform distribution of the impurity in the highly alloyed layers was accounted for. In the long diodes ($W/L_p > 2$) the drift segment of the volt-ampere characteristics of Rashba-Tolpygo-Lampert-Rose that preceded the diffusion segment was detected and investigated. (auth)

45679 REMOVABLE CURRENT LEAD FOR POWER SUPPLY IN SUPERCONDUCTING MAGNETIC SYSTEMS. Anashkin, O. P.; Kurnosova, L. V.; Razorenov, L. A.; Sidiyakina, T. M.; Fradkin, M. I. (Inst. of Physics, Moscow). *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 4, 1230-1 (Jul-Aug 1971).
Translated from *Prib. Tekh. Eksp.*; 14: No. 4, 233-4 (Jul-Aug 1971).

The construction of a removable current lead for power supply in short-circuited superconducting magnetic systems is described. The use of a removable current lead in equipment designed for operation on board an artificial earth satellite ensures reliable operation of the magnetic system in the short-circuit mode and allows a substantial reduction of the overall heat influx to the helium. (1 figure) (auth)

45680 INTRODUCTION OF CORRECTION FOR THE NON-LINEARITY OF A DIRECT-CURRENT Q-METER. Kovalev, A. I.; Shklyarevskii, V. M. (Physical-Technical Inst., Leningrad). *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 4, 1142-4 (Jul-Aug 1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 4, 165-7 (Jul-Aug 1971).

The corrections for the nonlinearity of a direct-current Q-meter are calculated by means of the exact formula for the impedance of the measurement tank circuit. (2 figures, 1 table) (auth)

45681 USE OF THE ADMISSION SYSTEM OF THE MI-1309 INSTRUMENT FOR PRECISION ISOTOPIC ANALYSIS. Golubchina, M. N.; Prilutskii, R. E.; Toksubaev, A. I. (All-Union Scientific-Research Geological Inst., Leningrad). *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 4, 1148-51 (Jul-Aug 1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 4, 170-2 (Jul-Aug 1971).

A study was made of the fractionation of isotopes when gaseous substances are introduced into a mass spectrometer using an admission system typical of instruments belonging to a unified series. It was shown experimentally that the greatest distortion of the isotope composition occurs in the ampule-container section and is associated with the inlet of gas into the container through a shutoff valve. Discrete admission of the sample is suggested. It is likewise shown that the difference between the intensities of the sample and standard gas flows has no effect on the results of the analysis within the limits $\pm 9\%$. (4 figures, 2 tables) (auth)

45682 SOME COMMENTS ON THE SIGNAL-TO-NOISE CHARACTERISTICS OF REAL PHOTOMULTIPLIER AND PHOTODIODE DETECTION SYSTEMS. Santini, R. E. (Purdue Univ., Lafayette, Ind.); Ingle, J. D. Jr.; Crouch, S. R. *Anal. Chem.*; 44: No. 9, 1708-11 (Aug 1972).

Comments are made by R. E. Santini on a previous paper (J. D. Ingle and S. R. Crouch, *Anal. Chem.*, Vol. 43, p. 1331) in which a comparison was made between the signal-to-noise characteristics of a high-gain photomultiplier and a photodiode when the overall gain of the total system including the amplifiers was the same. Additional variables which were not considered in the previous paper are presented and briefly discussed. Also, J. D. Ingle and S. R. Crouch reply to Santini's comments. (W.H.K.)

45683 MOESSBAUER DIFFRACTOMETER HAVING HIGH ANGULAR RESOLUTION. Artem'ev, A. N.; Aleshin, K. P.; Mirzababaev, R. M.; (and others) (Inst. of Atomic Energy, Moscow).

Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1656-8 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 64-6 (1971).

A Moessbauer diffractometer is described which allows measurement of the diffraction spectra of single crystals in a one- or two-crystal spectrometer with an x-ray tube or Moessbauer source. The x-ray crystal-monochromator and the γ -ray crystal-collimator, which are situated at a stipulated angle, allow alignment of the working crystal by an x-ray beam to be used in making the transition to operation with γ radiation. Measurement of the angular dependences may be carried out either automatically with a step $\approx 10''$, or manually with a step $\geq 0.5''$. The constant velocity of the Moessbauer source is used in the installation to carry out angular measurements and to perform motion with a constant acceleration; a multichannel analyzer is used to measure the Moessbauer spectra at constant diffraction angles. (auth)

45684 CALCULATION OF THE DEAD TIME OF HIGH-SPEED AMPLITUDE CONVERTERS. Glushkovskii, M. E. *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 6, 1671-5 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 78-82 (1971).

A procedure is described for the engineering calculation of the dead time and the determination of the counting characteristics of high-speed amplitude converters which operate according to the methods of weighting, A-t code conversion, and accelerating A-T code conversion. The dependences on the shape of the spectrum, the shape of the input pulses, the conversion method, the ratio between the conversion and recording times, and the input loading are considered. The shape of the spectrum is assumed to be exponential. An example is given of determining the counting characteristic of the BAP-6M converter combined with an AI-1024-4 recorder. (auth)

45685 AVERAGE-PULSE-AMPLITUDE METER. Selyutin, V. P. *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 6, 1693-4 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 96-8 (1971).

A network is described that allows measurement of the average value of a pulse amplitude with an accuracy $\sim 0.5\%$ which is achieved by using a control switch in the discharge circuit of the integrating cell. The network was used in investigating the efficiency of charge collection from semiconductor nuclear-particle detectors. (auth)

45686 COMPUTER SYSTEMS FOR PHYSICS RESEARCH CENTERS (REVIEW). Zabiyaikin, G. I.; Polyakov, V. N. (Joint Inst. for Nuclear Research, Dubna, USSR). *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 6, 1593-1606 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 5-19 (1971).

The premises underlying the creation of multimachine systems of electronic computers in physics research centers and the principal features of designing such systems for the purpose of handling experimental data in real time are considered. (79 references) (auth)

45687 ELECTRICAL INSULATION IN ELECTROSTATIC ANALYZERS (REVIEW). Ergakov, V. A.; Kryuchkov, I. I.; Trebukhovskii, Yu. V. (Inst. of Theoretical and Experimental Physics, M Moscow). *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 6, 1607-18 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 20-31 (1971).

The selection of the type of residual gas and its pressure, as well as the electrode material, in order to achieve the maximum electric field intensity in an interelectrode gap of 2 to 20 cm is considered for electrostatic analyzers having plane-parallel electrodes in a vacuum. Problems associated with the choice of insulator material and measures for equalizing the voltage along the insulator are discussed. Construction versions of the high-voltage input leads (for voltages up to 1 MV) into the vacuum chamber are presented. (36 references) (auth)

ISOTOPE AND RADIATION SOURCE TECHNOLOGY

Isotope Separation (Industrial)

Refer also to abstracts: 45289 and 47300.

45688 (MLM-1936(TR)) SEPARATION OF SULFUR ISOTOPES BY A CHEMICAL EXCHANGE METHOD. Egiazarov, A. S.; Kaviladze, M. Sh.; Kerner, M. N.; Oziashvili, E. L.; Esakiya, K. E. Translated for Mound Lab., Miamisburg, Ohio, from *Isotopenpraxis*; 7: No. 9, 379-83(1971). 15p. Dep. NTIS.

A comparison of methods for separating sulfur isotopes to obtain large quantities of highly concentrated sulfur isotopes showed that the most promising one is a chemical exchange method in a sulfur dioxide-bisulfite ion system. However, results obtained by various authors on the values of the elementary separation coefficient and on other parameters of the process are conflicting and inadequate for carrying out the technological process. The elementary separation coefficient, α , was determined by a method for the one-step equilibration of sulfur isotopes at temperatures of 25, 50, and 70°C and on a column with a known efficiency for isotopes $^{32}\text{S}/^{33}\text{S}$ and $^{32}\text{S}/^{36}\text{S}$ at a temperature of 70°C. (auth)

45689 (ORO-694) REPORT OF URANIUM ISOTOPE SEPARATION REVIEW AD HOC COMMITTEE. Benedict, Manson; Berman, Abraham S.; Bigeleisen, Jacob; Powell, Jack E.; Shacter, John; Vanstrum, Paul R. (Oak Ridge Operations Office (AEC), Tenn.). 2 Jun 1972. 68p. Dep. NTIS.

Results of committee efforts to evaluate the potential of uranium isotope separation methods other than gaseous diffusion and the gas centrifuge and to determine whether any other processes warrant AEC research and development are presented. It was concluded that many of the processes have such major qualitative disadvantages relative to gaseous diffusion that quantitative evaluation of their principal characteristics is not warranted. For a limited number of the relatively more promising processes, however, quantitative assessment has been made of appropriate process characteristics, to facilitate comparison with the gaseous diffusion process. This comparison is summarized and tabulated. For most of the processes, it has been possible either to estimate a separation factor or to set an upper bound for it. The number of stages or transfer units needed for a given degree of separation is inversely proportional to the separation factor tabulated, $\alpha - 1$. Of all the processes considered, only the Becker separation nozzle process has a higher separation factor than gaseous diffusion. Liquid-liquid chemical exchange between NOUF_6 and UF_6 has the highest separation factor of all the potentially reversible processes examined. The magnitude of the separation factor is at best only suggestive of the relative merit of a process. Indexes more directly related to the cost of separative work are the specific energy consumption, the specific uranium inventory, and the unit capital cost. (J.R.D.)

45690 DEUTERIUM SEPARATION PROCESS. Clarke, E. C. W.; Glew, D. N. (to Dow Chemical Co.). Canadian Patent 893,684. 22 Feb 1972.

A liquid mixture of hydrogen and deuterium sulfides is distilled at a pressure of from about 30 to about 1,300 psia and a temperature of from about -45 to about 100°C, and the gaseous deuterium sulfide-enriched overheads are collected. Refluxing or recycling steps can be incorporated in the process. The enriched distillate can subsequently be processed to yield deuterium oxide. (Canada)

45691 URANIUM ISOTOPE SEPARATION STUDIES IN FRANCE. Lebrun, C. (Societe de Construction d'une Usine de Separation Isotopique, Le Plessis, France); Mezin, M. *Energ. Nucl. (Paris)*; 14: No. 1, 5-16(1972). (In French).

The construction and operation of the Pierrelatte enrichment plant represents an important part of the program undertaken by France to meet the enriched uranium requirements of nuclear power reactors. An R and D program was completed on the well established gaseous diffusion technique. Results of basic economic studies indicate that this is the only method certain of fulfilling the enrichment requirements of the end of the present decade, which necessitate, in the western world, a construction rate capable of providing between 6 and 10 SWU/Y. These facts have led to the study of a gaseous diffusion production plant able to produce between 6 and 10 SWU/Y. The final plans will be ready in 1973. Other methods are also being reviewed; the most economic will be chosen for subsequent plants. (France)

45692 PROCESS AND DEVICE FOR THE SEPARATION OF MOLECULES OF DIFFERENT MASSES. Campargue, R. (to Commissariat a l'Energie Atomique). Canadian Patent 895,492. 14 Mar 1972

A process for the separation of molecules of different masses in a gaseous mixture is described. The mixture is circulated within a casing, into which an auxiliary gas jet is introduced at a supersonic velocity, creating a closed shock structure. The gas which is extracted from the interior of the shock structure contains a fraction of the mixture having a high concentration of light molecules, and the gas extracted from the outside of the shock structure has a low concentration of light molecules. The gaseous mixture to be processed can contain molecules of a different nature, isotopes of the same simple element (for example argon or hydrogen), or isotopic varieties of the same substance (for example hexafluorides of uranium-235 and uranium-238). (Canada)

45693 EVALUATE EUROPEAN SITING FOR NEXT ENRICHMENT PLANT. Grenon, Michel. *Power Eng.*; 76: No. 7, 35-7 (Jul 1972).

In Feb. 1972 an organization of six member nations, Belgium, France, Italy, Netherlands, UK, and West Germany, was formed to study the possibilities of building an international civilian gaseous diffusion plant in Europe. The experience of each country in separation work, the schedule for making decisions about the combined effort, and details on the design and performance of the French gaseous diffusion plant at Pierrelatte which has been operating for eight years are presented. (L.C.L.)

45694 ISOTOPE SEPARATION RATE IN HAEFFNER EFFECT. Sinha, Om P. (Simon Fraser Univ., Burnaby, B. C.). *Appl. Phys. Lett.*; 21: No. 1, 14-15(1 Jul 1972).

A simple theory is presented for Haeffner effect based on the ideas of Fiks. An exact formula derived for the isotope separation rate appears to be in good agreement with the experimental data on several systems. (auth)

45695 BRUCE AUXILIARY STEAM SUPPLY. Kolisnyk, Z. (Montreal Engineering Co. Ltd.); Spray, P. H. G. *Eng. J.* (Montreal); 55: No. 4, 22-6(Apr 1972).

An auxiliary steam plant for the Bruce Heavy Water Plant near Douglas Point, Ontario is nearly completed, and will produce 2.4 million lb/hr of steam at 180 lb/in² gage (360°F). The Bruce Heavy Water Plant, which will use the GS $\text{H}_2\text{O}/\text{H}_2\text{S}$ exchange process to produce 30 to 45% D_2O (followed by distillation), could be run on the entire steam output of Douglas Point Generating Station, but then 220 MW of electrical energy would be lost. After 1977, Bruce Nuclear Generating Station will supply most of the steam required, but the auxiliary supply will still be useful. The auxiliary plant will burn Bunker C residual oil supplied by unit train. Because of the danger of leakage of corrosive hydrogen sulphide into the three boilers, steam will be generated at 600 lb/in² gage, and the lower pressure steam generated from this primary steam in four reboilers. No power will be generated. The design is compared with that of a similar plant at Point Tupper, Nova Scotia. (auth) (Canada)

45696 ISOTOPIC ENRICHMENT OF HYDROGEN FOR HEAVY WATER PRODUCTION. Lambert, Irma; Ravoire, Jean; Roth, Etienne (to Commissariat a l'Energie Atomique). French Patent 2,079,708. 18 Oct 1971. Filed 10 Feb 1970. (In French).

A method is presented for the isotopic enrichment of hydrogen in heavy water production by isotopic exchange between the hydrogen and a primary or secondary aliphatic amine in the presence of a catalyst. A tertiary amine having an $\text{RR}'\text{R}''\text{N}$ form (R, R' and R'' = aliphatic radicals containing 1 to 4 carbon atoms) was added to the system. Trimethylamine and triethylamine were used in particular. (France)

45697 ULTRACENTRIFUGE ROTOR. Poubeau, Pierre (to Societe Nationale Industrielle Aerospatiale). French Patent 2,082,274. 15 Nov 1971. Filed 9 Mar 1970. (In French).

An ultracentrifuge for separating isotopic fluid elements is described that consists of a cylindrical vessel containing a radial vane system about a control axis. The radius of curvature of the sectors between consecutive vanes is greater than the outer radius of the vanes. Under the effect of the centrifugal rotation force, the radius of curvature of these sectors tends to approach the outer radius of the vanes. In this way the propeller-vane-wall system is given great rigidity. (France)

45698 VERSATILE HEAVY WATER. Swaroop, Janardhan (Bhabha Atomic Research Centre, Bombay). *Vaigyanik*; 3: No. 1, 8-12(Mar 1971). (In Hindi).

The importance of heavy water (D_2O) in the nuclear industry as a moderator to control the chain reaction in a reactor is discussed. Its physical and chemical properties, which distinguish it from ordinary water are outlined. Heavy water production

methods such as electrolysis of water, isotopic exchange process, partial distillation of water, low-temperature distillation, and hydrogen sulfide-ammonia dual temperature process are briefly described. Uses of heavy water in fields other than nuclear industry are enumerated. Experimental results of studies on the effects of heavy water on living organisms are also briefly described. (INIS)

Isotopic Power Sources

Refer also to abstract: 45677.

45699 (CEA-CONF-1998) RADIOISOTOPIC PACEMAKER DEVELOPMENTS IN FRANCE. Laurens, P.; Alais, M.; Berger, R.; Boucher, R.; Hixson, J. D.; Pivnicka, A. [nd]. 30p. (CONF-711008-5). Dep. NTIS (U. S. Sales Only).

From nineteenth conference on remote systems technology; Miami Beach, Fla. (17 Oct 1971).

A review of the French program for development of nuclear-powered cardiac pacemakers is presented. Developments in this program are described in a presentation supported by a series of slides. (J.R.D.)

45700 (COO-2155-4) BIOMEDICAL ENGINEERING SUPPORT. Progress Report. Kolff, W. J.; Donovan, F. M. Jr. (Utah Univ., Salt Lake City (USA). Dept. of Surgery). 15 Apr 1972. Contract AT(11-1)-2155. 42p. Dep. NTIS.

Results of research on power requirements for artificial hearts for humans, efficiencies of artificial hearts, and aspects of mating these hearts to nuclear power sources are reported. One animal experiment is described in which a Kwan-Gett heart made of silastic was implanted in a bovine calf. (J.R.D.)

45701 (DP-1300) SAVANNAH RIVER LABORATORY ^{60}Co POWER AND HEAT SOURCES. Quarterly Progress Report, January-March 1972. Angerman, C. L. (comp.) (Du Pont de Nemours (E. I.) and Co., Aiken, S. C. Savannah River Lab.). May 1972. Contract AT(07-2)-1. 19p. Dep. NTIS.

Galvanic corrosion on Inconel 600, nickel-plated cobalt metal, and couples between cobalt metal and superalloys during 61 days exposure to sea water; negligible corrosion occurred on the other superalloys and the ceramic cobalt compounds. None of the materials corroded significantly when exposed to bottom mud. Results of preliminary oxidation tests of iridium confirm data reported in the literature and indicate that a secondary capsule of an oxidation resistant material will be required to ensure containment of cobalt under accident conditions. Inactive superalloy capsules being heated at 850 to 1000°C are intact and performing satisfactorily after 39,400 h (4.5 yr). Fabrication of iridium capsules for ceramic cobalt compounds was begun; initial tests showed that sound welds could be made by TIG welding. The Hastelloy X being used to fabricate the fuel capsules and fuel pins for the WANL heat source exhibited satisfactory oxidation resistance during 5000-h tests at 1000°C; the Nickel 201 being used to fabricate the core also exhibited satisfactory resistance in 500-h tests at 1000 and 1125°C. The thermoelectric generator fueled with ^{60}Co continued to operate satisfactorily after 13 months with an electrical output of 58.5 W. (auth)

45702 (LA-4978) SPACE ELECTRIC POWER R AND D PROGRAM. Quarterly Report for the Period Ending April 30, 1972. (Los Alamos Scientific Lab., N. Mex.). Jun 1972. Contract W-7405-eng-36. 15p. Dep. NTIS.

Ceramic-metal multilayers containing Y_2O_3 as the electrical insulator were fabricated by arc-plasma-spraying Y_2O_3 onto metal collectors, followed by isostatic pressing of the trilayer assembly. Some properties of sprayed Y_2O_3 before and after isostatic pressing were determined. Arc-plasma-spraying of Y_2O_3 results in a dense, small-grained ceramic of low porosity after isostatic pressing at 10,000 psi and 1550°C. The electrical resistivity of a Nb-1% Zr/ Y_2O_3 trilayer was accurately measured, and long-term tests at temperature are in progress. The post-irradiation inspection and evaluations of alumina specimens irradiated by Oak Ridge National Laboratory in Capsule ORNL EB-2 are nearing completion. Volume expansions of Lucalox, Coors AD-995, and WESGO Al-995 were determined. The swelling of Lucalox was generally less than previously observed for similar material. The Coors AD-995 swelled considerably more than Lucalox and was very fragile; and the WESGO Al-995 swelled more than either of the other materials and more than previously observed for similar material. Results of leak tests indicated very low probability that cylindrical shells will retain vacuum integrity if radioinduced swelling exceeds 3 vol %. An isothermal irradiator test unit was constructed which duplicates cooling conditions in the Experimental Breeder Reactor (EBR-II) better than has been possible in the past. Initial tests of the prototype heat pipe for

Isothermal Irradiator K-5 were completed. Results of force-deflection measurements on ceramic bodies indicate decreases in moduli of elasticity and rupture of about a factor of two for irradiated alumina with a volume swelling of 3% compared to unirradiated material. No apparent decreases in these moduli due to irradiation were noted for Y_2O_3 for neutron fluences of $\sim 4.5 \times 10^{21}$ n/cm². Data on thermal diffusivity of one specimen of irradiated AD-995A alumina indicate no change due to irradiation at specimen temperatures above 500°C. A fracture sample of the tungsten cladding from General Electric Thermionic Test Assembly SCIP-513 was examined using a scanning-electron microscope. Numerous voids (0.2 to 2.5 μm dia) were detected in the cladding near the fuel/cladding interface. These voids decrease rapidly in size and number with increasing distance from this interface; they occur predominantly on grain-boundary surfaces. The locations of the voids could not be correlated with the locations of fission products detected at the fuel/cladding interface. (auth)

45703 (ORNL-4796) ORNL ISOTOPIC POWER FUELS QUARTERLY REPORT FOR PERIOD ENDING MARCH 31, 1972. Lamb, Eugene; Donnelly, R. G. (Oak Ridge National Lab., Tenn.). Jun 1972. Contract W-7405-eng-26. 38p. Dep. NTIS.

A vacuum hot-press for pressing pellets up to 1-in. diameter was assembled and tested. A process and the necessary hot-cell equipment to make $\sim 250\text{-}\mu\text{m}$ -dia $^{244}\text{Cm}_2\text{O}_3$ microparticles have been developed. These particles are strong enough to be blended with metal powder to form cermet. The curium source test is being disassembled and the equipment salvaged. Causes of the transfer of curium fuel from the pellet channels to the upper interior sections of the capsule have been analyzed. Four samples of the curium fuel were taken during the disassembly for spark source mass spectrometry, x-ray diffraction, and helium release analyses. Two cylindrical SrTiO₃ pellets and four spherical Gd₂O₃ pellets were used to test the operation of the impact test unit. Three sabot heaters have been designed to heat the specimens to their expected operating temperature before impacting them. Construction of the laser system for velocity measurement is expected to be completed in April. The Pt-Rh-W alloys were found to be relatively insensitive to atmospheric contaminants such as those found in radioisotope thermoelectric generators. Hemispheres of Pt-26% Rh-8% W and Pt-30% Rh-8% W-0.25% Hf-0.1% Ti have been successfully fabricated. Iridium has been zone refined into single-crystal bar, which can be cold swaged 40% without cracking. Much higher oxygen levels can be tolerated by T-111 without embrittlement at temperatures higher than present Pioneer operating temperatures. T-111 embrittled by oxygen contaminants at 825°C can be ductilized by a heat pulse equivalent to that received by a Pioneer heat source upon reentry into the earth's atmosphere. Molybdenum alloys and coatings on T-111 continue to show relative insensitivity to contaminating environments. (auth)

45704 NEUTRON RADIATION CHARACTERISTICS OF PLUTONIUM DIOXIDE FUEL. Taherzadeh, Mojtaba (Jet Propulsion Lab, Pasadena, Calif.); Gingo, Peter J. Nucl. Technol.; 15: No. 3, 396-410 (Sep 1972).

The major sources of neutrons from plutonium dioxide nuclear fuel are considered in detail. These sources include spontaneous fission of several of the Pu isotopes, (α, n) reactions with low Z impurities in the fuel, and (α, n) reactions with ^{18}O . For spontaneous fission neutrons a value of $(1.95 \pm .07) \times 10^3$ n/sec/gm PuO₂ is used. The neutron yield from (α, n) reactions with oxygen is calculated by integrating the reaction rate equation over all α -particle energies and all center of mass angles. The results indicate a neutron emission rate of $(1.14 \pm .26) \times 10^4$ n/sec/gm PuO₂. The neutron yield from (α, n) reactions with low Z impurities in the fuel is presented in tabular form for one part per million of each impurity. The total neutron yield due to the combined effects of all the impurities depends upon the fractional weight concentration of each impurity. The total neutron flux emitted from a particular fuel geometry is estimated by adding the neutron yield due to the induced fission to the other neutron sources. 22 references. (auth)

45705 RESULTS OF AN ENQUIRY INTO THE SOURCE REQUIREMENTS OF RADIOISOTOPE GENERATORS IN THE ENEA MEMBER COUNTRIES FOR 1971 TO 1975. Boutaine, M. (CEN de Saclay, Gif-sur-Yvette, France). Bull. Inform. Generateurs Isotop.; No. 11, 3-7 (Dec 1971). (In French).

This study indicated that the largest and most probable demands will be for sealed ^{238}Pu and ^{90}Sr sources, and that the demands envisaged in 1971 are not very realistic when the current development in techniques is taken into consideration. The results obtained were analyzed. The following sources are considered: sealed sources for use in thermoelectric, ionic or dynamic conversion; α or β emitting sources for use in radio-voltaic conversion; and radioelements in the crude form for use in energy conversion. (France)

45706 FRENCH PROGRAM ON THE PRODUCTION AND USES OF ^{238}Pu . Berger, R.; Boucher, R.; Derian, J. C. (CEN, Fontenay-aux-Roses, France). Bull. Inform. Generateurs Isotop.; No. 11, 8-36 (Dec 1971). (In French).

The studies undertaken in France on the production of ^{238}Pu are presented. The preparation of ^{237}Np and ^{241}Am based targets, the irradiation conditions and the chemical processing are described. Studies relating to the medical applications of ^{238}Pu are reviewed, in particular those concerning intracorporeal cardiac pacemakers and artificial intracorporeal hearts. The Delphi inquiry demonstrates that French cardiologists and specialists consider the artificial heart as a realistic objective, and expect that more widespread applications of such techniques will take place during the coming decade. (France)

45707 PROCESS OF FORMING AN ISOTOPIC HEAT SOURCE. Sump, Kenneth R.; Robinson, Ramon K.; Howard, Boyd D.; Drumheller, Kirk. (to United States Atomic Energy Commission). U. S. Patent 3,666,846. 30 May 1972. Filed 11 Apr 1969.

A process of encapsulating an alpha-emitting isotope to form an isotopic heat source is described. Microspheres are prepared, each containing its own helium gas plenum, an alpha-emitting isotope, and containment cladding. These microspheres are sealed in a capsule with or without a metal matrix. (Official Gazette)

45708 RADIOISOTOPIC FUEL CAPSULE. Selle, James E.; Kokenge, Bernard R. (to United States Atomic Energy Commission). U. S. Patent 3,659,107. 25 Apr 1972. Filed 29 Jul 1970.

The design of a radioisotopic fuel capsule for use in thermoelectric or thermionic generators which utilizes substoichiometric plutonium dioxide enclosed within a refractory container is described. (Official Gazette)

Radiation Sources

Refer also to abstracts: 45432, 45446-45448, 45467, 45471, 45473, 45555, 45942, and 46008.

45709 (INIS-mf-256) RADIOISOTOPES, RADIOACTIVE PHARMACEUTICALS, RADIOACTIVE CHEMICALS, RADIOACTIVE SOURCES, SERVICES. (Institut for Atomenergi, Kjeller (Norway)). 1972. 40p. INIS.

Specifications, prices and delivery times for eighty-eight radioactive pharmaceuticals and chemicals produced by Institute for Atomic Energy are given. ^{60}Co , ^{192}Ir and ^{198}Au sources for laboratory and industrial use and a variety of beta and gamma sources for training and demonstration purposes are described and prices given. Irradiation services are also available and containers, flux, etc. are specified. Standardized sources and solutions and absolutely standardized solutions are also available. The Institute also offers services in activation analysis and in the industrial applications of radioisotopes. (auth)

45710 USE OF RADIONUCLIDES FOR MEASUREMENTS IN MECHANICAL ENGINEERING, A FIELD OF WORK IN THE LABORATORY FOR ISOTOPE TECHNOLOGY. Gerve, A. (Gesellschaft fuer Kernforschung, Karlsruhe, Ger.). Kerntechnik; 14: No. 5, 201-4 (May 1972). (In German and English).

On Jan. 1, 1971, the Laboratory for Isotope Technology was extended by the creation of the Department for Physics in Mechanical Engineering. The work of this new Department is subdivided into basic work for the further development of isotope technology, application experiments for research problems in the field of mechanical engineering, and work on industrial contracts which can now also be carried out based on requests of the customers with the aid of a mobile measuring unit. The work is concerned mainly with activation techniques, wear investigations, and the use of computers for automating experiment control. (auth)

45711 IMPORTANT WEAR MEASURING METHODS USING RADIONUCLIDES. Gerve, A. (Gesellschaft fuer Kernforschung, Karlsruhe, Ger.). Kerntechnik; 14: No. 5, 204-9 (May 1972). (In German and English).

Thin layer activation of machine parts, i.e., irradiation with charged particles, makes it possible to carry out wear measurements without any hazard even in ordinary industrial operations. The thin layer difference method of measuring wear is based on measuring the reduction of the activity of wear parts activated by the thin layer method. The achievable accuracy is better than 0.5 μm abrasion. The continuous flow method of measuring wear is based on measuring the total wear by measuring the time dependence of the variation in the activity concentration of abraded particles in the lubricating oil. The achievable accuracy is of the

order of 10^{-8} g/h loss of material, even when several wear components are measured simultaneously. (auth)

45712 INVESTIGATION OF FREON FIRE-EXTINGUISHING SYSTEMS WITH A NUCLEONIC GAGE. Notea, A.; Segal, Y. (Technion-Israel Inst. of Tech., Haifa). Mater. Eval.; 30: No. 7, 153-6 (Jul 1972).

The dynamic properties of the fire-extinguishing system in an aircraft were examined using a three-channel gamma attenuation gage connected to each outlet of the system. The extinguishing agent was Freon 13B-1 (CBr F₃), and the tube internal diameter was 0.774 cm. The gamma source, ^{241}Am , was selected by optimization of the standard deviation due to statistical counting fluctuations. The collimator arrangement allowed checking of 65% of the tube cross section. The results of mass changes with time during discharge of the system are shown. The changes of characteristic parameters—discharge time and full width half maximum time—are presented for different nozzle configurations. (auth)

45713 INVESTIGATION INTO THE APPLICATION OF POST-CUTTING CHIP ACTIVATION ANALYSIS TO THE STUDY OF TOOL WEAR. Prasad, Kadambi Narasimha. University Park, Pa.; Pennsylvania State Univ. (1971). 96p. University Microfilms Order No. 72-19,365.

Thesis.

Post-cutting chip activation analysis is a technique for measurement of tool wear rate that is analogous to the technique that employs radioactive tools. The essential difference between the two techniques is that in the former neutron activation is performed after the machining operation, using samples of chips, whereas in the latter the tool is activated before machining. The main advantage of the chip activation technique over activation of the tool is that the levels of radioactivity to be handled are lower by about a factor of 10^4 . Gamma spectrometry is employed to identify and determine the tool material on the chips after activation. Two alternative approaches were examined in the application of this technique: one approach used the element tungsten as the neutron activatable tracer, tungsten being an inherent constituent of the tools that were used; the second approach, examined for use where the first was not possible, employed a tracer inserted into the tool material specifically for the purpose of wear measurement. The tools used were made of the T-1 type of high speed steel for both approaches. Because of its favorable nuclear properties, europium was chosen as the neutron activatable tracer for the second approach. The titanium alloy Ti-6Al-4V was chosen as the work material for the investigation into the chip activation technique due to the relatively low level of trace tungsten concentration in the material. The precision of the activation analysis method to determine wear rates was found to be less than 3% at the 95% confidence level. The results of the tests showed that a correlation could be obtained between the break-in wear rate and tool life that could predict tool life to within the same margin of error that was obtained from the Taylor plot (cutting speed versus tool life) for the conditions used. (Diss. Abstr. Int., B)

Design and Fabrication

45714 (CONF-720902-2) FABRICATION OF ^{252}Cf NEUTRON SOURCES AT OAK RIDGE NATIONAL LABORATORY. Van Cleve, J. E. Jr.; Knauer, J. B.; Williams, L. C.; Bigelow, J. E. (Oak Ridge National Lab., Tenn.). [nd]. 5p. Dep. NTIS.

From National topical meeting on the applications of californium-252; Austin, TX. (11 Sep 1972).

The method of loading the californium in the aluminum liner and source encapsulation in the primary container are described. Identification numbers inscribed on the sources are prefixed with NSS or NSD to distinguish singly and doubly contained sources. The qualification of the source involves both a helium leak test and a x-radiograph. In helium leak testing, the leak rate of the source is compared with a standard leak. The ^{252}Cf content of each source is determined by measuring the fast neutrons emitted. This method is quick, accurate, and nondestructive. The counting rate is corrected for geometry, background, scattering due to miscellaneous objects, and scattering due to the source containers. The system has a reproducibility of $\pm 1\%$ in comparing sources with a standard source assayed during the same test period. The source is decontaminated in a water-shielded, portable decontamination facility. The clean source is shipped to the customer in a specifically designed 12-ton carrier. (auth)

45715 DETERMINATION OF ISODOSE CURVES OF SEALED γ -SOURCES BY CALCULATION AND FILM DOSIMETRY. Pavlicsek, I.; Csueres, M.; Harsanyi, Gy. (MTA Isotop Intezet, Budapest). Izotopteknika; 15: No. 1, 25-36 (Apr 1972). (In Hungarian).

The calculation of source distribution from dose rate distribution is discussed from the point of view of source construction. Isodose curves for gynecological sources in body-equivalent media are shown. (auth)

Industrial Uses

Refer also to abstracts: 45456 and 45457.

45716 (BFL-2/1972) INDUSTRIAL PROCESSING OF IRRADIATED POTATOES. Penner, H.; Gruenewald, Th.; Rumpf, G.; Gehse, M.; Wolf, W. (Bundesforschungsanstalt fuer Lebensmittelrischhaltung, Karlsruhe (West Germany)). Mar 1972. 51p. (In German). Dep. NTIS (U. S. Sales Only).

In the framework of a joint experiment organized by the Bureau Eurisotop, Brussels, in several countries of the Common Market 10 tons of potatoes (varieties Bintje and Saturna) were treated with ionizing radiation in order to inhibit sprouting. The doses were 8.5 and 15 krad. After storage of 6 and 8 months (i.e., 7 and 9 months after harvest) these potatoes were processed into potato chips and dried potatoes. Seven tons of the same variety and origin were sprout inhibited by a mixture of Chlorpropham (CIPC) and Propam (IPC) or only IPC and stored and processed under the same conditions. Peelability showed no difference between irradiated and chemically-treated potatoes. Some of the radiation processed potatoes showed a grey discoloration. Due to a higher content of reducing sugars chips from irradiated potatoes were darker than controls of the same moisture content. After 6 months of storage the irradiated potatoes showed a higher amount of rot than the chemically-treated ones. (auth)

45717 RADIOISOTOPIC THICKNESS MEASUREMENTS IN THE MANUFACTURE OF WIRE. Plesch, Rolf (Siemens Aktiengesellschaft, Karlsruhe, Ger.). Siemens Rev.; 39: No. 5, 223-6 (May 1972).

The use of radioactive isotopes enables wire thicknesses to be measured without physical contact during the production process and in this way the manufacturer is no longer forced to test random specimens of the end product. The method can be used to avoid the production flow and lowers rejection rate. A further advantage of this new method is a considerable saving on the materials sector, for the set point for the wire thickness can be brought nearer the minimum tolerance limit. (auth)

45718 RADIATION POLYMERIZATION OF IMPREGNATED POROUS MATERIAL. Hills, P. R.; McGahan, D. J. (Atomic Energy Research Establishment, Harwell, Eng.). Nucleus (Karachi) 8: No. 4, 33-42 (1971).

The polymerization of impregnated porous material through the application of radiation is examined. (auth)

45719 FLUID IRRADIATING APPARATUS USING GAMMA RAYS AND HELICAL PASSAGEWAYS. Woodbridge, D. D.;

Mann, L. A. (to Energy Systems, Inc.). Canadian Patent 897,930. 18 Apr 1972.

A gamma irradiator for the sterilization and treatment of liquids such as sewage and industrial wastes is described. The irradiator has a hollow center chamber and a spiralling input chamber. The liquid spirals around the center chamber and flows into it, passes through and into an elongated core member located in the center chamber, out a pipe which is helically wrapped around the hollow member, and exits from the irradiator. Gamma radiation sources are located in the center chamber and around the elongated core member and may be placed in two parallel circles, one on each side of the helical exit pipe. Baffles may be located throughout the system to increase the turbulence, improving uniformity of radiation and preventing settling of solids. The irradiator is enclosed in shielding sides; inlets and outlets are curved or bent, and the cover for the shielding chamber is stepped in the usual manner. (Canada)

45720 INSTALLATION OF RADIOACTIVE SOURCES IN THE NO. 4 BLAST FURNACE AT CIA, FERRO BRASILEIRO FOR STUDYING LINING WEAR. Marri, Luisio. Metalurgia (Sao Paulo); 27: No. 163, 443-7 (Jun 1971). (In Portuguese).

Thirty ^{60}Co sources with individual activities ranging from 0.5 to 55.0 mCi and a total installed activity of 391.0 mCi were installed in the lining while the furnace was down for repairs. The sources were placed at depths of $\frac{1}{3}$, $\frac{1}{2}$, and $\frac{2}{3}$ the distance from the exposed inside surface of the lining on the assumption that $\frac{1}{3}$ of the thickness would be lost during the first 6 to 8 months of the new campaign, 50% by the end of $\sim 2\frac{1}{2}$ years and the remainder in ~ 6 years. Periodic measurement of the activity remaining at the various locations will determine whether these predictions are valid. Details are given concerning the locations of the sources in the lining, costs of installation and the program of measurement of radioactivity. Calculated figures are presented on the possible activity in a piece of steel pipe made from a heat in which one of the most active sources (55.0 mCi) may have been entrained and the absence of a safety problem attributable to the use of such a pipe in a main supplying water for human consumption. (Metals Abstr.)

45721 RADIOISOTOPES AND TURBINE FLOW MEASUREMENTS. Schuster, Jack S.; Hansen, Robert L. (Bureau of Reclamation, Denver). J. Power Div., Amer. Soc. Civil Eng.; 98: PO-1, 1-9 (Jun 1972).

Results are reported on research directed to measuring discharge through high-head turbines and pumps using radioisotope tracers. A series of measurements was made to compare the relative accuracy of methods involving radioisotope—velocity measurements, integrated sample measurements, dilution measurements, and total-count measurements. Results of these investigations did not show a major advantage to one particular method. Each method has its advantages and disadvantages and can be considered for a particular situation. (J.R.D.)

LIFE SCIENCES

Radiation Effects on Biochemicals (External Source)

Refer also to abstract: 45349.

45722 (COO-3232-1) MOLECULAR BASIS FOR THE MUTAGENIC AND LETHAL EFFECTS OF ULTRAVIOLET IRRADIATION. Progress Report, June 16, 1971-May 31, 1972. Grossman, Lawrence (Brandeis Univ., Waltham, Mass.). Contract AT(11-1)-3232. 6p. Dep. NTIS.

Research was conducted on elucidation of the enzyme mechanism by which uv-damaged DNA is repaired. The following steps in the excision repair cycle are discussed: incision, catalyzed by an endonuclease from *Micrococcus luteus*; reaction with formaldehyde; excision, catalyzed by an exonuclease; and reinsertion mechanisms involving DNA polymerase. (H.L.W.)

45723 RADIATION EFFECTS ON PLASMA ELECTRON-SPIN-RESONANCE (ESR) SPECTRA OF CANCER PATIENTS. Swartz, Harold M.; Wiesner, James (Medical Coll. of Wisconsin, Milwaukee). Radiology; 104: No. 1, 209-10 (Jul 1972).

In studies of blood plasma of cancer patients, the average magnitudes of several electron-spin-resonance peaks were seen to be significantly increased, apparently as the result of increased amounts of paramagnetic trace elements. Thirty patients studied during radiation therapy showed a trend back toward control levels, with the greatest response found in patients with squamous-cell carcinoma. (auth)

45724 EFFECT OF GAMMA IRRADIATION ON FUERTE AVOCADO FRUITS. Kamali, A. R.; Maxie, E. C.; Rae, H. L. (Univ. of California, Davis). HortScience; 7: No. 2, 125-6 (Apr 1972).

Gamma irradiation up to 100 Krad caused no immediate visual damage to fruits of avocado (*Persea americana* Mill.). Twenty Krad or more stimulated a climacteric-like rise in respiratory rate. Rates of ethylene (C_2H_4) production were related directly to dose up to 40 Krad. Ten and 40 Krad delayed the climacteric peak for 3 days. Softening was delayed by 10 Krad and hastened by 40 and 100 Krad. At 100 Krad, ripening was lacking and the fruit severely injured. No benefit remained from any dose after 3 weeks of storage. Storage reduced the time required for ripening by 2 days, and enhanced the development of injury symptoms and pigment degradation. (auth)

45725 EFFECT OF WATER ON THE RADIOSENSITIVITY OF PLASMATIC FATTY ACIDS. Dousset, N.; Douste-Blazy, L. (Univ., Toulouse). Compt. Rend. Soc. Biol.; 165: No. 6, 1462-5 (1971). (In French).

Water prevents the appearance of liquid peroxides in plasma subjected to x radiation (20,000 roentgens). Lyophilized plasma is more radiosensitive than liquid plasma; the irradiation induces the formation of lipidic peroxides, principally at the expense of the linoleic and oleic acids without modifying the free fatty acids. (France)

45726 POSSIBILITIES IN STUDYING THE MECHANISM OF RADIOSENSITIVITY OF AND RADIATION DAMAGE TO ALBUMINES AND NUCLEIC ACIDS. Damjanovich, S. (Debreceni Orvostudományi Egyetem Biofizikai Intézete, Hungary). Izotop-technika; 15: No. 3-4, 143-53 (1972). (In Hungarian).

A survey of literature and results of experiments prove that a single irradiation does not alter considerably the conformation of enzyme, or lower tolerance to heat. The effect can be observed in the regulation and, to a smaller extent, in the activity of the enzyme. (auth)

45727 IN VITRO REPAIR OF UV-INACTIVATED TRANSFORMING DNA FROM *BACILLUS SUBTILIS*. Heijneker, H. L.; Pannekoek, H.; Oosterbaan, R. A.; (and others) (Medical Biological Lab., TNO, Rijswijk, Netherlands. Univ., Groningen, Netherlands). Genen Phaenen; 15: No. 1, 31-2 (Jun 1972). (In Dutch).

Three specific enzymes, uv-specific endonuclease, DNA polymerase, and DNA ligase, were found to be involved in the in vitro repair of uv-induced damage in DNA. (auth)

45728 ULTRAVIOLET INACTIVATION AND MISCODING OF IRRADIATED R17-RNA IN VITRO. Remsen, Joyce F.; Cerutti, Peter A. (Univ. of Florida, Gainesville). Biochem. Biophys. Res. Commun.; 48: No. 2, 430-6 (25 Jul 1972).

Ultraviolet irradiation of the single-stranded Coliphage R17 leads to inactivation of the in vitro messenger activity of its RNA.

Uridine-photohydration is the major reaction occurring in the RNA under these conditions. At a residual activity of the synthetase gene of 37%, there are 11.2 uridine-photohydrates present in the total R17 genome, while only 5.3 photohydrates are necessary to decrease the translation of the coat gene to the same value. De novo incorporation of histidine into the coat protein is observed with R17-RNA from uv-irradiated phage, indicating that uridine-photohydrates can miscode as cytidine in an in vitro translation system. (auth)

45729 RESPIRATORY QUOTIENTS OF IRRADIATED FRUITS. Ahmed, E. M.; Dennison, R. A.; Green, G. F. (Univ. of Florida, Gainesville). J. Amer. Soc. Hort. Sci.; 97: No. 3, 373-6 (1972).

An automatic sampling system for monitoring CO_2 evolution and O_2 uptake of irradiated fruits under a continuous air flow was used. All fruits showed increased respiratory activities immediately following irradiation. The respiratory activities of irradiated tomatoes and strawberries decreased within 24 hr following irradiation, while those of oranges and grapefruit remained at elevated levels. The respiratory quotients (RQ) of irradiated citrus fruits were higher than those for control fruits, while those of tomatoes and strawberries were not. All RQ values were less than unity. (E.R.B.)

45730 CHEMICAL AND PHYSICAL CHANGES OF DEXTRAN 4000 IRRADIATED WITH VARIOUS DOSES OF γ -RADIATION. Derlikowski, J.; Szyszko, E.; Zagorski, Z. P. (Inst. for Drug Research and Control, Warsaw. Inst. of Nuclear Research, Warsaw). Diss. Phar. Pharmacol.; 23: 253-7 (1971).

Preparation "Dextran" of mean molecular weight 40000 (dry powder) decomposes under the influence of γ rays. The preparation cannot be sterilized by γ irradiation. (NSA of Poland)

45731 SEMI-IN VITRO REPAIR OF RADIATION-INDUCED DAMAGE IN TRANSFORMING DNA OF *BACILLUS SUBTILIS*. Noguti, T.; Kada, T. (National Inst. of Genetics, Shizuoka, Japan). J. Mol. Biol.; 67: No. 3, 507-12 (28 Jun 1972).

Toluene-treatment at 0°C was effective in making *Bacillus subtilis* cells accessible to substrates and co-factors of enzymes involved in the repair of damage induced by ionizing radiation. Transforming activity in the lysate of γ -irradiated (30 Krad) and toluene-treated bacteria increased remarkably in the presence of four deoxynucleoside triphosphates, nicotinamide adenine dinucleotide, and Mg^{2+} at 37°C. Under similar conditions, γ -ray induced DNA single-strand breaks were rejoined to a significant extent. Involvement of DNA-polymerase and DNA-ligase functions was strongly suggested in the repair of DNA damage induced by ionizing radiation. (auth)

45732 BIOCHEMICAL MECHANISMS IMPLIED IN THE SENSITIZATION OF LIVING ORGANISMS BY CHEMICALS UNDER THE EFFECT OF RADIATIONS AND CYTOSTATICS. III. THE EFFECT OF HMB, IAA, AND CAFFEINE UPON THE DNAase ACTIVITY IN THE LIVER OF NORMAL OR IRRADIATED RATS. Furnica, M.; Ganea, A. Rev. Roum. Biochim.; 8: No. 3, 211-18 (1971). (In French).

p-Hydroxymercuribenzoic (HMB) acid in 15 mg/kg body weight and iodoacetamide (IAA) in 15 mg/kg body weight were given to normal or 800 R irradiated rats and the activity of liver DNAase was followed. Three days after the treatment, DNAase inhibition in normal animals and its relative activation (compared to the irradiated controls) in the irradiated animals were reported. DNAase activation in the irradiated animals and radiosensitivity could be involved in the inhibition and irradiation damaged DNA restoration. Caffeine in doses to 100 mg/kg body weight did not alter liver DNAase activity. (Rom. Sci. Abstr.)

45733 EFFECT OF X-RAY IRRADIATION ON GLUTATHIONE PEROXIDASE. Olinescu, R. Rev. Roum. Biochim.; 8: No. 3, 231-8 (1971).

X-ray effect upon glutathione peroxidase extracted from swine blood and 2500 times purified was investigated. Irradiation was conducted in atmospheric air at 0°C with a 2 KR/min dose. Hydroperoxidase used as substrate protected the regulating and catalytic functions of glutathione peroxidase. The regulating function of GSH peroxidase is more radiosensitive than its catalytic one. The regulating function is affected at different places. It being however, in the neighborhood of the catalytic one. The stimulation of peroxidase activity governed by NADPH is counteracted upon irradiation contamination. GSH peroxidase in the swine liver, even less purified, is not as radiosensitive as the blood enzyme, where either the catalytic or allosteric functions are concerned. (Rom. Sci. Abstr.)

Radiation Effects on Cells (External Source)

Refer also to abstracts: 45611 and 45612.

45734 EFFECT OF TAURINE AS A PROTECTOR OF HUMAN MARROW CELLS AGAINST RADIATION. Marchionni, R.; Bersi, M. (Verbania Hospital, Italy). *Arch. Sci. Med.*; 126: No. 1, 20-2(1969). (In Italian).

Chromosome fragmentation counts on irradiated human marrow cells were used to determine the radioprotective effect of taurine. The results were unfavorable in the sense that the number of fragmentations observed in cultures to which taurine had been added was only slightly less than that in the control cultures. (auth)

45735 QUANTITATIVE AND QUALITATIVE ASPECTS OF RADIOBIOLOGY AND THEIR SIGNIFICANCE IN RADIATION THERAPY. Wideroe, R. (Brown, Boveri and Co. Ltd., Baden, Switzerland). *Acta Radiol., Ther., Phys., Biol.*; 10: No. 6, 605-24(Dec 1971).

The survival of mammalian cells in proliferating cell cultures exposed to various types of radiation is described and analyzed mathematically. Such investigations neglect the influence of milieu and different qualitative cell reactions. The results should therefore not be transferred indiscriminately to clinical treatment. However, general directions for treatment schemes using various types of radiation can be determined and cell parameters chosen that correspond well to present clinical experience. (auth) (Sweden)

45736 USE OF A MODEL OF TWO CLASSES OF TRYPTOPHAN RESIDUES FOR INTERPRETING THE RESULTS OF ULTRAVIOLET FLUORESCENCE OF CELLS EXPOSED TO X-RAYS. Ostashevskii, I. Ya.; Yagunov, A. S. (Central Scientific-Research Roentgeno-Radiological Inst., Leningrad). *Biophysics (USSR) (Engl. Transl.)*; 16: No. 5, 834-40(1971).

Translated from *Biofizika*; 16: No. 5, 802-7(1971).

The ultraviolet fluorescence spectra of x-irradiated (500 rads) and unirradiated Ehrlich ascites carcinoma cells were studied. An increase in the intensity (I_{max}), and a longwave shift was demonstrated in the fluorescence spectrum in irradiated cells. Analysis of this experimental fact and the reciprocal relation between the slope of the dose relation and the value of the initial intensity within the framework of the model of two classes of tryptophan gave preference to a mechanism whereby there is increase in the quantum yield of the red tryptophans with no change in the fractions of the residues of each. (auth)

45737 INDUCTION OF UNSCHEDULED DNA SYNTHESIS BY HALF-NUCLEUS IRRADIATION OF HeLa CELLS WITH AN UV-MICROBEAM IN COMBINATION WITH HYDROXYUREA TREATMENT. Yatani, Ryuichi (Mie Prefectural Univ., Tsu, Japan). *Mie Med. J.*; 20: No. 2, 93-104(Sep 1970).

Radiobiological properties of half-nucleus irradiation in HeLa cells treated with hydroxyurea were observed. The rates of unscheduled DNA synthesis during different phases of the cell cycle of HeLa cells were calculated from autoradiographic results obtained by technique of half-nucleus irradiation with an uv-microbeam in combination with hydroxyurea treatment. It was suggested that the S-phase was more sensitive to uv-irradiation and subsequently to the induction of unscheduled DNA synthesis than the other phases. (Japan)

45738 EFFECT OF X-IRRADIATION ON MACROPHAGES AND TUMOR CELLS CO-CULTIVATED IN VITRO. Vorbrott, A.; Grabska, A.; Steplewski, Z. (Inst. of Oncology, Gliwice, Poland). *Folia Histochem. Cytochem. (Krakow)*; 9: 153-8 (1971).

The effect of x-ray (1300 R) on the phagocytic activity of macrophages co-cultivated with tumor cells (ELD and C3HM) was studied. It was found that after irradiation phagocytic activity of macrophages was stimulated, as evidenced by their enlargement, increased number and size of lysosomes, engulfing, and phagocytosis of x-irradiated tumor cells. The significance of these observations is discussed in relation to the mechanisms involved in the phagocytosis of irradiated tumor cells. (NSA of Poland)

45739 FACTORS AFFECTING RADIATION SENSITIVITY. Quintiliani, Marcello. *Ann. Ist. Super. Sanita*; 7: 647-60 (1971).

An attempt is made to organize the available data on the lethal effects of radiation on single cells into a coherent pattern. A large body of experimental evidence demonstrates that radiosensitivity depends on a variety of factors, ranging from the stage in the cell cycle to the oxygen pressure and from the degree of hydration to the presence of a variety of added substances. Some of these intrinsic and environmental modifications are discussed. (E.R.B.)

45740 SPONTANEOUS AND X-RAY INDUCED MUTATION IN CULTURED, DIPLOID HUMAN CELLS. Albertini, Richard Joseph, Madison, Wis.; Univ. of Wisconsin (1972). 248p. University Microfilms Order No. 72-15,335.

Thesis.

A system was developed to detect somatic cell mutation to 8 azaguanine resistance in normal, cultured, diploid human fibroblasts. Hypoxanthine-guanine phosphoribosyltransferase (HG-PRT) deficient, 8 azaguanine (8AG) resistant fibroblasts from boys with the Lesch-Nyhan Syndrome served as prototype mutant cells. Such cells are known to be mutant at an X-linked genetic locus—the hpt locus. Both spontaneous and x-ray induced mutation were studied. Preliminary x-ray studies are described that determined clonal survival of cultured human fibroblasts as a function of x-ray dose. 250 R allowed 20% survival. Numerous reconstruction experiments assessed efficiency of recovery of prototype mutant cells from majority populations of normal cells. Recovery was a function of both cell density and 8AG concentration, with optimum recovery achieved at an initial inoculum of 2×10^4 cells per 60 mm plastic dish selected at a concentration of 8×10^{-6} M 8AG. Efficiency of recovery was between 60 and 90%. Internal reconstruction experiments were included in all subsequent mutagenesis experiments. Attempts to purify normal cell populations of pre-existing 8AG resistant mutant cells by HAT (hypoxanthine, aminopterin, thymidine) reverse selection against the mutants failed. Such pre-existing mutants were therefore carried into mutagenesis experiments where they biased estimations of spontaneous mutant clone frequencies. Five mutagenesis experiments are reported. Irradiated and unirradiated cell populations were studied. In two experiments, one dose of x rays was administered (250 R) while in the remainder two or three doses were given. Spontaneous mutation rate was estimated to be 4.5×10^{-6} per cell generation which was an overestimation because of pre-existing mutants. Spontaneous mutant clone frequency was subtracted in irradiated populations to determine induced mutant clone frequency. Induction of mutation by x rays was shown to be dose-dependent with increases occurring in a non-linear cumulative fashion. Induced mutation rates were estimated at 1.13×10^{-6} per R per locus for 75 R, 8.5×10^{-8} per R per locus for 125 R, 6.87×10^{-8} per R per locus for 150 R, and 2.6×10^{-7} per R per locus for 250 R. Study of derived mutant substrains showed these to be no more x-ray resistant than parental strains. HG-PRT activity was determined to be deficient in over 80% of them, although not as deficient as in L-N prototype mutant cells. All derived substrains studied could grow in both 8AG and in HAT medium. The triad of HG-PRT deficiency, 8AG resistance, and HAT growability described a class of mutant human fibroblasts heretofore unrecognized. Studies are also described that attempt to detect 8AG-resistant lymphocytes in human blood samples. Detection can be made with a sensitivity approaching 1%. A minority population of 8AG resistant cells comprising 5 to 10% of the total was found in young females heterozygous for the Lesch-Nyhan mutation. Since such cells are not detected in populations of lymphocytes from normals, the system may have use as an adjunctive heterozygous detection test. Other applications are discussed. (Diss. Abstr. Int., B)

45741 REGULATION OF GRANULOCYTE AND MACROPHAGE FORMATION IN DIFFUSION CHAMBER CULTURES OF MOUSE HAEMATOPOIETIC CELLS. Breivik, H. (Norwegian Defence Research Establishment, Kjeller); Benestad, H. Breien. *Exp. Cell Res.*; 70: 340-8(Feb 1972).

Bone marrow and blood cells from mice were cultured in peritoneal diffusion chambers in homologous recipients. Granulocytes and macrophages were formed in the chambers. About a quarter of the blood lymphocytes were recovered after the culture period, whilst bone marrow lymphocytes almost disappeared. The number of granulocytes harvested was linearly related to the number of cells implanted, indicating that growth regulatory cell interactions did not take place in the chambers. Regenerating bone marrow produced twice as many granulocytes per progenitor cell as did normal marrow. The cell yield was also increased when irradiated or cyclophosphamide pretreated chamber hosts were employed. Thus properties of the progenitor cells as well as humoral factors from the hosts did influence chamber granulopoiesis. In contrast to the growth results for granulocytes, the formation of macrophages showed a relative decrease when the cell inocula were increased. Pretreatment of the host mice with irradiation or with cyclophosphamide resulted in increased macrophagopoiesis in bone marrow cultures. (auth)

45742 VARIOUS LEVELS OF DNA REPAIR SYNTHESIS IN XERODERMA PIGMENTOSUM CELLS EXPOSED TO THE CARCINOGENS N-HYDROXY AND N-ACETOXY-2-ACETYL-AMINOFLUORENE. Stich, H. F.; San, R. H. C.; Miller, J. A.; Miller, E. C. (British Columbia Univ., Vancouver). *Nature (London) New Biol.*; 238: No. 79, 9-10(5 Jul 1972).

A study was made, by ^3H TdR incorporation methods, of the levels of DNA repair synthesis in Xeroderma pigmentosum (XP) cells of 5 patients exposed to the carcinogenic and mutagenic compounds N-acetoxy- or N-hydroxy-2-acetyl-aminofluorene, which are ultimate and proximate carcinogenic forms of 2-acetylaminofluorene (AAF). Experiments with normal human fibroblasts showed that at equimolarity, N-acetoxy-AAF was more active than N-hydroxy-AAF in producing DNA lesions resulting in repair synthesis. AAF alone showed a very weak capacity for eliciting DNA repair synthesis. Reduced repair capacities were observed in all cultured XP cells exposed to N-hydroxy-AAF or N-acetoxy-AAF. The levels of impairment between unrelated patients differed significantly, but the results between brother and sister were comparable. A comparison was made of the degree of DNA repair synthesis in XP cell cultures exposed to uv irradiation (1,000 ergs/mm²), 4-nitroquinoline-1-oxide (4 NQO) and N-methyl-N-nitroso-N-nitroguanidine (MNNG). The level of DNA repair synthesis was specific for each XP cell culture and barely affected by the type of initiating agent used. (UK)

45743 CONFIRMATION OF LESIONS HAVING THE POTENTIAL FOR FORMING CHROMOSOMAL ABERRATIONS. Dewey, W. C. (Colorado State Univ., Ft. Collins). *Int. J. Radiat. Biol.*; 22: No. 1, 95-7(Jul 1972).

Late S phase synchronized Chinese hamster ovary cells were x-irradiated with doses of 600 rads of 49 keV energy, and subsequently examined for chromosomal aberrations under different temperature conditions. Chromosomal damage was the same when the cells were irradiated at either 20° or 37°C, if the cells were placed at 37°C immediately after irradiation. When the temperature was lowered to 20°C after irradiation, the amount of chromosomal damage was reduced as the repair of potential damage was reduced, with a half-time of about 15 min. When the cells were placed at 20°C for 2 hr after a 2 min period at 37°C, confirmation of potential damage appeared to have been completed. (UK)

45744 EFFECTS OF X-IRRADIATION IN VITRO ON THE TRANSPORT OF POTASSIUM IONS IN THYMOCYTES. Chapman, I. V.; Sturrock, M. G. (Dundee Univ., Scotland). *Int. J. Radiat. Biol.*; 22: No. 1, 1-9(Jul 1972).

Rat thymocytes were exposed to x radiation in the dose-range 0.2 to 20 Krads in vitro. The decrease in intracellular potassium levels produced is attributed to two separate mechanisms: inhibition of active transport and increased membrane permeability. Evidence for both mechanisms is sought by ^{42}K tracer studies and flame photometry. The influence of ouabain, incubation at 6° to 10°C, and choline chloride media is studied, both in relation to total potassium changes and in the inhibition of influx rates of potassium ions. (auth)

45745 REPAIR OF A DNA COMPLEX FROM X-IRRADIATED CHINESE HAMSTER CELLS. Elkind, M. M.; Chang-Liu, C.-M. (Hammersmith Hospital, London). *Int. J. Radiat. Biol.*; 22: No. 1, 75-90(Jul 1972).

Using velocity sedimentation in alkaline gradients of the DNA from Chinese-hamster cells, damage and repair processes were studied as a function of x-ray dose. DNA-containing material released by alkaline lysis is initially contained in a complex-containing lipid, the sedimentation of which was anomalous relative to denatured DNA from unirradiated cells. Doses of x rays small enough to be in the range that permits high survival (i.e., 100 to 800 rads) accelerated the resolution of single-stranded DNA from this DNA complex, giving rise to a species having a number average molecular weight of 2×10^8 dalton. Large doses, greater than 1000 to 2000 rads, resulted in degradation of these DNA strands. Incubation for 20 min at 37°C or 25°C after irradiation resulted in the rapid repair of both forms of damage, although the rate of repair of damage to the complex decreased with increasing dose. Repair of radiation damage to the complex resulted in a re-association of lipid and DNA. The evidence supported the possibility that a large DNA-membrane structure was the principal target of radiation. (auth)

Radiation Effects on Microorganisms (External Source)

Refer also to abstracts: 45728, 45731, 45859, and 45899.

45746 LETHAL AND MUTAGENIC EFFECT OF IONIZING RADIATIONS WITH DIFFERENT LET ON EXTRACELLULAR BACTERIOPHAGES. Shelegedin, V. N.; Bresler, S. E.; Kriviskii, A. S.; Rybchin, V. N.; Serdyuk, L. G. (Polytechnic Inst., Leningrad. Inst. of Genetics of Micro-organisms, Moscow). *Stud. Biophys.*; 28: No. 3, 213-21(1971). (In Russian).

The inactivating and mutagenic effects of high energy radiations were studied using the extracellular phages C_{d} and λ_{11} as objects. The experimental data show that both phages in the state of dry films are inactivated exponentially and almost equally by γ rays and high energy protons and with increased D_{37} by the accelerated heavy ions ^{12}C and ^{20}Ne . The inactivation rates are roughly proportional to the molecular weight of the phage DNA. In broth, the equality of both γ radiation and protons inactivating effects is also found, but the figures for inactivation rates are increased more than four times in comparison with the dry samples. Probably it is due to the effect of reaction cage well known in radiation chemistry. This was confirmed by the measurement of inactivation rate in frozen broth at the temperature of liquid nitrogen. The inactivation rate by heavy ions (^{12}C and ^{20}Ne) is 4 to 5 times lower than by γ quanta or protons. This is probably due to inhomogeneity of ionization. But the yield of mutants is the same for all three radiations if plotted against inactivation rate. This shows that the mechanism of damage is substantially identical. (auth)

45747 EFFECT OF COMBINED METHODS ON CLOSTRIDIUM SPOROGENES INOCULATED IN SAUSAGE. Incze, K. (Orszagos Husipari Kutatointezet, Budapest); Farkas, J. *Izotoptechnika*; 15: No. 3-4, 172-8(1972). (In Hungarian).

Combination of mild heat treatment and irradiation leads to a two to threefold lengthening of storage time of tinned sausage. Nitric brine improves storability. (auth)

45748 DETERMINATION OF MICROBIAL CONTAMINATION OF DISPOSABLE MEDICAL DEVICES AND THE ESTIMATION OF RADIOSENSITIVITY OF ISOLATED MICROFLORA. Gazso, L.; Igali, S.; Varteresz, V. (Frederic Joliot-Curie Sugarbiológiai és Sugáregeszegügyi Kutató Intézet, Budapest). *Izotoptechnika*; 15: No. 3-4, 187-91(1972). (In Hungarian).

Needles, connected to radiosterilized syringes, are washed with 10 ml 0, 1% peptonic solution; the needle is shaken for 30 min in the wash; which, after filtration through 0, 45 μ membrane filter, is incubated on agar plate with trypton-glucose-yeast concentrate at 32°C for 3 days. The number of colony-producing units is determined for 3 days. (auth)

45749 INVESTIGATIONS ON THE RADIOSENSITIVITY OF YEASTS. Kiss, I. (Kozponti Elelmiszeripari Kutató Intézet, Budapest). *Izotoptechnika*; 15: No. 4, 574-4(1972). (In Hungarian).

The D_{10} value of yeasts can reach those of bacterial spores. The lethal radiation dose can be determined with the help of survival curves and the factors modifying them. (auth)

45750 MICROBE-GENETICAL PROBLEMS OF RADIOSTERILIZATION. Igali, S. (Frederic Joliot-Curie Sugarbiológiai és Sugáregeszegügyi Kutató Intézet, Budapest). *Izotoptechnika*; 15: No. 3-4, 160-6(1972). (In Hungarian).

In case of avoiding mass resterilization and mass reproduction of microorganisms in the sterilized product the probability of appearance of radioresistant spore populations is very low. (auth)

45751 BIOCHEMICAL FOUNDATIONS OF SPORE RESISTANCE. Csuzi, S. (Orvosi Vegytani Intézet, Budapest). *Izotoptechnika*; 15: No. 3-4, 154-9(1972). (In Hungarian).

According to literature data on the radioresistance of spores, a significant role is played by the lowered water content, the enrichment of spore mantle in disulfides, and the non-existence of DNA synthesis. (auth)

45752 IMMUNOGENICITY OF PURIFIED VENEZUELAN EQUINE ENCEPHALITIS VIRUS INACTIVATED BY IONIZING RADIATION. Gruber, Jack (Medical Sciences Lab., Frederick, Md.). *Infect. Immunity*; 3: No. 4, 574-9(Apr 1971).

Purified and concentrated Venezuelan equine encephalitis (VEE) virus derived from tissue cultures, rendered noninfectious by ionizing radiation with retention of in vitro serological activity, also retained a high level of immunogenicity. In mice, fluid vaccines afforded excellent protection against lethal challenge with homologous Trinidad strain VEE virus. A direct relationship was observed between concentration of vaccine or number of injections and survival. One intraperitoneal inoculation of undiluted vaccine protected essentially all mice challenged 21 days later with 100,000 mouse intraperitoneal LD_{50} of virus. Similarly, mice receiving three injections of vaccines diluted 1:100 were completely protected. Noninfectious VEE virus preparations combined with adjuvant 65, a nontoxic metabolizable vehicle, were likewise very effective in protecting mice immunized intraperitoneally or subcutaneously against lethal challenge. Guinea pigs immunized subcutaneously with adjuvant-combined vaccine survived lethal challenge of 1,000,000 guinea pig intraperitoneal LD_{50} . (auth)

45753 DNA SYNTHESIS IN GAMMA-IRRADIATED RECOMBINATION DEFICIENT STRAINS OF ESCHERICHIA COLI.

Gray, W. J. H.; Green, M. H. L.; Bridges, B. A. (Sussex Univ., Brighton, Eng.). *J. Gen. Microbiol.*; 71: No. 2, 359-66 (Jul 1972).

DNA synthesis in *recA* and *recA* *extra* strains of *Escherichia coli* was strongly inhibited without delay by low doses of gamma-radiation (0 to 10 Krads). The synthesis that did occur in these strains appeared to take place mainly on chromosomes that had not received a lethal hit. Low doses produced much less inhibition of DNA synthesis in *extra*, *recB*, *recA recB*, and wild-type strains. It was concluded that the *recA*⁺ gene product acted immediately, certainly before DNA replication, in the repair of lesions produced by gamma irradiation. In *recB* bacteria, the *recA*⁺ gene product was not required for continued DNA synthesis, although the synthesis that occurred in *recA recB* strains may have been abnormal since it was not reflected in increased survival. The *extra*⁺ gene product was not necessary for continued DNA replication, suggesting that its time of action in repair may be at a later stage. (UK)

45754 CHANGES IN SENSITIVITY OF CELLS DURING EXPOSURE TO RADIATION AT LOW DOSE RATE. Bryant, P. E. (Hammersmith Hospital, London). *Int. J. Radiat. Biol.*; 22: No. 1, 67-73 (Jul 1972).

A large sparing effect was observed for *Chlamydomonas reinhardtii* when dose-rate was reduced. Cells given radiation (5 MeV electrons) at low dose-rate (20 Krads/min) were subsequently given series of graded doses at a high dose-rate (200 Krads/min). Values of *D*₀ were larger, and extrapolation numbers lower for survival curves at high dose-rate, following low dose-rate pre-treatment, than if no pre-treatment had been given. The processes that led to a change in sensitivity of cells, and lower extrapolation numbers for survival curves at high dose-rate following low dose-rate pre-treatment, may have contributed to the overall sparing effect of a reduction in dose-rate. Because the shape of the split-dose survival curve at high dose-rate was different from that of the single-dose survival curve, current models, that relate changes in shapes of survival curves as dose-rate is reduced to recovery of cells from sublethal damage, could not be applied. (auth)

Radiation Effects on Plants (External Source)

45755 (TID-25954) RADIOECOLOGY AND ECOPHYSIOLOGY OF DESERT PLANTS AT THE NEVADA TEST SITE. Wallace, A.; Romney, E. M. (California Univ., Los Angeles, Lab. of Nuclear Medicine and Radiation Biology. California Univ., Riverside). 1972. Contract AT-04-1-GEN-12. 446p. Dep. NTIS.

Separate abstracts were prepared on 18 papers. Additional papers discuss the characteristics, ecology, and mineral cycling in plants indigenous to the Nevada Test Site. (C.H.)

For abstracts of individual papers see: 45192, 45193, 45756-45769, 45914, and 45923.

45756 (TID-25954, pp 5-11) EFFECTS OF A CHRONIC LEVEL OF GAMMA RADIATION FOR 5 YEARS ON SHRUBS IN THE NORTHERN MOJAVE DESERT. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

A 33,600 curie ¹³⁷Cs γ source which was differentially shielded to increase the uniformity of the distribution of radiation was mounted on a 15-m tower in the center of a 9-hectare plot in January 1964 in the Rock Valley area of the northern Mojave Desert. In the spring of 1969 the large majority of approximately 5000 *Ephedra nevadensis* within the plot failed to produce flowers and very little vegetative growth occurred in contrast to a control plot and to other nonirradiated areas. The cumulative radiation exposure throughout the plot ranged from 3 to 29 kR. Radiation effects on *Lycium andersonii* were uncertain and effects were doubtful on other species. *E. nevadensis* and *L. andersonii* have higher nuclear volume than do the other shrub species. Considerable resistance to γ radiation was experienced for several of the other species given acute exposure under laboratory conditions. (auth)

45757 (TID-25954, pp 12-48) AN ASSESSMENT OF THE SEASONAL GROWTH ON SOME DESERT SHRUBS SUBJECTED TO CHRONIC LOW INTENSITY GAMMA RADIATION. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

A study to assess the effects of low intensity chronic γ irradiation on desert shrubs was conducted adjacent to the Nevada Test Site. From February 1964 to September 1967 a careful record was kept concerning the qualitative and quantitative responses of six dominant species (*Ephedra nevadensis*; *Franseria dumosa*; *Grayia spinosa*; *Larrea divaricata*; *Lycium andersonii*; and *Lycium pallidum*) in an irradiated and a control plot. The data

reported represent over one-half million observations of tagged terminal shoots on these shrubs. Although definite radiation effects on these species were inconclusive at the time when the study was concluded, valuable information relative to seasonal dynamics in these plants is presented. Specific data are presented relative to desert community composition and the climatic and edaphic profiles which characterize the area. (auth)

45758 (TID-25954, pp 49-53) EFFECTS OF GAMMA RADIATION ON SEED GERMINATION AND SEEDLING SURVIVAL. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Seeds of several species of desert plants were irradiated with a ⁶⁰Co γ source at various exposures and characteristics of germination and seedling survival were recorded. In general the radiation had a greater effect on seedling survival than on germination. The sensitivity of the species to the radiation was greatly related to nuclear volume. Few *Artemisia tridentata* seedlings survived for radiation levels above 15 kR and for *Yucca schidigera*, above 20 kR. *Larrea divaricata* was little affected by levels up to 40 kR while *Atriplex canescens*, *Franseria dumosa*, *Yucca brevifolia*, and *Salsola pestifera* were much more resistant to the radiation. (auth)

45759 (TID-25954, pp 54-7) RADIATION DAMAGE ON NATURAL VEGETATION IN CLOSE-IN FALLOUT AREAS. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Both β and γ radiation contributed to the damage of vegetation in close-in fallout areas at the Nevada Test Site inasmuch as their effects were additive. However, because of its inherent nature, beta radiation contributed the most damage when fallout debris was deposited upon and retained by plant foliage. Where fallout radiation levels were sufficiently high, complete defoliation and kill of natural vegetation occurred, and, in areas of fallout patterns peripheral to where all or most plants were killed, the radiation damage to shrubs was often manifested in a particular fashion which depended upon the manner in which fallout debris was deposited. (C.H.)

45760 (TID-25954, pp 65-8) SENSITIVITY OF DESERT PLANTS TO ACUTE GAMMA RADIATION. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

The radiation sensitivity of some 30 different species of desert plants was compared in the laboratory using a 10,000 curie ⁶⁰Co γ source. Some deciduous plant species were difficult to score because of their inherent tendency to become dormant under environmental stress. Dormant plants survived much higher levels of radiation than actively growing plants. The findings reported here are for plants irradiated in a succulent, vegetative state. Based upon a limited amount of data, the order of decreasing sensitivity to acute gamma radiation generally was the same as the order of decreasing nuclear volume or ICV of the species. (auth)

45761 (TID-25954, pp 73-4) EFFECT OF ⁶⁰Co RADIATION ON EMERGENCE AND SEEDLING GROWTH OF LYCIUM PALLIDUM AT TWO LEVELS OF HYDRATION. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Dry and moist seeds of *Lycium pallidum* were exposed to doses of 0, 5 kR, 10 kR, and 20 kR of ⁶⁰Co γ radiation. Control seeds were oven dried at 105°C for 24 hours to establish the moisture content of the seed. Immediately after radiation treatment the seeds were sown in fine vermiculite in clay pots and placed in a glasshouse (temperature range 15 to 25°C) to observe emergence of seedlings. Indications of this study are that dry seed of *L. pallidum* will tolerate 10 kR of radiation and that moist seed will tolerate 5 kR, without significantly diminishing the emergence of seedlings or their early survival. At radiation levels much in excess of these, however, the rate of emergence declined sharply and the emerged seedlings had a low survival rate. (C.H.)

45762 (TID-25954, pp 75-86) REVEGETATION IN AREAS DAMAGED BY CLOSE-IN FALLOUT FROM NUCLEAR DETONATIONS. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Revegetation changes were followed in the close-in fallout area of a plowshare nuclear test at Nevada Test Site which killed most of the native vegetation in one area and severely damaged it in others. The *Artemisia* species dominate the vegetation of this area at Pahute Mesa. In the partially damaged area, *Artemisia* seedlings are reestablishing themselves but the area has become a grassland and few *Salsola* are present. Allelopathy may be involved in germination of new *Artemisia* seedlings. In the com-

pletely damaged area, *Salsola* populations have been dense and a new population of *Artemisia* seedlings have died. *Grayia spinosa* and to a lesser extent *Eurotia lanata*, *Ephedra nevadensis*, *Tetradymia axillaris* and *Chrysothamnus nauseosus* have successfully reestablished themselves. At the disturbed area around the Sedan crater only *Atriplex canescens* of the original perennial shrubs has germinated and survived. Some *Lycium andersonii* have survived near the crater (300 m away) with new growth sprouting from old crowns. *Salsola iberica* and *Salsola paulsenii* have been the principal plant species in this area since the event. There is evidence that browsing animals present could prevent recovery of shrubs in the area by eating any new seedlings which germinate or new shoots which sprout from old crowns. (auth)

45763 (TID-25954, pp 87-101) CHARACTERISTICS OF ARTEMISIA SPECIES (SAGEBRUSH). 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

The *Artemisia* species in general belong to the Great Basin Desert, and our soil temperature studies indicate that *A. tridentata* prefers cool soil temperatures (13 to 24°C) to hot soil (28 to 35°C). Optimum growth was obtained under controlled air and root temperature conditions of 24°C root temperature and 16°C air temperature. *A. tridentata* seeds readily germinate and under glasshouse conditions the roots of seedlings elongate at the rate of about 2.5 cm per day. Stem cuttings were easily rooted. *A. tridentata* in our Nevada Test Site study areas apparently live an average of about 30 years, but some individuals do live much longer, especially under optimum edaphic and moisture conditions. About one-third of the plants in the stand were dead but there are sufficient new seedlings present in the area to replace the old stand in 20 to 30 years if the same rate of new seedlings is maintained each year. Rabbits graze on new *A. tridentata* seedlings and this may be an important factor influencing seedling survival. Stands of *A. tridentata* use soil moisture sufficiently to curtail the growth of grasses. In areas in which the shrubs had been destroyed by irradiation from fallout debris, the first stage of revegetation was from grasses. However, new seedling development has occurred in these areas in more than sufficient amounts to reestablish the original stands of *A. tridentata*. Possibilities of allelopathy between the species present is of continued interest to us. *A. tridentata* appears to be sensitive to chlorides and to changes in soil pH. It is commonly found growing on calcareous soils, yet it grows well on non-calcareous soil. Low calcium levels of about 10^{-4} N in nutrient solution inhibited its growth. Field grown *A. tridentata* absorbs little sodium and it generally has lower phosphorus than many other shrub species. (auth)

45764 (TID-25954, pp 102-9) CHARACTERISTICS OF EPHEDRA SPECIES (MORON TEA). 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Four species of *Ephedra* occur at the Nevada Test Site and these separate into cold and hot desert species. *Ephedra* species grow slowly and in some years do not produce flowers or seeds. *E. nevadensis* has a clumping habit due to underground stolon production. Seedlings and cuttings of *E. viridis* and *E. nevadensis* were prepared and used for several studies. *E. viridis* grew somewhat better at a root temperature of 21°C than at either 28 or 16°C. Shoots of *E. nevadensis* do not accumulate sodium, are low in phosphorus and high in aluminum. Very young shoots are low in potassium and high in calcium in contrast to most plant species. *Ephedra* species are extremely sensitive to γ radiation. Cuttings of *E. viridis* were exposed to a ^{60}Co γ source and the LD₁₀₀ dose from acute exposure was determined to be 3 kR. (C.H.)

45765 (TID-25954, pp 115-32) CHARACTERISTICS OF LARREA DIVARICATA (CREOSOTE BUSH). 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Larrea divaricata is the most common shrub on a large portion of the Nevada Test Site. It is extremely drought resistant and well adapted for survival in the desert, reaching ages probably of well over 100 years. Shoots continually die and new ones arise from the crown. Leaves are functional at relatively low water contents. Germination of seed occurs under relatively warm and moist conditions, but survival is rare. Rapid depth of rooting is one of its survival mechanisms. It branches laterally and occupies much of the soil not covered by other shrubs. It responds to summer rains and may set seed a second time in one year. Limiting factors for its growth under natural conditions include water and mineral nitrogen. Good soil aeration is necessary and it does not spread into areas where water stands for periods of time. It is widely spaced but is in clumps with other species and does not seem to compete with itself by chemical

means. Spacing can be random, clumped, or regular. Being a hot desert species, it prefers warm to hot soil; but is not completely dormant during the winter season when air temperatures drop below freezing. It can be injured by frost but can recover. It is adapted to a wide range of soil pH even though it is native to calcareous areas high in pH and seems to prefer calcareous soil. Its mineral analysis revealed that it is low in zinc and copper when grown in calcareous soil. It tends to exclude sodium and tolerates little of it. It responds vigorously to severe pruning, but in the first season flowering was decreased by pruning. Ecotypes contribute to its variability. Rooted cuttings were made from one-year-old seedlings, but not from much older plants. Adventitious shoots grow from old severed roots. One-year-old *L. divaricata* wood is used for food by jackrabbits. *L. divaricata* is more resistant to ionizing radiation than are many other species. (auth)

45766 (TID-25954, pp 133-46) CHARACTERISTICS OF LYCIUM SPECIES. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Lycium andersonii, *Lycium pallidum*, and *Lycium shockleyi* appear to have different soil requirements. *Lycium* species are winter dormant and leafless because of extremes in soil moisture and soil temperature. Adventitious shoots form on roots of *L. andersonii* and cuttings of this species root readily. Seedlings of *L. andersonii* and *L. pallidum* germinate readily in the glasshouse when viable seed is obtained. Cuttings of *L. pallidum* seedlings have rooted; but some individual plants have been propagated by layering. Glasshouse-grown leaves of *L. andersonii* and *L. pallidum* are much different from those grown in the field. The ease of leaf drop in *L. andersonii* may be a survival mechanism. Shoots of rooted cuttings of *L. andersonii* grow several feet in a single flush in the glasshouse compared with a few cm in the field. *L. pallidum* grows poorly in the glasshouse unless soil temperature is low, at least initially. Growth of *L. pallidum* appears to be dependent upon soil temperature while *L. andersonii* does not. The mineral composition of *L. andersonii* is characterized by high calcium and potassium and low sodium and zinc contents. *L. pallidum* and *L. shockleyi* have lower calcium and potassium and higher sodium contents than *L. andersonii*. *Lycium* species appear to accumulate lithium. *L. andersonii* appears to have a low phosphorus requirement. *L. andersonii* is highly sensitive to ionizing radiation in relation to other shrub plant species found at the Nevada Test Site and it has a high nuclear volume. Grafting compatibility occurs between *L. andersonii* and *L. pallidum*. (auth)

45767 (TID-25954, pp 147-52) CHARACTERISTICS OF SALSOLA SPECIES. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Salsola species (commonly known as tumbleweed or Russian thistle) are abundant at the Nevada Test Site, especially where soil has been disturbed or where the native vegetation has been killed which results in a more favorable water balance. The species appear to have several adaptive mechanisms which result in its survival under arid conditions. These include rapid germination and penetration into soil, increased resistance to water loss through transpiration as soil temperature increases, and high photosynthetic efficiency through the C-4 dicarboxylic fixation pathway. *Salsola* is very resistant to ionizing radiation. Exposure of seeds to up to about 60 kR from a ^{60}Co γ source resulted in no apparent effect on germination or growth of the seedlings. Exposure to between 60 and 400 kR resulted in germination and elongation of shoots and roots to a certain limit only. Exposures above 500 kR resulted in progressively less elongation and in death of seedlings. The first response curve obviously relates to cell division. The implication is that the seed contains the fully differentiated seedling which grows by cell elongation when exposed to water at the proper temperature. No cell division is needed for this phase of growth; hence there is no effect of the radiation at levels which inhibit nucleic acid synthesis and function. (C.H.)

45768 (TID-25954, pp 153-61) CHARACTERISTICS OF FRANSERIA DUMOSA (BURRO BUSH OR BURR SAGE). 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Franseria dumosa is a shrub which dominates or associates with *Larrea divaricata* at the Nevada Test Site. It is relatively short lived, compared with other shrubs. It is deciduous depending more upon soil moisture than on hot or cold temperatures. Late summer rains of sufficient magnitude cause it to resume growth, flower, and set seed, thus sometimes completing two productivity cycles in a given year. It sets fruits in any season of the year. Leaf drop under conditions of stress may be a survival mechanism under desert conditions. Root growth of

seedlings is extremely rapid. *F. dumosa* requires warm conditions for growth, especially warm soil and air temperature, and it is not found at high altitudes. Low soil temperature (11°C) greatly decreased growth. Freezing injured the vegetative growth. It is tolerant of a moderate degree of salinity and is relatively insensitive to soil pH. It accumulates large quantities of boron and of calcium. It is moderately resistant to ionizing radiation. Acute γ irradiation with 4 kR was lethal to *F. dumosa*. (auth)

45769 (TID-25954, pp 188-98) CHARACTERISTICS OF YUCCA SPECIES *Y. SCHIDIGERA*, *Y. BREVIFOLIA*, AND *Y. BACCATA*. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

Age of *Yucca schidigera* and *Yucca brevifolia* can be estimated reasonably well by either counting vascular bundle rings or by counting numbers of blades, or from measuring height. The first method requires sacrificing the plant. On plants sprayed with water-base paint to differentiate between previous and new year's growth, an average of 18 new leaf-blades in 3 sets of 6 normally developed each year for both species. More or less of these sets of leaves may develop in the unusually wet or dry years. There is some evidence that the *Y. brevifolia* population is advancing down the bajada onto the playa in Yucca Flat. These *Yucca* plants do not flower each year. *Y. brevifolia* is more tolerant of salt than is *Y. schidigera* and it absorbs less sodium-factors which may be involved in the natural distribution of these species. Both species of *Yucca* seedlings grew best at root temperatures near 18°C and without CaCO_3 in the soil. Acute levels of gamma radiation prior to a subsequent year of continued growth in soil resulted in a slight growth stimulation from 4 to 12 kR, followed by a suppression of growth as the dosage levels were increased to 36 kR. Gamma radiation treatments had great influence on the ion transport processes in these plants. (auth)

45770 SOME SPECIFIC QUESTIONS OF RADIATION MUTAGENESIS OF POTATO. Tarasenko, N. D. (Inst. of Cytology and Genetics, Novosibirsk, USSR). *Izv. Sib. Otd. Akad. Nauk SSSR*, No. 5, Ser. Biol. Nauk; No. 1, 99-108 (Apr 1971). (In Russian).

The results of the three year investigation of variability and heritability of productivity, starch, and raw protein contents of potato are given. Phenotypic correlations between the above mentioned features and genotypic correlations between productivity and starch content were determined. Genotypic correlations were higher than phenotypic ones. Some data are given showing the possibility of changing the correlations between the bulb size and starch content under the influence of gamma-rays and fast neutrons in a series of mutants as compared to the initial varieties, as well as the possibility of experimental dechimerization of genetically chimeric varieties by means of radiation. (auth)

45771 RECESSIVE GENES PRODUCING ALBINO AND LIGHT GREEN SEEDLINGS IN SUGI, *CRYPTOMERIA JAPONICA* D. DON. Ohba, Kihachiro (Inst. of Radiation Breeding, Omiya, Japan); Mural, Masafumi. *Nippon Rin Gakkai-Shi*; 53: No. 6, 177-80 (Jun 1971). (In Japanese).

Out of commercial sugi (Japanese cedar) saplings planted in the gamma field of the Institute of Radiation Breeding in 1962, four heterozygous saplings for a recessive albino gene, two heterozygous saplings for a recessive gene producing light green or yellowish green seedlings, and one doubly heterozygous sapling for both genes were found. The recessive homozygotes are lethal soon after germination under natural conditions. These recessive genes were not expected to be produced by gamma ray irradiation, and they could have been transmitted to the progeny from the heterozygous mother trees from which commercial seeds were collected. After two years' crossing experiments including selfing with gibberellin-induced flowers, it was found that singly heterozygous saplings segregated the seedlings in the ratio of normal vs albino or light green as 3:1. Meanwhile, one doubly heterozygous sapling segregated normal vs albino vs light green as 9:4:3, because albinism has an epistatic effect on the light green. No allelic relation was proved between the recessive gene responsible for light green seedlings and other recessive genes which produce light green seedlings in a sugi individual and two cultivars, namely, G-5, Kumotoosi and Kuma-sugi, respectively. (auth) (Japan)

45772 CYSTEINE PROTECTION AGAINST ETHYLENIMINE DAMAGE IN BARLEY SEEDS. Kaul, B. L.; Choudhary, D. K. (Regional Research Lab., Jammu-Tawi, India). *Radiat. Bot.*; 12: No. 2, 57-61 (May 1972).

Experiments were performed to find out if cysteine could provide protection against the toxic and mutagenic effects of EI, a mono-functional alkylating agent. Cysteine was administered either pre or post EI treatment and in combination with it. It was found that in all three cases cysteine provided a strong protection against EI

damage. Protection was, however, maximum when both EI and cysteine were administered in a mixture. It is pointed out that cysteine being very sensitive to alkylation, mops up all of the excess of mutagen in post treatments and alkylates mol for mol in other treatments, thus, removing the excess of mutagen that otherwise would have acted detrimental to the cell, causing a host of physiological damages. (auth)

45773 INFLUENCE OF NEUTRON IRRADIATION ON SEED GERMINATION AND PLANT GROWTH OF TOMATOES. Rusu, F.; Luca, I.; Matei, V.; Savitchi, O. *Lucr. Stiint. Agron. Ion Ionescu De La Brad, Iasi*; 427-36 (1970). (In Romanian).

Experiments were carried out on tomato plants from seeds treated with different neutron doses, in dry or moist state. In the first stage, doses between 500 and 1500 rem were used to establish the neutron treatment. In the second stage the prolonged effect of the treatment applied in the first generation was studied. The conclusion was reached that 500 rem doses applied to moist seeds stimulate growth and development. The plants were more hardy and surpassed control plants by 1 to 2 stages of organogenesis. The positive effect was observed to be maintained also in second generations, as the nursery transplants obtained surpassed the control in growth and development. (Rom. Sci. Abstr.)

45774 GAMMA RADIATION-INDUCED CHANGES IN THE ANATOMIC STRUCTURE AND CHEMICAL COMPOSITION OF SPRUCE WOOD. Katkevich, Yu. Yu.; Odintsov, P. N.; Pendere, M. K. (Inst. of Wood Chemistry, Riga). *Khim. Drev.*; No. 5, 11-22 (1970). (In Russian).

Spruce seedlings were irradiated with ^{60}Co gamma radiation during the vegetative period (60 to 14,700 R). That year and during the two following years at the end of the vegetative period the anatomical structure and the chemical composition were studied. Small doses evoked no changes; at 2,000 R (5 R/hr) the cambium completely ceased to divide. Radiation damage to cambium by doses greater than 1,000 R (at 0.3 R/hr) was still evident two years after irradiation. Tracheids forming during exposure to radiation were of small dimensions and irregular form, and the wood had low specific weight. Shifts in the ratios of the chemical components constituting the wood were observed. The synthesis of cellulose and lignin were the processes most sensitive to radiation; the formation of hemicellulose, and especially the polyuronides, was affected the least by radiation. (tr-auth)

45775 STUDIES OF INDUCED MUTATIONS IN PEAS. XXVI. GENETICALLY CONDITIONED DIFFERENCES IN RADIATION SENSITIVITY. 4. Blixt, Stig. *Agri Hort. Genet.*; 28: 55-116 (1970).

Four pea lines at three different levels of gamma-ray sensitivity were crossed, and the sensitivity of the lines in the F_1 and F_2 -generations was studied. The criteria of radiation effect used have been the dose for fifty per cent germination inhibition (DRG 50), growth inhibition (DRE 50), survival of main shoot (DRP 50) and of main and secondary shoot (DRT 50), and the dose producing spotting degree 2.5 (DS 2.5). The characteristics studied in relation to sensitivity have been the number of recessive genes (NR and NR^A), flowering time (S-F) and the genes A, Cor, D, Di, F, Fs, Gri, I, Kp, Kpa, Le, Mifo, O, Pl, R, U and V. Of the four lines used, line 180 was the most radiation-resistant, the lines 1147 and 1241 the most radiation-sensitive, while line 1405 was intermediate. The F_1 -generations showed that the different criteria used for measuring sensitivity probably reflected radiation effects on different mechanisms. Also, the different crosses differed with respect to sensitivity: crosses involving line 180 were generally more resistant than those not involving this line. Of these latter crosses, the crosses involving line 1405 were more resistant than the cross not involving it. The results indicate that the gene A is of the first importance in determining the sensitivity; recessivity for this gene results in a low level of resistance to gamma-radiation. In genotypes dominant or heterozygous for A, the sensitivity is influenced by the number of double-recessive genes present in the genotype. The results from the analysis of the F_2 -generations showed a clear effect of the gene R in chromosome VII, genotypes recessive for this gene being more sensitive. Chromosome I was also found to be connected with sensitivity. Evidence seems to support the hypothesis that heterochromatic regions in these two chromosomes are involved in the gamma-sensitivity differences found in *Pisum*. Radiation sensitivity, its mechanisms and causes in micro-organisms and higher developed organisms, and the relation of these results to the results from *Pisum* are discussed. (auth)

45776 MUTATION GENETICS IN *PISUM*. Blixt, Stig. *Agri Hort. Genet.*; 30: 1-293 (1972).

A review of the literature on the genetics of the genus *Pisum* is presented. Some topics reviewed are systematics, ontogeny, cytology, genetics, spontaneous mutations, induced mutations, chromosome aberrations, leaf aberrations, genetically conditioned

sensitivity of the M_1 generation, factors modifying the sensitivity of the M_1 generation, chimaerism, the mutation rate, the mutation spectrum, linkage relations of mutations, and the role of mutations in plant breeding. A correlation was found between toxic and mutagenic effects of ionizing radiation, and it was found that radio-sensitivity was genotypically conditioned. (H.L.W.)

45777 (TT-54002/4, pp 442-50) **RADIORECOVERY EFFECT OF YEAST EXTRACT ON CARNATION CUTTINGS (DIANTHUS CARYOPHYLLUS L., WHITE SIM) IRRADIATED WITH ^{60}Co γ RAYS. PRELIMINARY MORPHOLOGICAL STUDY.** Fouletier, Bernadette; Brun, Therese; Jonard, Robert; Pereau-Leroy, Pierre (CEN de Cadarache, Saint-Paul-les-Durance, France. Univ., Montpellier, France). *Compt. Rend., Ser. D*; 274: No. 9, 1315-18 (28 Feb 1972). (In French).

Yeast extract has a radiorecovery effect, and increases the percentage of white petals of White Sim carnations subjected to γ irradiation. These facts indicate that the recovery is due to a distribution of cells damaged at the epidermic cellular base that determines the characteristics of the white flower. The processes occurring are thus more involved than the simple recovery by compensation resulting from the less damaged underlying cells carrying the red flower characteristics, as is the case with spontaneous recovery. (France)

Radiation Effects on Animals (External Source)

Man

Refer also to abstracts: 45290, 45732, and 45911.

45778 (NP-tr-1941) **RADIATION HAZARDS AND RADIATION PROTECTION.** Hug, O. Translated by D. H. Garrett (State Electricity Commission of Victoria), from Atomwirt., Atomtech.; 16: No. 6, 294-300 (Jun 1971). 17p. Dep. NTIS (U. S. Sales Only).

An unrealistic over-estimation of the hazards of exposure to radiations, that is no longer consistent with the findings of experience, can be just as harmful as the view that the risks are more or less a matter of chance. All forms of radiation associated with the operation of nuclear power plants must be strictly monitored. If the recommendations and existing regulations are correctly understood and properly applied in the practical measures for protection against radiation, according to the known facts the reduction of the threshold values to a tenth of those currently in force appears to be unnecessary. The intensity of the radiation corresponding to these values is appreciably lower than that of the dose equivalent from natural radiation sources. As the utilization of nuclear energy is unavoidably associated with certain radiation effects, not only on personnel employed in the various plants but also on large groups of the population, it is necessary to determine the ways in which the risk attendant on the different forms of radiation depends on the degree of exposure, and to ascertain the maximum permissible doses that are commensurate with the safety of the individual and of the community at large. The theoretical principles of the relative biological effectiveness of radiations are reviewed. Delayed manifestations of radiation injuries, particularly malignant diseases, are discussed. A critical evaluation of the maximum dosage currently accepted as permissible is presented. (auth)

45779 (TT-54002/4) **POLISH REVIEW OF RADIOLOGY AND NUCLEAR MEDICINE.** Bimonthly of the Polish Radiological Association, Warsaw, Poland. Translated Reprint of Pol. Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971). 123p. NTIS.

Separate abstracts were prepared for seven of the 20 articles appearing in the journal. The other articles discuss radiographic finding in various diseases, and the effectiveness of various diagnostic procedures. (E.R.B.)

For abstracts of individual papers see: 45780-45784, 45906, and 45927.

45780 (TT-54002/4, pp 430-6) **EVALUATION OF THE CAUSES OF FAILURES IN ROENTGENTHERAPY OF CARCINOMA OF THE LARYNX.** Ulkowski, Mirosław; Hartel-Ulkowska, Narcyza (Inst. of Radiology, Warsaw). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

Causes of failures in roentgentherapy of carcinoma of the larynx were analyzed. The main cause was late clinical stage and local advancement of the tumor. Another important cause, according to the authors, is connected with histological type of the tumor, namely anaplastic carcinoma. (auth)

45781 (TT-54002/4, pp 437-41) **SELECTED PROGNOSTIC ELEMENTS IN CARCINOMA OF THE LARYNX.** Światnicka, Jolanta; Heliński, Andrzej; Vorbrodtowa, Janina; Godawski, Jerzy. Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

On the basis of observations of 265 patients with carcinoma of the larynx treated with roentgen rays, the possibility of curing patients with T_3 , N_1 , N_2 , immobilization of the vocal cords, edema of the laryngeal mucosa during treatment, and patients with late response to irradiation was evaluated. (auth)

45782 (TT-54002/4, pp 442-50) **REMARKS ON LOCAL METASTASES TO LYMPH NODES IN TUMORS OF THE LARYNX, TONSILS, AND NASOPHARYNX AFTER X-RAY THERAPY.** Borejko, Juliusz (Inst. of Radiology, Warsaw). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

Results of x-ray therapy of tumor metastases to the lymph nodes in carcinoma of the larynx (105 patients), tonsils (39 patients), and nasopharynx (23 patients) are discussed. For various reasons, these patients were not operated. The reactions of the lymph nodes to irradiation were divided into three groups: favorable, palliative, and unfavorable. In the group of favorable reactions, the chances of survival in all three groups according to anatomic localization were about 50%. Patients with palliative and unfavorable reactions had small chances of survival. A striking and statistically significant observation concerns the number of positive reactions in tumors of the tonsil and nasopharynx, compared with tumors of the larynx. (auth)

45783 (TT-54002/4, pp 451-5) **PROGNOSTIC ELEMENTS IN INOPERABLE CARCINOMA OF THE BREAST.** Hliniak, Andrzej; Wojcieszek, Zbigniew; Ślomska, Janina; Petryna, Alicja; Godawski, Jerzy (Inst. of Oncology, Gliwice, Poland). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

On the basis of observations on 569 women with inoperable carcinoma of the breast treated with x rays in the years 1951 to 1963, the prognostic significance of the length of clinical history and the "T" and "N" characteristics of the tumor was analyzed. (auth)

45784 (TT-54002/4, pp 456-61) **DISTRIBUTION OF DOSES IN THE LESSER PELVIS IN THE TREATMENT OF CARCINOMA OF THE CERVIX UTERI. CALCULATION OF DOSES FROM RADIUM APPLICATORS AT SELECTED POINTS OF THE LESSER PELVIS.** Szymczyk, Wiesława; Siuda, Jerzy; Medvey, Wojciech; Polak, Bogumił; Polok, Bogdan (Inst. of Oncology, Cracow). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

A graphic and an analytical method of determination of the coordinates of points A are presented. It was assumed that the points and the first container of the uterine probe lie in a plane vertical to the sagittal plane. The points corresponding to the position of the obturator lymph nodes were referred to the skeleton (point "O"). Calculations were carried out on 90 patients. The error in the evaluation of doses in points A is calculated at $\pm 8\%$, in points O at $\pm 5\%$. The radiation absorption in the tissues was taken into consideration. (auth)

45785 (TT-54002/5) **POLISH REVIEW OF RADIOLOGY AND NUCLEAR MEDICINE.** Bimonthly of the Polish Radiological Association, Warsaw, Poland. Translated Reprint of Pol. Przegl. Radiol. Med. Nukl.; 35: No. 5, (1971). 138p. NTIS.

Separate abstracts were prepared for two of the 22 papers appearing in the journal. The other papers discuss radiologic techniques and the effectiveness of various radiotherapeutic methods. (E.R.B.)

For abstracts of individual papers see: 45786 and 45787.

45786 (TT-54002/5, pp 590-3) **DISTRIBUTION OF DOSES IN PATIENTS WITH HODGKIN'S DISEASE DURING CONVENTIONAL RADIOTHERAPY.** Głinska, Halina; Skolyszewski, Jan (Inst. of Oncology, Cracow). Translated from Pol. Przegl. Radiol. Med. Nukl.; 35: No. 5, (1971).

The differences between the measured and planned doses in both areas examined depend on the physical properties of radiation initiated by a voltage of 200 to 250 kV, and technical conditions of therapy. Tables of doses and diagrams of isodoses, on the basis of which therapy is planned, are based on results of measurements made under model conditions, differing from clinical practice. When irradiating the mediastinum, the most important cause of the difference is the increased absorption of radiation by the bones of the vertebral column, ribs, and sternum. The difference in the dose between the central and peripheral parts of the irradiated tissue block is a result mainly of a smaller contribution of dispersed radiation in the marginal part of the beam. When irradiating the cervico-supraclavicular area, the main factor is connected with the irregular shape of the tissue block and consequently insufficient contribution of the dispersed dose. From the clinical point of view, these phenomena are clearly unfavorable, constituting an obstacle to application of a uniform dose to the whole irradiated area. Since the results of roentgentherapy in malignant lymphogranulomatosis characteristically depend on the magnitude of dosage, development of a method permitting better distribution of doses should have a favorable effect on the therapeutic results.

Megavolt therapy offers such a possibility. The observed discrepancies between actual and planned doses were sometimes as high as 30%. Hence, therapy must be guided in a large degree by evaluation of the patient's clinical state, and not only by the nominal dosage. Individualization of therapy taking into account the intensity of the reaction to radiation and regression of the tumor is an important factor. (auth)

45787 (TT-54002/5, pp 602-4) TRIAL OF CORRELATING THE BLOOD PICTURE WITH ABSORBED DOSES IN PERSONS EXPOSED PROFESSIONALLY TO IONIZING RADIATION. Lobodzic, Włodzimierz; Pinkwart, Harry; Czaplewski, Stanisław (Inst. of Oncology, Gliwice, Poland). Translated from Pol. Przegl. Radiol. Med. Nukl.; 35: No. 5, (1971).

Analysis of the data does not show correlation between the blood pictures and doses absorbed individually. In the range from 0 to 20 rems absorbed during 7 yrs, progressive deterioration of the blood picture was not observed. We are aware that the material of 98 persons is too small for statistical conclusions; however, the doses lie within the limits of annual permissible absorption. In spite of the small material, the long period of observation of over 7 yrs justified the analysis. A similar analysis in institutions employing a larger number of personnel exposed to ionizing radiation would be interesting. The data indicate that in personnel exposed to doses not exceeding the permissible annual norms, or only sporadically in some months, the blood picture is not impaired proportionally to the individual doses absorbed. (auth)

45788 ESTIMATION OF SULPHUR CONTENT IN HUMAN HAIR AND NAIL FOR USE IN FAST NEUTRON DOSIMETRY. Amin, S. R.; Gupta, B. L. (Bhabha Atomic Research Centre, Bombay). Health Phys.; 23: No. 2, 243-4 (Aug 1972).

Sulfur activation in hair and nail has been used for the evaluation of the fast neutron dose received by persons involved in criticality accidents, by measuring the beta activity arising from the $^{32}\text{S}(n, p)^{32}\text{P}$ reaction. Work reported here was undertaken to investigate whether the sulfur content in hair shows any dependence on age, sex, or food habits. An increase was found with age up to 35 and then a decrease. The content was found to be higher in non-vegetarians than in vegetarians, and male hair content was higher than female. Some variations between the content were found in people of different races. In view of these divergences in values it is thought that the possible variation of sulfur content should be considered when fast neutron doses are evaluated from sulfur activation in hair. (UK)

45789 QUANTITIES FOR THE DESCRIPTION OF NEUTRON IRRADIATIONS OF MAN. A LOOK AT THE MECHANISM OF OVER-CONSERVATISM. Pretre, S. (Research Inst. for Protective Construction, Zurich). Health Phys.; 23: No. 2, 169-81 (Aug 1972).

The conversion from neutron fluence into either absorbed dose or dose equivalent is discussed. It is shown that the quality factor can have four different meanings which should be separated properly from each other. It is proposed to distinguish between conservative quantities which are used for preventive purposes (radiation protection) and which refer to late somatic and genetic effects of chronic exposures at low dose-rate (conditions within ICRP limits), and realistic quantities which are used to estimate the actual damage (radiation accidents, radiotherapy, civil defense, animal experiments) and which refer to early somatic effects of an acute exposure at high dose-rate (conditions mostly beyond ICRP limits). This distinction is urgently needed, particularly in the case of dose-equivalent, because the unit "rem" is not unequivocally defined. There are in fact two different "rem", one of them being the old but realistic "Röntgen Equivalent Man", while the newer one includes comparatively more conservatism. Among the many interactions neutrons can undergo in tissue, the capture $\text{H}(n, \gamma)\text{D}$ is of great importance to man, due to the large dimensions of his body. Therefore, a substantial part of any neutron dose is absorbed by man in the form of a gamma dose. With this in mind, it is felt that the internationally recommended quality factors for neutrons are generally too high. The question is raised why the protection against neutrons is more conservative than the protection against gamma quanta. (auth)

45790 ASPECTS OF RADIATION CONTROL FOR CLINICAL IMPLANTS OF ^{252}Cf . Oliver, G. D. Jr. (Anderson Hospital and Tumor Inst., Houston, Tex.). Health Phys.; 23: No. 2, 187-92 (Aug 1972). (CONF-710710-12).

From 16th annual Health Physics Society meeting; New York, N. Y. (12 Jul 1971).

The safe use of neutron-emitting isotopes, such as ^{252}Cf for interstitial and intracavitary radiotherapy, raises new problems of radiation protection in a hospital. Techniques associated with conventional radiotherapy sources have been adapted for the neutron sources. The use of source afterloading reduces the dose to personnel by eliminating source handling in sterilization pro-

cedures and in the surgical theater. In addition, the time required for source manipulation into implants can be minimized by using specialized instruments. Doses to nurses involved in routine care of the implanted patient can be at an acceptable level if a portable barrier is utilized when possible and if special training and practice are conducted. Based on these approaches, the use of californium has proven feasible in the treatment of more than 20 neoplastic lesions of patients. The maximum probable dose received by radiotherapists has been 1.1 mrem per microgram of ^{252}Cf implanted in regions of the head and neck, skin, and pelvis. Each attending nurse is estimated to have received 2 mrem per microgram implanted. (auth)

45791 RADIATION ACCIDENTS IN UNIVERSITY LABORATORIES. Hughes, D. Accident Anal. Prev. (Elmsford, N. Y.); 2: No. 1, 29-34 (May 1970).

Details are presented of 10 accidents that occurred at the University of Leeds during the period 1958 to 1968, involving radioactive substances (^{60}Co and tritium-labelled substances) or electrical machine sources of ionizing radiation. The University does not operate a reactor and most radiation work is performed with radioactive substances in tracer amounts, two gamma-ray irradiation facilities, 40 crystallographic x-ray sets, and a Van de Graaff generator. For each case described, appropriate safety measures to prevent repetition are indicated. (Occup. Safety and Health Abstr.)

45792 EFFECTS OF AGE AND RADIATION EXPOSURE ON CHROMOSOMES IN A MARSHALL ISLAND POPULATION. Demoise, Charles F. (Microbiological Associates, Inc., Bethesda, Md.); Conard, Robert A. J. Gerontol.; 27: No. 2, 197-201 (Apr 1972).

The effects of aging and long-term radiation exposure on chromosome counts in cultured peripheral blood lymphocytes and the influence of plasma from irradiated subjects were tested in a Marshall Island population. Both hypodiploid and polyploid levels were found to be related to the patients' age. Females over 50 years of age and older previously exposed males had approximately 1.5 times as many hypodiploid cells as did the younger subjects, while polyploid levels were sharply reduced in all subjects over 50 years. Radiation was also found to affect hypodiploid levels. This was most pronounced in the males where those exposed had 2.8 times as many hypodiploid cells as the unexposed. Similarly, radiated females had 1.3 times as many hypodiploid cells as the unexposed. No relationship was found between polyploid level and radiation. Both sex and chromosome size appeared implicated as possible factors contributing to hypodiploid levels. In all subjects, regardless of sex or exposure to fallout radiation, the largest and most frequent loss of chromosomes occurred in the G(Y) group (2.3 times the expected loss). In the C(X) chromosome group the females lost 15.2% more chromosomes than expected, and the males lost 12.6% fewer than predicted. No apparent sex or radiation effect occurred in the other five chromosome groups. A series of additional cultures indicated the presence of a chromosome breakage factor in the plasma from exposed subjects. Although chromosome aberrations were nearly doubled in cultures containing the exposed plasma, aneuploid levels were not affected. (auth)

45793 ACNEFORM CHANGES IN AREAS OF COBALT IRRADIATION. Trunnell, Thomas N.; Baer, Rudolf L.; Michaelides, Paraskevas (New York Univ., N. Y. Veterans Administration Hospital, New York). Arch. Dermatol.; 106: No. 1, 73-5 (Jul 1972).

Two patients developed an acneform eruption in skin areas on the trunk. Both patients had received γ radiation through the involved skin areas for treatment of internal tumors. The cutaneous changes in both cases responded to conventional therapy for acne. (auth)

45794 CHROMOSOME ABERRATIONS IN HUMAN BONE-MARROW CELLS AFTER A SINGLE LOCAL GAMMA-IRRADIATION. I. STUDY OF TYPES AND FREQUENCIES OF ABERRATIONS IN DYNAMICS AFTER IRRADIATION AT DOSES OF 250 AND 500 RDS. Petrova, S. A.; Pyatkin, E. K.; Suskov, I. I.; Alexandrov, N. N.; Vorob'ev, A. I. (Research Inst. for Oncology and Medical Radiology, Minsk. Biophysical Inst., Moscow). Stud. Biophys.; 28: No. 3, 157-64 (1971). (In Russian).

Direct chromosome studies of bone-marrow cells were made in a control group of 14 tumor patients and in a group of 88 patients at different intervals of time (1 to 168 hrs) after a single local irradiation with ^{60}Co gamma rays at doses of 250 and 500 rads. The highest frequency of aberrant cells was observed 18 and 21 hrs after irradiation with a dose of 250 rads and 24 hrs after irradiation with a dose of 500 rads. It was shown that following exposure to 250 rads the number of chromosome aberrations in cells entering mitosis soon after irradiation (apparently G_2 cells) was much

greater than that in cells entering mitosis in the subsequent periods. A sharp decrease in the number of aberrant cells and chromosome aberrations was observed 72 and 96 hrs after irradiations. (auth)

45795 AGE-SEX SPECIFIC ACHILLES TENDON REFLEX TIME IN JAPANESE: A STUDY OF ATOMIC BOMB SURVIVORS, HIROSHIMA. Omori, Yoshiaki (Atomic Bomb Casualty Commission, Hiroshima); Ashley, Frantz W.; Belsky, Joseph L. J. Chron. Dis.; 25: 111-20(1972).

Achilles tendon reflex in man was studied with the aid of photomograph on 2100 (ages 18 to 89 yr) atomic bomb survivors and nonexposed controls of the Atomic Bomb Casualty Commission in Hiroshima to delineate effects of A-bomb radiation and aging. The reflex time was significantly prolonged, progressively, with age and it was more delayed in females than in males. The mean tap-to-half relaxation time was 290 ± 40 msec in males (mean age 46) and 321 ± 49 msec in females (mean age 46), which are similar to figures reported previously in Europe and the United States. In relation to radiation and obesity, no statistically significant difference in reflex time was found. These studies suggest that neurologic components of the reflex arc may be resistant to delayed effects of A-bomb radiation. Problems in interpreting reflex tracings and on differences in reflex time among races are discussed. Attention is called to age as an extrathyroidal factor in affecting achilles tendon reflex. (auth)

45796 RADIOBIOLOGICAL PROBLEMS CAUSED BY SUPERSONIC TRANSPORT. (WITH A SURVEY OF THE FIRST RESULTS ESTABLISHED BY TESTS PERFORMED ON BOARD THE CONCORDE PROTOTYPE). Delahaye, R. P.; Kaiser, R.; Pfister, A. (Nuclear Research Center, Strasbourg). Aerosp. Med.; 43: No. 7, 782-4(Jul 1972).

The present subsonic commercial aircraft flights do not present any major problems as far as radiobiology is concerned. The aircraft flying at altitudes higher than 15,000 m are subjected to irradiation which are different from those generally observed in the lower part of the atmosphere. A good many military pilots are already been subjected to this new environment. The entry in service of commercial supersonic transport aircraft (SST) will increase the number of exposed persons. At cruise altitudes (16,000 to 18,000) the screen constituted by terrestrial atmosphere and protecting the earth from cosmic radiation particles is considerably decreased, and it proved quite necessary to conduct a study on the possible radiobiological hazards sustained by SST passengers and crew. Results of radiation dose measurements made with nuclear emulsions and photographic film detectors located throughout the Concorde prototype are reported. (C.H.)

45797 RADIATION INJURY OF THE SPINAL CORD. Solheim, P. Oe. (Norske Radiumhospital, Oslo). Acta Radiol., Ther., Phys., Biol.; 10: No. 5, 474-80(Oct 1971).

Five cases of late radiation myelopathy are reported. Pareses and sensory disturbances progressed for some time after latent periods of from 5 to 19 months. Four of the five cases were then considerably improved. The clinical course of late radiation myelopathy thus seems not always to be chronic or progressive. (auth)

45798 IRRADIATION IN THE EPIDEMIOLOGY OF LEUKEMIA AMONG ADULTS. Gibson, R. (State Univ. of New York, Buffalo); Graham, S.; Lilienfeld, A.; (and others). J. Nat. Cancer Inst.; 48: No. 2, 301-11(Feb 1972).

The relation between leukemia and diagnostic medical irradiation was examined in 1,414 adult leukemia cases from Upstate New York and the Minneapolis and Baltimore metropolitan areas and 1,370 adult controls comprising a probability sample of the same areas from which the cases were drawn. Details of exposure by site and numbers of films were compiled from records of hospitals and physicians. The association of irradiation with leukemia was examined by the histologic type, sex, and age of the subject. Irradiation had an enhanced risk for males for both acute and chronic myeloid leukemia but not for females. The effect was more pronounced for irradiation to the trunk than to all other sites. Neither irradiated males nor females had an elevated risk for any other histologic type of leukemia. However, only a small proportion of the irradiated population develops leukemia, and irradiation accounts for only a small proportion of leukemia cases. (auth)

45799 CUMULATIVE DOSE FROM DIAGNOSTIC IRRADIATION. Norwood, W. D. (Hanford Environmental Health Foundation, Richland, Wash.); Rising, F. L.; Kirklin, C. W.; (and others). Amer. J. Roentgenol., Radium Ther. Nucl. Med.; 115: No. 3, 644-8(Jul 1972).

Using a relatively simple method, diagnostic exposures could be recorded as a unit indicating the dose in kilogram-rads, which

is proportional to the total energy received by the tissue. Diagnostic doses could then be made additive with other such procedures involving different parts of the body and with therapeutic, occupational, or natural irradiation doses. Diagnostic roentgen-ray units might be compared and accounting methods used to indicate where reductions in exposures might be made optimally. The physician would be helped in his decision to order or not to order a diagnostic procedure by having knowledge of the patient's accumulated lifetime dose from radiation. The results of the study indicate that when appropriately averaged over the whole-body mass, the diagnostic exposures from medical roentgen rays are not as large relative to occupational whole-body exposures as previously believed. (auth)

45800 PULL-THROUGH OPERATION FOR RADIATION-INDUCED RECTOVAGINAL FISTULA. REPORT OF A CASE. Thomford, Neil R.; Smith, David E.; Wilson, Wayne H. (Ohio State Univ., Columbus). Dis. Colon Rectum; 13: No. 6, 451-3(1970).

A patient in whom a large radiation-induced rectovaginal fistula was successfully repaired by a surgical operation is described. Normal bowel and vaginal function were restored. If repair of a rectovaginal fistula is indicated, and the fistula is large and located in the proximal vagina, the abdominal endorectal operation should be considered. (auth)

45801 HARD AND SOFT TISSUE NECROSES FOLLOWING RADIATION THERAPY FOR ORAL CANCER. Beumer III, John; Silverman, Sol Jr.; Benak, Steve B. (Univ. of California, San Francisco). J. Prost. Dent.; 27: No. 6, 640-4(Jun 1972).

Three hundred and fifty-four patients, who underwent radiation therapy for oral cancer (excluding lip) at the University of California Medical Center at San Francisco between 1961 and 1969, were reviewed as to site of tumor, mode of therapy, tumor dose, and occurrence of tissue necrosis. The study group was limited to those 278 patients who were available for follow-up for at least 6 months. Osteonecrosis occurred in 3.6%: 8 of 141 dentulous patients (5.7%) and 2 of 137 edentulous patients (1.5%). All cases of bone necrosis occurred in the mandible and were within the treatment field. There were 18 cases (6.5%) of soft tissue necrosis. Eleven of these patients were treated by needle implantation or intraoral cone technique. Fifteen were self limited (5.2%). (auth)

45802 STENOSIS OF THE SMALL BOWEL AS A COMPLICATION IN RADIATION THERAPY OF CARCINOMA OF THE UTERINE CERVIX. Joelsson, I.; Raef, L.; Soederberg, G. (Karolinska Sjukhuset, Stockholm. Serafimerlasarettet, Stockholm). Acta Radiol., Ther., Phys., Biol.; 10: No. 6, 593-604(Dec 1971).

Radiation-induced injury of the small bowel was observed in 15 of 3,075 cases from the Stockholm area, treated for carcinoma of the uterine cervix during 1954 to 1969. Factors that might increase the risk of this complication are discussed. Symptoms and signs of the intestinal complications are described and the results of surgical treatment presented. (auth)

45803 RADIATION PROTECTION CONSIDERED FROM THE MEDICAL VIEWPOINT. Ohlenschlaeger, L. Zentralbl. Arbeitsmed. Arbeitsschutz; 21: No. 9, 292-8(Sep 1971). (In German).

Following an introductory review of the interaction of ionizing radiation with irradiated material, and of the physical units adopted for radiation measurement, the author considers the maximum allowable doses fixed by German legislation and the medical supervision of persons occupationally exposed to radiation by means of clinical examinations and blood and urine analysis. He indicates a number of diseases disqualifying persons from work involving possible exposure to radiation. Particular attention is given to the problem of the medical supervision of personnel who are members of mobile teams employed in the maintenance and repair of nuclear reactors. (Occup. Saf. Health Abstr.)

45804 RADIOPROTECTION PRACTICE AT THE MARCOULE CENTER. Chassany, J. (Centre de Production du Plutonium de Marcoule, Bagnols-sur-Ceze, France). Rayonnements Ionis. Tech. Mes. Prot.; No. 3, 20-41(Jul 1971). (In French).

The Marcoule center consists of essentially four piles, a fuel reprocessing plant, their subsidiaries, a tritium extraction plant, and a plant for the reprocessing of irradiated fuels. A total of 2,078 personnel are employed, 1204 of which are directly concerned with work involving radiations. Fourteen years of experience have benefited various branches of study, the following being cited: development and improvements in efficiency of methods and means of protecting and controlling installations and personnel, environmental control, intervention, dosimetry, and protection of the respiratory tract. Experience demonstrates that the effectiveness of the means existing is very dependent upon the attitude of the individual and the psychological environment. (France)

45805 EFFECTS OF CEPHARANTHINE ON FLUCTUATION IN WHITE BLOOD CELL COUNT BY RADIOTHERAPY FOR GYNECOLOGIC CANCER. Ito, Haruhide; Hosaka, Koji; Toriumi, Tatsuo; Kubo, Sohei (Jikei Univ., Tokyo). *Shinryo To Shinyaku*; 8: No. 5, 191-4 (May 1971). (In Japanese).

In the radiotherapy of malignant tumor, since a sufficient dose is expected, various side effects, especially hemopoietic hypofunction, appear in almost all cases and as its countermeasures, improvement of the procedure of radiation, blood transfusion, and administration of various drugs are being performed. Following a single administration of cepharanthine, marked effects were observed in 14 (70%) of the 20 cases. In particular, an effect was found in 13 (81.3%) of the 16 cases administered with 50 mg of cepharanthine. This indicates that the administration of 50 mg of cepharanthine has a far greater significance in the prevention of side effect than that of 10mg. Comparison of the group administered with cepharanthine with groups administered with other drugs revealed no significant difference, but this result included the administration of both 10mg and 50 mg, and more marked effect was observed in the group administered with 50 mg of cepharanthine than groups administered with other drugs. It is thought that the above results were obtained by the synergistic actions of reticuloendothelial hyperfunction and activation of the hemopoietic function of the bone marrow or liver function by cepharanthine on hemopoietic hypofunction particularly observed in the cancer-bearing organism. It was thought that leukocytopenia interfering with radiotherapy clinically could be avoided by the single administration of cepharanthine, especially the dose of 50mg. (Japan)

45806 SUPERSONIC RADIATION RISKS. Brit. Med. J.; 3: No. 5819, 130 (15 Jul 1972).

Literature is briefly reviewed and editorial comment made on estimates of doses received by passengers in supersonic flights. A representative dose for passengers at 17,500 m is quoted as 0.6 to 1 mRem/hr and this is related to estimated doses of radiation received from other sources. The need for adequate solar flare detection is mentioned, and it is pointed out that in 1956 a flare gave a calculated dose rate of 2 rads/hr at 18,000 m. (UK)

45807 RADIATION MENOPAUSE. Doherty, R. Brit. Med. J.; 3: No. 5819, 176 (15 Jul 1972).

A brief summary is made of a 1962 survey of the results of the use of radiation to induce menopause in female patients at the Plymouth Hospitals from 1948 to 1957. During this period 678 patients were treated with deep x-ray doses of between 850 and 900 rads, to the midpoint of the pelvis. Mention is made of side effects, effectiveness of treatment in inducing a menopause in dysfunctional bleeding, subsequent treatment, and later mortality. (UK)

45808 RADIATION PROTECTION MEASUREMENT. Kiefer, H.; Maushart, R. Oxford: Pergamon Press (1972). 577p.

Translated into English by R. Friese.

The subject is considered under the headings: introduction (development of radiation protection measurement techniques in the general field of radiation protection, radiation protection measurement in nuclear engineering, subdivision into special fields), dose determination, local dose monitoring, personal dose monitoring, activity determination (general remarks, measurement of activity in water, measurement of activity in air, measurement of activity in food and other biological substances, measurement external contamination, measurement of internal contamination), radionuclide determination survey of methods, physical methods; determination of the type of radiation, determination of half-life, determination of radiation energy, radiochemical methods, determination of specific radionuclides (tritium and low-energy β -emitters, iodine, radium and radon, uranium) measurements for the protection of the population (environmental survey of nuclear facilities, measurement after accidents in nuclear facilities, measurement in the case of a nuclear war) and trends in development (semiconductor detectors, problems of radiation measurement in space flight). (UK)

45809 HEPATIC INJURIES FOLLOWING IRRADIATION. Sakamoto, Yoshio; Kasuga, Toshio; Ohata, Takeo; Watanabe, Toshikazu; Nakanishi, Fumiko (Shinshu Univ., Nagano, Japan). *Nippon Igaku Hoshasen Gakkai Zasshi*; 30: No. 11, 120-30 (Feb 1971). (In Japanese).

In the telecobalt therapy to malignant diseases of the neighboring organs of the liver, a part of the liver was often included in the irradiated fields, and hepatic changes were found in these irradiated areas. These hepatic changes were investigated by liver scanning, isotopic liver function tests, and liver histopathologic examination and the mechanism of radiation hepatitis was discussed. In the early period of hepatic irradiation, the function of liver parenchymal cells increased temporarily or congestion of liver acini around the central veins occurred in the irradiated areas, therefore liver blood flow increased. Gold-198 colloid liver scanning in this

period suggested swelling of the liver in the irradiated area. When hepatic irradiation was continued or with the passing of time after completion of the irradiation, the liver blood flow decreased and the function of liver parenchymal cells, particularly of Kupffer's cells fell. As circulatory insufficiency of the liver following irradiation became severe, destruction and disappearance of the liver parenchymal cells and hemorrhage of liver acini began from the central portion of liver lobules and fibrosis surrounding central veins became remarkable. The reduction of photodensity in the irradiated area of the liver occurred in photoscintigram after irradiation less than 4,000 R or one month after beginning of radiation therapy. (auth)

45810 TISSUE MAST CELLS IN HUMAN BLOOD. PRE-LIMINARY REPORT. Csaba, G.; Hodinka, L.; Bozsoky, S.; Farkas, K. (Simmelweis Univ., Budapest. National Inst. of Rheumatology and Physiotherapy, Budapest). *Haematologia* (Budapest); 4: No. 3-4, 415-17 (1970).

A total of 125 blood smears from 39 patients suffering from autoimmune diseases, degenerative processes of the locomotor system and tumors, and those of 10 healthy subjects were examined. Mast cells were counted in 1500 microscopic fields of each smear under 600-fold magnification. Mast cells were present in the blood of each examined subject, whether healthy or ill (98% of the cells were of lymphoid character and similar in size to the erythrocytes). In different diseases the blood mast cell count varied. The results showed an increase in the number of blood mast cells in autoimmune diseases involving the thymus, as compared to the controls or other subjects suffering from degenerative diseases. In contrast, in subjects with tumor the mast cell count displayed a decreasing trend. Mediastinal x irradiation, by affecting the thymus, was found to influence the blood mast cell count as well. The eventual diagnostic importance of the blood mast cell count is indicated by the findings. However, in order to evaluate correctly the shifts, further examinations of a large patient material are needed. (auth)

45811 WORKLOADS AND USE FACTORS IN DIAGNOSTIC X-RAY INSTALLATIONS. Campbell, E. M.; Essenburg, A.; Holloway, A. F. (Manitoba Cancer Treatment and Research Foundation, Winnipeg). *Health Phys.*; 23: No. 1, 67-71 (Jul 1972).

A report is presented on the magnitude and distribution of workloads observed during the course of radiation surveys in diagnostic x-ray departments in hospitals in the Province of Manitoba. The data show that the magnitudes of the majority of these workloads were considerably lower than those quoted in tables relating to protective barrier requirements, and that the distribution of the primary beam workloads about an installation was such that suitable use factors deviated substantially from those recommended in the NCRP Report No. 34 and the Report of the Medical X-Ray Advisory Committee on Public Health Considerations in Medical Diagnostic Radiology (X-Rays). (auth)

45812 RADIATION HAZARDS TO MAN IN SPACEFLIGHTS. Altkofer, Otto Claus; Heinrich, Wolfgang; Simon, Manfred (Univ., Kiel). *Umschau*; 72: No. 9, 284-7 (May 1972). (In German).

The radiation hazards to astronauts in manned space flights by cosmic radiation are surveyed. In a short review the fundamental facts are reviewed and the basic principles of dosimetric units are described. The radiation effects due to the different components of cosmic radiation are discussed showing that the galactic and solar radiation, and the radiation belts are contributing in a different manner to the radiation hazards. Although the abundance of the heavy ions is relatively small, the radiation effects have to be evaluated separately due to the high ionization along the particle tracks. The possible radiation injuries are enumerated. (auth)

Vertebrates

Refer also to abstract: 45733.

45813 (COO-1427-41) CELLULAR PROLIFERATION AND REGENERATION FOLLOWING TISSUE DAMAGE. Harding, Clifford V. (Oakland Univ., Rochester, Mich. Dept. of Biological Sciences). Feb 1972. Contract AT(11-1)-1427. 31p. Dep. NTIS.

Results obtained during the past seven year period on radiation effects on eyes of rabbits, are summarized. Mechanical injury to the lens, in vivo, initiates the cell cycle and is accompanied by propagated waves of DNA synthesis and mitosis that move out slowly from the site of injury. The injury response, and the effect of x irradiation on the propagated injury response and wound healing was determined. Biochemical and microscopic methods used in the studies are described. A determination of the in vitro culture conditions that bring about a reproducible stimulation of DNA syn-

thesis and mitosis in the cultured mammalian lens was accomplished. It was found that the lens in culture provides an excellent model system for studies on the response of tissue to injury under carefully controlled and defined conditions. (TSS)

45814 (COO-3017-43) RADIATION CARCINOGENESIS. Progress Report VI, May 1, 1971-April 30, 1972. Warren, Shields; Gates, Olive (New England Deaconess Hospital, Boston, Mass. Cancer Research Inst.). 30 Apr 1972. Contract AT(11-1)-3017. 29p. Dep. NTIS.

Investigations have been continued during this period on the following projects: Studies of the extraordinarily high incidence of cancer, including cancer developing in unusual sites, in supra-lethally irradiated parabiont rats. The incidence has continued appreciably higher than in single rats irradiated at a less degree. One unusual tumor developed is the sarcoma that appears in the anastomotic line. Detailed studies of this tumor have been carried out in the course of the year. Special studies have been made of the induction of osteogenic sarcoma and uterine carcinoma in parabiont and single rats using focal internal radiation from ^{60}Co γ sources. Pathologic and pathophysiologic studies of the transplantable pheochromocytoma mentioned last year are continuing. Studies on the effects of perinatal radiation on mice so far as induction of leukemia and solid tumors is concerned. Variation in strain and life period of radiation produce important differences in the neoplastic response. Study of the mice intensively inbred for a number of generations following radiation. Completion of the study of disordered calcium metabolism mentioned in the last report. (auth)

45815 (COO-3486-1) EFFECTS OF IONIZING RADIATION ON SALAMANDER ORIENTATION. Progress Report. Shoop, C. Robert (Rhode Island Univ., Kingston (USA). Inst. of Environmental Biology). 1 May 1972. Contract AT(11-1)-3486. 19p. Dep. NTIS.

Two species of salamanders, *Ambystoma maculatum* and *A. opacum*, were used in field studies in western Rhode Island. Ionizing radiation was administered to males for studies on effects of radiation on migration. Effects of gamma radiation on salamanders and other amphibians were investigated with regard to species differences, temperature effects, and dose rate effects. No increase in death rate occurred over a period of one year with doses of 25 to 50 Rad. It was found that radioresistance corresponds inversely to relative liver-cell nuclear DNA levels of the species involved. Studies on migration of females were conducted using ^{181}Ta and ^{182}Ta tags; determination of dose from the tags varied with placement of dosimeters in the body. Studies were conducted on uptake and turnover of radioactive sodium, cesium, praseodymium, and europium by larval and hatchling amphibians and reptiles. (H.L.W.)

45816 (FRNC-R-7) BIOLOGICAL EFFECTS OF THE INTENSITY OF IRRADIATION. Legeay (Laboratoire Central de l'Armement, Arcueil (France)). 17 Dec 1971. 58p. (In French). Dep. NTIS (U. S. Sales Only).

The biological effects of the time distribution of the absorbed energy, i.e. the intensity of irradiation, were investigated in various mammals. Before studying the recovery from a total and continuous irradiation, the radiosensitivity of various mammals (such as mice, rats, dogs, monkeys, guinea pigs, sheep, donkeys) was determined by measuring their mean lethal doses. The effects of the intensity of irradiation for low and high LET radiations were evaluated. The information obtained on cell and tissue recovery kinetics is summarized. Recovery models are proposed. The effects of the intensity of irradiation are estimated. (France)

45817 (ORO-2690-12(Pt.B), Paper 12) STUDIES IN THE USES OF IONIZING RADIATIONS AS STIMULI. Smith, James C. 1p.

Progress is reported on studies on conditions under which gamma rays can be used as stimuli to elicit reactions in a variety of animals. The role of visual gamma ray detection in rats and monkeys was studied and it was found that gamma ray detection is possible in dark adapted animals when they have been deprived of the sense of smell. It was demonstrated that gamma rays acted as aversive stimuli when paired with a novel taste solution; aversion of that solution resulted. Comparisons were made of necessary conditions for developing aversions when using gamma rays and lithium chloride. The development of aversions was studied by a computer analysis of licking behavior. (H.L.W.)

45818 (UCLA-12-867) IRRADIATION AND VIRAL LYMPHOMAGENESIS IN THYMIC GRAFTS. Hays, Esther F. (California Univ., Los Angeles. Lab. of Nuclear Medicine and Radiation Biology). 1971. Contract AT(04-1)-Gen-12. 19p. Dep. NTIS.

The thymus of AKR neonatal mice was subjected to 850 and 5000 R ^{60}Co irradiation in vivo prior to its removal for grafting to thymectomized syngeneic hosts. These irradiated grafts placed in hosts inoculated as sucklings with Gross murine leukemia virus (GMuLV) developed lymphoma. The incidence was slightly lower and the latent period longer than that found in irradiated control grafts in similar hosts. When grafts were exposed to GMuLV in vitro just prior to grafting or in vivo 24 hr before grafting, but placed in hosts not given neonatal virus inoculation, lymphoma developed with reduced incidence in the 850 R irradiated grafts and did not develop in the grafts receiving 5000 R. A dose of 5000 R was not found to alter the lymphomagenic properties of filtrates containing GMuLV. These studies show that heavily irradiated thymic grafts can develop lymphoma in virus infected hosts, but that irradiation of grafts exposed to virus in vitro and in vivo inhibits the ability of GMuLV to produce lymphoma probably by reducing the numbers of cells available for virus replication, thus, indirectly lowering virus concentration in the thymic grafts. (auth)

45819 (UR-3490-63) IONIZING RADIATION-INDUCED THYROID CARCINOGENESIS IN THE DOG. Michaelson, S. M.; Lu, Shin-tsu; Quinlan, W. J. Jr. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). [nd]. 16p. (CONF-720505-3). Dep. NTIS.

From twelfth Hanford biology symposium on radionuclide carcinogenesis; Richland, WA. (11 May 1972).

The response of the thyroid gland of the dog to x ray or ^{131}I , either one alone or in combination, manifests itself in a temporal manner, functionally, by transient increased activity and progressive hypothyroidism and pathologically, by degeneration, hyperplasia, and carcinogenesis. Eleven cases of thyroid tumor were observed among 23 dogs that received external, internal radiation, or both survived 3 yr post exposure. One of four dogs received ^{131}I alone revealed bilateral follicular adenocarcinoma with anaplastic changes. This animal received ^{131}I in doses ranging from 17 to 50 μCi over a 5 yr-period. The calculated radiation exposure was 1580 rads. The animal died at age 13 yr 10 months or 5 yr 8 months after the initial dose of ^{131}I . Radioiodine uptake was markedly depressed corresponding to the non-differentiated nature of the neoplasm. The functional and histopathologic data on this dog is compared with other dogs that received either ^{131}I and/or x ray. Two of three dogs exposed to x irradiation alone had follicular adenoma and eight of 16 dogs that received ^{131}I in addition to x irradiation showed various types of thyroid tumors. The thyroid gland neoplasia seems to be more related to aging than to the onset of puberty as observed in man. Vascular alteration may initiate and amplify the visible and functional thyroid derangement. Increase in circulating TSH appears to be a basic factor in the sequence of events leading to permanent hyperplasia or carcinogenesis. The post-irradiation latency for induction of thyroid neoplasia in the dog (4 to 10 yr), man (9 to 14 yr), and the rat (1 to 2 yr) shows a linear relationship with basal metabolic rate. Although this relationship may be only fortuitous, it may be indicative of a fundamental factor related to tissue metabolism and/or thyroid cell turnover rate. (auth)

45820 (UR-3490-121) EFFECT OF RAT TUMOR ALLOGRAFTS ON THE DISTRIBUTION OF PASSIVELY TRANSFERRED ALLOANTIBODIES STUDIED BY HEMAGGLUTINATION AND ISOTOPIC TRACER TECHNIQUES. Buchsbaum, Donald Jay (Rochester Univ., N. Y.). 1972. 110p. Dep. NTIS.

Thesis.

The experimental studies reported here are concerned with transplantation antibodies produced in Buffalo strain rats following immunization with a tumor induced in a Fischer strain rat with methylcholanthrene. It is shown, with a hemagglutination technique for antibody titration, that when such antibodies are passively transferred to immunosuppressed Buffalo rats carrying this same Fischer tumor, the antibody titer in the blood declines more rapidly than in control Buffalo rats not carrying the tumor. A method was developed for obtaining radiolabeled transplantation antibody from immune Buffalo gamma globulin. When this labeled antibody was injected into Buffalo rats carrying the Fischer tumor, substantial preferential localization of radioactivity occurred in the tumor. With a few exceptions, serum taken from x ray and cortisone treated Buffalo rats bearing the Fischer tumor had hemagglutination titers lower by a factor of four or more than serum from control rats without tumors 3 or 4 days after immune Buffalo gamma globulin injection. Serum samples taken from tumor and control rats 6 hrs after injection appeared to have the same antibody titers. Radiolabeled transplantation antibodies were isolated from immune Buffalo gamma globulin by a twice repeated adsorption-elution procedure starting with Fischer red blood cells. The percentage of ^{125}I -antibody and ^{131}I -normal Buffalo gamma globulin present in eight growing

tumors 48 hrs after injection of the paired label mixture ranged from 1.6 to 9.3% and 0.4 to 1.1%, respectively. In general, the lower ^{125}I values were obtained with tumors weighing less than 0.4 grams, whereas the values were greater than 6% for tumors weighing 0.9 to 1.6 grams. The corresponding values for four growing tumors 96 hrs after injection were 1.9 to 4.0% and 0.3 to 1.5%. The average ratio of the fraction of the injected dose of antibody and normal gamma globulin was approximately one in all tissues except tumor, where the ^{125}I was more abundant by a factor of eight, 48 hrs after injection; and by a factor of six, 96 hrs after injection. (auth)

45821 CHOLERHEIC ENTEROPATHY AS A CAUSE OF DIARRHEA AND DEATH IN RADIATION ENTERITIS AND ITS PREVENTION WITH CHOLESTYRAMINE. Berk, Robert N.; Seay, David G. (Univ. of California, La Jolla, San Diego). *Radiology*; 104: No. 1, 153-6(Jul 1972).

Cholerheic enteropathy, a syndrome of diarrhea due to the effects of excessive bile salts in the colon, occurs when bile salt absorption from the terminal ileum is impaired. Cholestyramine, a nonabsorbable anion-exchange resin, prevents diarrhea by binding the bile salts, thus rendering them inactive. Evidence that cholerheic enteropathy has a significant role in the pathogenesis of diarrhea and death from radiation enteritis is reviewed, and experiments with rats treated with cholestyramine prior to abdominal irradiation are described. (auth)

45822 LYMPHOID-CELL RENEWAL UNDER LOW-LEVEL IRRADIATION. XI. THE CELL CYCLE AND GROWTH KINETICS DURING RADIATION LEUKEMOGENESIS IN C57BL MICE. Fabrikant, Jacob I.; Foster, Bessie R. (Univ. of Connecticut, Hartford). *Radiology*; 104: No. 1, 203-4(Jul 1972).

Studies of thymus-cell population kinetics suggest that during leukemogenesis, changes in the control of cell growth and division occur in cells undergoing differentiation and in those that are functionally mature as well as in proliferating cells, thus disturbing the rate of cell flux and transition from one phase of cell maturation to another. Progressive breakdown of cellular control mechanisms may be involved. Cell and tissue kinetics in the thymus are described in relation to the changing pattern of the lymphoid-cell cycle and growth fraction during the development of radiation-induced murine thymic lymphoma and disseminated leukemia. (auth)

45823 EFFECT OF RADIOPROTECTIVE PREPARATIONS ON THE RESPIRATORY EXCHANGE OF RATS. Novak, L.; Fedorov, B. A.; Gusek, B.; Semenov, L. F. (Inst. of Experimental Pathology and Therapeutics, Sukhumi, USSR. Inst. for Biophysics, Brno). *Radiobiol.-Radiother.*; 12: No. 6, 751-6(1971). (In German).

In experiments with 72 unirradiated white rats the influence of thiol-radioprotectors (cystamine, MEA, thiocarbamide, acetylcholine) on the adaptive respiratory exchange was examined and measured over the course of three hours. The sulfur preparations did not cause a decreased consumption of oxygen. Adrenaline and acetylcholine increased the consumption of O_2 ; indolylalkylamines decreased it. Histamine did not affect the respiratory exchange. The effect of the protectors is not caused by their influence on the oxygen consumption of the animal organism. (auth)

45824 INFLUENCE OF AET ON CERTAIN IMMUNOLOGIC PROCESSES IN RABBITS IRRADIATED BY RADIOACTIVE COBALT. Gidaly, M.; Gorgan, V.; Bilcea, M.; Uray, Z. (Inst. of Medicine, Cluj. Inst. of Oncology, Cluj). *Aggressologie*; 12: No. 4, 269-73(Apr 1971). (In French).

A study was made of the protective effect of AET on the immunologic capacity of rabbits irradiated with 400 Rad of ^{60}Co γ and vaccinated with *Salmonella typhimurium*. The doses used have a strong immunosuppressive effect on the formation of agglutinins as well as on phagocytosis. AET has a pronounced radioprotective effect on phagocytosis but an undetectable effect on the formation of agglutinins. (tr-auth)

45825 EFFECTS OF X-IRRADIATION ON OVARIAN FUNCTION IN THE PREGNANT HAMSTER. Greenwald, Gilbert S. (Univ. of Kansas, Kansas City). *Endocrinology*; 91: No. 1, 75-86(Jul 1972).

Whether antral follicles are essential for luteal progesterone secretion in the hamster was assessed by eradicating follicles in an exteriorized ovary with 3000 rads on day one of pregnancy. Various parameters were compared between the irradiated (I) ovary and non-irradiated (N) ovary at different stages of gestation. When both ovaries were present, the N ovary showed an excessive increase in weight because of compensatory follicular development and the growth of corpora lutea (CL) at their normal rate. In contrast, the I ovary was reduced in size because of the stunted growth of the CL and the absence of antral and large multilayered follicles. Despite their different size in the second half of pregnancy, the CL

and the I and N ovaries had the same content of progesterone, suggesting that under these circumstances, size per se of the CL did not reflect their functional capacity to produce the steroid. When the N ovary was removed at day 10 of pregnancy, the I CL commenced to grow and almost doubled their weight by day 16. The increased weight of the I CL occurred regardless of whether the semispayed hamsters were injected with estrone or the oil vehicle and was reflected in a significantly higher luteal content of progesterone than either the N or I ovaries when they were both in situ at day 16 of pregnancy. In hamsters, bilaterally ovariectomized at day 4 of pregnancy, it was necessary to inject estrogen as well as progesterone to maintain the optimal weight of embryonic swellings at day 8. In a group of irradiated hamsters, removal of the N ovary at day 4 resulted in reduced weight of embryonic swellings at day 8, comparable to the effects of injecting progesterone alone in the bilaterally ovariectomized animals. Thus, the I ovary (lacking antral follicles) secreted progesterone but none or insufficient estrogen to produce optimal decidualization. When the semispayed hamsters were given a single injection of estrogen, the embryonic swellings were restored to normal size. In animals bilaterally ovariectomized at day 10 of pregnancy, between 0.5 and 1 μg of estrone plus 1 mg progesterone was required on a daily basis to maintain pregnancy in all animals. When the N ovary was removed from irradiated animals on day 10 of pregnancy, pregnancy was terminated unless estrone was injected or fetuses were exteriorized. When the pregnant cornu was removed along with the N ovary, estrone was ineffective in maintaining the CL in the I ovary. This indicates that the effect of estrogen was not mediated directly on the CL. Collectively, these experiments demonstrate that antral follicles are the major (if not the only) source of estrogen in the pregnant hamster but that, in their absence, progesterone is still secreted by the CL. The essentiality of estrogen for the maintenance of pregnancy in the hamster is at the uterine level for interactions with progesterone. (auth)

45826 EFFECTS OF PARTIAL BODY IRRADIATION OF MICE ON THE QUANTITY OF 5-HYDROXYINDOL ACETIC ACID IN THE URINE, THE QUANTITY OF WATER CONSUMED, AND THE VOLUME OF URINE EXCRETED. Nakamura, W.; Nishimoto, Y. (National Inst. of Radiological Sciences, Chiba, Japan). *Compt. Rend. Soc. Biol.*; 165: No. 6, 1497-1501(1971). (In French).

Total, partial, or simulated irradiation produces variations in the following three biological criteria in mice: the quantity of 5-hydroxyindol acetic acid in the urine, the quantity of water consumed, and the volume of urine excreted. A 600R irradiation of the head accentuates these variations. A total body irradiation significantly reduces the curves in relation to these criteria. Protection of the head ensures a return to the initial values. (France)

45827 EFFECT OF RADIATION ON THE REGENERATION OF MINCED AUTO-TRANSPLANT MUSCULAR TISSUES OF THREE-WEEK-OLD RATS. Bulyakova, N. V. *Dokl. Akad. Nauk SSSR*; 202: No. 3, 729-31(21 Jan 1972). (In Russian).

The process of muscular tissue regeneration following trauma (crushing) immediately and one to four months after local irradiation at 2,000 R was investigated to ascertain if the capacity of the irradiated muscle was restored to a myoblastic mode of regeneration. Four sets of experiments were conducted using different irradiation schedules. It was shown that local irradiation of the muscles of three-week old rats with 2,000 R suppressed the process of post-traumatic regeneration as demonstrated by the formation of granular tissue, myoblasts, and myosmyoplasts. These observations agreed with those of other workers on fully-grown rats. (K.S.W.)

45828 CHANGES IN THE LOCALIZATION OF ACETYLCHOLINESTERASE IN NEUROMUSCULAR JUNCTION AFTER ^{60}Co IRRADIATION. Faes, M. H.; Aubert, C.; Rosselle, N.; van der Schueren, G. (CEN, Mol, Belg.). *Arch. Int. Physiol. Biochim.*; 79: 269-77(1971). (In French).

The posterior appendages of 30 frogs (*Rana esculenta*) were exposed to 10, 20, 40, and 100 kR of ^{60}Co gamma radiation and the animals were killed at intervals up to 29 days after irradiation. The sartorius muscle was removed from both legs of each frog and prepared for histochemical study. Methods for identifying and localizing acetylcholinesterase are described and macro- and microscopic changes in the muscle and neuromuscular junction are discussed. It was found that radiation modified the localization and the microscopic structure of the supporting elements of the acetylcholinesterase. The junction did not appear to be more radiosensitive than the nerve or muscle. Alterations in the motor plate were accompanied by degeneration of muscle fibers. (H.L.W.)

45829 CHANGE OF CONTENT OF SULFHYDRYL GROUPS IN RAT HEART AND LIVER MITOCHONDRIA AND SUPERNATANT AFTER HIGH ENERGY IRRADIATION. Pegel, V. A.; Dokshina, G. A.; Dement'eva, T. A. (Scientific-Research Inst.

of Biology and Biophysics, Tomsk, USSR). *Izv. Sib. Otd. Akad. Nauk SSSR*, No. 5, Ser. Biol. Nauk; No. 1, 108-13 (Apr 1971). (In Russian).

The contents of thiol groups (total, protein, non-protein) were analyzed at 10, 30 min, 1, 2, 4, 6, 24, 48, and 72 hrs after irradiation (betatron 25 MeV) at a dose of 1000 rad in rat liver and heart mitochondria. A decrease in SH-group content in rat liver and heart mitochondria caused by irradiation was determined to proceed mainly on account of protein thiols. The non-protein sulphhydryl groups changed insignificantly. Decrease in titre of thiol groups in mitochondria was accompanied by its increase in supernatant. The latter indicates textural disturbance of subcellular fractions in the course of irradiation. (auth)

45830 REGENERATION OF LYMPHOID BONE MARROW IN THE THYMECTOMIZED ADULT MOUSE SUBJECTED TO A SUBLETHAL DOSE OF X RAYS. Schmitz, A.; Haot, J. (Univ., Liege). *Compt. Rend. Soc. Biol.*; 165: No. 6, 1467-9 (1971). (In French).

Young mice were subjected to a 400 R x ray dose one month after they had been thymectomized. The absence of the thymus did not prevent the appearance of a large quantity of lymphoid cells in the bone marrow. The importance of this sudden production of medulla lymphoid cells in thymectomized animals suggests the possible regulatory function of the thymus in the regeneration of the bone marrow lymphoid cells after sublethal irradiation. (France)

45831 EFFECTS OF THYROIDIAN HORMONES AND γ IRRADIATION ON THE ADENYLCYCLASE OF THE PLASMA MEMBRANES OF THE RAT LIVER. Marc, J.-C.; Junien, J.-L.; Rocquet, G.; Michel, R. (Centre de Recherches du Service de Sante des Armees, Clamart, France. Univ., Paris). *Compt. Rend. Soc. Biol.*; 165: No. 6, 1257-62 (1971). (In French).

The adenylylase system of the liver plasma membranes is activated by several substances, and in particular by thyroid hormones. It is known that γ radiation modifies the functions of certain biological molecules. This fact was utilized in locating the adenylylase activation sites in the presence of NaF and glucagon, and in studying the behavior of 5' nucleotidase. After the first hour of irradiation (25740 rads) the effect of the glucagon is considerably reduced and after two hours of irradiation (51480) rads the effect was totally inhibited; NaF, however, retains a stimulating effect after two hours. The nucleotidase is not significantly modified. (France)

45832 PHENOL RED UPTAKE BY IRRADIATED AND NON-IRRADIATED CHICK MESONEPHROS IN VITRO. de Ridder, Leo I.; Mareel, Marc M. (State Univ., Ghent). *Lab. Invest.*; 26: No. 5, 600-3 (1972).

Ionizing irradiation has a stimulating effect on phenol red uptake in the proximal tubules of the 9-day chick mesonephros. The uptake and the radiobiologic effect on it are temperature-dependent. The latent period between irradiation and effect is only 30 minutes as shown by liquid scintillation counting. The selective uptake in the proximal tubules can be demonstrated by the autoradiographic technique for water-diffusible compounds. However, by this morphologic method, no quantitative difference can be seen between irradiated and control cultures. (auth)

45833 EFFECTS OF X IRRADIATION UPON THE RESPONSE OF MONONUCLEAR PHAGOCYTES TO AN INDUCED STERILE INTRAPERITONEAL INFLAMMATION. Harper, John R.; Wolf, Norman S. (Univ. of Washington, Seattle). *Res. J. Reticuloendothel. Soc.*; 11: No. 4, 368-82 (1972).

The radiosensitivities of the populations of mononuclear phagocytic cells (and their precursors) which respond to intraperitoneal stimulation with thioglycollate broth were examined in the mouse. X irradiation (500 to 1000 R) given 24 hours previous to the stimulation reduced the number of such cells collected from the peritoneal cavity four days poststimulation to the level of unstimulated animals. Similar irradiation given 24 hours poststimulation had no or little effect under otherwise similar circumstances. With time represented on the abscissa and cell numbers collected on the ordinate, progressively shortened poststimulus irradiation times resulted in a falling asymptotic curve through the coincidence point of stimulation and irradiation times. Intraperitoneal mitosis did not account for cellular accrual. Two populations of cells, based on radiosensitivity and the mitotic event, are suggested on the basis of the relationships of the numbers of cells collected at varying times before and after irradiation and known data for post-irradiation mitosis among cell populations. (auth)

45834 SUPPRESSION AND RECOVERY OF COLONY FORMING UNITS IN THE RAT FOLLOWING SUB-LETHAL X IRRADIATION. Okunewick, James P.; Phillips, Eleanore L. (Allegheny General Hospital, Pittsburgh). *Contract AT(11-1)-3097. Exp. Hematol.*; 22: No. 1, 26-7 (Mar 1972).

Rats were exposed to a single whole-body 200 R dose of x rays and were killed at selected intervals following irradiation. Marrow was removed from the excised femurs and cell suspensions were prepared. The cells were then injected into irradiated mice, the spleens excised, fixed, and the colonies counted. Graphs are presented to show results. The data indicated that recovery of the total nucleated marrow proceeds at a faster rate than recovery of transplantable colony-forming units. The colonies developed normally after transplantation regardless of when the donor cells were taken from rats. The prolonged repression of the CFU compartment suggested that factors other than residual effects of radiation act on the bone marrow CFUs after low level x radiation exposure. (H.L.W.)

45835 VASCULAR ALTERATION IN THE IRRADIATED BONE MARROW OF THE RABBIT MANDIBLE. A MICROANGIOGRAPHIC STUDY. Mori, Ryoichi (Kanazawa Univ., Japan). *Kanazawa Irigaku Soshu*; 89: 51-69 (Jun 1971). (In Japanese).

The mandible bones of mature rabbits were exposed to repeated ^{60}Co γ ray doses. The total irradiated doses of five groups were 3,000 R, 5,000 R, 7,000 R, 9,000 R and 12,000 R, respectively. The morphological changes of irradiated minute blood vessels were observed microangiographically, and the pathological changes of blood vessels and marrow tissues were investigated histologically. Dilations and constrictions of minute arterial vessels were observed under 5,000 R irradiation. The capillaries were constricted and were represented by only fine threads at irradiation over 5,000 R. The venous sinus showed a tendency to dilatation at radiation doses over 7,000 R. Dilatations, constrictions, and the presence of thrombi in the veins were investigated under conditions of over 7,000 R irradiation. Edema of the mandibular marrow occurred at doses over 7,000 R. Reticular cells and fibers increased in proportion to the dose. A comparison of radiation effects on blood vessels and bone marrow showed that changes appeared in blood vessels at doses under 5,000 R, but that changes in bone marrow appeared only at doses over 7,000 R. (H.L.W.)

45836 EFFECT OF IONIZING RADIATION ON ENZYME ACTIVITY IN THE NEUROGLIA OF THE DEVELOPING RABBIT BRAIN. Wender, M.; Kozik, M.; Owsiankowski, M. (Medical Academy, Poznan, Poland). *Folia Histochem. Cytochem.* (Krawkow); 9: 299-304 (1971).

The activity of acid and alkaline phosphatase, thiamine pyrophosphatase, adenosine triphosphatase, non-specific esterase, acetylcholinesterase, and butyrylthiocholinesterase was examined in the developing neuroglia of the rabbit brain exposed to a single dose of 150 R of x-rays on the 18th day of fetal life. The investigation led to the following conclusions: The results of histochemical examinations of the activities of acid and alkaline phosphatases, TPPase, and ATPase in the irradiated animals do not differ perceptibly from those obtained for normal development. The activity of butyrylthiocholinesterase in the myelinated fibres of the animals irradiated in fetal life does not vary from that observed for physiological conditions. On the other hand, no increase in the activity of this enzyme can be recorded in the process of maturation of the interfascicular oligodendroglia. This divergence becomes very distinct as soon as the myelinization process is concluded. The activity of butyrylthiocholinesterase in the interfascicular oligodendroglia does not play any substantial role in the metabolism of the myelin components. (NSA of Poland)

45837 ROLE OF TRANSPLANTED SLOWLY PROLIFERATING BONE MARROW CELLS FOR REGENERATION OF LETHALLY X-IRRADIATED RAT BONE MARROW. Haas, R. J.; Meyer-Hamme, K.; Flidner, T. M.; (and others) (Univ. of Ulm, Ger.). *Scand. J. Haematol.*; 9: No. 2, 121-9 (Apr 1972).

Inbred Lewis rats were labeled with ^3H -thymidine (^3H -TdR) in utero and for 6 weeks after birth to obtain 100% labeling of all the cells in the animals. Six weeks after the last ^3H -TdR injection only cytogenetically resting cells were still labeled. These cells comprised of two types of reticular cells, endothelial cells and a small fraction of bone marrow lymphocytes. At this time the animals were treated with 4×500 mg/kg body weight hydroxyurea at 6 hrs intervals. A profound marrow hypoplasia was obvious 6 hrs after the last HU injection. All proliferating precursor cells of the bone marrow were destroyed, whereas the cytogenetically resting cells were not changed in their absolute number. Successful transplantation of the marrow treated with HU to 1200 R x-irradiated recipients provided evidence for the presence of stem cells in the bone marrow of the donors and indicated that in rats, bone marrow regeneration can be achieved in the absence of cells morphologically defined as hemocytoblasts. (auth)

45838 EFFECTS OF IONIZING RADIATION ON THE STRUCTURE AND METABOLISM OF MITOCHONDRIA IN RAT LIVER CELLS. Zyss, Ryszard; Michalska-Kaszczynska, Maria

(Inst. of Postgraduate Medical Education, Warsaw). *Acta Med. Pol.*; 13: No. 2, 161-75(1972).

Using ultrastructural, cytochemical, and biochemical methods the character and intensity of morphological and structural changes in liver cell mitochondria of rats receiving a whole-body radiation dose of 1000 R were studied. It was found that swelling of varying intensity is the main form of morphological mitochondrial damage. With marked swelling, the matrix of the mitochondria and their membranes are damaged with disappearance of cristae and with focal disruption of membranes. The damaged mitochondria undergo sequestration—they are enveloped with membranes of the endoplasmic reticulum with development of the so-called cytoesosomes. Ultrastructural changes of mitochondria were associated with changes in their enzymatic activity, with a fall in the activity of succinic dehydrogenase and with a rise in the activity of ATP-ase. This suggests that structural damage to the mitochondria varies in its influence on the direction of changes in their enzymatic activity with the activity of ATP-ase being more dependent on the structural damage of mitochondria than the activity of succinic dehydrogenase. Explaining the mechanism of development of radiation-induced disturbances in the structure and functions of mitochondria, the authors consider that, most probably, these disturbances are due to changes in the physico-chemical characteristics of lipoprotein membranes caused by radiation. Damage to the lipoprotein membranes of mitochondria seems to be due mainly to disturbances in phospholipid metabolism. (auth)

45839 LYSOSOMAL SYSTEM OF LIVER CELLS IN RATS FOLLOWING A RADIATION DOSE OF 1000 r TO THE WHOLE BODY. Zyss, Ryszard; Michalska-Kaszczynska, Maria (Inst. of Postgraduate Medical Education, Warsaw). *Acta Med. Pol.*; 13: No. 2, 177-92(1972).

Following an x-ray dose of 1000 R to the whole body of rats, a progression of morphological changes including mainly an increase in the number of lysosomal structures—cytosomes, cytoesosomes, multivesicular bodies, and microbodies was observed. The increase in the number of these structures is a measure of the intensity of radiation-induced damage to the cells. Using histochemical, biochemical, and cytochemical methods, the level of acid phosphatase in the lysosomes was investigated. A progressing increase in the activity of this enzyme was observed. In biochemical investigations, this increase was reflected in the increasing total activity of this enzyme in the lysosomal fraction. A similar increase in the activity of this enzyme observed at the same time in the supernatant is explained by increased permeability of lysosomal membranes. A considerable role in the development of radiation-induced damage to the cellular structures is attributed to the observed changes in the metabolism of phospholipids. (auth)

45840 EFFECTS OF RADIOPROTECTIVE AGENTS ON HEMATOLOGICAL INDEXES AND IRON METABOLISM IN MICE THAT WERE EXPOSED TO MIXED RADIATIONS FROM A REACTOR. Santha, Andras; Sztanyik, Laszlo; Mandi, Erika; (and others). *Honvedorvos*; 24: No. 1, 46-60(1972). (In Hungarian).

To assess the radiation injury caused by mixed γ -neutron radiations from an atomic reactor in mice, peripheral blood picture and reticulocyte count were not sensitive as indicators. Incorporation of ^{59}Fe in erythrocytes, however, proved to be a sensitive test. Iron incorporation closely reflected both radiation dose and neutron- γ ratio. AET protected against γ as well as neutron radiation; 5-MOT protected only against γ radiation. It was concluded that preventive medications for mixed radiation injury should not contain Mexamine (5-MOT). (H.B.G.)

45841 BODY WEIGHT DECREASES IN SOME PROTON EXPOSED PRIMATES. Kirk, John H. (School of Aerospace Medicine, Brooks AFB, Tex.). *Space Life Sci.*; 3: No. 3, 271-7(Jun 1972).

Immature *Macaca mulatta* were exposed to varying doses of whole body irradiations utilizing 2 MeV x-rays; 55, 138, and 400 MeV protons; and 2.3 GeV protons. Over 200 primates remain in a life-long study group for irradiation effects. Mean body weight decreases as a function of age, sex, dose, and irradiation type were found in the 55 MeV proton exposed group but not in the other groups. The cause of the decreased weights is unknown at present, as is the reason for the lack of this effect in the more penetrating irradiations. (auth)

45842 DEOXYRIBONUCLEIC ACID METABOLISM IN THE SMALL INTESTINE IN VIVO AND IN VITRO. A STUDY OF THE NORMAL AND 180-MeV PROTON OR ^{60}Co GAMMA IRRADIATED MOUSE. Johanson, Karl J. *Abstracts of Uppsala Dissertations from the Faculty of Science*, No. 204. Uppsala; Univ. of Uppsala (1972). 26p.

Thesis.

Iodine-125-labeled IUdR was found to be a very specific precursor for intestinal and splenic DNA with little labeling by other molecules. No significant differences in the kinetics of the catabolism of $^{125}\text{IUdR}$ were seen in irradiated mice, as compared with control mice. In estimating the radioinduced decrease in $^{125}\text{IUdR}$ incorporation into DNA, similar results were obtained by biochemical and autoradiographic methods and also by measurements of the total ^{125}I retention in the whole small intestine. The relative biological effectiveness, based on the incorporation of $^{125}\text{IUdR}$ into intestinal DNA, was 1.2 ± 0.3 , as found by comparison of 180-MeV protons with ^{60}Co gamma radiation, and the ^{125}I retention after $^{125}\text{IUdR}$ injection was similar. Changes in the thymidine pool due to flooding by thymidine from dead cells seem to be of minor importance, judging from comparison of fluorouracil-treated control and irradiated mice. Segments of the small intestine were viable after 180 minutes culturing on a Millipore filter in the interface between T8 medium and an atmosphere of 95% oxygen and 5% carbon dioxide. This technique should be a valuable tool for radiobiological studies of the small intestine. (auth)

45843 VARIATIONS IN THE BONE MARROW LYMPHOID POPULATION IN THE SUBLETHALLY IRRADIATED GERM FREE MOUSE. Haot, J.; Gustafsson, B.; Boniver, J.; Revesz, L. (Karlinska Inst., Stockholm. Univ., Liege). *Compt. Rend. Soc. Biol.*; 165: No. 6, 1470-2(1971). (In French).

Apart from a slight delay in the thymic regeneration, the hematopoietic regeneration occurring in mice, which have received a sublethal dose of irradiation, does not appear to be very different from that of nonirradiated animals of the same strain, age, and sex. An accumulation of lymphoid cells, characteristic of the regeneration after irradiation, is found in the bone marrow of the germ-free mice. These lymphoid cells present the morphological characteristics of the x cells at the 12th day. (France)

45844 PROTECTION AGAINST GASTROINTESTINAL EFFECTS OF WHOLE-BODY X-IRRADIATION BY A BILE ACID SEQUESTRANT IN RATS. Parkinson, T. M.; Drake, N. A. (Upjohn Co., Kalamazoo, Mich.). *Experientia*; 28: No. 5, 553-4(15 May 1972).

Previous studies showed that cannulation of the bile duct reduced the loss of sodium and prolonged survival of rats exposed to ionizing radiation. These observations suggested that oral administration of a bile acid sequestrant, DEAE Sephadex, might increase survival times of x-irradiated rats. Therefore, rats were exposed to 1250 to 2100 R of x radiation and were allowed free access to food and water following irradiation. DEAE Sephadex was administered by stomach tube; controls were given distilled water. All rats died within nine days following irradiation. However, the survival time of rats treated with DEAE Sephadex was increased by 1.2 days. It is pointed out that bile acid sequestrant therapy might provide a means of reducing the severity of secondary effects of ionizing radiation in the human gastrointestinal tract. (H.L.W.)

45845 SENSIBILISATION IN RADIOBIOLOGY AND RADIOTHERAPY. Pohlitz, W. (Gesellschaft fuer Strahlen- und Umweltforschung mbH, Frankfurt am Main). *Strahlentherapie*; 143: No. 3, 348-52(Mar 1972). (In German).

A review is presented of various aspects of sensitization of living cells to ionizing radiation. The possibilities for radiosensitization of the different structures of the cell, tissues, and the entire organism are discussed. The role of DNA as a target molecule is emphasized and the reactions of DNA with radio-induced hydroxyl radicals with a high redox potential are discussed. Other topics reviewed are: effects of cell phase on sensitization; relation of sensitization to cell repair of radiation injuries; therapy with fast neutrons and mesons; effects of radio-induced radicals on nucleotides; radioinduced strand breaks in DNA; and immune reactions of living cells against radiation and chemicals. (H.L.W.)

45846 ACUTE DEATH OF MICE DURING MASSIVE GAMMA IRRADIATION. 3. Takeshita, Kenji (Hiroshima Univ.); Umayahara, Akira; Fujii, Kyoichi. *Nippon Igaku Hoshasen Gakkai Zasshi*; 31: No. 2, 133-42(May 1971).

An electrocardiographic observation was conducted to study the effect of massive irradiation of ^{60}Co gamma rays given to ICR female mice in a short period of time. The relation between exposure rate and the total exposure leading to death during irradiation was studied in the range of 10 R/min to 40 kR/min. The result indicated that the exposure rate was in proportion to the total exposure leading to death. The relation between the exposure rate and the time until death during irradiation represented almost a direct line in the log-log graph, although dose dependency was observed to some extent. Intraperitoneal injection of 2.5 mg phenobarbital was given 10 minutes before irradiation with 40 kR/min. The exposure rate leading to death was 2.6 times larger. An electrocardiographic observation was conducted continuously

from the beginning of irradiation until death was confirmed. The total exposure delivered was 18 kR which exceeded the 3.5 day effect in the relation to mean survival time and dose. However, the effect was equal to the delay of acute death during irradiation. The heart rate and electromyographic frequency in convulsions during irradiation were determined. The heart rate generally increased with irradiation; no significant change was observed with barbital injection. The electromyographic frequency indicated no definite pattern in convulsions during irradiation. Mean survival time of irradiated mice was determined and expressed as an exponential function of the given dose. (Japan)

45847 EFFECT OF VARIOUS CELL-FREE ORGAN EXTRACTS ON HEMATOPOIETIC STEM CELLS. Dons, R. F.; Fried, W.; Trobaugh, F. E. Jr.; Knospe, W. H. (Univ. of Illinois, Chicago. Rush Presbyterian St. Luke's Hospital, Chicago). *Exp. Hematol.*; 22: No. 1, 30-2(Mar 1972).

Tissue extracts were prepared by homogenizing mouse organs, centrifuging, and collecting the supernatant; each of the supernatants was then diluted with saline. Mice were injected with one ml of supernatant or one ml of normal saline and 2 hrs later the mice received 200 R whole-body x radiation. Four days later the mice were killed and cells were flushed from the femoral marrow, and the number of nucleated cells per ml of suspension was determined. One half ml of each suspension was then injected into each of 20 mice that had received 750 R x radiation. Nine days later the number of spleen colonies was counted. A significantly greater number of CFU was observed in mice treated with spleen, kidney, liver, lung, or stomach extracts than in saline-treated mice. (H.L.W.)

45848 GLUCOSE-6-PHOSPHATE PHOSPHOHYDROLASE ACTIVITY IN ERYTHROCYTES AFTER X-IRRADIATION. Moisiu, M.; Chisleag, C.; Ababei, L. (Inst. of Medicine and Pharmacy, Iasi). *Stud. Biophys.*; 28: No. 3, 199-206(1971).

After a single whole-body x irradiation with 600 R, glucose-6-phosphate phosphohydrolase activity fell in rabbit erythrocytes from 4.20 to 3.12 $\mu\text{mole} \cdot \text{ml cells}^{-1} \cdot \text{h}^{-1}$ and in rat erythrocytes from 3.56 to 2.40 $\mu\text{mole} \cdot \text{ml cells}^{-1} \cdot \text{h}^{-1}$. Cysteine and reduced glutathione (GSH) had no effect on G-6-Pase activity in normal erythrocytes, but restored enzymatic activity in the irradiated erythrocytes. Parachlormercury-benzoate (PCMB) inhibits G-6-Pase activity more strongly in the erythrocytes of normal than of irradiated animals. Inhibition was reversible with cysteine and GSH. (auth)

45849 PROTEIN SYNTHESIS IN NORMAL, X-IRRADIATED, AND FRIEND VIRUS INFECTED MOUSE SPLEEN NUCLEI. Munson, B. R.; Fiel, R. J. (Park Memorial Inst., Springville, N. Y.). *Res. Commun. Chem. Pathol. Pharmacol.*; 1: No. 4, 517-25(Jul 1970).

Nuclei were isolated and purified from normal, x-irradiated, and Friend virus (FV) infected mouse spleen and analyzed for their relative incorporation of ^3H -labeled amino acids. The data show that incorporation into FV infected nuclei was less than into normal spleen nuclei. Conversely, incorporation into spleen nuclei from x-irradiated mice was found to be stimulated although the level returned to normal in regenerated spleen. (auth)

45850 NEUROPATHOLOGICAL EVALUATION OF MONKEYS EXPOSED TO BODY-ALONE X-RADIATION. Haymaker, Webb (Ames Research Center, Moffett Field, Calif.); Sturrock, David; Kirk, John H.; Casey, Harold W.; Call, Nancy A. *Space Life Sci.*; 3: No. 3, 210-25(Jun 1972).

Twelve monkeys (*Macaca mulatta*) were x-irradiated and six controls received sham treatment. The radiation dose was 600 rad to the trunk and 6 rad to the head. Hematological studies before and after exposure included total leukocyte counts, platelet counts, hemoglobin concentrations, microhematocrits, and total red cell counts. At autopsy, specimens were removed from the cerebrum, cerebellum, and brain stem, placed in fixatives, and prepared for histological study. Photographs and tables describing abnormalities in the central nervous system are presented. The most striking change consisted of hypertrophy and multiplication of microglial cells in the molecular layer of the cerebellum. (H.L.W.)

45851 OCULAR SQUAMOUS CELL CARCINOMA (CANCER EYE) IN HEREFORD CATTLE: RADIATION REPAIR PROCESSES AND A COMPARISON OF CULTURED CELLS WITH XERODERMA PIGMENTOSUM IN MAN. Cleaver, J. E. (Univ. of California, San Francisco); Kainer, R. A.; Zelle, M. R. *Amer. J. Vet. Res.*; 33: No. 6, 1131-6(Jun 1972).

Epidermal cells and fibroblasts were cultured from skin biopsy samples taken from the lower eyelids of Hereford and black Aberdeen Angus cows to determine if the predisposition of Herefords to ocular squamous cell carcinoma (cancer eye) is due to an enzymatic defect in the repair synthesis of deoxyribonucleic acid

(DNA) damaged by ultraviolet (uv) light. This reparative defect is present in xeroderma pigmentosum (xp), a human disease in which ocular squamous cell carcinoma occurs. Differences were not observed between the Hereford and Aberdeen Angus cows in sensitivity of their cells to uv light or in the level of DNA repair synthesis. Cancer eye in Hereford cattle appears to be distinct from xp in man and not attributable to a defect in DNA repair. (auth)

45852 WHOLE-BODY GAMMA IRRADIATION OF NEWBORN PIGS: HEMATOLOGIC CHANGES. Case, Marvin T.; Simon, Joseph (Coll. of Veterinary Medicine, Urbana, Ill.). *Amer. J. Vet. Res.*; 33: No. 6, 1217-22(Jun 1972).

Newborn Yorkshire pigs were exposed to 150, 250, 300, 350, 375, 400, and 450 R of ^{60}Co γ radiation. Lymphopenia developed in the irradiated pigs usually within 24 hours after irradiation. Bone marrow damage caused neutropenia and anemia to develop 10 to 14 days after irradiation. Lymphopenia and neutropenia were responsible for the leukopenia in the irradiated pigs. Pigs exposed to a lethal dose of radiation had a gradual decrease of erythrocytes and leukocytes until death occurred. (auth)

45853 WHOLE-BODY GAMMA IRRADIATION OF NEWBORN PIGS: PATHOLOGIC CHANGES. Case, Marvin T.; Simon, Joseph (Coll. of Veterinary Medicine, Urbana, Ill.). *Amer. J. Vet. Res.*; 33: No. 6, 1223-30(Jun 1972).

Newborn Yorkshire pigs were exposed to 150, 250, 300, 350, 375, 400, and 450 R of ^{60}Co γ radiation. Hemorrhage was the principal lesion in the pigs which died 10 to 29 days after exposure to lethal doses of radiation (350, 375, 400, and 450 R). Hemorrhages were always found in the heart, lymph nodes, and skin and were often present in skeletal muscle, intestine, kidney, and brain. Hemopericardium, hemoperitoneum, or hemorrhage into the intestinal lumen were often severe enough to cause death. Destruction of hematopoietic cells occurred in lymph nodes, Peyer's patches, spleen, and bone marrow. Peyer's patches and bone marrow returned to histologic normalcy more rapidly than did lymph nodes and spleen. Radiation pneumonitis developed in the irradiated pigs and was most severe at 35 days after irradiation. (auth)

45854 EFFECTS OF RADIATION AND HYPOXIA ON THE METABOLIC FATE OF ERYTHROPOIETIN. Carmena, A. O. (Tufts Univ., Boston). *Acta Radiol., Ther., Phys., Biol.*; 10: No. 5, 497-503(Oct 1971).

Sprague-Dawley rats were kept at 19,000 feet for various periods. The animals that received 200 R whole-body irradiation had a significantly increased plasma erythropoietin. It is concluded that the levels of erythropoietin are in inverse relationship to the cellularity of the erythroid precursors. (auth)

45855 CULTURE OF MURINE LEUKEMIC CELLS INDUCED BY X-IRRADIATION. Iwai, Yoshiaki; Takamori, Yasuhiko; Okumoto, Masaaki. *Annu. Rep. Radiat. Center Osaka Prefect.*; 12: 113-19(1971).

Ascites leukemic cells derived from splenic reticular sarcoma induced by x-irradiation in RF mice were successfully cultivated in vitro. At early passages in vitro, the culture cells adhered less firmly to the glass and tended to float into the culture fluid. The culture cells altered epithelial like cells with increasing time of cultivation. The change in cell shape was accompanied by the changes in enzymatic activities of the cells, and by loss of retransplantability in the isologous mice. (auth)

45856 COMPARATIVE CHARACTERISTICS OF NUCLEOTIDE BLOCKS OF DNA IN RADIATION INJURY AND LEUKOSIS. Kritskii, G. A.; Batishchev, A. I.; Aleksandrov, S. V.; Fedorov, N. A.; Abramov, R. E. (Inst. of Biochemistry, Moscow). *Dokl. Akad. Nauk SSSR*; 203: No. 1, 233-6(1 Mar 1972). (In Russian).

The distribution of nucleotide blocks of DNA was studied under conditions of radiation damage and leukosis. White rats were given 500 R of x radiation, and after decapitation the bone marrow of the limbs was frozen, thawed, and examined. The change in the ratio of the nucleotide blocks in DNA of the hematopoietic tissues of animals during radiation injury and during leukosis possessed similar character. (K.S.W.)

45857 VASCULARIZATION OF MUSCULAR AUTOTRANSPLANTS DEVELOPED IN AN IRRADIATED SITE. Umnova, M. M. (Inst. of Biological Physics, Pushchino-na-Oka, USSR). *Dokl. Akad. Nauk SSSR*; 203: No. 1, 248-50(1 Mar 1972). (In Russian).

The effect of irradiation on the development of the blood supply in an autotransplant of the gastrocnemial muscle of the rat was studied. Experiments were conducted with 70 white rats weighing 120 to 130 g and unidentified as to strain. The right rear leg was

given a dose of 2,500 R (type of radiation not specified). The gastrocnemial muscle was removed, and in its place the contralateral unirradiated muscle was transplanted. The proximal part of the muscle and Achilles tendon were sewn to fragments of tendon of the removed muscle. The tibial nerve was sewn underneath to the mid-part of the transplant. Observations of the animals were made over a period of two months, and revealed that the rate of vascularization and reorganization of the unirradiated transplants varied in accordance with the radiation dose received by the transplant site. As a result of the radiation damage, the regeneration capacity of the blood vessels in the transplants was suppressed to a significant extent. In spite of this, anastomosis occurred and assured vascularization in the transplant. (K.S.W.)

45858 REMOTE EFFECTS, IN TIME AND PLACE, OF X IRRADIATION OF THE VISUAL CORTEX IN THE MONKEY. Caveness, William; Kato, Motohiro; Hess, Kenneth H.; Carsten, Arland L. *Trans. Amer. Neurol. Ass.*; 96: 217(1971).

Six to eight months after subjecting the right occipital lobe to a single exposure of 3,500 rads by 250 KV x-rays, eight young adult *Macaca mulatta* showed alterations in the photic evoked response from the visual cortex. Two showed unilateral depression in amplitude over the irradiated area, and six showed bilateral depression in amplitude, that is, over the homologous non-irradiated as well as the irradiated area. Of the latter, four showed behavioral loss in vision, and funduscopic evidence of optic atrophy. Upon sacrifice, in addition to the expected degenerative and proliferative changes in the right occipital lobe, there was diffuse brain swelling, severe in one instance, with intra- and extracellular fluid accumulation. But more intriguing was a partial loss in fibers in the optic nerves with the loss being predominantly on the temporal side in the right eye and nasal side in the left eye. (auth)

45859 RADIOMODIFYING CHEMICALS FOR BACTERIA AND MICE. Mulligan, David Dwight. Gainesville, Fla.; Univ. of Florida (1971). 200p. University Microfilms Order No. 72-16,637.

Thesis.

Chelating compounds were tested for their ability to modify the x-ray sensitivity of suspensions of stationary phase *Escherichia coli* B/r bacteria, and of female mice, age 12 to 15 weeks, strain BC3F₁/C. The bulk of the toxicity and radiomodification experiments were conducted using the following para-substituted, phenyl-containing chelating structures: thiocarboxylic acids, sulfenyl- and sulfonylcarboxylic acids, mercaptoalkylamines, and ammonium dithiocarbamates. In general for bacteria, the toxicity decreased and the radioprotection increased for a corresponding increase in electron-donating character of the para-substituents. Some electron-withdrawing groups caused radiosensitization for bacteria. On a per chemical dose basis, protection of mice usually increased as the electron-donating power of the substituents increased. Electron-withdrawing groups generally decreased protection for mice, but did not cause radiosensitization. The compounds that were the most protective were generally the least toxic to the bacteria and the mice and could therefore be employed in large amounts, affording greater protection against x rays. The powerful chelating agent, 8-mercaptoquinoline, was protective to mice but was also quite toxic. When it was used with an effective radioprotectant, AET, the combined effect was protection at considerably higher x ray doses than with either compound alone. A group of thiol-sulfonates were found whose low toxicity to mice and high toxicity to bacteria recommends that they be further investigated as possible antimicrobial agents for mammals. Several compounds ($\text{BrC}_6\text{H}_4\text{SCH}_2\text{COOH}$, $\text{ClC}_6\text{H}_4\text{NHCCSSNH}_4$, $(\text{NO}_2\text{C}_6\text{H}_4)_2\text{S}$, and $(\text{CH}_3)_2\text{NCSS-Na}$) were found to be particularly radiosensitizing to bacteria but at least moderately radio-protective to mice, indicating a possible use in selectively potentiating the toxicity of radiotherapy for tumors while protecting the whole animal from radiation. (Diss. Abstr. Int., B)

45860 X-IRRADIATION OF THE SKIN OF THE BEAGLE. Park, Richard Dee. Davis, Calif.; Univ. of California (1971). 173p. University Microfilms Order No. 72-17,922.

Thesis.

Single-dose and fractionated-dose x-irradiation studies were performed on the skin of Beagle dogs. In the single-dose study, five 48 cm² areas of skin were irradiated over the neck and back of the dogs. The doses used were 610, 1220, 1830, 2440, and 3660 rads. In the fractionated-dose study, 17 schedules were used varying fractionation dose, fractionation interval, and protraction time. Total dose was constant at 5800 rads. All animals were sacrificed approximately 120 days after the initial radiation treatment. Irradiation was done with a 200 kVp, 15 ma beam having a HVL of 5 mm Al, a FSD of 50 cm and delivering 195 R/minute. Gross observation, histopathology, autoradiography, and angiography were used to evaluate the radiation induced changes in skin. Thermoluminescent dosimetry with LiF crystals was used to confirm that

the radiation absorbed dose to the skin over the irradiated field was homogeneous; the backscatter factor was lower than reported; and skin over the thorax received less absorbed radiation than other areas with similar exposures because of a lower backscatter from the air filled lungs. Acute radiodermatitis from single-dose irradiation was proportional to the dose level. Chronic radiodermatitis was dose and time dependent. The following acute dose effects were observed; 610 rads produced erythema, 1220 rads produced dry desquamation, 1830 rads produced a moist desquamation; and 2440 or 3660 rads produced ulcers. Large, slow healing ulcers were seen with 3660 rads. Chronic radiodermatitis dose effects, observed no sooner than 50 days after irradiation, were; 610 rads produced slight or no changes, 1220 rads produced gray hair, 1830 rads produced patchy epilation, and 2440 and 3660 rads produced atrophic skin with some residual ulcers and tendencies for reulceration. Chronic histopathologic changes observed with 1220 rads and above consisted of dermal hyperplasia, increased dermal connective tissue, and infiltrating phagocytic mononuclear cells around dead skin structures. The hair follicles were the most radiosensitive skin structure followed by the sebaceous glands. The synthesis of DNA by epidermal cells was not directly affected by radiation as determined by the epidermal uptake of thymidine-³H immediately after irradiation and subsequent autoradiographic studies. Some areas of skin that received 1830 rads or more were hypovascular. Chronic vessel changes consisted of narrowing of the vascular lumen as the vessels entered the irradiated skin, proliferation of small vessels around the periphery of the irradiated skin and some large hypertensive peripheral vessels. It was proposed that the vascular changes predisposed the skin to reulcerate. To evaluate the final chronic effects of 2440 and 3660 rads an observation time of 6 months to 1 yr should be allowed. The epidermis was sloughed from all areas in the fractionated-dose study. The total dose of 5800 rads was concluded to be too high. A trend toward skin sparing, both acute and chronic, was noticed in areas irradiated with relatively intermediate fractionated levels of 600 and 800 R and a protraction time of 20 days. More studies are needed to establish safe radiation dose fractionation schedules in the dog. (Diss. Abstr. Int., B)

45861 DOSE TO THE EPITHELIUM OF THE UPPER RESPIRATORY TRACT FROM RADON AND ITS DAUGHTER PRODUCTS. Bykhovskii, A. V.; Khachirov, Dzh. G.; Shishkanov, N. G.; Dubrovin, S. A.; Klyuch, V. E. *Health Phys.*; 23: No. 1, 13-22(Jul 1972). (In Russian).

The study of the effects of ²²²Rn and its daughter products on the organism is one of the most important problems both of radiation hygiene and radiobiology. Tests were carried out on animals for the more accurate definition of the irradiation dose to the epithelium of the upper respiratory tract and to study its dependence on the concentration of radon and its daughter products. The direct radiometry of the alpha-activity in different parts of the respiratory tract enabled calculation of the dose in the basal epithelium layer of the trachea and main bronchi. It was shown in the course of the study that the dose production is essentially due to the daughter products of ²²²Rn which do not combine with aerosol particles of the air of the test chamber. (auth)

45862 EXPERIMENTAL STUDIES ON THE PERMEABILITY OF LINGUAL MUCOSA IN RADIATION GLOSSITIS. Takeda, Ryuichi (Kanazawa Univ., Japan). Kanazawa Iirigaku Shosho; 88: 33-60(Apr 1971). (In Japanese).

Progress of the experimentally irradiated lingual mucosa of rabbits from onset of inflammation to healing was observed macro- and microscopically. In the group irradiated with a single dose of 1800 rads of β -ray, the radiation glossitis showed the following processes: incubation period within 24 hrs, incipient stage for 3 to 5 days, middle stage for 6 to 9 days, and the final stage after 10 days. In the group irradiated with a single dose of 2500 rads of β -ray, the incipient inflammation was manifested after 2 days followed by the middle stage for 6 to 12 days, and the final stage. The latter group showed more severe inflammatory process and the prolonged period of the inflammation. In two groups irradiated with a single dose of 2500 rads x rays and ⁶⁰Co γ rays, the inflammatory process appeared almost the same. A comparative study between the radiation glossitis and its related permeability with ²⁴Na was carried out. Three days after irradiation, the permeability was slightly reduced in all groups regardless of radiation sources or dosages. In the severe ulcerated glossitic stage during 7 to 10 days, the ²⁴Na uptake was the highest. In autoradiograms, a deeper permeability was observed in particular at the area of ulceration with less epidermis. The ²⁴Na uptake decreased day by day in the recovering stage. The permeability came back nearly to the level of preirradiation 31 days after the irradiation. (Japan)

45863 EXPERIMENTAL STUDIES ON THE PULMONARY RADIATION INJURY. Oikawa, Masaru (Iwate Medical Coll.,

Morioka, Japan). Nippon Igaku Hoshasen Gakkai Zasshi; 31: No. 4, 426-42(Jul 1971). (In Japanese).

When irradiated, lungs of rats showed histologic changes similar to those seen in man with radiation pneumonitis, while lungs of rabbits did not show histologic change. The alveolar sac stability ratio tended to decrease at the early stage of the pulmonary radiation injury in rats suggesting that the primary manifestation of the injury was in the pulmonary surfactant systems. The early histologic changes characteristic of the injury in rats were edema in the alveolar space and interstitium, fibrin deposition, bizarre cells, foamy cells and atelectasis. As for the intermediate changes, capillary thrombus, the Masson body, and onionskin appearance like allergic inflammations are seen. Those changes were also seen in the non-irradiated lung though less frequent. In the present study, there was no qualitative difference in the pulmonary damages between irradiated and non-irradiated areas. These results favor the allergic involvement in the pathogenesis of pulmonary radiation injury. (auth)

45864 STUDY ON THE RBE OF VARIOUS TYPES OF RADIATION. Takahashi, Akira (Hiroshima Univ.). Nippon Ganka Gakkai Zasshi; 75: No. 7, 1661-9(Jul 1971). (In Japanese).

One eye of normal young albino rabbits, 10 to 15 weeks old, was irradiated with one of the following types of radiation and follow-up observation was made for a period of more than 60 weeks. The types of radiation used were 180 kv x rays and gamma rays (^{60}Co energy source) administered in the six different doubling doses of 50, 100, 200, 400, 800, and 1600 rad, and 14.1 MeV fast neutrons and mixed neutrons and gamma rays were administered in the five different doubling doses of 50, 100, 200, 400, and 800 rad. The relative biological effectiveness (RBE) of 14.1 MeV fast neutrons, gamma-rays (^{60}Co), and mixed rays required to produce mild disturbances that resulted in permanent changes in the crystalline lenses of young albino rabbits at the end of 20 weeks was 2, 1, 1 respectively when compared with 180 kv x rays, while that needed to produce permanent changes and cause such severe disturbances as possible decreased vision was calculated to be 2, 1, and incalculable. The RBE at the end of 40 weeks was 8, 1, and 2 respectively for mild disturbances that resulted in permanent changes, while that needed to produce permanent changes and cause the severe disturbance of possible decreased vision was calculated to be 2, 1, and 1. The RBE at the end of 60 weeks was 8, 1 to 2, and 2 respectively for mild disturbances that resulted in permanent changes, while that needed to produce permanent changes and cause such severe disturbances as possible decreased vision was calculated to be 2, 1, and 1 respectively. (auth)

45865 EFFECT OF IRRADIATION ON THYROID GROWTH IN MOUSE FOETUSES AND GOITROGEN CHALLENGED ADULT MICE. Walinder, G.; Sjoeden, A.-M. (Foersvarets Forskningsanstalt, Stockholm). Acta Radiol., Ther., Phys., Biol.; 10: No. 6, 579-92(Dec 1971).

The radiosensitivity of the goitrogen challenged thyroid gland was investigated in fetal and adult mice both after irradiation with ^{131}I accumulated in the gland and after x irradiation. The similar effect in the normally growing thyroid in the fetuses and young mice and the goitrogen-stimulated adult animals suggested that the rate of thyroidal growth rather than the age of the mouse was the decisive factor in determining the degree of the reaction. X irradiation proved to be two to four times more effective than internal irradiation with ^{131}I . (auth)

45866 ACUTE DEATH OF MICE IRRADIATED WITH PULSED X-RAYS FROM A LINEAR ACCELERATOR. Antoku, Shigetoshi; Takeshita, Kenji (Hiroshima Univ.). Hiroshima Daigaku Genbaku Hoshano Igaku Kenkyusho Nenpo; No. 12, 63-8(1971). (In Japanese).

The comparative lethal effects on mice of 25 MeV pulsed x rays from a linear accelerator and ^{60}Co gamma rays were studied. The $\text{LD}_{50(30)}$ of mice for pulsed x rays and ^{60}Co gamma rays was 809 rads and 902 rads, respectively. According to this biological response, the relative biological effectiveness (RBE) of pulsed x rays to ^{60}Co gamma rays was 1.1. The patterns of daily mortality and body weight changes of mice irradiated with pulsed x rays were similar to those irradiated with ^{60}Co gamma rays. These facts indicated no qualitative differences in biological effects between pulsed and sustained radiations. However, it is necessary to study dose rate effects in various biological materials, because significant differences have been reported for intestinal and cerebral deaths between pulsed and sustained radiations. (auth)

45867 STATISTICAL ANALYSIS OF RADIOINDUCED ABNORMAL DEVELOPMENT. Satow, Yukio; Miyabara, Shinichi; Ikeda, Takayoshi; Ueno, Takehisa (Hiroshima Univ.). Hiroshima Daigaku Genbaku Hoshano Igaku Kenkyusho Nenpo; No. 12, 190-6(1971). (In Japanese).

The irradiation of pregnant rats with 80 to 150 rad of 14.1 MeV fast neutrons at 7 to 11 days after conception induced malformations of the hearts and blood vessels in the fetuses. In this study some methods were carried out such as regression analysis or variance analysis to examine the relation between the malformation rate in the fetuses and the date or the dose of irradiation. Although the malformations are induced by the synthetic effects of the dose and date of irradiation, the results showed that the malformation rate depended on the date, rather than dose, suggesting the importance of "sensitive periods" for the abnormal development. The data obtained from previous experiments indicate that irradiation with the dose of 130 rad of neutrons at 8 days after conception was most effective to induce the cardiovascular anomalies. In other words, the irradiation of 130 rad at 8 days after conception induced a high incidence of the anomalies with low incidence of absorption of the embryos. The analysis of the relation between the anomaly and absorption using the χ^2 -test or experimental formula was carried out on this data. It was proved that there were some significant differences between anomalous and absorbed rate. (auth)

45868 RIBONUCLEIC ACID SYNTHESIS AND IONIZING RADIATION. Olkowski, Z. (Silesian Medical Academy, Katowice, Poland). Postepy Astronaut.; 4: No. 4, 67-71(1970). (In Polish).

The subject of these studies was the motoneurons of spinal cords of mice in acute radiation sickness. The synthesis of RNA was studied by autoradiography using ^3H -uridine. Besides the following enzymes, connected with RNA metabolism, were studied cytochemically: glucose-6-phosphate, 6-phosphogluconate and succinate dehydrogenases according to Pearse, adenosine-3-phosphatase according to Wachstein and Meisel, thiamine pyrophosphatase nucleosidediphosphatase according to Novikoff, Goldfisher, and Essner. It was shown, that the ionizing radiation inhibits RNA synthesis both in the cell nucleus and cytoplasm and inhibits the activity of investigated enzymes. The changes in the enzyme activity and localization are compared with the events that occur in incorporation of ^3H -uridine in experimental and control animals. (NSA of Poland)

45869 SENSITIZATION OF HYPOXIC CELLS IN VIVO WITH TRIACETONEAMINE N-OXYL (TAN). Hornsey, S. (Hammer-smith Hospital, London). Int. J. Radiat. Biol.; 22: No. 1, 91-14(Jul 1972).

Female mice were whole-body irradiated with 7 MeV electrons from the MRC Linear Accelerator, over a dose range of 2400 to 3600 rads while breathing N_2 and over a dose range of 1200 to 1300 rads while breathing O_2 . Counts were made of the number of survivors 5 days after irradiation. Animals given non-toxic intraperitoneal injections of triacetoneamine N-oxyl (TAN) at a rate of 0.238mg TAN/g mouse 5 min before irradiation while breathing N_2 were more sensitive than animals given no TAN. The dose modifying factor was 1.08. Animals given TAN 20 min before irradiation, while breathing N_2 , showed the same sensitivity as animals without TAN. TAN given to animals 5 min before irradiation while breathing O_2 had no effect on their sensitivity. (UK)

45870 RADIATION EFFECTS ON THE BONE GROWTH AND EFFECTS OF SALIVARY GLAND HORMONE (PAROTIN) ON IRRADIATED BONE TISSUES. Iwasaki, Zensuke (Iwate Medical Coll., Morioka, Japan). Nippon Igaku Hoshasen Gakkai Zasshi; 31: No. 1, 78-95(Apr 1971).

Single irradiation doses of 5,000 R, 2,000 R, and 1,000 R, respectively, were administered to three groups, the whole hind limbs (Group WB), the epiphysis alone (Group E), and the diaphysis alone (Group D), of Wistar rats aged one month or 3 months. An injection of 0.1 mg/kg of parotin was given twice a week over 24 weeks (Group P). Measurements by x-ray pictures were performed after irradiation. In the 1st, 3rd and 6th months after irradiation, the animals were sacrificed and the left hind leg was resected for histopathological examination. The younger the animals were and the larger the dose was, the more intensely the radiation disturbance in the bone appeared. The degree of disturbance was the highest in Group WB, followed by Group E and was the most mild in Group D. In the young rats aged one month, a growth retardation of the tibia was observed in the group irradiated with 1,000 R and was the mildest disturbance in comparison to the contrast group, the growth rate of the tibia 6 months after irradiation being 86.4% in Group WB, 89.7% in the Group E, and 95.5% in Group D. Therefore, the lower limit for manifestation of disturbance was estimated to be below 1,000 R. In Group P, in case of irradiation with 1,000 R, the arrangement of the cells in the epiphyseal cartilage was more regular than the group with irradiation alone and disappearance and necrosis of the nucleus of cartilaginous cells were less. There was no great difference from the control group in extension of the calcification layers and the number of osteocytes, showing an effect of reparation of distur-

bance to some degree. In the irradiated groups with more than 2,000 R the disturbance was greater not allowing for the effect of parotin. (Japan)

45871 STEM CELL RENEWAL IN SPLEEN AND BONE MARROW OF MICE AFTER REPEATED TOTAL-BODY IRRADIATIONS. Vos, O. (Rijksverdedigingsorganisatie TNO, Rijswijk, Netherlands). *Int. J. Radiat. Biol.*; 22: No. 1, 41-50 (Jul 1972).

To study the effect of repeated sublethal irradiations on the multiplication of haemopoietic stem cells, mice were irradiated one, two, or three times with an x-ray dose of 440 rads. The time-interval between the first and the second irradiation was 7 days. When three irradiations were given, the interval between the last two exposures varied between 7 and 100 days. The growth-rate of CFU's in femoral bone-marrow after two irradiations was slower than after one irradiation; at around the 15th day the overshoot in the spleen was smaller after two irradiations than after one. These findings were even more pronounced when a third dose was given within a few weeks after the second. When the third dose was given longer than 5 weeks after the second, the CFU growth curve in bone-marrow and spleen was closer to the curve obtained after one irradiation. The significance of the findings in respect of the capacity of stem cells for self-renewal is discussed. (auth)

45872 EFFECTS OF DIETARY ANTIOXIDANTS ON LIPID PEROXIDE FORMATION IN ANIMAL TISSUES AFTER WHOLE-BODY IRRADIATION. Dawes, E. A.; Wills, E. D. (Saint Bartholomew's Hospital, London). *Int. J. Radiat. Biol.*; 22: No. 1, 23-40 (Jul 1972).

Rates of lipid peroxidation in mouse-liver homogenates increased linearly with dose 24 hr after whole-body x irradiation (500 to 2000 rads), and reached a maximum 3 days after irradiation. Smaller increases in the rates of peroxidation occurred in heart, kidney, or spleen homogenates. Vitamin E supplementation of the diet depressed the rate of lipid peroxidation in liver, heart, spleen, and kidney homogenates after irradiation with 1000 to 2000 rads. Hydroperoxide and malonaldehyde increased significantly in spleen lipids after irradiation but much smaller changes occurred in heart or liver lipids. Dietary vitamin E inhibited increase of malonaldehyde but had little effect on hydroperoxide increases. Dietary BHT (2,6-di-tert-butyl-p-cresol) was less effective than vitamin E but depressed the rate of peroxidation in liver homogenate and microsomal fraction. In the liver microsomal fraction, the maximum rate of peroxidation and minimum antioxidant concentration occurred concurrently. Tissue and subcellular antioxidant concentrations are considered to play a major role in regulating the rate of lipid peroxidation after irradiation. (auth)

Invertebrates

Refer also to abstract: 45724.

45873 (TID-26078) EFFECTS OF GAMMA RADIATION ON THE SPECTRAL SENSITIVITY OF THE NOCTUID MOTH, HELIOTHIS ZEA. Baker, James F. (Florida State Univ., Tallahassee, Coll. of Arts and Sciences). Jun 1972. 14p. Dep. NTIS.

Ionizing radiations, such as x rays, beta rays, and gamma rays, can evoke a visual response in many organisms. In the noctuid moth, *Heliothis Zea*, constant rates of exposure to gamma radiation were shown to act in a manner similar to appropriately chosen levels of ambient illumination in the visible range (400 to 700nm). This study investigates the effects of gamma radiation on the spectral sensitivity of *Heliothis Zea*. If gamma rays were to depress visual sensitivity differentially with respect to wavelength, they would be acting in a manner analogous to selective chromatic adapting stimuli as observed in organisms with more than one visual pigment. Gamma radiation was found to depress visual sensitivity equally, as measured by the ERG response, regardless of wavelength, suggesting that *Heliothis Zea* does not have two or more populations of visual receptors with different pigments. (auth)

45874 NUCLEAR-CYTOPLASMIC RELATIONS IN THE MITOSIS OF SEA URCHIN EGGS. III. gamma-RAY-INDUCED DAMAGE TO WHOLE EGGS AND NUCLEATE AND ANUCLEATE HALF-EGGS. Rustad, Ronald C.; Yuyama, Shuhei; Rostad, Lynne C. (Case Western Reserve Univ., Cleveland). *J. Cell. Biol.*; 49: No. 3, 906-12 (Jun 1971).

Sea urchin eggs were cut into halves. The nucleate and anucleate halves and whole eggs were γ irradiated and then fertilized with normal sperm. The first mitosis of the diploid half-egg was more delayed than the division of the whole egg. There was a small, but highly significant, delay of the mitosis of the haploid half-egg, thus demonstrating cytoplasmic sensitivity to ionizing radiation. Since

the sensitivity of nucleate cells is influenced by cytoplasmic volume, the problem of the role of cytoplasm in repair is considered in relation to these data and other reports in the literature. (auth)

45875 HISTOPATHOLOGY OF IRRADIATED YELLOW MEALWORM ADULT AND LARVAL MIDGUTS. Brower, John H.; Ashraf, Mohammad (Dept. of Agriculture, Savannah). *Ann. Entomol. Soc. Amer.*; 65: No. 4, 950-4 (Jul 1972).

Gamma radiation doses of 5 to 100 kilorads had a pronounced effect on the midgut of both larval and adult yellow mealworms, *Tenebrio molitor* L. (Coleoptera: Tenebrionidae). Degenerative changes were positively correlated with both dose and time elapsed after exposure. The regenerative cells that renew the epithelial lining of the midgut were particularly sensitive to radiation damage; the damage eventually resulted in loss of the midgut epithelium. (auth)

45876 DIFFERENCES IN SYNAPTIC AFFINITY OF CHROMOSOME ARMS OF DROSOPHILA MELANOGASTER REVEALED BY DIFFERENTIAL SENSITIVITY TO TRANSLOCATION HETEROZYGOSITY. Roberts, Paul A. (Oregon State Univ., Corvallis). *Genetics*; 71: No. 3, 401-15 (Jul 1972).

The chromosome arms of *Drosophila melanogaster* differ in the degree to which translocation heterozygosity suppresses crossing over both regionally, and as whole arms. Using x-irradiated flies, the region of maximum sensitivity to structural heterozygosity in each arm of the long autosomes $\frac{1}{3}$ to $\frac{1}{2}$ the length of the arm in from the tip was identified. The regional differences are interpreted as being further indication that synapsis usually begins distally, but not at the telomeres. The different sensitivity of arms 3R, 2R, 3L, and 2L (ranked in order of increasing sensitivity) is evidence of intrinsic differences in the synaptic affinity of chromosome arms. (auth)

45877 MODIFICATIONS INDUCED BY GAMMA IRRADIATIONS IN THE OVIPOSITION OF WORKERS AND FEMALES OF THE ANT DOLICHODERUS QUADRIPUNCTATUS. Torossian, C.; Causse, R. (Faculte des Sciences, Toulouse. INRA, Montfavet, France). *Ann. Zool.-Ecol. Anim.*; 3: No. 4, 427-41 (1971). (In French).

Experiments on workers of *D. quadripunctatus* exposed to various doses of gamma rays result in reducing the fecundity and in a final sterility. Females of the same species subjected to gamma radiations offer a great resistance, and their mortality for doses of median value (5000 to 50,000 rads) remains low; nevertheless their longevity is shortened, especially for doses above 50,000 rads. The fecundity of these females is lowered to a residual level, all the eggs laid being abortive. A further series of experiments was made to disturb the balance in the ant society between female and workers. When only the queen is irradiated, the workers begin to lay eggs, even when the queen is still laying eggs. Irradiation with low doses resulting in sterile workers, followed by the introduction of young larvae, induces the production of workers, of females and of some winged females. (auth)

45878 FECUNDITY AND LONGEVITY OF F₁ FEMALES OF HABROBRACON FROM SPERM X-RAYED WITH 3000 r. Heidenthal, Gertrude (Russell Sage Coll., Troy, N. Y.); Nelson, Wayne; Clark, Leonard. *Genetics*; 71: No. 3, 349-65 (Jul 1972).

The fecundity and longevity of two groups of F₁ diploid virgins of *Habrobracon* the fathers of which were x rayed with 3000 R were compared. Sorting of wasps into two groups was based on low or high hatchability of their F₂ haploid eggs. Wasps with low hatchability (below 60%) are heterozygous for induced embryonic lethals such as recessives and translocations; wasps with high hatchability (100 to 60%) are free of lethals or possibly heterozygous for mild detrimentals. The F₁s showing high hatchability laid significantly more eggs on the average; the average difference in lifespan was almost statistically significant (7% level). A more sensitive analysis by regression indicated a highly significant linear regression of survival of all F₁s on hatchability of their eggs: F₁s whose eggs showed 100% hatchability. Furthermore, the former laid 4.99 more eggs on the average than the latter. The data also showed a very highly significant positive correlation coefficient between eggs laid and survival of mothers (0.122) and a very highly significant positive partial correlation coefficient (0.105) with hatchability held constant. The latter indicates that the F₁s that laid more eggs also lived longer, regardless of the hatchability of their eggs and hence of the heterozygosity of the mothers for lethals and other detrimentals. (auth)

45879 INFECTIVITY AND DEVELOPMENT OF X-IRRADIATED METACERCARIAE OF LEUCOCHLORIDIOMORPHA CONSTANTIAE (TREMATODA) IN THE CHICK AND ON THE CHORIOALLANTOIS. Fried, Bernard; Davis, James E. (Lafayette Coll., Easton, Pa.). *Trans. Amer. Microsc. Soc.*; 91: No. 2, 208-12 (Apr 1972).

Infectivity and development of *L. constantiae* metacercariae x-irradiated at 500 to 20,000 R were studied in the bursa of Fabricius of the chick and on the chorioallantois. X-irradiation significantly affected infectivity in the chick and on the chorioallantois. Control flukes and those irradiated up to 2,000 R were ovigerous. Egg development was suppressed at 3,000 R and postmetacercarial development did not occur at 5,000 to 20,000 R. Radiation did not affect pairing or clustering of worms in the bursa or on the chorioallantois. (auth)

45880 NEW METHOD FOR THE QUANTITATIVE STUDY OF INDUCED AUTOSOMAL NON-DISJUNCTION IN *DROSOPHILA MELANOGASTER*. Clark, A. M.; Sobels, F. H. (Univ., Leiden). *Genen Phaenen*; 15: No. 1, 29-30 (Jun 1972).

A new method for determining the relative frequency of non-disjunctional progeny in *Drosophila melanogaster* was developed. The dose-effect relationship was studied for the induction by x rays of non-disjunction of the second chromosome in stage 7 oocytes; no significant departure from linearity was observed over a range of exposures from 250 to 3000 R. Thus no evidence for a threshold, even at the low exposure dose of 250 R was obtained. (H.L.W.)

45881 CLONAL PARAMETERS OF TERGITE DEVELOPMENT IN *DROSOPHILA*. Garcia-Bellido, A.; Merriam, J. R. (CSIC, Madrid. Univ. of California, Los Angeles). *Develop. Biol.*; 26: 264-76 (Oct 1971).

A clonal analysis of abdominal development in *Drosophila* was carried out using cell marker mutants. The uncovering of these recessive markers in heterozygous individuals occurred after spontaneous loss of a X ring chromosome (gynandromorphs) or by somatic recombination induced at different developmental stages by 1000 R of x-ray. The minimum spot sizes resulting from preblastodermic gynandromorphs indicate that the least number of primitive anlage cells is about 8. This is also the number of imaginal cells per anlage present throughout the larval stages of development. Only with the onset of metamorphosis do the imaginal cells start dividing and spreading over the presumptive tergite area, each pair of bilateral anlage giving rise to a complete tergite. All the tergites studied (II-VI) behaved uniformly in the present experiments. Corresponding with the constant number of imaginal cells during the larval development there is a constant frequency of induced spots or a constant cell sensitivity to SCO. This sensitivity varies depending on the chromosome considered, being higher in the X chromosome than in the 3L chromosome arm. The data on gynandromorphs, as well as those of somatic spots, indicate that the adult tergites have an indeterminate pattern of growth. The cell determination to become trichome or micro- or macrochaeta occurs sometime during the pupal expansion of the growing imaginal cell population. Trichome- and chaeta-forming cells may then be separated to construct the adult cuticular pattern. (auth)

45882 ROLE OF THIOLS IN THE RADIOSENSITIVITY OF SEA URCHIN EGGS. Smirnova, I. B.; Latsis, R. V. (Inst. of Biological Evolution, Moscow). *Dokl. Akad. Nauk SSSR*; 203: No. 1, 254-6 (1 Mar 1972). (In Russian).

The change in radiosensitivity of sea urchin eggs and their content of SH-compounds at various stages of mitosis during the first three weeks of cleavage were studied. Experiments were conducted on artificially fertilized mature eggs of the sea urchin developing at 19 to 21°C in fresh sea water. Doses of 1,500 R of x rays were administered at different stages of mitosis, and the survival rate of the embryos was obtained. Eggs irradiated during the metaphase exhibited the greatest survival rate; survival was lowest among the telophase-prophase-irradiated. The metaphase was also richest in thiol content; the telophase-prophase was poorest. A 1% solution of ether in water was employed as a holding agent for temporarily stopping mitosis, for making comparisons in ensuing development. Sufficient correlation was then possible to suggest that the change in radiosensitivity during the course of embryogenesis was associated with a fluctuating level of endogenic thiols. (K.S.W.)

45883 EFFECT OF IONIZING RADIATION ON THE STABLE FLY, *STOMOXYS CALCITRANS* (LINNAEUS). White, Sheldon Andrew. Gainesville, Fla.; Univ. of Florida (1971). 147p. University Microfilms Order No. 72-16,670.

Thesis.

The stable fly is an important economic pest that requires continual and costly control measures. Eradication of this pest would be a far more worthwhile goal. The ultimate objective of this investigation was to determine if radiation-sterilization of stable fly pupae could produce males that would be of practical value in a sterile-male-release program. During this investigation, flies were irradiated with either 1000 R, 1200 R, 1500 R, 1750 R, or 2000 R of gamma radiation as 48-, 72-, 96-hour-old pupae. The effects of these dosages on eclosion, reproduction,

and longevity of stable flies were determined. The results indicated that, regardless of the age of pupae when irradiated, apparently none of the dosages adversely affected the time at which maximum adult eclosion occurred, the percent eclosion, or the ratio of males to females produced in each age group. The effect of gamma radiation on oviposition was dependent upon the age of the pupae when irradiation occurred. Females irradiated as 48-hour-old pupae failed to oviposit when treated with 1250 R. In those exposed as 72-hour-old pupae, oviposition was completely inhibited at 1750 R dosage level. Oviposition in females irradiated as 96-hour-old pupae was completely inhibited by a dosage of 2000 R. Oviposition in untreated females was both reduced and delayed when mated with males irradiated with 1500 or more roentgens as 48-hour-old pupae. This did not occur when untreated females were crossed with treated males irradiated as 72- or 96-hour-old pupae. Percent sterility in treated males and females, crossed with untreated flies, increased as radiation dosage increased. In the 48-hour-old age group, complete sterility in females occurred at 1250 R, while male sterility was complete at 1500 R. Complete sterility in females irradiated as 72- or 96-hour-old pupae occurred at a dosage of 1750 R. Virtually complete sterility in males from both age groups occurred at 2000 R. With few exceptions, the mean longevity of treated flies within an age group decreased as radiation increased. Further, longevity increased as the age of the pupae at irradiation increased. Longevity was most severely affected in flies irradiated as 48-hour-old pupae. Further, flies from this age group were apparently intolerant to cold and demonstrated severe wing deformations when exposed to dosages exceeding 1250 R. The mean longevity of males treated with 2000 R as 72- or 96-hour-old pupae was 9.8 and 17.0 days, respectively. The latter treatment produced males that were fully competitive with untreated males in each of the five mating competitiveness ratios tested. The results of these tests apparently indicate that, unless the age of pupae can be determined, the use of sterilized pupae in a successful stable fly sterile-male-release program will be limited. Pupation must be highly synchronous to achieve efficiency in mass-rearing and to facilitate the proper timing of the appropriate sterilizing dosage. Two methods for achieving synchronous pupation in stable flies are suggested. The results of sterility, longevity, and mating competitiveness tests appear to indicate that males exposed to 2000 R gamma radiation as 96-hour-old pupae are satisfactory in every respect for use in a sterile-male-release program for stable fly population control. A method for obtaining these flies from a mixed age group of pupae is suggested. (Diss. Abstr. Int., B)

45884 EFFECTS OF GAMMA RADIATION ON THE RESPIRATION OF *ACARUS SIRO*. Hemenway, Robert Clark, Jr.; Davis, Robert (Univ. of Georgia, Athens). *J. Ga. Entomol. Soc.*; 7: No. 2, 140-51 (Apr 1972).

The rate of oxygen uptake was determined for adult male and female, tritonymphal, and protonymphal stages of the grain mite, *Acarus siro* L., following exposure to gamma radiation doses of 0, 5, 15, 25, 50, and 100 krad. Respiratory rates were measured at temperatures of 15 and 30°C. The results seem to indicate a higher rate of oxygen consumption following exposure to a dose of 25 krad. In mites tested at 15°C, significant stimulation of oxygen consumption was observed only at 25 krad. In mites tested at 30°C, significant depression in oxygen consumption was observed at 5, 50, and 100 krad. Oxygen consumption was examined per individual mite and as a function of body weight. When individual mites were compared, females used the most oxygen, followed in descending order by males, tritonymphs, and protonymphs. Differences in oxygen consumption among life stages were less pronounced when comparisons were made on the basis of body weight. All stages exhibited increased oxygen consumption when the temperature was increased from 15 to 30°C. However, the proportional increase in oxygen uptake in females was not as great as in other stages. (auth)

45885 EFFECT OF ^{137}Cs RADIATION ON GROWTH, DEVELOPMENT, AND MORTALITY OF THE LARVAE OF THE TUSSOCK MOTH, *HEMEROCAMPA LEUCOSTIGMA*. Rossmore, H. W.; Hoffman, E. A. (Wayne State Univ., Detroit). *J. Invertebr. Pathol.*; 17: No. 2, 277-81 (Mar 1971).

Larvae and pupae of *Hemerocampa leucostigma* were irradiated in a ^{137}Cs γ source and observed for effects on growth, development, and mortality. Irradiated larvae were half of control weight 10 days postirradiation. Although pupae were extremely resistant, relatively low doses to larvae prevented emergence. LD₁₀₀ of first instars was 30 kR, while that of third instars was 85 kR. No delay in median time to pupation was noted. (auth)

45886 DISINFECTING MITE INFESTED HAM BY IONIZING RADIATION. Diehl, J. F. (Bundesforschungsanstalt fuer Lebensmittelfrischhaltung, Karlsruhe, Ger.). *Fleischwirtschaft*; 52: No. 1, 81 (Jan 1972). (In German).

Ham infected by mites was irradiated with 10 MeV electrons. A radiation dose of 200 krad destroyed the mites in all stages of development. This dose caused no changes in sensory properties during a storage period of 3 months. (auth)

45887 ENHANCING EFFECT OF LUCANTHONE (MIRACIL D) ON THE FREQUENCY OF X-RAY-INDUCED CHROMOSOME LOSS IN *DROSOPHILA* MALE GERM CELLS. U, Raymond (Duke Univ., Durham, N. C. Veterans Administration Hospital, Durham, N. C.). *Mutat. Res.*; 14: No. 3, 315-24 (Mar 1972).

The effect of lucanthone (miracil D), an inhibitor of RNA synthesis, plus x-irradiation on *Drosophila melanogaster* males Xc^{2yB}/Ysc^{y+} and on the subsequent production of chromosome loss via breakage has been investigated. Lucanthone feeding plus 495 R induced a significantly higher frequency of chromosome loss when irradiation was given in three equal fractionated doses at 3-h intervals than the same dose given acutely. On the other hand, the difference in frequency of non-disjunctional females was not significant. The enhancing effects of this chemical were found only in the fractionated series but were absent in acute x-irradiation series. This effect was found primarily in those cells in spermatid and spermatocyte stages at the time of irradiation. A pertinent point of interest presented was that not only protein synthesis but also RNA synthesis may play a significant role in the development of radiation damage and in postirradiation repair processes at the chromosomal level, since inhibition of RNA synthesis may eventually inhibit protein synthesis. (auth)

Radionuclide Effects (Internal Source)

Man

Refer also to abstract: 45905.

45888 (CONF-720520-1) PATHIOPHYSIOLOGY AND DIAGNOSIS OF INTERNAL CONTAMINATION. Lincoln, T. A. (Oak Ridge National Lab., Tenn.). 1972. 14p. Dep. NTIS.

From Medical planning and care in radiation accidents seminar; Oak Ridge, TN, (21 May 1972).

The hazards from internal irradiation following the uptake of radioisotopes by man are reviewed. It is pointed out that the most serious consequence of radioisotope uptake is the development of neoplasms but, since it may take years for this effect to be manifest, a large exposure at age 55 may be of little consequence in relation to life expectancy while exposure at age 25 may lead to neoplasia within the lifetime of the individual. The possible impact on public health in the event of increased amounts of Pu being released to the environment as a consequence of the expansion of the nuclear power industry is considered. It is pointed out that during the past 25 years no cancers have been reported in personnel working with Pu. Data are reviewed from studies on dogs and rodents that show the development of bone sarcomas and lung carcinomas following the inhalation or injection of ^{239}Pu . The lowest lung burden associated with neoplasm induction in dogs was .05 μCi at death nine years following exposure. Data on the occurrence of neoplasms in man following the injection of Ra showed no cases occurring at doses below 1200 rads. Animal studies have shown Pu to be 5 to 10 times as effective for neoplasm induction as Ra. This would indicate the threshold dose for neoplasm induction by Pu in man is between 50 and 200 rads. The relevance of animal data for estimation of the radiation response of man is discussed. Methods for the evaluation of the radiation dose to personnel following the uptake of ^{239}Pu or ^{241}Pu are discussed briefly. The importance of saving all urine and feces for radiometric analysis and of whole-body counting is emphasized. (C.H.)

45889 LYMPHOCYTE STIMULATION BY THYROGLOBULIN AFTER LOCAL IRRADIATION OF THE THYROID GLAND IN MAN. Einhorn, N.; Packalen, T.; Wasserman, J. (Karolinska Sjukhuset, Stockholm. Sjukvardsstyrelsens Bakterologiska Centrallaboratorium, Stockholm). *Acta Radiol., Ther., Phys., Biol.*; 10: No. 5, 481-7 (Oct 1971).

Lymphocyte stimulation by thyroglobulin was investigated in 46 cases of hyperthyroidism before and after local irradiation of the thyroid gland by ^{131}I . A statistically significant increase in the stimulation occurred at six weeks to three months. (auth)

Animals

Refer also to abstracts: 45814, 45819, and 45865.

45890 (NYO-3970-29) ENVIRONMENTAL RADIOACTIVITY: RADIATION EFFECTS AT THE CELLULAR LEVEL. Fabrikant, Jacob I. (Johns Hopkins Univ., Baltimore, Md. Dept. of Radiology). [nd]. Contract AT(30-1)-3970. 71p. Dep. NTIS.

The effects of internally deposited radioisotopes on man are not well understood, primarily because of the nonuniformity of dose distribution throughout the body, the heterogeneous composition of the various energies from radiations of different LET, and the biological variability in the rate of excretion of the radioisotopes and their daughter products. Data are reviewed from laboratory studies on mammals, primarily on mice, on the effects of internal irradiation on cells in various tissues. These studies provide evidence that during protracted radiation exposure the cells in rapidly renewing cell populations, such as bone osteoblasts, hematopoietic tissues, and epithelial linings of periosteal and endosteal membranes, are subject to radiation injury, depopulation, and regeneration and those cells with the greatest proliferative potential are most subject to radioinduced neoplasia. Data indicate that all ionizing radiations induce the same biological effects but the radiation injury may be distributed throughout the body in different ways, depending on the properties of the radiation and the chemical nature of the emitter. There is abundant evidence that the delayed mutagenic effects from internal emitters are similar to those observed following exposure to external radiation sources. Applications of this data to the prediction of the hazards to public health in the event of an increase in the levels of radioisotopes in the environment are considered. (C.H.)

45891 (ORO-2690-12(Pt.B)) RESEARCH PROPOSAL. Annual Report No. 12, Part B: September 1, 1971-August 31, 1972. (Florida State Univ., Tallahassee. Inst. of Molecular Biophysics). 1 Jun 1972. Contract AT(40-1)-2690. 75p. Dep. NTIS.

Separate abstracts were prepared on four sections of this report. Additional sections discuss RNA metabolism in eukaryotic organisms, the biological significance of tRNA modifications, effects of electronic interactions on molecular excitation processes, and physical production and detection of singlet oxygen in model biological systems. A list is included of publications during the report period. (H.L.W.)

For abstracts of individual sections see: 45817 and 45892-45894.

45892 (ORO-2690-12(Pt.B), Paper 4) BIOLOGICAL TOXICITY OF INTRACELLULAR RADIOISOTOPE DECAY BY BETA EMISSION AND ELECTRON CAPTURE. Hofer, K. F. 3p.

Lymphoid leukemia cells were labeled with various doses of ^{125}I -iododeoxyuridine, ^{131}I -iododeoxyuridine, or ^3H -thymidine and inoculated into new, nonradioactive hosts to facilitate evaluation of toxic effects resulting from the intranuclear decay of the three radioisotopes. Based on radioisotope incorporation data, number of tumor cells present at the time of labeling, labeling fraction, and nuclear diameter of cells, the rate of intracellular energy deposition per labeled cell was calculated from the known decay properties of each isotope. The toxic manifestations of ^{125}I exceeded the effects of tritium and ^{131}I by a factor of 12 to 15. Theories to explain the excessive toxicity of intranuclear ^{125}I are discussed; the theories are based on the fact that the electron capture decay of ^{125}I is accompanied by the release of Auger electrons. (H.L.W.)

45893 (ORO-2690-12(Pt.B), Paper 13) SPECIAL BIOPHYSICS PROJECTS. Smith, J. Marshall. 3p.

Studies on the biophysical evaluation of the radiotoxicity of intracellular ^{125}I are discussed. The unusually high radiotoxic effect of DNA-associated ^{125}I is pointed out. Progress is reported on investigations for determining the consequences of electron capture decay of ^{125}I and the radiolysis of labeled DNA. A theoretical analysis of the microdosimetry of the Auger effect in a DNA- H_2O matrix is being made. Progress is also reported on the development of a scanning electron microscope for the detection and mapping of trace elements in biological specimens. (H.L.W.)

45894 (ORO-2690-12(Pt.B), Paper 14) CHROMOSOMES OF EUKARYOTES: REGULATION OF DNA REPLICATION, ORGANIZATION AND INTEGRATION OF DNA AND EFFECTS OF THE DECAY OF INCORPORATED ISOTOPES (^3H AND ^{125}I). Taylor, J. Herbert. 3p.

Studies on organizational properties of the genome and the regulatory systems which normally operate in cells are being conducted in order to increase understanding of radiation damage and repair as well as evaluation of long-term effects of other mutagenic agents. Four functional units of DNA were identified and these are described. Studies on replicating segments of DNA were conducted using pulse-labeled cells. Electron microscope studies of partially melted DNA revealed small regions separated by 0.3 to 0.5 μ in contour length with quite low melting points. A study of the relative efficiencies of incorporated ^3H and ^{125}I in breaking DNA was initiated. Neither isotope was observed

to produce double-chain breaks, but both produced single-chain breaks with relatively high efficiencies. (H.L.W.)

45895 (UR-3490-102) PATHOGENESIS OF RADIONUCLIDE-INDUCED TUMORS. Casarett, George W. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). [nd]. 25p. (CONF-720505-4). Dep. NTIS.

From twelfth Hanford biology symposium on radionuclide carcinogenesis; Richland, WA. (11 May 1972).

A review is presented of carcinogenic mechanisms and radio-induced carcinogenesis following the uptake of radioisotopes by experimental animals. Discussions are presented of the pathological changes occurring during induction of the following classes of tumors: pituitary and thyroid tumors induced by ^{131}I ; bone tumors induced by ^{90}Sr , ^{226}Ra , ^{228}Ra , ^{238}Pu , and ^{228}Th ; lung tumors induced by ^{210}Po , ^{238}Pu , ^{239}Pu , $^{232}\text{ThO}_2$, ^{186}Au , ^{59}Fe , ^{35}S , ^{32}P and ^{144}Ce ; tumors of the gastrointestinal tract induced by ^{137}Cs , ^{85}Nb , ^{144}Ce , ^{90}Y , and ^{106}Ru ; liver tumors induced by $^{232}\text{ThO}_2$ and ^{238}Pu ; and kidney tumors induced by ^{210}Po . (H.L.W.)

45896 HEMATOPOIETIC AND IMMUNOLOGICAL ALTERATIONS IN MICE PRODUCED BY ^{89}Sr -INDUCED MARROW HYPOPLASIA. Gurney, C. W.; Klassen, L.; Birks, J.; Allen, E. (Univ. of Kansas, Lawrence). Exp. Hematol.; 22: No. 1, 27-30 (Mar 1972).

Results are summarized from a number of experiments utilizing intraperitoneally injected ^{89}Sr to produce destruction of bone marrow cells in mice. This method offers promise for a variety of hematologic and immunologic investigations. (H.L.W.)

45897 RADIATION-INDUCED TRANSPLANTABLE MYELOMONOCYTIC LEUKEMIA IN BEAGLES: HEMATOLOGIC PARAMETERS. Wilson, F. D.; Shifrine, M. (Univ. of California, Davis). Exp. Hematol.; 22: No. 1, 41-4 (Mar 1972).

A radiation-induced canine disease resembling myelomonocytic leukemia in man was serially transplanted to fetal puppies using viable tumor cells. A sequence of alterations in neoplastic expression was observed with repeated transplantations. In the earliest recipients the disease closely resembled that in the original donor dog and was characterized by myelomonocytic morphology and frank leukemic manifestations in peripheral blood. With further transplantation, the monocytic component became less apparent, and the disease appeared as a more pure granulocytic leukemia. In later passages, the leukemic signs in peripheral blood were lost (aleukemic leukemia). Evidence for decreased bone marrow involvement was seen in the last group of recipients. All puppies had normal myeloid: erythroid ratios although marrow values continued to show significant elevations in percentages of blasts and primitive granulocytes. Concurrent with peripheral blood and bone marrow changes there was a relative increase in a tendency for chloromatous tumor involvement with subsequent passages. This finding suggests the tendency for malignant cells to adapt to an ascitic environment. (auth)

45898 INCREASED ANDROGEN PRODUCTION IN FISH FOLLOWING IRRADIATION. Rathi, S. K. (Univ. of Rajasthan, Jaipur, India). Sci. Cult. m(Calcutta); 37: No. 12, 583-4 (Dec 1971).

Phosphorus-32 was administered internally to immature male *Heteropneustes fossilis* and controls were given an equal dose of stable H_3PO_4 . Ten days later the seminal vesicles of the irradiated fish exhibited a 34% increase in weight and hypertrophy was observed in histological preparations. These changes were attributed to an increased production of androgen. There was no change in the seminal vesicles of the control fish. (H.L.W.)

45899 HYGIENIC INVESTIGATION ON CONSUMPTIBILITY OF IRRADIATED COMPLETE RAT FEED. Barna, J. (Kozponti Elelmiszeripari Kutató Intézet, Budapest). Izotoptechnika; 15: No. 3 3-4, 217-24 (1972).

Malignant or pathological effects or changes in weight increase, organ weight, feed consumption, number of descendants, and blood picture of rats fed with LATI feed irradiated with 2.5 and 4.5 Mrad, respectively, could not be observed. (auth)

45900 EFFECTS OF ^{90}Sr AND ^{137}Cs ADMINISTERED CONTINUOUSLY UPON MICE. VIII. ABNORMALITY OF REPRODUCTIVE FUNCTION. Nishio, Kazuo. Annu. Rep. Radiat. Center Osaka Prefect.; 12: 111-12 (1971).

Reproduction of mice administered continuously with ^{90}Sr and ^{137}Cs through drinking water was observed with regard to litter size and reproduction-span (period between 1st and last birth). Propagation of mice was done by brother-and-sister mating being contained in the same cage during the entire life. Concentration of the isotopes in water is composed of three levels. High level is $0.1 \mu\text{Ci}$ of ^{90}Sr and $0.4 \mu\text{Ci}$ of ^{137}Cs per ml; intermediate level is $1/10$ and low level is $1/100$ respectively. In the low group, total litter size slightly decreased and reproduction-span was delayed and

lengthened. Intermediate and high group showed decrease of total litter size and shortening of reproduction-span except partial increase of litter size in first birth in intermediate group. (auth)

45901 BONE AND SOFT-TISSUE NEOPLASMS IN CATS EXPOSED TO RADIOSTRONTIUM. Ward, Jerrold M.; Wright, James F.; Nelson, Neal S.; (and others) (Twinbrook Research Lab., Rockville, Md. National Cancer Inst., Bethesda, Md.). J. Nat. Cancer Inst.; 48: No. 5, 1543-5 (May 1972).

Four neoplasms of bone and soft tissues adjacent to bone developed in three cats from 274 to 609 days after initial oral administration of strontium-89. Neoplasms included a giant cell tumor, osteosarcoma, hemangiosarcoma, and fibrosarcoma. Cumulative dose to the skeleton of each cat varied from 1700 to 4850 rads. Feline leukemia virus was not detected in the neoplasms by a complement fixation test or ultrastructural examination. Comparative aspects of radionuclide carcinogenesis are discussed. (auth)

45902 DETERMINATION OF THE ^{131}I DOSE TO THE MOUSE THYROID. Walinder, G. (Foersvarets Forskningsanstalt, Stockholm). Acta Radiol., Ther., Phys., Biol.; 10: No. 6, 558-78 (Dec 1971).

The radiation dose to the thyroid glands in fetal and adult mice after injections of ^{131}I was calculated by a mathematical model based on film measurements in excised glands and determinations of the thyroidal uptake and retention of the nuclide. The present measurements confirmed Loevinger's semitheoretic determinations of the β -dose to the adult mouse thyroid. (auth)

Plants

Refer also to abstract: 45529.

45903 MONITORING OF THE RADIOACTIVE CONTAMINATION OF WHEAT FLOUR. Coulon, R. (CEN, Fontenay-aux-Roses, France). Union Agr.; No. 325, 53-5 (Nov 1971). (In French).

A regional monitoring of the radioactive pollution of flours has been carried out in France since 1960. Samples from ten production regions were ground and studied. The results given the ^{137}Cs and ^{90}Sr concentration (in picocuries) per kg of flour, and per gram of K or per gram of Ca respectively. The values obtained for the 1968 and 1969 crops, and the mean values for the 1960-1969 period are presented. (France)

Nuclide Kinetics and Toxicology

Refer also to abstract: 45553.

Man

Refer also to abstract: 45538.

45904 (COO-3013-1) RADIATION HAZARDS IN ENVIRONMENTAL MEDICINE. Fabrikant, Jacob I. (Johns Hopkins Univ., Baltimore, Md. Dept. of Radiological Sciences). 5 Aug 1969. Contract AT(30-1)-3970. 12p. Dep. NTIS.

Radiation hazards to the health of human populations following the uptake and internal deposition of radioisotopes released to the atmosphere are considered from the standpoint of the contribution of radioisotopes released during the operation of nuclear power reactors. It is pointed out that during 1969 the atomic energy industry of the United States comprised a vast network of uranium mines, industrial plants, and laboratories that employed more than 100,000 people. While all these applications of nuclear energy have the capacity to contaminate the environment, with very few exceptions, the amounts of radioactive substances released so far have been so small that there has been no significant exposure of the public. Data are summarized on the uptake of ^{137}Cs , ^{131}I , and ^{90}Sr by man following release in fallout during nuclear weapons testing, and on the contribution of natural background radiation and diagnostic and therapeutic medical uses of radiation sources and radioisotopes to the radiation dose to man. It is concluded that, while the development of more nuclear reactors for power generation will involve more sources for potential radioactive contamination of man's environment, under normal operating conditions a nuclear reactor is a source of relatively insignificant amounts of environmental radioactivity and the safety devices built into the reactors and the stringent restrictions on the release of radioactive substances to the environment should ensure that the nuclear power industry does not present a threat to public health in the United States. (C.H.)

45905 (RD/B/N-2291) MULTI-COMPARTMENT MODEL FOR STUDYING THE DOSE-INTAKE RELATIONSHIP FOR RADIOACTIVE ISOTOPES. Clarke, R. H. (Central Electricity

Generating Board, Berkeley (England). Berkeley Nuclear Labs.). Apr 1972. 38p. Dep. NTIS (U. S. Sales Only).

In identifying the organs at risk after the intake of radioactive material and in order to evaluate the doses in those organs, it is necessary to use a model of the human body in which the transfer of radioactive materials is amenable to mathematical analysis. Such a model was developed in which the various organs of the body were represented by single or multiple compartments, coupled to provide a simplified metabolism. The method of entry of radioactive material could be by inhalation, ingestion, through contaminated skin or a wound, using standard models for the respiratory and gastrointestinal tracts. The distribution of a parent nuclide and all its daughter products throughout the compartments was followed as a function of time by the solution of the set of first order differential equations that represent transport within the body. Either acute or chronic exposures were treated and facilities existed for varying metabolic parameters with time. The physical basis of the model is described together with the techniques used to solve the coupled equations to obtain retention and dose data. (auth)

45906 (TT-54002/4), pp 462-70) STATISTICAL DATA ON IODINE UPTAKE OF THE THYROID GLAND. PART II. ^{131}I UPTAKE AFTER 6 AND 24 HOURS ($T_{6/24}$). Doroszewski, Jan; Niewiadomski, Michał (Postgraduate Medical Education Centre, Warsaw. Inst. of Radiology, Warsaw). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

When interpreting the results of iodine uptake tests, not only the general level of ^{131}I uptake by the thyroid, but also its dynamics is significant. Accurate determination of the changes in ^{131}I content of the thyroid would require repeated measurements of radioactivity of the gland at intervals after administration of the diagnostic dose of the indicator, which would not be feasible in ambulatory practice. Some information about ^{131}I content of the thyroid can be obtained, however, by making only two measurements at different times. A widely used method for studying iodine uptake on the basis of two measurements consists in measuring ^{131}I activity in the thyroid gland 6 and 24 hrs after administration of the diagnostic dose $T_{6/24}$. (auth)

45907 HUMAN FAECAL ^{226}Ra EXCRETION SHORTLY AFTER A RaSO_4 DUST CONTAMINATION, WITH ALGINATE TREATMENT. Vander Borgh, O.; Colard, J.; Boulenger, R. (Centre d'Etude de l'Energie Nucleaire, Mol, Belgium). Health Phys.; 23: No. 2, 240-3 (Aug 1972).

Data are presented on the excretion of ^{226}Ra after oral uptake by a 50 yr old healthy woman following the breakage of a 17 yr old therapeutic ampoule containing 50 mCi $^{226}\text{RaSO}_4$. A regular diet of alginated bread, resulting in an intake of 15g alginate per day, was eaten until the 20th day after contamination. The fecal ^{226}Ra contents per day for 160 days are tabulated and compared with previous data presented by Marinelli (1958). Measurements made in a whole-body counter are given. Much data on the case were lost and the reasons are given. On day 12 a residual body burden of 1.3×10^4 pCi ^{226}Ra was measured representing about 10% retention. (UK)

45908 LONG-TERM STUDY OF THE URINARY ^{90}Sr AND CALCIUM IN CHILDREN. Czosnowska, W. (Inst. of Nuclear Research, Warsaw); Pietrzak-Flis, Z.; Grabowski, D. Health Phys.; 23: No. 2, 215-21 (Aug 1972).

The intake and the excretion of ^{90}Sr and Ca in the urine of seven children, aged 4 to 7 yr, were observed during 28 consecutive months. The average intake was 12.4 pCi/day of ^{90}Sr and 539 mg/day of Ca. The average urinary ^{90}Sr was 0.71 pCi/day, while urinary Ca amounted to 33.0 mg/day. The average urine-diet observed ratio was 1.33. It was concluded that the urine-diet observed ratio and urinary Ca was connected with the same relationship for each subject studied and that the equation obtained for children could be an extension of the relationship obtained for adult Polish people. (TSS)

45909 ^{22}Na IN THE FOOD-CHAIN: LICHEN-REINDEER-MAN. Soeren Mattsson, L. J. (Univ. of Lund, Sweden). Health Phys.; 23: No. 2, 223-30 (Aug 1972).

The ^{22}Na content of samples of lichen, reindeer-blood, -flesh and -bone collected between 1961 and 1969 in the area of Funäsdalen in central Sweden (62.5°N, 12.5°E) was measured by means of a sum-coincidence γ -spectrometer. After the thermonuclear detonations in 1961 and 1962, the radioactive concentration of ^{22}Na in reindeer-flesh rose rapidly from about 1 to 10 pCi per kg to a maximum value of 200 pCi per kg wet weight in late 1964. Early in 1969 the same value had decreased to 25 pCi per kg wet weight. The radioactive concentration of ^{22}Na in reindeer-bone reached a maximum of 600 pCi per kg wet weight in mid-1965. The biological half-time of sodium incorporated in reindeer-bone was estimated to be about 0.5 yr. Analyses of γ -spectra from the annually per-

formed whole-body measurements on Lapps give a ^{22}Na -body burden of less than 2 nCi at the time of assumed maximum. The presence of ^{22}Na in the food-chain: lichen-reindeer-man has given a small contribution to the absorbed dose in man in comparison to many other bomb-produced and naturally occurring radionuclides. (auth)

Animals

Refer also to abstracts: 45545 and 45902.

45910 (CONF-720505-2) INCREASED EFFECTIVE HALF-LIFE OF INTRATRACHEAL ADMINISTERED $^{51}\text{Cr}_2\text{O}_3$ BY RESPIRATORY INFECTION. Creasia, D. A.; Nettesheim, P.; Hammons, Anna S. (Oak Ridge National Lab., Tenn.). 1970. 22p. Dep. NTIS.

From twelfth Hanford biology symposium on radionuclide carcinogenesis; Richland, WA. (11 May 1972).

The effect of lower respiratory tract infections on the clearance of radioactive materials from the lungs was studied. Respiratory infections were produced in pathogen-free mice by the intranasal inoculation of influenza virus. Chromium-51 was administered as chromium oxide by intratracheal injection and the effective half-life in the lungs was determined. It was found that the $T_{1/2}$ was increased in infected mice 1.3 to 2.1 times over that in uninfected mice when the $^{51}\text{Cr}_2\text{O}_3$ was administered prior to the acute stages of infection. When $^{51}\text{Cr}_2\text{O}_3$ was administered after the acute infectious episode, the $T_{1/2}$ values were lower but still represented an increase over those in uninfected mice. It is pointed out that the degree of retention was determined, not by the infection per se, but rather by the extent and persistence of inflammatory and scarring processes. Other types of respiratory infections and injuries induced by noninfectious agents may have similar effects. (E. R. B.)

45911 (COO-3013-25) RADIOBIOLOGY AND THE PUBLIC HEALTH. Fabrikant, Jacob I. (Connecticut Univ., Farmington. Dept. of Radiology). [1968]. Contract AT(30-1)-4244. 48p. Dep. NTIS.

Results are reviewed from a number of studies on the effects of radiation delivered at low doses and low dose rates on cells, tissues, and organisms. Emphasis is placed on the effects of chronic internal irradiation at the cellular and molecular level and repair mechanisms that ameliorate the effects of radiation injuries. It is pointed out that the radiosensitivity of mammalian cells varies enormously depending on species, organ, mitotic activity, physiological state, and environmental factors, as well as radiological variables such as dose, dose rate, and LET. It is emphasized that direct quantitative extrapolation of results of radiobiological studies on laboratory animals to man is not possible. Experimental evidence indicates that exposure to any radiation at any dose and dose rate involves some risk of deleterious effects. However, levels of natural radioactivity existed during the early evolution of man that were of higher intensity than levels existing at present. The control of environmental levels of radioactivity has become a sociological and political problem and unless man wishes to dispense with medical and industrial activities that involve low dose exposure to ionizing radiations he must evaluate the risks and limit the radiation exposure to a dose level at which the risk is acceptable to the individual and to society. (C. H.)

45912 (ORO-3728-5) MUTAGENIC EFFECT OF TRANSMUTATION FROM RADIONUCLIDES INCORPORATED INTO DNA OF DROSOPHILA MELANOGASTER. Progress Report, September 1970-May 1972. Lee, William R. (Louisiana State Univ., Baton Rouge. Dept. of Zoology and Physiology). Contract AT(40-1)-3728. 8p. Dep. NTIS.

Experiments were conducted to determine the genetic effects of transmutation of ^{33}P and this was compared with previous results of genetic effects of transmutation of ^{32}P and ^{14}C . It was found that approximately 50% more ^{33}P can be fed to *Drosophila* than ^{32}P without causing excessive sterility. In experiments with ^{33}P an increase in the mutation rate was observed in the non-stored as well as stored spermatozoa due to beta radiation during the labeling process. This elevation to about a 4% sex-linked recessive lethal rate observed in the F_2 at the beginning of storage makes the 0.13% spontaneous rate during storage insignificant. Therefore, with neither a significant increase in mutation due to beta radiation nor spontaneous mutations during storage, any increase in mutations observed during storage is attributed to transmutation of the incorporated radionuclide. The negative results obtained for lethals observed in the F_3 are in contrast to previous results with ^{32}P . The significance of this is that the mutational effect observed in earlier experiments with ^{32}P is due to the recoil of the ^{32}S nuclide and not due to either change in atomic number or excitation; these two events should be the

same for both ^{33}P and ^{32}P . A series of doubly labeled experiments using ^3H and ^{33}P was conducted to determine what percentage of ^{33}P in the cell was located in the DNA molecule. (H.L.W.)

45913 (RLO-2227-T-8-3) RADIONUCLIDE CYCLING IN NATURAL POPULATIONS OF AMPHIBIANS. Annual Progress Report, June 16, 1971–June 15, 1972. Willis, David L. (Oregon State Univ., Corvallis. Dept. of General Science). 1972. Contract AT(45-1)-2227. 17p. Dep. NTIS.

Progress is reported on radionuclide metabolism studies in the newt (*Taricha granulosa*). Whole-body retention and tissue distribution of ^{51}Cr , ^{54}Mn , ^{59}Fe , and ^{181}W were investigated. Results are presented in the form of graphs and the radioecological significance is discussed. Studies were also conducted on effects of chronic exposure to ^{134}Cs on the developing frog (*Rana pretiosa*). (H.L.W.)

45914 (TID-25954, pp 58-64) PERSISTENCE OF RADIONUCLIDES IN SOIL, PLANTS, AND SMALL MAMMALS IN FALL-OUT CONTAMINATED AREAS. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

The persistence of radionuclides in soil, plants, and small mammals was investigated in 1966 in areas that had been contaminated with fallout from above-ground nuclear detonations at the Nevada Test Site during the period 1952–1959. Study sites were established at various locations out to about 225 km from ground zero. Emphasis was placed upon the movement of ^{90}Sr and ^{137}Cs from abiotic to biotic components. Several neutron activation products were also studied in fallout areas located within 5 km of recent nuclear excavation tests. Radionuclides continued to be taken up through plant roots in small amounts as time progressed, and some continued to be deposited on foliage as resuspended dust particles. The inhalation route of entry became less important with passing time, whereas ingestion continued to be the most important route through which radionuclides entered small mammals living in old fallout areas. Long lived ^{90}Sr accumulated primarily in bone tissue while ^{137}Cs accumulated in muscle and soft tissue. Most of the neutron activation products are short-lived, but among those found in animal tissues were isotopes of cobalt, manganese, and tungsten. Findings indicate that ^{90}Sr and ^{137}Cs will continue to move in small amounts from abiotic to biotic components in fallout-contaminated areas with passing time. (auth)

45915 (UR-3490-115) EXPERIMENTAL INHALATION OF METALLIC SILVER. Phalen, Robert F.; Morrow, Paul E. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1971. 33p. Dep. NTIS.

Dogs were tracheally intubated and exposed to airborne particles produced by exploding ^{110m}Ag wires. The particles were agglomerates of spherical primaries and had activity median aerodynamic diameters near 0.5 microns. The deposition fraction in the respiratory tract was 0.17 ± 0.05 ; absolute deposition averaged about 1 $\mu\text{g}/\text{kg}$ of body weight. Translocation and clearance of the deposited material was followed by external gamma-ray detectors collimated to measure activity at several sites in each animal. Clearance data were analyzed into exponential components. The lung had biological clearance half-lives of 1.7, 8.4, and 40 days accounting respectively for about 59%, 39%, and 2% of the total amount deposited. Corresponding values for the liver were 9 days (97%) and 40 days (3%). Excretion was predominantly fecal, the major portion of which was believed to have been secreted into the bile. Sacrifice measurements showed that the liver was a major site for the deposition of silver. Lung, brain, and muscle also contained relatively large accumulations. Results showed that existing ICRP biological values for evaluating the hazards from radioactive silver are unreliable for inhalation. For example, considerably more silver (perhaps 90% of that initially deposited in the lungs) was absorbed into the body for general internal distribution, than would be calculated from the ICRP figures. (auth)

45916 (UR-3490-159) RETENTION OF ^{210}Pb IN BEAGLE DOGS. Hursh, John B. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1971. 20p. Dep. NTIS.

A study involving four beagle dogs injected intravenously with ^{210}Pb and observed for two years has yielded data indicating a marked similarity between the retention of lead and the retention of strontium by the beagle dog. (auth)

45917 METABOLISM OF MONOMERIC AND POLYMERIC PLUTONIUM IN THE RABBIT. COMPARISON WITH THE MOUSE. Rosenthal, M. W.; Lindenbaum, A.; Russel, J. J.; Moretti, E.; Chladek, D. (Argonne National Lab., Ill.). Health Phys.; 23: No. 2, 231-8 (Aug 1972).

Young adult female Dutch Belted rabbits were given a single intravenous injection of polymeric ^{239}Pu -nitrate (15% ultrafilter-

able) or of monomeric ^{239}Pu -citrate (90% ultrafilterable) and killed at 3 days. The plutonium content of the intact femur (bone plus marrow) was higher after the polymeric form. Physical removal of marrow from bone in the tibia showed the plutonium content of the bone to be slightly lower and of the marrow to be 17-fold higher after polymeric than after monomeric plutonium. Polymeric plutonium was also more concentrated in spleen, ovaries, intestines (with contents), skeletal muscle, and urine but less concentrated in liver. About the same levels of the two forms of plutonium were found in blood, bile, and lung. For the first 30 min after injection, the polymeric plutonium was cleared from the circulation more rapidly than the monomeric. After about 60 min, the two forms were cleared at generally comparable, decreasing rates. Polymeric plutonium was as much as 10 to 30 times more concentrated in predominantly red (erythropoietic) marrow than in fatty marrow. Monomeric plutonium was 1.3 to 2.8 times more concentrated in the red marrow. Mice injected with the same plutonium solutions as the rabbits showed, in comparison, a higher percent of the injected monomeric plutonium in the femurs and a lower percent in the liver; they showed a higher uptake of polymeric plutonium in the marrow, and a higher uptake of both forms in the spleen and lungs than did the rabbits. (auth)

45918 EFFECT OF ALUMINUM PHOSPHATE GEL ON WHOLE-BODY RETENTION OF SIMULTANEOUSLY ADMINISTERED ^{226}Ra , ^{85}Sr , AND ^{47}Ca IN MICE. Keslev, D.; Van Puymbroeck, S.; Van der Borgh, O. (SCK/CEN, Mol, Belgium). Experientia; 28: No. 5, 524-5 (15 May 1972).

A mixture of ^{226}Ra , ^{85}Sr , and ^{47}Ca was administered by gastric tube to mice and each animal received aluminum phosphate gel by gastric tube; control animals received agar gel. A table is presented to show total body retention of the administered dose four days after gastric intubation. It was found that aluminum phosphate gel, administered simultaneously with the isotopes, reduced the ^{226}Ra burden by a factor of 800, the ^{85}Sr burden by a factor of 10, and the ^{47}Ca burden by a factor of 3. (H.L.W.)

45919 VOPROSY PRODUKTSIONNOI, SANITARNOI I TEKHNICHESKOI GIDROBIOLOGII YUZHNYKH MOREI. (Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas). Zorina, T. A. (ed.). Kiev: Izdatel'stvo Naukova Dumka (1971). 196p.

This book is a collection of 19 papers by various authors on primary productivity in biological systems, and the hydrologic and hydrochemical conditions which prevail in the southern seas (Black Sea, Red Sea, and Mediterranean Sea). Two papers are concerned with the effects of ^{137}Cs and ^{90}Sr on the development of embryos of fish and sea urchins, and the cytogenetic effects resulting from the uptake of artificial radionuclides (^{137}Co , ^{90}Sr , ^{90}Y , ^{91}Y , and ^{14}C) by marine organisms. (K.S.W.)

45920 MEASUREMENT OF ^{226}Ra IN MICE. Lenger, V. (Inst. of Radiation Hygiene, Prague). Acta Radiol., Ther., Phys., Biol.; 10: No. 5, 488-96 (Oct 1971).

Low Compton scattering affords means of direct measurement of the ^{226}Ra 187 keV line, the quantitative evaluation being based on the phantom calibration procedure. The nonequilibrium state is an advantage. The retention curves of ^{226}Ra in mice during a period of up to 352 days are presented and the kinetics of daughter products, especially of ^{222}Rn , proved. (auth)

45921 DIFFUSION DEPOSITION OF SMALL-SIZED PARTICLES IN THE BRONCHIAL TREE. Martin, D.; Jacobi, W. (Hahn-Meitner-Institut fuer Kernforschung, Berlin). Health Phys.; 23: No. 1, 23-9 (Jul 1972).

The deposition pattern of radioactive aerosols ($\text{AMI} = 0.2-0.4 \mu\text{m}$) was studied in a plastic model of the upper bronchial tree, whose geometry corresponded with a dichotomous lung model. The deposition probabilities w in the different bronchial regions for inspiration and expiration were measured as a function of air flow rate v in the range of 0.1 to 50 l/min. Radioautographs were also prepared to study the fine structure of the deposition pattern. The measured w -values can be approximated by the relation $w = k \cdot v^{-n}$ ($n > 0$), where k is a characteristic factor for each generation and nearly equal for inspiration and expiration. The slope-coefficient n amounts to 0.65 ± 0.07 for all bronchial generations. The results indicate that deposition due to turbulent diffusion plays an important role in the trachea and larger bronchi. With increasing depth in the bronchial tree, however, the measured w -values approach the theoretical values, calculated with the theory of Gormley and Kennedy (Proc. Proj. Irish Acad. 52A: 163 (1949)), for diffusion in a laminar air stream. (auth)

45922 EFFECT OF MASS ON PROMETHIUM RETENTION AND REMOVAL FROM THE RAT. Smith, V. H. (Battelle-Northwest, Richland, Wash.). Contract AT(45-1)-1830. Health Phys.; 23: No. 1, 31-9 (Jul 1972).

Seven days after intravenous administration to rats of 3.32×10^{-4} to 5.36×10^3 mmole of isotopic or carrier-diluted PmCl_3 the fraction of injected monomeric promethium retained in the skeleton was greatest at the lowest mass administered. Generally no mass effect was observed in kidney, liver or spleen at the monomeric mass levels administered. The effectiveness of 0.3 mmole diethylenetriaminepentaacetic acid (DTPA) in decreasing the fraction of injected promethium retained in the skeleton following prompt (1 hr) or delayed (48 hr) treatment was less for the lowest mass injected than for other monomeric masses. In the soft tissues a relatively constant fraction of the control rats' promethium tissue burden was removed by DTPA treatment at all monomeric mass levels. At the highest mass level particulate formation typically altered the distribution as well as effectiveness of DTPA treatments. While the amount of promethium removed from most of the examined tissues by DTPA treatments at each monomeric level was an almost constant fraction of that retained in the control tissues, the moles of promethium removed increased with injected mass as the molar ratio of DTPA : Pm varied from 10^9 to 10^5 . In extending the results of animal experiments and tracer-level human experiments as predictors of radionuclide retention and of the efficacy of removal procedures in accidental human exposure situations, due regard should be given to the possible effect of mass as well as to the physical state of the incorporated radionuclide. (auth)

Plants

Refer also to abstracts: 45550 and 45755.

45923 (TID-25954, pp 69-72) ^{90}Sr UPTAKE BY DESERT VEGETATION. 1972.

In Radioecology and ecophysiology of desert plants at the Nevada Test Site.

The levels of ^{90}Sr uptake varied among different species of desert vegetation. A species of range grass, *Oryzopsis hymenoides*, had the lowest ^{90}Sr content of the species investigated and, perhaps significantly, the lowest specific activity of ^{90}Sr per stable strontium of all the species. Highest uptake of ^{90}Sr occurred in *Atriplex confertifolia*, *Atriplex hymenelytra*, and *Lycium andersonii*. Nearly the same levels of ^{90}Sr concentrated in leaf and stem tissues of *Artemisia tridentata*, *Astragalus lentiginosus*, and *Larrea divaricata*. In *Atriplex confertifolia* and *Atriplex hymenelytra* the ^{90}Sr was concentrated highest in stem tissue. The other species studied concentrated as much as three times more ^{90}Sr in leaf tissues as in stems. A similar distribution pattern occurred for the stable strontium levels in leaf and stem tissues of these desert plant species. A higher correlation coefficient was obtained for $^{90}\text{Sr} \times \text{strontium}$ than for $^{90}\text{Sr} \times \text{other cations}$ or sum of cations. The leaf-stem ratio for the ^{90}Sr was more correlated with the leaf-stem ratio for stable strontium than with other cations. There was a positive correlation between specific activity of ^{90}Sr and the cation sum of leaves. (auth)

45924 PLUTONIUM AND POLONIUM INSIDE GIANT BROWN ALGAE. Wong, K. M.; Hodge, V. F.; Folsom, T. R. (Univ. of California, San Diego). Nature (London); 237: No. 5356, 460-2 (23 Jun 1972).

Certain marine algae accumulate Pu so effectively that they might be used to detect small concentration changes of the element in seawater, where direct determination would be difficult. In a survey made in 1971, near a coastal power reactor after reported accidental burnup of fuel rods carrying 17,000 Ci of ^{238}Pu , Pu and other nuclides were compared in various types of algae, and large accumulations were found in several species; concentrations, however, ranged widely within species and suggested a correlation between activity and surface area. Attention was turned to comparing different tissues of one giant brown algae, *Pelagophycus porra*. Samples dissected for Pu analysis by α -spectrometry were also analyzed for ^{210}Po ; it was hoped that the distribution of ^{210}Po might serve as a model for predicting distributions of the less easily analyzed ^{239}Pu and certain other heavy metals. Concentrations of ^{210}Po varied significantly between different parts of the plant, and even between different sections of the same part, but were always much higher in the outermost layers; this also applied to ^{239}Pu concentrations, but to a lesser degree. Another finding was that for comparison of two different environments it is not sufficient merely to sample a given species, but identical sample tissue must be compared. (UK)

45925 EFFECT OF SOIL CONDITIONS AND POTASSIUM SUPPLY ON THE UPTAKE AND DISTRIBUTION OF ^{86}Rb IN THE ORGANS OF CORN PLANTS. Ivanov, S. N.; Nikitenko, N. F. Dokl. Akad. Nauk BSSR; 16: No. 2, 141-3 (Feb 1972). (In Russian).

The uptake of ^{86}Rb by the entire corn plant and by its tissues in less fertile soils with low exchange potassium content proceeds at a higher rate than in soils which are well-fertilized and containing considerable exchange potassium. In well-limed soils ^{86}Rb uptake by corn is 1.5 to 2 times lower per unit weight than in un-limed. In the staling phase the ^{86}Rb content in the stems is 1.5 to 2 times greater than in the leaves. With the appearance of the reproductive organs the ^{86}Rb content per unit weight of biomass is distributed as follows: highest in the panicle, then the leaves, and least in the stems. (tr-auth)

45926 STATISTICS OF RADIOISOTOPE ACCUMULATION IN HYDROBIONTS. Piskunov, L. I. (Sverdlovsk Regional Health-Epidemiological Station). Hidrobiol. Zh.; 6: No. 5, 70-2 (1970). (In Russian).

Hydrobionts, especially fresh-water plants, are distinguished by an asymmetric exchange of elements which leads to an accumulation of certain elements until a steady-state condition is reached. A statistical study was made of this type of accumulation in hydrobionts, and certain regularities in the process were found. The correlating connection in the complex pattern of micro-element exchange should be regarded as only one of the possible mechanisms for radioisotope migration within the biological chain. (K.S.W.)

New Tracer Techniques

45927 (TT-54002/4, pp 471-4) METHOD FOR LABELING HUMAN ALBUMIN WITH $^{99\text{m}}\text{Tc}$. Sadowska, Barbara; Zelenay, Krystyna (Inst. of Nuclear Research, Swierk, Poland). Translated from Przegl. Radiol. Med. Nukl.; 35: No. 4, (1971).

A simple method for labeling human albumin with technetium- $^{99\text{m}}$, which can be easily adapted for clinical purposes, is described. The yield of the radiochemical method, after purification, was 60%. Clinical investigations showed a complete usefulness of the obtained preparation and controls of radiochemical purity proved that the content of technetium- $^{99\text{m}}$ not bound to albumin is much below 5%. (auth)

45928 SIMULTANEOUS RADIOIMMUNOASSAY OF PLASMA TESTOSTERONE AND DIHYDROTESTOSTERONE. Coyotupa, Juan; Parlow, Albert F.; Abraham, Guy E. (Univ. of California, Torrance). Anal. Lett.; 5: No. 6, 329-40 (1972).

A method is described for the simultaneous radioimmunoassay of plasma testosterone (T) and 5α -dihydrotestosterone (DHT), using as binding reagent a highly specific antiserum which reacted significantly with T and DHT. Utilizing a one step celite microcolumn to separate T from DHT, we have measured these two androgens accurately in small aliquots of plasma from both male and female subjects. Plasma concentrations of T and DHT expressed as mean $\pm$ S.D. in ng/ml were respectively: in male subjects; normal adult: 4.9 ± 1.6 (T) and 1.3 ± 0.8 (DHT); hypogonadal: 0.24 ± 0.08 (T) and 0.20 ± 0.10 (DHT); prepubertal: 0.14 ± 0.02 (T) and 0.05 ± 0.02 (DHT); in female subjects; normally menstruating, irrespective of time of cycle: 0.22 ± 0.07 (T) and 0.34 ± 0.17 (DHT); postmenopausal: 0.29 ± 0.08 (T) and 0.15 ± 0.07 (DHT); ovariectomized: 0.24 ± 0.06 (T) and 0.09 ± 0.01 (DHT). When plasma levels of T and DHT were measured during a complete menstrual cycle, the levels of these steroids were found to be higher during the luteal than during the follicular phase. The ratio of plasma T : DHT concentrations was five times higher in normal adult male (3.1) than in menstruating female subjects (0.6). (auth)

45929 BLEOMYCIN LABELED WITH ^{57}Co . APPLICATION TO THE DIAGNOSIS OF MALIGNANT TUMORS AND THEIR GROWTH. Nouel, J. P. (Centre Henri Becquerel, Rouen); Renault, H.; Robert, J.; Wicart, L. Nouv. Presse Med.; No. 2, 95-8 (8 Jan 1972). (In French).

Intravenous injection of ^{57}Co labeled bleomycin enables primitive cancers to be detected and unsuspected metastasis to be revealed. This method is superior to existing techniques. (France)

MATERIALS

Ceramics and Cermets

Refer also to abstracts: 45311, 45407-45417, 45452, 45973, 45979, 45992, 45997, 46002, 46013, 46039, 46074, 47244, 47249, and 47255.

45930 (HEDL-TME-72-81) EFFECTS OF PORE STRUCTURE ON THE MELTING HEAT RATING OF OXIDE FUELS. Lawrence, L. A.; Gibby, R. L. (Hanford Engineering Development Lab., Richland, Wash.). May 1972. Contract AT(45-1)-2170. 41p. Dep. NTIS.

Mixed uranium-plutonium oxide fuels prepared by different fabrication techniques can have significantly different thermal properties. These differences, which stem from differences in pore shape and distribution characteristics resulting from the fabrication process, were investigated using thermal diffusivity techniques and in-reactor experiments. Fuel characterized by large amounts of linear porosity was found to have a thermal conductivity 30 to 40% lower than fuel of equivalent density with uniformly distributed fine spherical porosity and large isometric pores. Results of the irradiations confirmed a lower melting heat rating for fuel characterized by large amounts of linear porosity. Results of the irradiations were compared to temperature profile predictions based on the laboratory thermal diffusivity data. Reasonable agreement is obtained if expected effects of oxygen redistribution down the thermal gradient in the fuel are considered. (auth)

45931 (KAPL-3164) FABRICATION OF ALUMINUM OXIDE FILMS. Hance, A. B.; Hanrahan, L. R. (Knolls Atomic Power Lab., Schenectady, N. Y.). 27 Jul 1966. Contract W-31-109-eng-52. 8p. Dep. NTIS.

A simple low-cost procedure for the forming of Al_2O_3 films from common household aluminum foil is described. A method for accurately controlling the film thickness is described for self-supporting films of Al_2O_3 ranging from several hundred to 1000 Å thick. The method can be used for anodizing a smooth aluminum surface to form an Al_2O_3 surface layer of uniform thickness ranging from ~100 Å to 1000 Å. (auth)

45932 (LBL-450) HIGH-TEMPERATURE SPECTROSCOPY AND THERMODYNAMICS OF TITANIUM OXIDE AND VANADIUM OXIDE. Chang, Shih-Ger (California Univ., Berkeley. Lawrence Berkeley Lab.). Mar 1972. Contract W-7405-eng-48. 143p. Dep. NTIS.

Thesis.
Absolute emission measurements of (0,0) band of TiO α -system and VO green system were made on photographic plates. The TiO vapors were in equilibrium with $TiO_{1.6}$ liquid phase at 2140 to 2400°K and VO vapors were in equilibrium with VO solid and liquid phases at 2000 to 2150°K in an electric furnace. The integrated radiance of the band was determined by matching the observed profiles with the profiles determined theoretically. The radiance measurements and equilibrium partial vapor pressures of TiO and VO from mass spectroscopy and thermodynamic data yield: $f_{em} = 0.144$ for TiO (0,0) band $C^3\Delta-X^3\Delta$ transition, and $f_{em} = 0.0058$ for VO (0,0) band $C^4\Sigma^-X^4\Sigma^-$ transition. These correspond to an upper bound of radiative lifetime of 2.82×10^{-8} sec for $v' = 0$ $C^3\Delta$ state of TiO and 8.65×10^{-7} sec for $v' = 0$ $C^4\Sigma^-$ state of VO. (auth)

45933 (N-72-15609) SHOCK AND VIBRATION TESTS OF URANIUM MONONITRIDE FUEL PELLETS FOR A SPACE-POWER NUCLEAR REACTOR. Adams, Donald W. (National Aeronautics and Space Administration, Cleveland, Ohio. Lewis Research Center). Feb 1972. 31p. (NASA-TM-X-2493; E-6584). NTIS.

Shock and vibration tests were conducted on cylindrically shaped, depleted, uranium mononitride (UN) fuel pellets. The structural capabilities of the pellets were determined under exposure to shock and vibration loading which a nuclear reactor may encounter during launching into space. Various combinations of diametral and axial clearances between the pellets and their enclosing structures were tested. The results of these tests indicate that for present fabrication of UN pellets, a diametral clearance of 0.254 millimeter and an axial clearance of 0.025 millimeter are tolerable when subjected to launch-induced loads. (auth) (STAR)

45934 (N-72-18585) DEVELOPMENT OF DISPERSION-STRENGTHENED Ni-Cr-ThO₂ ALLOYS FOR THE SPACE-SHUTTLE THERMAL PROTECTION SYSTEM. Blankenship, Charles P.; Saunders, Neal T. (National Aeronautics and Space Administration, Cleveland, Ohio. Lewis Research Center). 1972. 33p. (CONF-720301-6; NASA-TM-X-68024; E-6830). NTIS.

From Spring conference of the American Society for Nondestructive Testing; Los Angeles, Calif. (13 Mar 1972).

Manufacturing processes were developed for TD-NiCr providing small sheet (45 × 90 cm), and larger sheet (60 × 150 cm) and foil. The alternate alloy, DS-NiCr, was produced by pack-chromizing Ni-ThO₂ sheet. Formability criteria are being established for basic sheet forming processes, which are brake forming, corrugation forming, joggling, dimpling, and beading. Resistance spot welding (fusion and solid state), resistance seam welding, solid state diffusion welding, and brazing are included in the joining programs. Major emphasis is centered on an Al-modified Ni-Cr-ThO₂ alloy development. These alloys, containing 3 to 5% Al, form the protective Al_2O_3 scale. This enhances oxidation resistance under reentry conditions. Both TD-NiCrAl and DS-NiCrAl alloys are included. A tentative composition of Ni-16Cr-3.5Al-2ThO₂ was selected based on oxidation resistance and fabricability. (auth) (STAR)

45935 (SC-RR-710884) OXIDATION AND ETCHING OF 80 PERCENT Si-20 PERCENT Ge. Fuller, C. T.; Loescher, D. H. (Sandia Labs., Albuquerque, N. Mex.). Dec 1971. 27p. Dep. NTIS.

Ellipsometry has been used to study the oxidation and chemical etching of 80% Si-20% Ge alloy. Oxidations were performed at 25, 50, and 800°C in air and at 85°C in water. The rate of oxidation at 25 and 50°C is low and is similar to that of Si. The rate of oxidation is rapid enough in 85°C water that the use of lower temperatures for washing samples is indicated. At 800°C the oxidation behavior of the alloy is like that of Si. For oxidation at 800°C and for times longer than about 30 minutes, the indices of refraction of the alloy surface and the oxide are almost identical to those of silicon. Estimates of oxide thickness on the alloy can be obtained from published data for Si. The data for oxidation of the alloy suggest that the Ge concentration at the alloy surface and in the oxide is depleted during oxidation but the mechanism for depletion is not understood. It was found that chemical etching is best accomplished with solutions of HF and HNO₃ containing at least 70% HNO₃. (auth)

45936 (UCRL-Trans-10632) CRYSTALLIZATION OF BORON-RICH BERYLLIUM BORIDES AND ALUMINUM BERYLLIUM BORIDES FROM ALUMINUM MELTS. Becher, H. J. Translated for Univ. of California Lawrence Livermore Lab., Livermore, from Acta Crystallogr., 17: No. 5 617(1964). 4p. Dep. NTIS.

In experiments with the system Be/Al/B, the compound Be₂B was finely powdered and fused with ten times its weight of Al for two hours in a sintered corundum crucible under argon at 1450°C. After treatment of the cooled regulus with HCl, a crystalline, dark residue consisting mainly of octahedral and icosahedral crystals having an average size of 0.05 mm remained. Analysis of the crystals gave the formula BeB₂. In other experiments, the B-rich compound BeB₁₂ was fused with 15 times its weight of Al. Together with impurities, the product was found to contain many reddish crystals, some being octahedral and other hexagonal leaflets. The identity periods of the α AlB₁₂ lattice were found from rotating-crystal photographs. From density determinations, it was found that on crystallization from the Al melt, BeB₁₂ and Al formed solid solutions having the composition (BeB₆)₂AlB₁₂, corresponding to a reaction $2BeB_{12} + Al \rightarrow (BeB_6)_2AlB_{12}$. It was noted that despite the large excess of Al, Be is not displaced from its union with B. (P.C.H.)

45937 THERMODYNAMIC STUDY OF ORDER-DISORDER TRANSITIONS IN VANADIUM CARBIDE. Emmons, George Herbert. Urbana, Ill.; Univ. of Illinois (1972). 175p. University Microfilms Order No. 72-19,824.

Thesis.

Differential thermal analysis was used to measure the thermodynamic properties of V_6C_5 and V_6C_7 at the order-disorder transitions of their respective carbon sublattices. The critical temperatures and enthalpy increases due to disordering were determined to be: $1250^\circ C \pm 2^\circ C$ and $6.65 \text{ cal/gm} \pm 2.90 \text{ cal/gm}$ for V_6C_5 and $1152^\circ C \pm 2^\circ C$ and $7.04 \text{ cal/gm} \pm 3.07 \text{ cal/gm}$ for V_6C_7 . The Bragg-Williams approximation was adapted to the carbon sublattice of V_6C_5 and was used to estimate properties of the order-disorder transition. An enthalpy increase due to disordering of 8.75 cal/gm was predicted using the critical temperature determined by DTA and the B-W approximation. The temperature range over which disordering is observable in V_6C_5 powders using DTA is smaller than predicted by B-W but in good agreement with the temperature range over which the effects of disordering are observable through resistivity changes in single crystal samples. The additional enthalpy increase of V_6C_5 below T_c is 33% of the total additional en-

thalpy increase of the transition. The latent heat of the transition $V_6C_5 + VC_{0.83}$ is 3.3 ± 1.0 cal/gm. The behavior of the isobaric specific heat below the critical temperature is consistent with a critical behavior described by $C_V = A \ln |\epsilon| + B$. Compositions of the V-C system between the compounds V_6C_5 and V_8C_7 (VC_x , $0.833 < x < 0.875$) undergo transitions at the critical temperatures of V_6C_5 and V_8C_7 . The ratio of the enthalpies of these two transitions is in good agreement with a ratio predicted from the carbon-to-metal ratio of these compositions, x , assuming that they are mixtures of the ordered compounds V_6C_5 and V_8C_7 . (Diss. Abstr. Int., B)

45938 TRANSPORT PROPERTIES OF THORIA AND THORIA-BASED SOLID SOLUTIONS. Etsell, T. H. (Max-Planck-Institut fuer Chemie, Mainz). *Z. Naturforsch.*; 27a: No. 7, 1138-49 (Jul 1972).

The methods for obtaining transport numbers in oxides are described and associated experimental difficulties are discussed. Their reliability is assessed and used in conjunction with related data to offer a consistent picture for thoria and thoria-based solutions between 500 and 1600°C. Both are ionic conductors within certain ranges of oxygen pressure. It is concluded that, for ThO_2 , and, for $ThO_2-Y_2O_3$ solutions (10 to 20 mole % $YO_{1.5}$), $\log P_\oplus = -4.5$, $\log P_\ominus = -60.5 \times 10^3/T + 23.3$, and $\log P_\oplus = 1.8$, $\log P_\ominus \sim -57.9 \times 10^3/T + 12.4$ where $P_\oplus$ and $P_\ominus$, respectively, are the oxygen pressures (atm) where the ionic and p-type and the ionic and n-type conductivities are equal. Ionic conductivity and charge carrier mobilities in thoria are briefly considered. (auth)

45939 FISSION RECOIL FROM SURFACE OF NUCLEAR FUELS BY IRRADIATION. Bloem, P. J. C.; Schurink, F. Ned. *Tijdschr. Natuurk.*; 37: 448-52 (11 Nov 1971). (In Dutch).

When reactor fuel is irradiated by neutrons, fission fragments with high energies are released and cause, by leaving the oxide surface, a damage of the lattice. The extent of this damage in ThO_2/UO_2 fuel, fabricated as spheres with a diameter of 5μ by the KEMA sol-gel process, was investigated. Irradiation took place in water at different temperatures, in air, in organic liquids, and in solutions. Radiation damage was measured by the amount of material which was released from the surface and which can be easily etched off from the disturbed surface. It was found that oxygen had a great influence on the damage of the oxide lattice, while KI additions to the water greatly reduced the damage by eliminating the OH-radicals. A few possible damaging mechanisms of UO_2/ThO_2 upon irradiation in water are discussed. (Netherlands)

45940 METHOD FOR THE MANUFACTURE OF FISSILE OXIDES AND THE APPARATUS DEVELOPED. Bazin, Roger; Delpech, Andre (to Commissariat a l'Energie Atomique). French Patent 2,088,170. 7 Jan 1972. Filed 25 May 1970. (In French).

A method for directly preparing uranium oxide suitable for sintering and subsequent use as nuclear fuels, from uranyl nitrate solutions, is described. A uranyl nitrate/formic acid solution is atomized in an enclosure maintained between 150 and 300°C, the product obtained is then transformed into oxide by calcination. Formic acid is added to a uranyl nitrate solution at a concentration equal to at least 0.4 ml/gram of uranium. The atomization of the solution is obtained using a nitrogen/hydrogen mixture at a temperature equal to at least that of the enclosure. (France)

45941 IMPROVEMENTS IN OR RELATING TO WELDING FLUX COMPOSITIONS FOR SUBMERGED ARC WELDING AND METHODS OF MAKING THE SAME. (to Council of Scientific and Industrial Research). Indian Patent 114,500. 13 Aug 1969.

The flux is comprised of lime, silica, magnesia, alumina, manganese oxide, chromium sesquioxide, molybdenum oxide, titanium dioxide, zirconium dioxide, and fluorine. (Indian Sci. Abstr.)

45942 HYDROGEN GENERATORS FOR NEUTRON TUBES. Bobrowski, Czeslaw; Gutowski, Ryszard; Korytowski, Adam; Milosz, Jacek. *Elektronika*; 8: No. 4, 148-51 (1972). (In Polish).

Information concerning hydrogen and metal reactions is presented especially in the case of Ti and Zr. The particular requirements applying to hydrogen generators for neutron tubes are formulated and design methods are described. (auth)

45943 CRYSTALLIZATION OF BeO , Be_2SiO_4 , AND SiO_2 FROM $Li_2MoO_4-MoO_3$. Osmer, Judith A.; Chase, A. B. (Aerospace Corp., El Segundo, Calif.). *J. Amer. Ceram. Soc.*; 55: No. 6, 292-5 (Jun 1972).

Single crystals of phenacite (Be_2SiO_4), bromellite (BeO), and tridymite (SiO_2) were grown from an $Li_2MoO_4-MoO_3$ flux. Phenacite, with rhombohedral symmetry, grew in three distinct shapes with aspect ratios (length/width) as follows: needles (>3), rods (>1.1 to 1.5), and rhombohedral-faced crystals (≈ 1). The latter grew as single crystals; the others were twinned on the (1010).

For most experiments the temperature was held constant at 1165°C and the Li_2MoO_4/MoO_3 ratio at 1/16. The growth mechanism for crystallization was the evaporation of MoO_3 . The system produced one to three phases, depending on the BeO/SiO_2 ratio. Bromellite grew until a BeO/SiO_2 ratio of 0.8 was attained. It grew as a hemipyramidal crystal having a short prism with a curved (0001) top or as a hexagonal plate. The pyramid- and prism-shaped crystals were twinned, although a few hexagonal plates were single. Tridymite grew in small hexagonal plates when the BeO/SiO_2 ratio was less than 1.5. The effects of temperature, nucleation, and flux composition on crystal shape, twinning, and occurrence are discussed. (auth)

45944 CRITICAL SURFACE ENERGIES OF Al_2O_3 AND GRAPHITE. Rhee, S. K. (Bendix Corp., Southfield, Mich.). *J. Amer. Ceram. Soc.*; 55: No. 6, 300-3 (Jun 1972).

Based on the wetting properties of $\alpha-Al_2O_3$ and pyrolytic graphite by liquid metals, the critical surface energy for spreading was determined and compared with the surface energy of the ceramics. The surface energy γ_s (ergs/cm²) of the (0001) surface of $\alpha-Al_2O_3$ was estimated to be 892–0.12T (°C), that of the c plane of pyrolytic graphite 1139–0.13T (°C), and that of the α plane of pyrolytic graphite 1300–0.17T (°C). Also, the liquid-solid interfacial energies were calculated and found compatible with the values reported in the literature. The surface energy of liquid Ag or liquid Cu was not significantly affected by carbon. (auth)

45945 PHASE ANALYSIS IN ZIRCONIA SYSTEMS. Garvie, Ronald C.; Nicholson, Patrick S. (McMaster Univ., Hamilton, Ont.). *J. Amer. Ceram. Soc.*; 55: No. 6, 303-5 (Jun 1972).

Linear calibration curves were developed for determining the content of free ZrO_2 in partially stabilized zirconia ceramics by x-ray diffraction techniques. The systems $CaO-ZrO_2$ and $MgO-ZrO_2$ were studied by two methods. The matrix method, in which free ZrO_2 was considered to be distributed in a matrix (the cubic phase), gave approximately equal mass absorption coefficients for the monoclinic and cubic phases. The polymorph technique, in which the cubic phase was considered to be a polymorph of ZrO_2 and in which integrated intensities were used, gave the better results. (auth)

45946 DIFFUSION IN THE SYSTEM $K_2O-SrO-SiO_2$. III. INTERDIFFUSION COEFFICIENTS. Varshneya, A. K.; Cooper, A. R. (Case Western Reserve Univ., Cleveland). *J. Amer. Ceram. Soc.*; 55: No. 6, 312-17 (Jun 1972).

The matrix of interdiffusion coefficients, [D], was determined by measuring spatial composition distribution in couples with compositions (in weight fractions) near 0.21 K_2O –0.17 SrO –0.62 SiO_2 after they were heat-treated for various times at 630 to 810°C. The results were consistent with the Onsager hypothesis in that [D] at a given temperature was independent of diffusion time and the direction (in composition space) of the diffusion couple. The eigenvectors and eigenvalues of the [D] matrix were determined; the diffusion paths in composition space are thus easily understood. As is consistent with intuition, the "fast" eigenvector nearly paralleled the direction of constant SiO_2 concentration. (auth)

45947 KINETICS AND BURST PHENOMENON IN ZrO_2 TRANSFORMATION. Maiti, H. S.; Gokhale, K. V. G. K.; Subbarao, E. C. (Indian Inst. of Tech., Kanpur, India). *J. Amer. Ceram. Soc.*; 55: No. 6, 317-22 (Jun 1972).

The monoclinic-tetragonal transformation in ZrO_2 was studied by DTA. The reverse transformation occurred at 850 to 1000°C, depending on the crystallite size, which was a function of the experimental variables. Isothermal kinetics were exhibited by material containing small crystallites; athermal behavior began to appear as the crystallite size increased. On cooling, strain-free samples underwent a very rapid transformation which was evident as a burst. (auth)

45948 PREPARATION OF TRANSPARENT Y_2O_3 -DOPED ThO_2 . Greskovich, C.; O'Clair, C. R.; Curran, M. J. (General Electric Corp., Schenectady, N. Y.). *J. Amer. Ceram. Soc.*; 55: No. 6, 324-5 (Jun 1972).

A powder preparation and processing method is described for production of Y_2O_3 -doped ThO_2 (YDT) ceramics with high optical quality. Microstructural and optical features of the sintered product are also presented. The results showed that grain size and porosity in YDT ceramics increased as the Y_2O_3 content increased from 5 to 15 mole %. Therefore, it is proposed that the induction period required for the onset of discontinuous grain growth is longer in ThO_2 –5 mole % Y_2O_3 than in ThO_2 –15 mole % Y_2O_3 , and fewer sites for discontinuous grain growth are present in the latter material than in the former. (P.C.H.)

45949 SURFACE TENSION, DENSITY, AND VOLUME CHANGE ON MELTING OF Al_2O_3 SYSTEMS, Cr_2O_3 , AND Sm_2O_3 . Rasmussen, J. J. (Battelle-Northwest, Richland, Wash.). *J. Amer. Ceram. Soc.*; 55: No. 6, 326 (Jun 1972).

An x-radiographic method was used to obtain data on surface tension, density, and volume change on melting of Al_2O_3 , Al_2O_3 - MgO , Y_2O_3 - Al_2O_3 , Sm_2O_3 - Al_2O_3 , Sm_2O_3 , and Cr_2O_3 . Measurement of the volume change on melting by the radiographic method had the advantage that the thermal expansion of the sample was measured in both the solid and the liquid states. Oxide additives to Al_2O_3 reduced the volume change on melting significantly. The Sm_2O_3 and Y_2O_3 were much more effective in reducing the volume change than the MgO . Data are tabulated. (P.C.H.)

45950 ACOUSTIC VELOCITY MEASUREMENTS IN THE METAL HYDRIDES, $\text{ScH}_{1.99}$, $\text{YH}_{1.93}$, AND $\text{ErH}_{1.81}$. Beattie, A. G. (Sandia Labs., Albuquerque, N. Mex.). J. Appl. Phys.; 43: No. 7, 3219-21 (Jul 1972).

Longitudinal and shear velocities were measured as a function of temperature in polycrystalline samples of $\text{ScH}_{1.99}$, $\text{YH}_{1.93}$, and $\text{ErH}_{1.81}$. From these measurements values of the limiting Debye temperature, θ_0 , the bulk modulus, Young's modulus, the shear modulus, and Poisson's ratio are calculated. (auth)

45951 NEW ELECTRICAL PROCEDURE FOR THE COMPACTION OF TECHNICAL POWDERS. Jacobi, Siegfried; Schneider, Rolf; Tradowsky, Klaus. Elektrotech. Z., A; 92: No. 8, 486-8 (Aug 1971). (In German). (KFK-1492).

A compaction procedure through which powder particles are exposed to a high, electro-dynamically-produced acceleration of 20×10^5 g is described. In contrast to usual procedures, compaction occurs without external application of pressure. By this method, relatively large rods can be obtained with homogenous and high densities approaching approximately 70% of theoretical values. The procedure was tested on powdered Al_2O_3 and UO_2 as well as on metallic powders of carbonyl iron and molybdenum. (tr-auth)

45952 DYSPROSIUM BOROCARBIDES. Bauer, Josef; Debuigne, Jean (Institut National des Sciences Appliquées, Rennes, France). Compt. Rend., Ser. C; 274: No. 13, 1271-4 (27 Mar 1972). (In French).

A ternary Dy-B-C system was studied using x-ray diffraction and metallography on samples quenched after melting. The existence of the DyB_2C_2 , DyB_2C , DyBC and Dy_2BC_2 phases was verified. The structure of the three first phases and that of the YB_2C_2 , YB_2C , and YBC phases are isotypic. A unit cell is proposed for Dy_2BC_2 . (France)

45953 HEXAVALENT W BRONZES CONTAINING Th IN INTERSTITIALS: $\text{Th}_{x/4}(\text{B}_x\text{W}_{1-x})\text{O}_3$ PHASES (B = Nb, Ta). Feyenrol, J. Y.; Sabatier, Rene; Baud, Gilbert (Univ., Clermont-Ferrand, France). Compt. Rend., Ser. C; 274: No. 11, 1059-62 (13 Mar 1972). (In French).

Two new bronze type phases were observed in both the $\text{Th}_{x/4}(\text{Nb}_x\text{W}_{1-x})\text{O}_3$ and the $\text{Th}_{x/4}(\text{Ta}_x\text{W}_{1-x})\text{O}_3$ systems. A tetragonal phase with a WO_3 -like structure occurs at high temperature along with a cubic phase of the perovskite-type. Their domains of homogeneity were investigated and the cubic structure of the perovskite phase, having the space group $\text{Pm}3\text{m}$, was crystallographically verified. (France)

45954 PHASES OF SCHEELITE STRUCTURE IN THE SYSTEMS OF PLUTONIUM MOLYBDATE-LITHIUM OR SODIUM MOLYBDATE. Tabuteau, A.; Pages, M.; Freundlich, W. (Institut du Radium, Paris). Mater. Res. Bull.; 7: No. 7, 691-7 (Jul 1972). (In French).

A molybdate of tetravalent plutonium has been identified and the systems $\text{Pu}(\text{MoO}_4)_2$ - Na_2MoO_4 and $\text{Pu}(\text{MoO}_4)_2$ - Li_2MoO_4 have been studied by means of liquid-solid equilibrium diagrams and x-ray diffraction. Three phases have been found: $\text{Na}_2\text{Pu}(\text{MoO}_4)_3$ with a scheelite structure, and $\text{Na}_4\text{Pu}(\text{MoO}_4)_4$ and $\text{Li}_4\text{Pu}(\text{MoO}_4)_4$ with a scheelite superstructure. These compounds were isostructural with the corresponding neptunium compounds. (auth)

45955 SPECIAL TECHNIQUES FOR THE METALLOGRAPHIC EXAMINATION OF HARD METALS. Arndt, Renate; Hillnhagen, Emil (Krupp Forschungsinstitut, Essen). Prakt. Metallog.; 9: No. 6, 309-17 (Jun 1972). (In German and English).

With the help of a continuous band interference filter, phase pictures can be produced of hard metal specimens coated with ZnSe having particular advantages for quality control. By vapor-phase deposition of C and Au and an appropriate astigmatism correction applied individually to the relevant points on the specimen, satisfactory scanning electron micrographs of hard metals can be produced. An electrolytic polishing technique for hard metals (polished specimens) has been developed. Using an acidic polishing solution in the double jet technique, thinned specimens were produced of WC-Co and WC-(Ti,W) C-Co hard metals containing regions transparent to electrons in all existing phases. Experiments on the thinning of hard metals by ion bombardment produced very promising results. (auth)

45956 STUDY OF ACTIVATED SINTERING MECHANISM OF UO_2 POWDER BY HIGH TEMPERATURE X-RAY DIFFRACTOMETRY. Lee, Byoung Whie; Suh, Kyung Soo (Atomic Energy Research Inst., Seoul). J. Korean Nucl. Soc.; 4: No. 2, 132-6 (Jun 1972).

The mechanism for activated sintering of UO_2 by an addition of 0.05 wt % TiO_2 was investigated using a high temperature x-ray diffractometer. The diffraction pattern of UO_2 pellets was studied in a temperature range from room temperature to 1200°C in hydrogen atmosphere. At 1200°C , the expansion of UO_2 lattice was 1.448 and 1.354% greater when it was compared with those at room temperature for pellets with and without the 0.05 wt % TiO_2 addition, respectively. The effects of the TiO_2 addition were to increase the lattice constant of UO_2 by 0.094% at 1200°C . The lattice constant at 1200°C without the TiO_2 addition was equal to that at 1080°C with the 0.05 wt % TiO_2 addition. This temperature difference could be well compared with the suppression of sintering temperature by TiO_2 that had been observed previously. It is believed that the increase in lattice expansion due to the TiO_2 addition would give rise to the activated sintering of UO_2 by the lattice-expansion-induced-enhancement of self diffusion. (auth)

45957 NUCLEAR FUEL COMPRISING URANIUM DIOXIDE IN A POROUS CERAMIC OXIDE MATRIX. Burke, Thomas J.; Belle, Jack; Clayton, John C. (to United States Atomic Energy Commission). U. S. Patent 3,657,137. 18 Apr 1972. Filed 4 Jun 1964.

A new ceramic-type nuclear fuel consisting of particles of uranium dioxide embedded in a matrix of zirconium oxide, in which the uranium dioxide particles are partially separated from the matrix by an annular void is prepared by mixing uranium trioxide with calcium-oxide-containing zirconium dioxide powder and a small quantity of a binder, pressing the mixture, and firing the mixture in an atmosphere of hydrogen. (Official Gazette)

45958 METAL OXIDE FILM STRUCTURE. Aladjem, Abraham (to Israel Atomic Energy Commission). U. S. Patent 3,675,021. 4 Jul 1972. Priority date 26 Sep 1968, Israel.

The invention provides metal oxide film structures for use as radiation windows, membranes of electrolytic cells, and other purposes. The structures consist of an annular, or other closed frame of tantalum, niobium, tungsten, or zirconium integral with a thin film of an oxide of the same metal, which spans the area enclosed by the frame. The structure is manufactured by subjecting a blank of the metal concerned, centrally masked on the larger part of the surface, to an anodizing treatment producing the oxide on all unmasked parts, removing the mask and subjecting the oxide-covered blank, as anode, to an electrolytic process by which the unoxidized metal except the frame, is dissolved. (Official Gazette)

45959 COATED-PARTICLE FUEL AND FERTILE ELEMENT FOR NUCLEAR REACTORS AND METHOD OF OPERATING SAME. Boettcher, Alfred. U. S. Patent 3,669,832. 13 Jun 1972. Priority date 17 Jan 1967, Germany.

A fuel element is described for nuclear reactors which comprises coated fuel and breeder particles in which coated particles containing the fuel have diameters of at most half the diameters of coated particles containing the fertile material. The coated particles are embedded in a graphite shell or received in a vacuum-tight metal shell. The different sizes of the coated particles containing the fuel and fertile material is the basis for mechanical discrimination between them. (Official Gazette)

45960 NITRIDE FUELS: PROPERTIES AND POTENTIALS. Bauer, Arthur A. (Battelle Memorial Inst., Columbus, Ohio). Reactor Technol.; 15: No. 2, 87-104 (1972).

Interest in nitride fuels centers principally on the use of (U,Pu)N in fast breeder reactors. A review of nitride fuel properties is presented including properties and irradiation behavior of UN, whose major potential application is as a space reactor fuel. Information on UN as presented here serves principally as background to, and to supplement, data on mixed-nitride, (U,Pu)N, fuels. Although the mixed nitrides have received less concentrated study, a sufficient technology base appears to exist that will permit an evaluation of mixed nitrides as liquid-metal-cooled fast breeder reactor fuels. The properties of (U,Pu)N related to its irradiation performance, safety characteristics, and fuel-cycle costs are reviewed. Fuel-pin design is discussed, and fuel-performance potential and limitations are outlined. Emphasis is placed on potential advantages resulting from the use of (U,Pu)N to fuel an advanced liquid-metal-cooled fast breeder reactor. (auth)

45961 THERMODYNAMICS OF RARE-EARTH-CARBON SYSTEMS. IV. THE LUTETIUM-CARBON SYSTEM. Guido, M.; Balducci, G.; De Maria, G. (Univ., Rome). J. Chem. Phys.; 57: No. 4, 1475-9 (15 Aug 1972).

The vaporization of the lutetium dicarbide-carbon system was investigated by the Knudsen effusion-mass spectrometric technique over the temperature range 2100 to 2600°K. Lu(g) partial pressures, dissociation energies for $\text{LuC}_2(\text{g})$ and $\text{LuC}_4(\text{g})$ molecules, and the standard heat of formation for $\text{LuC}_2(\text{s})$ were obtained. (auth)

45962 FERROELECTRICS IN THE $\text{K}_2\text{O-SrO-Nb}_2\text{O}_5$ SYSTEM. Ainger, F. W. (Allen Clark Research Centre, Towcester, Eng.); Beswick, J. A.; Porter, S. G.; Clarke, R. *Ferroelectrics*; 3: 321-5(1972).

Research into the tungsten bronze phase field of the $\text{K}_2\text{O-SrO-Nb}_2\text{O}_5$ system (KSN) has shown that ferroelectricity occurs over a wide range of compositions close to the $\text{KNbO}_3\text{-SrNb}_2\text{O}_6$ binary line. The approximate location of the congruently melting composition was found by chemical and x-ray analyses of crystals produced from vertical float zoning and Czochralski pulling experiments and was centered on $\text{K}_2\text{Sr}_3\text{Nb}_{20}\text{O}_{80}$. This composition gave crystals of much improved optical quality—free from striae but containing vertical structure. $\text{K}_2\text{Sr}_3\text{Nb}_{20}\text{O}_{80}$ crystals have distinctive dielectric properties, the dielectric constant of poled crystals being half that of the as-grown ones. The true lattice spacing in the "c" direction was double that of the basic tetragonal unit cell and may be explained in terms of tilted NbO_6 octahedra. The low half-wave voltage and dielectric properties of $\text{K}_2\text{Sr}_3\text{Nb}_{20}\text{O}_{80}$ make it potentially useful for electro-optic and pyroelectric applications. However, the reproducibility of striae free crystals pulled from the melt has not been proven. (auth)

45963 Al-Fe-Mo OXIDE CATALYSTS. I. THE OXIDE SYSTEM $\text{Al}_2\text{O}_3\text{-Fe}_2\text{O}_3$ AND $\text{Fe}_2\text{O}_3\text{-MoO}_3$. Khakimov, U. B.; Talipov, G. Sh.; Samigov, K. A.; Rakhmatullaev, N.; Abidova, M. F. *Zh. Obshch. Khim.*; 42: No. 2, 259-61(Feb 1972). (In Russian).

Infrared spectroscopy, EPR, and x rays were used in an examination of the systems $\text{Fe}_2\text{O}_3\text{-MoO}_3$ and $\text{Al}_2\text{O}_3\text{-MoO}_3$ after heating the binary systems to 600°C for 6 hr. It was established that the compounds $\text{Al}_2(\text{MoO}_4)_3$ and $\text{Fe}_2(\text{MoO}_4)_3$ had been formed. In an x-ray examination of the system $\text{Al}_2\text{O}_3\text{-Fe}_2\text{O}_3$ traces of the spinel FeAl_2O_4 were found. This evidence suggests that Fe-Al-Mo catalysts represent complex, multiphase systems. (K.S.W.)

45964 BOROCARBIDES OF THULIUM, LUTETIUM, TERBIUM, YTTERBIUM, AND SOME OF THEIR PROPERTIES. Bezruk, E. T.; Markovskii, L. Ya. *J. Appl. Chem. USSR (Engl. Transl.)*; 45: No. 1, 1-3(Jan 1972).

Translated from *Zh. Prikl. Khim.*; 45: No. 1, 3-6(Jan 1972).

The existence of thulium and lutetium borocarbides having the composition LnB_2C_2 , with a tetragonal structure, was demonstrated. The only ternary compounds formed by the metals Tm, Lu, Tb, and Yb with carbon and boron at 1500 to 2100° are borocarbides of the LnB_2C_2 type. No other borocarbides were found for these rare earth elements. (auth)

45965 STRUCTURE OF BINARY SILICIDES. Matyushenko, N. N. *Rev. High-Temp. Mater.*; 1: No. 2, 203-53(1972).

The table of structure data for binary silicides MeSi_x was published in book form (*Kristallicheskie Struktury Dvoynykh Soedinenii*) by the Metallurgiya Publishing House, Moscow, in 1969. The figure showing the relative change in volume as a function of the atomic number of the second component is from the same source. The table comprises data on the composition, the second-component content, the structure type and system, the number of formula weights per unit cell, the lattice parameters, the theoretical and experimental densities, the formula volume of the binary compound and its difference from the formula volume of the second component Me. The temperature at which the measurement was made (or the temperature range of stability) is given, where available. (auth)

Metals and Alloys

Refer also to abstract: 45311.

Corrosion and Erosion

Refer also to abstracts: 45461, 45487, and 45935.

45966 (BMI-1927) STRESS-CORROSION CRACKING OF SENSITIZED STAINLESS STEEL IN OXYGENATED HIGH-TEMPERATURE WATER. White, Earl L.; Berry, Warren E.; Boyd, Walter K. (Battelle Columbus Labs., Ohio). Jun 1972. Contract W-7405-eng-92. 66p. Dep. NTIS.

The intergranular stress-corrosion cracking behavior of sensitized austenitic stainless steels has been studied in water at 288°C containing 0.2, 1, 10, and 100 ppm dissolved oxygen and to a lesser extent in 288°C water first adjusted to pH 10 with NH_3 and then to pH 5 with CrO_3 . Sensitized Type 304 stainless

steel exposed to an accelerated test containing 100 ppm dissolved oxygen cracked within a few days at stress levels of 2 Sm and 3 Sm and after 83 days at 5 Sm (5 Sm = 0.9 yield strength). Mill-annealed material did not fail at 3 Sm loadings after 84 days' exposure to this same environment. On the other hand, only highly stressed (3 Sm) sensitized Type 304 stainless steel failed (217 days) in long-term exposures of up to 450 days in simulated BWR water containing 0.2 ppm oxygen. Sensitizing treatments of 7 or 24 h at 621°C or a furnace cool from 1038°C produced a high degree of susceptibility to cracking in Type 304 stainless steel in oxygenated water. Agglomerating or locally homogenizing the Type 304 steel at 816°C prior, or subsequent, to sensitizing at 621°C improved its resistance to cracking in oxygenated water, although abrading the surface of these specimens to remove a heavy tarnish film may have influenced the results. Abrading or liquid honing the surface of the Type 304 steel improved its resistance to cracking in oxygenated water, while pickling in HF-HNO_3 made it more susceptible to cracking under these conditions. Prestraining the Type 304 stainless steel did not increase its susceptibility to cracking in oxygenated water. The addition of 1 or 10 ppm fluoride to water containing 10 ppm oxygen had no effect on the susceptibility of sensitized stainless steel to stress-corrosion cracking. Sensitized, wrought, Types 304, 309, and 316 stainless steel were about equally susceptible to stress-corrosion cracking in oxygenated water. On the other hand, "sensitized" Type 304L, Type 308L weld deposit, Type 347, Incoloy 800, and Inconel 600 were resistant to the same conditions that produced rapid stress-corrosion failures of Type 304 in oxygenated water. Inconel 625 bellows failed by stress-corrosion cracking at the crevice formed by the inside bend of the bellows. Aerated or deaerated simulated off-chemistry PWR water, where the initial pH 10 (NH_3) was adjusted to pH 5 with CrO_3 , did not produce stress-corrosion cracking of sensitized Type 304 stainless steel at the same stress levels and exposure times that produced rapid cracking in oxygenated water (100 ppm oxygen). (auth)

45967 PASSIVITY OF TYPE 304 STAINLESS STEEL: EFFECT OF PLASTIC DEFORMATION. Elayaperumal, K.; De, P. K.; Balachandra, J. (Bhabha Atomic Research Centre, Bombay). *Corrosion*; 28: No. 7, 269-73(Jul 1972).

The passivity of Type 304 austenitic stainless steel, as affected by plastic deformation, was studied by obtaining the potentiostatic anodic polarization curves of annealed and cold worked materials in hydrogen saturated 1N H_2SO_4 at room temperature. While the effect of 26% cold work is not significant, 50%, and 68% cold work make the attainment of passivity more difficult by shifting the open circuit potential in the active direction and increasing the critical current density. Studies on passive current versus time indicate that cold work does not affect the initial passive current, but the extent of decrease of passive current with time is less for the cold worked material than the annealed material. These results seem to be related to the strain-induced bcc martensite which forms when the metastable austenitic stainless steel is cold worked. This martensitic phase, the presence of which has been checked by x-ray and metallography, has a high critical current density for passivity. Also, the active dissolution of a ferritic stainless steel, which does not transform to martensite, is not affected by cold work. (auth)

45968 EROSION OF A TUNGSTEN CATHODE IN NITROGEN. Zhukov, M. F.; An'shakov, A. S.; Dandaron, G.-N. B.; Sazonov, M. I. (Inst. of Heat Physics, Novosibirsk, USSR). *Izv. Sib. Otd. Akad. Nauk SSSR*, No. 8, Ser. Tekh. Nauk; No. 2, 7-12(Jun 1971). (In Russian).

The effectiveness of a protective nitrogen blanket on the destructive action of large plasmatron currents on a thoriated tungsten cathode was investigated. Tungsten rods 4 to 9 mm in diameter functioned as the cathode for currents ranging from 200 to 1,000 A. In experiments lasting for 0.5 to 2.5 h, specific erosion of a 4mm diameter tungsten cathode in N_2 lay in the interval 3×10^{-9} to 1×10^{-8} g $\text{A}^{-1} \text{sec}^{-1}$. (K.S.W.)

45969 KINETICS OF SOLID VANADIUM-NITROGEN REACTION AT HIGH TEMPERATURES. Muije, Pieter. Pullman, Wash.; Washington State Univ. (1972). 72p. University Microfilms Order No. 72-18,479.

Thesis.

Thermogravimetric, metallographic, and x-ray diffraction studies were made of the kinetics and mechanisms of the reaction between high-purity vanadium sheet and purified nitrogen gas. Nitriding was carried out isothermally in the temperature range of 900 to 1450°C at pressures between 700 and 7×10^{-2} torr for 160 hours maximum. The kinetic data obeyed either the parabolic, cubic, or linear empirical rate equation during limited time intervals. The general expression for the integral form of these rate equations is: $(\Delta W/A)^m = b T e^{-Q/RT} t + C(T)$ where $\Delta W/A$ is the weight gain per unit surface area, b a constant, T the absolute

temperature, Q the activation energy, R the universal gas constant, t the elapsed time, and C the integration constant. The exponent m is 1, 2, or 3 for the linear, parabolic, or cubic rate equation. During nitriding below 1150°C at 700 torr, the weight gain showed a parabolic time dependence except for the first few minutes. At 1150°C and above, the reaction was first parabolic, but was then followed by a cubic stage. Lowering the pressure from 700 to 70 torr did not affect the reaction kinetics. However, lowering the pressure further to 7×10^{-1} torr caused the cubic reaction to disappear and reduced the parabolic reaction rates. During nitriding at the lowest pressure of 7×10^{-2} torr, the linear in addition to the parabolic time dependence of weight gain was observed. The weight gains during the parabolic reactions were primarily reflected by the diffusion of nitrogen into the vanadium to form a solid solution, with a small amount of nitrogen combining with vanadium to form a thin compact V_3N layer. The activation energy for the parabolic reaction was independent of temperature and pressure and was 38,700 cal/mole. The cubic reaction was related to the formation of a compact layer of VN at the surface. The VN formed on the V_3N which in turn formed on the saturated solid solution of nitrogen in vanadium. The activation energy for the cubic reaction was also independent of temperature and pressure and was 49,800 cal/mole. The linear reaction, which was sometimes observed initially at 7×10^{-2} torr, seemed to be related to a rate-controlling surface mechanism. (Diss. Abstr. Int., B)

45970 EFFECT OF HEAT TREATMENTS ON THE CORROSION OF 18/10 STAINLESS STEELS WITH LOW C CONTENTS IN NITRIC MEDIUM IN THE PRESENCE OF FERRIC IRON. Blanchet, J.; Coriou, H.; Grall, L.; Pelras, M.; Nakajima, N. (CEN, Fontenay-aux-Roses, France). Mem. Sci. Rev. Met.; 69: No. 3, 193-8 (Mar 1972).

In the presence of boiling 2N nitric acid containing 40 g l⁻¹ of iron nitrate, 18/10 stainless steels with low C contents (0.012 and 0.023%) exhibited intergranular corrosion after annealing; the corrosion increased with the carbon contents. This phenomenon, which is due to the presence of ferric ions, was not encountered when the steels were adequately superquenched; a generalized dissolution of the metal was observed in this case, which increased when ferric ions were added to the medium. Such corrosion must be taken into account in welded structures. (France)

45971 VANADIUM OXIDATION IN THE PRESENCE OF MOISTURE. Varhegyi, Gyozo; Fekete, Istvanne; Vitanyi, Palne; Imre, Aladarne. Femip. Kut. Intez. Kozlem.; 10: 203-13 (1971). (In Hungarian).

The oxidation process on vanadium sponge produced by magnesium reduction during storage was investigated. On the basis of model experiments, gas analysis, and x-ray diffraction it has been stated, that the oxidation of vanadium sponge is caused first of all not by the oxygen content, but by the water vapor content of the air. Rules to prevent oxidation are listed. (auth)

Preparation and Fabrication

Refer also to abstract: 45406.

45972 (COO-3311-1) FORMING OF COMPOSITE FOR SUPERCONDUCTING ROD. Technical Progress Report. Hahn, W. C. Jr.; Avitzur, Betzalel (Lehigh Univ., Bethlehem, Pa. Dept. of Metallurgy and Materials Science). 25 Feb 1972. Contract AT(11-1)-3311. 7p. Dep. NTIS.

Research directed to production of Nb-44 wt% Ti wire clad with an Al sleeve is described. Effort during the report period was devoted to determining the effect of core-to-sleeve ratio on production of sound specimens. Processes were developed for production of an Al single-cored rod in which the hard wire core is inserted. (J.R.D.)

45973 (ORNL-RMIC-12) COMPILATION OF CRYSTAL GROWERS AND CRYSTAL GROWTH PROJECTS. Connolly, T. F. (Oak Ridge National Lab., Tenn.). Jun 1972. Contract W-7405-eng-26. 142p. Dep. NTIS.

A compilation of organizations distributed throughout the world engaged in activities associated with crystal growth is presented. Materials and applications indexes are included. Four hundred forty-one entries are included. (J.R.D.)

45974 PLUTONIUM FUEL PRODUCTION PLANT AND BREEDER PROJECTS. Vandem Bemden, E.; Lafontaine, I. (Societe Belge pour l'Industrie Nucleaire, Brussels). Energ. Nucl. (Paris); 14: No. 1, 29-32 (1972). (In French).

In conjunction with the CEN-MOL laboratories, Belgonuclear has studied the problems connected with the use of Pu as a nuclear fuel. This company has also developed a pilot production plant. In addition a Pu fuel production plant has been built at Dessel and will

be put into service in the near future. It is envisaged that this plant will provide the Pu enriched rods for the SNR core. (France)

45975 ELECTRON TRANSFER DURING ZONE MELTING OF YTTRIUM. Ivantsov, I. I.; Mashkarova, V. T.; Lavlov, V. S.; Saenko, E. M. Izv. Akad. Nauk SSSR, Metal.; No. 1, 114-18 (1972). (In Russian).

The distribution of impurities during the refining of Y by zone melting was studied. All impurities analyzed, both metallic and gaseous, migrated to the anode portion. Zone melting of Y in an electric field removed most impurities, the degree of purification increased with the number of passages and the time of electrotransfer. It was concluded that zone melting can result in coarse single Y crystals of technical purity. (K.S.W.)

45976 MECHANISM OF SINTERING IN COBALT-RARE-EARTH PERMANENT-MAGNET ALLOYS. Benz, M. G.; Martin, D. L. (General Electric Research and Development Center, Schenectady, N. Y.). J. Appl. Phys.; 43: No. 7, 3165-70 (Jul 1972).

Observations of composition, temperature, and time effects during the sintering of cobalt-samarium permanent-magnet alloys are presented. A correlation between the conditions which lead to the onset of rapid shrinkage during sintering and the conditions which lead to a high H_{ci} is noted. A model for the mechanism of shrinkage during sintering is postulated where the slow step is the diffusion of samarium atoms in the grain-boundary regions via a samarium-atom-cobalt-vacancy cluster exchange mechanism. Furthermore, it is suggested that the "high" H_{ci} structures achieved in these alloys would appear to be those with a relatively high concentration of cobalt vacancies in the grain-boundary regions. (auth)

45977 STUDY OF ELECTROLYTIC DEPOSITION OF TUNGSTEN-COBALT ALLOYS. Yur'ev, B. P.; Kiseleva, L. S.; Zabelin, I. V. (Leningrad Polytechnic Inst.). J. Appl. Chem. USSR (Engl. Transl.); 45: No. 1, 79-82 (Jan 1972).

Translated from Zh. Prikl. Khim.; 45: No. 1, 85-9 (Jan 1972).

A study of cathodic polarization curves for joint reduction of tungsten, cobalt, and hydrogen in mixed $Na_2WO_4 + CoSO_4$ solutions with additions of ammonium citrate, in relation to the solution composition and temperature, showed that formation of W-Co electrolytic alloy on the cathode is accompanied by considerable depolarization of both components simultaneously. Current efficiency for the alloy increased with decrease of current density and with rise of temperature. The tungsten content in the alloy increased appreciably with rise of temperature; current density had considerably less influence on the alloy composition. The value of α for tungsten was close to the theoretical if it was assumed that reduction of WO_4^{2-} ions to the metal proceeds by stages, the one-electron process being the determining (slowest) step. It was found by x-ray structural and microscopic investigations of the electrolytic alloys that alloys containing 50% tungsten (by weight) contain in their structure an ϵ' phase (solid solution of tungsten in cobalt) and the chemical compound Co_3W , and are consistent in structure with the phase diagram of the W-Co system. It is suggested that alloy formation at the cathode is one of the determining factors in depolarization and determines the possibility of reduction of tungsten jointly with other metals at the cathode. (auth)

45978 ELECTROCHEMICAL REDUCTION OF URANIUM(VI) TO URANIUM(IV) IN SULFURIC ACID SOLUTIONS. Stabrovskii, A. I.; Slepchenko, I. G. J. Appl. Chem. USSR (Engl. Transl.); 45: No. 1, 101-6 (Jan 1972).

Translated from Zh. Prikl. Khim.; 45: No. 1, 110-15 (Jan 1972).

It was found with the aid of polarization curves that, dependent on the cathode potential, formation of U(IV) in solution may proceed by reduction of U(VI) to U(IV) at the cathode as the result of disproportionation of U(V), or by interaction of U(VI) with U(III) in the bulk solution. (auth)

45979 Zr Hf NEWSLETTER, JUNE 1972. Akron, N. Y.; AMAX Specialty Metals Division (1972). 18p.

Developmental and other information is summarized on Hf and Zr metal, Zr-base alloys, use of Zr as an additive to other alloys, and production of Zr oxides and silicates. Information is also included on Zr borides, carbides, nitrides, germanides, hydrides, sulfides, selenides, silicides, and tellurides. 316 references are included. (J.R.D.)

45980 SURFACE PREPARATION OF URANIUM PARTS.

de Vaulchier du Deschaux, Louis; de Fouchier, Alain; de Vaucelles, Emmanuel; Facquet, Louis (to Commissariat a l'Energie Atomique). U. S. Patent 3,674,655. 4 Jul 1972. Priority date 24 Jan 1968. France.

Prior to electroplating parts of uranium or uranium base alloys, the parts are subjected to a surface preparation, the preliminary steps involve chemical degreasing, electrolytic degreasing, and if necessary chemical etching and an anodic electrolytic treatment in an aqueous solution of an organic salt or a sulphate of lithium and/or magnesium. (Official Gazette)

45981 VACUUM INDUCTION MELTING. Schlatter, R. J. *Metals*; 24: No. 5, 17-25(May 1972).

The increase of vacuum-induction melting (VIM) capacity is closely related to the continuing growth in the use of superalloys for thermal engines, which in turn stimulated the development of a large variety of wrought and cast alloys. The presence of substantial amounts of reactive elements and the requirements of complete chemistry control (alloy composition, beneficial trace elements, and harmful impurities) to very narrow tolerances as well as exceptional cleanliness justify the application of this melting technique. VIM is an expensive operation, even in large furnaces of advanced design with many operational improvements. But there is no other proven primary industrial melting process providing the exact metallurgical conditions necessary for the reliable and consistent production of the great variety of these complex alloys. The physicochemical aspects of the VIM process are well understood today, but for various reasons the practice does not generally appreciate the importance of this valuable knowledge, and its full utilization is lacking. Much progress has been made in the area of refractory technology, but more work is needed, particularly in developing new types of refractory materials or ceramic systems to extend the metallurgical potential of the process and improve the operational efficiency. The shortcomings of the VIM process regarding casting and solidification control are difficult to overcome, hence the three-step manufacturing process with final consumable electrode vacuum arc remelting—or electroflux remelting—will remain technically, metallurgically, and economically the most flexible approach, providing exceptional overall quality and integrity in these sophisticated superalloys. (auth)

45982 RECRYSTALLIZATION OF METALS OF HIGH PURITY. Chaudron, Georges (Univ., Paris). J. Less-Common Metals; 28: 175-85(1972). (In French).

It is known that studies on the recrystallization of metals have generally had the practical purpose of preparing single crystals. In the very ingenious method of Burgers and Tiedema, it is possible to make single crystals of pre-determined orientation. Professor Burgers has studied the progress of recrystallization by various methods: x-ray diffraction and optical microscopy. The influence of purity on the conditions under which the polygonized state appears is pointed out. The existence of a critical polygonization temperature was first found in the case of aluminum. The polygonization of iron has been studied as a function of purity. In the case of zone-refined iron, the subgrains formed were highly perfect. Studies on the polygonization of copper, which were made possible by the availability of samples with a range of increasing purities, are described. By thermal cycling, the polygonized structure underwent modifications above a certain critical temperature. This temperature was strongly purity-dependent. In conclusion, the polygonized state appeared to occupy a domain of metastability which was distinct from the recrystallized state. (auth)

45983 INFLUENCE OF TEMPERATURE ON ELECTRO-DEPOSITION OF TUNGSTEN-COBALT ALLOY. Zabelin, I. V.; Yur'ev, B. P. J. Appl. Chem. USSR (Engl. Transl.); 45: No. 1, 197-9(Jan 1972).

Translated from Zh. Prikl. Khim.; 45: No. 1, 205-7(Jan 1972).

Studies were made on the effects of temperature, carried out for determination of the activation energies and kinetic characteristics of the partial processes of reduction of WO_4^{2-} and Co^{2+} ions at the cathode, on the electrolytic deposition of W-Co alloys. Relatively low activation energies were found in both cases, which indicates that the course of the processes is controlled by diffusion. It was concluded that reduction of Co and W under the specific conditions under consideration proceeds in the region of mixed control, i.e., is controlled both by diffusion and by electrochemical factors. (P.C.H.)

45984 GRAIN REFINING OF MAGNESIUM ALLOYS. Emod, Gyula. *Fennip. Kut. Intez. Kozlem* 10: 163-9(1971). (In Hungarian).

Grain refining of magnesium and its alloys was achieved by treating them in a melted state. Treatments were as follows: overheating, chlorination, addition of carbon compounds, etc. Grain refining can be attained by adding certain alloying elements (Zr, Al, Zn etc.). The mechanism of grain refining methods and formation of crystal nuclei on a theoretical basis were also studied. (auth)

Properties, Structure, and Phase Studies

Refer also to abstracts: 45466, 45471, 47248, 47250, and 47263.

45985 (BARC-577) CHEMICAL DIFFUSION IN THE BETA PHASE OF ZIRCONIUM-TITANIUM SYSTEM. Raghunathan, V. S.; Tiwari, G. P.; Sharma, B. D. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 25p. Dep. NTIS (U. S. Sales Only).

Interdiffusion studies were carried out on the bcc phase of zirconium-titanium alloy system using diffusion couples of zirconium and titanium-zirconium alloys. A geoscan electron microprobe was employed to establish the concentration-penetration profiles in the diffusion zones, and the interdiffusion coefficients ($\bar{D}_{12}$) were evaluated by Boltzmann-Matano analysis. The concentration and temperature dependence of $\bar{D}_{12}$ were examined. The results showed that the interdiffusion data for the Zr-Ti system in β -phase can be expressed by $\bar{D}_{12} = A \exp [-(11.2\Delta T_s^2/RT)]$ where A is a function of the limiting diffusivity values of two components at infinite dilution, and ΔT_s^2 is the difference between the solidus temperature and the temperature that would be computed if solidus line varied linearly with composition. This equation is a modified version of Vignes-Birchall equation applicable to vacancy-controlled interdiffusion in fcc solid solutions. The diffusivity values of Zr in β -Ti and Ti in β -Zr have also been evaluated by extrapolation. The results indicated that interdiffusion in the zirconium-titanium system in the low-temperature range of bcc phase occurs by a vacancy mechanism. (auth)

45986 (BARC-585) ABSORPTION OF CATHODIC HYDROGEN BY IRON AND STEEL. Radhakrishnan, T. P. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 40p. Dep. NTIS (U. S. Sales Only).

Delayed failure and embrittlement due to hydrogen absorption pose intricate problems and limit the application of many high-strength steels in industry and technology. Hydrogen pickup occurs concomitantly during various chemical and electrochemical metal finishing processes and also during galvanic corrosion of the plated steel components. The behavior of cathodic hydrogen in iron and steel and, in particular, the various factors which influence absorption were examined. The mechanism of entry and release of hydrogen was considered on the basis of electrochemical kinetics with due emphasis on the location and state of hydrogen in steel. Methods for studying hydrogen absorption and techniques for evaluating the diffusion coefficient of hydrogen in iron and steel are summarized. A comparison of the data for the diffusivity of hydrogen, obtained by various methods, indicates wide discrepancy and suggests trapping of hydrogen in voids. The effects of pH, poisons, inhibitors, and oxidizing agents on the absorption of hydrogen by steel were examined. The influence of metallurgical factors, such as composition, cold-work, annealing, microstructure of the steel, and applied tensile-stress on hydrogen absorption were considered in some detail. Delayed failure of hydrogenated steel and the theories of embrittlement were also studied. (119 references). (auth)

45987 (CONF-710601-, pp 605-12) VOID FORMATION IN QUENCHED STAINLESS STEEL. Rowcliffe, A. F. (Westinghouse Electric Corp., Madison, Pa. Advanced Reactors Div.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Disk shaped specimens of five austenitic stainless steels were solution treated at temperatures of 1300 and 1100°C and directly quenched into salt or lead baths held at aging temperatures between 400 and 700°C. In one of the steels, which contained an unusually high phosphorus concentration (18 Cr 10 Ni 0.31 P), vacancy loops and voids were produced; the nature of the defects formed depended upon the solution treatment temperature and on the aging temperature. It was found that there was a critical temperature below which it was necessary to quench in order to nucleate voids. Following solution treatment at 1300°C the critical temperature was between 650 and 700°C; quenching to lower temperatures increased the concentration and decreased the average size of the voids. After solution treatment at 1100°C it was necessary to quench to below 500°C in order to nucleate voids. The observation of voids in quenched stainless steels has not been reported previously. The phenomenon is believed to be the result of a strong phosphorus atom-vacancy interaction which increases the equilibrium vacancy concentration at the solution treatment temperature and greatly modifies vacancy clustering behavior at the aging temperature. (auth)

45988 (CONF-710601-, pp 630-46) SOLUTE SEGREGATION AND STRESSES GENERATED AROUND GROWING VOIDS IN METALS. Anthony, T. R. (General Electric Co., Schenectady, N. Y. Research and Development Center). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

During cooling from a high-temperature anneal, the vacancy supersaturation in a zone neighboring the surface of an aluminum sample reaches the critical point where vacancies begin to precipitate heterogeneously at the metal-oxide interface. The resulting vacancy condensation cavities provide large natural vacancy sinks ($1000\mu^3$) around which any impurity segregation generated by vacancy currents forming these cavities can be directly determined. Vacancy currents were found to cause either enrichment, depletion, or no segregation around the cavities depending on the particular solute addition. From the vacancy and vacancy-generated impurity currents directly measured in these experiments, the ratios of vacancy jump frequencies away from and around impurity atoms in aluminum are determined. Conversely, in other alloy systems in which vacancy jump frequencies are available from isotope diffusion experiments, the impurity segregation generated by vacancy currents around growing voids can be calculated. (auth)

45989 (CONF-710601-, pp 647-62) ANNEALING OF LOOPS AND TETRAHEDRA. Washburn, Jack (California Univ., Berkeley, Lawrence Berkeley Lab.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

A discussion of the mechanisms for annealing of dislocation loops and tetrahedra is presented. The growth of vacancy defects in metals is reviewed with emphasis on vacancy supersaturation. The role of dislocation climb in the shrinkage of loops and tetrahedra is discussed together with the effect of stacking-fault energy. Dislocation-loop migration by diffusion along the dislocation core is shown to be a possible mechanism for coarsening of small dislocation loops at temperatures just below that required for ordinary self-diffusion. (auth)

45990 (CONF-720806-2) ANALYTICAL APPLICATIONS OF THE J INTEGRAL. Merkle, J. G. (Oak Ridge National Lab., Tenn.). 1972. 29p. Dep. NTIS.

From sixth national symposium on fracture mechanics; Philadelphia, PA. (28 Aug 1972).

It has recently been shown by experiment that a quantity known as the J Integral may be a useful fracture criterion in the inelastic range. The mathematical definition of the J Integral is used to derive a relationship between the conventional elastic stress concentration factor of a sharp notch and the elastic stress intensity factor for a crack of the same shape. This relationship has previously been derived only by assuming some particular shape of cracked body, but this assumption is shown to be not necessary. The same assumptions, together with Neuber's equation for the inelastic stress and strain concentration factors for a sharp notch, are used to derive a relationship between the J Integral and the parameter K_{Ic} of the Equivalent Energy Method. A graphical procedure for estimating upper and lower bounds on the full restraint value of the fracture toughness is also discussed. (auth)

45991 (COO-916-25) STRUCTURE AND PROPERTIES OF SOLID SOLUTIONS. Technical Progress Report, September 1, 1971-June 15, 1972. Hendrickson, A. A.; Koss, D. A. (Michigan Technological Univ., Houghton, Dept. of Metallurgical Engineering). 1972. Contract AT(11-1)-916. 27p. Dep. NTIS.

Results of studies on bcc solid solution hardening in high-purity Nb-Hf, Nb-W, and Nb-Ta alloys are presented. Data from measurements of a thermal yield stress in substitutional Nb-W-Mo alloys deformed at 900°K are included. Results of fcc solid solution hardening studies are also presented which indicate that serrated yielding occurred in single crystals of Ag-base-In alloys. (J.R.D.)

45992 (LBL-659) SOME PROPERTIES OF PROTACTINIUM METAL AND ITS COMPOUNDS. Dod, Raymond Lloyd (California Univ., Berkeley, Lawrence Berkeley Lab.). May 1972. Contract W-7405-eng-48. 98p. Dep. NTIS.

Thesis.

A modification of the usual metallothemic method of metal production is described, utilizing a molten salt crucible which substantially reduces the probability of contaminating the protactinium metal with crucible material. The method may have general applicability to the preparation of small quantities of high melting point metals. A redetermination of the magnetic susceptibility of protactinium metal was made, resulting in a substantial change in value although not in temperature dependence from that previously reported. The new value is $(190 \pm 5) \times 10^{-6}$ cgs units/mole and is independent of temperature from $\sim 7^\circ\text{K}$ to 300°K . This value holds for both the normal tetragonal form of the metal and also the fcc high temperature modification. The magnetic susceptibility of PaF_4 is also reported. The observed

behavior can be interpreted to follow the Curie-Weiss law with an effective moment of 1.81 Bohr magnetons and a Weiss constant of 22.5° . The reactions of protactinium metal with several common gases at temperatures up to 700° are described. Standard x-ray powder diffraction techniques were used to analyze the results. New compounds reported include two nitrides prepared by reaction of the metal with nitrogen and with ammonia. Both exhibited a cubic fluorite type lattice. The observed lattice parameters were $a_0 = 5.436 \pm 6 \text{ \AA}$ and $a_0 = 5.395 \pm 3 \text{ \AA}$. A new hydride structure for PaH_3 was observed, isomorphic with α -uranium hydride with $a_0 = 4.152 \pm 2 \text{ \AA}$. Additionally, an unidentified phase was observed in reaction of hydrogen with Pa metal at 400°C , which was presumed to be a hydride since thermal decomposition regenerated the metal. The hydride previously reported for protactinium was not observed, nor was any evidence for the existence of the reported monoxide, PaO , seen in any reaction, including those with air, oxygen, and water. The melting point and thermal expansion coefficients for a high purity sample of the metal have shown that the previous values, derived from samples of dubious quality, were more accurate than had been thought. (auth)

45993 (ORNL-tr-2377) BRITTLE-DUCTILE TRANSITION OF BODY-CENTERED CUBIC METALS WITH SPECIAL CONSIDERATION OF TUNGSTEN AND MOLYBDENUM. PART I. EXPERIMENTAL METHODS. Pink, E.; Sedlatschek, K. Translated for Oak Ridge National Lab., Tenn., from Metall; 23: No. 12, 1249-57(1969). 22p. Dep. NTIS.

The most advantageous method to determine the brittle-ductile transition of tungsten and molybdenum sheets consists of the bending test conducted at different temperatures. The results changed when important dimensions of the equipment were varied. The sharpest indication of the transition was possible with the smallest possible gage length. The ratio of punch radius to specimen thickness i.e., the so-called T-value, was important to the transition temperature. The experiences collected led to suggestions for standardization of the method. The transition temperatures for work-hardened rolled molybdenum and tungsten were -80 to -70°C and 100 to 200°C , respectively, and for the recrystallized metals to 20 to 100°C or about 360°C , respectively. Recrystallized molybdenum and tungsten with a distinct stacked structure were still ductile at room temperature. The variation of the transition temperature of the system molybdenum-tungsten was not linear. The substitution effect produced an increase in the transition temperature in the recrystallized state and a decrease in the work-hardened metal. In work-hardened dispersions of $\text{Mo-Al}_2\text{O}_3$ the transition temperature decreased. An embrittling effect was produced only by larger amounts of dispersate ($>1 \text{ vol}\%$). (P.C.H.)

45994 (UCLA-34-P-177-10) REACTOR STEELS STUDIES. Quarterly Progress Report, October-December 1971. Ono, K.; Tetelman, A. S.; Ardell, A. J. (California Univ., Los Angeles, School of Engineering and Applied Science). Mar 1972. Contract AT(04-3)-34. 13p. Dep. NTIS.

Research progress is reported on structure and strengthening of 2.25 Cr-1 Mo steels, effects of microstructure on fracture modes of Types 304 and 316 stainless steels, and precipitation reactions in stainless steels. (J.R.D.)

45995 (UJV-2802-M) MECHANICAL CHARACTERISTICS OF URANIUM ALLOYS OF THE TYPE U-Mo(Fe, Al, Si) AND U-Fe-Al-Si(Cr). Saxl, I. (Ceskoslovenska Akademie Ved, Rez, Ustav Jaderneho Vyzkumu). 1972. 46p. (In Czech). Dep. NTIS (U. S. Sales Only).

The mechanical properties of uranium alloys containing low additions of Al, Fe, Mo, Si, and Cr were measured in the temperature interval 20 to 640°C . The mutual correspondence between phenomenological quantities (strain rate sensitivity m and logarithmic hardening at the maximum load n_P) and ductility parameters (inhomogeneous elongation ϵ_n and homogeneous elongation ϵ_P) was confirmed. Above 200°C , the relations $\epsilon_P = n_P$ and $\epsilon_n = f(m)$ were held independently of the experimental conditions (e.g., temperature, deformation rate, and composition). Below 200°C , both the relations were disturbed by the embrittling phenomena connected with the presence of the nonmetallic inclusions. The temperature dependence of the total elongation (with its characteristic intermediate temperature minimum) was the direct consequence of the temperature dependences of hardening and strain rate sensitivity parameters. The amount of alloy hardening and its temperature characteristics are discussed on the basis of recent dispersion hardening theories. (auth)

45996 (WCAP-7841) DISCUSSION ON FRACTURE MECHANICS CONCEPTS. Mager, T. R.; Buchalet, C.; Enrietto, J. F. (Westinghouse Electric Corp., Pittsburgh, Pa.). Mar 1972. 24p. Dep. NTIS.

Results of brief review of four fracture mechanics concepts applicable for developing a fracture prevention criterion for heavy-section nuclear reactor pressure vessels are presented. It is concluded that linear-elastic fracture mechanics technology (L.E.F.M.) is applicable to heavy-section components, such as nuclear reactor vessels, when plane strain conditions prevail and the plastic zone remains small. The two basic assumptions and the technical consequences of the stress concentration theory (SCT) were examined with reference to heavy sections. It was found that the SCT cannot be applied to heavy-section steel vessels. The crack opening displacement (COD) approach is compatible with both linear-elastic fracture mechanics and transition concepts. The path independent J-Integral apparently can be used to predict fracture in low- to high-strength steels for conditions ranging from linear-elastic to fully plastic. However, the concept has not been fully developed to date, thus practical application appears to be several years away. (J.R.D.)

45997 CONSTITUTION OF BINARY ALLOYS, SECOND SUPPLEMENT. Shunk, Francis A. McGraw-Hill Series in Materials Science and Engineering. New York, McGraw-Hill Book Company (1969). 759p. \$37.50.

Systems are arranged in alphabetical order of the chemical symbols of the elements. For each system, the existence of a review in Hansen, Elliott, or this Supplement is indicated by the appropriate page number, and the inclusion of a diagram is also indicated. Other references pertinent to the systems are also given. (P.C.H.)

45998 THERMAL PROPERTIES OF LIQUID CERIUM. Mardynkin, I. P.; Vertman, A. A. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 95-8(1972). (In Russian).

The properties of liquid cerium were studied in an apparatus designed for the simultaneous measurement of thermal conductivity and heat capacity at high temperatures. The results indicated that the heat conductivity and thermal capacity of liquid Ce were only weakly dependent on the temperature over the temperature range 1,100 to 1,600°K. (K.S.W.)

45999 EFFECT OF MICRO-ADDITIVES OF RARE EARTH METALS ON THE GRAIN BOUNDARY ENERGY OF NICKEL. Koval, A. D.; Ol'shanetskii, V. E. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 124-32(1972). (In Russian).

The effects of adding small amounts of Nd, Pr, Ce, and La on the grain boundary energy and on the migration of the boundary of pure nickel was studied. Such additives were disposed chiefly on the grain boundary, changing the value of its excess free energy. The concentration dependence of the grain boundary energy of 99.9999% pure Ni was a curve with a maximum at 0.5×10^{-3} atom fraction of rare earths. The velocity of migration of a grain boundary at some particular rare earth metal content was associated with both additive interaction and the formation of chemical compounds with "unaccounted-for metals" (with the refining of the grain boundary). The four metals affected the grain boundary energy and the velocity of boundary migration in the following order (greatest effect last): La, Ce, Pr, Nd. (K.S.W.)

46000 CREEP OF NIOBIUM UNDER CONDITIONS OF IN-LEAKAGE OF OXYGEN. Abramyan, E. A.; Ivanov, L. I.; Kazilin, E. E. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 177-82(1972). (In Russian).

The effects of a current of oxygen at an intensity of 5×10^{17} mole/sec on creep of niobium during twisting were investigated over the temperature range 1,200 to 1,850°C and under a stress of 6 to 200 kg/cm². A sharp increase in the creep rate occurred simultaneously with contact with the oxygen stream, this phenomenon being temperature-dependent. Several reasons for the sudden increase in creep rate are discussed. (K.S.W.)

46001 THERMODYNAMIC PROPERTIES AND PHASE COMPOSITION OF La-Zn ALLOYS. Kovalevskii, A. V.; Lebedev, V. A.; Nichkov, I. F.; Raspopin, S. P. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 183-7(1972). (In Russian).

The La-Zn equilibrium diagram was investigated with the aid of emf measurements over the range 723 to 1,123°C using a galvanic cell of the type La-Zn/LiCl-KCl + 2 wt % LaCl₃/La-Zn_{sat. soln.}. The existence of the following eight metallic compounds was established: LaZn₁₁, LaZn₉, LaZn₈, LaZn₇, LaZn₆, LaZn₄, LaZn₂, and LaZn. Thermodynamic properties of the alloys attested to the energy of interaction between La and Zn. (K.S.W.)

46002 REACTION OF Nb-Mo ALLOYS WITH BORON. Zhurkovskii, G. L.; Braun, S. M. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 188-92(1972). (In Russian).

A study of the reactions of B with Nb-Mo alloys indicated that the reaction rate was determined by the percent of a component in the alloys. Over the range 1,100 to 1,400°C the rate of increase of the boride phase in alloys with 20 to 50% of Mo was four to six

times less than pure Nb and Mo. The prospects of using this type of alloy in boron-containing media are discussed. (K.S.W.)

46003 SURFACES OF THE SOLIDUS AND THE LIQUIDUS IN THE SYSTEM Th-Zr-U. Badaeva, T. A.; Kuznetsova, R. I. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 196-200(1972). (In Russian).

The isothermal surfaces of the solidus and liquidus of the ternary system Th-Zr-U were constructed from data from thermal analysis, differential heating curves, and the microstructure of these alloys. The results were found to agree with previous work. Solidus and liquid curves in the binary systems Th-U and Th-Zr from 20 to 70 at. % U were established with greater precision than before. For the compositions studied the temperature of the solidus ranged from 1,070 to 1,485°C; that of the liquidus from 1,260 to 1,855°C. (K.S.W.)

46004 THERMODYNAMIC PROPERTIES OF SOLUTIONS IN THE SYSTEM In-Sn-Hg. Nigmatova, R. Sh.; Sudakov, V. A.; Kosin, L. F. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 201-7(1972). (In Russian).

Pressure measurements of Hg vapor from In-Sn-Hg alloys in the temperature interval 150 to 340°C were used to calculate thermodynamic functions of amalgams in this ternary system. Calculated values for the activity of Hg in this system were in good agreement with experimental values. Significant negative and small positive deviations of thermodynamic quantities from those expected for ideal solutions were realized. (K.S.W.)

46005 TERNARY SYSTEM Y-Fe-Al IN THE REGION 0 TO 33.3 PERCENT Y. Zarechnyuk, O. S.; Rykhal, R. M.; Ryabov, V. P.; Vivchar, O. I. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 208-10(1972). (In Russian).

Ninety ternary alloys of the system Y-Fe-Al up to a Y content of 33.3% were investigated with x-ray structural analysis, and an isothermal section was constructed at 500°C. Compounds and phases identified were: a compound YFe_{4.0-5.8}Al_{8.0-8.2}, a compound YFe_{1.0-1.2}Al_{1.0-0.8}, a compound YFe₂Al₁₀, a phase $\lambda_2(\text{Fe})$ (a solid solution of Al in YFe₂), a phase $\lambda_2(\text{Al})$ (a solid solution of Fe in YAl₂), a phase ϕ_1 (a solid solution of Al in Y₂Fe₁₇), and a phase ϕ_2 (a solid solution of Al in Y₂Fe₁₇, with a structure different from ϕ_1). (K.S.W.)

46006 EFFECT OF MOLYBDENUM ON AGING OF Fe-Co-W ALLOYS. Ershova, L. S. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 211-15(1972). (In Russian).

The alloying of Fe-Co-W alloys with Mo lead to an increase in the micro-deformation of the crystal lattice of the quenched α -solid solution. A comparison of the aging kinetics in alloys V16K23 (15.52% W, 23.52% Co) and V16M8K23 (16.05% W, 22.50% Co) showed that the addition of Mo significantly raised the rate of dispersed solidification. The effects of Mo on structural changes during aging are explained by a change in the composition of the quenched solid solution and an increase in the imperfection of the crystalline lattice. (K.S.W.)

46007 CALORIMETRIC STUDY OF Nb-Fe-Si-Al SYSTEM. Gorelkin, O. S.; Dubrovina, A. S. *Izv. Akad. Nauk SSSR, Metal.*; No. 1, 216-21(1972). (In Russian).

The heats of solution of Al and Si in Fe, and of Nb in Nb-Fe alloys were determined with a large capacity isothermal calorimeter. The necessity for making allowance for the heat of solution of elements in estimating thermal balances in niobium alloys was demonstrated. (K.S.W.)

46008 STUDIES OF THE ORIGIN OF THE EXOGENOUS OXIDE INCLUSIONS IN KILLED AND UNKILLED CAST STEELS THROUGH SUBSEQUENTLY ACTIVATED TRACER ELEMENTS. Benkoe, Gyulane; Simon, Sandor; Szarka, Gyula. *Neue Huette*; 17: No. 1, 40-3(Jan 1972). (In German).

Oxide inclusions in steel originating from heat-resistant material were studied. Barium-140, in the form of BaCl₂, was applied to the heat-resistant cladding of the casting vessel and various other parts of the casting equipment. By electrolysis, inclusions were then separated from the steel and their activity measured using a Geiger-Mueller counter. Results showed that inclusions originated mainly from those sections of the heat-resistant material that were mechanically stressed during the casting process. In an alternate method, stable barium was applied to the heat-resistant cladding. After isolation of the steel inclusions by electrolysis, they were activated with 14-MeV neutrons and the γ spectrum was obtained using a crystal detector and a multichannel analyzer. (H.B.G.)

46009 EFFECT OF MECHANICAL SURFACE HARDENING ON THE LOW-CYCLE FATIGUE STRENGTH OF TITANIUM ALLOYS. Kudryavtsev, I. V.; Vainshtein, V. G. *Metal Sci. Heat Treat. (USSR) (Engl. Transl.)*; 13: No. 11-12, 1038-40(1971).

Translated from *Metalloved. Term. Obrab. Metal.*; 13: No. 12, 44-46 (Dec 1971).

Surface hardening effectively increased the low-cycle fatigue strength of titanium alloy VT3-1 particularly with stress concentrators. Hardening a notch by burnishing substantially increased the fatigue limit under load to a value corresponding to the yield strength of the material and could completely eliminate the unfavorable effect of the concentrator. Hardening of the area near the stress concentrator without hardening the notch itself had comparatively little effect. (auth)

46010 FORMATION OF INTERMETALLIC PHASES IN THE SYSTEM Ir-Ta-Ni. Bernard, V. B.; Kuprina, V. V.; Burnashova, V. V. *Vestn. Mosk. Univ., Ser. II. Khim.*; 13: No. 2, 240-1 (1972). (In Russian).

The system Ir-Ta-Ni was investigated over a full range of concentrations. New intermetallic phases denoted as X, Y, A were observed. (auth)

46011 STUDY OF ZIRCONIUM EVAPORATION FROM OPEN SURFACE. Golubtsov, I. V.; Mikul'skaya, G. Yu.; Pavlova, L. M. *Vestn. Mosk. Univ., Ser. II. Khim.*; 13: No. 2, 206-10 (1972). (In Russian).

A new apparatus was built for studying the evaporation of construction materials in vacuum at high temperature. The evaporation rate of β -zirconium was measured. (auth)

46012 SINTERED MAGNESIUM COMPOSITES, CLADDING MATERIAL OF URANIUM FUEL ELEMENTS COOLED BY CARBON DIOXIDE. PART II. ANISOTROPY OF PROPERTIES OF EXTRUDED AND RECRYSTALLIZED PRODUCTS. Volf, Frantisek; Sokol, Ludvik; Cech, Bohuslav (Nuclear Research Inst., Rez, Czech.). *Jad. Energ.*; 18: No. 5, 151-7 (May 1972). (In Czech).

Discussion is presented on the anisotropy of properties of sintered magnesium products, especially of extruded and recrystallized rods and tubes. The effect of processing, conditions on the formation of preferred orientation has been investigated. The preferred orientation of the magnesium hexagonal unit cell in green compacts of magnesium powder is parallel to the axis of compacting, whereas in extruded and recrystallized products it is almost perpendicular to the extrusion axis. (auth)

46013 EXPLOITATION OF SOLID ELECTROLYTES IN RESEARCH AND TECHNOLOGY OF NUCLEAR POWER. Jakes, Dusan (Nuclear Research Inst., Rez, Czech.). *Jad. Energ.*; 18: No. 5, 158-63 (May 1972). (In Czech).

Methods of application are explained on the basis of indispensable theoretical inferences. Experimental conditions are outlined for investigating the thermodynamics of solid bodies, making solubility measurements, and determining the diffusion coefficients in various materials used in nuclear technology. Solid electrolytes such as ZrO_2 -CaO, ZrO_2 - Y_2O_3 , ZrO_2 -MgO, La_2O_3 -CaO, and CaF_2 - YF_3 have been used with oxygen at different pressures as the anode and cathode, respectively. The use of electrolytes as analytical detecting elements is discussed. (tr-auth)

46014 PHYSICO-CHEMICAL MEASUREMENTS IN METALS RESEARCH. PART 1. Rapp, R. A. (ed.). *Techniques of Metals Research*. Volume IV. New York; John Wiley and Sons, Inc. (1970). 575p.

Information on physicochemical measurements on metals is included under the chapters: physicochemical measurements in metals research; vapor pressure methods, including measurement of the gross equilibrium vaporization rate, free vaporization measurements (Langmuir technique), measurement of vapor pressures by transpiration, isopiestic, and other techniques, and mass spectrometry in the measurement of vapor pressure; calorimetry, including reaction calorimetry and bomb calorimetry; chemical equilibria, including the absorption of unary gases by condensed phases, equilibria and transport phenomena involving gas mixtures and condensed phases, and equilibria and transport phenomena between solid and liquid phases and between two liquid phases; and phase equilibria and transformations, including phase equilibria in metallic systems and techniques for investigating phase-transition kinetics in metals. Vapor pressure data for the solid and liquid elements are appended. (P.C.H.)

46015 PHYSICO-CHEMICAL MEASUREMENTS IN METALS RESEARCH. PART 2. Rapp, R. A. (ed.). *Techniques of Metals Research*. Volume IV. New York; John Wiley and Sons, Inc. (1970). 718p.

Information on physicochemical measurements on metals is included under the chapters: electrochemical measurements, including aqueous electrolyte cells, molten salt electrolyte cells, and solid electrolyte galvanic cells; viscometry and densitometry,

including viscosity of liquid metals and liquid density measurements; surface energy measurements, including measurement of liquid-vapor and solid-vapor surface energies and interfacial energies between condensed phases; diffusion measurements, including volume diffusion in solids, surface, grain boundary, and dislocation pipe diffusion, and diffusion measurements in liquid metals; and oxidation and corrosion, including oxidation of metals and corrosion of metals. (P.C.H.)

46016 X-RAY AND NEUTRON DIFFRACTION STUDY OF THE HOLMIUM-IRON SYSTEM. Simmons, Michael Fred. Rolla, Mo.; Univ. of Missouri (1971). 74p. University Microfilms Order No. 72-18,205.

Thesis.

Neutron diffraction and x-ray powder studies have been made of holmium-iron intermetallic compounds. The magnetic moments of Ho and Fe are in good agreement with measured spontaneous magnetization assuming a ferrimagnetic model with the Ho moments antiparallel to the Fe moments. The magnetic moments of the Fe atoms are found to be localized and thus independent of the rare earth content in the alloys. In contrast to $HoCo_3$, (001) type magnetic peaks in the neutron diffraction patterns are still prevalent at room temperature and gradually decrease as the Curie temperature is approached. This fact indicates that the easy direction of magnetization remains in the basal plane. The easy direction of magnetization at room temperature is found to be parallel with the cube edge in $HoFe_2$ and to lie in the basal plane in $HoFe_3$ and Ho_2Fe_{17} . The easy direction is parallel to the a-axis in Ho_2Fe_{17} . It is parallel to the b-direction, i.e., 30° from an a-axis, in the $HoFe_3$ hexagonal cell. (Diss. Abstr. Int., B)

46017 GENESIS OF CELLULAR PRECIPITATION IN COPPER RICH COPPER-INDIUM ALLOYS. Fournelle, Raymond Albert. Rolla, Mo.; Univ. of Missouri (1971). 148p. University Microfilms Order No. 72-18,190.

Thesis.

The morphology of cellular precipitation in a Cu-9.5 at. %In alloy has been investigated by light and electron microscopy and x-ray and electron diffraction. Both cellular and general precipitation were observed to occur simultaneously in quenched and aged alloys while only cellular precipitation was observed to occur in isothermally aged alloys. Because of the presence of wide, vacancy depleted precipitate free zones in the vicinity of grain boundaries, general precipitation was observed to have no effect on the early development of cellular precipitation. Hence, the early development of cellular precipitation was found to be identical for both types of heat treatment. By light and electron microscopy of the early stages of precipitation a mechanism for the development of cells from an initially unoccupied grain boundary was developed. According to this mechanism the grain boundary allotriomorphs begin to form, the boundary interacts with them by bowing out between them and trailing precipitate behind to eventually develop a steady state cellular structure. Assuming that the critical step in the development of a cell is the ability of the boundary to bow between initial allotriomorphs, a criterion for the occurrence of cellular precipitation was developed. (Diss. Abstr. Int., B)

46018 PHASE DIAGRAM OF THE NIOBIUM-GALLIUM SYSTEM. Pan, V. M.; Latsheva, V. I. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 38, 95-100 (1971). (In Russian).

The phase equilibrium diagram of the Nb-Ga system was constructed from data from differential thermal analysis, tempered microstructure, and x ray analysis. A rather wide region of solid solutions (up to 20 atom % Ga at $1750^\circ C$) was discovered. Six compounds existed at room temperature: Nb_3Ga , Nb_5Ga_3 , Nb_5Ga_4 , Nb_4Ga_5 , Nb_5Ga_{13} , and $NbGa_3$. The single compound $NbGa_4$ existed at high temperatures and in a narrow temperature interval. For each of the compounds the temperature and character of formation were determined. The critical temperature of the superconducting compound Nb_3Ga , annealed at $800^\circ C$ for 10 h, at half-transition was $16^\circ K$. (tr-auth)

46019 PECULIARITIES IN THE STRUCTURAL SENSITIVITY AND MECHANICAL CHARACTERISTICS OF NIOBIUM. Mil'man, Y. V.; Udovenko, A. A. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 35, 34-49 (1971). (In Russian).

A survey was made of the literature on the structure, substructure, and mechanical characteristics of Nb and its alloys, and of the factors that affect them, such as plastic deformation and heat treatment. The influence of grain dimensions, crystal dislocations, and annealing processes are also discussed. (58 references). (K.S.W.)

46020 SEPARATION PROCESSES IN SUPERCONDUCTING ALLOYS OF NIOBIUM-ALUMINUM. Ganzhula, N. N.; Patysheva, V. I.; Pan, V. M.; Chuistov, K. V. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 36, 107-11 (1971). (In Russian).

X-ray studies were made on polycrystalline samples and micro single crystals of transformation processes occurring during heating of quenched alloy Nb-18% Al. Before transformation of a second phase (Nb₃Al) in the narrow temperature interval 750 to 800°C, there occurred ordering of the original solid phase with the formation of a crystal of the CsCl type. Breakdown of the Nb-Al alloy quenched from 1750°C was noticed after heating at 800°C (1 h), and was accompanied by significant hardening. Maximum hardness was achieved after aging for 1 h at 1200°C. During disintegration the intermetallic phase Nb₃Al separated out, with a composition essentially depleted of Al in comparison with the stoichiometric. In an alloy with 15% Al after 70% deformation and annealing at 900°C for 10 h, a diffused transition to the superconducting state began at 14°K, and the critical current density at 4.2°K depended on the magnetic field. (tr-auth)

46021 VARIATIONS IN THE CRITICAL TEMPERATURE OF THE SUPERCONDUCTING COMPOUNDS Nb₃Al AND Nb₃Al_xGe_{1-x}. Pan, V. M.; Latysheva, V. I.; Sudovtsov, A. I. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 36, 104-7 (1971). (In Russian).

The dependence of the critical temperature T_c of the superconducting compounds Nb₃Al and Nb₃Al_xGe_{1-x} on the composition and previous thermal history was investigated. Changes in T_c were contrasted with data on the location of phase boundaries of stability regions of these compounds and data on changes in the density of electron states. The latter were found by electron-positron annihilation. An increase in T_c in the compound Nb₃Al during an elevation on the concentration of Al was associated with a growth in the density of electron states. The highest T_c was obtained in non-equilibrium alloys in which high Al and Ge contents were fixed by rapid cooling. At low-temperature (700 to 750°C) annealing, approaches to an equilibrated composition depleted of Al and Ge did not occur through aging, but the degree of long-range order of atoms increased. (tr-auth)

46022 ORIENTATION DEPENDENCE OF DAMPING OF ELASTIC VIBRATIONS IN TUNGSTEN SINGLE CRYSTALS. Benieva, T. Ya.; Golub, T. V. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 36, 90-4 (1971). (In Russian).

The amplitude dependence of internal friction in single crystals of tungsten was determined over the temperature range 100 to 700°C by a method of damping of free bending movement of the crystals. Tungsten crystals with orientations of [100], [110], and [111] exhibited considerable damping, pronounced amplitude dependence, and an absence of amplitude dependence of internal friction in that region of amplitude deformation observed in special orientation. A critical shearing micro-stress of breaking and its temperature dependence were determined from data on amplitude-dependent internal friction. The smallest value for the critical micro-stress was observed for a W crystal with the orientation [100]; and the greatest value with the orientation [111]. Such behavior for crystals with three orientations apparently stems from anisotropic mobility of boundary dislocations in the W lattice. (tr-auth)

46023 DETERMINATION OF THE ENERGY OF DISLOCATION OF AN IMPURITY ATOM BY AN INTERNAL FRICTION METHOD. Benieva, T. Ya.; Polotskii, I. G.; Skopin, V. S. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 36, 86-90 (1971). (In Russian).

The energy of dislocation of a foreign atom in solid solutions of Cu-Mg, Mo, and W was determined from experimental data on the amplitude dependence of internal energy. The results of the computation were compared with the energy of interaction in Cu-Mg solid solutions from other methods of calculation. (tr-auth)

46024 COMMENTS ON DIFFUSION OF COBALT IN PLUTONIUM BY M. J. DAPTH, M. P. DARIEL AND A. ROSEN IN *SCRIPTA MET.*, 5, 191-194 (1971). Million, B.; Kucera, J. (Inst. of Physical Metallurgy, Brno). *Ser. Met.*; 6, No. 7, 617-19 (Jul 1972).

Comments are given on a previous study on the Co hetero-diffusion in the fcc δ -phase and in the bcc ϵ -phase in Pu with the use of the serial sectioning method. Use of the least squares method applied to 363°C data given previously led to the value $D = 1.96 \times 10^{-11}$ cm²/sec for the volume hetero-diffusion coefficient, to the value $D_1 = 5 \times 10^{-8}$ cm²/sec for the grain boundary diffusion with the grain width $2a = 6 \times 10^{-4}$ cm and to the effective grain size $b = 0.026$ cm. Comparing this Co volume hetero-diffusion coefficient with the Pu self-diffusion coefficient at the same temperature $D = 2.86 \times 10^{-11}$ cm²/sec, it was found that these values are rather close to each other, their portion being 1.46. Thus Pu appears to be different from other polyvalent metals, as Co does not diffuse in Pu with significantly higher diffusion coefficients than is the case for Pu self-diffusion. It was concluded that there is not an especially strong vacancy-cobalt atom impurity bonding in δ -Pu. (P.C.H.)

46025 VACANCY DEFECTS IN QUENCHED VANADIUM. Boostrom, D. R.; Chen, C. W. (Iowa State Univ., Ames). *Ser. Met.*; 6, No. 7, 571-8 (Jul 1972).

Transmission electron microscopy (TEM) was used to examine V foils after quenching and annealing. Use of the Kuhlmann-Schultz method (using superfluid properties of liquid helium(II)) for quenching high-purity V produced defects only when the quenched foils were annealed between 300 and 500°C. The defects were probably complexes of vacancies and interstitial impurity atoms, presumably of oxygen. No evidence was found to indicate the possibility of quenching vacancies into V under the conditions of the quenching technique used and at the present impurity level. The results thus reconfirmed the predictions of others in regard to the effective trapping of vacancies by interstitial impurity atoms in bcc transition metals during quenching. (P.C.H.)

46026 THERMODYNAMIC PROPERTIES OF THE PHASE In₂Bi₃. Singh, H. P.; Rao, M. H.; Misra, S. (Banaras Hindu Univ., India). *Ser. Met.*; 6, No. 7, 621-6 (Jul 1972).

The heats of formation and fusion of the equilibrium phase, In₂Bi₃, were determined calorimetrically. The free energy of formation and the excess free energy of the corresponding liquid alloy were also calculated. Data are tabulated. (P.C.H.)

46027 FREE VOLUME AS A GUIDE TO GRAIN BOUNDARY PHENOMENA. Aaron, H. B.; Bolling, G. F. (Ford Motor Co., Dearborn, Mich.). *Ser. Met.*; 6, No. 7, 553-62 (Jul 1972).

It is suggested that the entire class of properties associated with the presence of static grain boundaries can be rationalized primarily in terms of the amount of free volume without need for a detailed model of the grain boundary structure. A second class of properties associated with the presence of moving boundaries requires knowledge of the distribution of free volume and thus a model for their rationalization. Also, free volume is related to diffusion, and the contribution of free volume to the activation enthalpy for diffusion is quantitatively developed. Two examples are given that suggest how the concept of a free volume and a measure of its amount might be used without need for a detailed model of the boundary. (20 references) (P.C.H.)

46028 TEMPERATURE DEPENDENCE OF THE FRICTION STRESS FOR BASAL DISLOCATIONS IN BERYLLIUM IN THE RANGE 300 TO 500°K. Turner, G. I.; White, J. S. (Atomic Weapons Research Establishment, Aldermaston, Eng.). *Acta Met.*; 20, No. 8, 997-1000 (Aug 1972).

The temperature dependence of the friction stress for basal dislocations in beryllium has been measured in the range 300 to 500°K. The friction stress values were evaluated from the analysis of closed mechanical hysteresis loops developed by the cyclic compression of single crystals. The results showed that the friction stress for this system decreases in an almost linear fashion with increasing temperature from 0.038 kg mm⁻² at 300°K to 0.018 kg mm⁻² at 500°K. For each temperature investigated, the maximum loop width and the corresponding irreversible work data obtained from the family of closed hysteresis loops were fitted to a power law of the type $W = A \epsilon_{MLW}^N$, where W = the irreversible work done in forming a loop, ϵ_{MLW} = the maximum loop width in microstrain and A and N are constants. The friction stress, $dW/d\epsilon_{MLW}$, was then determined using the power law relationship at $\epsilon_{MLW} =$ one microstrain and also from the alternative method involving the use of the elastic limit as obtained by extrapolation of the maximum shear stress versus the maximum loop width curve to zero loop width. Both methods of analysis gave essentially the same values. The significance of the results in terms of the influence of basal slip on the fracture of beryllium is discussed. (auth)

46029 CONTRAST PATTERNS PRODUCED BY SELF-INTERSTITIAL ATOMS IN FIELD ION MICROSCOPE IMAGES OF A B.C.C. METAL. Seidman, D. N.; Lie, K. H. (Cornell Univ., Ithaca, N. Y.). *Acta Met.*; 20, No. 8, 1045-59 (Aug 1972).

A general method is presented for calculating the contrast pattern to be expected from a single self-interstitial atom (SIA) in a field ion microscope (FIM) image. It was assumed that the SIA was embedded inside a FIM spectrum well below the surface. The SIA was uncovered and eventually removed by the field evaporation of successive atomic planes. Atoms on the surface were displayed outwards from the surface as a result of the dilatational field of the SIA. This outward displacement of atoms resulted in 3 possible contrast effects. These 3 contrast effects were: a bright spot; an extra bright spot; and a vacant lattice site. A single SIA atom can produce all these contrast effects and the totality of these effects was termed a contrast pattern. The atom displacements were obtained from the α -iron interatomic potential of Johnson, without allowing for surface relaxation effects. Detailed atomic displacement maps were presented for the {111} planes with a {110} split and a {111} split crowdion SIA configura-

tions located many interplanar spacings underneath these surfaces. It was shown that the resulting atomic displacement maps contained sufficient information to allow the configuration of the SIA to be determined. An atom displaced normal to a surface plane by a SIA was represented by an electric dipole whose strength was a function of the magnitude of the displacement. A local field enhancement factor was then calculated (the possibility of charge redistribution was allowed for in the enhancement factor) and converted to an extra ion current via the Chen-Seidman data on the ion current vs electric field characteristic curves of individual planes. An application of the above principles was presented for a SIA contrast pattern detected in a (111) plane of tungsten irradiated with 20 keV W^+ ions at 18°K (Scanlan et al.). This contrast pattern contained the three contrast effects suggested by the model. (auth)

46030 INITIAL REACTION OF NITROGEN WITH ZIRCONIUM AT 1440°. Sallach, R. A. (Sandia Labs., Albuquerque, N. Mex.). *J. Phys. Chem.*; 76: No. 15, 2156-8 (20 Jul 1972).

The reactions of nitrogen with zirconium at 1440° and pressures between 2 and 10 mtorr were followed using a microbalance and a radiant-heat furnace. At constant pressure there was an initial linear reaction whose duration was inversely proportional to the square of the pressure and which was followed by a transition to a parabolic regime. The linear portion corresponded to the reaction of nitrogen with a β -Zr surface and continued while the α -Zr phase was formed, ending when the α -Zr surface saturated and a thin ZrN layer formed. In the linear regime the sticking coefficient was independent of gas pressure and had the same value for both the β -Zr and α -Zr surfaces—between 0.05 and 0.08. (auth)

46031 SYSTEM NIOBIUM-ALUMINUM-GERMANIUM, ITS PHASE DIAGRAM, AND SUPERCONDUCTING PROPERTIES. (A REVIEW). Pan, V. M.; Patysheva, V. I. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 33, 38-50 (1971). (In Russian).

The Nb angle of the phase equilibrium diagram is presented for the ternary system Nb-Al-Ge in the form of isothermal sections at 1,000 to 2,000°C, polythermal sections for various component combinations, and the surface of the liquidus. The existence of two four-phase peritectic equilibria was established: 1,910 and 1,930°C. There was observed in the solid state a series of solid solutions between the isomorphous compounds β -Nb₃Al and β -Nb₃Ge, whereas the stoichiometry of the compound Nb₃Al_xGe_{1-x} was maintained only in the compositions Nb₃Al_{0.8}Ge_{0.2}–Nb₃Al_{0.7}Ge_{0.3}. Gradual substitution of Al by Ge produced a gradual increase on T_K of the compounds Nb₃Al_xGe_{1-x}, but only during the stoichiometry of this composition. For the alloys of the composition Nb₃Al_{0.75}–Nb₃Al_{0.7}Ge_{0.3} rapidly cooled without extraction and annealed at 700°C, T_K attained 20.1°K with a transition width of 0.3°K. The increase of T_K for these alloys during low-temperature annealing was associated with regulation, and occurred due to disintegration of metastable compounds of the stoichiometric composition which began at a somewhat higher temperature. (41 references) (tr-auth)

46032 DISINTEGRATION OF SUPERSATURATED SOLID SOLUTIONS ACCORDING TO A HETEROGENEOUS MECHANISM. (A REVIEW). Larikov, L. N.; Shmatko, O. A. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 33, 5-37 (1971). (In Russian).

The dependence of the changes in a series of properties of aging alloys on the mechanism of separation was examined, and the simultaneous formation and growth of centers of new phases during heterogeneous decomposition were considered. Aging alloys were classified, and those in which decomposition of supersaturated solid solutions occurred based on a heterogeneous mechanism are listed, which includes alloys of W, In, Mo, and Ti. (301 references) (tr-auth)

46033 ALLOYS OF CHROMIUM WITH TERBIUM. Svechnikov, V. N.; Kobzenko, G. F.; Ivanchenko, V. G. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 33, 93-5 (1971). (In Russian).

The alloys Cr-Tb were investigated by differential thermal analysis, metallography, and hardness testing. An equilibrium phase diagram was constructed over the temperature range 600 to 1,900°C. The results showed that three non-variant transformations may occur, depending on the temperature and composition: $1_1 \rightleftharpoons \alpha_{Cr} + 1_2$ at 1,775°C (monotectic); $1(1_2) \rightleftharpoons \alpha_{Cr} + \beta_{Tb}$ at 1,165°C (eutectic); and $\beta_{Tb} \rightleftharpoons \alpha_{Cr} + \alpha_{Tb}$ at 1,045°C (eutectoid). (tr-auth)

46034 SOLUBILITY OF CERIUM, PRASEODYMIUM, AND NEODYMIUM IN CHROMIUM CONTAINING OXYGEN. Kobzenko, G. F.; Kholodov, A. G.; Ivanchenko, V. G.; Martynchuk, E. L.; Kashevskaya, O. N. (Inst. of Metal Physics, Kiev). *Metallfizika*; No. 33, 95-9 (1971). (In Russian).

The solubilities of Ce, Pr, and Nd in Cr were investigated by x-ray studies, metallographic, and hardness tests, and curves of solubility were constructed for the temperature interval 1,000 to 1,800°C. The limiting solubility of these rare earth metals in Cr amounted to 1.2 (Ce), 0.9 (Pr), and 1.1 at % (Nd). The influence of each metal present as a solid solution in Cr on the microhardness of the alloy was determined. During alloying of the rare earth metals in Cr which contained oxygen, primary and secondary processes of reduction were evident, brought about by the change in solubility of oxygen and rare earth metal in α_{Cr} . The primary process led to a more complete deoxidation of Cr. (tr-auth)

46035 STRUCTURE, PHASE COMPOSITION, AND PROPERTIES OF CAST COMPLEX ALLOYS OF NICKEL-NIOBIUM. Belinskaya, L. A.; Braun, M. P.; Ionkina, E. M.; Lebedev, K. P.; Khilchevskaya, T. V. (Inst. of Foundry Problems, Kiev). *Metallfizika*; No. 33, 102-7 (1971). (In Russian).

The structural changes and certain properties of Ni-Nb alloys up to 8% Nb were studied in connection with small additions of W (3%) and Mo (5.1%). Phase compositions were studied after isolating precipitates in specially selected electrolytes. The microstructures were determined on samples in the cast state and after a thermal treatment, which consisted of quenching from 1,200°C and subsequent annealing at 900°C for 16 hours. (tr-auth)

46036 STRUCTURE AND DECOMPOSITION OF Zr AND Ti B.C.C. SOLID SOLUTIONS. Sass, S. L. (Cornell Univ., Ithaca, N. Y.). *J. Less-Common Metals*; 28: 157-73 (1972).

The literature concerning the decomposition of Zr and Ti alloys is reviewed with particular emphasis on the bcc (β) $\rightarrow$ athermal omega (ω) transformation and the structure of Zr and Ti bcc solid solutions. Illustrative examples were taken principally from the Zr-Nb and Ti-Nb systems. High resolution dark-field electron microscopy, diffraction, and ultrasonic techniques have been used to follow the $\beta \rightarrow \omega$ phase transformation in Zr and Ti alloys. The ω morphology was found to consist of 10 to 15 Å diameter particles arranged in {111} rows. As well, clusters of rows were presented in low solute content alloys and the cluster size decreased with increasing solute content. The observed morphology can explain the presence of {111} planes of diffuse intensity in reciprocal space and the change in appearance of the ω reflections with increasing solute content. The mechanism of the transformation is discussed in terms of instability or metastability of the bcc structure relative to a $\frac{2}{3}$ {111} longitudinal displacement wave. This mechanism can be thought of as involving the {111} displacement of 2 neighboring close-packed rows of atoms relative to each other. Single crystal elastic constant measurements on Zr-Nb alloys showed that C_{44} decreases in the vicinity of the $\beta \rightarrow \omega$ transformation. This is attributed to a softening of the bcc structure which is a precursor to the actual transformation. Alloys that are known to be well above the M_s temperature for the $\beta \rightarrow \omega$ transformation, and hence must be single phase bcc solid solutions, exhibit diffuse ω reflections. It is suggested that the diffuse intensity is due to the presence of a {111} linear fault (or disorder) in the bcc phase. The atom movements associated with this fault are similar to the displacements required for the $\beta \rightarrow \omega$ transformation. (auth)

46037 SOME COMMENTS ON THE BURGERS AND BOGERS: BURGERS TRANSFORMATION MECHANISMS AND THEIR RELATION TO MATHEMATICAL THEORIES OF MARTENSITE CRYSTALLOGRAPHY. Christian, J. W. (Oxford Univ.). *J. Less-Common Metals*; 28: 67-74 (1972).

The relation between single and double shear descriptions of lattice transformations and the current theories of martensite crystallography is discussed in terms of a formalism developed by Acton, Bevis, Crocker, and Ross. The relevance of these theories to nucleation problems is also examined briefly. Particular attention is paid to the bcc $\rightarrow$ orthorhombic transformations and it was concluded that in certain titanium alloys the lattice deformation is a close approximation to an invariant plane strain. The transformation mechanism may thus be similar to Burgers' original proposal, published in 1934. (auth)

46038 IMAGING PROCESS IN FIELD ION MICROSCOPY. Mueller, Erwin W. (Pennsylvania State Univ., University Park). *J. Less-Common Metals*; 28: 37-50 (1972).

Imaging by field ionization is a complex process. Only recently has the original concept of field desorption led to high resolution self-imaging of a surface. The better established use of field ionization in field-ion microscopy revealed experimental facts which in existing quantum mechanical discussions of the basic process are still quite incompletely understood. Problematic are the sharply localized origin of the ions a few Å away from the metal surface, regional and local brightness features indicating chemical specificity within the bcc, fcc, and hcp lattice groups of the pure metals, and selective imaging of constituents of alloys. A new aspect of the ionization process was introduced with the

discovery, by the atom-probe, of field adsorption of image gas atoms at the apex of the substrate atoms. It is proposed that field ionization probability is greatly enhanced when the electron can tunnel through the intermediate apex-adsorbed atom. This hypothesis is applied to the qualitative explanation of several experimental features. (auth)

46039 ELECTRON MICROSCOPIC INVESTIGATION OF NIOBIUM-OXYGEN ALLOYS WITH ADDITIONS OF ZIRCONIUM AND HAFNIUM. Lobodyuk, V. A.; Loktionov, V. A. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 34, 45-9(1971). (In Russian).

Alloys of Nb-O with small additions of Zr and Hf were studied by electron microscopy. Small particles (40 to 120 Å) were observed which were apparently ZrO_2 or HfO_2 , and were occasionally formed as dislocations. The amount of material thus separated decreased with a lowering of the oxygen content. (tr-auth)

46040 TENSILE PROPERTIES OF α -ZIRCONIUM. Reed-Hill, Robert E. (Univ. of Florida, Gainesville). *Rev. High-Temp. Mater.*; 1: No. 2, 97-154(1972).

The mechanical properties of zirconium are considered in terms of tensile test measurements. The tensile properties of metals are widely used for practical design purposes as well as for analyses of the dislocation reactions associated with work hardening and the rate controlling flow-stress mechanism. For this reason emphasis was placed on the fact that the stress-strain behavior can be strongly influenced by a number of factors that have been largely ignored. These include dynamic strain aging as well as dynamic recovery, and in a metal such as zirconium, deformation twinning. It was found that dynamic strain aging tends to make the yield stress temperature (or strain rate) insensitive, to produce anomalous work hardening phenomena, and to cause ductility anomalies such as the "blue brittle" effect. Except for a maximum at elevated temperatures associated with strain aging, the work hardening rate in zirconium as a result of dynamic recovery, decreased continuously with increasing temperature, in a manner analogous to that observed in pure face-centered cubic metals. It was found that twinning is only of significance in zirconium when the specimen texture contains a large fraction of grains with basal planes steeply inclined to the tensile stress axis. A high degree of twinning tends to basically alter the shape of the stress-strain curve. (71 references). (auth)

46041 STRUCTURE OF INTERMEDIATE LAYERS IN DIFFUSION COATINGS. Gabe, D. R. (Univ. of Sheffield, Eng.). *Rev. High-Temp. Mater.*; 1: No. 2, 155-202(1972).

Factors affecting the structure and constitution of metal diffusion coatings are reviewed with particular emphasis on the coating of iron and steel. Diffusion theory and kinetic data are considered in-so-far as they provide evidence of the intermediate alloyed layers. Emphasis is laid upon the fact that most intermediate layers are non-equilibrium in constitution, and the reasons for this feature are discussed in terms of the thermodynamic factors. (167 references). (auth)

46042 TEXTURES OF ROLLING AND ANNEALING IN TUNGSTEN FOILS OF VARIOUS COMPOSITIONS. Zasimchuk, E. E.; Isaichev, V. I. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 34, 62-5(1971). (In Russian).

The predominant orientations in tungsten foils of different compositions were studied by x-ray diffraction after rolling, original recrystallization, and after secondary recrystallization. Orientation of anomalous grains were noted, an increase in these grains being attributed to a decrease in the surface free energy of the system, and occurring at the expense of grains whose crystallographic boundaries coincided with the foil surface and possessed a large surface energy. Impurities may decrease the difference in the surface energies of different boundaries and lead to an increase in the process of secondary recrystallization of certain orientations. (tr-auth)

46043 INFLUENCE OF IMPURITIES AND TEMPERATURE ON THE DEFORMATION OF MOLYBDENUM DURING DYNAMIC LOADING. Benieva, T. Ya.; Golub, T. V. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 34, 56-61(1971). (In Russian).

The temperature dependence of critical micro-stress was determined in Mo single crystals of varied purity, using measurements of the amplitude dependence of internal friction by the method of damping of free bending vibrational motion. The critical micro-stress was determined by the magnitude of the deformation at the moment of breaking, and by the value of Young's modulus allowing for the orientation factor. The height of the potential barrier which had been overcome by dislocations during small shifts was established. The critical micro-stress and Young's modulus were measured as a function of temperature up to about 700°K. (tr-auth)

Radiation Effects

Refer also to abstracts: 45702, 46918, 47229, 47231, and 47303.

46044 (AI-AEC-13033) STUDIES OF VOID FORMATION IN PROTON-IRRADIATED TYPE 316 AND TITANIUM-MODIFIED 316 STAINLESS STEELS. Keefer, D. W.; Pard, A. G.; Kramer, D. (Atomics International, Canoga Park, Calif.). 30 Jun 1972. Contract AT(04-3)-824, 31p. Dep. NTIS.

Void formation was studied as a function of helium content in annealed Type 316 stainless steel irradiated with 1-MeV protons at 500°C. Increasing the helium content resulted in an increased void density, with a concurrent decrease in void size. Swelling was unaffected by changes in helium between 0.28 and 28 appm. In a sample containing no helium, the onset of observable voids was delayed to much higher displacement damage. The effect of periodic increases in helium content in a sample irradiated at 500°C was also studied. This procedure altered the void size and density from that observed in singly-injected samples. Below ~20 displacements per atom (dpa), the swelling was also affected. Void formation was studied in annealed Type 316 stainless steel irradiated at 400, 500, and 600°C for amounts of damage up to 50 dpa. The volume increase at each temperature varied exponentially with displacement damage, and the exponent increased with temperature. Volume increases exceeding 20% were associated with 50 dpa at 500°C and 20 dpa at 600°C; no limits on volume increases were apparent at any temperature. Void number density decreased, and average void size increased, with increasing temperature for a constant damage. A titanium-modified Type 316 stainless steel developed a very inhomogeneous void distribution when proton irradiated at 600°C. However, the addition of titanium did not appear to diminish the volume increase at damages between 5 and 10 dpa, when compared to ordinary Type 316 stainless steel. (auth)

46045 (CONF-710601) RADIATION-INDUCED VOIDS IN METALS. Proceedings of the 1971 International Conference Held at Albany, New York, June 9-11, 1971. Corbett, James W. (ed.) (State Univ. of New York, Albany); Ianniello, Louis C. (ed.) (Atomic Energy Commission, Washington, D. C.). Apr 1972. 884p. Dep. NTIS.

A total of 40 papers were presented at the conference. A separate abstract was prepared for each of 23 papers. Sixteen papers were previously abstracted and appeared in NSA as 26: 31556, 26: 5600, 26: 574, 26: 530, 25: 58300, 25: 58299, 25: 58298, 25: 49486, 25: 49482, 25: 48256, 25: 47143, 25: 47142, 25: 47141, 25: 45103, 25: 37982, and 25: 35503. (J.R.D.)

For abstracts of individual papers see: 45987-45989 and 46046-46065.

46046 (CONF-710601-, pp 31-83) REPORT ON THE CEA PROGRAM OF INVESTIGATIONS OF RADIATION-INDUCED CAVITIES IN METALS: PRESENTATION OF SOME RESULTS. Adda, Y. (Commissariat a l'Energie Atomique, Saclay (France). Centre d'Etudes Nucleaires). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

To establish the fundamental basis of void formation in irradiated metal and to correlate the damage obtained by different nature of particles, different theoretical and experimental studies were undertaken at the C.E.A. These are essentially: computer simulation of atomic displacement cascades and their thermal annealing, theoretical calculation of the nucleation and growth of voids; experimental studies of the swelling and defect supersaturation levels obtained in pure metals under neutron irradiation, and ion bombardment. An overall review of these programs is presented along with the main results. (auth)

46047 (CONF-710601-, pp 125-41) VOID FORMATION IN CLADDING AND STRUCTURAL MATERIALS IRRADIATED IN DFR. Bramman, J. I.; Bagley, K. Q.; Cawthorne, C.; Fulton, E. J. (United Kingdom Atomic Energy Authority, Dounreay (Scotland). Reactor Group). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Immersion density measurements and transmission electron microscopy studies have been carried out on specimens of four stainless steels (AISI M316 and M316L, FV548 and EN58B) and of a nickel-based precipitation-hardening alloy (Nimonic PE16), irradiated in DFR to fluences up to 8.7×10^{22} n/cm² in the temperature range 300 to 650°C. Void formation has been shown to be very sensitive to small differences in composition and pre-irradiation condition. Volume changes for a given material, condition and irradiation temperature can be adequately represented by a linear function of fluence following a threshold exposure during which no swelling of technological significance occurs. Each material studied shows a characteristic temperature dependence of swelling. (auth)

46048 (CONF-710601-, pp 156-73) FLUENCE AND TEMPERATURE DEPENDENCE OF VOID FORMATION IN HIGHLY IRRADIATED STAINLESS STEELS. Appleby, W. K.; Sandusky, D. W. (General Electric Co., Sunnyvale, Calif. Breeder Reactor Dept.); Wolff, U. E. (General Electric Co., Pleasanton, Calif. Vallecitos Nuclear Center). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Void formation in the 300 series austenitic stainless steels has been examined by transmission electron microscopy and the bulk swelling measured by immersion density. Specimens of Types 316 and 347 were obtained from fuel pin clads while the Type 304 was from a materials capsule. Volume increases of up to 10% were measured after fluences of up to 7.3×10^{22} n/cm² E > 0.1 MeV. Void size was found to increase with irradiation temperature while the void number density decreased. The upper temperature limit of void formation in Type 304 stainless steel was found to be $650 \pm 15^\circ\text{C}$. At the higher fluences it was found that void nucleation had effectively ceased and that continued swelling was primarily due to void growth. Extensive intragranular precipitation in Types 316 and 347 at 525 to 550°C encouraged void nucleation. (auth)

46049 (CONF-710601-, pp 174-86) SUPPRESSION OF VOID FORMATION IN STAINLESS STEEL. Laidler, J. J. (Hanford Engineering Development Lab., Richland, Wash.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Work is in progress at HEDL with the goal of developing a cladding material which is resistant to swelling, has good elevated-temperature mechanical properties, and meets the established criteria for LMFBR materials selection. The primary material presently under investigation is AISI Type 348 stainless steel, a niobium-stabilized austenitic alloy. By means of thermomechanical treatments, precipitation of niobium carbide can be caused to occur in association with extrinsic stacking faults. The extensive NbC precipitation which takes place not only enhances the elevated temperature tensile and creep properties of the material, but additionally provides a very high concentration of sinks and traps for point defects. The existence of a highly specific sink for vacancies appears to sufficiently alter the normal void nucleation and growth scheme that it may be possible to suppress void formation completely. Screening experiments utilizing high energy heavy ion bombardment have demonstrated the effectiveness of the microstructure of thermomechanically-treated 348 stainless steel in reduction of the tendency for swelling. Neutron-irradiated specimens have been examined recently and these results are presented. (auth)

46050 (CONF-710601-, pp 230-47) HELIUM MIGRATION IN METALS. Bauer, Walter; Wilson, W. D. (Sandia Labs., Livermore, Calif.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Calculations and experiments on helium migration in several metals are reported. A modification of the Wedepohl method for the determination of interatomic potentials between closed-shell atoms and ions was developed and applied to helium. Both Slater and Kohn-Sham exchange approximations have been employed in the calculation of the atomic charge distributions and also in the overlap region between the atoms. Using five different potentials to describe the copper lattice, the minimum energy configuration of an interstitial He atom was determined. The (1, 0, 0) octahedral position was found to be the most stable He interstitial position, independent of the five potentials used. The activation energy for interstitial He migration was found to lie within the narrow range, .45 eV to .71 eV, and the path of migration was found to be along a (110) crystallographic direction. The energy required to activate a He atom from a substitutional site was found to be ~2 eV. A number of metals such as Ni, Cu, Al, Pd, and stainless steel have been implanted with 300 keV He⁺ ions at 80°K. After implantation, the He outgassing was measured using isochronal and isothermal anneal schedules. Low temperature measurements were carried out in situ using a high sensitivity He mass spectrometer connected to a positive ion accelerator. At high doses (~10¹⁸ He atom/cm²) significant low temperature He outgassing occurs. Most of the outgassing occurs above ~400°C as is the case at lower implantation doses. (auth)

46051 (CONF-710601-, pp 397-429) DEPENDENCE OF RADIATION EFFECTS ON THE PRIMARY RECOIL ENERGY. Robinson, Mark T. (Oak Ridge National Lab., Tenn.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

The status of the analytical theory of the displacement cascade is reviewed with emphasis on those topics of importance for neutron damage in metals. These include atomic cross sections and interatomic potentials, electronic stopping cross sections, the energy lost from displacement cascades by electron excitation, the number of defects produced in a cascade, the effects of target crystallinity, the spatial extent of cascades, and the primary recoil spectrum in neutron irradiations. (auth)

46052 (CONF-710601-, pp 430-48) VOID FORMATION IN METALS DURING ION BOMBARDMENT. Nelson, R. S.; Hudson, J. A.; Mazey, D. J.; Walters, G. P.; Williams, T. M. (Atomic Energy Research Establishment, Harwell (England)). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

The fact that fast-neutron damage in metals can be qualitatively simulated by irradiation with energetic ions from particle accelerators is well known. The technique is discussed, and the conditions necessary for obtaining quantitative design data reviewed. Results are presented which point out the equivalence between the reactor and accelerator irradiations. Furthermore, a selection of results that illustrate the potential usefulness of the technique is shown. For instance, the void swelling obtained as a function of dose for nickel, 316 stainless steel, and Nimonic PE16 is compared to damage doses equivalent to well over 10²³ neutrons/cm². (auth)

46053 (CONF-710601-, pp 479-98) DEUTERON-GENERATED VOID FORMATION IN AUSTENITIC STAINLESS STEELS. Armijo, J. S.; Lauritzen, T. (General Electric Co., San Jose, Calif. Breeder Reactor Dept.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Simulation of in-reactor radiation damage in typical fast reactor core components by high-energy ion bombardment is discussed. The study was performed with several grades of austenitic stainless steel which were impregnated with helium to create an environment similar to that occurring during neutron irradiation and subsequently bombarded with deuterons. Initial experimental conditions were selected to provide a damage state comparable to that caused by a neutron fluence of 1×10^{23} . The specimens were irradiated in target assemblies instrumented to permit constant temperature monitoring and close beam focusing control. Results have revealed the existence of voids in specimens irradiated within the temperature range 400 to 665°C. The voids were observed to be predominantly polyhedral, ranging in size from 52 to 425 Å, and homogeneously distributed in any plane parallel to the bombarded surface. Void volumes as high as 6.5 percent were observed, this quantity having been measured in a region that experienced 83 displacements per atom. (auth)

46054 (CONF-710601-, pp 511-21) SWELLING AS A FUNCTION OF DISPLACEMENT DAMAGE IN PROTON-IRRADIATED 316 STAINLESS STEEL. Keefer, D. W.; Pard, A. G.; Kramer, D. (Atomics International, Canoga Park, Calif.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Specimens of 316 stainless steel, pre-injected with about 5 appM helium, were proton irradiated at 500 and 600°C. Electron microscopy was used to determine volume swelling in specimens having undergone between 0.5 and 50 displacements per atom. Displacements were calculated from the Kinchin-Pease model with a threshold energy of 25 eV. Swelling at 500°C increased with displacement damage up to about 25% for 50 displacements per atom. No evidence of saturation of swelling with increasing damage was observed. The minimum void size increased with damage, and the void density increased with atom displacement rate. These results imply occurrence of nucleation early in the irradiation, followed by continued void growth. A specimen having undergone 100 displacements per atom at 500°C exhibited a void array which was so complex that it is doubtful that a swelling determination can be made. Ferrite, transformed from austenite during the irradiation, was observed in this specimen as small crystallites within the austenite grains. Evidence of austenite-to-ferrite transformation was obtained for specimens with about 5 or more displacements per atom. A specimen irradiated to 18 displacements per atom at 600°C exhibited frequent joining of voids to form a complex array. Preliminary examination indicates swelling at 600°C is greater than at 500°C. (auth)

46055 (CONF-710601-, pp 522-32) OBSERVATION AND THEORY OF THE VOID LATTICE IN MOLYBDENUM. Evans, J. H.; Bullough, R.; Stoneham, A. M. (Atomic Energy Research Establishment, Harwell (England)). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Experimental observations of the void lattice in molybdenum irradiated with 2 MeV N^+ ions are presented together with a theoretical explanation of its formation. Electron micrographs are presented that show the void lattice is body centered cubic and oriented precisely parallel to the underlying molybdenum atomic lattice. The theoretical explanation of this lattice is based on the conjecture that its existence must be a direct consequence of the elastic interaction between pairs of voids in the (cubic) anisotropic molybdenum. The pairwise interaction is shown to be non-central and long-ranged. This interaction has been used to show that such a stable body centered cubic void lattice can exist at a definite ratio of void lattice parameter to void radius. This result has been obtained by both explicit summation of the pairwise interaction and by a novel and exact reciprocal lattice procedure. (auth)

46056 (CONF-710601-, pp 533-49) APPLICATION OF HIGH-VOLTAGE ELECTRON MICROSCOPY TO VOID STUDIES. Buswell, J. T.; Fisher, S. B.; Harbottle, J. E.; Norris, D. I. R.; Williams, K. R. (Central Electricity Generating Board, Bristol (England), Berkeley Nuclear Labs.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Electron irradiation with the beam of a high voltage electron microscope can produce void growth in metals, simulating the effect of high dose neutron irradiation. Thin foil specimens are employed and account must be taken of surface effects. Advantages of the technique are the high dose rate, the relative simplicity of electron irradiation and the ability to make observations continuously during irradiation. In nickel, electron irradiation produces voids only in specimens previously either implanted with gas ions or neutron irradiated. In stainless steel, voids are produced by electron irradiation alone and may be surrounded by strain fields. Direct experimental evidence is provided for dislocations acting as preferential sinks for interstitials, so promoting void growth. Comparative studies in some nickel alloys suggest that void growth depends on dislocations being free to climb and on the interstitials being free to migrate rapidly to them. For stainless steel, the effect of precipitates on void growth is discussed. In both nickel and stainless steel, voids grow at varying rates, sometimes coalescing with each other or shrinking. An analysis is given of the effects observed when neutron-irradiated nickel is thinned and subsequently electron irradiated. (auth)

46057 (CONF-710601-, pp 550-62) APPLICATION OF HIGH-VOLTAGE ELECTRON MICROSCOPY TO THE STUDY OF VOIDS AND OTHER POINT-DEFECT AGGLOMERATES. Urban, K.; Wilkens, M.; Seeger, A. (Max-Planck-Institut fuer Metallforschung, Stuttgart (West Germany)). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

High-voltage electron microscopy at high temperatures provides a powerful tool for studying the nucleation and growth of defect clusters associated with void formation. This method has advantages such as very high damage rates without creation of displacement cascades, and direct observation of defect clusters and their growth. The point-defect concentrations in thin metal foils under electron irradiation were calculated. Using these calculations, it is shown how results obtained in the high-voltage electron microscope may be used to interpret effects in bulk material of technically relevant dimensions. In nickel, the formation and growth of interstitial and vacancy dislocation loops and voids during irradiation in the temperature range from 200 to about 600°C were observed. Vacancy agglomerates grew only near edge dislocations that acted as preferential sinks for interstitials. Voids were observed only in samples containing gaseous impurities. From the fact that voids occurred in samples in which the impurities were not introduced by prebombardment with high-energy gas ions it may be concluded that displacement cascades and presence of gases of extremely low solubility are not a necessary pre-requisite for void formation. (auth)

46058 (CONF-710601-, pp 563-604) VOID FORMATION IN QUENCHED OR IRRADIATED METALS. Balluffi, R. W.; Seidman, D. N. (Cornell Univ., Ithaca, N. Y. Dept. of Materials Science and Engineering). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

The detailed mechanisms responsible for the formation of voids in both quenched and irradiated metals are discussed from a critical point of view. Attempts are made to correlate void formation phenomena under quenching and irradiation conditions whenever possible. Particular attention was given to the void nucleation problem and to the question of whether the nucleation is homo-

geneous or heterogeneous. The importance of an increased understanding of the conditions for interstitial dislocation loop nucleation in the irradiation case was emphasized. Aspects of void growth problem are considered, and the effectiveness of voids (and dislocations) as point defect sinks is discussed. A number of areas where useful experiments can be performed are pointed out. (auth)

46059 (CONF-710601-, pp 613-29) PRODUCTION OF VACANCY CONDENSATES BY LASER BOMBARDMENT. Smidt, F. A. Jr.; Metz, S. A. (Naval Research Lab., Washington, D. C.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Bombardment of thin foils by a laser pulse of 3.5×10^{-8} sec duration and 1.5 cal/cm^2 total energy input has been found to produce high concentrations of vacancies in 99.98% pure nickel and vanadium. The vacancies condense upon annealing the foils at $0.31 T_m$ to form voids in nickel and faulted loops and a few isolated tetrahedra in vanadium. Although it was initially believed the vacancies formed as a result of quenching, concentrations as high as 1% or 10^2 times that expected at the melting point have been observed. The generation of vacancies by nonconservative motion of jogs does not appear feasible either because the dislocation density is too low. Interaction of shock waves generated by the laser pulse are suggested as a possible alternative explanation. The technique offers a supplementary approach to the study of vacancy condensates and the factors that affect their nucleation. (auth)

46060 (CONF-710601-, pp 672-83) EFFECT OF IRRADIATION TEMPERATURE ON SIZE OF DEFECT AGGREGATES IN COPPER. Larson, B. C.; Young, F. W. Jr. (Oak Ridge National Lab., Tenn.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Nearly perfect copper crystals have been irradiated with 10^{18} fast neutrons per cm^2 ($E > .6 \text{ MeV}$) at temperatures varying from 43 to 400°C, and the resulting decrease in the integrated anomalously transmitted intensities has been measured for $\text{CuK}\alpha$, $\text{MoK}\alpha$, and $\text{AgK}\alpha$ x-rays. The decreases have been expressed in terms of effective absorption coefficients which are proportional to the point defect concentration and the first and second moments of the defect aggregate size distribution. These data indicate little change in the defect structure for irradiation temperatures from 43 to 200°C; however, for temperatures above 200°C the absorption coefficients rapidly increased by a factor of about 20 at 300°C, indicating a rather large increase in the sizes of the defect aggregates. This conclusion is supported by the wavelength dependence of the coefficients, line broadening of the Bragg profiles, and electron microscopy and x-ray topography. The maximum effect occurred at 300 while at 400°C (which is above the onset of stage IV annealing) there was no damage detectable for these neutron doses. Results of similar measurements made on nearly perfect aluminum crystals that had been irradiated at 43 and 127°C for doses up to $7 \times 10^{18} \text{ n/cm}^2$ indicated no such effect in aluminum for these temperatures. (auth)

46061 (CONF-710601-, pp 739-56) COMPUTER SIMULATION OF ATOMIC DISPLACEMENT CASCADES IN METALS. Torrens, Ian M. (Commissariat a l'Energie Atomique, Saclay (France). Centre d'Etudes Nucleaires); Robinson, Mark T. (Oak Ridge National Lab., Tenn.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

A comprehensive computer code has been developed to simulate atomic displacement cascades in a wide variety of crystal lattices generated from primary atoms of energy up to more than 100 keV. The model includes a realistic potential for elastic atomic collisions, inelastic energy losses to electrons, and an adiabatic approximation to finite temperature effects. Results of a statistical analysis of cascades in copper and iron are presented. These include the energy variation of the total number of point defects produced, the effect of thermal vibrations on cascade development, and information on vacancy cluster distribution at the end of the cascade. Comparison is made with previous simulation results and with analytical theory. The results are used as initial data for a separate cascade thermal annealing code with the aim of obtaining information on the nucleation of voids in irradiated metals. (auth)

46062 (CONF-710601-, pp 769-97) THEORY OF VOID FORMATION AND GROWTH IN IRRADIATED MATERIALS. Bullough, R.; Perrin, R. C. (Atomic Energy Research Establishment, Harwell (England)). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

A detailed theory of void formation and growth is presented and correlated to the appropriate accelerator measurements in nickel and steel. The complex dependence of the swelling phenomenon on the various physical and structural properties of the irradiated materials is discussed with particular emphasis on the various mechanisms that can prevent, slow down, or completely stop the swelling process. (auth)

46063 (CONF-710601-, pp 798-824) THEORETICAL STUDY OF THE SWELLING OF FAST-NEUTRON-IRRADIATED MATERIALS. Harkness, S. D.; Li, Che-Yu (Argonne National Lab., Ill.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

Possible mechanisms for limiting void growth under fast neutron irradiation are analyzed. It is found that the void volume at which saturation should occur is dependent on the void number density. Higher void number densities lead to lower void volumes at saturation. The importance of knowing the generation rate of Frenkel pairs that can participate in diffusional processes rather than the total number of displaced atoms is discussed. It is stressed that this could be especially important in translating results gained by proton bombardment. The relationship between the development of voids under heavy ion and under neutron bombardment is considered using a theoretically based computer model. In this model void nucleation is taken as the sum of a homogeneous nucleation rate and a heterogeneous term. The homogeneous nucleation rate is taken as that recently derived by Russell and by Katz and Wiedersich. The heterogeneous nucleation rate is an empirically fitted term based on the form expected from nucleation by helium. It is found that homogeneous nucleation can adequately explain the results from heavy ion bombardments where the target was not previously injected with helium. Homogeneous nucleation in EBR-II irradiations is only found to be important for short exposures ($nvt < 10^{22}$ n/cm²) at irradiation temperatures below 400°C. (auth)

46064 (CONF-710601-, pp 825-40) NUCLEATION OF VOIDS IN MATERIALS DURING HIGH-ENERGY IRRADIATION. Katz, Joseph L. (Clarkson Coll. of Tech., Potsdam, N. Y. Dept. of Chemical Engineering); Wiedersich, Hartmut (North American Rockwell Corp., Thousand Oaks, Calif. Science Center). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

High energy radiation simultaneously produces vacancies and interstitials in crystalline materials. The nucleation of defect precipitates under this circumstance is fundamentally different from the usual nucleation problem because of the competing effects of interstitials and vacancies on the size distribution of precipitate embryos. The rate of homogeneous nucleation of voids in such a material, in which interstitials as well as vacancies are supersaturated, is derived. It is shown that from thermodynamics only $n^0(x)$, the distribution of void sizes when the interstitials are in equilibrium with the vacancies can be obtained. The constrained equilibrium distribution of voids $n(x)$ when the interstitials are also supersaturated is determined by the kinetics of incorporation of vacancies and interstitials into the embryos. Explicit expressions are given for $n(x)$ in terms of kinetic parameters in $n^0(x)$. The rate of nucleation is then derived from $n(x)$ along lines similar to classical nucleation theory. It is found that the supersaturation of defects which occur during irradiation in a high flux of energetic neutrons or ions is sufficient for readily observable nucleation rates of voids at intermediate temperatures but insufficient at high temperatures. (auth)

46065 (CONF-710601-, pp 841-52) EFFECT OF STRESS ON RADIATION-INDUCED VOID GROWTH. Garner, F. A.; Wolfer, W. G.; Biancheria, A.; Boltax, A. (Westinghouse Electric Corp., Madison, Pa. Advanced Reactors Div.). Apr 1972.

From International conference on radiation-induced voids in metals; Albany, N. Y. (9 Jun 1971).

The influence of applied stress on the radiation-induced density changes observed in fast reactor structural materials is currently under study. Candidate mechanisms being investigated as potential contributors to density changes include the growth of grain boundary cavities or helium bubbles under tension; the volume change associated with the hydrostatic component of irradiation creep; and the growth of voids under conditions of externally applied stress. Emphasis is placed on the last mechanism. The changes in point defect currents that arise in the vicinity of voids in a stressed matrix is also investigated. (auth)

46066 (COO-3431-1) STUDY OF RADIATION INDUCED DEFECTS IN METALS. Final Report, January 15, 1966-January 14, 1972. Townsend, John R. (Pittsburgh Univ., Pa. Dept. of Physics). 5 Jun 1972. Contract AT(11-1)-3431. 6p. Dep. NTIS.

The topics discussed fall under three major categories: (1) effects of low temperature proton irradiation on the Young's modulus and internal friction of copper and tungsten, (2) computer simulation of point lattice defects and the resulting changes in the elastic bulk modulus, and (3) piezo-resistance in metals arising from stress-induced ordering of point defects. (auth)

46067 (HEDL-TME-71-126) RECOVERY AND RECRYSTALLIZATION OF PROTOTYPIC FTR FUEL CLADDING: 20 PERCENT CW 316 STAINLESS STEEL TUBING (RM-11). Paxton, M. M.; Holmes, J. J. (Hanford Engineering Development Lab., Richland, Wash.). Sep 1971. Contract AT(45-1)-2170. 25p. Dep. NTIS.

Design, performance and safety analyses of FTR core components are presently based on conservative property estimates because knowledge concerning the thermal stability of cold-work in Type 316 stainless steel is meager. Information concerning the retention of cold-work properties in 316 SS after exposure to the FTR core environment is presented. It appears that at all operating temperatures below 1200°F, use of a factor representing 75% of the tensile strength of 20% CW 316 SS as a design assumption is warranted. (auth)

46068 (ORNL-TM-3733) POSTIRRADIATION MECHANICAL PROPERTIES OF TYPES 304 AND 304 + 0.15 PERCENT TI STAINLESS STEEL. Bloom, E. E.; Stiegler, J. O. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 30p. Dep. NTIS.

The tensile and creep-rupture properties of Types 304 and 304 + 0.15% Ti stainless steels have been determined after irradiation at temperatures in the range 400 to 820°C to maximum neutron fluences of 3.5×10^{22} neutrons/cm² (>0.1 MeV). Changes in mechanical properties were related to the microscopic observations of irradiation-produced defects. When irradiated in the annealed condition in the neighborhood of 450°C, Type 304 stainless steel exhibited an increased yield stress, reduced strain hardening coefficient, and reduced uniform and total elongation. The increased yield stress could be correlated with the strengthening expected from irradiation-produced voids and dislocations. With increasing irradiation temperature the concentration of these defects decreased and thus the magnitude of the yield stress increase became less. At 500 to about 600°C irradiation and test temperature a pronounced reduction in creep-rupture ductility was observed. Fractures were intergranular. It is suggested that in this temperature range the void-dislocation structure together with the transmutation-produced helium were responsible for the intergranular fractures and low ductilities. At higher temperatures no void-dislocation structures were formed. For these conditions the ductilities were higher than in the 500 to 600°C range but still significantly below the unirradiated value. Variation in alloy composition and preirradiation microstructure had a strong influence on the postirradiation properties. Type 304 + 0.15% Ti stainless steel exhibited significantly higher tensile and creep-rupture ductilities than the standard alloy when irradiated and tested above 450°C. The strength properties of specimens irradiated in the 10% cold-worked condition were similar to those of material irradiated in the annealed condition, but the total elongation and reduction in area were slightly lower. (auth)

46069 (UJV-2795-M) ROLE OF STATISTICAL FLUCTUATIONS IN TEXTURE DURING THE RADIATION GROWTH IN URANIUM. Novak, J. (Ceskoslovenska Akademie Ved, Rez, Ustav Jaderneho Vyzkumu). 1972. 11p. (In Czech). Dep. NTIS (U. S. Sales Only).

The effects of statistical fluctuations of the preferred orientation of uranium grains on the fuel element distortion under irradiation were examined. It has been supposed for simplicity that uranium grains grow under irradiation in direction b and contract isotropically in perpendicular directions. The model yielded a value of mean square of the angle between directions of the axis of the distorted cylindrical specimen determined at both basal planes of the cylinder as a function of the radiation growth of individual grains, of the grain size, and of geometry of the cylindrical specimen. Numerical values obtained using the model were in accordance with the result of an irradiation experiment. (auth)

46070 (UJV-2798-M) DEFORMATION OF FUEL ELEMENT RODS DURING IRRADIATION. Novak, J.; Pelcik, E. (Ceskoslovenska Akademie Ved, Rez, Ustav Jaderneho Vyzkumu). 1972. 25p. (In Czech). Dep. NTIS (U. S. Sales Only).

Temperature and stress dependence of the axial deformation rate of the metallic uranium cylindrical fuel elements during irradiation in a gas loop of the reactor VVRS has been studied. The temperature of uranium was below 350°C. The results showed an influence of the can on fuel element deformation. (auth)

46071 IRRADIATION EFFECTS ON STRUCTURAL MATERIALS. I. RADIATION HARDENING. Bement, A. L. Jr. (Massachusetts Inst. of Tech., Cambridge). *Rev. Roum. Phys.*; 17: No. 3, 361-80(1972).

A fundamental discussion on neutron radiation hardening is given under the following topics: dispersed barrier hardening, temperature dependence, impurity effects, strain hardenability, and grain size effects. (66 references) (P.C.H.)

46072 CORRELATION OF IRRADIATION DATA USING ACTIVATION FLUENCES AND IRRADIATION TEMPERATURE. Lynch, John H. (Lewis Research Center, Sandusky, Ohio). *Nucl. Technol.*; 15: No. 3, 411-21(Sep 1972).

A new method of correlating radiation damage data is tested using data taken from the open literature. This method, the activation fluence method, was found to be as accurate as other contemporary models with which it was compared. The new method also has several advantages over the other methods. The method employs a new entity, the activation fluence (time integrated specific activation rate), as the independent variables in a regression model. Temperature at which the irradiation takes place is also a variable. Although the method was treated for a specific type of damage (change in nilductility transition temperature for A302-B steel) it has no inherent restrictions and is limited only by the imagination of the user. (auth)

46073 ROOM TEMPERATURE INTERNAL FRICTION OF NEUTRON-IRRADIATED ALUMINUM. Bose, S. K.; Mishra, R.; Bhattacharya, D. L. (Banaras Hindu Univ., India). *Indian J. Pure Appl. Phys.*; 10: No. 1, 21-2(Jan 1972).

It has been observed that at low frequencies (~1 Hz) the amplitude-independent room temperature internal friction (Δ) of a neutron-irradiated aluminium wire specimen has always a value higher than that of an annealed specimen. It is possible to explain the observed effect in terms of an increase in the mean distance (L_c) between two pinning points on dislocation lines. The break-away stress value of an irradiated and annealed specimen has also been explained in terms of a slightly modified Granato-Luecke theory. (auth)

46074 RANGES OF FISSION FRAGMENTS IN URANIUM-ALUMINUM ALLOYS AND URANIUM OXIDES. Saeki, Masakatsu; Ishimori, Tomitaro (Japan Atomic Energy Research Inst., Ibaraki). *Nippon Genshiryoku Gakkaishi*; 14: No. 6, 278-82(Jun 1972). (In Japanese).

The mean ranges of 6 fission fragments (^{91}Y , ^{95}Zr , ^{99}Mo , ^{132}Te , ^{140}Ba and ^{144}Ce) have been determined in U-Al alloys (4.66, 16.20, 20.15 wt% U), UO_2 , and U_3O_8 by radiochemical measurement. The results obtained are tabulated. (auth)

46075 CHANGE IN LINEAR DIMENSIONS OF METALS UNDER THE INFLUENCE OF γ RADIATION. Gurevich, M. E.; Shalaev, A. M. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 36, 113-15(1971). (In Russian).

The change in length, $\Delta l/l$ of samples of Al, Ni, Cu, and Mo, which had been irradiated with 10^9 R of ^{60}Co gamma radiation, was experimentally measured during subsequent isochronous annealing. Dimensional change was associated with the annealing of point defects, which had been produced by γ irradiation. (tr-auth)

46076 FORMATION OF DEFECTS IN A CRYSTALLINE LATTICE DURING IRRADIATION OF METALS AND ALLOYS (A REVIEW). Shalaev, A. M. (Inst. of Metal Physics, Kiev). *Metallofizika*; No. 34, 5-20(1971). (In Russian).

A literature survey was made of theoretical and experimental work conducted on radiation defects in metals and alloys. Primary and secondary effects are differentiated. The displacement of atoms by neutrons, gamma radiation such as that from ^{60}Co , and electrons is considered. (52 references). (K.S.W.)

Other Materials

Refer also to abstracts: 45338, 45944, 46013, and 47225.

46077 (GULF-GA-A-12035) IRRADIATION-INDUCED DIMENSIONAL CHANGE IN HTGR CONTROL MATERIALS. Stansfield, O. M. (Gulf General Atomic Co., San Diego, Calif.). 28 Apr 1972. Contract AT(04-3)-167. 27p. (CONF-720420-8). Dep. NTIS.

From Conference on reactor materials performance; Richland, WA. (24 Apr 1972).

Boronated graphites containing 22 to 43 wt % boron as B_4C and representing current warm-pressing or extrusion technology were irradiated at $650 \pm 100^\circ\text{C}$ to fast neutron fluences up to 7×10^{21} n/cm ^{2} ($E > 0.18$ MeV). The irradiation caused an increase in thermal expansivity, a decrease in thermal conductivity, and an anisotropic dimensional change related to the preferred orientation of

the graphite crystallites in the graphite matrix. Although irradiation-induced B_4C swelling has been reported, the dimensional change in boronated graphite is not significantly influenced by that effect. Dimensional change in boronated graphite is controlled by the fast neutron and ^{10}B fission damage irradiation properties of the graphite-binder matrix. At a constant ^{10}B -isotope concentration, the irradiation-induced dimensional change of the boronated graphite increases with increasing ^{10}B -isotope enrichment of the boron in the B_4C addition. This effect, which may result from more severe ^{10}B fission fragment damage in the matrix surrounding ^{10}B -enriched B_4C , leads to distortion directly related to the ^{10}B burnup gradient. The use of natural boron in the fabrication of boronated graphite results in dimensional changes independent of the ^{10}B burnup gradient and correlated with fast neutron fluence under HTGR service conditions. (auth)

46078 (NYO-1710-106) RESEARCH ON GRAPHITE. Final Report for Work Completed During the Period 1954 to 1972. Walker, P. L. Jr.; Kline, D. E.; Thrower, P. A. (Pennsylvania State Univ., University Park, Dept. of Materials Science). 30 Jun 1972. Contract AT(30-1)-1710. 28p. Dep. NTIS.

Major areas of research on carbons and graphite under the contract include carbon formation and graphitization (1954 to 1972), gas-graphite interaction and solid state studies (1954 to 1972), dynamic mechanical properties of graphites and carbons (1966 to 1972), and radiation damage to graphite (1968 to 1972). Based on research results, over 100 technical publications were written. A complete list of these publications is given along with a list of papers submitted for publication. In this way the extent of text material has been kept to a minimum, with only what are considered the major accomplishments being emphasized. (J.R.D.)

46079 (UCRL-Trans-10622) EQUATION OF STATE FOR LIQUID AMMONIA. Planck, R. Translated for Univ. of California Lawrence Livermore Lab., Livermore, from Kaeltetechnik; 12: No. 10, 282-3(1960). 4p. Dep. NTIS.

The specific volume of liquid ammonia as a function of pressure and temperature may be calculated very accurately by means of an equation given by Eucken. The numerical constants of this equation are presented. The numerical evaluation is relatively easy. (auth)

46080 USE OF REINFORCED PLASTIC IN A URANIUM PRODUCTION FACTORY. Walz, Albert. *Corros.-Trait., Prot., Finition*; 20: No. 2, 155-8(Mar 1972). (In French).

Various equipment of uranium production factory was exposed to severe corrosion conditions: installations for obtaining uranium oxide from ore by solvent extraction methods; installations for the processing of uranium oxide. The Cother corporation factory of Canon City extensively utilizes SVR reinforced plastics in their chemical plants for corrosion protection. The manufacture of SVR ATLAC 382 and its application to the development of solvent extraction tanks is described. (France)

46081 COMPARATIVE EFFECTS OF ION BOMBARDMENT AND PARTIAL PRESSURES ON SHEET POLYETHYLENE. Antoniou, Antoine; Bui Ai; Lacoste, Robert; Mayoux, Christian (Ecole Nationale Supérieure d'Electrotechnique d'Informatique, et d'Hydraulique, Toulouse). *Compt. Rend., Ser. B*; 274: No. 16, 955-8(17 Apr 1972). (In French).

The damage caused on polyethylene by ion beams (200 to 2000 eV) was studied and compared to that induced by partial discharges. Observations by multiple and specular reflexions in infrared spectroscopy revealed structural changes. The existence of a proportional law between the particle mass and the damage rate being established, two hypotheses to explain the material erosion by the discharges are proposed. (France)

46082 AEROSOLS PRODUCED BY X-RAYS. Keng, Edward Y. H.; Chu, Richard R. C.; Knight, James A. Jr.; Orr, Clyde Jr. (Georgia Inst. of Tech., Atlanta). *J. Colloid Interface Sci.*; 39: No. 1, 94-102(Apr 1972).

Small proportions of organic vapors in air produce aerosols under the influence of x-rays. The particulates are generally spherical in shape and less than a micron in diameter. Those formed from acetylene and benzene vapor appear yellow and brown, respectively. They are essentially nonvolatile at room temperature and are very stable even when exposed to the high vacuum of an electron microscope. However, a substantial size reduction was attained by several hours of heating at 250°C under atmospheric pressure. The size and quantity of the particulates formed under x-irradiation depend on the organic vapor, its concentration, the humidity of the air, the intensity of the x-rays, irradiation time, and aging time after exposure. The growth and formation of particulates do not stop immediately after exposure ceases. Infrared spectrum analysis provides information on the chemical structure of the particulates. (auth)

46083 METHOD AND APPARATUS FOR COUNTING RADIATION DAMAGE TRACKS IN SHEET MATERIAL USING A CAPACITIVE DISCHARGE DEVICE. Cross, William Gunn; Tommasino, Luigi (to Atomic Energy of Canada Ltd.). U. S. Patent 3,663,815. 16 May 1972. Priority date 18 Feb 1969, Canada.

Fission fragment tracks produced in sheet mica and other electrical insulating materials of radiation dosimeters are counted by leaching the tracks so that they provide electrical discharge paths through the sheet, placing an aluminum foil on top of the sheet, applying a voltage between an electrode covering the underside of the sheet and the foil to sequentially puncture the foil by electrical discharges through the tracks and counting the discharges. (Official Gazette)

46084 MORPHOLOGY AND MECHANICAL PROPERTIES OF CELLULOSE ACETATE GRAFTED WITH POLYACRYLONITRILE. Siahkolah, Mohammed Ali. Raleigh, N. C.; North Carolina State Univ. (1972). 201p. University Microfilms Order No. 72-17,740.

Thesis.

Stretched and normal (unstretched) cellulose acetate yarns were grafted with acrylonitrile monomer from the vapor-phase using γ irradiation as the graft initiator. The grafted samples, in general, were characterized by measuring and studying their tenacity, orientation, amount of the actual graft and the parent homopolymers, moisture regains before and after saponification, homogeneity, compatibility, graphitizability, and finally their electron micrographs. The dynamic mechanical properties of the grafted samples were measured by Vibron and were thoroughly studied. Attempts were made to correlate the experimental results to those calculated based on certain mechanical models. Finally, the molecular relaxation time, τ , a characteristic of the system, was found by subjecting the grafted and ungrafted samples to successive interval sorption tests with acetone. A new method was found to calculate the relaxation times throughout the whole course of the experiment, with a more accurate value. These and other experimental methods showed that the cellulose acetate grafted with polyacrylonitrile is a rather homogeneous system. These two polymers construct the grafted system such that it may be considered to be in the form of two continuous interpenetrating matrices. These matrices not only share the external load based on their relative compositions, but due to the presence of a special kind of crosslinking the modulus tends to be higher than the calculated value. The segmental environments of the grafted samples are believed to be relatively heterogeneous. However, evidence for the existence of any aggregates of like-molecules to an appreciable size that could scatter the light and lead to an opaque film was lacking. (Diss. Abstr. Int., B)

46085 SYNTHESIS AND STUDY OF LANTHANIDE PEROXIDES. Bogdanov, G. A.; Selivanova, M. N.; Garkushenko, T. L. (Moscow Technical Inst.). Zh. Neorg. Khim.; 17: No. 3, 616-20 (Mar 1972). (In Russian).

New peroxides of La, Sm, Er, and Lu were synthesized, and it was noted that the ability to form peroxide compounds increased from La to Lu. The stability of a given lanthanide was inversely proportional to the quantity of peroxide groups attached to it. Differential heating curves suggest a complex character; the peroxides decompose gradually, releasing oxygen and water and causing some exo- and endo-effects. (tr-auth)

46086 RESEARCH AND DEVELOPMENT ON GRAPHITE IN THE FEDERAL REPUBLIC OF GERMANY. Fitzer, E. (Univ., Karlsruhe, Ger.). Ber. Deut. Keram. Ges.; 49: No. 5, 141-9 (May 1972). (In German).

Research institutions as well as the most important research and development results with respect to artificial carbon products are surveyed. The trend for future application of artificial carbon products is outlined. In addition to the broad application in filaments, felts, and fibers for reinforcement of netted or thermoplastic polymers, special application of carbon fiber reinforcement of compound materials is discussed with respect to ultracentrifuges and rotor blades of helicopters. Of great interest is

the construction of a carbon fiber-reinforced glider with wings of 30 m span. Such wingspans were not considered before. (H.B.G.)

46087 EFFECT OF NUCLEAR RADIATIONS ON RUBBER JOINTS. Monnier, Bernard; Van de Voorde, Marcel (CERN, Geneva). Ind. At. Spatiales; 26: No. 1, 39-50 (Jan 1972). (In French).

The CERN undertook studies for the determination of the most suitable rubber for vacuum techniques in a radioactive environment. The changes in the various mechanical characteristics with the different absorbed doses were used to determine the limits of each type of rubber. (France)

46088 PULSED METHOD FOR IMPREGNATION OF GRAPHITE. Beatty, Ronald L.; Kiplinger, Dale V. (to United States Atomic Energy Commission). U. S. Patent 3,664,859. 23 May 1972. Filed 4 Mar 1969.

An improved method for preparing carbon-impregnated graphite bodies is provided comprising pulsing the graphite bodies between a vacuum atmosphere and a pressurized hydrocarbon atmosphere at elevated temperatures. The resulting graphite bodies, which are impregnated with pyrolytic carbon deposits, have helium permeabilities of less than 1×10^{-11} cm²/sec. (AEC)

46089 ADHERING RESINS TO SUBSTRATES, ESPECIALLY METAL, BY RADIATION. Hall, Roger P.; Pratt, Ivor; Young, Richard A. (to SCM Corp.). U. S. Patent 3,669,796. 13 Jun 1972. Filed 3 Oct 1968.

A process for radioinduced coating of a substrate, and especially one having a metallic surface, with a substantially catalyst-free system containing a polymerizable organic unsaturated resin susceptible to free-radical catalysis; and the resulting product are described. In one form, a film of the resin is superimposed on the substrate while a facing side of either the resinous film or substrate is contacted at any time prior to such radiation with an organic substituted, radiation-responsive silane, or derivative thereof such as a siloxane or polysiloxane of the silane. Thereafter, the film and substrate are subjected to high-energy radiation to induce adherence of one to the other. The process is also adapted for coating articles with normally air-inhibited, thermosetting resins by a two-step process, wherein the resin film is first passed through one treating zone effective to impart mass integrity and thereby define a sheet, and the sheet together with the silane and the substrate is then passed through another treating zone effective substantially to complete the cure of the resin and simultaneously adhere the sheet to the substrate, at least one of the treating zones comprising exposure to high energy radiation. (Official Gazette)

46090 USE OF IRRADIATION IN PLASTIC WASTES DISPOSAL. Tanaka, Masaru; Mifune, Tsuyoshi (Wayne State Univ., Detroit). J. Sanit. Eng. Div., Amer. Soc. Civil Eng.; 98: SA-3, 561-7 (Jun 1972).

Radioinduced embrittlement of plastics for promotion of waste disposal by consolidation was investigated. Results of embrittlement reactions caused by fast electrons are reported for widely used plastic materials such as polyvinyl chloride and polyethylene. It was found that extended irradiation is required to render these plastics brittle enough for mechanical consolidation. The method cannot be recommended for commercial use. (J.R.D.)

46091 SURFACE TENSION OF STRESS-ANNEALED PYROLYTIC GRAPHITE. Morcos, Ikram (Hydro-Quebec Inst. of Research, Varennes, Quebec). J. Chem. Phys.; 57: No. 4, 1801-2 (15 Aug 1972).

Previous measurements and some of the assumptions made in calculating γ_s for cleavage graphite are discussed. Neumann's derivation of Good's equation is given: $\cos \theta = [(0.015 \gamma_s - 2)(\gamma_s \gamma_L)^{1/2} + \gamma_L] / \gamma_L [0.015(\gamma_s \gamma_L)^{1/2} - 1]$. Advancing and receding meniscus measurements were made with five liquids on stress-annealed highly oriented type Z Y H graphite at 25°C. The results indicate that γ_s is probably ~ 34.6 dyn/cm. The lower γ_s values (26.5 and 28.75 dyn/cm) obtained with CCl₄ and acetonitrile are due to the γ_L being lower than γ_s and the consequent adsorption of vapors on the graphite surface. (D.L.C.)

NUCLEAR MATERIALS AND WASTE MANAGEMENT

Accountability and Safeguards

Refer also to abstracts: 45183-45186, 45190, 45191, 45587, and 46148.

46092 (BNL-16930) CONTINUOUS SAMPLING PLAN FOR THE TAKING OF PHYSICAL INVENTORIES. Epel, L. G.; Marcuse, W.; Zucker, M. S. (Brookhaven National Lab., Upton, N. Y.). Jun 1972. 13p. (CONF-720529-3). Dep. NTIS.

From thirteenth annual meeting of the Institute of Nuclear Materials Management; Boston, MA. (31 May 1972).

One method of performing a periodic physical inventory involves the stopping of flows into and out of the material balance area being inventoried, determining that all items appearing in the records are actually in inventory, and weighing and sampling a subset of these items on a random basis for chemical and/or isotopic analyses. This procedure may interfere with production, will intermittently load the analytical laboratories, and results in delays in establishing the amount of special nuclear material that is in inventory. These undesirable effects become more onerous as the period between physical inventories is shortened. A continuous sampling plan is described that eliminates these effects and in addition makes possible a more continual knowledge of the inventory. Further, with this plan and under proper conditions of flow and inventory size, a potential for reducing the total number of samples required to achieve equivalence with the usual method of establishing an inventory exists. The continuous sampling plan is based on a stochastic process of shipping and sampling items that yields a probabilistic model for the number of items in inventory that have been sampled at any time. The expected value of the number of sampled items can be made equal to the number of sampled items in the usual inventory method by a suitable choice of parameters. Further, any function that depends on the number of sampled items in inventory also becomes a random variable and can be made to have the same expected value as it has in the periodic physical inventory method. (auth)

46093 (ICP-1014) EVALUATION OF INSTRUMENTATION FOR NUCLEAR FUELS REPROCESSING PLANT INPUT WEIGHT MEASUREMENTS. Groth, F. M.; Cartan, F. O. (Allied Chemical Corp., Idaho Falls, Idaho, Idaho Chemical Programs Operations Office). Jul 1972. Contract AT(10-1)-1375. 33p. Dep. NTIS.

Results of a study at the Idaho Chemical Processing Plant were used to evaluate instrumentation for measurement of plant input solution weight. A system to directly weigh up to five tons of radioactive solution in an input accountability tank was installed and calibrated but found to be unsatisfactory in this installation. Two methods to determine liquid weight using its pressure in a tank were also investigated, including complete error analyses. A precision pressure transducer measurement of solution weight gave a 0.10% standard deviation—the best among those instruments evaluated. Recommendations for improving input weight measurements are given. (auth)

46094 (WASH-1210) ATTRIBUTE SAMPLING INSPECTION PROCEDURE BASED ON THE HYPERGEOMETRIC DISTRIBUTION. Sherr, Theodore S. (Division of Nuclear Materials Security (AEC), Washington, D. C.). May 1972. 215p. Dep. GPO \$1.75.

A procedure for attribute sampling inspection is described. Tables are presented which indicate the minimum sample size and associated acceptance number for lot sizes from 1 to 5600 which satisfy specified combinations of risks. Tables are included for sixteen choices of risk combinations. Also presented are computer programs which, first, enable generation of tables for combinations of risks not included and second, provide a basis for investigating implications of varying sample sizes and acceptance numbers. (auth)

46095 URANIUM AND THE BALANCE OF POWER. World Miner. Metals; No. 7, 15-21(1972).

Power policy rather than politics may prove to be the biggest problem facing governments throughout the world as the 20th century reaches its closing years. This article examines the special case of uranium, at present the "Cinderella" of the energy industry, which seems destined very soon to meet its political and financial "Prince Charming." Among matters discussed are: availability of uranium ore and uranium enrichment progress. (auth)

46096 ATOMIC SAFEGUARDS: A STUDY IN INTERNATIONAL VERIFICATION. McKnight, Allan. UNITAR Series No. 5. New York; United Nations Institute for Training and Research (1971). 323p. \$6.50.

The safeguards system of the IAEA is examined in detail. Information is presented on the political development of the IAEA safeguards from 1946 to 1970 and the executive administration of the IAEA safeguards. A 42-item bibliography and a 9-item supplement containing: notes on international laws relevant to IAEA safeguards, on treaties and their verification, and on nuclear industry and commerce; statute of the IAEA; IAEA document INFCIRC/66/Rev. 2 on IAEA safeguards system; Treaty on the Non-Proliferation of Nuclear Weapons; Treaty for the Prohibition of Nuclear Weapons in Latin America; and the agreement between the IAEA and Mexico are included. (L.C.L.)

46097 FUEL ASSEMBLY IDENTIFICATION. ANSI N18.3-1972. Hinsdale, Ill.; American Nuclear Society (1972). 2p.

A standard prescribing requirements for fuel assembly identification (ANSI N 18.3-1972) for nuclear and associated applications is presented. (J.R.D.)

Disposal and Storage

Refer also to abstracts: 45565, 46090, and 46238.

46098 (AAEC/TM-628) MANAGEMENT OF WASTES FROM IRRADIATED NUCLEAR FUELS. Hespe, E. D.; Hardy, C. J. (Australian Atomic Energy Commission Research Establishment, Lucas Heights). 1972. 19p. (CONF-720627-1). Dep. NTIS (U. S. Sales Only).

From Symposium on the role of nuclear energy in Australia's development; Canberra, Australia (1 Jun 1972).

The fuel of present and future proposed power reactors has to be withdrawn from the reactor before all the fissile material is used, and it is economically desirable to recover this material. About 99.9% of the fission products present in the irradiated fuel is usually isolated in an aqueous nitrate solution in the reprocessing plant which recovers the unused uranium and the newly-created plutonium. It is this waste which, without proper management, could present a significant radiological hazard to this and future generations. Because of their abundance and relatively long half-lives, the fission product isotopes of most concern are cesium-137 and strontium-90. Small amounts of transuranic elements are also of concern because they have very long half-lives and high radiotoxicity. The amount of waste produced is proportional to the total nuclear power production and it will require surveillance for thousands of years. The wastes release up to 20 kilowatts of energy per ton of fuel reprocessed and hence may require cooling in the early years of storage. The main concern in waste management is the possibility that the wastes may become dispersed in the environment, be ingested by man, and thus expose him to internal radiation injury. All the major producers of these wastes use some form of storage in tanks as an interim measure. All have worked, or are working, on the development of processes to convert the wastes to insoluble high melting-point solids for final storage. The methods of ultimate disposal of the solids that have been developed so far, or have been proposed, are reviewed. Significant amounts of irradiated fuel are unlikely to arise from an Australian nuclear power program before the 1980s and the various factors which will have to be considered before a decision is taken to build a reprocessing plant are listed. An important question is where the long-term responsibility for waste management should lie, and this is discussed briefly. Responsible people in Western Europe have recently expressed confidence that the use of nuclear power need not be limited by waste management problems. Australia is fortunate in being able to observe and profit from the experiences of other countries, and it is concluded that the role of nuclear energy in Australia's development will not be limited by these problems. (auth)

46099 (ARH-SA-117) REMOTE WELDING PROGRAM FOR ATLANTIC RICHFIELD HANFORD COMPANY'S WASTE ENCAPSULATION AND STORAGE FACILITY. Wormell, John C. (Atlantic Richfield Hanford Co., Richland, Wash.). 4 May 1972. 22p. (CONF-720903-11). Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

Features of nondestructive testing capabilities incorporated into welding equipment for remote operations are described. Descriptions are included of the waste encapsulation and storage

facility, capsule materials, capsule fabrication for storage of ^{137}Cs and ^{90}Sr , and in-cell welding equipment and operation. (J.R.D.)

46100 (CONF-720903-4) BURIAL OF RADIOACTIVE WASTE IN SALT. Frederick, E. J. (Oak Ridge National Lab., Tenn.). 1972. 15p. Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

The use of natural salt formations for the long-term storage of radioactive waste has been actively studied at ORNL since 1957. Results of these studies, which included Project Salt Vault, a field-scale experiment, confirmed the feasibility of this concept and provided data for the design of an actual waste repository. A conceptual design of a repository for demonstrating storage of both solidified high-level fuel reprocessing wastes and alpha-contaminated solid wastes was completed in April 1971. The rationale of using salt deposits for storage of radioactive waste, siting considerations in locating a repository, the operating concepts developed during the conceptual design for the handling and burial of the solidified high-level wastes, and the long-term geological effects on the area surrounding a repository are discussed. (auth)

46101 (IAEA-136, pp 10-17) TECHNICAL STUDIES CONNECTED WITH THE RELEASE OF RADIOACTIVE WASTES INTO THE SEA. (International Atomic Energy Agency, Vienna (Austria)). 1971. INIS.

The Laboratory convened a Panel on Procedures for Establishing Limits for Radionuclides in the sea at which 15 papers were presented. A review of subjects related to marine radioactivity is under preparation. Investigations are under way on studies of non-radioactive pollutants by methods used in marine radioactivity studies, particularly in connection with mercury pollution; the horizontal diffusion of pollutants on the sea surface is being investigated in collaboration with the German Hydrographic Institute in Hamburg and the possible role of euphausiids in the biological transport of radionuclides is being investigated by the Laboratory. (auth)

46102 (ORNL-tr-2369) SOME PROBLEMS OF THE DISCHARGE OF RADIOACTIVE GASES AND AEROSOLS FROM NUCLEAR FACILITIES INTO THE ATMOSPHERE. Zindler, H. (Staatliche Zentrale fuer Strahlenschutz, Berlin (East Germany)). Translated for Oak Ridge National Lab., Tenn., from report SZS-17/69. 12p. Dep. NTIS.

An abstract of this paper, prepared from the original language, appeared as NSA 24: 25252.

46103 RADIOACTIVE WASTE STORAGE SYSTEM AND METHOD. Sayers, Robert P. (to FMC Corp.). U. S. Patent 3,663,817. 16 May 1972. Filed 28 Jul 1969.

A system and method for storing radioactive waste are described, wherein the waste is placed in receptacles and the receptacles are moved selectively along a plurality of storage paths. The receptacles are retained selectively on the paths for periods of time dependent upon the periods required for the radioactivity of the contained waste to decay to predetermined levels. (Official Gazette)

Handling Equipment and Procedures

Refer also to abstract: 47011.

46104 (LA-DC-72-556) CELLS FOR NONDESTRUCTIVE EXAMINATIONS OF IRRADIATED FUEL PINS. Fitzgibbon, F. J.; Montgomery, C. D.; Trujillo, J. R. (Los Alamos Scientific Lab., N. Mex.). 1972. 7p. (CONF-720903-3). Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

In October 1970 a decision was made to reactivate four LASL plutonium hot cells to conduct nondestructive postirradiation examination (PIE) of irradiated fuels sent to LASL at the request of the USAEC. These cells had been designed for research and development work on reprocessing irradiated plutonium reactor fuel and had been operational for a number of years. To make the cells suitable and efficient for the PIE of irradiated fuels, a number of modifications had to be considered and implemented to satisfy the planned examinations. The changes and additions made to the existing building are described; operational procedures are not discussed. (auth)

46105 (ORNL-tr-2371) WAK: SPECIFICATIONS AND FINISHING CONTROL FOR EQUIPMENT. Clauss, W.; Issel, W.; Warchold, L.; Meininghaus, H. Translated for Oak Ridge

National Lab., Tenn., from Atomwirt., Atomtech., 15: 84-9 (1970). 15p. Dep. NTIS.

All supplies were individually ordered by following the principle of "safety before economy." The suppliers were responsible for observing the specifications which were set in about 100 technical delivery and acceptance conditions (TDA). The documents submitted for examination were inspected primarily by Institute of Reactor Safety (IRS) and Technical Control Association (TUV) (Baden), while quality control was the responsibility of TUV of Baden and the local TUV. The limitation in the number of TDA's also leads to the highest quality conditions for less highly stressed components. (auth)

46106 (ORNL-tr-2372) SPECIAL MACHINES AND EQUIPMENT OF WAK. Geier, H.; Hoffmann, W.; Huppert, K. L.; Zimmermann, G.; Roth, B. F. Translated for Oak Ridge National Lab., Tenn., from Atomwirt., Atomtech., 15: No. 2, 89-94 (1970). 15p. Dep. NTIS.

The experimental nature of WAK justifies the remaining necessary development and testing of special equipment. Equipment which is specific for a process and which differs from conventional chemical engineering equipment is described. For the head-end process, this involves the disassembly machine on which the irradiated fuel assemblies are disassembled by sawing and grinding into the individual pins. Other paragraphs deal with the various extractors utilized as well as with a newly developed system for remote sampling from radioactive process solutions. (auth)

46107 (UCRL-73756) SHIELDED CAVES FOR DUAL-MODE OPERATION. Hanson, C. L.; Coops, M. S. (California Univ., Livermore, Lawrence Livermore Lab.). 8 May 1972. 10p. (CONF-720903-5). Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

A family of one-foot-thick water shielded facilities was constructed to permit rapid, convenient manipulations on intensely alpha active materials. Although nominally a manipulator operated containment enclosure, each shield contains the feature of allowing gloved operations in the enclosure on demand with minimal expenditure of time or effort. The basic shield encloses a replaceable alpha confinement box and provides shielding on four faces of the box, the top and rear not being shielded. Rear shielding is provided by locating the units against a structural concrete wall or heavy cell wall. (auth)

46108 (UCRL-73757) CONTAMINATION CONTROL HARDWARE FOR REMOTE HOT CELL TRANSFERS. Coops, M. S.; Hanson, C. L.; Voegele, A. L. (California Univ., Livermore, Lawrence Livermore Lab.). 8 May 1972. 11p. (CONF-720903-2). Dep. NTIS.

From twentieth remote systems technology conference; Idaho Falls, ID. (17 Sep 1972).

Remotely-operated plastic-bag sealers have been installed in the Livermore Neutron Cells to minimize the spread of contamination during transfers of highly alpha contaminated materials from the confinement boxes to the underfloor transfer system. An in-cell welding system has been added to one of the rear cell areas to seal active waste materials into steel canisters to aid in meeting "special form" specifications. The operation of these devices is controlled remotely in the semi-automatic mode to increase the reliability of the welded seals. (auth)

46109 DOUBLE ELECTROMAGNETIC VALVE FOR SIMULTANEOUSLY CONTROLLING DANGEROUS FLUIDS AND VACUUMS. Fremion, Lucien (to Commissariat a l'Energie Atomique). French Patent 2,088,094. 7 Jan 1972. Filed 15 May 1970. (In French).

An electromagnetic valve intended for applications in conjunction with dangerous fluids (corrosive or radioactive) is described. The valve possesses a magnetic part, which is attracted by an excited electromagnet and shuts-off the flow by an axial displacement. Two different circuits can be simultaneously controlled: a circuit for the fluid to be processed and a circuit containing the processing atmosphere. The displacement of the magnetic part also controls the opening of the two circuits. This opening can be carried by remote control, contamination risks thus being avoided. (France)

46110 SAMPLE-CONVEYING MECHANISM FOR SCINTILLATION COUNTING. Scherzer, Walter O.; Thomas, Edward W. (to Beckmann Instruments, Inc.). U. S. Patent 3,663,816. 16 May 1972. Filed 20 May 1966.

A conveying mechanism for transporting a sample vial from an external loading platform into a light-tight counting chamber of a scintillation detector is described. The invention comprises a wall defining a counting chamber having an elongated passage, the wall

having a shoulder. A sample platform is also incorporated having an attached shaft to position the platform through the passage and into the counting chamber, and a sleeve movable about the shaft for engaging the shoulder to prevent external light from entering the counting chamber. (Official Gazette)

46111 EQUIPMENT FOR CONCENTRATING RADIOACTIVE SOLUTIONS. Arnal, Thierry; Ganiwet, Michel; Somon, Roland (to Commissariat a l'Energie Atomique). French Patent 2,081,129. 3 Dec 1971. Filed 6 Mar 1970. (In French).

Equipment is described for concentrating (continuously) large volumes of radioactive solutions e.g., plutonium nitrate elution, oxalic mother liquor, and uraniferous effluents. A vertical axis cyclone is used to separate the liquid and vapor phases of the solution. The cyclone has an output for the liquid phase situated at its lower end, an output for the vapor phase situated at its upper end, and at least one input for the solution to be concentrated; this input consists of an admission tube tangentially joined to the wall of the cyclone. This tube is associated at the exterior to a heating and vaporization assembly. The equipment is also provided with a tube for evacuating the liquid phase, tubing for feeding the apparatus with the solution to be treated, this tubing being connected to the admission tube, a heat exchanger, and a device for estimating the position of the solution level in the cyclone. (France)

46112 PNEUMATIC MACHINE FOR THE REMOTE CONTROLLED SEALING OF FLASKS IN A SHIELDED CELL. Cassimatis, Jean; Vie, Robert (to Commissariat a l'Energie Atomique). French Patent 2,088,806. 13 Dec 1971. Filed 24 Apr 1970. (In French).

A pneumatic machine for the remote controlled sealing of flasks (in a shielded cell) using skirted supple plugs is described. The machine is provided with a receptacle for the flasks, which pivots about an axis and receives, at the appropriate moment, a flask with its plug. A hollow casing above the receptacle contains a series of pistons, fitted one into the other. A compressed air system, which is remotely controlled by a handle device, is used to advance these pistons towards the flask held in the receptacle or to recede them from it, and to ensure the sealing of the flasks. (France)

46113 HOLDER FOR RADIOACTIVE MATERIAL. Glasser, Herman (to Nuclear Associates, Inc.). U. S. Patent 3,673,411. 27 Jun 1972. Filed 3 Mar 1970.

A holder is described for radioactive material fabricated of high-density material to reduce transmission of energy, either by radiation or particle bombardment, either inwardly or outwardly thereof while yet enabling observation of material within an inner container or vial. (Official Gazette)

Transportation

46114 (MLM-1911) DOCUMENTATION OF DOT-SP-5915 FOR 55/30 AND 83/55 DRUM COMBINATION PACKAGING. Madding, R. D. Jr.; Vallee, R. E.; Griffin, J. F. (Mound Lab., Miamisburg, Ohio). 28 Apr 1972. Contract AT-33-1-GEN-53. 45p. Dep. NTIS.

A report is presented to make available to all potential users the technical information and limits pertinent to the construction and use of the 55/30 and 83/55 drum combination packages for which the Department of Transportation has issued special permit

No. 5915. A complete physical and technical description of the packages is presented. The authorized packaging consists of either of two sizes of metal, drum-type "bird cage" packaging, for single trip use only. The Model 55/30 consists of a 55-gal outer drum and a 30-gal inner drum, and the Model 83/55 consists of an 83-gal outer drum and a 55-gal inner drum. For both models, the annular space between the drums is filled with moistened vermiculite. A nuclear criticality safety analysis is included. This analysis showed that for Fissile Class I packages, uranium-235, uranium-233, and plutonium-239 must not exceed 200, 190, and 175 g, respectively for container Model 83/55, and 180, 150, and 138 g, respectively, for container Model 55/30. Design and development considerations required for proof of the containers' ability to withstand a sequence of four hypothetical accident conditions are discussed. The four tests conducted in sequence on the same container were free drop, puncture, thermal, and water immersion tests. The tests showed that dampened vermiculite provided good insulation around the inner containers. It was calculated that no corrosion inhibitor was required in the damp vermiculite unless the containers were to be stored for long periods of time. Sponge rubber seals performed better than other gasket materials tested for the drums. In addition to the development data required for special permit No. 5915, data are presented pertaining to designs for disposable containers for radioactive sludge, for solid plutonium-238 compounds, and for radioactive gas. Test results on a design modification to the reusable DOT Specification 6M container are also reported. Tables, graphs, drawings, pictures, technical references, and a copy of permit DOT-SP-5915 are included. (auth)

46115 PACKAGING FOR RADIOACTIVE MATERIALS. Blum, Paul (to Transnucleaire Societe pour les Transportes de l'Industrie Nucleaire). French Patent 2,088,028. 7 Jan 1972. Filed 8 May 1970. (In French).

A packaging for radioactive materials emitting neutrons and producing heat is described. The packaging is characterized by the fact that good conducting metallic elements capable of absorbing neutrons and having a U or I shaped section, are buried in a bonding material provided in the space between the inside and the outer shell of the packaging. (France)

46116 SHIPPING CONTAINER FOR FISSIONABLE MATERIAL (25 May 1972) (Engineering Materials). (Dow Chemical Co., Denver, Colo. Rocky Flats Div.). (CAPE-1657). 4 drawings.

This container may be used to ship up to 5 kg of ^{233}U or ^{239}Pu or 11 kg of ^{235}U . The container consists of a 30 gallon drum with vented lid, fiberboard insulation and a light-gage stainless steel containment vessel. (For details see TID-4100 Suppl. 72) (F.S.)

46117 TRANSPORT OF RADIOACTIVE MATERIALS BY RAIL. Schaefer, H. (Bundesbahn-Zentralamt, Minden, Ger.). Eisenbahn-Ing.; 22: No. 5, 141-4(1971). (In German).

The regulations for the transport of radioactive materials are the same for transport by rail or road; they are based on the "Regulations for the Safe Transport of Radioactive Materials" (IAEA 1967). Their aim is to create a framework of generally accepted regulations and instructions for such transports in order to ensure that these transports differ as little as possible from that of other goods. The principle was to achieve this by the type of packing. (INIS)

PARTICLE ACCELERATORS

Design, Development, and Operation

Refer also to abstracts: 45432, 46146, and 46873.

46118 (KFKI-71-47) NEW 5 MeV VAN DE GRAAFF ION ACCELERATOR AT THE CENTRAL RESEARCH INSTITUTE FOR PHYSICS OF THE HUNGARIAN ACADEMY OF SCIENCES, BUDAPEST. Pasztor, E.; Kostka, P.; Klopfer, E.; Berecz, Gy.; Buerger, G.; Gombos, P.; Horvath, B.; Kiralyhidi, L.; Riedl, P. (Magyar Tudományos Akadémia Kozponti Fizikai Kutató Intézete, Budapest), 1972, 122p. Dep. NTIS (U. S. Sales Only).

The new 5 MeV Van de Graaff ion accelerator designed and built at the Central Research Institute for Physics of the Hungarian Academy of Sciences and started in normal operation on March 12, 1970, is described. In its present state, the equipment is suitable for accelerating protons, deuterons and He^+ ions. Analyzed ion beams are of max $10 \mu\text{A}$ intensity and the stability of energy of ions is 1.5 to $2.5 \cdot 10^{-4}$ at the target. Design is also described, main technical data are given, and special problems of the high-pressure gas insulation and the charging system are dealt with in detail. Original measuring results are given on the harmful effects of humidity and dust contents. The mechanical and electrical designs of the ion source and its auxiliary equipment are dealt with, technical data listed, and the system engineering is described. Technological and ion-optical development of the accelerator tube is outlined from a conventional homogeneous-field porcelain tube to a PVA-cemented inclined-field glass accelerator tube. Details of the ion-optical design, as well as conditioning and initial operation of the completed tube, are also discussed. Design of the complex energy stabilization system, the NMR fluxmeter developed in the Institute, and the impulse width modulated stabilization system of the magnetic field are also treated. Finally, experiences gained during 6,253 hours of operation completed so far are summed up. (auth)

46119 (LA-5000) ORIGINS AND HISTORY OF THE LOS ALAMOS MESON PHYSICS FACILITY. Livingston, M. Stanley (Los Alamos Scientific Lab., N. Mex.). Jun 1972. Contract W-7405-eng-36. 68p. Dep. NTIS.

A history of the Los Alamos Meson Physics Facility (LAMPF) from initiation of preliminary plans and proposals in 1962 to the present is presented. It includes the year-by-year story of the actions of the Atomic Energy Commission, the United States Congress, and the Bureau of the Budget relative to funding the project. It also discusses the discovery, subsequent interest, and proposed applications of the pi-meson. (auth)

46120 ELECTRON ACCELERATION IN AN ONDULATOR SYNCHROTRON. Ziyakaev, R. G.; Fomenko, G. P.; Chuchalin, I. P.; Yushkov, Yu. G. (Tomsk Polytechnic Inst., USSR). JETP Lett. (USSR) (Engl. Transl.); 15: No. 6, 212-13 (20 Mar 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 6, 301-2 (20 Mar 1972).

The 2-MeV electron undulator synchrotron at Tomsk Polytechnic Institute is briefly described. The electrons are accelerated in a 10-cm-band resonator in which the H_{014} mode is excited. In each passage through the resonator, the electrons increase their energy by 350 keV at the maximum of the electric field and are then returned to the resonator at the next maximum of the electric field with the aid of electron-mirror turning magnets installed in line with the resonator. The output beam current is 50 mA, and the electron energy spread is 6%. (W.H.K.)

46121 DOSE ESTIMATES OUTSIDE THE SHIELDING OF THE CERN INTERSECTING STORAGE RINGS BY THE MOYER METHOD. Routti, Jorma T.; Van de Voorde, M. H. (CERN, Geneva). Nucl. Eng. Des.; 21: No. 3, 421-34 (1972).

Formulas and results are presented for a convenient estimation of dose levels outside the transverse shielding of a high-energy proton accelerator. They are obtained by applying the Moyer method to uniform, local, and exponential beam-loss distributions. The techniques are used in shielding considerations of the CERN Intersecting Storage Rings. The layout of the main shielding is described, and estimates of the beam losses during various machine operations are summarized and used to evaluate doses outside the main shield. (8 figures, 2 tables) (auth)

46122 Q-VALUE EFFECTS OF ACCELERATING STRUCTURE IN TRAVELING-WAVE LINACS. Irie, Koichi (Mitsubishi Electric Corp., Amagasaki, Japan). Jap. J. Appl. Phys.; 11: No. 7, 1056 (Jul 1972).

Quantitative evaluation of the Q-value effect on accelerator performance is important as it affects the cost of construction. The results of such effects on the electron energy of a traveling-wave linac are given. It is found that the Q-value of the accelerating structure has little effect on electron energy. What effect there is is less in the constant-gradient machine than the constant-impedance one. The smaller the total attenuation parameter, the smaller the effect. (1 figure) (R.W.R.)

46123 COMPUTERIZED CONTROL OF AN ACCELERATOR AND THE ASSOCIATED EXPERIMENTS. Bluet, J. C.; Fardeau, P.; Arnaud, A. (CEN de Cadarache, Saint-Paul-les-Durance, France). Bull. Inform. Sci. Tech. (Paris); No. 168, 88-90 (Mar 1972). (In French).

The CII 90-10 computer associated with the Van de Graaff accelerator at Cadarache is described. This computer is programed in order to ensure the monitoring and possible energy adjustments of the accelerator. (France)

46124 ACCELERATION OF ARGON IONS TO 1.17×10^{10} ELECTRON VOLTS. Isaila, M. V. (Princeton Particle Accelerator, N. J.); Schimmerling, W.; Vosburgh, K. G.; (and others). Science; 177: No. 4047, 424-5 (4 Aug 1972).

Argon ions were accelerated to 1.17×10^{10} electron volts in the Princeton Particle Accelerator. The synchrotron was tuned by use of a neon beam with a charge-to-mass ratio equal to that of the argon ions. The fully accelerated argon ions were detected by the observation of etched tracks in cellulose nitrate sheets and also by the use of scintillation counters. Predictions of the range and of the characteristics of argon tracks in plastics were confirmed. (auth)

46125 DIELECTRIC STRENGTH OF ACCELERATOR TUBES. Almazov, A. V.; Myntsov, F. F. Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1619-22 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 32-5 (1971).

The results of investigations of the influence of various factors on the dielectric strength of elements of accelerator tubes are expounded. It is shown that the greatest influence is produced by the quality of the contact between the dielectric and the electrodes. For a good contact the minimal strength of the elements in a vacuum for an element length of 25 mm is 44 kV/cm. (auth)

Auxiliaries and Components

46126 (ORNL-tr-2617) INTRANUCLEAR CASCADE IN THE CONCRETE SHIELD OF A 70-GeV PROTON SYNCHROTRON. Britvich, G. I.; Getmanov, V. B.; Lebedev, V. N.; Mal'kov, V. V.; Sychev, B. S. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoi Energii SSSR, Serpukhov. Institut Fiziki Vysokikh Energii). Translated for Oak Ridge National Lab., Tenn., from report IFVE-ORZ-71-81. 11p. Dep. NTIS.

The transport of the high energy component in a concrete shield in the neutral particle channel of the Serpukhov Synchrotron was studied. The internal target of the accelerator bombarded by protons of 70 GeV served as the radiation source using steel and concrete collimators. The neutral particles were magnetically separated from the charged particle background and were incident on the concrete block in which the hadron flux attenuation was studied. The density of the concrete was 2.3 to 2.35 g/cm³. The beam profile at the point of incidence on the concrete block, the flux density distribution in the beam plane, and the flux density distribution in planes perpendicular to the beam axis are shown graphically. Also, the experimental data on the integrated transverse flux density distribution are compared with the calculated results of the internuclear cascade in iron. (W.H.K.)

46127 AUTOMATIC FREQUENCY CONTROL AGAINST TEMPERATURE CHANGE IN LINAC'S ACCELERATING STRUCTURE. Irie, Koichi (Mitsubishi Electric Corp., Amagasaki, Japan). Jap. J. Appl. Phys.; 11: No. 7, 1057 (Jul 1972).

The phase slip resulting from temperature changes in the accelerating structure of an electron linac causes a reduction in energy gain. This is ordinarily prevented by two independent control systems, one regulating temperature and the other regulating microwave frequency. However, it is shown that regulation is sufficient if the microwave frequency follows the change of the resonant frequency of the structure due to temperature change, and a method to accomplish this type of control is proposed. (3 figures) (R.W.R.)

46128 COMPENSATED CURRENT-DRIVEN MAGNETIC CHANNEL IN A 240-cm ISOCHRONOUS CYCLOTRON. Belyakov, V. A.; Litunovskii, R. N.; Minyaev, O. A. (Electrophysical Equipment Research Inst., Leningrad). *Sov. Phys.-Tech. Phys.* (Engl. Transl.); 17: No. 1, 154-6 (Jul 1972).

Translated from *Zh. Tekh. Fiz.*; 42: No. 1, 194-7 (Jan 1972).

The basic theoretical and experimental characteristics of a compensated current-driven channel in the beam extraction system in the 240-cm isochronous cyclotron of the Institute of Nuclear Research, Academy of Sciences of the Ukrainian SSR (IYaI AN UkrSSR) are reported. The channel lowers the magnetic field to 2200 Oe, with a positive radial gradient to 150 Oe/cm, at roughly 80-kW power input. The amplitude of the first harmonic of the field disturbance introduced by the channel in the beam acceleration region is less than 10 Oe. (3 figures) (auth)

46129 MAGNETIC SHIELD FOR A 240-cm ISOCHRONOUS CYCLOTRON. Belyakov, V. A.; Vasil'ev, V. I.; Litunovskii, R. N.; Minyaev, O. A. (Electrophysical Equipment Research Inst., Leningrad). *Sov. Phys.-Tech. Phys.* (Engl. Transl.); 17: No. 1, 157-9 (Jul 1972).

Translated from *Zh. Tekh. Fiz.*; 42: No. 1, 197-9 (Jan 1972).

The basic experimental characteristics of the passive magnetic shield of the beam extraction system of the 240-cm isochronous cyclotron at the Nuclear Research Institute, Academy of Sciences of the Ukrainian SSR (IYaI AN UkrSSR) are given. The shielding factor in the working range of the externally applied magnetic field, 5000 to 7000 Oe, varies from 0.03 to 0.5. A positive radial field gradient of 350 Oe/cm is established by means of the cutoff angles on the channel ends and the variable internal aperture. The perturbations in the magnetic field in the beam acceleration zone due to this field are compensated to the level of several oersteds (first harmonic) with the aid of two pairs of iron shims. (3 figures, 2 tables) (auth)

46130 DIELECTRIC STRENGTH OF INSULATORS IN A VACUUM. Solodovnikov, A. P. *Instrum. Exp. Tech.* (USSR) (Engl. Transl.); 14: No. 4, 1156-7 (Jul-Aug 1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 4, 176-7 (Jul-Aug 1971).

The joining of insulators to electrodes and the choice of the method of treating their lateral surface are considered; application is made to the construction of accelerator tubes for electrostatic generators. (1 figure) (auth)

46131 CONTROL SYSTEM FOR BEAM FORMATION AFTER SINGLE DEFLECTION AT THE EXIT OF THE 300 MeV ELECTRON LINEAR ACCELERATOR. Egiyan, K. Sh.; Bochek, G. L.; Mocheshnikov, N. I.; Alanakyan, K. V.; Sitenko, M. L. (Inst. of Physics, Erevan. Physical-Technical Inst., Kharkov). *Izv. Akad. Nauk Arm. SSR, Ser. Fiz.*; 6: No. 3, 214-18 (1971). (In Russian).

The control system designed for beam formation after single deflection in the 300-MeV Kharkov linear accelerator is described. The principle of operation is based on a comparison of the intensities of the beam at two points (at the exit of the accelerator and immediately before the target of the experimental arrangement. (5 figures) (auth)

46132 IMPROVEMENT OF THE ELECTRON BEAM SHAPER OF THE B5M-25 MODEL BETATRON. Kozlov, A. P.; Shishov, V. A. (Leningrad Oncology Inst.). *Med. Radiol.*; 17: No. 5, 70-5 (May 1972). (In Russian).

An improved design of the B5M-25 betatron electron beam shaper is proposed. It enables creating uniform dose fields measuring $18 \times 18 \text{ cm}^2$ with a variation factor not in excess of 10%. The use of the improved shaper enables the skin radiation load to be cut down by 15 to 20% by comparison with the former version, and also makes it possible to reduce appreciably the background radiation outside the radiation field. (5 figures) (auth)

Ion Sources, Injection, and Extraction

46133 PRODUCTION AND ACCELERATION OF MULTI-CHARGED IONS OF PHOSPHORUS AND ZINC. Pasyuk, A. S.; Tret'yakov, Yu. P.; Zager, B. A. *At. Energ. (USSR)*; 32: No. 6, 505-8 (Jun 1972). (In Russian).

Experiments were conducted for the purpose of developing ion sources in the solid phase for use in a cyclotron. A stainless steel crucible loaded with zinc or red phosphorus was protected by a heat shield during operation as an ion source. The technical grade of zinc yielded Zn^{10+} , Zn^{2+} , and other ions, and the Zn^{10+} ion current was investigated as a function of the pressure in the vacuum chamber. Zinc consumption was 0.035 g/hr. A charge spectrum of phosphorus ions showed the presence of P^+ , P^{2+} , P^{3+} , P^{4+} , P^{5+} , and P^{6+} , in addition to oxygen ions. Phosphorus consumption was 0.13 g/hr. (3 figures) (K.S.W.)

46134 MULTIPACTORING ELECTRON GUN FOR HIGH DUTY LINACS. Liska, Donald J. (Los Alamos Scientific Lab., N. Mex.). *Proc. IEEE (Inst. Elec. Electron. Eng.)*; 59: No. 8, 1253-4 (Aug 1971).

A high duty factor electron gun suitable for linac applications is described. This gun operates on the secondary emission principle to generate a pulsed, monochromatic electron output beam in excess of 20 mA average current. It has been operated at 14 percent duty factor and shows a high degree of stability and ease of control. (auth)

46135 SEPARATE CONTROL OF THE INJECTOR HEATING AND OF THE CURRENT AMPLITUDE IN TRANSFORMER ACCELERATORS. Goncharov, N. F.; Egorov, A. A. (Inst. of Nuclear Physics, Novosibirsk, USSR). *Instrum. Exp. Tech. (USSR)* (Engl. Transl.); 14: No. 6, 1623-4 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 36-7 (1971).

An automatic pulse network is described for separate control of the operating mode of two loads which are fed from a single source to a single transmission line. The loads differ in character: the operating modes of one of them is determined by the rms value of the source voltage; the operating mode of the other is determined by the amplitude value. A circuit is given for comparing the rms value with a reference voltage. The network described is used for separate control of the heater and the modulating-pulse amplitude for the injector in the ELIT transformer accelerator. (auth)

Ion Optics and Field Calculations

46136 (JINR-P9-6421) INFLUENCE OF THE ELECTRO-MAGNETIC FING ACCELERATION ON THE FOCUSING BY IMAGE FORCES. Zhabitskii, V. M.; Ivanov, I. N. (Joint Inst. for Nuclear Research, Dubna (USSR)). 1972. 13p. (In Russian). Dep. NTIS (U. S. Sales Only).

The focusing of the accelerated electron ring by the image forces in a perfectly conducting cylindrical screen is considered. The corrections to the focusing force induced by the acceleration are found, and their influence is analyzed. (auth)

46137 (JINR-P9-6514) STABILITY OF THE SCREENED RING AT THE CENTER SHIFT. Zhabitskii, V. M.; Zolina, I. A.; Kaminskaya, A. M.; Ivanov, I. N.; Perel'shtein, E. A. (Joint Inst. for Nuclear Research, Dubna (USSR)). 1972. 18p. (In Russian). Dep. NTIS (U. S. Sales Only).

The problem is considered on the electron ring stability when its center is shifted with respect to the axis of the screening tube. The particle motion in the shifted ring is analyzed. The limitations are found for the admissible deviations of the ring center depending on the system parameters. (auth)

46138 LINEAR STABILITY OF A CHARGED BEAM IN AN INDUCTION SYSTEM. Grishin, V. K. *Sov. Phys.-Tech. Phys.* (Engl. Transl.); 17: No. 1, 6-8 (Jul 1972).

Translated from *Zh. Tekh. Fiz.*; 42: No. 1, 9-12 (Jan 1972).

The kinetic equation is used to study the longitudinal stability of a charged beam in a ferromagnetic channel. The growth rates of instabilities in high-current devices are estimated for the example of a linear induction accelerator. (auth)

46139 PARTICLE ACCELERATION IN A H_{111} CYLINDRICAL CAVITY IN AN AXIAL MAGNETIC FIELD. Gryzlov, A. V.; Panasyuk, V. S.; Ryzhkov, V. M.; Sokolov, A. A.; Tereshkin, Yu. M. *Sov. Phys.-Tech. Phys.* (Engl. Transl.); 17: No. 1, 9-13 (Jul 1972).

Translated from *Zh. Tekh. Fiz.*; 42: No. 1, 13-18 (Jan 1972).

Certain aspects of particle acceleration in the electric field of the H_{111} mode in a cylindrical cavity are discussed for the case of a pulsed magnetic drive field. The "resonant" time dependence is found for increasing magnetic field. The limiting energy is determined for particles accelerated by a homogeneous high-frequency electric field in a constant, homogeneous magnetic field. The maximum rate of rise of a linearly changing magnetic drive field is found. Acceleration rates are found for a linearly increasing magnetic field and for the "resonantly" increasing magnetic field. (3 figures) (auth)

46140 APPROXIMATIONS OF THE SPACE CHARGE FIELD FOR DIFFERENT BEAM SHAPES IN THE CENTRAL REGION OF A SYNCHROCYCLOTRON. Besnier, Gilbert; Thouroude, Daniel (Univ., Rennes, France). *Compt. Rend., Ser. B*; 274: No. 15, 927-30 (10 Apr 1972). (In French).

The space charge field was calculated for two simple forms of the injected beam. It is assumed that the beam breaks away from the source in a charge sector (cyclotron phase) which generates a charge column in the azimuthal direction. (France)

46141 ORTHOGONALITY RELATIONS IN PERIODIC WAVEGUIDES HAVING ANISOTROPIC SURFACE IMPEDENCE. Garault, Yves; Jecko, Bernard (Univ., Limoges, France). *Compt. Rend., Ser. B*; 274: No. 12, 769-72 (20 Mar 1972). (In French).

Iris-loaded rectangular periodic structures and circular periodic structures such as iris-loaded waveguides and shielded-ring lines are studied. The simultaneous excitation of two hybrid modes in one of these guides is considered. The waveguide is considered to be smooth with a uniform anisotropic surface joining the ends of the iris or rings. Calculations are simplified, only the fundamental mode of the electromagnetic field excited (in the coaxial region in the case of a ring line) is considered. The dependence of the field on z is demonstrated. The corresponding function is integrated over the cross section of the guide (of circular or rectangular cross section). This integral is demonstrated to be zero for all structures. (France)

46142 INSTABILITY OF AXIAL OSCILLATIONS IN AN ELECTRON-RING ACCELERATOR. Zenkevich, P. R.; Koshkarev, D. G.; Perel'shtein, E. A. *At. Energ. (USSR)*; 33: No. 1, 567-71 (Jul 1972). (In Russian).

The oscillations of an electron-ring column result in additional forces acting on the local center of gravity of the electrons owing to nonstationary charges and currents. Taking into account these forces, the authors consider the stability of an electron and electron-ion ring relative to lateral wiggles directed parallel to the ring axis. It is shown that in an electron ring there may arise a dipole axial instability related to the quasi-static and radiation fields of the electrons. Expressions are obtained for the increment and threshold of this instability in empty space and in a waveguide having longitudinal cuts. There is investigated the behavior of an electron-ion ring with the interaction forces between electrons and ions as well as the forces acting between electrons taken into account. It turns out that for a proper choice of ring parameters the presence of ions decreases the dipole instability increment of an electron-ring. (1 figure) (auth)

Storage Rings

Refer also to abstracts: 46121, 46426, and 46546.

46143 CERN PROTON STORAGE RING. Keil, E. (CERN, Geneva). *Naturwissenschaften*; 59: No. 5, 197-204 (May 1972). (In German).

The interest in proton-proton colliding-beam storage rings is due to the fact that they allow the observation of the strong proton-proton interaction at much higher center-of-mass energies than conventional accelerators, and at similar counting rates. The CERN Intersecting Storage Rings (ISR) are described with particular emphasis given to the differences between storage rings and accelerators. The results of the first few months of commissioning are discussed in detail. Some space charge phenomena which limited the ISR performance have been eliminated; some others are still present. 40% of the design current, and 5% of the design luminosity have already been reached. (auth)

Experimental Facilities

Refer also to abstracts: 45638 and 46848.

46144 (CONF-720607-22) 150-METER TIME-OF-FLIGHT FACILITY AT ORELA. Silver, E. G.; Ingle, R. W.; de Saussure, G.; Perez, R. B.; Weaver, H.; Robson, R. J. (Oak Ridge National Lab., Tenn.). 1972. 4p. Dep. NTIS.

From eighteenth annual American Nuclear Society conference; Las Vegas, NV. (18 Jun 1972).

The modification of the ORELA neutron time-of-flight facility by increasing the flight path from 40 to 150 meters is described. The collimators added to the system are briefly described, and advantages of the increased flight path are discussed. The system employs an 800-gallon liquid scintillator gamma detector. (W.H.K.)

46145 DECREASE OF THE OFF-DUTY FACTOR IN THE OPERATION OF A LINEAR ACCELERATOR. Adeishvili, D. I.; Grishaev, I. A.; Mocheshnikov, N. I.; Tolstoi, A. E. *At. Energ. (USSR)*; 33: No. 1, 593-4 (Jul 1972). (In Russian).

The conversion of the pulsed electron beam of a linear electron accelerator into a continuous beam of photons by utilizing an internal target in a storage ring is described. The intensity and duration of the photon beam bremsstrahlung are capable of being varied within wide limits; and for target distances of 4, 8, and 16 mm from the equilibrium orbit the bremsstrahlung intensity was measured as a function of the relative spacing of the circulating beam and the target. (3 figures) (K.S.W.)

46146 CYCLOTRON LABORATORY AT THE TECHNICAL UNIVERSITY OF EINDHOVEN. Poppema, O. J. (Technische Hogeschool, Eindhoven, Netherlands); Verhaar, B. J.; Hagedoorn, H. L.; Bruininx, E. *Ned. Tijdschr. Natuurk.*; 37: 55-63 (18 Feb 1971). (In Dutch).

A survey is given of the experimental and theoretical work at the Cyclotron laboratory of the Technical University of Eindhoven. Nuclear reactions are studied with beams of polarized protons and deuterons. Applied work is performed on elemental analysis by means of nuclear reactions, on production of radioisotopes and on the use of short-living nuclides in the study of gas discharge. A few remarks are made regarding the automatic regulation of the cyclotron with a POP-9 computer. (Netherlands)

46147 LINE-FOCUS ELECTRON BEAM LENS. Kalibjian, R. (Univ. of California, Livermore). *Proc. IEEE (Inst. Elec. Electron. Eng.)*; 60: No. 2, 244 (Feb 1972).

An electrostatic lens system has been developed to transform the circular beam from a conventional electron gun into a line-focus beam. The line-focus beam is uniform throughout its projected length on a phosphor screen. (auth)

46148 BETATRON BREMSSTRAHLUNG FOR PHYSICAL RESEARCH. Baci, G. (Inst. of Atomic Physics, Bucharest). *Stud. Cercet. Fiz.*; 24: No. 1, 65-79 (1972). (In Rumanian).

Nuclear research at the Bucharest betatron using bremsstrahlung beams is outlined. Topics include photonuclear reactions, the giant resonance, the dynamic collective nuclear model, photo-fission experiments, safeguards technology, nondestructive nuclear materials analysis, and other applications of the betatron in nuclear technology. (W.H.K.)

46149 INVESTIGATION OF POSITRON SPECTRA OBTAINED ON A LINEAR 8 TO 30 MeV ELECTRONIC ACCELERATOR. Tarutin, I. G.; Pilyavets, V. I. (Scientific-Research Inst. of Oncology and Medical Radiology, Minsk). *Instrum. Exp. Tech. (USSR) (Engl. Transl.)*; 14: No. 6, 1625-8 (1971).

Translated from *Prib. Tekh. Eksp.*; 14: No. 6, 38-40 (1971).

The results are presented of investigations of the energy distribution of positrons which are formed in Ta conversion targets of 0.2, 0.5, and 0.8 radiation length units (X_0) thick. The dependence of the conversion coefficient at the maximum of the positron energy distribution on the energy of the incident electrons in the 12 to 30 MeV range were determined. It was shown that fluxes of positrons obtained on an LUE-25 linear electron accelerator amount to $\sim 10^7$ particles/sec. The optimal values of target thickness lie in the 0.7 to 0.9 X_0 range when the electrons incident on the converter have an energy of 25 to 30 MeV. A semi-empirical formula is derived for determining the yield of positrons having a stipulated energy as a function of the target thickness and the energy of the incident electrons. (auth)

PHYSICS (ASTROPHYSICS AND COSMOLOGY)

Cosmic Radiation

Refer also to abstracts: 45215, 46219, 46257, and 46461.

46150 (NYO-3038-13) DETECTION OF LARGE COSMIC-RAY SHOWERS BY ATMOSPHERIC FLUORESCENCE. Final Report. Greisen, Kenneth (Cornell Univ., Ithaca, N. Y. Lab. of Nuclear Studies). 18 May 1972. Contract AT(30-1)-3038. 25p. Dep. NTIS.

Cosmic particles with energy on the order of 10^{20} eV arrive at the Earth with extremely low frequency, about one per square mile per decade. In 1962, a new experimental method was proposed to try to detect such particles with greater frequency by viewing them from afar (up to 20 miles from their path), using the fluorescent light emitted by some 10^{15} excited molecules along the path of such a particle and its progeny in the air. The proposed scheme also was expected to give new types of data by permitting each shower to be viewed throughout its history. In addition, its energy was to be directly measurable from the total number of molecules excited in the atmosphere. The direction of each shower relative to the stars would be revealed by the coordinates of the fluorescent light registered as a function of time. Initially, on the basis of previous reports on the fluorescent efficiency of air, the project looked easy; but when the spectrum and absolute yield of optical photons under fast particle bombardment of air were accurately measured it was found that the true yield was much smaller than anticipated. Still the project looked possible, though more difficult. In the years since 1963, several increasingly complex versions of optical detection systems have been set up on remote hilltop sites in the Ithaca area. With painstaking effort, they have been operated on many clear, moonless nights—while on many more nights the weather prevented operation. Marginal indications of possible showers were received, but no events stood out clearly enough from the background of starlight to permit unambiguous analysis as air showers. Each period of fruitless measurement was followed by study of weaknesses of the detection system and attempts to improve it. At one point a major change was initiated in which a detecting station was designed in the form of a "bug's-eye"; that is, hundreds of individual phototubes looking through large lenses at different parts of the sky. In addition to these efforts to improve the detector, other studies were made of the atmosphere. Among other minor projects, a way was developed of simulating showers by projecting a beamed pulse of light into the sky. The scattered light from the head of this beam is geometrically and kinematically like the source of light in an air shower, except for travelling upwards in time instead of downwards. Despite all these efforts, success was not achieved in development of a fruitful shower detector. In January 1972, the project was closed down. (auth)

46151 CHARGE COMPOSITION AND ENERGY SPECTRUM OF VERY HEAVY COSMIC RAYS. Fernandez, Joseph Ignatius. St. Louis; Washington Univ. (1971). 145p. University Microfilms Order No. 72-17,95. Thesis.

A large area [0.92 m² steradian] electronic detector system for very heavy primary cosmic rays was flown on a high-altitude [~4.5 mg residual atmospheric pressure] balloon, from Sioux Falls, South Dakota in September 1970. The main elements of the detector were two pulse type gas filled ionization chambers and a Lucite Cherenkov counter. Measurements were made on the charge composition of VH [20 ≤ Z ≤ 30] and VVH [Z ≥ 30] cosmic rays above 600 MeV/n and the energy spectrum of VH cosmic rays from 600 MeV/n to 1.28 GeV/n. In-flight calibration confirmed iron as the most abundant element in this region of the charge spectrum. (TSS) (Diss. Abstr. Int., B)

46152 SOLAR MODULATION OF GALACTIC PARTICLES. NUMERICAL INTEGRATION OF THE STEADY FOKKER-PLANCK EQUATION. Martinic, Nicolas J. Abstracts of Uppsala Dissertations from the Faculty of Science, No. 4. Uppsala; University of Uppsala (1972). 46p. Thesis.

Galactic particles are modulated by the solar atmosphere showing, as a general feature, a decrease in the flux in $-v$. From numerical integration of the steady, spherically symmetric Fokker-Planck equation, which governs this phenomenon as a first approximation, it is shown that the galactic flux measured at the Earth presents the effect of: a) convection; b) diffusion; and c) adiabatic deceleration. The effect of convection is always a positive modulation at all kinetic energies (i.e. decrease of density of galactic particles vis-à-vis the galactic density), while the other type

of modulation can be positive or negative depending on the particle's energies; pointing out that the integrated modulation due to adiabatic deceleration remains zero when all energies are taken into account. The density of galactic particles for a wide interval of energies (10^3 BeV to 10^{-3} BeV) is presented for all values of the radial distance in interplanetary space, whose range was taken 10 and/or 25 AU and for rigidity, kinetic energy, and total energy power laws, for protons; and the Webber law for electrons in the same range of energies. It is concluded that the proton spectrum presented in 1965 (Gloeckler and Jokipii, 1967) agrees fairly well with that calculated here when the galactic spectrum is assumed to be of the form E^{-7} , where E is the total energy of the protons. The electron spectrum presented in 1965–68 (Lezniak and Webber, 1971) shows the same topology as that calculated here. The effect of nongalactic sources which can be taken into the formalism is discussed. The numerical analysis, which is discussed in detail, follows the implicit scheme of the finite differences of the Fokker-Planck equation. (auth)

46153 ANGULAR DISTRIBUTION AND ENERGY SPECTRUM OF COSMIC-RAY SINGLE NEUTRONS WITH ENERGIES HIGHER THAN $5 \cdot 10^{11}$ eV AT MOUNTAIN ALTITUDES. Babayan, Kh. P.; Boyadzhyan, N. G.; Mamidzhanyan, E. A. (Inst. of Physics, Erevan). Izv. Akad. Nauk Arm. SSR, Ser. Fiz.; 6: No. 3, 168–73 (1971). (In Russian).

The results of the study of single neutrons with energies higher than 5×10^{11} eV at mountain altitudes by means of an ionization calorimeter supplemented with proportional counters are given. The angular distribution and the integral energy spectrum of single neutrons were obtained. (4 figures, 1 table) (auth)

46154 GALACTIC GAMMA RADIATION BETWEEN 200 MeV AND 10 GeV. Bennett, K.; Penengo, P.; Rochester, G. K.; Sanderson, T. R.; Sood, R. K. (Imperial Coll. of Science and Tech., London). Nature (London); 238: No. 5358, 31–3 (7 Jul 1972).

It is stated that it has not yet been established whether the γ radiation from the Galaxy at energies >100 MeV is a diffuse line flux from the galactic disk or arises from a number of weak point sources, and the production mechanism is poorly understood. In principle, the uncertainty about the production mechanism could be resolved by examining the energy spectrum of the γ rays with high-angular-resolution detectors, and a study is described of the high-energy part of the spectrum between 200 MeV and 10 GeV, using balloon-borne detectors together with a spark chamber. Details are given of the techniques employed. The experiment was flown from Lusaka in Zambia on Jan. 25 and Feb. 9, 1971. The sky map obtained was scanned for possible point sources, and only one position suggested evidence for a source; there was, however, an apparent line flux from the galactic disk above the expected count rate. The data obtained provided evidence for the existence of a galactic line source of high-energy γ rays in the region of the galactic center. (UK)

46155 ³⁷Ar/³⁹Ar ACTIVITY RATIOS IN METEORITES AND THE SPATIAL CONSTANCY OF THE COSMIC RADIATION. Begemann, F. (Max-Planck-Institut fuer Chemie, Mainz). J. Geophys. Res.; 77: No. 20, 3650–9 (10 Jul 1972).

Argon 37 and argon 39 were measured in a bulk sample and a metal-rich fraction of the Lost City stone meteorite. In the bulk sample the activity of both isotopes was in fair agreement with the activities reported previously for this meteorite. The activities calculated for the pure metal phase (³⁷Ar = 15.1 ± 2.5 dpm/kg, ³⁹Ar = 20.0 ± 1.5 dpm/kg), however, were different. In particular, the activity ratio of 0.75 ± 0.10 was considerably higher than the previously published values. Consequently the radial heliocentric gradient of the galactic cosmic radiation was found to be much smaller. The data available for the activity ratios in the metal of different meteorites are discussed. Because the wide scatter in the ratios observed in meteorites that fell at comparable times during the solar cycle is due to different decay rates of ³⁷Ar, it is argued that at least during solar maximum the ratios depend essentially on the perihelions of the meteoroid orbits and on whether the meteorites fall on Earth when they are on their inward-bound trip or when they are going out. Except in special cases like Lost City, different aphelions will have a noticeable influence only for meteorites falling near solar minimum, provided that the spatial gradient of the cosmic-ray intensity is small or zero at solar minimum. (auth)

46156 HEAVY-ION RADIATION STUDIES IN TERRESTRIAL AND EXTRATERRESTRIAL MATERIALS. Seitz, Martin George.

St. Louis; Washington Univ. (1972). 132p. University Microfilms Order No. 72-17,959.

Thesis.

Analysis of radiation effects is undertaken to identify changes from solar and galactic radiation that might be expected in extra-terrestrial samples. X-ray observations reveal macroscopic crystal distortion in terrestrial samples of pyroxenes, feldspars, quartz, apatite, epidote, and muscovite irradiated with 10.4-MeV/amu argon ions to doses of 2.0×10^{13} cm². No distortion is seen in olivine. The distortion occurs to a lesser degree with 10.4-MeV/amu neon and not at all with 7-MeV/amu helium irradiations to doses of 4.0×10^{13} and 9×10^{15} ions/cm², respectively. The distortion increases with decreasing energy of the irradiation, and, at any energy, is proportional to dose. Using a model of crystal distortion, the experimental results are better explained by the damage being proportional to the primary rate of ionization of the incident ions rather than to the number of displacements produced by close nuclear encounters. Some irradiated minerals show lattice-parameter changes. Sometimes, crystal fracturing results from severe distortion or from strain caused by the ions that have come to rest in the minerals. The minerals show different sensitivities to radiation and are classified accordingly. Except for fracturing, radiation effects recover by annealing. Both lunar and meteoritic materials show crystal distortion similar to that produced by heavy-ion irradiations. However, the distortion does not anneal, demonstrating that it is not of radiation origin. The other radiation effects are not observed in any of the lunar materials, including the very fine material (~20 microns) of the lunar soil where the largest radiation effects are expected. This lack of pronounced radiation effects is consistent with current ideas of the radiation history of lunar and meteoritic materials. In the second part of the thesis (fission tracks in lunar samples), apatite, whitlockite, and zircon appear to be of equal importance as uranium sources in the heterogeneous rock 12013. A new titanium-zirconium phase is estimated to contain about 10% of the rock's uranium. The more chemically equilibrated rock 12040 has about 80% of the uranium in whitlockite. Baddeleyite and the new zirconium-titanium phase contain most of the remaining uranium. The potassium-rich residual liquids of rock 12040, with ~7 ppM uranium, contain most of the uranium-bearing grains; thus the well-defined potassium-uranium correlation in whole lunar rocks is valid for 12040 on a microscopic scale as well. Thorium-uranium ratios in minerals of 12013 range from 0.2 to 20 while whitlockites of 12040 have thorium-uranium ratios near 3.7, close to the natural abundance ratio. If plutonium acts chemically like thorium, some previously measured plutonium-uranium ratios for the early solar nebula may be too high. Fossil-track densities in whitlockite of four lunar rocks are found to be higher than expected from primary cosmic rays and spontaneous fission of uranium. Proton irradiations at 28 GeV show that the excess tracks are accountable for from high-energy spallation products. Plastic replicas of the fossil densities are able to isolate primary cosmic rays and fission tracks and allow fission-track dating. This dating technique promises to be an important tool for the search for exotic rock fragments of the lunar soil. (Diss. Abstr. Int., B)

46157 STOPPING PARTICLES UNDERGROUND. Grupen, C.; Wolfendale, A. W.; Young, E. C. M. (Univ. of Durham, Eng.). *Nuovo Cim.*; 10B: No. 1, 144-54 (11 Jul 1972).

An attempt was made to interpret the experimental results on stopping muons underground, which appear to show an anomalously high stopping rate, in terms of atmospheric muons, muons from the decay of locally produced pions, and muons from neutrino interactions. The results of calculations of the various contributions are given and the necessary corrections to the experimental data are discussed. It is shown that the experimental results are in reasonable agreement with the expected contributions from the various sources and that no new process is required. (auth)

46158 DIURNAL EFFECT OF THE COSMIC RAYS DURING THE PERIOD 15 OCTOBER 1965-30 JUNE 1966. II. THE EQUATORIAL COSMIC-RAY ANISOTROPY AND THE INTERPLANETARY MAGNETIC FIELD. Bussoletti, E.; Iucci, N. (Univ., Rome). *Nuovo Cim.*; 10B: No. 1, 56-65 (11 Jul 1972).

Using the values of the "diurnal variation waves", obtained with the method described in Part I, possible correlations between the parameters of the interplanetary magnetic field B and the equatorial cosmic-ray anisotropy in periods of magnetically quiet space were studied. Attention was mostly devoted to the study of the morphology of the diurnal variation connected with the evolving "sector" structure of interplanetary space. Rather big differences from the results obtained in the December 1963-February 1964 period were found. This discrepancy can be attributed to the different phase of the solar activity. (auth)

46159 DIURNAL EFFECT OF THE COSMIC RAYS DURING THE PERIOD 15 OCTOBER 1965-30 JUNE 1966. I. METHOD OF ANALYSIS AND STATISTICAL DISTRIBUTION. Bussoletti, E.; Iucci, N.; Villorosi, G. (Univ., Rome). *Nuovo Cim.*; 10B: No. 1, 43-55 (11 Jul 1972).

An improved method of analysis of the diurnal variation of cosmic rays has been developed starting from the one presented by Bachelet et al (1968). Instead of the usual numerical filtering techniques the properties of the asymptotic cones of acceptance of high-latitude cosmic-ray stations were used. Then all the results were collected into a matrix which could give a compact view of the equatorial anisotropy on an hourly basis. Because of the lack of stations in the Asiatic and Pacific area, the study was limited to an "average diurnal variation wave". The statistical distribution of daily values of both amplitudes and phases was analyzed. They agreed with the preceding results of various authors, and also showed an unexpected trend for the direction of the minimum for days with half-peak amplitudes of the anisotropy less than or equal to 0.5%. A preliminary investigation of the north-south anisotropy during Forbush decreases was also done. (auth)

Stars

Refer also to abstracts: 46212, 46466, and 46749.

46160 CORE OF ETA CARINAE. Gehrz, R. D.; Ney, E. P. (Univ. of Minnesota, Minneapolis). *Sky Telesc.*; 44: No. 1, 4-5 (Jul 1972).

The 60-inch telescope at Cerro Tololo was used to obtain a series of high-quality exposures of Eta Carinae on high-speed Ektachrome (ASA-160) color film. An isophote representation was constructed in which structures visible in the Gaviola chart (1950) were identifiable. The two condensations in Thackeray's (1949) elliptical ring were clearly seen. The velocities of the knots and distance were derived. (W.D.M.)

46161 RELATIVISTIC VELOCITY-POTENTIAL HYDRODYNAMICS AND STELLAR STABILITY. Schutz, Bernard Frederick Jr. Pasadena, Calif.; California Inst. of Tech. (1972). 188p. University Microfilms Order No. 72-17,650.

Thesis.

The equations of relativistic, perfect-fluid hydrodynamics are considered in Eulerian form using six scalar "velocity-potential" fields, each of which has an equation of evolution. These equations determine the motion of the fluid. Einstein's equations and the velocity-potential hydrodynamical equations follow from a variational principle. These equations are also cast into Hamiltonian form, with Hamiltonian density $-T_0^0(-g^{00})^{-1/2}$. The second variation of the action is used as the Lagrangian governing the evolution of small perturbations of differentially rotating stellar models. In Newtonian gravity this leads to linear dynamical stability criteria already known. In general relativity it leads to a new sufficient condition for the stability of such models against arbitrary perturbations. By introducing three scalar fields the Newtonian stability criteria are greatly simplified for the purpose of practical applications. The relativistic stability criterion is not yet in a form that permits practical calculations, but ways to place it in such a form are discussed. (Diss. Abstr. Int., B)

46162 INTERPRETATION OF THE NEUTRAL HELIUM TRIPLET SPECTRUM IN PLANETARY NEBULAE. Persson, Sven Eric. Pasadena, Calif.; California Inst. of Tech. (1972). 184p. University Microfilms Order No. 72-17,648.

Thesis.

The theory of the He I triplet spectrum is modified to include the effects of solid particles within the ionized regions of planetary nebulae. New observations of the emission lines $\lambda\lambda 5876$, 7065, and 10830 are presented for over twenty nebulae, extending the list of objects studied previously by O'Dell. The empirical ratios $R_1 \equiv I(\lambda 10830)/I(\lambda 5876)$ and $R_2 \equiv I(\lambda 7065)/I(\lambda 5876)$ are combined with the theory to study the discrepancies between the He I 2^3S depopulation rates D_1 (derived from R_1), D_2 (from R_2), and D_t (theoretical). If the presence of dust is neglected, the discrepancies are such that $D_1 > D_t > D_2$, in most cases. This is essentially the result found earlier by Capriotti for a smaller sample of objects. Estimates of $\tau_d(\lambda 10830)$, the $\lambda 10830$ dust optical depths which force $D_1 = D_t$, are derived by accounting for the resonance trapping and dust absorption of the $\lambda 10830$ radiation. These estimates are consistent, on an order-of-magnitude basis, with values implied by observed infrared excesses in planetary nebulae, and by simple models of reradiating dust shells. The important effects of nebular structural features upon the He I triplet spectrum are studied theoretically and observationally. Overall correlations be-

tween the apparent 2^3S depopulation rate discrepancies and nebular parameters are shown to have a connection with a wide range of nebular problems. (Diss. Abstr. Int., B)

46163 VISCIOUS EFFECTS IN RAPIDLY ROTATING STARS WITH APPLICATION TO WHITE-DWARF MODELS. Durisen, Richard Henry. Princeton, N. J.; Princeton Univ. (1971). 84p. University Microfilms Order No. 72-18,767.

Thesis.

Methods were developed for constructing viscous evolutionary sequences of baryotropic equilibrium configurations with particular application to differentially rotating zero-temperature white-dwarf models. Contributions to the viscosity due to both ions and electrons are discussed. Such sequences were constructed using degenerate electron viscosity. For $M \approx 1.4M_{\odot}$ and $Z = 7$, single timesteps typically show viscous time scales $\sim 10^7$ to 10^8 years, luminosities $\sim 10^1$ to $10^{-1} L_{\odot}$ and effective temperatures $\sim 100,000$ to $25,000^{\circ}K$ respectively. For $Z \geq 10$, ion viscosity probably becomes important for all masses, resulting in shortened time scales. Extensive evolutions show that some stars do not approach uniform rotation. The central density for stars with $M \approx 1.5M_{\odot}$ increases without limit and may even diverge in a finite time, suggesting a Type-I supernova theory. (Diss. Abstr. Int., B)

46164 TYPE A STARS IN THE URSA MAJOR STREAM. Latyshev, I. N. (Leningrad State Univ.). Sov. Astron. AJ (Engl. Transl.); 15: No. 6, 1010-17(May 1972).

Translated from Astron. Zh.; 48: No. 6, 1269-79(Nov 1971).

The existence of the Ursa Major stream was investigated on the basis of the radial velocities and proper motions of type A stars brighter than 6m, following a method developed by the author. About 80 of the 1036 stars considered do belong to the stream. (auth)

46165 GUM NEBULA. de Latil, Pierre. Sci. Avenir; No. 303, 362-9(May 1972). (In French).

After a brief review of the different stages of the discovery of the Gum Nebula and its characteristics, hypotheses currently made about its origin and evolution are presented. (France)

46166 BIRTH OF STARS. Bok, Bart J. Sci. Amer.; 227: No. 2, 48-61(Aug 1972).

The general scheme developed here is one favoring the formation of protostars through the process of gravitational collapse in clouds of interstellar gas and dust. The author points out that it seems quite natural that star birth should be occurring now in the spiral arms of the Milky Way and of neighboring galaxies. Many dark nebulas and globules composed of interstellar gas and dust are seen almost in the act of collapsing into protostars or their close relatives. Objects that are either protostars or very young stars have also been observed. Infrared objects provide a natural link between small dark clouds and relatively normal stars; they may be cool, dense dust clouds with a star or cluster of stars near the center. The cocoon nebulas may also supply newborn stars. The T Tauri stars seem to be the next stage and help bridge the gap between the protostars and the young stars. (L.M.T.)

46167 ANGULAR MOMENTUM LOSS FROM CONTACT BINARY SYSTEMS. Moss, D. L. (Univ. of Manchester, Eng.). Mon. Notic. Roy. Astron. Soc.; 157: No. 4, 433-41(1972).

The effect of angular momentum loss on zero-age main sequence contact binary systems constructed according to the Lucy model was investigated. It was found that the separation of the mass centers decreases, the mass ratio decreases, and the system adopts a dumb-bell configuration. If the system reaches a configuration in which it fills its outer closed common equipotential surface it is unstable to mass loss with the local angular momentum through the outer Lagrange point. (auth)

46168 PHOTOELECTRIC MEASUREMENTS OF THE 6284 Å DIFFUSE INTERSTELLAR LINE. Murdin, Paul (Royal Greenwich Observatory, Hailsham, Eng.). Mon. Notic. Roy. Astron. Soc.; 157: No. 4, 461-75(1972).

A photoelectric index of the $\lambda 6284$ diffuse interstellar line was observed in 48 stars. It is shown that this line is probably exactly correlated with the line at $\lambda 4430$. There are galactic variations in the ratio of $\lambda 6284$ to interstellar reddening, correlated with similar variations of $\lambda 4430$ to reddening. There is no clear evidence whether these variations are correlated with variations of the ratio of total-to-selective absorption. (auth)

46169 INFLUENCE OF LOCAL CONDITIONS IN THE INTERSTELLAR MEDIUM UPON STAR FORMATION. Hunter, James H. (Univ. of South Florida, Tampa). Mon. Notic. Roy. Astron. Soc.; 157: No. 4, 17P-19P(1972).

The gravitational stability of a spherical cloud surrounding a

stellar core in the interstellar medium is discussed utilizing a modified form of the virial theorem. It is argued that the masses of the circumstellar shells surrounding most infrared objects should not exceed $\sim 10^{-2} M_{\odot}$. (auth)

46170 WHITE DWARFS AND NEUTRON STARS. Sutherland, Peter Gordon. Urbana, Ill.; Univ. of Illinois (1972). 148p. University Microfilms Order No. 72-19,943.

Thesis.

The equation of state of zero-temperature matter in complete nuclear equilibrium is given for mass densities below 5×10^{14} g/cm³. Taking into account the effect of the Coulomb lattice and using more recent nuclear mass extrapolations, the sequence of equilibrium nuclides present at mass densities between 10^4 g/cm³ and 4.3×10^{11} g/cm³ was redetermined, the point of neutron drip; and the equation of state was calculated. For densities between 4.3×10^{11} g/cm³ and 5×10^{14} g/cm³ the equation of state was taken from the calculations of Baym, Bethe, and Pethick; included were extensions of the equation of state to still higher densities, taken from the calculations of Pandharipande, which include the effects of hyperons. Zero-temperature white dwarf and neutron star models were calculated for the equation of state. The maximum stable white dwarf mass, the Chandrasekhar limit, was $1.00 M_{\odot}$. The lightest stable neutron star has a mass $0.0925 M_{\odot}$ and a central intensity of 1.55×10^{14} g/cm³; neutron stars between $0.0925 M_{\odot}$ and $0.11 M_{\odot}$ are entirely solid. Moments of inertia, surface deformations and mass quadrupole moments were calculated for slowly rotating neutron stars. Some of the effects of strong magnetic fields (B in the range 10^4 to 10^8 gauss) on the line spectra and polarization of light from white dwarfs are discussed. Exact formulae for line centers and rms widths for the Balmer series in hydrogen are given, neglecting mixing between states of different principal quantum numbers. The range of validity of this approximation and deviations from the Van Vleck and Schiff-Snyder formulae are discussed. A general framework for discussing the origin of the polarization of light from white dwarfs is described and the polarization features produced by the hydrogen bound-free opacity are considered within this framework. The consequences of these calculations for the interpretation of white dwarf spectra are discussed. (Diss. Abstr. Int., B)

46171 INTERSTELLAR DUST AND DISTANCES TO PLANETARY NEBULAE. Lutz, Julie Haynes. Urbana, Ill.; Univ. of Illinois (1972). 108p. University Microfilms Order No. 72-19,872.

Thesis.

UBV and H_γ data were used to construct diagrams of color excess versus distance for early-type stars within 1.5° of eight planetary nebulas. These diagrams and the observed extinctions of the planetaries were used to derive distances for six of these nebulas. The diagrams were also used in conjunction with Cahn and Kaler (Astrophys. J. Suppl., 22, 319, 1971) distances for the program nebulas to derive a value of R, the ratio of total to selective absorption in the visual region, for the line of sight to each nebula. (Diss. Abstr. Int., B)

46172 CONVECTION IN STELLAR ATMOSPHERES. Travis, Larry Dean. University Park, Pa.; Pennsylvania State Univ. (1971). 220p. University Microfilms Order No. 72-19,393.

Thesis.

A survey of the principal theories of convection in stellar atmospheres was made for purposes of comparison as well as detailed examination of the deficiencies in the theories and numerical procedures. Since one of the major problems encountered was the instability of the standard model iteration techniques when convection was included, an improved iterative procedure was developed. This procedure, a linear relaxation method similar to that employed by Auer and Mihalas (1970 M. N., 149, 65) for non-LTE model calculations, has an advantage of extreme flexibility in that it can be applied to nearly any formalism for including convection. (TSS) (Diss. Abstr. Int., B)

46173 EQUILIBRIUM STRUCTURE OF CLOSE BINARY STARS. Naylor, Mark Dennis Thomson. Toronto; Univ. of Toronto (1971). National Library of Canada, Ottawa.

Thesis.

Two hundred and forty-five interior models were obtained for uniformly rotating synchronous close binary systems, using a modified double-approximation technique. Primaries of 10, 5, and $2 M_{\odot}$ with mass ratios of 0.0 to 1.0 in steps of 0.1 were considered, and a primary of $1 M_{\odot}$ with a mass ratio of unity for the case of small to moderate distortion was studied. (TSS) (Diss. Abstr. Int., B)

46174 TEN-MICRON SPECTROSCOPY OF CIRCUMSTELLAR SHELLS. Gammon, R. H.; Gaustad, J. E.; Treffers, R. R. (Univ.

of California, Berkeley). *Astrophys. J.*; 175: No. 3, 687-91(1 Aug 1972).

The infrared spectra of several stars with $10\text{-}\mu$ "silicate" excesses were observed at high resolution ($\Delta\nu = 4\text{ cm}^{-1}$). A broad emission feature extending from 850 to 1100 cm^{-1} was found, but no fine structure was seen. The excess was consistent with radiation from dust composed of basic silicates but not with that from acidic minerals such as quartz. (auth)

46175 OBSERVATIONS OF PLANETARY NEBULAE AT 1.65 TO 3.4 MICRONS. Willner, S. P. (Smithsonian Astrophysical Observatory, Cambridge, Mass. Harvard Coll. Observatory, Cambridge, Mass.); Becklin, E. E.; Visvanathan, N. *Astrophys. J.*; 175: No. 3, 699-706(1 Aug 1972).

Photometric measurements at $2.2\text{ }\mu$ of 15 planetary nebulae are presented, along with measurements of 12 at $1.65\text{ }\mu$ and seven at $3.4\text{ }\mu$. The measurements agree with the predicted thermal emission from ionized hydrogen and helium except for four nebulae. IC 418 and BD + 30°3639 are brighter than predicted at all three wavelengths, and NGC 40 shows excesses at 1.65 and $2.2\text{ }\mu$, but was not measured at $3.4\text{ }\mu$. NGC 7027 shows an excess only at $3.4\text{ }\mu$. All the excesses are interpreted in terms of emission from dust distributed throughout or surrounding the nebulae. Dust temperatures as high as 1000°K are needed to explain the excess at $1.65\text{ }\mu$ (auth)

46176 STUDIES OF HYDRODYNAMIC EVENTS IN STELLAR EVOLUTION. II. DYNAMIC INSTABILITIES IN STELLAR ENVELOPES. Sparks, Warren M. (Goddard Space Flight Center, Greenbelt, Md.); Kutter, G. S. *Astrophys. J.*; 175: No. 3, 707-15(1 Aug 1972).

For an evolved red giant of $1.1\text{ }M_\odot$ (He and heavy-element core of $1.0\text{ }M_\odot$, H-rich envelope of $0.1\text{ }M_\odot$) the dynamic instabilities that result when the luminosity exceeds the limiting value L_0 for a static envelope were investigated. Assuming a constant opacity, $L_0 \approx 10^{3.3}[(M/M_\odot)/\kappa] L_\odot$. When $L > L_0$, the H-rich envelope expands and detaches itself as a shell from the stellar core with final velocities of a few km s^{-1} and rates of mass flow of 10^{-5} to $10^{-4}\text{ }M_\odot\text{ year}^{-1}$. There exists an infrared phase which lasts until the entire escaping envelope is optically thin and becomes heated by the ultraviolet radiation from the stellar remnant. If T- and ρ -dependent opacities were used, the value of L_0 would increase somewhat and the ejection of the H-rich envelope would be preceded by pulsation. These instabilities may be responsible for the pulsation of long-period variables, the ejection of planetary nebulae, and the formation of certain infrared objects. (auth)

46177 RED SUPERGIANTS AND NEUTRINO EMISSION. II. Stothers, Richard (Goddard Space Flight Center, New York). *Astrophys. J.*; 175: No. 3, 717-30(1 Aug 1972).

The variation with stellar mass of the ratio of the numbers of blue and red supergiants was investigated. Statistical data for supergiants in young open clusters and subgroups of associations were collected to supplement a more restricted list in Paper I. Improved methods were used to identify hydrogen-burning supergiants, as well as faint supergiant remnants of binary mass exchange, and to arrange the bright evolved supergiants in order of their masses. Neither of these two operations requires knowledge of stellar distances or luminosities. Relevant published work on stellar evolution, rotation, mass loss, and duplicity was used to predict upper and lower limits on the blue-to-red ratio. It was concluded that (1) the observed paucity of very massive red supergiants (and of carbon stars) confirms and extends the trend observed in Paper I, and thereby supports the idea of neutrino-induced acceleration of the carbon-burning and later phases of evolution; (2) the h and χ Persei association shows the same dependence of the blue-to-red ratio on stellar mass as do other clusters and associations; and (3) a moderate decrease of the blue-to-red ratio is observed with increasing galactocentric distance in the Galaxy (as in M33) and seems to be due to a relative scarcity of extremely young stars in distant galactic regions. (129 references) (auth)

46178 HYDROSTATIC OXYGEN BURNING IN STARS. II. OXYGEN BURNING AT BALANCED POWER. Woosley, S. E.; Arnett, W. David; Clayton, Donald D. (Inst. of Theoretical Astronomy, Cambridge, Eng. Rice Univ., Houston, Tex.). *Astrophys. J.*; 175: No. 3, 731-49(1 Aug 1972).

The nucleosynthesis that occurs during hydrostatic oxygen burning in a "typical" stellar zone was examined in detail under the assumption that such stellar zones are in an approximate state of balanced power from $^{16}\text{O} + ^{16}\text{O}$ energy generation and from neutrino (or photon) energy loss. Special attention was paid to the amount of neutron enrichment that occurs and the implications for nucleosynthesis theory. Accuracy of a common approximation for $^{16}\text{O} + ^{16}\text{O}$ energy generation is discussed also. Theoretical models which

assume the existence of a direct $e - \nu$ coupling in the weak interaction give results that are more consistent with current ideas regarding stellar nucleosynthesis than models which do not. (auth)

46179 STELLAR REACTION RATES FOR THE $^{12}\text{C} + ^{12}\text{C}$, $^{12}\text{C} + ^{16}\text{O}$, AND $^{16}\text{O} + ^{16}\text{O}$ REACTIONS. Michaud, Georges (Univ. of Montreal). *Astrophys. J.*; 175: No. 3, 751-6(1 Aug 1972).

Thermonuclear reaction rates were calculated for the $^{12}\text{C} + ^{12}\text{C}$, $^{12}\text{C} + ^{16}\text{O}$, and $^{16}\text{O} + ^{16}\text{O}$ reactions using optical models for the extrapolation from the recent measurements. At $T_9 \leq 0.7$ the thermonuclear reaction rate for $^{12}\text{C} + ^{12}\text{C}$ is larger than that of Patterson, Winkler, and Zaidins by a factor of 10 but uncertain by a factor of 4. (auth)

46180 OBSERVATIONS OF THE He II $\lambda 10124$ LINE IN O AND OF STARS. Mihalas, Dimitri (National Center for Atmospheric Research, Boulder, Colo.); Lockwood, G. W. *Astrophys. J.*; 175: No. 3, 757-64(1 Aug 1972).

Observed equivalent widths of He II $\lambda 10124$ in absorption-line O stars are found to be in agreement with non-LTE computations assuming no overlap of H and He II lines, and in disagreement with the predictions of LTE and of non-LTE calculations assuming exact coincidence of even-even He II lines and the corresponding hydrogen lines. It is therefore inferred that pumping of He II transitions by H lines does not actually occur in the O stars. The discovery of the $\lambda 10124$ line in emission in ζ Pup (O5f) argues further in this direction; and with the often-cited selectivity of the He II emission mechanism now much weakened, it is suggested that the causes of the emission must involve chromospheric phenomena. Analysis of source-sink terms in equivalent-two-level-atom source functions suggests trends that are compatible with the observed behavior of various lines. (auth)

46181 METALLICISM IN BORDER REGIONS OF THE Am DOMAIN. I. EXTREMELY YOUNG Am STARS IN THE ORION 1c ASSOCIATION. Smith, Myron A. (Univ. of California, Santa Cruz). *Astrophys. J.*; 175: No. 3, 765-77(1 Aug 1972).

A search for metallic-line stars by means of 127 and $63\text{ }\text{\AA}\text{ mm}^{-1}$ spectrograms and uvby photometry resulted in the discovery of five such stars in the extremely young Orion 1c association. A number of arguments in favor of their membership are made. It appears that Am stars in the pre-main-sequence-contraction phase occupy the same domain of the H-R diagram as do evolving Am stars. From the general properties of these stars it is concluded that the Am mechanism becomes operative and stabilizes within about 10^6 years. Evidence is also presented that in open clusters the characteristic rotation of stars, and not age directly, is important in determining the probability that an A star will become metallic lined. These conclusions are in best accord with an interpretation of metallicism in terms of subsurface diffusion. (auth)

46182 ROTATIONAL VELOCITIES OF Ap STARS. Abt, Helmut A.; Chaffee, Frederic H.; Suffolk, Graham (Kitt Peak National Observatory, Tucson, Ariz.). *Astrophys. J.*; 175: No. 3, 779-85(1 Aug 1972).

Projected rotational velocities were measured on coudé spectra for essentially all the northern bright Ap stars. Differences in these rotational velocities between subgroups are found to be marginal, except that Guthrie's discovery of a deficiency of very narrow-lined Si stars is confirmed. Ap stars have one-quarter the mean projected rotational velocity of normal stars of the same types and luminosities, but it is likely that a low rotational velocity is an insufficient criterion to produce a peculiar spectrum. Ap stars cannot be explained simply by an aspect effect. A comparison of projected rotational velocities with equatorial rotational velocities computed from periods of variations in light, magnetic field, or spectrum leads to evidence for random orientations of rotational axes. (auth)

46183 LUMINOSITIES AND MOTIONS OF THE F-TYPE STARS. II. METAL-DEFICIENT STARS. Eggen, O. J. (Australian National Univ., Canberra). *Astrophys. J.*; 175: No. 3, 787-807(1 Aug 1972).

An improved method of computing luminosities of F- and early G-type metal deficient stars from intermediate-band photometry is applied to the available photometry for some 250 objects with $\Delta[m]_1 > +0.025$, including all of those brighter than visual magnitude 6.5 for which accurate apparent motions are available. Twenty halo stars, for which $\Delta[m]_1 > +0.080$, are included. The distribution of the resulting space motions is in excellent agreement with that found from previous studies, based on UVB photometry, and used in discussing the dynamics of a collapsing galaxy. (auth)

46184 PRELIMINARY ANALYSIS OF THE ATMOSPHERIC ECLIPSE OF 32 CYGNI. Galatola, Anthony (La Salle Coll., Philadelphia). *Astrophys. J.*; 175: No. 3, 809-18(1 Aug 1972).

The blue and ultraviolet light curves of 32 Cygni were analyzed by using as a model an opaque core surrounded by a semitransparent envelope. The effects of the atmospheric eclipse were removed from the light curve to obtain a light curve of the body eclipse. In order to get a solution the body eclipse was assumed to be total. This led to an orbital inclination of 82° and $1.87 R_\odot$ for the radius of the B star. The radius of the K star was $120 R_\odot$ in blue light and $123 R_\odot$ in ultraviolet light. It was found that the light curves could be calculated if the opacity-density relation were written in the form $k\rho = Ae^{-\alpha r} + Be^{-\beta r} + Ce^{-\gamma r}$, where r is the radial distance and A , B , C , α , β , and γ are constants. Close to the surface of the K star the opacity in the blue was greater than that in the ultraviolet. This led to the conclusion that the cumulative effect of line absorption was a possible source of opacity for the light of the B star. (auth)

46185 MEASUREMENT OF POLARIZED 10-MICRON RADIATION FROM COOL STARS WITH CIRCUMSTELLAR SHELLS. Capps, R. W.; Dyck, H. M. (Kitt Peak National Observatory, Tucson, Ariz.). *Astrophys. J.*; 175: No. 3, 693-71 (Aug 1972).

Observations are reported of the linear polarization of 8 to $14\text{-}\mu$ radiation from 10 cool stars. It was found that the level of measurable polarization is small, about 1%, but that a possibly significant detection was achieved for several stars. The implications of the data are discussed briefly. (auth)

46186 EARLY DATA FROM THE ULTRAVIOLET SKY-SCAN TELESCOPE IN THE TD1 SATELLITE. Wilson, R. (Astrophysics Research Unit, Culham, Eng.); Gardier, S.; Jamar, C.; (and others). *Nature (London) Phys. Sci.*; 238: No. 81, 34-6 (17 Jul 1972).

An initial examination of data from the TD1 satellite telescope shows it to be functioning satisfactorily. The spectra observed are in good agreement with information obtained from rocket-borne telescopes. (auth)

46187 OPTICAL OBSERVATIONS OF THE SUPERNOVA IN NGC 5253. Osmer, P. S. (Cerro Tololo Inter-American Observatory, La Serena, Chile); Hesser, J. E.; Kunkel, W. E.; (and others). *Nature (London) Phys. Sci.*; 238: No. 80, 21-2 (10 Jul 1972).

The Type I supernova in NGC 5253, discovered recently, is stated to be of particular interest because of its brightness and because it is the second supernova to occur in this peculiar galaxy during the past 100 years. Some rather extensive studies made from Cerro Tololo are described, including UVB photometry and spectroscopic observations. The colors were found to be those of an F-type star, suggesting that the emitting region is very large and fairly cool. No rapid variations in total light were observed. The spectroscopic observations showed the broad emission features characteristic of Type-I supernovae, as well as narrow interstellar features in the H and K lines. Additional bands in the uv and near-infrared, apparently not previously detected, were observed. Observations of the fluxes of the supernova in the wavelengths between the emission lines showed peaks near $5000 \text{ \AA}$ and, although the spectrum does not have a blackbody shape, the fluxes imply that the temperature of the emitting region is of the order 6 to $8 \times 10^3 \text{ K}$. If the region is optically thick it has a radius of $\sim 10^4 R$. The upper limit for the expansion velocity appears to be $10,000 \text{ km/sec}$. (UK)

46188 EVIDENCE FOR THE HYDRODYNAMIC CHARACTER OF MICROTURBULENCE. Van Paradijs, J. A. (Univ., Amsterdam). *Nature (London) Phys. Sci.*; 238: No. 81, 37-8 (17 Jul 1972).

The validity of determinations of the chemical composition of stellar atmospheres from observations of absorption lines in stellar spectra has been questioned (Worrall and Wilson, *Nature*, 236: 15 (1972)), the main argument being directed against the concept of "microturbulence." This concept, which has been used for many years, is discussed, and observations are presented that support the concept and indicate that "microturbulence" is a real hydrodynamic effect. (UK)

46189 RELATIVISTIC ROCHE PROBLEM. Fishbone, Leslie G. (Univ. of Maryland, College Park). *Astrophys. J.*; 175: No. 3, L155-9 (1 Aug 1972).

Bodies in orbit around Kerr black holes will experience tidal gravitational forces. The Roche limit for fluid bodies in stable circular orbits is qualitatively like that in Newtonian situations; for highly energetic circular orbits, the fluid density required for the existence of a body in equilibrium is magnified by the square of its energy-at-infinity per unit mass. (auth)

46190 CIRCUMSTELLAR MOLECULES IN THE PLEIADES. Hobbs, L. M. (Univ. of Chicago). *Astrophys. J.*; 175: No. 3, L145-7 (1 Aug 1972).

Both old and new observations of the interstellar $\lambda 4232$ line of CH^+ are considered for the brightest members of the Pleiades. They suggest that the molecules are circumstellar in some sense, perhaps resembling in this respect the micron-sized grains inferred to be present in this region. (auth)

46191 PRIMARY SPECTRUM OF THE ECLIPSING BINARY LR CENTAURI. Bessell, M. S. (Australian National Univ., Canberra). *Astrophys. J.*; 175: No. 3, L133-6 (1 Aug 1972).

Coudé spectra and photoelectric scans were obtained during maximum light for LR Cen, an eclipsing binary of period 2.095595d variously suspected and discounted as being the pulsating x-ray source Cen X-3. These observations indicate the primary to be a reddened $[E(B-V) \approx 0.30]$, shallow-lined ($v \sin i = 300 \text{ km s}^{-1}$) B8.5 III star with several shell features, strong broad absorption at Ca H and K and Na D, strong narrow absorption at Na D, and H α partly filled in. If one assumes synchronous rotation and orbital revolution, a radius of $12 R_\odot$ is determined for the primary, in agreement with the x-ray results of Cen X-3. The star is filling its Roche lobe and possibly losing mass. (auth)

46192 BLACK HOLES IN THE BRANS-DICKE. THEORY OF GRAVITATION. Hawking, S. W. (Cambridge Univ., Eng.). *Commun. Math. Phys.*; 25: No. 2, 167-71 (1972).

It is shown that a stationary space containing a black hole is a solution of the Brans-Dicke field equations if and only if it is a solution of the Einstein field equations. This implies that when the star collapses to form a black hole, it loses that fraction (about 7%) of its measured gravitational mass that arises from the scalar interaction. This mass loss is in addition to that caused by emission of scalar or tensor gravitational radiation. Another consequence is that there will not be any scalar gravitational radiation emitted when two black holes collide. (auth)

46193 BLACK HOLES IN GENERAL RELATIVITY. Hawking, S. W. (Cambridge Univ., Eng.). *Commun. Math. Phys.*; 25: No. 2, 152-66 (1972).

It is assumed that the singularities which occur in gravitational collapse are not visible from outside but are hidden behind an event horizon. This means that one can still predict the future outside the event horizon. A black hole on a space-like surface is defined to be a connected component of the region of the surface bounded by the event horizon. As time increases, black holes may merge together but can never bifurcate. A black hole would be expected to settle down to a stationary state. It is shown that a stationary black hole must have a topologically spherical boundary and must be axisymmetric if it is rotating. These results together with those of Israel and Carter go most of the way towards establishing the conjecture that any stationary black hole is a Kerr solution. Using this conjecture and the result that the surface area of black holes can never decrease, one can place certain limits on the amount of energy that can be extracted from black holes. (auth)

46194 NEAR-INFRARED PHOTOMETRY OF MIRA VARIABLES. Lockwood, G. W. (Kitt Peak National Observatory, Tucson, Ariz.). *Astrophys. J., Suppl. Ser.*; 24: No. 209, 375-419 (Jun 1972).

A five-color narrow-band photometric system is described which has been used for determining magnitudes at 1.04μ , near-infrared colors, and molecular band-strength indices for M stars. The system gives accurate spectral types from M4 to M10 and useful, though less accurate, spectral types from M0 to M4. Mean 1.04μ magnitudes, near-infrared colors, and spectral-type indices are given for photometric and spectroscopic standard stars, and 1795 individual sets of five-color measurements are given for 292 M- and S-type Mira variables. The Miras are shown to map out large loops in a diagram of molecular band strength as a function of color, and large differences of band strength can occur at a given color and phase in different stars or in the same stars during different cycles. The Miras with periods less than 300d are warmer by several hundred degrees at a given spectral type than either the normal giants used for calibration or the Miras with periods longer than 300d. The mean phase of earliest spectral type is ~ 0.03 , while maximum light at 1.04μ occurs at ~ 0.11 . Minimum light at 1.04μ and latest spectral type are reached almost simultaneously with minimum visual light. (auth)

46195 LINEAR PULSATIONS AND STABILITY OF DIFFERENTIALLY ROTATING STELLAR MODELS. I. NEWTONIAN ANALYSIS. Schutz, Bernard F. Jr. (California Inst. of Tech., Pasadena). *Astrophys. J., Suppl. Ser.*; 24: No. 208, 319-46 (May 1972).

A systematic method is presented for deriving the Lagrangian governing the evolution of small perturbations of arbitrary flows

of a self-gravitating perfect fluid. The method is applied to a differentially rotating stellar model; the result is a Lagrangian equivalent to that of Lynden-Bell and Ostriker. A sufficient condition for stability of rotating stars, derived from this Lagrangian, is simplified greatly by using as trial functions not the three components of the Lagrangian displacement vector, but three scalar functions. This change of variables saves one from integrating twice over the star to find the effect of the perturbed gravitational field. (auth)

46196 LINEAR PULSATIONS AND STABILITY OF DIFFERENTIALLY ROTATING STELLAR MODELS. II. GENERAL RELATIVISTIC ANALYSIS. Schutz, Bernard F. Jr. (California Inst. of Tech., Pasadena). *Astrophys. J., Suppl. Ser.*; 24: No. 208, 343-74 (May 1972).

The author has previously given an Eulerian velocity-potential variational principle for relativistic perfect-fluid hydrodynamics. The second variation of the principle is here used as the Lagrangian density governing the evolution of small perturbations of fully relativistic, differentially rotating stellar models. Noether's theorem is used to construct a globally conserved angular-momentum density, whose integral over a spacelike hypersurface is the second-order correction to the star's total angular momentum. From the Hamiltonian is constructed a globally conserved energy density, whose integral is the second-order correction to the star's active gravitational mass. By Liapunov's second theorem, positive-definiteness of the energy density guarantees stability of the star. In the Newtonian limit and in the special case of relativistic radial pulsations, this is equivalent to stability criteria already known. Means are discussed whereby the general criterion might be made more suitable for practical applications. (auth)

46197 STARS WITH DARK COMPANIONS. Prosen, Marijan. *Obz. Mat. Fiz.*; 18: No. 1, 20-3 (Mar 1971). (In Slovenian).

This contribution deals with a short survey of investigations of the stars supposed to have dark companions. Periodic perturbations of the proper motion of the near star may be explained by existence of one or more dark companions moving around the star. The perturbations of the proper motion of BD + 43561 are explained by the possibility that two dark companions, with masses like Jupiter and Saturn, could be moving around it. (auth)

Quasi-Stellar, Radio, and X-Ray Sources

46198 FINE STRUCTURE IN H II REGIONS. Balick, Bruce. Ithaca, N. Y.; Cornell Univ. (1971). 184p. University Microfilms Order No. 72-15,390. Thesis.

Interferometric studies at $\lambda 3.7$ and 11.1 cm are reported for the H II regions DR 21, W51 components G49.5-0.4 and G49.3-0.3, W33, NGC 2264S, and IC 410. At these wavelengths the resolutions are $3''$ and $10''$, respectively; the former is the best resolution yet achieved for the mapping of H II regions. Some of the nebulae are shown to exhibit strong variations in their radio intensity called fine structure, with scale sizes of $10^{-1.5}$ pc. Other H II regions show either intermediate structure ($10^{-0.5}$ pc) or no structure at all. An accurate method is developed to estimate the emission measure and electron temperature based on the spectral behavior of the structure emission. Typical fine-structure emission measures of $10^{7.5}$ pc cm $^{-6}$ and electron temperatures of 10^4 K were found. From this, electron densities of $10^{4.5}$ cm $^{-3}$ and nebular masses of $10^{-1} M_{\odot}$ can be deduced. Since pressure gradients associated with the high densities will tend to dissociate the structure, mechanisms which might confine or resupply the density are explored. Two models emerge as possible explanations. One is based on flows from high-density neutral material imbedded within the H II region. The second depends on neutral gas of a protostellar cocoon to confine a high-density H II region in its interior. The first of these models is shown to be slightly preferable on the basis of its compatibility with other properties of H II regions. In addition, a model for the well-studied core of the Orion nebula based entirely on flows is constructed; it is shown that this model is the first to provide an interpretive framework for all available radio and optical observations of Orion. (Diss. Abstr. Int., B)

46199 OBSERVATION OF PULSED GAMMA RAYS WITH ENERGIES GREATER THAN 250 keV FROM THE CRAB NEBULA PULSAR NP 0532. Orwig, Larry Eugene. Durham, N. H.; Univ. of New Hampshire (1972). 231p. University Microfilms Order No. 72-19,028. Thesis.

Data from a balloon flight experiment using an $11\frac{1}{2}''$ diameter $\times$

4" thick NaI scintillation spectrometer were analyzed to search for gamma-ray pulsations from NP 0532. The flight was conducted on June 7, 1970, and the payload was carried to a nominal atmospheric depth of 3.5 g/cm ^{2} . The detector was pointing in the direction of the Crab Nebula for the entire time at float altitude. A superposed epoch analysis was performed on a 12,000 second portion of the data spanning a total time interval of 16,700 seconds at float altitude. A positive pulsed contribution was observed at the expected apparent frequency of NP 0532 having the double pulse structure typical of NP 0532. The results indicate a time-averaged pulsed photon flux of $(1.44 \pm 0.57) \times 10^{-3}$ photons cm $^{-2}$ sec $^{-1}$ in the energy interval from 250 keV to 2.3 MeV. This represents a time-averaged pulsed power of $I = (6.49 \pm 2.57) \times 10^{-4}$ keV cm $^{-2}$ sec $^{-1}$ keV $^{-1}$. The ratio of interpulse to main pulse was found to be 2.4 ± 1.9 . A further subdivision of the energy interval was performed to obtain spectral information above 250 keV. Although the data is statistically very marginal, the analysis indicates a positive photon flux from 250 keV to 725 keV of $(1.20 \pm 0.52) \times 10^{-3}$ photons cm $^{-2}$ sec $^{-1}$ or a time-average pulsed power of $(1.14 \pm 0.50) \times 10^{-3}$ keV cm $^{-2}$ sec $^{-1}$ keV $^{-1}$. Above 725 keV no pulsed contribution was found. Therefore, upper limits at the 95% confidence level were obtained for two energy bins, one extending from 725 keV to 2.3 MeV, and the second an integral channel above 2.3 MeV. No evidence was found for an enhancement in the pulsed flux at the 0.5 MeV annihilation-line region. (Diss. Abstr. Int., B)

46200 OBSERVATIONS OF 3C 272.1 AT 2.7 AND 5.0 GHz. Riley, J. M. (Cambridge Univ., Eng.). *Mon. Notic. Roy. Astron. Soc.*; 157: No. 4, 349-57 (1972).

The radio source 3C 272.1, associated with the elliptical galaxy M84 in the Virgo cluster, was observed at 2.7 and 5.0 GHz with half-power beam widths of $12 \times 52''$ arc and $6.5 \times 28''$ arc, respectively. The maps obtained show that the source has a spiral structure. Linear polarization of up to 20% in places was found, consistent with a magnetic field predominantly perpendicular to the axis of the spiral. (auth)

46201 RELATIVISTIC BEAMING. McCrea, W. H. (Univ. of Sussex, Brighton, Eng.). *Mon. Notic. Roy. Astron. Soc.*; 157: No. 4, 359-65 (1972).

Working in terms of photon-fluxes rather than other radiational parameters, a short self-contained treatment is given of the observable properties of a revolving light-source recently discussed by F. G. Smith (1970, 1971) in connection with a theory of the radiation of pulsars. (auth)

46202 ACCURATE POSITIONS OF RADIO SOURCES AT 408 MHz. Hunstead, R. W. (Univ. of Sydney). *Mon. Notic. Roy. Astron. Soc.*; 157: No. 4, 367-402 (1972).

The Molonglo radio telescope was used to measure positions and flux densities for 314 small-diameter radio sources between declinations of $+19^\circ$ and -80° . The mean estimated errors in position were less than $1.5''$ in right ascension and $2.0''$ in declination; this accuracy was confirmed by a limited series of comparisons with precise interferometer positions. A large fraction of previously suggested southern identifications was not supported by the more accurate radio positions. Source extensions were found in 25% of the listed sources by application of a sensitive method for detecting the direction of elongation. (auth)

46203 OBSERVATIONS OF THE SUPERNOVA REMNANT IC 443 AT 1.42 GHz. Hill, I. E. *Mon. Notic. Roy. Astron. Soc.*; 157: No. 4, 419-31 (1972).

The supernova remnant IC $_{443}$ was mapped at 1.42 GHz with a resolution of $1.05 \times 2.71''$ arc, using the Half-Mile telescope at Cambridge. The map shows that the source has a thick shell extending over the eastern half of its circumference with no evidence of a shell on the western edge. There is remarkably detailed agreement of optical and radio features, with some optical filaments clearly located on the outer edge of radio ridges. Comparison of the map with other high-resolution maps at higher frequencies shows an increase in the spectral index near the brightest optical features. Estimates of some physical parameters of the source are derived in the final section of the paper. (auth)

46204 POLARIZATION OF RADIO SOURCES AT $\lambda 49$ CM AND $\lambda 73$ CM. Conway, R. G.; Gilbert, J. A.; Kronberg, P. P.; Strom, R. G. (Univ. of Manchester, Jodrell Bank, Eng.). *Mon. Notic. Roy. Astron. Soc.*; 157: No. 4, 443-59 (1972).

Measurements are presented of the flux density at $\lambda 73$ cm and the linear polarization at $\lambda 49$ and $\lambda 73$ cm of extragalactic sources, mostly quasars. These measurements establish the degree to which the linear polarization decreases at long wavelengths. Evidence is presented that the depolarization does not occur in the galactic halo. Little or no depolarization seems to occur in radio

variables. The polarization at $\lambda 49$ cm of two sources (ζ C 454.3 and BL Lac) may have changed over a period of one year. (auth)

46205 VARIABILITY OF THE OH SOURCE W3. Wilson, A. J.; Davies, R. D.; Elder, J. (Nuffield Radio Astronomy Labs., Jordell Bank, Eng.). Mon. Notic. Roy. Astron. Soc.; 157: No. 4, 21P-6P(1972).

Variability was detected in the 1665 MHz spectrum of the W₃OH source. Observations extending back to 1965 show an average decrease of $8.9 \pm 0.9\%$ per year in the -41.7 km s^{-1} feature. Irregular variability on the time scale of a few years is also found. The implications of these observations are discussed. (auth)

46206 MICROWAVE CELESTIAL WATER-VAPOR SOURCES. Johnston, K. J.; Knowles, S. H.; Schwartz, P. R. (Naval Research Lab., Washington, D. C.). Sky Telesc.; 44: No. 2, 88-90 (Aug 1972).

Microwave emission from water vapor sources has been detected in two types of sources: OH—H II regions and cool, late M-type stars. These two sources are briefly discussed. (W.D.M.)

46207 POLARIZATION APERTURE SYNTHESIS OF FIVE GALACTIC NONTHERMAL RADIO SOURCES. Hermann, Bruce Raymond. Urbana, Ill.; Univ. of Illinois (1972). 329p. University Microfilms Order No. 72-19,846.

Thesis.

Five possible supernova remnants were mapped using the techniques of aperture synthesis interferometry. The objects 3C 10 (Tycho's supernova), 3C 58, DR 4 (the Gamma-Cygni radio source), 4C 28.16 (located in the shell of S-147), and 3C 358 (Kepler's supernova) were mapped at 11.1 cm wavelength with the National Radio Astronomy Observatory's three-element interferometer. Intensity maps, polarization intensity maps, and electric vector position angle maps were obtained. A special polarization calibration scheme for this instrument was developed which employs both the tensor approach of Ko (I.E.E.E. Transactions of Antennas and Propagation, AP-15, p. 10, 1967) and the empirical correction procedure of Conway and Kronberg (M.N.R.A.S., 142, p. 11, 1969). (Diss. Abstr. Int., B)

46208 ABILITY OF THE LUMINOSITY-VOLUME TEST TO REVEAL THE STATISTICAL EVOLUTION OF THE LUMINOSITY OF QUASI-STELLAR SOURCES. Lynds, Roger (Kitt Peak National Observatory, Tucson, Ariz.); Petrosian, Vahe. Astrophys. J.; 175: No. 3, 591-9(1 Aug 1972).

The principles and capabilities of the luminosity-volume test and source-count analysis as procedures for studying the statistical evolution of the radio and optical luminosities of quasi-stellar sources are reviewed. In contrast to the conclusions expressed by Longair and Scheuer, a comparison of the two methods reveals that the luminosity-volume test is substantially more powerful and lucid and that a knowledge of the individual redshifts of objects is required for the most comprehensive determination of the evolutionary history of quasi-stellar sources. (auth)

46209 VRS FOR VRO. Swenson, G. W. (Univ. of Illinois, Urbana). Nature (London) Phys. Sci.; 238: No. 81, 37(17 Jul 1972).

Reference is made to use of the designation VRO for cosmic radio sources. Originally, this designation referred to the Vermilion River Observatory at the University of Illinois, but subsequently it has been used to indicate "variable radio object." Of the large number of sources in the Vermilion River Observatory catalog very few are known to be "variable radio objects" and in order to avoid confusion it is suggested that the designation VRS ("variable radio source") would be better for this class of object. (UK)

46210 TON 155 AND 156: A DOUBLE QUASAR. Stockton, A. N. (Univ. of Hawaii, Honolulu). Nature (London) Phys. Sci.; 238: No. 81, 37(17 Jul 1972).

It has been shown that Ton 156 is a quasar (Janes and Lynds, Astrophys. J. Lett., 155: L47(1969)), although a satisfactory redshift could not be determined. Spectroscopic observations by the present author, using the Mauna Kea 224-cm telescope, indicated that Ton 155, located at about $35''$ arc north of Ton 156, is also a quasar, and also allowed redshifts to be determined for both objects. Thus the possibility of a double quasar arises. The observations also give support to the possibility that not all quasar redshifts are cosmological. (UK)

46211 OBSERVATION OF A HARD X-RAY FLARE IN CYG X-1. Agrawal, P. C. (Tata Inst. of Fundamental Research, Bombay). Nature (London) Phys. Sci.; 238: No. 80, 22-3(10 Jul 1972).

Data obtained during a balloon flight from Hyderabad on the night of April 6, 1971 (Nature, 232: 38(1971)) were reanalyzed

on a minute-to-minute basis to examine further possible structures in time variation of hard x rays. This analysis revealed the occurrence of a "flare" in Cyg X-1 similar in characteristics to those recorded in Sco X-1. Details of this flare are reported. This appears to be the first observation of a hard x-ray flare in Cyg X-1. (UK)

46212 ASSOCIATION OF X-RAY SOURCES WITH BRIGHT STARS. Gursky, Herbert (American Science and Engineering, Cambridge, Mass.). Astrophys. J.; 175: No. 3, L141-4(1 Aug 1972).

A comparison of selected Uhuru x-ray sources with the SAO star catalog reveals 10 stars within the positional error of the x-ray sources where only about two are expected on the basis of chance. Three of these are already under study as optical candidates for x-ray sources. The new results lend confidence to these candidates as being correct and yield candidate stars for seven other x-ray sources. The association of bright stars with x-ray sources is expected on the basis of the model which invokes mass transfer from a massive star to a compact companion. (auth)

46213 OPTICAL SEARCH FOR CENTAURUS X-3. Brucato, R. J.; Kristian, J.; Westphal, J. A. (Carnegie Institution of Washington, D. C. California Inst. of Tech., Pasadena). Astrophys. J.; 175: No. 3, L137-9(1 Aug 1972).

The optical eclipsing binary LR Cen has been eliminated as a candidate for Cen X-3 on the basis of a real discrepancy of orbital periods. It is believed that the position coincidence of Wray 795 with Cen X-3 is not statistically significant. (auth)

46214 DETECTION OF HIGH-ENERGY GAMMA RAYS FROM THE CRAB NEBULA. Fazio, G. G.; Helmen, H. F.; O'Mongian, E.; Weekes, T. C. (Smithsonian Astrophysical Observatory, Cambridge, Mass.). Astrophys. J.; 175: No. 3, L117-22(1 Aug 1972).

By means of the ground-based atmospheric Cherenkov technique, observations of the Crab Nebula, averaged over a 3-year period, indicate that a flux of γ rays ($4.4 \pm 1.4 \times 10^{-11}$ photons $\text{cm}^{-2}\text{s}^{-1}$) with energy $\geq 2.5 \times 10^{11}$ eV has been detected at the 3.1σ level. This flux corresponds to an emission of 6×10^{33} ergs s^{-1} , significantly less than the continuous x-ray emission. The γ -ray flux may vary with time, with the most significant flux ($1.21 \pm 0.24 \times 10^{-10}$ photons $\text{cm}^{-2}\text{s}^{-1}$) occurring 60 to 120 days after a major spin-up of the pulsar NP 0532. This increase was observed on three different occasions, and if the flux in only these intervals is used, the effect is at the 5σ level. The total γ -ray energy observed on each occasion was $\sim 10^{41}$ ergs, an energy approximately equal to the energy of the pulsar spinup. (auth)

46215 EXTRAGALACTIC ORIGIN OF THE TRANSIENT X-RAY SOURCES. Sofia, Sabatino (Univ. of South Florida, Tampa). Astrophys. J.; 175: No. 3, L113-15(1 Aug 1972).

It is shown that the current assumption that the transient x-ray source 2U 1543-47 is located within our Galaxy leads to serious energetic difficulties. However, all the presently available observations are consistent with the interpretation that the source was produced by a supernova event in a nearby galaxy. On the assumption that the nature of this source is similar to that of the other known transient sources, observational tests are proposed to verify their extragalactic origin. (auth)

46216 POLARIZED RADIO EMISSION FROM FIVE SUPERNOVA REMNANTS. Milne, D. K. (CSIRO, Epping, Australia). Aust. J. Phys.; 25: No. 3, 307-13(Jun 1972).

Maps are presented of the polarization and total power emission at 2700 MHz from five southern supernova remnants: MSH 09-32, 10-53, 14-63, 15-52, and 15-56. Observations of 14-63 and 15-56 at 1410 MHz are also presented. (auth)

Solar Phenomena

Refer also to abstract: 47143.

46217 SOLAR RADIO SPIKE BURSTS. Tarnstrom, Guy LaVern. College, Alaska; Univ. of Alaska (1971). 184p. University Microfilms Order No. 72-17,101.

Thesis.

Spectral observations of meter wavelength solar radio emission with ≈ 0.01 second time-resolution show three types of spike bursts: fast drift spike bursts, with $f \geq 100$ MHz; medium drift spike bursts, with $f < 100$ MHz; and simple spike bursts, of ~ 1 -MHz bandwidth and no apparent frequency drift. All types of spike bursts tend to occur in regular sequences of similar bursts. Rare, extensive sequences of quasi-periodic fast-drift spike bursts are observed. Fine structure of Type-III burst events reveal occa-

sional spike burst—Type-III burst associations which appear to fall into three classes. Type III bursts are observed with duration ~ 0.1 second approaching the range of spike-burst durations ($\lesssim 0.1$ second). Large-frequency-range spike bursts resemble such short-duration Type III bursts, suggesting a morphological continuity between spike bursts and Type-III bursts. A plasma hypothesis interpretation of spike bursts requires spike burst exciter of spatial extent less than $\sim 10^3$ to 10^4 km, density ~ 1 to 10 electrons/cm³, and coronal temperature $\sim 10^6$ K. It is suggested that the spatial extent of a given exciter determines whether a spike burst or a Type-III burst is produced. Unusual meter-wavelength fine structure observed includes spike burst and Type-I burst chains with a sharp high-frequency cutoff, complex patterns of parallel drifting bands, and ionospheric scintillation of noise-storm continuum. Several examples are given. (Diss. Abstr. Int., B)

46218 CONTINUUM NEAR THE H AND K LINES OF Ca⁺ IN THE SOLAR SPECTRUM. Matsushima, Satoshi; Kawabata, Kiyoshi (Pennsylvania State Univ., University Park). *Astrophys. Lett.*; 11: No. 3, 103-6 (Jul 1972).

A method is proposed for the determination of the true continuum for the spectral region where no continuum is visible. The method is based on the use of the characteristic weakening of absorption lines due to blending with a strong line and thereby eliminates various uncertainties involved in theoretical methods attempted earlier. The continuum determined by this method for the H and K lines of Ca⁺ in the solar spectrum is about 10 to 15% higher than the newest empirical determination, and is in good agreement with higher values suggested by earlier workers. (auth)

46219 EVIDENCE FOR SOLAR PARTICLE PRODUCTION ABOVE ~ 75 GeV. Schindler, S. M.; Kearney, P. D. (Colorado State Univ., Ft. Collins). *Nature* (London); 237: No. 5357, 503-5 (30 Jun 1972).

The first direct experimental evidence is presented for solar-particle production in the energy region above the maximum vertical geomagnetic cutoff; this was the outcome of an experiment specifically designed for the detection of solar-generated particles in the energy region above ~ 75 GeV. The detector was a narrow-angle muon telescope consisting of three in-line plastic scintillators and two narrow-gap spark chambers, mounted on a solar tracking equatorial mount. For specific resolution of a very high energy solar particle component the detecting system was located underground at a depth of 200 m at Idaho Springs, Colorado. On Oct. 3rd 1971 a positive correlation was obtained between a significant intensity increase and a specific solar flare event. Preliminary satellite data on the associated proton event are given. Only single station detection was observed, but this is not unique in the history of solar-related particle events. It is considered that the results constitute good evidence for particle acceleration occurring during the initial phase of a solar flare. No specific acceleration mechanism is postulated to account for the results. (UK)

46220 SIZE OF THE DUST PARTICLES OF A COMET ATMOSPHERE AND THE HELIOCENTRIC DISTANCE. Dobrovolski, O. V.; Ibadinov, Kh. (Inst. of Astrophysics, Dushanbe, USSR). *Dokl. Akad. Nauk Tadzh. SSR*; 15: No. 2, 15-18 (1972). (In Russian).

Possible factors that might affect the size of the dust particles in a comet were considered, e.g.: the free flight path of particles during the sublimation of ice, the collapse of the surface of the dust matrix, and the formation of particles in the near-nuclear region of the head by means of condensation, any of which may be related to the heliocentric distance of the comet. During the free escape of dust particles from the surface of the nucleus under the influence of gas pressure, the average diameter of comet particles and the magnitude of their effective acceleration at 4 AU from the Sun is basically a function of the structure of the nucleus and not of the heliocentric distance. Formation of dust by condensation in the near-nuclear region of the head leads to growth of particle dimensions with decreasing heliocentric distance. (K.S.W.)

46221 EXCITER OF TYPE III BURSTS AND CORONAL TEMPERATURE. Aubier, M.; Bolschot, A. (Observatoire, Meudon, France). *Astron. Astrophys.*; 19: No. 3, 343-53 (Jul 1972).

The time profiles of Type-III solar radio bursts are often used to determine some characteristics of the burst exciter, and the temperature of the corona at different heights. It is shown that, under certain hypothesis, the time profiles give independently the duration of the exciter and the decay time constant of the burst. New observations of Type-III bursts in the decimeter range are presented. They were obtained with the 1000-ft dish

of Arecibo (Puerto-Rico). The analysis of the time profiles leads to average total durations of the exciter function of 3.9, 6.2, and 7.5 s and average decay time constants of 1.2, 1.7, and 2.0 s at 60, 36.9, and 29.3 MHz, respectively. The values of these two parameters scatter around these averages from burst to burst and from day to day within a factor of approximately 3. A relation is found between the duration and the decay time constant derived with the above method. This relation brings some doubt on the significance of the coronal temperature derived from the decay time constant with the hypothesis of a collisional damping of the plasma waves. Other explanations are proposed. (auth)

46222 ACTIVE LONGITUDES OF SUNSPOTS AND SOLAR FLARES IN THE 20th CYCLE. Maris, Georgeta (Observatoriu Astronomic, Bucharest). *Stud. Cercet. Astron.*; 17: No. 1, 71-7 (1972). (In Rumanian).

The active longitudes of different indices for sunspots and solar flares were investigated by the isoline method, for the period July 1, 1964–December 31, 1969. It is shown that the active longitudes of sunspots and solar flares practically coincide. The rate of concentration in the active longitudes for different solar events is higher for more important events. The problems connected with north–south asymmetry of solar activity and of the concentration rate in the active longitudes are considered, also. (auth)

46223 RADIO AND X-RAY EMISSIONS ASSOCIATED WITH THE CHROMOSPHERIC FLARES OF NOVEMBER 15 AND 16, 1970. Maris, Georgeta (Observatoriu Astronomic, Bucharest). *Stud. Cercet. Astron.*; 17: No. 1, 79-84 (1972). (In Rumanian).

The radio bursts and x-ray emission associated with the chromospheric flares of November 15 and 16, 1970 are presented. The chromospheric flare of November 15 had two brightness maxima, one of 1B importance at 7h06mUT and one of 1N importance at 7h57mUT. There were two outstanding events in the x-ray range and very complicated radio bursts in the microwave range associated with this flare. The 2N importance flare of November 16 attained brightness maximum at 9h56mUT. There were not such complicated radio bursts as on November 15, but the x-ray flare was similar to the second x-ray flare on November 15. An explanation of these radio and x-ray emissions is proposed, in agreement with the comprehensive model of Takakura and Kai (1966). It is suggested that the differences between the associated phenomena of the two chromospheric flares were due to the different acceleration mechanism of the electrons. (auth)

46224 PHOTOSPHERIC FACULAE AND THE SOLAR OBLATENESS. Chapman, Gary A. (Aerospace Corp., El Segundo, Calif.); Ingersoll, Andrew P. *Astrophys. J.*; 175: No. 3, 819-29 (1 Aug 1972).

Photospheric faculae near the equatorial solar limb may provide the excess brightness which Ingersoll and Spiegel showed would explain Dicke and Goldenberg's oblateness measurement. Three lines of evidence support this statement: (1) the excess emission of faculae may arise in optically thin regions, as required by the Ingersoll–Spiegel hypothesis; (2) faculae are sufficiently widespread on the solar surface to account quantitatively for the observed signal; and (3) temporal fluctuations in the expected signal due to faculae in 1966 are correlated with fluctuations in the observed signal at the 1% level. (The probability of the correlation coefficient for uncorrelated data exceeding the observed value is less than 1%.) Although this evidence clearly demonstrates that faculae make a sizable contribution to the observed oblateness signal, it does not preclude an equally sizable contribution due to true gravitational oblateness. Evidence that faculae may not be the only source of oblateness signal comes from the apparent fact that the ratio of fluctuation amplitude to mean signal amplitude is greater for the facular signal than for the observed oblateness signal. However, this difference may be due to errors in reading the photographs from which the facular signal was derived, or to differences in processing the two sets of data. A better test of the hypothesis cannot be made until the daily oblateness signals and their standard deviations are available. In any case, it appears that further data analysis will be necessary before a reliable value of the solar oblateness can be inferred. (auth)

46225 FACULAE AND THE SOLAR OBLATENESS. Dicke, R. H. (Princeton Univ., N. J.). *Astrophys. J.*; 175: No. 3, 831-5 (1 Aug 1972).

Chapman and Ingersoll (see preceding paper) have suggested that the excess solar oblateness found by Dicke and Goldenberg is wholly or largely due to the presence of faculae near the solar limb. This contrasts with the author's earlier statement based on a statistical study that contributions from faculae are unimportant. A new statistical study based on Chapman and Ingersoll's own facular function supports the author's 1970 statement. (auth)

46226 TIME-DEPENDENT IONIZATION EQUILIBRIUM AND LINE RADIATION UNDER FLARELIKE CONDITIONS. Kafatos, Minas C. (Massachusetts Inst. of Tech., Cambridge); Tucker, Wallace H. *Astrophys. J.*; 175: No. 3, 837-41 (1 Aug 1972).

The results of calculations of time-dependent ionization equilibrium and line emission are presented and compared with the values obtained under the assumption that steady-state conditions prevail. In the models considered, it is assumed that the electron density is constant ($\approx 10^8 \text{ cm}^{-3}$) and that the temperature increases by a factor of 10 from $3 \times 10^6 \text{ K}$ on time scales ranging from 100 to 300 s and decays back to $3 \times 10^6 \text{ K}$ on a time scale ranging from 600 to 1400 s. Ions of oxygen and silicon are considered, and it is found that the spectrum is softer during the rise and harder during the fall than predicted by the steady-state approximation. (auth)

46227 DENSITY OF THE SOLAR CORONA FROM OCCULTATIONS OF NP 0532. Counselman III, C. C. (Massachusetts Inst. of Tech., Cambridge); Rankin, J. M. *Astrophys. J.*; 175: No. 3, 843-56 (1 Aug 1972).

The mean electron density of the solar corona was determined from measurements of radio-frequency dispersion for pulsar NP 0532. In 1969 and 1970 mid-June, the density at $10 r_{\odot}$ was $7000 \pm 600 \text{ cm}^{-3}$, only one-half the value which others have inferred from eclipse and scintillation measurements at sunspot maximum. Between 5 and $20 r_{\odot}$, the density varied with radius as the power -2.9 ± 0.2 . Simultaneous measurements of coronal multipath scattering show that the mean density does not always increase when scattering is enhanced. In June 1971, the coronal density inferred from pulsar measurements under the south pole of the Sun was much less than in 1969 and 1970, although equatorial values did not appear to be significantly different. Coronal multipath scattering delays were generally less than one-tenth of their 1969-1970 level. (auth)

46228 STRUCTURE AND HEATING OF THE CHROMOSPHERE-CORONA TRANSITION REGION. Moore, Ronald Lee. Stanford, Calif.; Stanford Univ. (1972). 141p. University Microfilms Order No. 72-20,718.

Thesis.

A broad endeavor of solar physics is to determine from the observed radiation the structure of the solar atmosphere and the nature of the energy balance this structure implies. A basic feature of the solar atmosphere, which is still not well understood, is the chromosphere-corona transition region in which, in a height of a few thousand kilometers, the temperature increases from about 10^4 K at the top of the chromosphere to about 10^6 K at the bottom of the corona. The problem investigated in this dissertation is that of the structure and heating (or energy balance) of the transition region and the role of the transition region in the structure and heating of the atmosphere as a whole. (TSS) (Diss. Abstr. Int., B)

46229 ACCRETIONARY PROCESSES IN THE EARLY SOLAR SYSTEM: AN EXPERIMENTAL APPROACH. Kerridge, John F.; Vedder, James F. (Ames Research Center, Moffett Field, Calif.). *Science*; 177: No. 4044, 161-3 (14 Jul 1972).

Micrometer-size silicate flakes do not accrete during impacts in the velocity range 1.5 to 9.5 kilometers per second. Conventional accretionary theories for silicate bodies are applicable only to particles whose orbits are similar. Metal-silicate fractionation in the solar system may have been affected by differences in the accretionary behavior of the metal and silicate particles. (auth)

46230 CONDITIONS FOR ACCELERATION OF PARTICLES IN SOLAR ATMOSPHERE. Velinov, P. I. (Geophysical Inst., Sofia). *Compt. Rend. Acad. Bulg. Sci.*; 25: No. 1, 35-8 (1972).

Certain problems related to conditions necessary for acceleration of charged particles in the solar atmosphere are examined, particularly the effect of ionization losses, including both single and multiple ionization. This effect is essential in the range of relativistic energies, i.e., at the initial stage of acceleration when the chemical composition and spectrum of the solar cosmic rays are formed. (auth)

46231 THEORETICAL STUDIES OF THE FLUX AND ENERGY SPECTRUM OF GAMMA RADIATION FROM THE SUN. Cheng, Chung-chieh. (Goddard Space Flight Center, Greenbelt, Md.). *Space Sci. Rev.*; 13: No. 1, 3-123 (May 1972).

The various γ -ray-production mechanisms in solar flares were studied and the flux, the spectrum, and the decay curves of γ radiation were calculated. From the continuity equation, taking into account the energy losses for solar-flare-accelerated particles, the time-dependent particle distribution and the time

behavior of the resulting γ rays were obtained. The important processes for producing γ rays in solar flares are found to be nonthermal electron bremsstrahlung, decay of neutral π mesons, positron annihilation, neutron capture, and decay of excited nuclei. The results are applied to several known solar flares. For a large flare such as the class 3^+ on February 23, 1956, continuous γ rays with energies up to 100 MeV from electron bremsstrahlung and neutral π meson decays are observable at the orbit of the Earth by existing γ -ray detectors. Line γ rays from positron annihilation (0.51 MeV), neutron capture (2.23 MeV), and deexcitation of excited nuclei ^{16}O (6.14 and 7.12 MeV) and ^{12}C (4.43 MeV) are particularly strong and well above the continuous γ -ray background due to electron bremsstrahlung. These lines can be detected at the Earth. (118 references) (auth)

46232 COMPOSITION AND ENERGY SPECTRA OF HEAVY NUCLEI WITH $0.5 < E < 40 \text{ MeV}$ PER NUCLEON IN THE JANUARY 24, AND SEPTEMBER 1, 1971 SOLAR FLARES. Crawford, H. J.; Price, P. B.; Sullivan, James D. (Univ. of California, Berkeley). *Astrophys. J.*; 175: No. 3, L149-53 (1 Aug 1972).

Energy spectra of O, Si, and Fe between 0.5 and 40 MeV per nucleon were measured with stacks of plastic detectors exposed in a rocket during the January 24, 1971 flare. The Fe/Si ratio decreased from 1.8 ± 0.3 to 0.42 ± 0.14 with energy. In both the January 24 flare and the September 1 flare, the abundances of elements $10 \leq Z \leq 28$ at $\sim 15 \text{ MeV}$ per nucleon were similar to high-energy cosmic-ray source abundances; S, Ne, and Ar were significantly lower than current solar values. (auth)

46233 SPACE OBSERVATIONS OF THE SOLAR CORONA. Brueckner, Guenter E. (Naval Research Lab., Washington, D. C.). *Astronaut. Aeronaut.*; 10: No. 7, 24-30 (Jul 1972).

Coronal observations made in the 20th century are briefly reviewed. Observing techniques in the uv region developed since the early 1960's are discussed in some detail and coronal structure is considered. (W.D.M.)

Galaxies

Refer also to abstracts: 46168 and 46761.

46234 SOME CURRENT STUDIES OF GALAXIES. Hodge, Paul W. (Univ. of Washington, Seattle). *Sky Telesc.*; 44: No. 1, 23-7 (Jul 1972).

Arguments on relationships between galaxies with different redshifts presented at the Seattle meeting of the AAS, April 1972, are summarized. Papers on peculiar galaxies, infrared studies, and normal galaxies are briefly reviewed. (W.D.M.)

46235 PREDOMINANT MOTION IN SPHERICAL SYSTEMS OF GALAXIES. Karachentsev, I. (Special Astrophysical Observatory, Moscow); Zonn, W.; Shcherbanovskii, A. *Astrophys. Lett.*; 11: No. 3, 151-4 (Jul 1972).

Statistical study of kinematics of some clusters and groups of galaxies supports the hypothesis of their instability. The study is based on the examination of regression curves of radial velocities of members of a cluster on the projections (on the celestial sphere) of their distances from the center of a cluster. (auth)

46236 TWO NEW LARGE-DIAMETER GALACTIC REGIONS OF FAINT $H\alpha$ EMISSION. Courtes, G.; Sivan, J. P. (CNRS, Marseille, Observatoire, Marseille). *Astrophys. Lett.*; 11: No. 3, 159-62 (Jul 1972).

In order to detect the limit of the general emission of the $H\alpha$ line in the Milky Way, a photographic survey was undertaken with a 55° field camera using a very narrow interference filter centered on $H\alpha$. Two of the plates revealed two new large-diameter regions of faint emission at high galactic latitudes. Coordinates, sizes, exciting stars, distances, and brightnesses are given. In one of these pictures, the whole Cygnus Complex is recorded. Morphological remarks are given. (auth)

46237 RADIO MAP OF THE SPIRAL GALAXY MAFFEI 2 AT 1415 MHz. Allen, Ronald J. (Sterrenkundig Laboratorium Kapteyn, Groningen, Netherlands); Raimond, Ernst. *Astron. Astrophys.*; 19: No. 3, 317-25 (Jul 1972).

A field of one degree square centered near the infrared object Maffei 2 was observed with the Westerbork Synthesis Telescope in the radio continuum at 1415 MHz. The angular resolution of $26'' \times 31''$ was sufficient to reveal a nuclear source $19'' \pm 4''$ in diameter with a flux of 0.21 ± 0.01 flux units located near the optical center of Maffei 2. The nucleus is surrounded by a disk of emission 7.8×3.5 in angular size with a mean surface brightness of about 5° K at 1415 MHz. The radio morphology is consistent with the inference from previous infrared and radio H I

observations that Maffei 2 is a spiral galaxy. A region of increased radio intensity in the disk of the galaxy apparently coincides with a concentration of H II regions in a spiral arm. The radio continuum data would seem to favor a Hubble type of Sbc or later, and a distance of about 5 Mpc for Maffei 2. A list of incidental radio sources in the field is given, as well as the positions of eleven faint stars in the immediate neighborhood of the galaxy. (auth)

46238 EXTENDED VAN WIJK SEQUENCE IN THE LARGE MAGELLANIC CLOUD. Ardeberg, A. (European Southern Observatory, Santiago de Chile). *Astron. Astrophys.*; 19: No. 3, 384-7 (Jul 1972).

Ten stars in the van Wijk sequence in the Large Magellanic Cloud were observed extensively in the UVB system using 5 E regions for standards. In V and (B-V) the results obtained were in very good agreement with previous results. However, for the (U-B) color indices the differences were considerable. With the present data the ten stars form a very suitable local photometric standard sequence for the Large Magellanic Cloud. (auth)

46239 STUDY OF THE CONTINUUM OF THE NUCLEI OF GALAXIES. II. Andrillat, Y.; Souffrin, S.; Alloin, D. (Observatoire, Meudon, France). *Univ. Languedoc, Montpellier, France*. *Astron. Astrophys.*; 19: No. 3, 405-16 (Jul 1972).

A method previously described (Alloin et al., 1971a), based on the study of equivalent widths of absorption lines, was applied to determine the stellar population of the galactic nuclei and to compute a "synthetic" continuum that is compared with observation. Nuclei of four galaxies were studied: NGC 1052, NGC 2655, NGC 2903, NGC 4569. It was verified that the thermal continuous emission of the gas is negligible except for NGC 2903. For NGC 1052, NGC 2655, NGC 4569, it was possible to construct models of the stellar population which were consistent with the intensities of the absorption lines. These models incorporated a large proportion of K and M dwarfs (in order to account for the intensities of the Mg I and Na I lines). No intrinsic reddening appeared in the visible range (4000 to 6000 Å). For NGC 2903, the observed intensities of absorption lines could not be accounted for in terms of stellar radiation alone but must include an important contribution of nonstellar radiation in the blue (30% at 4000 Å), thus explaining the abnormal blue color of the galaxy. The presence of super-metal rich stars in the galactic nuclei would not affect the relative stellar contributions, nor the conclusion concerning the presence of nonthermal radiation in NGC 2903. However, since the synthetic continua would be "bluer," an intrinsic reddening would be needed to account for the observations. (auth)

46240 INTERSTELLAR ORGANIC CHEMISTRY. Sagan, Carl (Cornell Univ., Ithaca, N. Y.). *Nature (London)*; 238: No. 5359, 77-80 (14 Jul 1972).

Complex organic molecules that are stable against radiation may pervade interstellar space, with their degradation products a source of those molecules observed. The connection between biological and interstellar organic chemistry is, however, analogical rather than substantive, and the prospect for interstellar biology is dim. (auth)

46241 CORRELATION OF REDSHIFT WITH MAGNITUDE AND MORPHOLOGY IN THE COMA CLUSTER. Tift, W. G. (Univ. of Arizona, Tucson). *Astrophys. J.*; 175: No. 3, 613-25 (1 Aug 1972).

Redshifts and magnitudes for nearly 100 galaxies in the core of the Coma cluster are now available. The sample is nearly complete to seventeenth magnitude for the nuclear regions (central 4.8 in. spot). Nearly all the higher-redshift galaxies in the Coma cluster are nonellipticals. The separation of the ellipticals from the nonellipticals is significant in both the redshift and magnitude coordinates and is shown to be a maximum in a linear combination of redshift and magnitude along a direction sloping close to 0.6 mag fainter per 1000 km s⁻¹ increase in redshift. The morphological separation is in the same sense as the pattern in the Virgo cluster. When the redshift-magnitude diagram is examined in the direction of the maximum morphological separation, it is shown to have a strongly banded structure which is not present in any other direction. The observed redshift-magnitude pattern is compared with an outlying distinct group of galaxies in the Coma cluster which shows a nearly identical pattern slightly shifted in redshift. The pattern is also used to predict the (m, log z)-diagram for field galaxies. Observed field galaxies are shown to fit the predicted pattern and to deviate from the normal velocity-dispersion model. After allowance for magnitude and redshift uncertainties it is shown that residual true Doppler velocity dispersion in the Coma cluster cannot exceed about 220 km s⁻¹, and the cluster is not dynamically unstable. In the presence of an intrinsic unexplained redshift which produced the observed pattern, the mean

redshift of a group of galaxies need not be the same as the Hubble velocity of the cluster. Systematic trends in the deviation of the Hubble velocity from the mean redshift are possible due to selection with distance and may explain the local apparent dependence of the Hubble constant on distance. (auth)

46242 STRUCTURE OF THE COMA CLUSTER OF GALAXIES. Rood, Herbert J. (Wesleyan Univ., Middletown, Conn.); Page, Thornton L.; Kintner, Eric C.; King, Ivan R. *Astrophys. J.*; 175: No. 3, 627-47 (1 Aug 1972).

Radial velocities in the Coma cluster, many of them new, are analyzed in conjunction with previous data on the density distribution. The cluster extends more than 200' in radius, with bright and faint galaxies distributed nearly the same. Some spirals are members. Several different determinations of M/L give values around 250. The cluster is stabilized by a "missing mass" some 7 times the mass of the galaxies, which must be distributed in the same way as the galaxies. (auth)

46243 RADIAL CYANOGEN-BAND GRADIENTS NEAR THE NUCLEI OF GALAXIES. Spinrad, Hyron (Univ. of California, Berkeley); Smith, Harding E.; Taylor, Donald J. *Astrophys. J.*; 175: No. 3, 649-57 (1 Aug 1972).

Very high spatial-resolution area scans of the nuclear regions of eight nearby galaxies, of types E0-E5, S0, Sa, and Sb were obtained in 1970-1971 with 33-Å spectral purity at the blue λ 4200 CN band. Partially confirming and extending previous work of McClure and Spinrad et al., all galaxies scanned to show either zero or negative radial CN-band-strength gradients. The strength of the CN gradient along the major axis was found to be inversely correlated with the degree of flattening of the system. Also, scans of NGC 3115 shows a substantial CN decrease along the minor axis while the CN strength is constant along the major axis. The interpretations of the CN gradients are briefly discussed in terms of differences in chemical composition relating to the dynamical histories of simple spherical and flattened systems. (auth)

46244 ABUNDANCES OF INTERSTELLAR MOLECULES. Watson, W. D.; Salpeter, E. E. (Cornell Univ., Ithaca, N. Y.). *Astrophys. J.*; 175: No. 3, 659-71 (1 Aug 1972).

Equilibrium abundances were calculated for various molecules in interstellar H I gas clouds using rates from a previous paper for molecule formation on grain surfaces, and rates from other authors for gas-phase reactions and for photodissociation. Results depend on a parameter $\xi \sim (n/100 \text{ H cm}^{-3}) \exp(2.5\tau_v)$, where n is gas density and τ_v is the visual optical depth from points inside the cloud to its surface. Molecules probably are not depleted by condensation if $\xi \lesssim 10^4$ (but the situation is uncertain if grains with ice mantles are present). For slightly shielded clouds ($\xi \lesssim 10$) where hydrogen is mainly atomic and carbon is mainly C⁺, the combined abundances of various molecules containing C, N, or O were estimated (relative to atomic C, N, and O) to be $\sim 10^{-4}\xi$. When ξ exceeds some critical value ~ 300 , the neutral C in the gas eliminates photons beyond 11.3 eV; because of their high binding energy, CO molecules are then not dissociated and are expected to be almost as abundant as C atoms. Abundances (with even greater uncertainties) are also estimated for the more complex molecules. (45 references) (auth)

46245 FORMATION OF CLOUDS IN A COOLING INTERSTELLAR MEDIUM. Schwarz, Joseph (Harvard Coll. Observatory, Cambridge, Mass.); McCray, Richard; Stein, Robert F. *Astrophys. J.*; 175: No. 3, 673-86 (1 Aug 1972).

An interstellar medium cooling from 8200°K after sudden heating (e.g., by soft x rays or by low-energy cosmic rays) favors the development of thermal condensations. One-dimensional hydrodynamic calculations show that the condensation process will yield interstellar clouds with densities of ~ 100 times the intercloud density, given a small ($\sim 10\%$) initial density perturbation, provided the scale length λ of the initial perturbation satisfies $\lambda_c \lesssim c\tau_c$, where c is the sound speed in the gas and τ_c is the radiative cooling time. The condensations reach maximum density in a few cooling timescales. For an initially uniform medium of density $n_H = 0.3$ hydrogen atoms per cm³ that is 5% ionized, a density perturbation of wavelength 3 pc will grow by a factor of 100 in about 10^6 years, reaching a final ionized fraction $x \approx 10^{-3}$ and temperature $T \lesssim 20^\circ\text{K}$. The intercloud medium at this time will have cooled to about 2000°K. In the one-dimensional case, the final extent of the cloud will be about 0.01 pc. Cloud dimensions more typical of those observed in the interstellar medium are a likely result of a three-dimensional treatment of this problem. An increase in the critical condensation wavelength λ_c , as well as higher temperatures for the time-dependent intercloud medium, will result if cooling agents such as C, Si, Fe, and O are more highly ionized than assumed in these calculations. Soft x rays are a plausible source of such anomalously high ionization. (auth)

- 46246** STUDY OF NEUTRAL HYDROGEN GAS MOTION WITHIN SPIRAL ARMS AND IN THE LOCAL REGION. Harten, Ronald Harold. College Park, Md.; Univ. of Maryland (1972). 132p. University Microfilms Order No. 72-20,492. Thesis.

Data from several recent high resolution 21-cm line studies was analyzed in a two-part study of the motions of the neutral-hydrogen gas within spiral-arm features. This study considers particularly the structure of the gas motions perpendicular to the galactic plane. (TSS) (Diss. Abstr. Int., B)

- 46247** HIGH-FREQUENCY RESOLUTION H I LINE OBSERVATIONS OF INTERSTELLAR DUST CLOUDS. Knapp, Gillian Revill. College Park, Md.; Univ. of Maryland (1972). 263p. University Microfilms Order No. 72-20,253. Thesis.

The density and temperature of atomic hydrogen in dense interstellar dust clouds were investigated by searching for narrow self-absorption dips in the 21-cm emission profiles in the directions of such clouds. The observations were made with the 140-foot telescope of the National Radio Astronomy Observatory. It was found that about half of the dust clouds observed showed self-absorption dips and that the greater the observed optical density of the clouds the greater the chance of their showing up in self-absorption. (TSS) (Diss. Abstr. Int., B)

- 46248** LUMINOSITY-SPECTRAL INDEX RELATIONSHIP FOR RADIO GALAXIES. Macleod, J. M.; Doherty, L. H. (National Research Council of Canada, Ottawa). Nature (London); 238: No. 5359, 88-9(14 Jul 1972).

It is stated that the relationship between spectral index and luminosity for the 3CR extragalactic sources suggests that sources with high radio luminosity have, on the average, steeper spectra than do low-luminosity sources, but the scatter in the relationship is quite large. Furthermore, correlation exists only for radio galaxies and not for quasars. It is shown here that, even for radio galaxies, the relationship holds only for a particular subgroup in the 3CR catalog. For those galaxies with power-law spectra there is a clear relationship between luminosity and spectral index, but for those galaxies that do not have simple power-law spectra luminosity is independent of spectral index. (UK)

- 46249** RELATIONSHIP BETWEEN X-RAY LUMINOSITY AND VELOCITY DISPERSION IN CLUSTERS OF GALAXIES. Solinger, Alan B.; Tucker, Wallace H. (American Science and Engineering, Cambridge, Mass.). Astrophys. J.; 175: No. 3, L107-11(1 Aug 1972).

The connection between the x-ray luminosity L_X and the dynamics of clusters of galaxies is considered. L_X is found to be strongly correlated with the velocity dispersion ΔV . The observed relationship can be explained most simply within the framework of a thermal-bremsstrahlung model with the x-ray luminous mass being a fraction of the cluster binding mass. This implies that the clusters are or were gravitationally bound. The x-ray intensities or velocity dispersions are predicted for several clusters. (auth)

- 46250** BRIGHT NEBULAE NEAR CONCENTRATIONS OF HIGH-VELOCITY GAS. Minkowski, R. (Univ. of California, Berkeley); Silk, Joseph; Siluk, Richard S. Astrophys. J.; 175: No. 3, L123-5(1 Aug 1972).

Some coincidences in position on the sky between apparent emission nebulae at $|b_{II}| > 10^\circ$ and concentrations of high-velocity gas ($|v| > 100 \text{ km s}^{-1}$) are reported. (auth)

- 46251** KINEMATICS OF MOLECULAR CLOUDS NEAR THE GALACTIC CENTER. Scoville, N. Z. (Columbia Univ., New York). Astrophys. J.; 175: No. 3, L127-32(1 Aug 1972).

A model is proposed in which most of the molecular clouds near the galactic center are situated in a radially moving ring of radius 250 pc. (auth)

- 46252** 21 cm OBSERVATIONS OF NGC 45. Lewis, B. M. (Australian National Univ., Canberra). Aust. J. Phys.; 25: No. 3, 315-28(Jun 1972).

Observations of NGC 45 were made at a wavelength of 21 cm using the radio telescope at Parkes. Line profiles were measured for a grid of 44 points spaced at intervals of $6'$ arc. From the measurements the mass of neutral hydrogen was calculated to be $8.2 \times 10^8 M_\odot$, assuming a distance of 3 Mpc. A simple self-consistent model was determined by computing line profiles to compare directly with the observations. This enabled the rotation curve to be corrected for the first-order effects of beam smoothing and resulted in a total limiting mass of $2.5 \times 10^{10} M_\odot$. (auth)

- 46253** APPROXIMATE SOLUTIONS OF THE RELATIVISTIC GRAVITATIONAL FIELD EQUATIONS TO DESCRIBE CLUSTERS OF GALAXIES. Cook, M. W. (Univ. of New England, Armidale, Australia). Aust. J. Phys.; 25: No. 3, 299-305(Jun 1972).

Approximate solutions to the Einstein field equations are found that describe a spherically symmetric inhomogeneity in a general Robertson-Walker model, i.e., one with an arbitrary equation of state. The approximation hypothesis is that the pressure deviates only slightly from uniformity, and it is found that the density may have quite large local fluctuations, e.g., by a factor of 10^6 over a region 10^{-2} Mpc in diameter. Reference is made to observed data to determine which categories of stellar objects may be described by the results. (auth)

Cosmology

Refer also to abstracts: 46596, 46749, 46755, 46758, 46759, and 46761.

- 46254** (COO-1112-51) TIME VARIATION OF THE FINE STRUCTURE CONSTANT AND SPECTRAL LINES FROM QUASISTELLAR OBJECTS. Tomozawa, Yukio (Michigan Univ., Ann Arbor, Harrison M. Randall Lab. of Physics). Jun 1970. 7p. Dep. NTIS.

It has been suggested that the gravitational constant (G) is likely to be an increasing function of time (t) in the model of an expanding universe. Assuming the $G \propto e$, the relation $\alpha^{-1} = \frac{3}{2} \pi \ln 1/t + \text{constant}$ is obtained, i.e., the fine structure constant (α) is slowly increasing in time. It is pointed out that a more accurate measurement of quasar spectra is needed to test the relation. (W.D.M.)

- 46255** MEASUREMENT OF THE ISOTROPY OF THE MICROWAVE BACKGROUND TEMPERATURE AT 6-cm WAVELENGTH. Davis, Karl C. Princeton, N. J.; Princeton Univ. (1971). 145p. University Microfilms Order No. 72-13,737. Thesis.

A survey of the sky brightness was made at 6-cm wavelength for the purpose of measuring the isotropy of the cosmic background radiation. The instrument used had an angular resolution of approximately 5 and measured the apparent sky brightness temperature difference between regions centered on the celestial pole and on points southward to -9.5 declination. The instrument was operated 24 hours a day for approximately one year, thus covering all values of right ascension. Galactic radio emission at 6 cm was estimated using a 75-cm sky survey and a temperature spectral index of -2.8 . The results of the 6 cm survey, after subtraction of galactic emission, were examined for several possible anisotropies. Because of a systematic error related to movement of the measuring beam in declination all information about anisotropies having the celestial pole as a symmetry axis was lost. An anisotropy with 24-hour spatial period with amplitude in the equatorial plane of 2.5 ± 1.5 milliKelvin (corresponding to a velocity of the local center of rest with respect to uniform expansion of the Universe of $250 \pm 150 \text{ km/s}$) with direction 10 ± 2 hours right ascension was observed. This result is in agreement with previous measurements at 3-cm wavelength. An anisotropy with 12-hour spatial period was observed but was found to depend strongly on a small fraction of the data near the galactic center and is believed not to be a feature of the cosmic background. The data in regions removed from the galactic center was examined for areas of angular diameter 15 and 30 with anomalous temperature. No evidence was found for such regions. Upper limits on the maximum departure from isotropy of 37 milliKelvin for 15 areas and 16 milliKelvin for 30 areas are given with 90% confidence for the area of the sky covered (47% of the total sphere). Details of the experiment and data analysis are given with suggestions for future isotropy measurements. (Diss. Abstr. Int., B)

- 46256** OBSERVATIONAL COSMOLOGY. Longair, M. S. (Mullard Radio Astronomy Observatory, Cambridge, Eng.). Rep. Progr. Phys.; 34: No. 12, 1125-1248(1971).

While a strong case can be made for the hot model of the universe, many other theories have been proposed which would drastically change this picture. The status of a number of these hypotheses, including alternative theories of gravitation, noncosmological redshifts, steady-state theory, and variations of the constants of nature with epoch, are discussed in the light of recent observations. (auth)

- 46257** HOT UNIVERSE, COSMIC RAYS OF ULTRAHIGH ENERGY, AND ABSOLUTE REFERENCE SYSTEM. Sato, Humitaka (Kyoto Univ.); Tati, Takao. Progr. Theor. Phys. (Kyoto); 47: No. 5, 1788-90(May 1972).

Interaction of the 3°K cosmic thermal radiation with cosmic particles with $E > 10^{20}$ eV should lead to an attenuation mean-free-path as small as $10^{8.0} \sim 10^{7.3}$ parsec. Analysis of a large air shower has revealed that the energy spectrum of cosmic rays extends smoothly beyond 10^{20} eV. Therefore, the nonexistence of the expected cutoff in the vicinity of 10^{20} eV offers a critical discrepancy between theory and observation. A way is proposed to get rid of this discrepancy. (W.D.M.)

46258 PHYSICAL ASSOCIATIONS BETWEEN QUASI-STELLAR OBJECTS AND GALAXIES. Burbidge, G. R. (Univ. of California, San Diego); O'Dell, S. L.; Strittmatter, P. A. *Astrophys. J.*; 175: No. 3, 601-11 (1 Aug 1972).

Further evidence is presented relevant to apparent associations between galaxies and quasi-stellar objects. It is shown that for the five 3C QSOs which lie very close to bright galaxies the QSO-galaxy angular separations are inversely proportional to the redshifts of the galaxies. This lends additional support to the association hypothesis. However, an analysis of an essentially disjoint sample of QSOs identified from the Parkes radio catalog, and fainter galaxies in the catalog of Zwicky and his associates, reveals no statistically significant tendency toward pairing of QSOs and galaxies. The results can be explained in one of two ways. One is that, despite the evidence to the contrary, the 3C results are due to chance. The other is that real associations in the second sample are masked by selection effects. It is shown that several selection biases do indeed discriminate against the detection of QSO-galaxy pairs, especially for fainter objects. Finally, on the assumption that the 3C QSO-galaxy associations are real, the relationship of QSOs to galaxies is discussed. (auth)

46259 MASS DIFFERENCE OF THE MUON AND THE ELECTRON. Hoyle, F.; Narlikar, J. V. (Cambridge Univ., Eng.). *Nature* (London); 238: No. 5359, 86-7 (14 Jul 1972).

It is stated that the fact that the muon and the electron have different masses can be understood in terms of a Machian theory of inertia. The masses of particles are not fixed and immutable but depend on the detailed particle composition of the Universe. (auth)

46260 CAN THE LUMPY DISTRIBUTION OF GALAXIES BE DETECTED BY X-RAY OBSERVATIONS. Webster, A. S. (Univ. of California, Berkeley). *Nature* (London); 238: No. 5358, 20-4 (7 Jul 1972).

It is stated that published data from x-ray telescope observations are not adequate to reveal the fluctuations in the surface brightness of the background radiation caused by clustering or superclustering of the sources giving rise to the background. (auth)

46261 HOMOGENEOUS ROTATING UNIVERSE WITH FLAT SPACE. Demianski, M. (Univ., Warsaw); Grishchuk, L. P. *Commun. Math. Phys.*; 25: No. 3, 233-44 (1972).

The homogeneous, anisotropic cosmological model is considered. It satisfies three physically reasonable conditions: it is space homogeneous, possesses flat space-like sections, and is filled with expanding, rotating, and shearing matter. The asymptotic solution is presented and general properties are discussed. The question how the rotation influences the behavior of matter near the singularity is investigated. (auth)

Planetary Phenomena

Refer also to abstract: 47016.

46262 POSSIBLE PRODUCTION MECHANISMS OF LUNAR MAGNETIC FIELDS. Cap, Ferdinand F. (Goddard Space Flight Center, Greenbelt, Md.). *J. Geophys. Res.*; 77: No. 19, 3328-33 (1 Jul 1972).

The impossibility of the production of local surface magnetic fields on the Moon by conduction currents in the lunar soil and in local lunar atmospheres by volcanic eruption is shown. However, it is suggested that convection currents produced by the ionization (by radiation and/or by triboelectric effects) of volcanic-ash-particle flows may produce the local magnetic fields of about 1000γ that are believed to have existed on the Moon about 3.5×10^5 years ago. A simple electrodynamic model for such flows and experiments for further investigation of this hypothesis are discussed. (30 references) (auth)

PHYSICS (ATMOSPHERIC)

Auroral and Ionospheric Phenomena

46263 (AD-734392) METHOD FOR RAPID COMPUTATION OF VLF, STEEP-INCIDENCE REFLECTIONS FROM THE IONOSPHERIC D REGION. Research and Development Report, October 1970-June 1971. Hill, J. R. (Naval Electronics Lab. Center, San Diego, Calif.). 19 Aug 1971. 25p. (NELC-TR-1784. NTIS.

A method for the rapid computation of coefficients for vlf reflections from the ionospheric D region is described. Previously calculated wave-field solutions are used to compute reflection coefficients for new electron-density profiles. The usefulness of the method is demonstrated by a study of several D-region profiles. The study shows the sensitivity of the reflection coefficients to small changes in the profiles. Experience indicates that three to six times as many profiles can be examined in a parametric study than was previously possible for a given computer cost. The resulting increase in computational speed greatly facilitates parametric studies of electron-density profiles. (auth)

46264 (COO-2195-5) CORRELATION OF D-REGION ABSORPTION AND THE 10 mb WINDS AND TEMPERATURE DURING THE IQSY. Wu, Mao-Pou; Newell, Reginald E. (Massachusetts Inst. of Tech., Cambridge, Dept. of Meteorology). [nd]. Contract AT(30-1)-2241. 18p. Dep. NTIS.

Relations between absorption and some meteorological parameters were studied for the entire period of the IQSY. The method is to calculate the correlation coefficient between the D-region hf radio-wave absorption at local noon and 10-mb level winds and temperature at 12 Z for nine stations in the Northern Hemisphere. Results show that the absorption correlates negatively with the zonal wind (positive toward east), but positively with the temperature for the majority of cases, and that the meridional wind (positive toward north) is correlated with absorption in the winter season. It is suggested that the northward transport of atomic oxygen by large-scale eddies is related to the winter D-region abnormal absorption and some correlations with airglow data are presented to support this suggestion. (auth)

46265 (N-72-18715) OGO-1 AND OGO-3 MIT PLASMA EXPERIMENTS S4903. Final Project Report. (Massachusetts Inst. of Tech., Cambridge, Center for Space Research). 30 Jun 1968. Contract NAS5-2053. 208p. (NASA-CR-122351). NTIS.

Plasma proton and plasma electron prototype and flight models were designed, fabricated, and tested. Ground-support equipment for the models was also prepared. The flight models were launched aboard the first and third Orbiting Geophysical Observatories on Sept. 4, 1964 and June 6, 1966. These experiments have generally functioned in accordance with the design specifications and useful data are still being received. (STAR)

46266 CRITICAL COMPONENT OF THE INTERPLANETARY MAGNETIC FIELD RESPONSIBLE FOR LARGE GEOMAGNETIC EFFECTS IN THE POLAR CAP. Friis-Christensen, E. (Danish Meteorological Inst., Charlottenlund, Denmark); Lassen, K.; Wilhjelm, J.; (and others). J. Geophys. Res.; 77: No. 19, 3371-6 (1 Jul 1972).

An observed influence of the interplanetary magnetic-sector structure on the geomagnetic variations in the polar cap appears to be due to the component of the interplanetary magnetic field near the ecliptic perpendicular to the Earth-Sun direction. This suggests that the observed effect on the ground originates in the front of the magnetosphere. (auth)

46267 FERMI ACCELERATION OF AURORAL PARTICLES. Sharber, J. R.; Heikkila, W. J. (Univ. of Texas, Dallas). J. Geophys. Res.; 77: No. 19, 3397-3410 (1 Jul 1972).

A number of nighttime acceleration mechanisms proposed in the literature are reviewed and rejected for the role of producing the key nighttime auroral-particle fluxes. Parallel electric fields are rejected for several reasons, but particularly because of the observed simultaneous precipitation of electrons and protons. Acceleration in the neutral sheet is inadequate for producing the particle energies, the observed field-aligned pitch-angle distribution at high latitudes, and the spectral hardening toward lower latitudes. Neutral-point mechanisms, although often suggested in principle, have never been demonstrated satisfactorily in theory or in practice. Pitch-angle scattering from a trapped population produced by transverse adiabatic compression is also incapable of producing the field-aligned distribution. Possibly longitudinal or Fermi acceleration, which results from the known magneto-

spheric convection, is the main nighttime auroral acceleration mechanism. The argument is supported by data obtained with the soft-particle spectrometer of Isis 1. In particular, the rotational scanning provided by the spinning satellite shows the regular change from a field-aligned pitch-angle distribution at high latitudes to a trapped distribution at low latitudes, reflecting a change from Fermi acceleration on the distant taillike field lines to betatron acceleration on the more dipolar field lines closer to the Earth. The operation of the Fermi mechanism requires multiple bounces between hemispheres, and therefore a loss cone must develop at both 0° and 180°; such distribution is shown by the Isis 1 data when the range of the pitch-angle scan is sufficiently wide. One other feature is evident in the data: a hardening of the spectrum at small pitch angles. This energy-dependent pitch-angle distribution has a characteristic signature on the energy-time spectrograms that is designated as a Λ structure; when both loss cones show, the characteristic signature is a topless Λ structure. Adiabatic compression thus plays a vital role in auroral and magnetospheric substorms; longitudinal or Fermi acceleration causes auroral-particle precipitation, whereas transverse or betatron acceleration is responsible for lower-latitude phenomena, such as the ring current. (45 references) (auth)

46268 OBSERVATIONS OF THE AURORAL OVAL BY THE ALASKAN MERIDIAN CHAIN OF STATIONS. Snyder, A. L.; Akasofu, S.-I. (Univ. of Alaska, College). J. Geophys. Res.; 77: No. 19, 3419-30 (1 Jul 1972).

The Alaskan meridian chain of stations "scans" the polar sky like an azimuth scan radar once a day between the corrected geomagnetic latitudes of 60° and 85°N. The all-sky photographs from the chain of stations have provided and clarified several important features of auroral morphology, in particular the dynamics of the night-sector auroral oval. Results are summarized. (41 references) (auth)

46269 THERMOSPHERIC MOLECULAR OXYGEN FROM SOLAR EXTREME-ULTRAVIOLET OCCULTATION MEASUREMENTS. Roble, R. G.; Norton, R. B. (National Center for Atmospheric Research, Boulder, Colo.). J. Geophys. Res.; 77: No. 19, 3524-33 (1 Jul 1972).

The molecular-oxygen number density distribution in the 95- to 120-km region of the atmosphere has been determined from solar-occultation measurements made by the Solrad-8 satellite. The solar intensity in the region 1040 to 1350 Å was monitored by a photometer aboard the satellite during occultation of the Sun by the Earth. The bulk of the energy in the wavelength interval lies in Lyman α , which is primarily absorbed by molecular oxygen. Therefore the solar-occultation measurements are directly inverted to give the number density distribution of molecular oxygen at the occultation tangent point. The data from 93 occultation scans obtained at midlatitudes and high latitudes in both hemispheres from December 1, 1966, to August 31, 1967, are presented. The results show a seasonal variation of molecular oxygen in the lower thermosphere. In general, the molecular-oxygen number density distribution in the summer is approximately 3 times greater than that in the winter. (47 references) (auth)

46270 GEOMAGNETIC EFFECT ON THE NEUTRAL TEMPERATURE OF THE F REGION DURING THE MAGNETIC STORM OF SEPTEMBER 1969. Blamont, J. E.; Luton, J. M. (CNRS, Verrieres-le-Buisson, France). J. Geophys. Res.; 77: No. 19, 3534-56 (1 Jul 1972).

A spherical Fabry-Perot interferometer on Ogo 6 (launched June 5, 1969) gives the profile of the 6300-Å oxygen line in the dayglow at different altitudes between 200 and 320 km. The temperature of the neutral components of the atmosphere is deduced from these measurements with an accuracy of $\pm 65^\circ\text{K}$. The results presented correspond to the storm period September 25 to October 5, 1969. During this period a severe geomagnetic storm occurred. The temperature measurements show a very strong effect in the two polar regions, while the equatorial regions present a smooth temperature variation. (auth)

46271 ANALYSIS OF OGO 6 OBSERVATIONS OF THE O I 5577-Å TROPICAL NIGHTGLOW. Thomas, R. J.; Donahue, T. M. (Univ. of Pittsburgh). J. Geophys. Res.; 77: No. 19, 3557-65 (1 Jul 1972).

Atomic oxygen green-line data from the horizon scanning photometer on Ogo 6 were examined. Unfolding the satellite data from the tropical F region yields altitude and latitude variations of the O(I) emissions. The spatial variations of the tropical F-region electron density are then calculated by assuming dissociative re-

combination and using a model atmosphere. Where comparisons to ground-based data are possible, the results are good. Thus, the satellite observations constitute a form of topside sounding of the ionosphere below the F peak and provide synoptic data about this portion of the ionosphere that are otherwise impractical to obtain. (auth)

46272 COLLISIONAL LOSSES IN A VERY-LOW FREQUENCY DUCT ASSOCIATED WITH THE LOWER-HYBRID-RESONANCE FREQUENCY. Gross, S. H. (Polytechnic Inst. of Brooklyn, Farmingdale, N. Y.). *J. Geophys. Res.*; 77: No. 19, 3589-91 (1 Jul 1972).

Calculations of loss in a multi-ion horizontal duct that may trap and guide vlf waves indicate that losses are small for some configurations and not for others. Under appropriate conditions signals can be guided horizontally for hundreds of kilometers without excessive attenuation. (auth)

46273 NUMERICAL EVALUATION OF CHAPMAN'S GRAZING INCIDENCE INTEGRAL CH (X, chi). Smith III, F. L.; Smith, Cody (Colorado State Univ., Fort Collins). *J. Geophys. Res.*; 77: No. 19, 3592-7 (1 Jul 1972).

A formula that is both numerically convenient to use and accurate to within 2% for all χ when $X > 50$ may be used to approximate $ch(X, \chi)$. The approximation may be applied to optical-depth calculations in nonexponential models of the terrestrial atmosphere, the resulting errors still remaining smaller than 4% for most ionospheric applications. (auth)

46274 RELATIONSHIP BETWEEN Pc 1 MICROPULSATIONS AND IONOSPHERIC SPREAD F. Fraser-Smith, A. C. (Stanford Univ., Calif.). *J. Geophys. Res.*; 77: No. 19, 3602-6 (1 Jul 1972).

Simultaneous micropulsations and ionosonde records obtained on the Stanford campus were compared for the 18-month interval from Jan. 1, 1970 through June 30, 1971 to study the relationship between spread F and the Pc 1 micropulsations. The 73 simultaneous observations were made at night. There was evidence for both an increase in the frequency of occurrence and in the strength of ground-recorded Pc 1 when spread F was present in the overhead ionosphere. A tendency was present for Pc 1 to occur most often in the time interval most favorable for spread F, although no correlation was found between the observed durations of associated activity. During the 18-month interval, occurrences of Pc 1 and spread F reached peaks simultaneously at 0100 to 0400 LST. Possible evidence for a correlation between Pc 1 and spread F also is provided by long-term observations made separately on their solar-cycle variations. (J.C.W.)

46275 SATELLITE AND GROUND-BASED OBSERVATIONS OF A RED ARC. Nagy, A. F. (Univ. of Michigan, Ann Arbor); Hanson, W. B.; Hoch, R. J.; Aggson, T. L. *J. Geophys. Res.*; 77: No. 19, 3613-17 (1 Jul 1972).

Simultaneous observations were performed during the red-arc period of Aug. 8-9, 1970. Photometric observations were made near Richland, Washington and Ogo 6 obtained data in the southern hemisphere. The formation of the arc at the electron-temperature peak and density trough supports the thermal-conduction theory of red-arc formation. The lack of the dc electric fields in the red-arc region ruled out dc electric fields as an excitation mechanism; particle precipitation as a direct source of the red-line emission can also be ruled out by the observations made. The formation of the arc in the plasmopause region and the detection of low-frequency ac fields are consistent with a theory that predicts wave-particle interactions at the plasmopause as a mechanism for supplying thermal energy to the plasmasphere electrons. (J.C.W.)

46276 $H\beta/N_2^+$ 4709-A INTENSITY RATIO AND VIBRATIONAL ENHANCEMENT OF N_2^+ FIRST NEGATIVE BANDS IN PROTON-EXCITED AURORAS. Degen, V.; Belon, A. E.; Romick, G. J. (Univ. of Alaska, Fairbanks). *J. Geophys. Res.*; 77: No. 19, 3618-20 (1 Jul 1972).

The N_2^+ first negative band system was examined for a possible vibrational enhancement that may be expected to occur in auroras excited by low-energy (<10 keV) protons. Vibrational enhancement has been well documented only in certain large auroras, but the role of protons in inducing anomalous vibration is still controversial because of possible contamination of the spectra by O II and other uncertain excitation conditions. Observational data are discussed concerning the faint aurora, also known as the hydrogen arc. Theoretical estimates made by Omholt (1971) and based on uncertain cross sections give $H\beta/N_2^+$ 4709-A intensity ratios in a pure proton-excited aurora ranging from 1.25 for protons with an initial energy of 100 keV to about 10 for 3-keV protons. Eather's 1967 earlier calculations gave ratios of 1 to 2.5 for this energy range. A search by Eather (1968) for maximum ratios in night-side auroras using a multichannel tilting filter photometer led to a statistical value for a maximum ratio of about 1.1 to 0.8. Eather

suggests a ratio of about 1 as a working criterion for a nighttime proton-induced aurora. Such auroras are subvisual or marginally visual. Ratios as high as 3 have been reported in extremely faint (<10 rayleighs in $H\beta$) dayside auroras. (J.C.W.)

46277 NONLINEAR SATURATION OF TYPE-I IRREGULARITIES IN THE EQUATORIAL ELECTROJET. Weinstock, Jerome; Sleeper, Arthur (National Oceanic and Atmospheric Administration, Boulder, Colo.). *J. Geophys. Res.*; 77: No. 19, 3621-4 (1 Jul 1972).

A nonlinear theory is presented for the saturation of the type-1 equatorial E-layer instability. Saturation is achieved by the transfer of wave energy to electrons (electron trapping). The irregularity is found to be about 2% of the background density. Formulas are derived for both the particle density and the electric field of the irregularity. (auth)

46278 DETERMINATION OF THE QUENCHING OF $O(^1D)$ BY MOLECULAR NITROGEN USING THE IONOSPHERIC MODIFICATION EXPERIMENT. Hernandez, G. (National Oceanic and Atmospheric Administration, Boulder, Colo.). *J. Geophys. Res.*; 77: No. 19, 3625-9 (1 Jul 1972).

The collisional frequency of quenching of $O(^1D)$ by molecular nitrogen is derived from measurements of the lifetime of O I 6300-A emission during the transient disturbances created by the ionospheric modification experiment (IME). The mean value obtained for the frequency of quenching (at 120 km) is 18 sec^{-1} , the spread being 17 to 20 sec^{-1} . The corresponding reaction rate coefficient is $9(\pm 2) \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$, based on the model atmosphere used, and is in excellent agreement with recent laboratory and aeronomic determinations. (auth)

46279 INVESTIGATION OF THE IONOSPHERIC D REGION AT SUNRISE. I. TIME VARIATIONS OF OZONE, METASTABLE MOLECULAR OXYGEN, AND ATOMIC OXYGEN. Turco, R. P.; Sechrist, C. F. Jr. (Univ. of Illinois, Urbana). *Radio Sci.*; 7: No. 7, 703-16 (Jul 1972).

As a step in determining the behavior of the D-region ionization at sunrise, the development of atomic oxygen and ozone over sunrise was investigated, taking into account the absorption of solar ultraviolet radiation by the time-varying, nonuniform ozone distribution over the mesosphere at sunrise. The dissociation rates of ozone and molecular oxygen are calculated and used to determine the time variations in the O and O_3 densities at sunrise for a model considering only the oxygen allotropes. Transport processes were neglected in the sunrise calculations because chemical time constants are shorter than those for transport. However, the assumed initial presunrise atomic oxygen and ozone distributions, based on an average of theoretical and experimental results, do include the effects of transport. It was found that at altitudes from 60 to 80 km there is a delay in the ozone sunrise, and thus a delay in the development of the atomic oxygen density. $[O]/[O_3]$ was computed during sunrise, and a similar delay in its sunrise increase was observed. The production of metastable oxygen $O_2(^1\Delta_g)$ by the photodissociation of ozone in the Hartley band at sunrise was also investigated. A simplified chemical scheme for $O_2(^1\Delta)$ is presented, from which a closed-form solution for the $O_2(^1\Delta)$ densities over sunrise was obtained. $O_2(^1\Delta)$ may be important in the ion chemistry at sunrise, and can contribute to the early appearance of electrons in the D region. (auth)

46280 INVESTIGATION OF THE IONOSPHERIC D REGION AT SUNRISE. II. ESTIMATION OF SOME PHOTODETACHMENT RATES. Turco, R. P.; Sechrist, C. F. Jr. (Univ. of Illinois, Urbana). *Radio Sci.*; 7: No. 7, 717-23 (Jul 1972).

The photodetachment rates of several negative ions believed to be present in the D region are estimated for sunrise conditions. The photodetachment cross sections of O^- and O_2^- which have been measured in the visible and infrared wavelength ranges are used. Since the photodetachment cross sections of NO_2^- and NO_3^- are not yet known, the behavior of their rates could only be estimated. Infrared band absorption by H_2O and CO_2 is included in the calculations. For atoms and molecules with large electron affinities, the ozone ultraviolet absorption is important. The effects of photodetachment on D-region behavior are discussed and found to be relatively unimportant. (auth)

46281 INVESTIGATION OF THE IONOSPHERIC D REGION AT SUNRISE. III. TIME VARIATIONS OF NEGATIVE-ION AND ELECTRON DENSITIES. Turco, R. P.; Sechrist, C. F. Jr. (Univ. of Illinois, Urbana). *Radio Sci.*; 7: No. 7, 725-37 (Jul 1972).

The growth over sunrise of the C and D layers of the ionosphere is investigated for a model which includes: the negative ion species O^- , O_2^- , O_3^- , O_4^- , NO_2^- , NO_3^- , CO_2^- , and CO_4^- ; ion-pair production by galactic cosmic rays, precipitated electrons, photoionization of NO by scattered Lyman- α radiation, and the photoionization of NO and

$O_2(^1\Delta)$ by direct solar radiation; the photodetachment of most of the negative ions; and the time variations of all these parameters. It is found that the inclusion of the reaction $NO_3^- + O \rightarrow NO_2 + O_2^-$ with a rate constant of $5 \times 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$ in combination with fast electron-ion recombination ($\alpha \sim 10^{-5} \text{ cm}^3 \text{ sec}^{-1}$) can accurately reproduce both the development of the C layer at sunrise and the correct magnitude and form for the D-layer development. The photodetachment of NO_3^- is not a suitable alternate source of sunrise electrons because of the large electron affinity of NO_3 . The photodetachment of lower-affinity negative ions, such as hydrated species, could replace the $NO_3^- + O \rightarrow NO_2 + O_2^-$ reaction if indeed the presunrise mesospheric ions are predominantly hydrated. The dominant ions in the present scheme are NO_3^- , CO_3^- , and CO_2^- . Atomic oxygen greatly limits the quantity of negative ions above 75 km at all times of the day by associative detachment and is responsible for the night-time D-layer ledge of electron density. The 98 and 94° VLF radio propagation effects are explicable on the basis of this model. (auth)

46282 SYNTHESIS OF JICAMARCA DATA DURING THE GREAT STORM OF MARCH 8, 1970. Woodman, Ronald F. (Jicamarca Radar Observatory, Lima); Sterling, D. L.; Hanson, W. B. *Radio Sci.*; 7: No. 7, 739-46 (Jul 1972).

Continuous measurements of electron temperature and electron concentration were made at Jicamarca from 0700 to 0830 LT and from 1130 to 2000 LT during the great magnetic storm of March 8, 1970, when the F2 peak rose as high as 800 km. Continuous measurements of the vertical component of $E \times B$ drift were made from 1130 to 1730 LT. A synthesis of the observed electron concentrations was made using a model that solved the time-dependent electron-ion continuity equation for O^+ and the molecular ions; the latter chemistry was included to improve the solutions at lower heights. It was found, using the observed values of vertical drift, that most of the observed features of the F2 region ionosphere were synthesized by the model. An interpretation of the synthesis that emphasizes the importance of horizontal diffusion at the equator is given. Observations are presented for March 7, 8, and 9, 1970, including the Huancayo magnetograms, which display a close correlation with the drift measurements. (auth)

46283 CHATANIKA, ALASKA, AURORAL-ZONE INCOHERENT-SCATTER FACILITY. Leadabrand, R. L. (Stanford Research Inst., Menlo Park, Calif.); Baron, M. J.; Petriceks, J.; Bates, H. F. *Radio Sci.*; 7: No. 7, 747-56 (Jul 1972).

The operational status of an auroral-zone incoherent-scatter radar is discussed. The Stanford Research Institute 1290-MHz incoherent-scatter radar, which was moved to Chatanika, Alaska ($L = 5.7$), an auroral-zone location near Fairbanks, is now producing interesting new results. The siting of the radar is such that D-, E-, and F-region incoherent-scatter measurements can be made without ground-clutter interference. Although coherent auroral-clutter echoes can at times be seen at relatively low elevation angles in the north, auroral echoes are sufficiently weak at 1290 MHz in the sidelobes of the antenna that they are not detectable during normal incoherent-scatter measurements. The capabilities of the radar system are outlined, and samples are presented of the type of incoherent-scatter results being obtained. (auth)

46284 RING CURRENT EFFECT ON MAGNETOSPHERIC ELECTRON DENSITY PROFILES DERIVED FROM PLASMAPAUSE WHISTLERS. Mathur, A. (Univ. of Southampton, Eng.); Rycroft, M. J.; Sagredo, J. L. *Nature (London)*; 237: No. 5357, 508-10 (30 Jun 1972).

It is stated that the longitudinal drift of Van Allen radiation belt charged particles, with opposite charges drifting in opposite directions in the geomagnetic field, constitutes a westward directed ring current at a distance of several Earth radii, and although at magnetically quiet times this current neither reduces to the geomagnetic field nor affects the parameters that determine the group velocity of a whistler mode-signal propagating along a particular field line, these can be changed significantly at magnetically active times, such as during the main and recovery phases of a magnetic storm, where the ring current is stronger. The assumption that is conventionally made when deriving magnetospheric electron density profiles from whistlers, that the Earth's magnetic field is a perfect dipole, is valid during magnetically quiet periods, and electron density profiles exhibiting the plasmapause have thus been deduced from whistlers at times of both very high and low planetary magnetic index. During geomagnetic storms, however, the enhanced ring current reduces the equatorial magnetic field strength and so reduces the electron gyrofrequency and the nose frequency of a whistler propagating along a field line that passes through a particular point on the Earth's surface; it is necessary, therefore, to include both the distortion of the field line due to the ring current and the change of the whistler mode refractive index when determining the equatorial density along the

actual path of propagation. Results are given that show quantitatively the effect of the ring current perturbation to the geomagnetic field on whistler propagation near the plasmapause. The ring current effect was pronounced outside the plasmapause, noticeable on the plasmapause, and insignificant within the plasmasphere. Also, when the effect of a ring current is allowed for, the electron density change at the plasmapause is increased, making the plasmapause an even more noticeable feature of the distribution of thermal plasma in the magnetosphere. (UK)

46285 VLF EMISSIONS DURING POST BREAKUP PHASE OF POLAR SUBSTORM. Hayashi, Kanji; Kokubun, Susumu (Tokyo Univ.). *Rep. Ionosphere Space Res. Jap.*; 25: No. 4, 369-82 (Dec 1971).

Fine structures of vlf emissions during substorms are examined, using data obtained at Syowa Station in Antarctica. It is found that a special type of chorus emission, denoted here by auroral chorus, is observed during the maximum or recovery phase of substorm. The spectral structure of auroral chorus is quite different from that of daytime polar chorus. Rising elements with duration of about 0.2 second intermittently appear in a group of several seconds' duration and the interval between these groups is usually of the order of ten seconds. The whole event lasting about 40 minutes consists of short sequences of risers' groups overlapping in time in a frequency of 0.5-2 kHz. These characteristic time scales of chorus emissions during substorms are almost in agreement with those of pulsating auroras, suggesting a common origin for the energetic electron precipitation that causes pulsating aurora. (auth)

46286 OBSERVATION OF THE INTENSITY RATIO BETWEEN (5,1) AND (9,4) BANDS OF OH EMISSION IN THE NIGHT AIR-GLOW. Yano, Kyo; Takahashi, Hisao (Niigata Univ., Japan). *Rep. Ionosphere Space Res. Jap.*; 25: No. 4, 301-16 (Dec 1971).

Observations were made of the intensity ratio of (5, 1) to (9, 4) bands of OH emission in the nightglow from November 1968 to October 1969 with an interference filter tilting photometer. The methods of evaluating the ratio from photometer readings are described. About 245 values of the ratio were obtained on 25 clear nights in the twelve months, scattering around 1.7 with the maximum of 2.1 and the minimum of 1.4. Although some consistency of its monthly variation was found, no regularity such as seasonal variation could be ascertained. It was revealed that on more than half of the observed nights the ratio was roughly constant in the course of night except in twilight conditions, while a nocturnal change from 1.4 up to 2.1 was observed on several occasions. These results are interpreted in terms of the vibrational population of OH molecules in the ground state. It is suggested that the OH molecules in the upper atmosphere could be produced through various kinds of chemical reactions with different amounts of energy being released, and that there might be a change in the rate of each reaction contributing to the OH formation with a characteristic time of only a few hours duration. (auth)

46287 ELECTRON CONTENT OF THE IONOSPHERE, SOLAR, AND GEOMAGNETIC ACTIVITY. Ebel, A. (Univ., Cologne). *Gerlands Beitr. Geophys.*; 81: No. 1-2, 11-20 (1972). (In German).

Observations of the ionospheric electron content N_T made at several stations of low and middle latitudes were analyzed with respect to solar and geomagnetic activity. The data cover the period 1965-1968. The coefficients of the regression equation $N_T = c_0 + c_S \cdot S + c_{AP} \cdot A_p$, S (flux of the solar 10.7-cm radiation) and A_p (geomagnetic-activity index) were computed for each month of the year and several hours of the day as far as data were available. The results show a clear semiannual variation of the coefficient c_S at all stations at noon and a systematic seasonal variation of c_{AP} with a minimum in summer and a maximum in winter at higher latitudes, whereas c_{AP} varies irregularly at lower latitudes. The night-time value of c_S reaches a maximum in summer or late spring at all stations with the exception of the equatorial one. The daily amplitude of c_S is smaller during summer than during winter. A comparison with a similar analysis of $(f_oF_2)^2$ shows that the electron content depends on the geomagnetic activity in a different way than does the maximum electron density. It should be noted that the coefficient c_0 of the regression equation nearly vanishes for all months and stations. The different dependence of the electron content on solar and geomagnetic activity, respectively, explains several observed peculiarities of the variation of N_T during one solar cycle, for instance the hysteresis effect, i.e. higher values during the descending than the ascending part of the solar cycle. The seasonal anomaly can be described by the coefficient c_{AP} for the higher latitudes. The semiannual and seasonal variations of c_S and c_{AP} are supposed to reflect various transport properties of the thermosphere with respect to the ion-

ized component. This may be a meridional transport shifting of the plasma from low to high latitudes during the equinoxes, thus producing a semiannual variation. On the other hand, the possible influence of a prevalent zonal wind system—superimposed on the daily wind variations in the thermosphere at middle and high latitudes—allows explanation of the changes of c_{AP} observed at the northern stations. (tr-auth)

46288 MODEL STUDIES OF DENSITY FLUCTUATIONS OF SMALL AND LARGE IONS IN THE ATMOSPHERE. Lundquist, S.; Murty, R.; Rosen, V.; Scuka, V. (Institutet foer Hoegspaeningsforskning, Uppsala). *Phys. Scr.*; 4: No. 6, 293-6(1971).

A simplified theoretical model for atmospheric ions has been used to study the variation of small-ion production rate. Linear equations have been derived to study the fluctuations of small and large ion concentrations with time. It is found that no oscillating solutions are possible. Using Bendixon's theorem it is shown that the same conclusion holds for the nonlinear equations. The conclusions have been compared with the results of other workers. (auth) (Sweden)

46289 SMALL-SCALE FLUCTUATIONS IN THE F-REGION CRITICAL FREQUENCY OBSERVED AT DELHI. Setty, C. S. G. K.; Gupta, Anurag B. (Univ. of Delhi). *Indian J. Pure Appl. Phys.*; 10: No. 2, 161-3(Feb 1972).

Accurate measurements of small fluctuations of F-region critical frequency have been made over a period of one year. Spectral analysis of these records shows periodic fluctuations in F-region critical frequency of periods between 10 and 100 min falling in the range of acoustic-gravity waves in the ionosphere with a sharp lower cutoff period in the neighborhood of 10 min. The mean amplitude in this range is of the order of 50 ± 10 kHz. As the height of reflection at the peak of the F-layer changes with season, the lower cutoff period of the spectral components is found to increase with the increase of height. It is also noted that the optimum period is in the range 24 to 30 min depending on height and that the total rms deviation of electron density is about 0.5%. Also, a preliminary study has been made to yield apparent horizontal and vertical velocity components in the plane of the magnetic meridian using the radio-ray tracing and the cross-correlation techniques. (auth)

46290 LARGE SCALE PERIODIC DISTURBANCES IN THE IONOSPHERE. Basu, S.; Guhathakurta, B. K. (University Coll. of Tech., Calcutta). *Indian J. Pure Appl. Phys.*; 10: No. 2, 164-6(Feb 1972).

Beacon satellite observations, carried out at Haringhata Field Station (geographic lat: $22^{\circ}58'N$, long: $88^{\circ}30'E$, Magnetic dip: $32^{\circ}N$) of the University of Calcutta, indicate the presence of large-scale irregularities of electron density, periodic on a spatial scale. These ionospheric disturbances occurring during summer months over the post-midnight period are often associated with anomalous nighttime gradients of electron density. Under such circumstances the spatial wavelengths are found to change suddenly from low to high wave numbers. (auth)

46291 COUPLING BETWEEN THE IONOSPHERE AND THE PROTONOSPHERE AND ITS IMPLICATIONS ON THE LONG-TERM VARIATIONS OF IONOSPHERIC ELECTRON CONTENT. Waldman, Helio. Stanford, Calif.; Stanford Univ. (1972). 185p. University Microfilms Order No. 72-16,812.

Thesis.

Recent experimental determinations of the flux of ionization between the ionosphere and the protonosphere have indicated that, during the day, the plasma will often flow upwards at a rate that by far exceeds the previously calculated theoretical limit of that flux. The theory of the coupling between the two layers is re-examined in the light of the new evidence, and the discrepancy is shown to be probably due to a previous underestimation of the neutral hydrogen concentrations in the thermosphere. In the upper F region, where diffusion dominates the behavior of the ionospheric oxygen ions, there is a maximum velocity with which such ions can flow upwards. This fact, combined with the condition of photochemical equilibrium prevailing in the lower ionosphere, gives rise to the existence of a limiting upward flux of oxygen ions from the ionosphere. In the topside ionosphere, a similar situation occurs with the protons: up to a certain height they are under chemical equilibrium, and above, they can escape upwards at a rate that is limited by collisions with the oxygen ions, the dominant charged species. Consequently, a maximum proton flux also exists, which limits the rate at which ionization can be transferred from the ionosphere to the protonosphere. Such flux is controlled by the abundance of oxygen ions in the topside ionosphere, which in turn is strongly dependent on the upgoing flux of this species. For continuity reasons, the flux of oxygen ions coming from the bulk of the F region must match the flux of protons escaping the ionosphere. It is shown that, on account of these simultaneous relationships,

the limiting exchange flux cannot be properly determined from a fixed model of the F region (as has been attempted in the past) particularly when the resulting flux value is comparable with the limiting upward flux of oxygen ions in the ionosphere. In order to circumvent this difficulty, a self-consistent limiting proton flux is derived. The effect of the exchange flux on the daytime ionosphere is studied. The long-term variations in the daytime exchange flux are estimated with the use of model hydrogen concentrations based on the well-known inverse relationship between the abundance of neutral hydrogen and the neutral temperature in the thermosphere. The present work also includes a review of the basic processes occurring in the ionosphere, with emphasis on the concepts of limiting velocity and limiting flux. In addition, an approach to the problem of numerical simulation of the ionosphere is presented and discussed. (TSS) (Diss. Abstr. Int., B)

46292 OPTICAL DETECTION OF RADIO-FREQUENCY HEATING OF ELECTRONS IN THE F REGION. Sipler, Dwight Passmore. Pittsburgh; Univ. of Pittsburgh (1971). 94p. University Microfilms Order No. 72-17,098.

Thesis.

Heating of the electrons in the F region of the ionosphere by a high-power beam of radio-frequency energy was detected by observation of attendant effects on the intensity of the 6300-Å line of the atomic-oxygen nightglow. Collisional absorption of rf energy propagated in the extraordinary mode produces a small (few percent) transient suppression of the intensity of the 6300-Å line as a result of the decrease in the dissociative recombination production of $O(^1D)$ atoms with increasing electron temperature. The magnitudes of the observed intensity suppression signals are consistent with electron temperature increases in the range 8 to 35%. A plasma instability triggered by the high-intensity rf fields propagated in the ordinary mode was observed to cause large increases in the 6300-Å line intensity by direct electron impact excitation of the ^1D state of atomic oxygen. Measurements of the time constants of these intensity changes have enabled an in-situ determination of the quenching coefficient for the $O(^1D)$ state assuming the quenching is due to N_2 . A value $k_q = (5 \pm 2) \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ has been obtained by this method. (Diss. Abstr. Int., B)

46293 DAILY VARIATION OF THE IONOSPHERIC F2 EQUATORIAL ANOMALY IN THE AMERICAN AND ASIAN SECTORS. Anderson, David Nels. Boulder, Colo.; Univ. of Colorado (1971). 157p. University Microfilms Order No. 72-17,231.

Thesis.

When observed noontime values of the maximum electron density in the ionospheric F2 region are plotted as a function of magnetic latitude, a curve is produced that has two peaks, one on either side of the dip equator at $\pm 16^{\circ}$ dip latitude. A theoretical investigation is made of the daily variation of the latitudinal distribution in NMAX(F2) (the so-called Appleton or equatorial anomaly) and specifically attempts to account for the longitudinal differences observed between the American and Asian sectors. Solution of the time-dependent electron continuity equation yields the electron density. Since the coefficients are complicated functions of space and time, the equation is solved numerically. Production, loss, and transport of ionization are taken into account, where the effects of neutral wind, ambipolar diffusion, and $\vec{E} \times \vec{B}$ drift are included in the transport term. By describing the magnetic field in two equivalent ways $\vec{B} = -\nabla\gamma$ and $\vec{B} = \nabla\alpha \times \nabla\beta$, where α , β , and γ are known magnetic scalar potentials, it is possible to transform the spherical r , θ , and ϕ space coordinates of the continuity equation to coordinates that define directions parallel and perpendicular to the magnetic field thus putting the equation in a form suitable for numerical integration. For each of the two sectors, results are presented indicating the extent to which the equatorial anomaly's daily variation is affected by changes in the geomagnetic field configuration, neutral wind, and $\vec{E} \times \vec{B}$ drift. It is found that development of the anomaly is most sensitive to the electric-field model assumed. When NMAX(F2) is plotted as a function of local time at the magnetic equator, the observed differences between the American and Asian sectors can be accounted for by an upward $\vec{E} \times \vec{B}$ drift that commences two hours earlier in the Asian. Other investigations includes the effects of the neutral wind in producing north-south asymmetry in the anomaly; the solar-cycle effect on the anomaly's daily variation; and the comparison of results between the centered tilted dipole approximation and the real magnetic field to estimate the differences introduced by the approximation. (Diss. Abstr. Int., B)

46294 CHARGED-PARTICLE TRANSPORT IN TURBULENT ASTROPHYSICAL PLASMAS. Newman, Charles Edward Jr. Stanford, Calif.; Stanford Univ. (1972). 79p. University Microfilms Order No. 72-20,719.

Thesis.

The effect of electromagnetic fluctuations, or plasma turbulence,

on the motion of the individual particles in a plasma is investigated; two alternative methods are used to find a general equation governing the time-evolution of a distribution of charged particles subject to both an external force field and the random fields of the fluctuations. It is found that, for the high-temperature, low-density plasmas frequently encountered in the study of astrophysics, the presence of even a small amount of turbulence can have a very important effect on the behavior of the plasma. Two problems in which turbulence plays an important role are treated in detail. First, the contribution to electrical and thermal conductivities by the turbulence is calculated; it is shown that for the solar wind at the orbit of Earth the turbulent effects dominate the effects of interparticle collisions in the determination of these transport coefficients. The implications of the transport coefficient calculation on models of the solar wind are discussed. Second, the problem of a distribution of relativistic electrons being stochastically accelerated by plasma turbulence in addition to emitting synchrotron radiation is discussed; this process has been proposed by other authors to be responsible for the observed electron energy spectra of synchrotron radio sources, but, based on arguments concerning the particle flux in phase space, this proposal is found to be very improbable. (Diss. Abstr. Int., B)

46295 DESIGN CONSIDERATIONS IN THE MEASUREMENT OF ELECTRON TEMPERATURE IN THE IONOSPHERE. Tyler, A. F. (Univ. of Birmingham, Eng.). *Radio Electron. Eng.*; 42: No. 7, 309-13 (Jul 1972).

Measurements of electron temperature in the ionosphere using rocket and satellite-borne probes were made, using an experimental technique involving an extension of Langmuir-probe principles. The apparatus, in its latest form, was developed to provide a more compact and reliable system by exploiting the advantages of linear integrated circuits over the discrete component circuitry used previously. In considering the performance of the experiment, particular attention was paid to the stability problems associated with the closed-loop system employed. (auth)

46296 NATURE OF A SEMI-SECULAR CYCLE IN THE TROPOSPHERE. Loginov, V. F. *Tr. Gl. Geofiz. Observ.*; No. 245, 25-30 (1969). (In Russian).

Data supporting the cosmic nature of a semi-secular cycle in certain characteristics of the troposphere are advanced (the circulation of the atmosphere in the region of the North Sea, a thermal cycle at high latitudes). Taking into account contemporary knowledge of the intensity and the structure of the magnetic field in the Universe, and the assumption of high-energy particles as a disturbing factor in tropospheric circulation, it is possible to comprehend many features of the Sun-troposphere relation in time, and in particular their instability. (tr-auth)

46297 CIRCULATION IN THE TROPOSPHERE AND THE STRATOSPHERE AND HIGH-ENERGY PARTICLES. Sazonov, B. I. *Tr. Gl. Geofiz. Observ.*; No. 245, 56-63 (1969). (In Russian).

Atmospheric circulation on the Earth under various conditions is considered, particularly the cyclones and anticyclones in the stratosphere. At the same time, circulation peculiarities in the atmosphere were different in these layers, and were traced not only in latitudinal, but also in longitudinal effects at levels H_{50} and H_{500} . Reasons are advanced supporting the hypothesis of high-energy protons absorbed in the troposphere and stratosphere for being different at different levels, and accordingly stimulate the cyclone-anticyclone pairs at these levels. Troughs and crests are generated in tropospheric and in the stratosphere regions of attenuation of the intensity of the Earth's magnetic field, which are favorable to the intrusion of protons. These are related to the transfer of air masses from west to east in the region of a magnetic anomaly under the influence of cosmic radiation. (tr-auth)

46298 NEW METHOD FOR THE DETERMINATION OF $N(h)$ IN THE D-REGION. Rumi, G.-C. (Istituto Elettrotecnico Nazionale Galileo Ferraris, Turin). *Nuovo Cim.*; 10B: No. 1, 240-54 (11 Jul 1972).

It is shown that an lf ((30 to 300) kHz) link operating essentially on a one-hop ionospheric propagation of an ordinary and quasi-transverse wave can be used for the determination of the electron density at some heights in the D region; it is just enough to detect certain peculiar doublets of single-strength breaks at the receiving site. Such breaks—fifteen of them were observed during February and March 1970 at the IENGf in Turin—are interpreted as the result of a swift increase of the electron density in the lower ionosphere and are shown theoretically to be characterized by the following values of the electron density N , and of the electron collision frequency ν : a) $N = 0.107 f^2$ and b) $\nu = 13.60 f$, where f is the operating frequency. Since $\nu = \nu(h)$ depends only on the height h , and is an available datum, a point of the electron-density pro-

file, namely $N = 0.107 f^2$ and $h = h(13.60 f)$, can be associated with each break. The breaks yield information concerning a disturbed electron-density profile since they relate to disturbed conditions in the ionosphere. A method whereby the electron-density profile for quiet conditions can be derived from the same for disturbed conditions is outlined. (auth)

46299 PHOTODETACHMENT OF ELECTRONS FROM MAJOR NEGATIVE IONS IN THE LOWER D REGION. Thomas, L.; Gondhalekar, P. M.; Bowman, M. R. (*Radio and Space Research Station, Slough, Eng.*). *Nature* (London); 238: No. 5359, 89-90 (14 Jul 1972).

It is stated that the relatively high atmospheric pressures in the lower D region of the ionosphere, at heights below about 80 km, permit the formation of negative ions by the three-body attachment of electrons to molecules or atoms, and the photodetachment of electrons from negative ions by solar radiations has been used to explain increases in electron concentrations occurring prior to ground sunrise. A theoretical study of the height distributions of electrons and negative ions shows the need, however, to consider the detachment of electrons from the major negative ions throughout the day in order to account for the daytime electron concentrations observed below 70 km. Reliable laboratory measurements of photodetachment efficiencies are available only for O^- , O_2^- and O_3^- , but the likely effects of including the photodetachment of electrons from other ions, such as CO_3^- , NO_2^- and NO_3^- are here considered. (UK)

46300 THEORETICAL MODELS OF IONOSPHERIC STORMS. Matuura, Nobuo (*Radio Research Labs., Tokyo*). *Space Sci. Rev.*; 13: No. 1, 124-89 (May 1972).

Ionospheric F-region disturbances, commonly called ionospheric storms, are characterized by changes in global electron-density distribution over a period of a few days accompanied by geomagnetic storms. The global morphology and disturbances in polar and high-latitude regions and equatorial regions are described. Processes affecting the ionosphere are discussed, including: corpuscular effects on the ionosphere, chemical reactions in the ionosphere, and effects of plasma transport. The theoretical models of ionospheric storms presented are for global disturbances in the upper atmosphere, polar and high-latitude ionosphere, middle- and lower-latitude ionosphere, and equatorial ionosphere. 350 references. (J.C.W.)

Particle Kinetics and Trapping

46301 (N-72-18362) LIGHT-ION TROUGH. Taylor, Harry A., Jr. (National Aeronautics and Space Administration, Greenbelt, Md. Goddard Space Flight Center). Feb 1972. 35p. (NASA-TM-X-65832; X-621-72-44). NTIS.

Several fundamental characteristics of LIT (light ion trough), including evidence of diurnal variation as well as response to magnetic disturbances, are identified. These identifications were made using the Bennett radio-frequency ion mass spectrometer onboard the OGO 6 from 1969 to 1970. It was shown that the LIT, rather than the total ionization trough, appears as a more meaningful indicator of the plasmopause. This makes the LIT a fundamental parameter for studies of the formation and maintenance of the plasmasphere. (STAR)

46302 ANALYSIS OF PIONEER 9 LOW-FREQUENCY WAVE OBSERVATIONS NEAR INTERPLANETARY DISCONTINUITIES. Scarf, F. L.; Greenstadt, E. W. (TRW Systems Group, Redondo Beach, Calif.). *J. Geophys. Res.*; 77: No. 19, 3317-27 (1 Jul 1972).

The sector boundary of February 2, 1969 was unusually complex, and within a single 7-hour period the Pioneer 9 magnetometer and plasma probe detected a reverse shock, a slow shock, and a conventional fast shock. During this interval the electron gyrofrequency and the ion plasma frequency each varied by more than a factor of 10, and the low-frequency (broad-band and 400-Hz) electric-field wave levels were high and variable. It is shown that, near the discontinuities, enhanced 400-Hz wave levels were not generally associated with field gradients or current layers, but elevated 400-Hz wave levels were detected when the ion density fell within a fairly well-defined window. This suggests that modes related to ion sound waves were detected, and the observations are assessed in terms of various theories for interplanetary-wave generation. (auth)

46303 EXPLORER 33 AND 35 PLASMA OBSERVATIONS OF MAGNETOSHEATH FLOW. Howe, Herbert C. Jr.; Binsack, Joseph H. (Massachusetts Inst. of Tech., Cambridge). *J. Geophys. Res.*; 77: No. 19, 3334-44 (1 Jul 1972).

The magnetosheath flow in the region $-60 R_E \leq X_{SE} \leq -20 R_E$ is mapped by using plasma data from the Massachusetts Institute of

Technology plasma experiments on the Earth-orbiting Explorer 33 and the lunar-orbiting Explorer 35. Temporal and spatial variations in the flow are separated by using 309 hours of simultaneous data from the two spacecraft, one spacecraft being in the solar wind and the other spacecraft in the magnetosheath. The overall flow pattern agrees well with the predictions of a simplified hydrodynamic theory. The measured ratio of magnetosheath to solar-wind densities is shown to have some dependence on the observed solar-wind sonic Mach number. This density ratio is consistently <1 in the region adjacent to the magnetotail boundary and thus reflects the irreversible nature of the flow across the bow shock. (auth)

46304 ENTRY OF HIGH-ENERGY SOLAR PROTONS INTO THE DISTANT GEOMAGNETIC TAIL. Durney, A. C. (European Space Tech. Centre, Noordwijk, Netherlands); Morfill, G. E.; Quenby, J. J. *J. Geophys. Res.*; 77: No. 19, 3345-60(1 Jul 1972).

During the solar proton events of November 18, 1968, and February 25, 1969, a detector on board the low-altitude polar-orbiting Esro-2 spacecraft observed marked structure in the enhanced intensities of 100- to 300-MeV protons at high latitudes. While concentrating mainly on the results of the northern hemisphere for November 18, this structure has been investigated in an attempt to discover the access mechanisms for these protons. A trajectory program was developed to discover the access positions at the magnetopause and the asymptotic directions: it was found that the main features of the structure could be accounted for by direct access of the anisotropic interplanetary flux and by a high-intensity flux of protons approaching the Earth from the tail. Secondary features of the structure need an improved trajectory program for interpretation. After a brief discussion of some tail models the conclusion was reached that the protons entering the tail were not consistent with either a diffusive or an adiabatic reconnection model. A tail access process that appears to satisfy the results and whose main requirements are a sharp magnetic-field boundary at the magnetopause and a slightly divergent tail field is discussed. (38 references) (auth)

46305 CHANGES IN MAGNETOSPHERIC CONFIGURATION DURING THE SUBSTORM GROWTH PHASE. Coroniti, F. V.; Kennel, C. F. (Univ. of California, Los Angeles). *J. Geophys. Res.*; 77: No. 19, 3361-70(1 Jul 1972).

Three calculations suggest that the magnetospheric configuration changes during the substorm growth phase. A flaring-tail model indicates that the observed increases in geomagnetic-tail field are explained if the dayside magnetopause shrinks by 1 to 2 R_E . Increased tail flaring also requires that the tail current sheet approach near the Earth during the growth phase. The motion of the inner edge of the plasma sheet, and consequently an equatorward shift of the nightside auroral oval, is consistent with the structural modifications mentioned above. (auth)

46306 ANISOTROPY OF HIGH-LATITUDE ELECTRON FLUXES DURING SUBSTORMS AND STRUCTURE OF THE MAGNETOTAIL. McDiarmid, I. B.; Hruska, A. (National Research Council of Canada, Ottawa). *J. Geophys. Res.*; 77: No. 19, 3377-83(1 Jul 1972).

High-latitude electron profiles ($E > 20$ keV) observed near midnight during the expansive phase of substorms frequently show pitch-angle distributions strongly peaked at 90° to the local magnetic field. Such distributions, which are characteristic of bouncing particles, are observed up to invariant latitudes of 80° , which is roughly 10° higher than normal. This phenomenon implies that the tail magnetic field up to distances of the order of $100 R_E$ is closed and sufficiently regular to allow electrons of these energies to bounce between mirror points. The results suggest that during the expansive phase of substorms the neutral line lies many tens of Earth radii beyond the Earth and that the thickness of the neutral sheet may be of the order of magnitude of the thickness of the plasma sheet. The high-latitude electron profiles also suggest the existence of large but continuous variations of density in the tail plasma sheet in the direction perpendicular to magnetic lines of force. (auth)

46307 HIGH-ENERGY ELECTRON SPIKES AT HIGH LATITUDES. Brown, J. W.; Stone, E. C. (California Inst. of Tech., Pasadena). *J. Geophys. Res.*; 77: No. 19, 3384-96(1 Jul 1972).

Over 750 spikes of precipitating electrons with $E \geq 425$ keV were observed aboard the low-altitude polar orbiter Ogo 4 between July 30, 1967, and December 31, 1967. The spikes may be divided into three distinct populations depending on whether they occur at latitudes below, at, or above the local limit of trapping. These spikes are designated type 1, 2, and 3, respectively. Type-3 spikes occur in a narrow latitude band about 3° wide, centered at invariant latitude $A \approx 78^\circ$ at 1000 MLT (magnetic local time) and 68° at 2000 MLT. Their relative frequency of occurrence, intensity, and

hardness do not depend significantly on magnetic local time. Type-3 spikes appear to be associated with spikes observed near the magnetopause and the neutral sheet. Type-2 spikes also occur in a latitude band about 3° wide, centered at about 71° at 1000 MLT and 67° at 2200 MLT. Their frequency of occurrence is highly dependent on magnetic local time, a large maximum occurring near 2300 and very few events occurring between 0600 and 1200 MLT. Type-2 spikes appear to be related to island fluxes in the neutral sheet, although they occur on closed field lines and may persist for many hours. Type-1 spikes occur in a wider band of latitudes, from about 62° to 68° near midnight and 66° to 68° near noon. The local-time dependence of their frequency of occurrence is similar to that of Type-2 spikes but less pronounced. Although they are observed on closed field lines, Type-1 spikes do not persist for periods longer than about 1 hour, and it was concluded that they are produced by strong pitch-angle scattering from the stably trapped population. The average spectral index (assuming a power-law spectrum) is 3 to 5, and the median flux (>425 keV) is about $150 \text{ el cm}^{-2} \text{ sec}^{-1} \text{ ster}^{-1}$, although Type-2 events near midnight tend to be larger and harder. All types tend to be more intense and to occur at lower latitudes when K_p is large. (auth)

46308 ELF NOISE BANDS ASSOCIATED WITH AURORAL-ELECTRON PRECIPITATION. Gurnett, D. A.; Frank, L. A. (Univ. of Iowa, Iowa City). *J. Geophys. Res.*; 77: No. 19, 3411-17(1 Jul 1972).

Observations of a new type of elf noise band that is closely associated with low-energy auroral electron precipitation are reported. These observations have been made at relatively low altitudes (<3000 km) with the polar-orbiting satellite Injun 5. The noise bands typically have a center frequency of 100 to 300 Hz and often appear to consist of many nearly monochromatic bursts, typically of a few seconds' duration, superimposed to produce the observed noise band. These elf noise bands are observed only in a relatively narrow range of latitudes (a few degrees) in the auroral zone and are almost always associated with intense fluxes, $10^8 \text{ el cm}^{-2} \text{ sec}^{-1} \text{ ster}^{-1}$, of precipitating electrons with energies from a few hundred electron volts to several kiloelectron volts. On the dayside of the magnetosphere the region where the elf noise bands and the associated low-energy electron precipitation are observed has been identified as the polar cusp. In considering the possible explanations of these elf noise bands it is noted that the spectral characteristics of this noise are very similar to a type of narrow-band electromagnetic noise called "lion's roar," which has been observed at much higher altitudes in the magnetosheath with the satellite Ogo 5. It is suggested that the elf noise bands observed at low altitudes with Injun 5 are caused by lion's roar emissions that have propagated down "open" magnetic-field lines to low altitudes from the magnetosheath region. (41 references) (auth)

46309 GEOMAGNETICALLY TRAPPED ALPHA PARTICLES. I. OFF-EQUATOR PARTICLES IN THE OUTER ZONE. Blake, J. B.; Paulikas, G. A. (Aerospace Corp., Los Angeles). *J. Geophys. Res.*; 77: No. 19, 3431-40(1 Jul 1972).

The population of trapped alpha particles in five contiguous energy channels between 0.85 and 9.0 MeV was measured in the B and L intervals $2 \leq L \leq 4$ and $0.06 \leq B \leq 0.35$ gauss during June and July 1969. The particle distribution functions of these particles suggest that the location of the source of these particles is at $L > 4$ and that inward radial diffusion determines the spatial and energy distributions. The particle distribution functions show that these alpha particles do not diffuse into the inner zone. These results provide no support for the concepts of in-situ acceleration or direct capture of energetic solar alpha particles. (auth)

46310 INNER-ZONE ENERGETIC-ELECTRON REPOPULATION BY RADIAL DIFFUSION. Tomassian, Albert D. (Univ. of California, Los Angeles); Farley, Thomas A.; Vampola, Alfred L. *J. Geophys. Res.*; 77: No. 19, 3441-54(1 Jul 1972).

A quantitative study of the intrusion of natural electrons into the inner radiation zone during and after the geomagnetic storm of September 2, 1966, shows that the transport is consistent with a radial-diffusion mechanism in which the first two invariants are conserved. Except for the 3-day period of the storm main phase when data were missing, the radial-diffusion coefficient is $D = 2.7 \times 10^{-6} \text{ L}^{7.9} \mu^{-0.5} \text{ day}^{-1}$ in the range $1.7 \leq L \leq 2.6$ and $13.3 \leq \mu \leq 27.4 \text{ MeV gauss}^{-1}$. This value could be produced by variation of a large-scale electric field across the magnetosphere having an amplitude of 0.28 mV/m and a period of 1600 sec. Electric fields having approximately these characteristics have been inferred from previous observations of the motion of whistler ducts within the plasmapause. If fields of this amplitude and period exist throughout the magnetosphere, the radial diffusion of all geomagnetically trapped particles except the high-energy inner-zone protons is strongly influenced by electric-field variations. A comprehensive review of previously reported radial-diffusion coef-

ficients shows reasonable agreement for L less than about 3.0, but serious discrepancies among reported values exist for determinations made in the outer zone. These discrepancies cannot be explained by the simple theory of radial diffusion due to variation of large-scale electric or magnetic fields. (auth)

46311 PITCH-ANGLE DIFFUSION OF RADIATION-BELT ELECTRONS WITHIN THE PLASMASPHERE. Lyons, Lawrence R.; Thorne, Richard Mansergh; Kennel, Charles F. (Univ. of California, Los Angeles). *J. Geophys. Res.*; 77: No. 19, 3455-74(1 Jul 1972).

An attempt is made to explain the formation of the quite-time electron slot, which divides the radiation-belt electrons into an inner and an outer zone. The pitch-angle diffusion of radiation-belt electrons resulting from resonant interactions with the observed plasmaspheric whistler-mode wave band is investigated. The effects of wave propagation obliquely to the geomagnetic-field direction with the resulting diffusion at all cyclotron-harmonic resonances and the Landau resonance are evaluated, along with the effects of interactions occurring at all geomagnetic latitudes. The results account for the long-term stability of the inner radiation zone, the location of its outer edge as a function of electron energy, and the removal of electrons to levels near zero throughout the slot. Computed pitch-angle distributions and precipitation decay rates are in good agreement with slot-region observations. (38 references) (auth)

46312 OBSERVATION OF VERY-LOW-FREQUENCY WHISTLER-MODE WAVES IN THE REGION OF THE RADIATION-BELT SLOT. Koons, H. C.; McPherson, D. A. (Aerospace Corp., El Segundo, Calif.). *J. Geophys. Res.*; 77: No. 19, 3475-82(1 Jul 1972).

Following a sudden-commencement magnetic storm, band-limited vlf emissions between 3.9 and 10.4 kHz were observed at high altitudes within the plasmasphere ($L \sim 2.5$ to 3.6) by a receiver aboard spacecraft OV1 14 (1968-26B). The maximum spectral intensities of 1.1 mV/Hz^2 and $2.8 \text{ } \mu\text{V/Hz}^2$ on the magnetic and electric antennas, respectively, occur in the 5.6-kHz channel. The waves can resonantly interact with energetic electrons above $\sim 50 \text{ keV}$ at $L = 2.5$, and the measured wave intensities are sufficiently high to account for the short lifetimes of medium-energy electrons in the slot region between the inner and outer Van Allen radiation zones. A comparison with ground-based vlf data suggests that the waves are generated by band-limited amplification of incoherent noise. (auth)

46313 TURBULENCE OF ELECTROSTATIC ELECTRON CYCLOTRON HARMONIC WAVES OBSERVED BY OGO 5. Oya, Hiroshi (Goddard Space Flight Center, Greenbelt, Md.). *J. Geophys. Res.*; 77: No. 19, 3483-94(1 Jul 1972).

Vlf emissions that have been observed near $3/2 f_H$, $5/2 f_H$, and $7/2 f_H$ by Ogo 5 in the magnetosphere (f_H is the electron cyclotron frequency) are analyzed in the light of the mechanism used for the diffuse plasma resonance f_{Dn} observed by Alouette 2 and Isis 1. The vlf emission is considered to be generated by nonlinear coupling mechanisms in certain portions of the observation as the f_{Dn} is enhanced by its association with nonlinear wave-particle interaction of the electrostatic electron cyclotron harmonic wave, including the instability due to the nonlinear inverse Landau damping mechanism in the turbulence. The difference between the two observations is in the excitation mechanism of the turbulence; the turbulence in the plasma trough detected by Ogo 5 is due to natural origins, whereas the ionospheric topside sounder makes the plasma-wave turbulence artificially by submitting strong stimulation pulses. Electron density values in the plasma trough are deduced by applying the $f_{Dn} - f_H/f_H$ relationship obtained from the Alouette-2 experiment as well as by applying the condition for the wave-particle nonlinear interactions. The electron density values reveal good agreement with the ion-density values observed simultaneously by the highly sensitive ion mass spectrometer. (auth)

46314 STABILITY OF OBLIQUELY PROPAGATING WHISTLERS. Brinca, A. L. (Stanford Univ., Calif.). *J. Geophys. Res.*; 77: No. 19, 3495-3507(1 Jul 1972).

By means of an energy conservation approach, an analysis was made of the growth rates of whistlers with arbitrary frequency and direction of propagation in a cold plasma permeated by a dilute energetic electron population. Numerical results are obtained for whistler frequencies and plasma characteristics prevailing at the onset of artificially stimulated emissions ($\omega/\Omega \sim 0.5$, $\omega_p/\Omega \sim 10$, the energetic electron population with varying energy and pitch angle). It is found that the growth rates do not, in general, maximize for propagation along the static magnetic field, and a criterion for the existence of maximal growth at this direction is derived. An application to artificially stimulated emissions is discussed. (auth)

46315 WHISTLER SIDE-BAND GROWTH DUE TO NON-LINEAR WAVE-PARTICLE INTERACTION. Brinca, A. L. (Stanford Univ., Calif.). *J. Geophys. Res.*; 77: No. 19, 3508-23(1 Jul 1972).

The distortion caused by a large-amplitude whistler (with wave normal parallel to the static magnetic field) on the energetic-electron velocity distribution of a predominantly cold magnetoplasma is derived analytically. Whistler test waves impressed on the perturbed magnetoplasma and within two narrow bands centered on the frequency ω_0 of the original wave may experience large consecutive growth at the early stages of the wave-particle interaction if the unperturbed energetic-electron distribution $f_0(\nu_{||}, \nu_{\perp})$ satisfies two conditions. First f_0 must have a certain degree of $\nu_{\perp}$ monochromaticity (a well-defined peak for some $\nu_{\perp} > 0$); second, f_0 either must be linearly unstable for the original whistler or, if it is stable, it must satisfy $2(\Omega/k_0) \int_0^\infty f_0 \nu_{\perp} d\nu_{\perp} \approx |\int_0^\infty \nu_{\perp}^2 (\partial f_0 / \partial \nu_{||}) d\nu_{\perp}|$ where the integrals are evaluated at $\nu_{||} = (\omega_0 - \Omega)/k_0$, Ω is the electron cyclotron frequency of the static magnetic field, and k_0 is the wave number of the original whistler. The relevance of these whistler side bands to magnetospheric phenomena is assessed; in particular, an identification of the onset of artificially stimulated emissions with the creation of whistler side bands gives good agreement with observations. (36 references) (auth)

46316 12-MONTH WAVE IN GEOMAGNETIC ACTIVITY. Meyer, Joachim (Freie Univ., Berlin). *J. Geophys. Res.*; 77: No. 19, 3566-74(1 Jul 1972).

Besides the well-known 6-month wave in geomagnetic activity and its recurrence tendency, no persistent 12-month wave has been detected by harmonic analysis of consecutive monthly values. Such a 12-month wave, however, can be shown to exist if, instead, single activity sequences are analyzed, i.e., daily values at intervals of one solar rotation. A harmonic sequence analysis of this kind has been carried out for the daily character figures C8 from 1884 to 1968. For years before and including sunspot minimums two clearly distinguished significant 12-month waves having nearly equal amplitudes and opposite phases are obtained. The maximum occurs shortly after one of the two equinoxes: at the end of March or at the beginning of October. The amplitude is almost 3 times that of the semiannual wave in the same sequences. For years around sunspot maximums the amplitude is smaller, and in the harmonic dial the clouds of points for both 12-month waves can no longer be clearly distinguished. The dates of maximum of the revealed 12-month wave again exclude an explanation by means of the classical axial hypothesis. The statistical result instead suggests an interpretation primarily by means of a temporary N-S symmetry in the distribution of the magnetic M regions within single solar sectors and the same magnetospheric modulation mechanism responsible for the 6-month wave. (auth)

46317 DAYTIME IRREGULAR GEOMAGNETIC PULSATION, PID, AND ITS RELATION TO MAGNETOSPHERIC SUBSTORM. Morioka, Akira; Saito, Takao (Tohoku Univ., Sendai). *Rep. Ionosphere Space Res. Jap.*; 25: No. 4, 343-59(Dec 1971).

The daytime irregular pulsation in the auroral zone, Pid, is studied from data obtained with induction magnetometer at Syowa Station in 1968. Results of the analysis show that (1) the waveform of Pid is expressed by a group of many impulses having approximate pulse-height with a fraction of gammas, pulsewidth between one and five seconds, a repetition period between five and twenty seconds; (2) Pid occurs most frequently in the daytime and tends distinctly to occur on magnetically disturbed days; (3) Pid is frequently associated with the daytime cosmic-noise absorption; (4) Pid appears in the day-side auroral zone with a systematic time lag behind the breakup of a magnetic substorm on the night side; and (5) the time lag of the onset time of Pid behind that of the substorm is proportional to the local time of the onset of Pid. A model of the generation mechanism of Pid is proposed: a part of the energized electrons produced during the onset of the magnetospheric substorm drifts eastwards around the Earth, and is precipitated into the day-side auroral zone, causing the daytime cosmic-noise absorption and x-ray microbursts; the high-energy electron microbursts contribute to the ionospheric E layer, giving rise to a series of temporal and localized enhancements of the ionospheric current, whose magnetic effect is observed as a group of the impulses when the observation is carried out with induction magnetometers on the ground. (auth)

46318 CHERENKOV INSTABILITY AND VLF EMISSIONS GENERATED OUTSIDE THE PLASMAPAUSE. Kimura, Iwane (Kyoto Univ.). *Rep. Ionosphere Space Res. Jap.*; 25: No. 4, 360-8(Dec 1971).

A coupling between electromagnetic waves and space-charge waves due to beaming electrons in an ambient plasma is reconsidered. Strong couplings take place in the close vicinity of the ambient plasma frequency and electron cyclotron frequency. A

whistler-mode wave can be amplified by such a coupling for frequencies a little less than the plasma frequency if the plasma frequency is less than the cyclotron frequency. This condition will probably occur just outside the plasmopause. Vlf emissions generated by such a mechanism should be closely associated with the location of the plasmopause. (auth)

46319 ASSOCIATION BETWEEN QUASI-PERIODIC VLF EMISSION AND MICROPULSATION. Kokubun, Susumu (Tokyo Univ.). Rep. Ionosphere Space Res. Jap.; 25: No. 4, 383-8 (Dec 1971).

Vlf emissions with periods from 2 to 260 s, called periodic emission and quasi-periodic emission (QP) or long-period vlf pulsation, are often observed at high and middle latitudes. QP emissions with periods longer than 6 s are considered to be the result of hydromagnetic modulation of the basic emission source in the magnetosphere. The associations between micropulsations and vlf emissions show experimental evidence for the hydromagnetic modulations of vlf emissions based on data at Syowa Station. Four examples of intensity-modulated emissions accompanying worldwide geomagnetic changes are shown and discussed. These include a quasi-periodic modulation of vlf emission intensity associated with a world-wide geomagnetic change; periodic vlf emission associated with a regular micropulsation with a period of 10.8 seconds; concurrent occurrence of quasi-periodic emission and quasi-sinusoidal micropulsations with a period of about 40 seconds; and an association between vlf chorus emissions and irregular micropulsations. (J.C.W.)

46320 IDENTIFICATION OF SOLAR-NEUTRON EVENTS BY OBSERVATION OF RADIATION-BELT PROTON FLUXES. Clafin III, Edward Scott. Riverside, Calif.; Univ. of California (1971). 186p. University Microfilms Order No. 72-17,053. Thesis.

A region of theoretically stable trapping but experimentally low flux exists at the outer edge of the trapped proton belt. Trapping lifetimes in this region are shorter than the 10 to 200 years typical of the inner belt. Trapping for periods exceeding $\frac{1}{2}$ hour may occur. A burst of neutrons arriving at the Earth following a solar flare will cause an increase in the trapped proton flux above 20 MeV. Quantitative prediction of the increase is made possible by calculation of the injection coefficient for solar neutrons. For a sufficiently intense neutron burst, an increase will be detectable in the low flux at the outer edge of the proton belt. Proton spatial and pitch-angle distributions will help to identify the solar neutron source. Lower limits on the required neutron flux are estimated and compared with upper limits established by experiment. It is found that observation of trapped-neutron-decay protons is potentially a factor of 10 more sensitive for detecting solar neutrons than past satellite neutron experiments. A possible example of detection of trapped-neutron decay protons occurred following the August 28, 1966 solar flare. An unusual count-rate increase was observed immediately after the flare by the UCSD detector aboard Explorer 26. Comparison with Vela neutron detector data shows that the increase could not have been caused by trapped neutron decay protons. Additional events of this kind have not been found. More sensitive proton detectors may be able to observe the predicted effect. (Diss. Abstr. Int., B)

46321 MAGNETOHYDRODYNAMIC EFFECTS AT THE MAGNETOSPHERE BOUNDARY. Verzariu, Pompiliu. Washington, D. C.; American Univ. (1971). 94p. University Microfilms Order No. 72-18,100. Thesis.

Interactions between the solar wind and the magnetosphere are discussed relative to the following two aspects: (a) variations in the compression of the magnetosphere responding to solar-wind pressure changes; and (b) reflection and refraction of magnetohydrodynamic (MHD) waves at the boundary of the magnetosphere. Concerning (a), it is found that solar-wind pressure changes and magnetic-field changes at the Earth's surface are linearly correlated during magnetically quiet conditions in accordance with the theoretical expectations from the pressure balance between the solar-wind plasma and the Earth's magnetic field. A constant of proportionality relating the two quantities is statistically determined. The average value for the proportionality constant agrees favorably with the theoretical value given by the Mead-Beard magnetosphere model. Magnetohydrodynamic (MHD) theory is used to study problem (b) above, namely the transmission of waves from the magnetosheath plasma into the magnetosphere under normal conditions of solar-wind flow. Two cases taken as representative of magnetopause conditions in the vicinity of the subsolar point and near the dusk and/or dawn meridian are considered. It is found that the total transmitted energy constitutes only about 1 to 2% of the energy flux for a distribution of isotropic incident waves. The flux transmitted into the magnetosphere from waves having

an amplitude of 1 gamma (1 gamma = 10^{-5} gauss) is roughly of the order 10^{-6} to 10^{-5} erg $\text{cm}^{-2} \text{sec}^{-1}$. Assuming a steady isotropic flux of waves with amplitudes of 1 gamma incident on a hemispherical sunward magnetospheric boundary of radius $15 R_E$, the total energy input into the magnetosphere is estimated to be of the order of 10^{21} erg per day. The wavelengths and amplitudes of the ripples induced by the MHD waves on the boundary surface are also examined. For the observed power density of MHD waves in the magnetosheath, the calculated wavelengths range from a few tenths of $1 R_E$ to several hundred Earth radii. (Diss. Abstr. Int., B)

46322 DISTURBANCE OF THE EARTH'S MAGNETIC FIELD AND ITS VARIATIONS IN THE ELEVEN-YEAR CYCLE. Ol, A. I. Tr. Gl. Geofiz. Observ.; No. 245, 35-48 (1969). (In Russian).

The more important indices of magnetic disturbances and their variations in the 11-year cycle of solar activity are discussed in detail. Different indices reflect different types of corpuscular radiation from the Sun, the two basic types being flare and recurrent. Flare types arise during the maximum solar activity of the 11-year cycle, and recurrent types occur at the minimum. Annual variations in corpuscular radiation are also considered. Values of the index M of the magnetic disturbance are traced over the years from 1884 to 1967. (tr-auth)

46323 CONJUGATE POINT PHENOMENA. Rothmuller, I. J.; Beer, T. Sci. Progr. (Oxford); 60: No. 238, 205-15 (Summer 1972).

A review is given of the magnetospheric and ionospheric manifestations of conjugate point phenomena. The model generally chosen in order to calculate conjugate points is discussed. The conjugate phenomena, visual aurora, x-ray aurora, whistlers, other vlf emission, micropulsations, magnetic variations and F-region effects are discussed. (UK)

Solar Wind and Cosmic Ray Interactions

Refer also to abstract: 46322.

46324 (N-72-18796) INTERACTION OF THE SOLAR WIND WITH A PLANETARY IONOSPHERE. Blank, J. L.; Sill, W. R. (Bellcomm, Inc., Washington, D. C.). 25 Jan 1972. Contract NASw-417. 37p. (NASA-CR-125673; TM-72-2015-1). NTIS.

An electrodynamic model for an ionosphere-solar wind interaction is developed based on the existence of a low-beta plasma below the anemopause. The currents for the interaction are driven by the solar-wind motional electric field and induce a stagnation magnetic field at the anemopause. For Venus and Mars, the lower region of the ionosphere near the electron density peak has the highest conductivity and, therefore, the tangential component of the induction current flows substantially in this region. The current paths close in the anemopause, which is a solar wind current sheath analogous to the magnetopause. Both the fraction of the undisturbed solar-wind motional electric field, which drives the induction current, and the required fraction of incident solar-wind particles, crossing the anemopause to produce this current, are shown to be small. (auth) (STAR)

46325 COSMIC-RAY ELECTRON SPECTRUM AND ITS MODULATION NEAR SOLAR MAXIMUM. Schmidt, P. J. (Univ. of Chicago). J. Geophys. Res.; 77: No. 19, 3295-3310 (1 Jul 1972).

A balloon-borne ionization spectrometer was used to measure the energy spectrum of cosmic-ray electrons in the range 20 MeV to 15 GeV. Balloon flights were made from Fort Churchill, Manitoba, Canada, in the summers of 1968, 1969, and 1970. Primary electron spectra were obtained for these three periods that show the effect of differing levels of solar modulation. In particular, a relatively strong decrease in the electron intensity between 300 MeV and 2 GeV was observed between 1969 and 1970, whereas neutron-monitor counting rates increased. It is shown that the data can be explained quantitatively by using the force-field approximation to the cosmic-ray transport equation if the rigidity dependence of the diffusion coefficient is generalized to be time-dependent. The parameters obtained in fitting the electron measurements, when they are applied to the 1968 proton energy spectrum, lead to general agreement with measurements under the assumption that the unmodulated proton spectrum is a power law in total energy. (43 references) (auth)

46326 HEAT CONDUCTION IN A TURBULENT MAGNETIC FIELD WITH APPLICATION TO SOLAR-WIND ELECTRONS.

Hollweg, Joseph V.; Jokipii, J. R. (California Inst. of Tech., Pasadena). *J. Geophys. Res.*; 77: No. 19, 3311-16(1 Jul 1972).

Random, long-wavelength fluctuations were considered in a turbulent magnetic field that can appreciably decrease the heat conductivity of a plasma along the magnetic field. In simple cases of interest, the reduction along the average field is approximately by the factor $(\cos \delta\theta)^2$, where $\delta\theta$ is the angle of the local magnetic field relative to the average field. Application to solar-wind electrons indicates that this reduction in heat conductivity due to observed fluctuations in the interplanetary magnetic field may be of the order of a factor of 2. This may help to explain recent measurements that indicate a rather low electron heat flux in the solar wind. (auth)

46327 RIGIDITY DEPENDENCE OF COSMIC-RAY MODULATION AT RIGIDITIES >2 GV IN AUGUST 1969. Stoker, P. H.; Raubenheimer, B. C.; van der Walt, A. J. (Univ. for C.H.E., Potchefstroom, S. Afr.). *J. Geophys. Res.*; 77: No. 19, 3575-82 (1 Jul 1972).

The modulation function $M = \ln(dN_0/dP)/(dN/dP)$ is determined relative to solar minimum (May 1965) from differential responses obtained from survey flights at 307 g/cm² in October 1964, August 1966, and August 1969. The results for 1964 and 1966 are compatible with a modulation function proportional to P^{-1} , but for August 1969 the modulation is a much stronger function of rigidity, being either $M = \Delta\eta/P^{1.6}$ or $M = \Delta\eta/P + 4.0/P^2$. However, the counting rates of ground-based neutron monitors at high, middle, and low latitudes relative to solar minimum, are better represented by the latter modulation function for the time after the June 1969 reduction in cosmic-ray intensity, the modulation term $\propto P^{-1}$ being solar-controlled as for the 11-year variation and the term $\propto P^{-2}$ being of a more persistent nature. (auth)

46328 VELOCITY AND FLUX DEPENDENCE OF THE SOLAR-WIND HELIUM ABUNDANCE. Hirshberg, J. (Ames Research Center, Moffett Field, Calif.); Asbridge, J. R.; Robbins, D. E. *J. Geophys. Res.*; 77: No. 19, 3583-8(1 Jul 1972).

The causes of the variations in the relative abundance of helium in the solar wind are not understood. To define the theoretical problem more precisely, empirical relationships between helium abundances and other pertinent solar-wind parameters are necessary. It is shown that the average percentage of helium increases with the solar wind velocity. The straight-line least-squares fit to the data is given by $n_{He}/n_p \times 10^2 = 1.02 \times 10^{-2} v - 0.39$, where $n_{He}/n_p \times 10^2$ is the mean percentage of helium averaged over a 25-km/sec data interval and v is the velocity of the center point of the interval in kilometers per second. It is confirmed that the abundance of helium tends to be higher for low solar-wind fluxes rather than lower as expected from theory. The interpretation of these results is discussed, and it is concluded that more realistic theoretical treatments of the problem are necessary before these results can be understood in terms of models of the solar corona and solar wind. (auth)

46329 DETECTION OF SOLAR-WIND ELECTRON PLASMA FREQUENCY FLUCTUATIONS IN AN OBLIQUE NONLINEAR MAGNETOHYDRODYNAMIC WAVE. Fredricks, R. W. (TRW Systems Group, Redondo Beach, Calif.); Scarf, F. L.; Russell, C. T.; Neugebauer, M. *J. Geophys. Res.*; 77: No. 19, 3598-3601(1 Jul 1972).

A nonlinear plasma-wave phenomenon was observed in the upstream interplanetary medium by Ogo 5 instrumentation on Jan. 30, 1969. Data are shown of the total ion flux, plasma-wave frequency, and total magnetic field. Source mechanisms for hf waves in the solar wind are discussed and show that changes in the wave growth conditions cannot readily account for significant frequency variations. The observed flux variations are related to changes in local density rather than to any large fluctuations in wind flow speed or direction. It is argued that the Doppler effect is always small for these wave modes and that this phenomenon cannot account for the wave frequency changes. It is concluded that the identification of the electron-plasma frequency emission whenever it is present in the solar wind provides a high-time-resolution measurement of local electron density. The technique represents an improvement over conventional probe techniques, which typically scan over energy on time scales of several seconds or longer. The availability of such quasi-instantaneous density measurements on common time scales with field measurements has a great potential impact on detailed theoretical interpretations. One such theoretical interpretation facilitated by the common time scale of density and magnetic field is exemplified by consideration of the large-amplitude oblique MHD wave and its associated large-amplitude density fluctuation. (J.C.W.)

46330 COMPARISON OF MEASUREMENTS OF THE CHARGE SPECTRUM OF SOLAR COSMIC RAYS FROM NUCLEAR EMUL-

SIONS AND THE EXPLORER 35 SOLID-STATE DETECTOR. Armstrong, T. P. (Univ. of Kansas, Lawrence); Krimigis, S. M.; Reames, D. V.; Fichtel, C. E. *J. Geophys. Res.*; 77: No. 19, 3607-12(1 Jul 1972).

Recent measurements relating to the He/M (medium) nuclei abundance ratio in solar cosmic rays from solid-state detectors carried aboard the lunar orbiting Explorer 35 are compared with earlier measurements using nuclear emulsions. The He/M ratio of 27 ± 9 obtained from Explorer 35 in the energy range of 0.5-~2.5-MeV/nucleon for solar events in 1967-1968 is lower than the value 58 ± 5 measured earlier for large events in the 1960-1969 period in the 12- to 200-MeV/nucleon range with emulsions. A mild energy dependence of the abundance ratio is one possible explanation of the observed differences. (auth)

46331 PROBLEMS IN SOLAR-SYSTEM PLASMA PHYSICS. Leer, Egil. San Diego, Calif.; Univ. of California (1972). 125p. University Microfilms Order No. 72-17,444.

Thesis.

The excitation of parametric instabilities by radio waves in a magnetoplasma is discussed. A uniform medium is assumed and linear approximations are used. A simple method, based on energy balance, is used first to re-derive the well-known threshold condition for the purely growing parametric instability in a homogeneous medium and later to estimate the effects of inhomogeneity in a semiquantitative manner. The threshold is then calculated in a more quantitative manner for the case where the effects of inhomogeneity dominate over those of collisions. The method has the advantages that its range of validity is easily checked, that it provides good physical insight and that it is easily applicable to electromagnetic instabilities. The flow of plasma from the Sun is discussed. A two-fluid model of the solar wind with anisotropic proton temperature and allowing for extended coronal proton heating is considered for the case of a purely radial and of a spiral magnetic field. Proton-proton Coulomb collisions together with a spiral interplanetary magnetic field are found to be sufficient to reduce the thermal anisotropy in the proton gas to a value in agreement with observations. Reasonable values are obtained for the flow velocity, number density and the proton temperature near the orbit of the Earth. Kinetic and hydrodynamic descriptions of a collisionless solar wind proton gas are compared. Heat conduction and viscosity are neglected in the hydrodynamic formulation but automatically included in the kinetic formulation. The results of the two models are very nearly the same, indicating that heat conduction and viscosity are not important in the solar wind proton gas beyond about 0.1 AU. It is concluded that the hydrodynamic equations provide a valid description of the collisionless solar-wind protons, and hence that future models of the quiet solar wind should be based on a hydrodynamic formulation. (TSS) (Diss. Abstr. Int., B)

46332 STATISTICAL TREATMENT OF THE RELATIONSHIP BETWEEN THE SOLAR FLARE-PRODUCING REGIONS AND THE SC + SI EVENTS. Moldovanu, A.; Bradu, E. B.; Tarnauceanu, D. (Centre of Technical and Physical Research, Iasi). *Gerlands Beitr. Geophys.*; 81: No. 1-2, 1-7(1972).

The validity of the hypothesis that both geomagnetic storms and Forbush decreases can be ascribed to the effects of interplanetary streams produced in the solar wind by solar-flare producing regions and not to the effects of individual solar flares (Ballif and Jones) was examined statistically. For this purpose, an adequate graphical representation is used for each solar rotation for the individual solar flares and the flare-producing regions. The dynamics of the flare-producing regions is more evident. The SC and SI events appear to be grouped morphologically as collective events, gr(SC + SI), supposing that each gr(SC + SI) event reflects the interaction between a tangential discontinuity and the magnetosphere. The main hypothesis assumes that if the tangential discontinuities are really related to the flare-producing regions, then for successive rotations, the variation of the number of flare-producing regions passing the solar central meridian must be followed by a parallel variation in the number of gr(SC + SI) events. The data obtained from the analysis of 19 solar rotations around the maximum of the solar cycle No. 20 (rotations 1839-1857) reveal a good correlation between the variation of the number of flare-producing regions, active at the central meridian, and the variation in the number of gr(SC + SI) events ($r \sim 0.9$). On such a statistical treatment, the working hypotheses are better proved than in the one-to-one treatment. Hence, the main cause of the geomagnetic disturbances, as the SC and/or SI events, appear to be the interplanetary streams or the tangential discontinuities, somehow related to the flare-producing regions. (auth)

46333 SOME STUDIES ON SOLAR MICROWAVE BURSTS OF DIFFERENT TYPES IN RELATION TO OPTICAL FLARES AND

OTHER ALLIED EVENTS. Das Gupta, M. K. (Univ. of Calcutta); Sarkar, S. K. *Indian J. Pure Appl. Phys.*; 10: No. 2, 153-60 (Feb 1972).

Occurrences of some common types of solar microwave bursts at 4995 and 8800 MHz were studied in relation to flares of different areas and intensities were examined for the period November 1966 to September 1969. It has been found that simple impulsive bursts have the maximum occurrences, and complex bursts show the maximum percentage association with solar flares. Further, both complex bursts and complex bursts with fluctuations show maximum association with bright flares while all other remaining types are in general associated with flares of normal intensity. Moreover, the variation of their percentage association with increase of both peak fluxes and energies of the bursts has also been examined. It has been noted that, except for post-burst increase, for all other types of bursts, percentage association values increase with increasing peak fluxes and energies. Complex bursts show relatively higher percentage associations in all the ranges. Of the different types of bursts, solar microwave great bursts are of vital importance for their close connection with different events. These great bursts during the period November 1966 to January 1970 have also been examined in relation to solar flares, solar meter-wave bursts, different types of sudden ionospheric disturbances,

and geomagnetic crochets. Distribution of the sources of great bursts over the solar disk shows a strong north-south asymmetry. Great bursts are highly associated with meter-wave type-III (fast drift) bursts. Great bursts show maximum association with bright flares. In general, these are highly associated with SIDs. Association with SPA is 100% and with SWF 97.9%. Association with crochets increases with the peak flux of the great bursts, the burst-associated crochets being mainly due to bright flares. Variation of the peak flux of great bursts with frequency has also been examined. (auth)

46334 SCIENTIFIC RESULTS FROM HEOS 1. Taylor, Brian G. (European Space Research and Tech. Centre, Noordwijk, Netherlands). *Raumfahrtforschung*; 16: No. 3, 101-7 (1972).

Selected major results of the eight scientific experiments carried aboard the HEOS-1 spacecraft were presented. This ESRO satellite was launched on December 5, 1968, into a highly eccentric orbit. Significant and unique results have been obtained during the following three years. The discussion of measurements especially regarding interplanetary shock and solar-wind interactions is complemented by an extensive list of literature comprising all papers about the scientific results published so far. (auth)

PHYSICS (ATOMIC AND MOLECULAR)

Atomic, Electron, Ion, and Molecular Beams and Their Reactions

Refer also to abstract: 47245.

46335 (FRNC-TH-198) CAPTURE OF LOW-ENERGY RARE GAS IONS BY METALLIC TARGETS. Roussel, Joseph (Paris Univ., Orsay (France)). 1971. 100p. (In French). Dep. NTIS (U. S. Sales Only).

Thesis.

Experimental results on the capture of the rare gases Ar and Ne by the surface of a monocrystalline Mo target bombarded with ions in a direction normal to the 110 plane are presented. Theoretical studies were also made. They consisted of determining the interaction potential between the atoms and of developing a capture model for the calculation of the capture coefficients. It is shown that a rare gas atom is captured, essentially, by a substitutional mechanism involving the ejection of the metal atom, the substituted atom being retained by the lattice if it is sufficiently surrounded. The spheres used to calculate the transparency of the atomic layers traversed (in the framework of a hard sphere approximation) correspond to the energy-loss cross sections; in particular the atom taking part in a collision too near to the head-on case does not have enough energy left to create for itself a capture site. (auth)

46336 SURFACE AND VACUUM PHYSICS INDEX. VOLUME 7, NO. 5, 1972. Bibliography, Author Index, Subject Index in Alphabetical Order, Cumulative Subject Index in Logical Order. Frankfurt am Main; Zentralstelle fuer Atomkernenergie-Dokumentation (1972). 74p.

This listing contains 256 references on various aspects of the subject. (M.O.W.)

46337 STUDY OF RELATIVE LEVEL POPULATIONS IN FOIL EXCITED He AND Li. Andersen, N.; Carriveau, G. W.; Glinska, A. F.; Jensen, K.; Melskens, J.; Veje, E. (Univ. of Copenhagen). Z. Phys.; 253: No. 1, 53-60 (1972).

A set of relative level population measurements was made for 40-keV beam-foil-excited He and Li. The relative populations are discussed as functions of the principal quantum number, the azimuthal quantum number, the spin, and the excitation energy. (auth)

46338 OPTIMIZATION OF LOW-ENERGY ELECTRON SCATTERING FOR PRODUCING A BEAM OF SPIN-POLARIZED ELECTRONS. Zeman, Herbert David. Stanford, Calif.; Stanford Univ. (1972). 99p. University Microfilms Order No. 72-16,821.

Thesis.

The means by which the highest possible output polarized electron beam quality may be achieved using low-energy electron scattering from gases are discussed. A particular source of polarized electron beams utilizing low-energy scattering of an electron beam produced by a thermionic arc electron gun from mercury atoms in an atomic beam is described. Measurements were taken with a primary beam energy of 50 eV at a scattering angle of 60° and with a primary beam energy of 170 eV at a scattering angle of 50°. Graphs of output polarization and intensity are given vs primary beam current for a variety of mercury atomic beam densities at both operating points. The best beam achieved had a current of 40 nA and a polarization of 20%. (Diss. Abstr. Int., B)

46339 ENERGY SHIFTS AND RELATIVE INTENSITIES OF K X RAYS PRODUCED BY SWIFT HEAVY IONS. Saltmarsh, M. J.; van der Woude, A.; Ludeman, C. A. (Oak Ridge National Lab., Tenn.). Phys. Rev. Lett.; 29: No. 6, 329-33 (7 Aug 1972).

The energy shifts of Fe, Co, Zr, and Sn K x rays produced by 5-MeV/amu He, C, O, Ne and 10-MeV/amu C ions depend nearly linearly on the stopping power for the bombarding ion. The shifts and the $K\beta/K\alpha$ ratios show a projectile energy dependence similar to that predicted for the L-ionization cross section. This suggests that these phenomena can be explained by existing Coulomb excitation theories. (auth)

Atomic and Molecular Properties

Refer also to abstracts: 46413, 46415, 46417, and 46523.

46340 (ORNL-TM-3820) PHOTOELECTRON CHEMICAL SHIFTS IN INORGANIC COMPOUNDS. McGuire, Gary Elder

(Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 103p. Dep. NTIS.

Thesis. Submitted to Tennessee Univ., Knoxville.

Selected inner-shell ionization energies of a variety of inorganic compounds were examined by using x-ray photoelectron spectroscopy. Attempts were made to relate the chemical shifts in the core level energies to structural and bonding properties of the compounds. Among the materials studied were UO_2 , UO_3 , and U_3O_8 . It was found that it was possible to distinguish the three oxides by photoelectron spectroscopy because the uranium core electrons have slightly different binding energies in each compound. It was also observed that the chemical shift from uranium metals to the uranium oxides was smaller than the shift observed in going from the metal to the metal oxide in the case of other metals. (A.G.W.)

46341 (UCRL-51231) THEORETICAL RADIATIVE TRANSITION RATES FOR K- AND L-SHELL X RAYS. Scofield, James H. (California Univ., Livermore, Lawrence Livermore Lab.). 1 Jun 1972. Contract W-7405-eng-48. 62p. Dep. NTIS.

Theoretical results covering the range of elements from $Z = 5$ to 102 are given for the radiative transition rates in filling vacancies in the K and L shells. The calculations use a relativistic Hartree model for the atoms. Given here are the ratios of the rates for the major K-shell x rays, the total radiative decay rate and theoretical binding energies for the K shell and L subshells, the transition rate and theoretical energy for each K x ray going by E-1, E-2, or M-1 radiation and each L x ray going by E-1 or E-2 radiation, the ratios of M-2 to E-1 rates, and the A_{22} angular correlation factor between K and L x rays. (auth)

46342 ELECTRIC FIELD EXPERIMENTS ON THE META-STABLE STATES OF HELIUM. Petrasso, Richard David. Waltham, Mass.; Brandeis Univ. (1972). 57p. University Microfilms Order No. 72-17,995

Thesis.

The atomic-beam magnetic-resonance technique was used to measure the relative Stark splitting between the $M = \pm 1$ and the $M = 0$ Zeeman sublevels of the 2^3S_1 state of helium. This shift should be zero in the LS coupling approximation. The shift is $(1.14 \pm 0.04) \times 10^{-1}$ Hz/(statV/cm)² and can equivalently be interpreted as giving rise to a tensor polarizability of $(3.41 \pm 0.11) \times 10^{-3}$ a.u. The shift decreases the normal magnetic splitting between the $M = +1$ and the $M = 0$ sublevels. Measurement of the transition rate of the normally metastable helium 2^1S_0 state in the presence of a uniform electric field by observing the attenuation of a metastable beam of helium passing between parallel electric field plates is also reported. This attenuation, when combined with measurement of the velocity distribution of the beam, yields a transition rate of $(0.926 \pm 0.020) F^2 s^{-1}$ where F is the field strength in kilovolts per cm. This agrees well with the calculated value of $(0.920 \pm 0.030) F^2 s^{-1}$ which is derived from a sum of oscillator strengths published by various workers. Both the experimental and theoretical values are in clear disagreement with the Bethe-Lamb theory, which predicts a transition rate of greater than $1.17 F^2 s^{-1}$ for the helium 2^1S_0 state. The Bethe-Lamb equations have been used extensively to calculate the electric-field quenching of hydrogen and hydrogen-like atoms in the $2^2S_{1/2}$ state. Applying the quench theory used for the helium calculation to the hydrogen $2^2S_{1/2}$ gives the same transition rate as the Bethe-Lamb theory to within 1%. The previous experimental checks of the Bethe-Lamb theory were made only on hydrogen and hydrogen-like atoms; the results of these experiments indicated that this theory was valid. However, the helium quench experiment shows unambiguously that the Bethe-Lamb equations are incorrect. (Diss. Abstr. Int., B)

46343 FINE STRUCTURE OF THE HELIUM NEGATIVE ION. Mader, D. L.; Novick, R. (Columbia Univ., New York). Phys. Rev. Lett.; 29: No. 4, 199-202 (24 Jul 1972).

Fine-structure transitions were observed in the metastable $(1s2s2p) ^4P_1$ state of the negative helium ion He^- using an rf resonance technique. Preliminary determinations were made of the energy separations: $|E_{3/2} - E_{1/2}| = 825.23 \pm 0.82$ MHz and $|E_{3/2} - E_{5/2}| = 8663 \pm 56$ MHz. It is shown that the $J = 3/2$ level lies between the $5/2$ and $1/2$ levels. These results are to be compared with the current theoretical estimates of $E_{3/2} - E_{1/2} = -2030$ MHz and $E_{5/2} - E_{3/2} = -9410$ MHz. (auth)

46344 HIGHER MULTIPOLE COEFFICIENTS OF THE HYDROGEN-HYDROGEN INTERACTION AS DETERMINED BY DYNAMIC MULTIPOLE POLARIZABILITIES. Venanzi, Thomas J. (Univ. of

California, Santa Barbara). *Phys. Rev. Lett.*; 29: No. 4, 197-9 (24 Jul 1972).

The Padé approximant is used to calculate the higher multipole coefficients of the hydrogen-hydrogen interaction, with excellent results. However, the rate of convergence is slower and is attributed to the fact that higher excited states contribute more in the higher multipole interactions than in the dipole-dipole case. (auth)

46345 ABSOLUTE MEASUREMENT OF THE LIFETIME OF THE 4^1D LEVEL OF THE HELIUM ATOM USING THE ELECTRON-PHOTON COINCIDENCE METHOD. Peresse, Joseph; Pochat, Amedee; Le Nadan, Andre (Univ., Brest, France). *Compt. Rend., Ser. B*; 274: No. 12, 791-4 (20 Mar 1972). (in French).

The coincidence method is applied to the inelastically scattered electron and the de-excitation photon in the measurement of the lifetimes of excited atomic states. The results obtained for the mean lifetime of the 4^1D level of helium are given. (France)

46346 ORIGIN OF ENERGY BARRIERS IN THE EXCITED STATES OF He_2 . Guberman, S. L.; Goddard III, W. A. (California Inst. of Tech., Pasadena). *Chem. Phys. Lett.*; 14: No. 4, 460-5 (15 Jun 1972).

The $A^1\Sigma_u^+$ and $C^1\Sigma_g^+$ states of He_2 were examined using self-consistent (spatially projected) generalized valence bond (GVB) wave functions. Both states were found to have humps (0.06 and 0.22 eV, respectively) at large R (3.1 and 2.1 Å, respectively). The repulsive nature of these states at large R results from non-bonding interactions between the singlet pairs of orbitals located on different centers. For R smaller than the size of the excited He orbital (2s or 2p), the state becomes attractive if the symmetry is such that the wave function can build in attractive He_2^+ ($^2\Sigma_u^+$) character. (auth)

46347 INVESTIGATION OF THE PRESSURE SHIFT OF THE GROUND STATE HYPERFINE LINES OF H , D , AND T . Morgan, Carlisle Lee Jr. New Haven; Yale Univ. (1971). 173p. University Microfilms Order No. 72-17,146.

Thesis.

An optical pumping method for measurement of hydrogen hyperfine pressure shifts was extended by improvements in sample preparation, temperature control, and magnetic field stabilization to yield results with approximately 0.1% accuracy in argon. The shifts for hydrogen were also measured for the first time in krypton and xenon. When the shift is expressed as $(1/\nu_0) [(d\nu)/d\rho] = (A + BT)$, the results in argon for hydrogen, deuterium, and tritium are $A_H = (-4.800 \pm .006) \times 10^{-9}/\text{torr}$ (0°C), $B_H = (0.956 \pm .011) \times 10^{-11}/\text{torr}$ (0°C)/ $^\circ\text{C}$, $A_D = (-4.802 \pm .022) \times 10^{-9}/\text{torr}$ (0°C), $B_D = (1.091 \pm .047) \times 10^{-11}/\text{torr}$ (0°C)/ $^\circ\text{C}$, $A_T = (-4.795 \pm .012) \times 10^{-9}/\text{torr}$ (0°C), and $B_T = (.929 \pm .021) \times 10^{-11}/\text{torr}$ (0°C)/ $^\circ\text{C}$. The fractional pressure shift for hydrogen is $(-10.4 \pm 0.2) \times 10^{-9}/\text{torr}$ (0°C) in krypton and $(-20 \pm 2) \times 10^{-9}/\text{torr}$ (0°C) in xenon between 32°C and 35°C . The results are compared with existing theories for the shift. The absence of any isotope dependence supports existing theories of the shift and encourages comparison with the measurements of muonium hyperfine splitting. The measured linear temperature dependence requires further development of the theoretical models of the shift. (Diss. Abstr. Int., B)

46348 RELAXATION PROCESSES OF THE $n = 3$ STATES OF HELIUM. Wellenstein, Hermann Fridenandt. Austin, Tex.; Univ. of Texas (1971). 107p. University Microfilms Order No. 72-15,851.

Thesis.

The positive column of a discharge through helium serves as a reactor for studying the relaxation processes of the $n = 3$ states. The critical conditions for appearance or disappearance of striations are expressed in terms of current, pressure, and tube radius as $I = A(pR)^{-5}$. The production mechanism of the molecular ion is studied by mass-analyzing the ion diffusion current through a hole in the side of the tube. Each of the $n = 3$ states is selectively and periodically modulated by absorption from the metastable states. By monitoring the in-phase changes of the excited states and molecular ion population over a range of discharge current and pressure, rate constants are determined for associative ionization, for excitation transfer by collision with neutrals and with electrons, and for the total neutral collisions. (Diss. Abstr. Int., B)

46349 ELECTRIC DIPOLE POLARIZABILITIES OF LANTHANIDES AND ACTINIDES. Saxena, K. M. S.; Fraga, S. (Univ. of Alberta, Edmonton). *J. Chem. Phys.*; 57: No. 4, 1800-1 (15 Aug 1972).

Electric dipole polarizabilities for the ground states of the lanthanides and actinides and their first three positive ions, calculated using numerical Hartree-Fock functions, are tabulated. Additional calculations were carried out for Ba and Rn for which experimental values are available. The agreement obtained between the present results and experimental values for the latter two elements indicates that the present values should represent a satisfactory prediction of the polarizabilities of the lanthanides and actinides. (A.G.W.)

46350 EFFECTS OF SCREENING AND POLARIZATION ON VIBRATION-ROTATION PROPERTIES OF H_2 . Hayes, Stuart F.; Beckel, Charles L. (Univ. of New Mexico, Albuquerque). *J. Chem. Phys.*; 57: No. 4, 1497-1502 (15 Aug 1972).

Eight simple models of H_2 in the ground state are examined through use of Dunham's formulas, numerical integration of the vibrational Schrödinger equation, and the WKB energy condition. These models provide four examples for studying the effects of variable screening and four examples for the effects of polarization on vibration-rotation properties. At small v , variability of the screening parameter α correlates with positive curvatures of the ΔG and B_v curves; i.e., for the four models with $\alpha = 1$ both curvatures are negative, and both curvatures are positive for the four cases with α variable. The dominance of the quartic potential coefficient a_2 in determining near-equilibrium ΔG and B_v curvatures is pointed out. Near dissociation, the presence of polarization causes positive curvature of these two curves. In showing this, large fictitious values of the reduced mass are employed to provide several vibrational levels very close to the dissociation limit. The D_v curves have a pronounced minimum, similar to that of the experimental curve, for α variable but not for $\alpha = 1$. In addition, the values of D_v and H_v are much closer to experiment for α variable than for $\alpha = 1$. (auth)

46351 LONG-RANGE INTERACTIONS BETWEEN TWO HYDROGEN ATOMS AND TWO HELIUM ATOMS. Yamabe, Tokio; Ishimaru, Shingo; Fukui, Kenichi (Kyoto Univ.). *Bull. Chem. Soc. Jap.*; 45: No. 5, 1384-7 (May 1972).

The numerical results of the interaction energy between two hydrogen atoms and two helium atoms are examined with the use of exact formulas for the second-order perturbation energy using the Unsöld approximation. It is found that the analytic expression gives a good long-range interaction energy between two hydrogen atoms comparable to the best variational results so far obtained. In the case of two helium atoms, the result lies in between experimental data for large nuclear distances; this seems to support the reliability of the present estimation. Finally, the effectiveness of the Unsöld approximation is remarked on. (auth)

46352 HYDROGEN ATOM IN A ROTATING ELECTRIC FIELD. Lisitsa, V. S. *Opt. Spectrosc. (USSR)* (Engl. Transl.); 31: No. 6, 468-9 (Dec 1971).

Translated from *Opt. Spektrosk.*; 31: No. 6, 862-5 (Dec 1971).

A calculation was performed of the emission spectrum of a hydrogen atom placed in a homogeneous electric field rotating with a constant angular velocity. The method used to solve the problem, based on the four-fold symmetry of the hydrogen atom, avoids the difficulties related to the secular-equation method and provides a solution for any line. It is shown that the method utilized here can be applied to the collision of hydrogen atoms with charged particles, in particular, to the theory of Stark broadening of hydrogen spectral lines in plasma. (auth)

46353 THEORETICAL STUDIES OF DIAMAGNETIC PROPERTIES OF THE HYDROGEN MOLECULE ION. I. APPROXIMATE VARIATION-PERTURBATION CALCULATION. Garrett, Thomas B.; Zeroka, Daniel (Lehigh Univ., Bethlehem, Pa.). *Int. J. Quant. Chem.*; 6: No. 4, 651-61 (Jul 1972).

A variation-perturbation method, analogous to a technique devised by Das and Bersohn for the study of the hydrogen molecule, is developed for calculation of the diamagnetic susceptibility χ and magnetic shielding σ and is applied to a study of the hydrogen molecule ion. Two approximations are investigated: in a first approximation, $\chi = -0.37814\alpha^2 a_0^3$ and $\sigma = 1.112 \times 10^{-5}$ cgs units and in a second approximation, $\chi = -0.37569\alpha^2 a_0^3$ and $\sigma = 1.128 \times 10^{-5}$ cgs units. (auth)

46354 THEORETICAL STUDIES OF DIAMAGNETIC PROPERTIES OF THE HYDROGEN MOLECULE ION. II. EFFECT OF VARYING INTERNUCLEAR SEPARATION. Garrett, Thomas B.; Zeroka, Daniel (Lehigh Univ., Bethlehem, Pa.). *Int. J. Quant. Chem.*; 6: No. 4, 663-8 (Jul 1972).

The diamagnetic susceptibility χ and magnetic shielding σ for H_2^+ are investigated in the range of internuclear separations $R = 1.6$ to 2.4 au according to a previously reported technique. From these data values of $\langle \chi \rangle$ and $\langle \sigma \rangle$, for which nuclear motion due to zero-point vibration and centrifugal stretching is taken into ac-

atoms are calculated at 300°K. These averages are $\langle \chi \rangle = -0.3902$ a.u., and $\langle \chi^2 \rangle = 1.096 \cdot 10^{-3}$ a.u., which are approximately 3.2% and 1.4%, respectively, smaller than the equilibrium values. (auth)

46356 SIMPLE PERTURBATION AND PERTURBATION-VARIATION TREATMENTS OF THE $1s\sigma_z$ AND $2p\sigma_z$ STATES OF H_2^+ . Tung, Eleanor W.; Sanders, William A. (Catholic Univ. of America, Washington, D. C.). *Int. J. Quant. Chem.*; 6: No. 4, 1972.

First-order wave functions and binding energies of the $1s\sigma_z$ and $2p\sigma_z$ states of H_2^+ are calculated by simple methods for internuclear separations between 0.2 and 10 a.u. An exact perturbation treatment of the $1s\sigma_z$ state with a zeroth-order function of the form $N \exp(-sR/2) \exp(sR/2)$ yields only fair results. An alternative method starts with zeroth-order functions of the form $N \exp(-sR/2) \exp(-sR/2) = \exp(sR/2)$ for the $1s\sigma_z$ and $2p\sigma_z$ states, respectively. An approximate first-order trial function is set up and the energies are determined variationally. For both states the results are comparable to those obtained by exact perturbation treatments of the same order. (auth)

46358 MULTIPHOTON TRANSITION IN HYDROGEN BETWEEN 1S AND 2P LEVELS. Rahman, Naseem K.; Reiss, Howard R. (American Univ., Washington, D. C.). *Phys. Rev. A*; 6: No. 3, 1252-5 (Sep 1972).

1S-2P multiphoton transition amplitudes are calculated for hydrogen atoms in the presence of an intense electromagnetic field. Nonlinear behavior of the transition amplitudes appears in the case of 1S-2S transitions considered earlier. (auth)

46357 RETARDED VAN DER WAALS POTENTIAL BETWEEN PAIRS OF SPINLESS ATOMS. Au, Chi-kwan E. (Columbia Univ., New York). *Phys. Rev. A*; 6: No. 3, 1232-8 (Sep 1972).

The retarded van der Waals potential between pairs of spinless atoms, H-H, H-He, He-He, Ne-Ne, Ar-Ar, and Ar-Ar, was computed with covariant formulas of Feinberg and Sucher. Electric and magnetic effects were treated on equal footing. For hydrogen, which possesses a large magnetic polarizability, magnetic effects dominate for large distance of separation, and for small distances, an additional contribution to the London constant of a pair of hydrogen atoms from the magnetic effects is found to be 0.136, about 0.24% of the currently accepted value. (auth)

46359 MOLECULAR-BEAM MAGNETIC-RESONANCE MEASUREMENTS OF THE ANISOTROPIES OF THE ELECTRIC POLARIZABILITIES OF H_2 AND D_2 . MacAdam, K. B.; Ramsey, N. F. (Harvard Univ., Cambridge, Mass.). *Phys. Rev. A*; 6: No. 3, 398-306 (Sep 1972).

The anisotropies $\gamma = \alpha_{||} - \alpha_{\perp}$ of the static electric polarizabilities of H_2 and D_2 were measured in a molecular-beam magnetic-resonance experiment. The quadratic Stark shifts of the separated-oscillatory-field resonances in the ground electronic and rotational and $J = 1$ rotational states were measured in parallel electric and magnetic fields. The results are $\gamma(H_2) = -0.16 \pm 0.005 \text{ Å}^3$ and $\gamma(D_2) = 0.2917 \pm 0.0004 \text{ Å}^3$, which agree very closely with theory. A previously measured combination of the spin-rotation and quadrupole-interaction constants of D_2 was independently remeasured, and the dependence upon orientation of the nuclear magnetic shielding of D_2 was found to be $\sigma_{\perp} - \sigma_{||} = -11.9 \text{ ppm}$. An efficient electron-bombardment detector was constructed for this experiment. (auth)

46360 ANALYTIC CALCULATION OF THE COHERENT X-RAY SCATTERING FACTOR FOR THE GROUND STATE OF THE HELIUM ISOELECTRONIC SEQUENCE. Sahni, V. (Polytechnic Inst. of Brooklyn); Krieger, J. B. *Phys. Rev. A*; 6: No. 3, 328-36 (Sep 1972).

The analytical method developed by the authors for the determination of the expectation value of single-particle operators $\hat{O} = \sum_i \hat{O}_i$ with respect to second order is employed in both the first and second decoupling approximations to obtain analytic expressions for the coherent x-ray scattering factor $F(\mathbf{k})$ for the ground state of the helium isoelectronic sequence valid for all values of momentum transfer. The trial wave function ψ_{trial} employed is an energy-minimized Hartree product of hydrogenic states. For helium the results for the form factor in the second decoupling approximation are found to be superior to those of the first, are within 1.2% of the highly accurate values calculated using a 120-parameter configuration-interaction wave function, and have an accuracy equivalent to that of an analytical Hartree product treatment. This error is further reduced as the atomic number is increased. In order to demonstrate the internal self-consistency of the technique, it is shown that the expectation value of a single-particle operator as obtained by direct use of the analytical method is the same as the expectation value

obtained employing the form factor, provided that the latter has also been calculated by the analytical method employing the same trial wave function. Finally, the calculations are extended to the infinite-momentum-transfer range and the results are studied in this limit by employing a cusp condition for the exact ground-state wave function of two-electron atomic systems written in terms of the logarithmic derivative of the electron density at the origin. It is observed that in the infinite-momentum-transfer limit the results for helium are in error by 0.57% and that this error is further diminished for each successive element of the isoelectronic sequence. In addition, it is noted that the cusp condition is exactly satisfied via the formalism of the second decoupling approximation and is independent of the variational parameter employed. (auth)

46360 SECOND-ORDER CORRECTIONS TO THE FINE STRUCTURE OF HELIUM. II. CONTRIBUTIONS FROM ^1P AND ^3D INTERMEDIATE STATES. Hambro, Lars (Univ. of California, San Diego, La Jolla). *Phys. Rev. A*; 6: No. 3, 865-70 (Sep 1972).

The contributions from the three spin-dependent Breit operators in second-order perturbation theory are calculated when the symmetries of the intermediate states are odd ^1P or ^3D . Standard Hylleraas expansions with up to 165 terms are used for the perturbations of the wave functions. The large interval of the fine structure of the 2^3P level in helium is increased by $2.17(2) \times 10^{-4} \text{ cm}^{-1}$ by ^1P and by $0.0089(6) \times 10^{-4} \text{ cm}^{-1}$ by ^3D . Two methods for handling angular integration over D tensors are described. (auth)

46361 LINEAR STARK EFFECT DUE TO RESONANT INTERACTIONS OF STATIC AND DYNAMIC FIELDS. Cohn, Arthur (Air Force Cambridge Research Labs., Bedford, Mass.); Bakshi, P.; Kalman, G. *Phys. Rev. Lett.*; 29: No. 6, 324-7 (7 Aug 1972).

The line shape of the Lyman- α transition of a hydrogen atom simultaneously subjected to a static and a perpendicular high-frequency electric field was investigated. It was found that resonance effects occur, producing a profile substantially different from the Stark spectrum of the fields acting independently. This result shows that the interpretation of turbulent hydrogen plasma spectra by the Blokhintsev (dynamic field only) theory is not generally valid. (auth)

Positronium, Muonium, and Mesic Atoms and Molecules

Refer also to abstracts: 46372, 46571, 46708, 47031, and 47245.

46362 (RISO-254) POSITRON LIFETIMES IN WATER AND ICE, AND IN FROZEN AQUEOUS SOLUTIONS. Eldrup, Morten (Danish Atomic Energy Commission, Risoe, Research Establishment, Tekniske Højskole, Lyngby (Denmark)). Nov 1971. 115p. Dep. NTIS (U. S. Sales Only).

Thesis.

An introduction to the subject is given, the principle of lifetime measurements is discussed, and the experimental apparatus is described. A computer program used for data analysis is described and the problems involved in the analysis are discussed. All lifetime spectra are resolved into three components. For H_2O the long lifetime, attributed to ortho-positronium, is 0.68 ns in ice between -196 and 100°C, increases to 1.12 ns at 0°C, and abruptly to 1.86 ns upon melting. Its relative intensity is 52% in ice and 27% in water. Spectra for heavy ice are identical with those for light ice, whereas heavy water gives 2.01 ns, 22%. The increase in lifetime in ice above -100°C is attributed to trapping of positronium in temperature-created defects. Spectra for frozen aqueous solutions of concentrations above 10^{-7} mole fraction contain at -160°C a long-lived component attributed to the trapping of positronium. Very good agreement with the trapping model is obtained for HF solutions, both fast frozen and annealed. An irreversible change in the spectrum takes place by heating a fast frozen HF solution above approximately -125°C. (112 references) (auth)

46363 PHASE AND TEMPERATURE DEPENDENCE OF POSITRON ANNIHILATION IN LIQUID CRYSTALS. Nicholas, J. Blair; Ache, Hans J. (Virginia Polytechnic Inst. and State Univ., Blacksburg). *J. Chem. Phys.*; 57: No. 4, 1597-1603 (15 Aug 1972).

Positron lifetime measurements were carried out in a series of liquid crystalline compounds. The resulting lifetime spectra were decomposed into two components. The short-lived component (lifetime τ_1 and intensity I_1) was attributed to free-positron and para-positronium annihilation where the long-lived component (τ_2, I_2)

was considered to be the result of orthopositronium annihilation. Changes of τ_2 and I_2 during mesomorphic phase transitions are generally very small and can be related to the transition entropies involved, supporting the contention that the positron annihilation process is affected by variations in the structural regularities of the environment. Various mechanisms that could account for the observed changes of the positron lifetimes and their intensities at the phase-transition points are discussed. (auth)

Collision Phenomena

Refer also to abstracts: 46280, 46348, and 46721.

46364 (ORNLM-TM-3839) LOW-ENERGY ELECTRON ATTACHMENT TO MOLECULES AT HIGH DENSITIES. McCorkle, D. L.; Christophorou, L. G. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 87p. Dep. NTIS.

Thesis. Submitted by D. L. McCorkle to Tennessee Univ., Knoxville.

Low-energy electron attachment to a number of polyatomic molecules was studied using the electron swarm method. Electron-attachment rates were measured for sulfur hexafluoride (SF_6) and molecular oxygen (O_2). Electron-attachment cross sections, determined by assuming an analytic form of the cross section, for anthracene, 1,2-benzanthracene, 1,4-naphthoquinone, and sulfur hexafluoride are reported and discussed. A new swarm-unfolding method for determining electron-attachment cross sections was developed. This was applied for the determination of the attachment cross sections for carbon tetrachloride, n-bromodecane, 1,4-naphthoquinone, and 1,2-benzanthracene, which increase sharply at thermal energies. Also, it was applied for the determination of the cross sections for the production of O^- from O_2 and O^- from N_2O , which lie at higher energies. The attachment cross section for the formation of O_2^- with increasing N_2 density in O_2 - N_2 mixtures was determined. At low densities this cross section exhibits distinct structure attributed to electron capture into vibrational states of O_2^- . (auth)

46365 (ORO-2911-14) INTERACTION OF HEAVY keV IONS WITH MATTER. Progress Report. Boring, John W. (Virginia Univ., Charlottesville, Research Labs. for the Engineering Sciences). Jun 1972. Contract AT(40-1)-2911. 27p. (AEEP-3930-109-72-U). Dep. NTIS.

Work during the period in four areas is summarized: (1) a critical review of calculations of the effects of heavy-particle penetration, including the relative importance of electronic and nuclear energy transfer processes; (2) investigation of the use of the binary-encounter method for calculating electronic stopping cross sections; (3) a study of the dependence of nuclear trajectories in heavy-particle collisions on the inelastic character of the events; and (4) use of the binary-encounter method for calculating the inactivation cross section for enzyme molecules. Figures show results on the following: comparison of three potentials (Bohr, Thomas-Fermi, and modified Thomas-Fermi) for calculation of average energy going to electronic processes in Ar-Ar and U-U collisions; electronic stopping cross sections for 0.1-, 0.8-, and 10.0-MeV helium as a function of target Z; calculated reduced cross sections for H^+ -Ar and H^+ -Kr scattering; cross sections for inactivation of RNase by protons at 0.1 to 100 keV. (A.G.W.)

46366 IONIZING COLLISIONS OF FAST ALKALI ATOMS WITH Cl_2 , Br_2 , AND O_2 . Kashihiro, Nobuyuki. Lafayette, Ind.; Purdue Univ. (1971). 147p. University Microfilms Order No. 72-7976.

Thesis.

A time-of-flight apparatus is described for measuring the inelastic energy loss incurred by fast (<50 to 350 eV) alkali atoms in ionizing collisions with diatomic molecules. The inelastic energy losses were found to be nearly energy-independent for collisions of K and Cs with Cl_2 , Br_2 , and O_2 , and it is furthermore shown that these energy losses are from about 8 to 10 eV, irrespective of the reactants. The collisions between alkali atoms and halogen molecules (Cl_2 and Br_2) yield intermediate alkali ions (M^+) and excited molecular ions (X_2^+) in the $2\Sigma_g$ state, followed by dissociation to the ground-state halogen ions and excited halogen atoms. In the case of oxygen, excited molecular ions in the $2\pi_u$ state are formed, and these species either remain in the bound state of the $2\pi_u$ or decay to vibrationally excited molecular states ($^b\Sigma_g^+$, $^a\Sigma_g^+$). The excitation functions (change of cross sections with incident energy variations) are interpreted in terms of curve crossings near or inside the equilibrium distance of the M- X_2 potential surfaces. A comparison of the results obtained at high energies (20 to 200 eV cm) is made to those known from collision energies between 1 and 20 eV (cm), and a composite excitation function for

the three most probable processes over the whole energy range is constructed. (Diss. Abstr. Int., B)

46367 ENERGY AND ANGULAR DISTRIBUTIONS OF SECONDARY ELECTRONS RESULTING FROM IONIZING COLLISIONS OF ELECTRONS WITH HELIUM AND KRYPTON. Oda, Nobuo; Nishimura, Fumio; Tahira, Sadamu (Tokyo Inst. of Tech.). J. Phys. Soc. Jap.; 33: No. 2, 462-7 (Aug 1972).

Cross sections differential in angle and energy were measured for secondary electrons resulting from ionizing collisions of electrons with helium at primary energies $E_0 = 500$ eV and 1 keV and with krypton at $E_0 = 1$ keV. The total spectra, consisting of the continuum spectra due to direct ionizations and the sharp line spectra due to indirect ionizations (autoionizations or Auger processes), were measured with an energy resolution of about 0.2 eV over the angular range from 10° to 130° . The continuum spectra are derived from the total spectra for electron energies from 25 to 45 eV for helium and from 24 to 52 eV for krypton and compared with calculations by the binary encounter theory. (auth)

46368 OSCILLATIONS OF THE CHARGE-EXCHANGE CROSS SECTION IN THE SYSTEM $\text{Li}^+\text{-He}$. Latypov, Z. Z.; Shaporenko, A. A. (Ioffe Inst. of Physics and Tech., Leningrad). Sov. Phys.-Tech. Phys. (Engl. Transl.); 17: No. 1, 114-16 (Jul 1972).

Translated from Zh. Tekh. Fiz.; 42: No. 1, 147-50 (Jan 1972).

The cross section for the charge-exchange process $\text{Li}^+ + \text{He} \rightarrow \text{Li} + \text{He}^+$ was studied experimentally over the range of Li^+ energies $600 \leq E \leq 2800$ eV; the cross section oscillates. The relationship between these oscillations of the charge-exchange cross section and those of the helium-atom excitation cross section in the process $\text{Li}^+(1s^2) + \text{He}(1s^2) \rightarrow \text{Li}^+(1s^2) + \text{He}(1s2s)$ is discussed. (auth)

46369 RELATIONS BETWEEN SOME EXPONENTIAL APPROXIMATIONS IN ROTATIONALLY INELASTIC MOLECULAR COLLISIONS. Pack, R. T. (Brigham Young Univ., Provo, Utah). Chem. Phys. Lett.; 14: No. 4, 393-5 (15 Jun 1972).

Relationships between the infinite-order sudden (or strong-coupling), the first-order sudden, and the exponential distorted-wave approximations are derived and discussed. The resulting 6 matrices for an atom-diatom rotationally inelastic collision are compared. (auth)

46370 COUPLED STATE CALCULATIONS FOR PROTON-HYDROGEN SCATTERING IN THE READING REPRESENTATION. Zimmerman, Ivan Harold. Seattle; Univ. of Washington (1972). 318p. University Microfilms Order No. 72-20,904.

Thesis.

The problem of proton scattering by atomic hydrogen is considered in the semiclassical impact-parameter approximation. The electron wave function is written as a sum over a specific set of expansion functions, which represent channels through which the electron probability may flow during the course of the collision. These channels are all closely coupled by the hamiltonian for the system and are centered on the two protons. An introductory chapter attempts to give both a historical and a theoretical background to the impact-parameter approximation. In the second chapter the coupled equations are derived. Symmetries of the system are utilized in reducing the amount of work to be done. The radial part of the expansion basis function $\phi_k(r)$ is chosen to be a Slater orbital $r^{\lambda_k} e^{-\lambda_k r}$, but with the parameter λ_k chosen according to particular criteria. With the set of λ_k 's so chosen, the radial functions are called Reading functions, and the representation of the electron wave function in terms of these orbitals is called the Reading representation. Using this representation, simplified expressions for the numerically evaluated exchange matrix elements may be obtained, which greatly increase the speed of computation. An additional refinement to the program is the utilization of the time-transition matrix $U^\pi(t, t_0)$, which relates $a_k^\pi(t)$, the probability amplitude for occupation of the k th orbital with parity symmetry $\pi = \pm 1$, to the earlier amplitudes $a_k^\pi(t_0)$. A complete set of preliminary results is obtained and compared to existing experimental and theoretical work. It is seen that a 6-state expansion, as was used in deriving the results in chapter IV, is not sufficient to guarantee good general accuracy and that convergence of calculated results is not apparent until some 7 states are included in the $l = 0$ angular momentum symmetry with perhaps 4 states in the $l = 1$ spectrum for each of $m = 0$ and $m = 1$. In chapter V an extensive program of large-scale computation is proposed, and refinements to the theoretical formulation of the problem are suggested and considered. (TSS) (Diss. Abstr. Int., B)

46371 OPTICAL EXCITATION FROM LOW-ENERGY $\text{He}^+ + \text{H}_2$ AND $\text{H}_2^+ + \text{He}$ COLLISIONS. Nathan, Richard David. Gainesville, Fla.; Univ. of Florida (1971). 135p. University Microfilms Order No. 72-16,638.

Thesis.

Observations of spectra and measurements of cross sections and polarization of emission arising from collisions of $\text{He}^+ + \text{H}_2$ and $\text{H}_2^+ + \text{He}$ were made in order to examine the processes of direct excitation, dissociative excitation, dissociative charge-transfer excitation, and charge exchange into excited states. Optical spectra from both collision processes, investigated in the wavelength region from 3500 to 7500 Å, revealed strong production of the Balmer series of atomic hydrogen and lines of neutral helium. Cross sections were determined from 30 to 700 eV for Balmer α and β emissions and for the $3^3\text{D} \rightarrow 2^3\text{P}^0$, $4^3\text{D} \rightarrow 2^3\text{P}^0$, and $3^1\text{D} \rightarrow 2^1\text{P}^0$ transitions of helium from $\text{He}^+ + \text{H}_2$ and for Balmer α , β , and γ and the $3^1\text{D} \rightarrow 2^1\text{P}^0$ and $3^3\text{D} \rightarrow 3^3\text{P}^0$ transitions of helium from $\text{H}_2^+ + \text{He}$ collisions. The Balmer α and β curves from the $\text{He}^+ + \text{H}_2$ reaction were the only ones that exhibited maxima in the low-energy region investigated. Measurements of the Balmer α emission cross section as a function of energy of the ionizing electrons in the case of $\text{H}_2^+ + \text{He}$ indicated the necessity of considering the vibrational state of the projectile in the dissociative excitation process. The polarization of Balmer α and β emission was determined for both reactions from bombarding energies of 40 to 700 eV. The detection of Balmer α polarization arising from $\text{He}^+ + \text{H}_2$ collisions is the first observation of polarized emission resulting from the process of dissociative charge transfer. (Diss. Abstr. Int., B)

46372 POSITRON-HYDROGEN COLLISIONS AND POSITRONIUM FORMATION. Abu-Wakid, Shukri Edward, Baton Rouge, La.; Louisiana State Univ. (1971). 120p. University Microfilms Order No. 72-17,741.

Thesis.

In Part I a modified close-coupling method is suggested in order to incorporate polarization effects in low-energy scattering. It is applied to $e^+ - \text{H}$ collisions for energies below the $n = 3$ threshold of hydrogen. S-, p-, and d-wave elastic phase shifts and inelastic cross sections are calculated. Results show a fair improvement over those of close coupling. In Part II the algebraic close-coupling method is used to calculate the s-wave phase shifts and positronium-formation cross section for various types of close-coupling expansions of the total wave function. The positronium channels are included explicitly, and polarization effects in both the hydrogen and positronium atoms are incorporated by pseudostates that give the full ground-state polarizabilities. (Diss. Abstr. Int., B)

46373 SECOND BORN ZERO-ANGLE DIFFERENTIAL CROSS SECTIONS IN HIGH-ENERGY LIMIT FOR ELECTRON-IMPACT EXCITATION OF He: THE $2^3\text{S} - 1^4\text{S}$ AND $2^3\text{P} - 1^4\text{S}$ TRANSITIONS. Bonham, R. A. (Indiana Univ., Bloomington). J. Chem. Phys.; 57: No. 4, 1604-12(15 Aug 1972).

The high-energy limit of the inelastic differential cross section for the excitation of the 2^3S and 2^3P states of He from the ground state in the forward direction is investigated by expanding the T matrix in powers of V about zero scattering angle and considering all terms proportional to V and V^2 where V is the interaction potential. It is shown that the leading terms in a power series in the reciprocal of the incident energy in this second Born infinite-channel close-coupling theory can qualitatively account for the observed enhancement of the exchange cross section in the forward direction over the predictions of the first Born theory. The ratio of the square of the leading second Born term to the square of the leading first Born term is shown to be proportional to the incident energy for the $2^3\text{S} - 1^4\text{S}$ transitions and of the same order in the incident energy as the first Born form in the case of the $2^3\text{S} - 1^4\text{S}$ transition. The value of the ratio at 500 eV for the $2^3\text{S} - 1^4\text{P}$ transition is estimated to be close to 100 and is relatively independent of the choice of an effective energy-loss parameter introduced to carry out the sum over intermediate states. (auth)

46374 IONIZATION OF ARGON BY ELECTRON IMPACT. Crowe, A.; Preston, J. A.; McConkey, J. W. (Univ. of Windsor, Ont.). J. Chem. Phys.; 57: No. 4, 1620-5(15 Aug 1972).

The ionization efficiency curves for Ar^+ and Ar^{2+} were measured for electron energies from threshold to 300 eV. Structure in the Ar^+ curve was confirmed and arguments and evidence are put forward to support the view that this structure is due to production of the ion via various channels (in particular the auto-ionization channels) rather than to "competition" between the Ar^+ and Ar^{2+} channels as has been suggested recently. (auth)

46375 OPTICAL MODEL FOR VIBRATIONAL RELAXATION IN REACTIVE SYSTEMS. Poulsen, Lise Lotte; Ross, John; Steinfeld, Jeffrey I. (Massachusetts Inst. of Tech., Cambridge). J. Chem. Phys.; 57: No. 4, 1592-7(15 Aug 1972).

The probabilities for vibrational transitions in the presence of open reactive channels are calculated with an optical model. A linear collision of an atom and a diatomic molecule is considered. The distorted-wave Born series is derived, and an analytic ex-

pression is given for the first-order result for an exponential repulsive potential. A semiclassical analysis is also presented. The results are applied to an electronically excited I_2 molecule colliding with an inert atom. The vibrational transition probabilities are found to be appreciably affected only in the few cases where the predissociation probability is very high, that is larger than about 0.3. (auth)

46376 ACOUSTIC MEASUREMENTS OF ROTATIONAL ENERGY TRANSFER. Bauer, H.-J.; Bass, H. E. (Univ. of Mississippi, University). J. Chem. Phys.; 57: No. 4, 1763-6(15 Aug 1972).

The problem of using acoustic measurements to determine rotational energy transfer rates was examined. A method to link the rate of energy transfer to the acoustic behavior of the gas was developed and applied to the collision model of Raff and Winter. The results in the case of H_2 demonstrate the importance of using an accurate acoustical treatment when testing a reaction model. (auth)

46377 RETARDATION EFFECTS IN LOW-ENERGY ATOM-ATOM SCATTERING. Konrady, John Andrew Jr. College Park, Md.; Univ. of Maryland (1972). 92p. University Microfilms Order No. 72-20,254.

Thesis.

The feasibility of detecting retardation effects in low-energy helium-helium scattering is studied. Differential and total cross sections for the elastic scattering of helium atoms at very low energies are computed with and without retardation effects considered. Differences between the retarded and unretarded total and forward differential cross sections exceed 3% in an energy range in which low-resolution experiments have already been performed. The calculations are based on the Bruch-McGee semi-empirical potential and a modified version of this potential utilizing the O'Carroll-Sucher arc-tangent approximation to the Casimir-Polder potential. New bounds on the second term in Born series for the partial wave amplitudes are presented. Also, a regularization scheme for the Born approximation to the scattering amplitudes for singular potentials is developed. (Diss. Abstr. Int., B)

46378 FORMATION OF EXCITED HYDROGEN ATOMS BY CHARGE-TRANSFER NEUTRALIZATION OF HIGH-ENERGY PROTONS. Ford, John Charles. Atlanta; Georgia Inst. of Tech. (1971). 182p. University Microfilms Order No. 72-19,703.

Thesis.

Absolute cross sections for the formation of atomic hydrogen in the 3s, 3p, and 3d states of excitation resulting from the impact of 75- to 350-kilovolt protons with various gaseous targets such as hydrogen, nitrogen, argon, and helium were measured with greater accuracy than heretofore possible. In addition, a number of more complicated molecular targets such as methane, ethane, propane, carbon dioxide, carbon monoxide, and nitric oxide were used to produce 3s-state excitation with the objective of establishing simple empirical rules for predicting molecular cross sections in terms of the cross sections for the constituent atoms. Considerable effort was directed toward providing a vivid understanding of many of the associated processes affecting the formation and decay of excited projectiles. Finally, a study of the formation of excited hydrogen by the dissociation of H_2^+ and H_3^+ projectiles was made. A collimated beam of protons was allowed to traverse a differentially pumped target cell. The Balmer alpha radiation from the emerging beam displayed a distinctive spatial profile. From this profile the relevant cross sections were inferred via a mathematical deconvolution procedure. The relative production of the 3s, 3p, and 3d states of excitation was found to be nonstatistical; further, it was found that production of lower angular momentum states is preferred. The ratios of the cross sections for producing 3s, 3p, and 3d states by charge transfer are in disagreement with existing theory for the neutralization of protons by helium. No simple rules were found for predicting the production of 3s hydrogen by the charge-transfer neutralization of protons by complicated molecular targets. The dissociation of H_2^+ and H_3^+ was found to produce an abundance of 3p and 3d state hydrogen; this contrasted strongly with the production of hydrogen by charge transfer. (Diss. Abstr. Int., B)

46379 QUANTUM-MECHANICAL SCATTERING OF ASYMMETRIC MOLECULES. II. COMPARISON BETWEEN THE SYSTEMS $\text{He}-\text{D}_2$ AND $\text{He}-\text{HT}$. Heukels, W. F.; van de Ree, J. (University of Tech., Eindhoven, Netherlands). J. Chem. Phys.; 57: No. 4, 1393-99(15 Aug 1972).

Exact quantum mechanical coupled-channel calculations were performed in order to compare the rotating potential model for heteronuclear molecules with an asymmetric mass distribution and the usual potential with anisotropy terms for homonuclear molecules. The scattering cross sections of $\text{He}-\text{D}_2$ and $\text{He}-\text{HT}$

show no difference in shape, apart from some quenching of the orbiting resources. There is, however, a significant difference in absolute magnitude, due predominantly to inelastic processes. In the inelastic cross sections, the effect of the shift of the center of mass from the center of interaction in asymmetric molecules counteracts the effect of a change in moment of inertia. (auth)

46380 SCATTERING BY AN r^{-2n} POTENTIAL IN THE BORN APPROXIMATION AND APPLICATION TO DIPOLE-QUADRUPOLE SCATTERING. Ganas, P. S. (California State Coll., Los Angeles). *J. Chem. Phys.*; 57: No. 4, 1406-7(15 Aug 1972).

By using the Born approximation, an analytic formula is derived for the phase shifts δ_l for scattering by a potential with an r^{-2n} tail. The low-energy properties of the phase shifts are discussed, and an expansion of $\tan \delta_l$ in powers of k is presented. The dipole-quadrupole r^{-8} potential is discussed as a special case. (auth)

46381 MULTIPLE-SCATTERING EFFECTS IN HIGH-ENERGY ELECTRON-MOLECULE COLLISIONS. II. POLY-ATOMIC MOLECULES. Yates, Albert C. (Indiana Univ., Bloomington). *J. Chem. Phys.*; 57: No. 4, 1686-93(15 Aug 1972).

The Glauber theory of multiple scattering is applied to the elastic collisions of fast electrons with polyatomic molecules in an effort to explain deviations between theory and experiment in recent molecular structure determination by electron diffraction. The full scattering amplitude is developed inclusive of all double-scattering contributions, with the differential cross section for the process given up to terms representative of the interference between single and double scattering. For sufficiently energetic incident electrons, the multiple-scattering contributions to the differential cross section are shown to reduce to simple analytic forms. In particular, the three-atom single-double interference term is shown to be an oscillatory function of momentum transfer. Application of the approximate formulas to the rhenium hexafluoride molecule gives excellent agreement with the experimental results of Jacob and Bartell. (auth)

46382 MONTE CARLO TRAJECTORY CALCULATIONS OF THE THREE-BODY RECOMBINATION AND DISSOCIATION OF DIATOMIC MOLECULES. Shui, Ven H. (Massachusetts Inst. of Tech., Cambridge). *J. Chem. Phys.*; 57: No. 4, 1704-17(15 Aug 1972).

The modified phase-space theory of reaction rates was applied to the problem of the three-body recombination and dissociation of diatomic molecules. The results illustrate the important influence of the weak attractive minimum in the third-body interaction potential and the effect of barrier penetration for recombination at low temperatures. The system $N_2 + Ar$ was used as a typical illustrative example, and good agreement was obtained between the theoretical predictions and the experimental measurements of the reaction rate coefficients over the temperature range 200 to 12,000°K. The recrossing factor and the nonequilibrium factor were obtained from Monte Carlo trajectory calculations for states near the dissociation limit. The trajectories were sampled within the reaction zone with a weight proportional to the equilibrium reaction rate and numerically integrated in both timewise directions to determine the complete histories of the collisions. A simple, separable function for the equilibrium transition rate $R(\epsilon_i, \epsilon_f)$ from initial energy states ϵ_i to final energy states ϵ_f , which could be characterized by three parameters, was obtained to fit the numerical data and was used to solve the steady-state master equation. Distributions of the trajectories with respect to energies and impact parameters are presented, and several typical trajectories are shown to illustrate the important features of the collisions. The contribution to the overall reaction rate from the "complex mechanism" was also obtained. (auth)

46383 COMPOUND ATOM STATE IN NEON. BOUND-STATE CALCULATION OF RESONANCE ENERGY. Foglia, C. (CNR, Parma, Italy). *Univ., Parma, Italy*. *Nuovo Cim.*; 10B: No. 1, 348-54(11 Jul 1972).

The first resonance energy in elastic electron scattering from neon is estimated from a single-configuration Hartree-Fock calculation of the resonance state. Radial wave functions and other related parameters for the Ne^- ion in the configuration $(2p)^5(3s)^22p$ are tabulated. (auth)

46384 PARTIAL IONIZATION CROSS SECTIONS OF CARBON DIOXIDE BY ELECTRONS (25 to 600 eV). Adamczyk, B. (Maria Curie-Skłodowska Univ., Lublin); Boerboom, A. J. H.; Lukasiewicz, M. *J. Mass Spectrom. Ion Phys.*; 9: No. 4, 407-12(Sep 1972).

Measurements were made of the partial ionization cross sections of CO_2^+ , CO^+ , O^+ , C^+ , and O_2^+ , originating from ionization of CO_2 by electrons of 25 to 600 eV. Total collection of these ions was achieved by using a cycloidal mass spectrometer. The relative measurements were normalized, using absolute values of the total cross sections from the literature. (auth)

46385 THIRD-DERIVATIVE METHOD FOR DETERMINING ELECTRON-IMPACT ONSET POTENTIALS. Finney, C. D.; Harrison, A. G. (Univ. of Toronto). *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 221-33(Aug 1972).

In an attempt to overcome the resolution problems of the Morrison second-differential method of determining ionization and appearance potentials from ionization efficiency data, the procedure of taking a third derivative was investigated both theoretically and experimentally. The results obtained indicate that the technique assists in revealing structure in the second-derivative curves and, in most cases studied, permits estimation of onset potentials in good agreement with photoionization and monoenergetic electron impact values. The limitations of the technique are discussed. (auth)

46386 MULTIPLE IONIZATION OF RARE GAS CLUSTERS BY ELECTRON IMPACT. Obert, W. (Univ., Karlsruhe, Ger. Kernforschungszentrum, Karlsruhe, Ger.). *J. Mass Spectrom. Ion Phys.*; 9: No. 3, 295-7(Aug 1972).

Condensed molecular beams (clusters beams) of the rare gases neon, argon, and xenon are ionized by electron impact. The mass-to-charge ratio of the cluster ions is determined in a retarding-field analyzer as a function of the electron energy. It is found that the probability of multiple ionization by one electron increases in the order Ne, Ar, Xe. This result is consistent with predictions based on the model for cluster ionization developed by Falter et al. (auth)

46387 INVERSION FROM THE ENERGY DEPENDENCE OF TOTAL CROSS SECTIONS. Guha, Evelyn; Ray, Swati; Barua, A. K. (Indian Association for the Cultivation of Science, Calcutta). *J. Phys. Soc. Jap.*; 33: No. 2, 468-72(Aug 1972).

The suitability of a recently suggested method for the inversion from the energy dependence of total-cross-section data at high energies was critically examined by considering He-He system. For this purpose, the data had to be generated partially by theoretical calculations. The results of the analysis show that the method has excellent potentialities, provided the required experimental data are available. Some suggestions for performing such experiments are given. (auth)

46388 ANGULAR DISTRIBUTION OF $H(2s)$ FORMED BY ELECTRON CAPTURE AND DISSOCIATION COLLISIONS. Fitzwilson, Roger Louis. Atlanta; Georgia Inst. of Tech. (1971). 189p. University Microfilms Order No. 72-19,702.

Thesis.

The differential cross section for the formation of hydrogen atoms in the metastable $2s$ state was measured for electron-capture and dissociation collisions. The reactions were studied for impact energies between 4 and 20 keV and scattering angles between 0.5 and 3 degrees. For the electron-capture reaction, the single projectile-target combination of H^+ and He was studied. For the dissociation work, collisions involving projectiles of H_2^+ and H_3^+ and targets of H_2 and He were studied. For comparison with previous theoretical and experimental work on electron capture, the differential cross sections for the formation of H^+ and H^0 were also measured. From the three cross sections, the probability P_{2s} for the formation of $H(2s)$ and the probability P_0 for the formation of H^0 were found. The fractional metastable content of the scattered neutral flux P_{2s}/P_0 was also evaluated and compared with existing theory. Similar probabilities were formed for the dissociation processes and compared with the electron-capture results. All differential cross sections for electron capture were found to be smoothly varying functions of scattering angle. The data fit curves of the form $K\theta^{-n}$ where n ranged from 2.5 to 3.2, depending on the energy and the particular cross section. The probabilities P_{2s} and P_0 were found to be fairly independent of scattering angle for angles greater than 0.7 degree but showed oscillatory structure as the impact energy was varied. The metastable fractional content of the scattered neutral flux, P_{2s}/P_0 , was found to rise from a small value at low angles to a high value at large angles; the magnitude of the rise depended on the impact energy. The dissociation data were also smoothly varying functions of angle. The differential cross sections for the dissociation of H_2^+ demonstrated a change in slope at scattering angles that could be correlated with known potential energy curves for the H_2^+ molecular ion if Franck-Condon excitations to antibonding states are assumed to be the mode of dissociation. The H_3^+ data showed no such breaks in the differential cross sections. (Diss. Abstr. Int., B)

46389 OPTIMIZED PSEUDOSTATES, INTERMEDIATE-ENERGY SCATTERING, AND COLLECTIVE MODES IN ATOMS. Mittleman, M. H. (City Coll., New York). *Phys. Rev.*, A; 6: No. 3, 879-84(Sep 1972).

In many of the usual methods of calculating scattering ampli-

tudes, a basis set is chosen in which to expand the wave function, and then the coefficients are optimized. The question of how to, in addition, optimize the basis set is considered here. A natural outgrowth of the formalism is the concept of an "average inelastic channel," which could, under certain circumstances, be interpreted as collective mode of excitation of the target. (auth)

46390 SLOW ELECTRONS FROM ELECTRON-IMPACT IONIZATION OF He, Ne, AND Ar. Grissom, J. T.; Compton, R. N.; Garrett, W. R. (Oak Ridge National Lab., Tenn.). *Phys. Rev., A*; 6: No. 3, 977-87(Sep 1972).

A trapped-electron technique is used to determine the integral cross sections for production of secondary electrons in the energy range from 0 to 1 eV resulting from electron-impact ionization of He, Ne, and Ar. For incident electron energies above 150 eV the slow-electron production cross section decreases approximately as the total-ionization cross section. A simple Born calculation for the case of atomic hydrogen also shows this general behavior. The measured absolute cross sections compare well with recently measured relative angular distributions and with Born and Born-exchange calculations. (auth)

46391 EXCITATION OF THE ELECTRONIC STATES OF THE NITROGEN MOLECULE BY ELECTRON IMPACT. Chung, S.; Lin, Chun C. (Univ. of Wisconsin, Madison). *Phys. Rev., A*; 6: No. 3, 988-1002(Sep 1972).

The electron-impact excitation cross sections of the $a^1\Pi_g$, $c^1\Sigma_u^+$, $a''^1\Sigma_u^+$, $\omega^1\Delta_u$, $b^1\Sigma_u^+$, $b^1\Pi_u$, $A^3\Sigma_u^+$, $B^3\Pi_g$, $C^3\Pi_u$, $D^3\Sigma_u^+$, $W^3\Delta_u$, and $E^3\Sigma_u^+$ states of N_2 were calculated over the range of 0 to 2000 eV for the singlet states and 0 to 40 eV for the triplet states by means of the Born approximation with Ochkur's and Rudge's scheme for treating the electron-exchange-scattering amplitude. The Franck-Condon-factor approximation was used to obtain the excitation cross sections to each vibrational level of the electronic states. The computation of the scattering-amplitude integrals was greatly facilitated by expressing the molecular wave functions in terms of atomic Gaussian-type orbitals. Four sets of self-consistent-field molecular wave functions were employed for the calculations in order to test the sensitivity of the calculated cross sections to the choice of the wave functions. With the exception of the $c^1\Sigma_u^+$ state, the cross sections based on three of the four sets vary by typically about 15%. As a test of the Born cross sections for singlet-singlet excitation, comparison between the theoretical and the available experimental values shows 25% agreement for the $a^1\Pi_g$ state at 900 eV and 50% for the $a''^1\Sigma_u^+$ state at 80 eV. In the case of triplet excitation, the theoretical cross sections of the $C^3\Pi_u$ state agree very well with the experimental data, but for the $A^3\Sigma_u^+$ and $B^3\Pi_g$ states the discrepancy is generally as large as a factor of 2. (auth)

46392 ANGULAR DISTRIBUTION OF ELECTRONS ELASTICALLY SCATTERED FROM N_2 . Shyn, T. W.; Stolarski, R. S.; Carignan, G. R. (Univ. of Michigan, Ann Arbor). *Phys. Rev., A*; 6: No. 3, 1002-12(Sep 1972).

The angular distribution of electrons elastically scattered from N_2 as a function of energy was measured utilizing a crossed-beam technique. Measurements were made of monoenergetic electrons with energies between 5 and 90 eV scattered from a collimated beam of N_2 at angles from -114° to $+160^\circ$. The wide angular range of the measurements enabled the determination of the total elastic-scattering and momentum-transfer cross sections. The measurements are relative and have been normalized to Fisk's theoretical calculation at 5 eV. The results generally agree well with other published measurements and with theory. (auth)

46393 INELASTIC COLLISIONS OF FAST CHARGED PARTICLES WITH ATOMS: IONIZATION OF THE ALUMINUM L SHELL. Manson, Steven Trent (Georgia State Univ., Atlanta). *Phys. Rev., A*; 6: No. 3, 1013-24(Sep 1972).

Born-approximation calculations of the ionization of aluminum L-shell electrons were performed using Hartree-Slater wave functions. Generalized oscillator strengths and proton energy-loss cross sections from threshold to ionized-electron energies of 128 Ry were calculated. The results show a delayed maximum above threshold, and the ramifications of this phenomenon are discussed. Comparison is made with hydrogenic results and good agreement is found at intermediate and large energy loss, but the hydrogenic calculation is totally inadequate near threshold. The implications of this for stopping-power calculations and sub-shell corrections is discussed. The Bethe asymptotic cross section is obtained, and the variation of parameters therein is investigated. (auth)

46394 DE-EXCITATION OF FAST He, Ne, AND Ar METASTABLE ATOMS IN VARIOUS GASES. Moseley,

John T.; Peterson, J. R.; Lorents, D. C.; Hollstein, M. (Stanford Research Inst., Menlo Park, Calif.). *Phys. Rev., A*; 6: No. 3, 1025-31(Sep 1972).

Absolute cross sections for the de-excitation of $He(2^1S)$ and of $He(2^3S)$ metastables by N_2 and Ar were measured over the energy interval from 150 to 1600 eV using an optical technique. The 2^1S cross sections are found to be as much as 50% larger than the 2^3S cross sections for de-excitation by N_2 , whereas for Ar the two cross sections are nearly equal at these energies. The cross sections all lie in the range of 7 to 20 $\text{\AA}^2$ and decrease with increasing energy. De-excitation cross sections were also measured by collecting slow ions for mixed beams of He metastables in N_2 , Ar, and C_2H_2 , and for mixed beams of Ar metastables in C_2H_2 and Ne metastables in Ar. These measurements are in the energy range from 50 to 600 eV. They do not discriminate between different metastable states, but the results exhibit an energy behavior similar to that of the state-separated measurements and are consistent with those measurements when comparison can be made. (auth)

46395 SEMIEMPIRICAL CALCULATIONS OF THE $He^*(2^3S \text{ AND } 2^1S) + Ar$ IONIZATION TOTAL CROSS SECTIONS. Olson, Ronald E. (Stanford Research Inst., Menlo Park, Calif.). *Phys. Rev., A*; 6: No. 3, 1031-6(Sep 1972).

The ionization (associative and Penning) total cross sections for $He^*(2^3S) + Ar$ and $He^*(2^1S) + Ar$ were calculated from 10^{-3} to 10^4 eV and compared with experimental data. The classical calculations used interaction potentials that agreed with ab initio theoretical calculations for the repulsive wall, possessed the correct long-range energy dependence, and were in agreement with glory-scattering experiments. The coupling width was an exponential with its free parameter determined by the thermal-energy ionization cross section for each system. The calculations were then extended over a wide range of energies. It was found that as the collision energy increased at the very low energies, the cross sections decreased until about 0.1 eV where they began slowly to rise. As the collision energy is further increased, the cross sections reach a maximum of about 15 $\text{\AA}^2$ around 10 eV, and then decrease again at higher energies. The cross sections predicted by the high-energy results, 100 to 1500 eV, are compared with the data of Moseley, Peterson, Lorents, and Hollstein (*Phys. Rev. A*, 6: 1025-31(1972)) and are found to agree within 20% and to have a similar energy dependence. (auth)

46396 OPTICAL EXCITATION AND POLARIZATION IN LOW-ENERGY COLLISIONS OF $He^+ + H_2$. Isler, R. C.; Nathan, R. D. (Univ. of Florida, Gainesville). *Phys. Rev., A*; 6: No. 3, 1036-48(Sep 1972).

Optical studies of spectral lines produced in collisions of $He^+ + H_2$ were made at energies below 700 eV. Lines of atomic hydrogen and helium are the only features observed. Emission cross sections were measured for Lyman- β , Balmer- α , Balmer- β , and for the following He I lines: 5876 $\text{\AA}$ ($3^3D \rightarrow 2^3P$), 6678 $\text{\AA}$ ($3^1D \rightarrow 2^1P$), 4471 $\text{\AA}$ ($4^3D \rightarrow 2^3P$). Emission cross sections are also listed at certain selected energies for all spectral lines observed. Polarization measurements were performed on the Balmer- α , the Balmer- β , and the visible helium lines. Only the Balmer- α line exhibits significant polarization, and the results are interpreted as indicating that the excited hydrogen atoms dissociate primarily in a direction perpendicular to the ion beam. (auth)

46397 ANGULAR DISTRIBUTION OF METASTABLE HYDROGEN FORMED BY ELECTRON CAPTURE IN A HELIUM TARGET. Fitzwilson, R. L.; Thomas, E. W. (Georgia Inst. of Tech., Atlanta). *Phys. Rev., A*; 6: No. 3, 1054-63(Sep 1972).

A study was made of the angular distribution of scattered particles induced by proton impact on a target of helium. Cross sections are presented for scattering of protons, of neutral hydrogen atoms, and of metastable hydrogen atoms. Impact energies range from 4 to 20 keV; scattering angles are from 0° to 2.0° . Measurements of the fractional metastable content of the scattered neutral flux rise from 1% or less at 0° to as much as 8% at angles of 1° . Theoretical predictions of the cross sections for scattering of all particles, neutrals and protons, show reasonable agreement with experiments. Theoretical predictions, by Colegrave and Stevens, of the probability for forming a neutral atom disagree with the experimental values by a constant phase factor. Measurements of the probability for forming a metastable atom do not agree with available theoretical predictions. (auth)

46398 CROSS SECTIONS FOR SINGLE-ELECTRON CAPTURE IN COLLISIONS OF PROTONS AND DEUTERONS WITH ATOMS. Tripathi, D. N.; Rai, D. K. (Banaras Hindu Univ., India). *Phys. Rev., A*; 6: No. 3, 1063-9(Sep 1972).

Electron-capture cross sections by protons (into 1s, 2s, and 3s

states) resulting from collisions with atomic systems were calculated using the simple classical formula of Gryzinski. Results obtained are compared with other theoretical calculations and experimental results. Comparison is made between the capture and ionization cross sections from collisions of heavy singly charged particles with atoms. (auth)

46399 DIFFERENTIAL- AND TOTAL-CROSS-SECTION CALCULATIONS FOR MODELS OF $e-H_2$ COLLISIONS. Sams, W. Neal (Univ. of Houston, Tex.); Frommhold, Lothar; Kouri, Donald J. *Phys. Rev., A*; 6: No. 3, 1070-6 (Sep 1972).

Differential and total cross sections for $e-H_2$ collisions were calculated by the close-coupling procedure using three different model interaction potentials. Particular attention is focused on the simplest possible model having the desired physical properties. This model uses square wells to describe short-range interactions and the known polarization and quadrupole interactions for long-range potentials. The results obtained for this square-well model for $e-H_2$ collisions are very encouraging considering its simplicity. Comparisons with results obtained using other model potentials are presented. (auth)

46400 METHOD OF EFFECTIVE POTENTIALS: SCATTERING OF POSITRONS AND ELECTRONS BY LIGHT ATOMS. Monaghan, William (New York Univ., N. Y. Staten Island Community Coll., New York); Rosenberg, Leonard *Phys. Rev., A*; 6: No. 3, 1076-86 (Sep 1972).

The theory of effective potentials is developed in a way that lends itself to variational formulations. The positron-helium and electron-helium systems are chosen to illustrate the method. The effective potential is defined in terms of the resolvent operator for a system modified by removal of the open-channel states. Rigorous maximum and minimum principles are derived, which should be useful in the computation of the effective potential and which are valid, for sufficiently low energies, even when the target bound-state wave function is not known exactly. A numerical application of this method to positron-hydrogen scattering has been made and results are reported. The effective-potential approach to the scattering problem leads naturally to a model for resonance calculations. This model is reviewed here and formulated variationally. An Appendix is devoted to further elaboration of this approach to resonance calculations in the context of potential scattering, with the aid of Jost-function theory. (auth)

46401 FERMI PRESSURE SHIFT AND ELECTRON-RARE-GAS-ATOM SCATTERING LENGTHS. Madan, Rabinder N. (North Carolina Agricultural and Technical State Univ., Greenboro). *Phys. Rev., A*; 6: No. 3, 1242-4 (Sep 1972).

The treatment of the Fermi shift for alkali metals in the paper by M. H. Choudhury (*Phys. Rev.* 186, 66 (1969)) is examined. Some difficulties are pointed out, which render Choudhury's results to be of doubtful value. It is seen that starting from Choudhury's Hamiltonian one may recover an equation considered earlier by Mittleman for studying the Fermi-shift phenomenon. Finally, the Mittleman result is employed to make new estimates for the electron-neutral-atom scattering lengths. (auth)

46402 INTERPOLATION OF VARIATIONALLY OBTAINED SCATTERING CROSS SECTIONS. Heller, Eric J. (Harvard Univ., Cambridge, Mass.). *Phys. Rev., A*; 6: No. 3, 1247-8 (Sep 1972).

Standard variational methods are shown to contain the information necessary to construct the Jost function, a quantity previously shown to yield readily to interpolation. This allows narrow resonances to be found and characterized without extensive explicit computations. (auth)

46403 EXCITATION OF HYDROGEN IN COLLISIONS WITH ALKALI-METAL ATOMS. Mathur, K. C.; Tripathi, A. N.; Joshi, S. K. (Univ. of Roorkee, India). *Phys. Rev., A*; 6: No. 3, 1248-52 (Sep 1972).

Excitation of hydrogen atoms from the ground state to 2s, 2p, 3s, 3p, and 3d states in passing through alkali metal atoms was studied using the Born approximation. The form-factor description of the target and the generalized-oscillator-strength representation of the projectile were used. (auth)

46404 CHEMI-IONIZATION IN COLLISIONS BETWEEN HELIUM METASTABLE ATOMS AND ARGON. Howard, J. S.; Riola, J. P.; Rundel, R. D.; Stebbings, R. F. (Rice Univ., Houston, Tex.). *Phys. Rev. Lett.*; 29: No. 6, 321-4 (7 Aug 1972).

Cross sections for chemi-ionization in thermal energy collisions of $He(2^1S)$ and $He(2^3S)$ metastable atoms with argon were mea-

sured using both crossed-beam and gas-cell techniques. The results indicate that several earlier measurements of these cross sections were in error. (auth)

46405 OBSERVATION OF A BROAD RESONANCE IN THE 2^3S EXCITATION OF HELIUM BY ELECTRON IMPACT. Crooks, G. B.; DuBois, R. D.; Golden, D. E.; Rudd, M. E. (Univ. of Nebraska, Lincoln). *Phys. Rev. Lett.*; 29: No. 6, 327-9 (7 Aug 1972).

A broad P-wave resonance was observed at about 50 eV primary energy in the angular distribution of electrons scattered after excitation of the 2^3S state of helium. The resonance has a width of about 15 eV and affects the differential excitation cross section by more than 3 orders of magnitude. It is suggested that this resonance is associated with the temporary formation of triply excited states of the negative helium ion. (auth)

46406 SCATTERING OF CsF BY ELECTRONS. Slater, Richard C.; Fickes, Michael G.; Stern, Richard C. (Columbia Univ., New York). *Phys. Rev. Lett.*; 29: No. 6, 333-6 (7 Aug 1972).

Absolute differential and total cross sections for the scattering of CsF by 0.6- to 6.5-eV electrons were measured using the molecular-beam recoil technique. The first-Born-approximation differential cross section is used as the theoretical model to fit the data. The experimental differential cross sections rise more sharply near 0° than predicted. Total cross sections derived from these measurements are a factor of 1.5 to 4 smaller than the first-approximation results. (auth)

46407 MANIFESTATION OF STRUCTURE IN ELECTRON ENERGY DISTRIBUTION FUNCTIONS AND DETERMINATION OF THE EFFECTIVE CROSS SECTIONS OF INELASTIC PROCESSES. Soldatov, A. N.; Prilezhaeva, N. A. (Tomsk State Univ., USSR). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 51-62 (1971). (In Russian).

Features of the energy distribution function for electrons in a discharge with two tubular coaxially oriented electrodes were experimentally investigated. Effective cross sections of inelastic collisions were determined using the structure in the distribution functions of electrons, this structure having appeared as a result of inelastic processes. The time of energy recoil by electrons in "elastic" and "inelastic" energy regions was examined. The effective cross section of de-excitation of the metastable $He_2^3S_1$ was determined during collisions of the second kind with slow electrons, and the cross section of two metastable $He_2^3S_1$, the result of which was the production of an ionized molecule of helium He_2^+ and an electron with an energy of 15 eV. Reasons for the manifestation of structure in the electron energy distribution functions are discussed. (tr-auth)

Atomic Theory

46408 STATISTICAL CONSIDERATION OF RETARDATION IN RELATIVISTIC HARTREE-FOCK EQUATIONS. Guruchumeliya, A. D. (Georgian Polytechnic Inst., Tbilisi, USSR). *Soobshch. Akad. Nauk Gruz. SSR*; 67: No. 1, 53-6 (Jul 1972). (In Russian).

A statistical calculation of exchange energy is presented, in which retardation appears when the atom is described in relativistic terms by the Hartree-Fock method. The contribution of retardation to the Hartree-Fock potential was found to equal $(\pi/2)\alpha^2\rho$, which leads to correlations of the order of α^2Z^2 for energy-term values of atoms. The ultrarelativistic limit is also considered. (auth)

46409 CORRELATION EFFECTS IN ATOMIC SYSTEMS. Larsson, Sven. Abstracts of Uppsala Dissertations from the Faculty of Science, No. 215. Uppsala; University of Uppsala (1972). 14p.

Thesis.

Eleven previously published papers on which a dissertation is based are summarized in three groups. The first group deals with calculations of wave functions for three-electron systems, the second with examination of correlation corrections to hyperfine structure expectation values, and the third with calculations of correlation corrections to the orbitals and a discussion of exchange and correlation potentials. (A.G.W.)

46410 INFINITE ORDER PERTURBATION THEORY WITH APPLICATIONS TO ATOMIC AND MOLECULAR SYSTEMS. Braendas, Erkki. Abstracts of Uppsala Dissertations from the Faculty of Science, No. 201. Uppsala; University of Uppsala (1972). 13p.

Thesis.

A dissertation based on seventeen previously published papers

is summarized. In the thesis, it is shown that substantial improvements in theory and computations of atomic and molecular properties and structure are obtained by incorporating the most attractive features of the variational and perturbation method into a unified theory. Inner projections and Pade approximants are used to demonstrate the merits of such a combination. (A.G.W.)

46411 THEORY OF COOPERATIVE EMISSION FROM TWO-LEVEL ATOMS. Rehler, Nicholas E. Rochester, N. Y.; Univ. of Rochester (1972). 153p. University Microfilms Order No. 72-18,832.

Thesis.

A new approach to the problem of cooperative emission from two-level atoms excited by a short plane-wave pulse is introduced. The treatment applies to an arbitrary number of atoms confined to a region with linear dimensions that may be large compared with λ , the wavelength associated with the atomic transition frequency. Explicit expressions are found, which describe the time evolution and directional character of the radiated power. The normally ordered relative intensity fluctuation is calculated for both the field radiated into any direction by a system with linear dimensions $\ll \lambda$ and the field radiated into the forward direction by a larger system. Large fluctuations when the system is nearly inverted indicate the discrete nature of spontaneous emission; the very small fluctuations found when only half of the atoms are excited show that the emission is then nearly classical. The far-field radiation patterns appropriate to the cases of a random distribution of atoms within a sphere and a right circular cylinder are considered in detail. Polar plots are presented of the radiation patterns produced when the atoms in a cylinder are excited by plane-wave pulses traveling both parallel and at an angle of 10° with the cylinder axis. All results are compared with related work by other authors. (Diss. Abstr. Int., B)

46412 PAIR CORRELATION ENERGIES AS DERIVED FROM THE ANALYSIS OF SEMI-EMPIRICAL VALUES OF CORRELATION ENERGIES OF ATOMS. Anno, Tosinobu; Sakai, Yoshiko (Kyushu Univ., Fukuoka). J. Chem. Phys.; 57: No. 4, 1636-47 (15 Aug 1972).

For the purpose of obtaining the semiempirical formulas for the pair correlation energy of various types of atoms and ions having the electron configurations $1s^2 2s^m 2p^n$ ($m = 0, 1, 2$; $n = 0, 1, \dots, 6$), semiempirical values of the correlation energies of such atoms are analyzed. It is pointed out that the correlation energy in the average of a configuration is more convenient than the correlation energy in each of the stationary states belonging to the configuration, for the purpose of this analysis. The correlation energy in the average of the configuration is shown to be equal to E_{AV}^{corr} , the average of the correlation energy in various states belonging to the configuration. The E_{AV}^{corr} values are obtained for a wide variety of the $1s^2 2s^m 2p^n$ atoms and regularities are found among these E_{AV}^{corr} values. From the analysis of these regularities, it is found that the E_{AV}^{corr} values of the $1s^2 2s^m 2p^n$ atoms can be reproduced very well with an equation $E_{AV}^{corr} = C_1 + C_2m + C_3n + C_4m^2 + C_5mn + C_6n^2 + C_7m^2n + C_8mn^2 + C_9m^2n^2 + C_{10}m(m-1)Z + C_{11}m(m-1)nZ + C_{12}m(m-1)Z^2$, where Z is the atomic number, along with the numerical values of the coefficients C_i 's ($i = 1, 2, \dots, 12$) determined from a least-squares fitting of this equation to various empirical E_{AV}^{corr} values. The concept of the average pair correlation energy of the configuration ϵ_{kl} is also proposed. Expressions for ϵ_{kl} 's are given in terms of the pair correlation energies of various type in each of the states belonging to the configuration. It is shown that E_{AV}^{corr} can be expressed in a simple form in terms of ϵ_{kl} 's. From this expression and the semiempirical expression of E_{AV}^{corr} mentioned above, a semiempirical formula for the average pair correlation energy is obtained for each type of electron pairs. It is found that these semiempirical formulas are in accord with the results of the Sinanoğlu theory for the ground and the excited states and of nonempirical calculations of pair correlation energies, rather than with those of a previous purely empirical analysis of the correlation energies by Allen, Clementi, and Gladney. (auth)

46413 UPPER AND LOWER BOUNDS TO EXCITED STATES OF TWO-ELECTRON ATOMS. Wang, P. S. C.; Weinhold, F. (Stanford Univ., Calif.). J. Chem. Phys.; 57: No. 4, 1738-45 (15 Aug 1972).

The excited 1^1S and 1^3P states of the helium isoelectronic sequence are treated by the Ritz and Bazley variational methods, which together provide rigorous upper and lower bounds to the true nonrelativistic energy levels. With compact variational functions constructed from about 25 hydrogenic basis terms, up to 20 or more states of each symmetry can be treated for each ion. The error limits finally approach a few parts in 10^5 or better for many of the more highly excited states (including many not yet observed), an accuracy sufficient to determine the ionization potentials to about 1%. (auth)

46414 NATURE OF TWO-PARTICLE CORRELATIONS IN ATOMS. Geller, Elaine N.; Geller, Murray (California Inst. of Tech., Pasadena). J. Chem. Phys.; 57: No. 4, 1814 (15 Aug 1972).

A method for analytical evaluation of radial integrals used by Ruffa (J. Chem. Phys., 52: 1512 (1970)) in discussion of the nature of two-particle correlations in atoms is suggested briefly. (A.G.W.)

46415 VARIATIONAL BOUNDS FOR IMAGINARY FREQUENCY POLARIZABILITY AND DISPERSION INTERACTION CONSTANT. Braun, P. A.; Rebane, T. K. (Leningrad State Univ.). Int. J. Quant. Chem.; 6: No. 4, 639-46 (Jul 1972).

A study is made of a variational principle that provides upper and lower bounds for the polarizability under a perturbation with imaginary frequency. As an example bounds for the imaginary frequency polarizability of atomic hydrogen and ensuing limits for the constant of van der Waals interaction of two hydrogen atoms are calculated. (auth)

46416 DECOUPLED HARTREE-FOCK METHODS. I. CALCULATIONS FOR ATOMS WITH ORTHOGONALIZED ORBITALS. Naray-Szabo, G. (Technical Sciences Univ., Budapest). Int. J. Quant. Chem.; 6: No. 4, 779-85 (Jul 1972).

By a proper approximation of the interaction term in a many-electron Hamiltonian the Hartree-Fock (HF) equations are decoupled. Making use of this simplification one obtains a good initial guess for the wave function with minimal computational work. Refining the procedure, the exact HF limit can be achieved. Total energies, ionization potentials and excitation energies for light atoms are calculated. (auth)

46417 VARIATIONAL PRINCIPLES FOR SINGLE-PARTICLE EXPECTATION VALUES IN THE HARTREE APPROXIMATION. Krieger, J. B. (Brooklyn Coll.). Phys. Rev., A; 6: No. 3, 919-27 (Sep 1972).

The application of Delves's variational principle for the calculation of the expectation value of single-particle operators W is investigated in the Hartree approximation. In this approximation the auxiliary function is taken to be of the form $\psi_{1T} = \Sigma f_i(\vec{r}_i) \psi_{0T}$, where ψ_{0T} is the trial ground-state wave function. For a given ψ_{0T} , a set of coupled integrodifferential equations satisfied by the $f_i(\vec{r}_i)$ is derived by minimizing an auxiliary functional containing ψ_{0T} , ψ_{1T} , W , and the Hamiltonian of the system. The resulting equations are uncoupled in two different approximations each valid for one- and two-particle systems and are compared to those employed by others and to those previously suggested by the authors on the basis of certain self-consistency considerations. The uncoupled equations are solved exactly for single-particle operators that depend on the radial distance only. The utility of the technique is demonstrated by showing that for a model hydrogen-atom problem it leads to highly accurate results for the electron density at the origin and the Fourier transform of the electron density even when the calculation of these quantities employing ψ_{0T} alone is substantially in error. The results of applying the technique to the helium atom for the calculation of $\langle \gamma^n \rangle$, $n = -2, -1, 1, 2$, and the electron density at the origin employing an energy-minimized product of hydrogenic wave functions for ψ_{0T} are reviewed and compared with those of Pekeris and found to have an accuracy equivalent to a numerical Hartree-Fock calculation. Finally, the decoupling approximations are extended to systems containing more than two particles. (auth)

46418 APPLICATION OF GREEN'S FUNCTIONS TO RADIATIVE COOPERATIVE EFFECTS IN MULTI-ATOM SYSTEMS. Duncan, G. Comer (Bowling Green State Univ., Ohio). Phys. Rev., A; 6: No. 3, 947-58 (Sep 1972).

Spontaneous-radiation processes associated with a number of multi-atom systems are studied. Green's-function techniques are used in conjunction with a model of N two-level atoms interacting with a quantized radiation field to investigate the assumption of the independence of the spontaneous-radiation properties of a given atom from the states of the other atoms in the system. It is shown that the natural linewidth of an excited atom in the presence of a de-excited atom is different from that of an isolated excited atom. For interatomic separations smaller than a critical separation, the two-atom system is best considered as a collective unit with regard to its spontaneous-radiation properties. The influence of these "radiative cooperative effects" is studied also for the systems of two initially excited atoms. Also the influence of radiative cooperative effects in the scattering of a photon by a system of two de-excited atoms is studied. It is shown that the frequency distribution of the scattered radiation exhibits a double-peak structure for interatomic separations smaller than $\frac{1}{2}\lambda$. The techniques of Green's functions are then applied to the study of radiative cooperative effects in several many-atom systems. It

is shown that within the context of the model certain information is obtained exactly via the Green's-function techniques as applied to the many-atom problems of a single excited atom in the presence of $N \gg 1$ de-excited atoms and the scattering of a photon by a system of $N \gg 1$ de-excited atoms. The necessity of treating many-atom systems from a "collective" point of view is easily seen from our results. (auth)

46419 COUPLED MULTICONFIGURATIONAL SELF-CONSISTENT-FIELD METHOD FOR ATOMIC DIPOLE POLARIZABILITIES. I. THEORY AND APPLICATION TO CARBON. Billingsley II, Frank P.; Krauss, Morris (National Bureau of Standards, Washington, D. C.). Phys. Rev., A; 6: No. 3, 855-65(Sep 1972).

A method for calculating static dipole polarizabilities of atoms

within a multiconfigurational self-consistent-field (SCF) framework is presented. The method involves the direct solution of the multiconfigurational SCF equations of an atom in the presence of a perturbing field simulated by a charged particle. The use of a multiconfigurational framework allows this technique to be applied straightforwardly to any given state of both degenerate and nondegenerate atoms, and also allows the explicit introduction of correlation effects. Sample calculations are reported for the static dipole polarizabilities of the carbon atom in its ^1S , ^1D , and ^3P states with partial inclusion of correlation. The results are compared with those obtained from many-body perturbation theory and other techniques. In addition, a prescription for specifying a sufficiently flexible set of polarization basis functions is described. (auth)

PHYSICS (ELECTROFLUID AND MAGNETOFLUID)

46420 WEAK FLUCTUATION MODES OF AN ELECTRICALLY AND THERMALLY CONDUCTING VISCOUS COMPRESSIBLE FLOW IN A HOMOGENEOUS MAGNETIC FIELD. Stanisic, Milomir M. (Purdue Univ., Lafayette, Ind.). *Z. Angew. Math. Phys.*; 22: No. 6, 1088-94(25 Nov 1971).

The propagation of small disturbances in a three-dimensional unsteady, viscous, compressible fluid with finite conductivity is considered in detail, so that earlier results appear as special limiting cases. It is noted that although the flow is not isentropic in the strict sense, the entropy change appears only as second order effect both in the pressure and in the magnetic field. It is concluded that, in the case of compressible, electrically and thermally conducting viscous flow with finite conductivity there is no corresponding group and phase velocity associated with given wave number resulting from finite conductivity. (M.H.R.)

46421 VORTICITY JUMP ACROSS A STATIONARY MAGNETOGASDYNAMIC DISCONTINUITY. Sharma, V. D.; Ram, Rishi (Banaras Hindu Univ., India). *Z. Angew. Math. Phys.*; 22: No. 6, 1126-34(25 Nov 1971).

The Hayes [J. Fluid Mech. 2, 595(1957)] analysis is generalized to an MHD discontinuity and it is shown that the vorticity jump across a magnetogasdynamic discontinuity is not derivable from the momentum equation alone but would also essentially require the use of the energy equation except in some particular cases of MHD flows. (M.H.R.)

46422 SECONDARY FLOW OF A CONDUCTING FLUID IN A ROTATING STRAIGHT ANNULAR PIPE. Rao, V. V. Ramana (Andhra Univ., Waltair, India). *Proc. Indian Acad. Sci., Sect. A*; 75: No. 4, 189-98(Apr 1972).

A theoretical analysis is made of the flow of a conducting viscous and incompressible fluid through a straight annular pipe of circular cross section flowing under a constant pressure gradient. The pipe is rotated about an axis perpendicular to it and also there is imposed a uniform magnetic field transverse to the motion. The secondary flow pattern was studied in detail for small values of angular velocity and Hartmann number. (auth)

46423 MAGNETOHYDRODYNAMIC ENTRANCE FLOW IN A CHANNEL. Flores, F.; Recuero, A. (Univ. of Madrid). *Appl. Sci. Res. (The Hague)*; 25: No. 6, 403-12(Mar 1972).

An MHD entrance flow in a channel is studied in the Prandtl approximation. It is shown that consistency with the approximation requires an appropriate change in the boundary conditions that hold for the original, unsimplified equations. The correct boundary conditions are established and used to repeat the numerical calculations in a few cases studied by other authors with the unmodified boundary conditions. In the cases considered here, the numerical differences do not amount in practice to more than 10%, although they grow with increasing Hartmann number. (auth)

46424 MOTION OF A SPHERE IN AN ELECTRICALLY CONDUCTING ROTATING FLUID. Sarma, L. V. K. V.; Seetharamaswamy, R. (Indian Inst. of Tech., Madras). *Appl. Sci. Res. (The Hague)*; 25: No. 6, 431-44(Mar 1972).

The combined effect of rotation and magnetic field is investigated for the axisymmetric flow due to the motion of a sphere in an inviscid, incompressible electrically conducting fluid having uniform rotation far upstream. The steady-state linearized equations contain a single parameter $\alpha = \frac{1}{2}\beta R_m$, β being the magnetic pressure number and R_m the magnetic Reynolds number. The complete solution for the flow field and magnetic field is obtained, and the distribution of vorticity and current density is found. The induced vorticity is $O(\alpha^4)$ and the current density is $O(R_m)$ on the sphere. (auth)

46425 TWO-DIMENSIONAL NONSTEADY-STATE SOLUTION OF MAGNETOHYDRODYNAMIC EQUATIONS. Imshennik, V. C. *Zh. Prikl. Mekh. Tekh. Fiz.*; No. 2, 3-7(1972). (in Russian).

A solution to MHD equations by A. G. Kulikovskii (1958) supports the case of a plane, cumulation of matter in the vicinity of a null line of a magnetic field, which may be related to certain plasma processes (z-pinch, solar flares). A certain class of simple solutions to MHD equations for a two-dimensional plane, nonsteady-state motion was examined. Two subclasses of solutions with completely different physical significance were found, and of fundamental interest is that subclass which comprises the solution for a cumulative plane compression of matter in the environment of the null line of a magnetic field. (K.S.W.)

PHYSICS (HIGH-ENERGY)

Particle Interactions and Properties (Experimental)

Refer also to abstract: 46126.

Electromagnetic

Refer also to abstract: 46733.

46426 (BNL-16968) PHOTOPRODUCTION EXPERIMENTS AT ISABELLE. Drickey, D. J. (Brookhaven National Lab., Upton, N. Y.). 16 Jun 1972. 7p. Dep. NTIS.

Trivial extensions are made to the ISABELLE e-p parameters of 5-GeV electrons on 200-GeV protons. The electroproduction total cross section as a function of equivalent laboratory photon energy is shown. It is concluded that photoproduction experiments in a new energy range are possible with the ISABELLE e-p option even for Compton scattering processes provided care is taken in the design of the interaction region. (J.F.P.)

46427 (CONF-710816-13) $\pi^+\pi^-\pi^0$ PRODUCTION BY e^+e^- ANNIHILATION AT THE ω MASS. Benaksas, D.; Cosme, G.; Jean-Marie, B.; Jullian, S.; Laplanche, F.; Lefrancois, J.; Liberman, A. D.; Parrour, G.; Repellin, J. P.; Sauvage, G. (Paris Univ., Orsay (France). Laboratoire de l'Accelérateur Lineaire). [nd]. 12p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

Cross sections for $e^+ + e^- \rightarrow \pi^+ + \pi^- + \pi^0$ annihilation in the Orsay storage ring, the ω width, and the width and branching ratio for $\omega \rightarrow e^+ + e^-$ decay are found using a large solid angle detector. The experimental set-up is described. (J.F.P.)

46428 (CONF-710816-14) $\pi^+\pi^-$ PRODUCTION BY e^+e^- ANNIHILATION IN THE ρ ENERGY RANGE WITH THE ORSAY STORAGE RING. Benaksas, D.; Cosme, G.; Jean-Marie, B.; Jullian, S.; Laplanche, F.; Lefrancois, J.; Liberman, A. D.; Parrour, G.; Repellin, J. P.; Sauvage, G. (Paris Univ., Orsay (France). Laboratoire de l'Accelérateur Lineaire). 1971. 16p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

A large solid angle detector was used to observe two body events produced by electron-positron collisions in the Orsay storage ring. From the $\pi^+\pi^-$ excitation curve in the ρ region the amplitude and the phase of the ω - ρ interference, and the ρ resonance parameters were deduced. (auth)

46429 (CONF-710816-15) HADRONIC PRODUCTION BY e^+e^- COLLISIONS AT 990 MeV WITH THE ORSAY STORAGE RING. Benaksas, D.; Cosme, G.; Jean-Marie, B.; (and others) (Paris Univ., Orsay (France). Laboratoire de l'Accelérateur Lineaire). 1971. 8p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

A large solid angle detector is used to measure the cross sections for $e^+ + e^- \rightarrow \pi^+ + \pi^- + \pi^0(2\pi^0)$ interactions at 990 MeV in colliding beams. Also an upper limit is found for $e^+ + e^- \rightarrow 2\pi^+ + 2\pi^-$ interactions. (J.F.P.)

46430 (COO-1195-241) MISSING-MASS SPECTRUM NEAR 960 MeV FROM $\pi\pi \rightarrow n + (\text{mm})$. Holloway, L.; Huld, B.; Koetz, U.; (and others) (Illinois Univ., Urbana). Jun 1972. Contract AT(11-1)-1195. 7p. (CONF-720409-6). Dep. NTIS.

From third international conference on experimental meson spectroscopy; Philadelphia, PA. (28 Apr 1972).

Supporting evidence for structure near 960 MeV in the reaction $\pi\pi \rightarrow n + (\text{missing-mass})$ at 2.5 GeV/c is presented. (auth)

46431 (NP-19256) SEPARATION OF σ_S AND σ_T IN THE REGION OF THE $\Delta(1236)$ RESONANCE AND DETERMINATION OF THE MAGNETIC DIPOLE TRANSITION FORM FACTOR. Baetzner, K.; Beck, U.; Becks, K. H.; (and others) (Bonn Univ. (West Germany). Physikalisches Institut). Feb 1972. 13p. (PIB-1-157). Dep. NTIS (U. S. Sales Only).

The longitudinal and transverse cross sections σ_S and σ_T are evaluated from measurements of inelastic ep scattering at scattering angles ranging from 20 to 81°. The magnetic dipole transition form factor G_M^* is determined and the reliability of the determination is discussed. (auth)

46432 (SLAC-Trans-142) PHOTOPRODUCTION OF NEUTRAL PIONS ON HELIUM NUCLEI. Tsarev, V. A.; Daion, M. I.; Rakov, Yu. A. (Akademiya Nauk SSSR, Moscow. Institut Fiziki). Translated for Stanford Linear Accelerator Center, Calif., from Preprint No. 170. 27p. Dep. NTIS.

The physics and proposed measurements of the cross sections for $\gamma + {}^4\text{He} \rightarrow {}^4\text{He} + \pi^0$ interactions at 2 to 5 GeV and pion emission angles above 10° using a streamer chamber are discussed. The π^0 lifetime can also be determined from this reaction. (J.F.P.)

46433 SCALE RELATION FOR PROTON FORM FACTORS. Bilen'kaya, S. I.; Bilen'kii, S. M.; Kazarinov, Yu. M. (Joint Inst. for Nuclear Research, Dubna, USSR). JETP Lett. (USSR) (Engl. Transl.); 15: No. 7, 297-9(5 Apr 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 7, 420-3(5 Apr 1972).

The validity of the scale relation $G_M(q^2) = \mu G_E(q^2)$, where $G_E(q^2)$ and $G_M(q^2)$ are the charge and magnetic form factors of the proton and μ is the total magnetic moment of the proton in nuclear magnetons, was investigated through an analysis of all available data on e-p scattering cross sections. The electromagnetic form factors were extracted directly from the data by assuming a certain functional dependence of the form factors on q^2 and obtaining the values of the corresponding parameters by minimizing the functional χ^2 . The experimental data of different groups were reduced by introducing normalization factors to take account of the possible errors in the normalization of the data. The analysis confirmed the validity of the scale relation within the appropriate errors. (R.W.R.)

46434 EXPERIMENTAL STUDY OF THE DECAYS $K_{S,L} \rightarrow \gamma\gamma$. Banner, M.; Coronin, J. W.; Hoffman, C. M.; Knapp, B. C.; Shochet, M. J. (Princeton Univ., N. J.). Phys. Rev. Lett.; 29: No. 4, 237-40(25 Jul 1972).

The branching ratio ($K_S \rightarrow \gamma\gamma$)/($K_S \rightarrow \text{all}$) was measured to be $(-1.9 \pm 2.4) \times 10^{-4}$ in an optical spark-chamber experiment. The result assumes no interference between $K_S \rightarrow \gamma\gamma$ and $K_L \rightarrow \gamma\gamma$. In addition, the branching ratio ($K_L \rightarrow \gamma\gamma$)/($K_L \rightarrow \text{all}$) was measured to be $(4.32 \pm 0.55) \times 10^{-4} \times |\eta_{00}/\eta_{+-}|^2$. (auth)

46435 FURTHER EVIDENCE FOR THE DECAY MODE $\phi \rightarrow \eta\gamma$ AND MEASUREMENT OF THE BRANCHING RATIO $\Gamma(\phi \rightarrow \eta\gamma)/\Gamma(\phi \rightarrow \text{total})$. Basile, M.; Dalpiaz, P.; Frabetti, P. L.; (and others) (CERN, Geneva. Univ., Bologna. Istituto Nazionale di Fisica Nucleare, Bologna. CNRS, Strasbourg). Nucl. Phys.; B44: No. 2, 605-17(1972).

The reaction $\pi^-p \rightarrow \eta\phi$ was studied at 1.8 GeV/c incident pion momentum using the Bologna-CERN NBC set-up, in order to investigate the electromagnetic decay mode $\phi \rightarrow \eta\gamma$. (27 $\pm$ 6) events were observed, yielding a branching ratio $\Gamma(\phi \rightarrow \eta\gamma)/\Gamma(\phi \rightarrow \text{total}) = (7.3 \pm 1.9)\%$. The theoretical implications of this result are discussed. (auth)

46436 POLARIZED PROTON AND DEUTERON TARGETS AND HIGH-ENERGY PHYSICS EXPERIMENTS. Borghini, M. (CERN, Geneva). pp 145-80 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Progress in polarized targets to be used for high energy physics experiments (above a few GeV) is reviewed. This progress consists of improvements in the nature of the substances containing the polarized protons (alcohol targets), the polarization of deuterons, a better understanding of the effects producing the polarization, and the use of more elaborate equipment as far as target geometry is concerned. Recent results obtained using these targets in lepton scattering, photoproduction and hadron scattering are briefly surveyed. (France)

46437 DIFFRACTIVE FRAGMENTATION IN HIGH ENERGY PHOTON REACTIONS. Fries, D. (Univ., Karlsruhe, Ger.). pp 267-88 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

A multipion photoproduction experiment was performed using an 18-GeV bremsstrahlung photon beam and a gaseous hydrogen target. Preliminary results from the analysis of film data are

presented. Experimental evidence was obtained between 4 and 18 GeV, for a ρ^0 signal having an energy dependence similar to that involved in diffractive elastic ρ photoproduction. A diffractive fragmentation of the isovector photon is shown to occur simultaneously with the fragmentation of the target. (France)

46438 TOPICS IN PHOTOPRODUCTION. Braccini, P. L. (Univ., Pisa, Italy). pp 527-49 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

The photoproduction reaction $\gamma + p \rightarrow \rho^0 + p$ was studied at 2.8 and 4.7 GeV. A t dependence of the mass distribution was observed and a dependence of the t distribution on the dipion mass was also found, the results being in agreement with a previous dual model. In the helicity system, for $-t < 0.4$ (GeV/c)², the density matrix has the same structure as the photon density matrix. Evidence for the validity of the Soeding model was obtained from the cross sections for ρ production. (France)

46439 REVIEW AND ANALYSIS OF THE LATEST DATA ON MEASURING FUNDAMENTAL PHYSICAL CONSTANTS (MAGNETIC MOMENTS OF AN ELECTRON, MUON, AND PROTON). Gorbatshev, S. V.; Kholin, V. M.; Nosal, V. N. Meas. Tech. (USSR) (Engl. Transl.); 15: No. 1, 20-31 (Jan 1972).

Translated from Izmer. Tekh.; 15: No. 1, 14-20 (Jan 1972).

This review and analysis of data covers the relation of the electron magnetic moment to the Bohr magneton, the relation of the magnetic moment of an electron and a proton, the ratio of the proton magnetic moment to the Bohr magneton, evaluation of the ratio of a muon magnetic moment to a muon magneton, and the ratio of the muon and proton magnetic moments. Also included are the ratio of the g -factors of a muon and electron, the ratio of the masses of a proton and an electron, and the ratio of the masses of a muon and an electron. (J.F.P.)

46440 MEASUREMENT OF THE MAGNETIC MOMENT OF THE ANTIPROTON. Fox, J. D. (Brookhaven National Lab., Upton, N. Y.); Barnes, P. D.; Eisenstein, R. A.; (and others). Phys. Rev. Lett.; 29: No. 3, 193-5 (17 Jul 1972).

Fine-structure splittings in atoms formed by antiprotons with both lead and uranium were measured. Since the fine structure is produced by an interaction between the magnetic moment of the antiproton and the Coulomb field of the nucleus, a value for the magnetic moment can be derived from the observed splitting. A magnetic moment of -2.83 ± 0.10 nuclear magnetons is found which agrees with the TCP prediction. (auth)

46441 PHOTOPRODUCTION OF CHARGED PIONS BY NUCLEI AT $E_0 = 1160$ MeV. Grishaev, I. A.; Lapin, N. I.; Nikiforov, V. I.; Pugachev, G. D.; Shramenko, B. I. (Physical-Technical Inst., Kharkov). Ukr. Fiz. Zh. (Russ. Ed.); 16: No. 9, 1550-2 (Sep 1971). (In Russian).

The yield of charged π -mesons formed by the nuclei C, Al, Cu, Ta, and Pb was measured during maximum energy of retarding gamma-quanta at $E_0 = 1160$ MeV at an angle of 150° . The kinetic energy of the pions was 40 and 65 MeV. Detection of pions was accomplished using photoemulsion. The total yield of charged mesons was proportional to $A^{0.71}$. (tr-auth)

46442 ASYMMETRY MEASUREMENTS FOR π^0 AND π^+ PHOTOPRODUCTION BY LINEARLY POLARIZED γ -RAYS IN THE ENERGY RANGE (400 TO 900) MeV. Zdarko, R. W. (Stanford Univ., Calif.); Dally, E. B. Nuovo Cim.; 10A: No. 1, 10-18 (1 Jul 1972).

The asymmetry ratio $A = (\sigma_{\perp} - \sigma_{\parallel}) / (\sigma_{\perp} + \sigma_{\parallel})$ was measured by means of linearly polarized γ -rays for π^0 and π^+ photoproduction. For the reaction $\gamma + p \rightarrow n + \pi^+$, measurements were taken for $\theta_{c.m.} = 135^\circ$ at γ -ray energies ranging from 390 to 909 MeV. For the reaction $\gamma + p \rightarrow p + \pi^0$, measurements were taken for $\theta_{c.m.} = 60^\circ$ and 90° at γ -ray energies ranging from 426 to 918 MeV. (auth)

46443 ϕ -MESON MASS DETERMINATION. Balakin, V. E.; Bukin, A. D.; Pakhtusova, E. V.; Sidorov, V. A.; Khabakhpashev, A. G. (Inst. of Nuclear Physics, Novosibirsk, USSR). Phys. Lett.; 40B: No. 3, 431-2 (10 Jul 1972).

Calibration of the electron-positron storage ring energy using the angular distribution of pions produced in the reaction $e^+e^- \rightarrow \phi \rightarrow K_L^0 K_S^0 \rightarrow \pi^+\pi^-$ allowed the ϕ -meson mass to be determined: $M_\phi = 1020.8 \pm 0.8$ MeV. (auth)

46444 ASYMMETRY FOR π^+ PHOTOPRODUCTION FROM POLARIZED PROTONS BETWEEN 300 AND 900 MeV AT 90° . Arai, S. (Nagoya Univ., Japan); Fukui, S.; Horikawa, N.; (and others). Phys. Lett.; 40B: No. 3, 426-30 (10 Jul 1972).

The asymmetry of the cross section for π^+ photoproduction from a polarized butanol target was measured at a c.m. angle 90° and photon energies between 300 MeV and 900 MeV by detecting only positive pions. (auth)

46445 COHERENT PHOTOPRODUCTION OF ρ MESONS FROM DEUTERIUM. Overman, Irving Dean. Stanford, Calif.; Stanford Univ. (1972). 81p. University Microfilms Order No. 72-16,766.

Thesis.

Measurements of the forward differential cross section for the process $\gamma + d \rightarrow \rho^0 + d$ were made at Stanford Linear Accelerator Center. The bremsstrahlung photon beam was made by passing the SLAC high power electron beam into an aluminum or copper radiator. A condensation type liquid deuterium target was used for large momentum transfers and a high pressure gas target was used for small momentum transfers. The recoil deuterons were detected and analyzed in momentum and direction by the SLAC 1.6 GeV/c spectrometer using a time-of-flight technique. A rotatable "missing mass" hodoscope was placed in the focal plane of the spectrometer to determine the kinematics. Measurements were made at three energies: 6, 12, and 18 GeV and covered a momentum transfer range of $-0.15 > t > -1.4$ (GeV/c)². The results were interpreted on the basis of the Glauber high energy multiple diffraction model and found to be in excellent agreement. From this analysis values for both the total and differential cross section for the reaction $\rho^0 + N \rightarrow \rho^0 + N$ were extracted in rough agreement with the quark model. The value of σ_T obtained in this experiment was used together with previous hydrogen photoproduction data and with total photoabsorption cross section measurements to determine the γ - ρ coupling constant in the vector dominance model. The results of these two calculations are in disagreement well outside experimental errors and indicate the vector dominance model does not describe all physical processes relating to photon-hadron interactions. (Diss. Abstr. Int., B)

46446 STUDY OF NON-STRANGE BOSON RESONANCES PRODUCED BY QUASI-MONOCROMATIC PHOTON BEAMS OF ENERGIES BETWEEN 4.3 AND 7.5 GeV IN A HYDROGEN BUBBLE CHAMBER. Menke, Michael Matthew. Stanford, Calif.; Stanford Univ. (1972). 187p. University Microfilms Order No. 72-16,755.

Thesis.

The photoproduction of resonances was studied in three exposures of a hydrogen bubble chamber to positron annihilation radiation of 4.3, 5.25 and 7.5 GeV nominal energies. The general analysis procedure and results on the three-prong event topology are presented. The highly constrained reaction $\gamma p \rightarrow \pi^+\pi^-\pi^0$ over the energy range 2 to 8 GeV and the reactions $\gamma p \rightarrow \pi^+\pi^-\pi^0$ and $\gamma p \rightarrow \pi^+\pi^+\pi^-n$ at the annihilation energies were studied. Cross sections are given. Using various models to parameterize ρ^0 photoproduction in $\gamma p \rightarrow \pi^+\pi^-\pi^0$, the mean t -slope found is 7.0 ± 0.4 GeV⁻², while the ρ^0 forward cross section decreases from about $130 \mu\text{b}/\text{GeV}^2$ at 3.3 GeV to about $100 \mu\text{b}/\text{GeV}^2$ at 7.5 GeV. The cross section for $\gamma p \rightarrow \omega p$ is decomposed into an OPE and a diffractive part, which is found to be $1.5 \pm 0.3 \mu\text{b}$. $(d\sigma(\rho)/dt)/d\sigma(\omega)/dt|_{t=0}$ is then 9.4 ± 2.3 at 7.5 GeV. In comparing $d\sigma/dt$ for vector meson production with that of Compton scattering via VDM for photon energies above 4 GeV, it is found that for $\gamma^2/4\pi = 0.32 \pm 0.03$ there is good agreement at all s and t . This same $\gamma^2/4\pi$ relates the forward ρ^0 cross sections to $\sigma_T(\gamma p)$ at all our energies. A search for ρ' $\rightarrow 2\pi$ finds, at the 90% confidence level, $\sigma_{\rho'}(1250) < 0.3 \mu\text{b}$ and $\sigma_{\rho'}(1650) < 0.1 \mu\text{b}$ per 100 MeV width. Inelastic peripheral ρ^0 production is seen in both the proton and the neutron channels, but with present statistics specific nucleon isobars cannot be associated with it. The quasi-two body reactions $\gamma p \rightarrow \pi^-\Delta^{++}$, $\rho^-\Delta^{++}$, and $A_1^0 n$ are observed, with cross sections decreasing with photon energy like E_γ^{-3} . For the first two $a = 1.74 \pm 0.16$ and 0.6 ± 0.2 respectively, indicating that they are unlikely to have the same production mechanism. It follows that if the reaction $\gamma p \rightarrow \rho^-\Delta^{++}$ is due to an OPE process, the required rho radiative decay width (~ 0.5 MeV) is much in excess of the value predicted by SU(3). (Diss. Abstr. Int., B)

46447 PI-ZERO PHOTOPRODUCTION FROM COMPLEX NUCLEI. Meyer, Walter Thomas. Ithaca, N. Y.; Cornell Univ. (1971). 139p. University Microfilms Order No. 72-15,424.

Thesis.

One of the consequences of the vector meson dominance theory for photon interactions is that at high energies a photon incident on a nucleus has its ability to interact inhibited by the interference from a wave of photoproduced vector mesons. The result of this interference is that at low energies, less than about 2 GeV, one would expect the interactions to occur over the entire volume of the nucleus, and that at high energies, above about 8 GeV, they

would occur only near the surface. It is convenient to refer to the nucleus in terms of the effective number of nucleons, Z_{eff} , N_{eff} , or A_{eff} , taking part in the interaction. Thus, excluding final state absorption, A_{eff} , for example, would be expected to vary as the nucleon number A at low energies and as $A^{1/2}$ at high energies. The transition between these two limits is investigated. The process measured was $\gamma + A \rightarrow \pi^0 + A$ where the energies and positions of the two decay photons from the pi-zero were measured. The photon detectors consisted of two arrays, each consisting of 28 lead glass total absorption counters $1\frac{3}{4}$ inches $\times$ $1\frac{3}{4}$ inches $\times$ 15 inches deep. By looking at the shower spread between adjacent counters we get spatial information of the photon and by summing the pulses from these cells we get energy information. At the highest energies the energy resolution was 15 percent fwhm and the spatial resolution was about 0.5 fwhm. A_{eff} was measured for beryllium, carbon, aluminum, copper, silver, and lead at mean photon energies of 3.5, 4.4, 6.3, 8.3 GeV. The range in four momentum transfer was $.1 < -t < .25$ (GeV/c)². Contrary to theoretical expectations no significant transition with energy is seen. The cause of this disagreement is not yet known. (Diss. Abstr. Int., B)

Weak

46448 (LBL-791) RANGE-MOMENTUM CORRELATION IN CARBON AND IRON FOR MUONS OF 500 TO 1500 MeV/c. Field, R. Clive (California Univ., Berkeley. Lawrence Berkeley Lab.), 14 Apr 1972. Contract W-7405-eng-48. 25p. Dep. NTIS.

The range and straggling of positive and negative muons in a stopping medium composed principally of lead, graphite, and steel were investigated experimentally with data from a Bevatron spectrometer. It is concluded that the Bethe-Bloch formula is capable of representing the range behavior of muons between 500 MeV/c and 1500 MeV/c within $\pm 0.3\%$. Values of the mean excitation energies of 79 ± 7 eV for graphite and 287 ± 13 eV for iron are obtained. (auth)

46449 (SLAC-Trans-143) K^0 DECAY IN A WILSON CHAMBER. Astier, A.; Blaskovic, L.; DeCourreges, M. M.; Equer, B.; Lloret, A.; Rivet, P.; Slaud, J. Translated for Stanford Linear Accelerator Center, Calif., from pp 227-34 of report NP-11592 (Vol.1). 10p. (CONF-610904-1). Dep. NTIS.

From Conference on elementary particles; Aix-en-Provence, France (14 Sep 1961).

A Wilson chamber was used to determine the branching ratios for the decays $K_S^0 \rightarrow 3\pi^0$, $K_S^0 \rightarrow \pi^+ + \pi^- + \pi^0$, $K_S^0 \rightarrow \pi^+(\pi^-) + e^-(e^+) + \nu$, and $K_S^0 \rightarrow \pi^+(\pi^-) + \mu^-(\mu^+) + \nu$. The mean K_S^0 lifetime was also found. (J.F.P.)

46450 INVESTIGATION OF LOW- q^2 INELASTIC NEUTRINO EVENTS. Masnou, J. L.; Six, J. (Laboratoire de l'Accelérateur Lineaire, Orsay, France). Nuovo Cim.; 10A: No. 1, 87-96 (1 Jul 1972).

Experimental results on low- q^2 inelastic neutrino interactions are presented and compared with the PCAC hypothesis. Agreement is found for $q^2 < 0.12$ (GeV/c)². Correction terms, coming from the transverse-current contributions, are estimated from a meson dominance model and are found to grow rapidly with q^2 . The PCAC hypothesis with these correction terms is only valid experimentally for $q^2 < 0.2$ (GeV/c)². (auth)

46451 INELASTIC INTERACTION INDUCED BY HIGH-ENERGY MUONS (6 GeV, 12 GeV) AT LOW MOMENTUM-TRANSFER IN NUCLEAR EMULSION. Montret, J. C.; Coupat, B.; Michel, B.; Vazeille, F. (Univ., Clermont-Ferrand, France). Nuovo Cim.; 10A: No. 3, 490-508 (1 Aug 1972).

Inelastic scattering of 6 and 12 GeV muons was studied in Ilford K5 nuclear emulsions. For energy transfers greater than 150 MeV (for π muoproduction threshold) and a four-momentum transfer smaller than 0.1 (GeV/c)², the cross sections are respectively (11.1 ± 1.4) $\mu\text{b/nucleon}$ at 12 GeV and (6.8 ± 0.9) $\mu\text{b/nucleon}$ at 6 GeV. The ratios of the mean values of the longitudinal and transverse cross sections are respectively 0.46 ± 0.15 and 0.25 ± 0.15 . These values are very useful for a comparison between electron and muon. (auth)

46452 INCLUSIVE REACTION $\mu + \text{NUCLEON} \rightarrow \pi + \text{ANYTHING}$ AT 10.1 AND 15.8 GeV/c. Jain, P. L.; Ahmad, Z.; Malucci, R. D.; Potoczak, M. J.; Girard, B. (State Univ. of New York, Buffalo). Lett. Nuovo Cim.; 4: No. 13, 601-7 (29 Jul 1972).

It is shown that the shape and the slope of the curves for p_T^2 and p_T distributions for inclusive pion production reactions in muon-nucleon interactions at 10.1 and 15.8 GeV are the same as those in hadronic reactions, indicating that the muon-nucleon interactions

behave like a hadron. The distribution $d\sigma/dY$, where Y is the rapidity variable, does not show the exact limited target fragmentation at these energies. (L.B.S.)

46453 EXPERIMENTAL LIMIT ON THE NEUTRAL CURRENT IN THE SEMILEPTONIC PROCESSES. Lee, W. (Columbia Univ., New York). Phys. Lett.; 40B: No. 3, 423-5 (10 Jul 1972).

A search was made for the reactions $\nu_\mu + n \rightarrow \nu_\mu + n + \pi^0$ and $\nu_\mu + p \rightarrow \nu_\mu + p + \pi^0$ in a large optical spark chamber as a test for the presence of neutral currents in semi-leptonic weak interactions. No candidate for either reaction was observed. With 90% confidence, the upper limit for the neutral current reactions was found to be: $\{\sigma(\nu_n + p \rightarrow \nu_\mu + \pi^0 + p) + \sigma(\nu_\mu + n \rightarrow \nu_\mu + \pi^0 + n)\} / 2\sigma(\nu_\mu + n \rightarrow \mu^- + \pi^0 + p) < 0.14$. (auth)

46454 EXPERIMENTAL SEARCH FOR SOME RARE K^+ DECAY MODES. Ljung, Donovan Allen. Madison, Wis.; Univ. of Wisconsin (1972). 140p. University Microfilms Order No. 72-15,367.

Thesis.

The stopped K^+ decays, $K^+ \rightarrow \pi^+\gamma\gamma$, $K^+ \rightarrow \pi^0\pi^+\nu$, and $K^+ \rightarrow \pi^+\pi^+\nu$ have been searched for in the 40" heavy liquid bubble chamber at Argonne National Laboratory. The chamber was filled with CF_3Br . No examples of the $K^+ \rightarrow \pi^+\gamma\gamma$ decay mode were found. The experiment was sensitive to pions with kinetic energy 6 to 102 MeV and 114 to 127 MeV. Due to the sensitivity at the lower end of the pion spectrum, a test could be made of the Fujii model. This model uses an off-mass-shell extrapolation of the $K\pi_2$ amplitude to make predictions for this decay. The experimental upper limit on the branching ratio is reported to be 2.2×10^{-5} at the 90% confidence level assuming the pion spectrum appropriate for the Fujii model. This result is inconsistent with the branching ratio predicted by the Fujii model. Assuming a phase space model for the pion spectrum, the upper limit is 3.5×10^{-5} at the 90% confidence level. The decay $K^+ \rightarrow \pi^0\pi^+\nu$ was observed for the first time. Two events were found in which all four converted gammas were seen. These two events give a branching ratio of $1.8^{+0.8}_{-0.6} \times 10^{-5}$ for this decay. The form factor f_1 for the decay is reported as $|f_1| = 0.97^{+0.09}_{-0.10}$. These results are in good agreement with $\Delta I = 1/2$ rule predictions. The study of $K^+ \rightarrow \pi^+\pi^0\gamma$ determined the strength of two direct contributions to this decay as compared with the internal bremsstrahlung contribution. The direct contributions are E1 and M1 dipole radiation, and their respective strengths are given by the parameters γ and β . Twenty-two radiative $K\pi_2$ events with three converted gamma rays were analyzed. The experiment was sensitive to positive pions with kinetic energy 55 to 102 MeV. (Diss. Abstr. Int., B)

46455 EXPERIMENTAL STUDY OF K^+ DECAY. Chiang, I-hung. Rochester, N. Y.; Univ. of Rochester (1972). 114p. University Microfilms Order No. 72-18,807.

Thesis.

An array of optical spark chambers was used to study the decay of 1.84 GeV/c K^+ in flight. The experiment was performed using a partially separated K^+ beam at the Argonne National Laboratory Z.G.S. Accelerator. The apparatus was sensitive to all decay modes, and had a high detection efficiency for the two γ rays from the π^0 . The six principal decay channels of the K^+ were observed in a relatively bias-free fashion. A total of 90,000 events having K^+ decays originating in a specified fiducial region were sorted among these modes. $K_{\mu 3}$ and $K_{\pi 2}$ decays were separated by kinematical analysis using events in which both π^0 γ rays were observed. The branching ratios obtained are: $K_{\mu 2} = (63.16 \pm 0.42)\%$, $K_{\pi 2} = (21.15 \pm 0.26)\%$, $\tau = (5.55 \pm 0.20)\%$, $\tau' = (1.84 \pm 0.06)\%$, $K_{\mu 3} = (3.26 \pm 0.16)\%$, and $K_{e 3} = (5.04 \pm 0.11)\%$. The 2γ opening angle distribution in the $K_{e 3}^+$ data was also studied. $\lambda_+ = 0.04 \pm 0.015$ was found. Using this value of λ_+ and the $K_{\mu 3}$ to $K_{e 3}$ branching ratio $\xi(0) = -0.36^{+0.09}_{-0.11}$ was obtained. (Diss. Abstr. Int., B)

Strong (Baryon-induced)

Refer also to abstracts: 45647, 46590-46592, 46629, 46861, and 46872.

46456 (COO-1112-75) 200-GeV PROTON-PROTON ELASTIC SCATTERING AT HIGH TRANSVERSE MOMENTUM. Ratner, L. G. (Argonne National Lab., Ill.); Krisch, A. D.; Roberts, J. B.; Terwilliger, K. M. (Michigan Univ., Ann Arbor). 5 Jun 1970. 22p. Dep. NTIS.

This is a proposal to study p-p elastic scattering at the highest possible P_T^2 at NAL, using a CH_2 or H_2 target placed directly in the extracted beam and a double arm spectrometer. It is expected to be able to set an upper limit at the level $(d\sigma/dt/d\sigma/dt)_{t=0} \approx 10^{-14}$. This would be sufficient to determine if there are exactly three regions in the p-p interaction with considerable precision. (auth)

46457 (KFK-1509) PROTON-PROTON FINAL STATE INTERACTION IN THE REACTION $d + p \rightarrow p + p + n$. Matthaei, H. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Experimentelle Kernphysik). Mar 1972. 77p. (In German). Dep. NTIS (U. S. Sales Only).

Thesis. Submitted to Karlsruhe Univ. (West Germany).

The proton-proton final state interaction in the reaction $d + p \rightarrow p + p + n$ was investigated systematically by kinematically complete experiments using the 52.3 MeV external deuteron beam of the Karlsruhe Isochronous Cyclotron. The analysis of the data in terms of a slightly modified Watson Migdal ansatz resulted in typical values of the p - p scattering length which are in good agreement with the value known from free p - p scattering. (The same holds for the value of n - p singlet scattering length measured under the same kinematical conditions, as compared with the result of free n - p scattering.) In a special set of experiments the angular distribution of final state interacting p - p and n - p pairs in their c.m. system was measured in order to test the applicability of the two step reaction model to the reaction under investigation. Finally a new approach to a simple theory of proton-proton final state interaction was developed. Within the framework of this model the reaction is assumed to proceed via a two step process with a zero range break-up mechanism in the first step which is followed by a p - p final state interaction. The model includes pure Coulomb final state interaction in all partial waves. The results of the calculation are in good agreement with experimental data. (auth)

46458 QUANTUM NUMBERS AND TOTAL AND PARTIAL WIDTHS OF THE RESONANCE $NN(1970)$. Dal'karov, O. D.; Kerbikov, B. O.; Shapiro, I. S. (Inst. of Theoretical and Experimental Physics, Moscow). JETP Lett. (USSR) (Engl. Transl.); 15: No. 7, 305-8 (5 Apr 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 7, 430-4 (5 Apr 1972).

The meson $NN(1970)$ has been observed in the interaction $\bar{p} + p \rightarrow K_1^0 + K_1^0$ and in backward elastic $\bar{p}p$ scattering. Based on these data, a value for the ratio Γ_{NN}/Γ was estimated in several ways. Since the result was of the order of magnitude of unity, it was concluded that the $NN(1970)$ meson is a resonant state in the nucleon-antinucleon system. It was determined to be a ${}^3D_3$ state with the quantum numbers $I^{PC} = 3^{--}$. (R.W.R.)

46459 PARTICLE PRODUCTION IN PROTON INTERACTIONS IN NUCLEI AT 24 GeV/c. Eichten, T. (Physikalische Inst., Aachen); Haidt, D.; Pattison, J. B. M.; (and others). Nucl. Phys.; B44: No. 2, 333-43 (1972).

Particle production by 24 GeV/c protons from Be, B_4C , Al, Cu, and Pb was measured. Pion, kaon, proton and antiproton production spectra measured over a range of angles from 17 to 127 mrad and momenta from 4 to 18 GeV/c are given in a table. (auth)

46460 EXPERIMENTAL DETERMINATION OF THE TWO-PARTICLE CORRELATIONS IN 60 GeV/c π^+p AND 67 GeV/c pp INCLUSIVE REACTIONS. Fialkowski, K.; Rybicki, K.; Wit, R. (Jagellonian Univ., Krakow). Nucl. Phys.; B44: No. 2, 509-19 (1972).

Two-particle correlations are studied for inclusive π^+p interactions at 60 GeV/c and pp collisions at 67 GeV/c. Positive correlations are observed for slow c.m.s. particles. (auth)

46461 COLLECTIVE INTERACTIONS OF NUCLEONS OF HIGH-ENERGY COSMIC HEAVY NUCLEI. Grigorov, N. L.; Zhuravlev, D. A.; Kozlov, B. D.; (and others) (Moscow State Univ.). Dokl. Akad. Nauk SSSR; 202: No. 1, 65-6 (Jan 1972). (In Russian).

Available experimental data are used in preliminary studies to show that nucleons which are bound in nuclei form pions of high energy with greater probability than free protons of the same energy. Two experimental approaches to a search for collective interactions of nucleons with complex nuclei are given: the study of pion spectra generated by nuclei of known energy, and the measurement of the distribution of the portion of energy transferred to all π^0 -mesons during the interaction of a nucleus with known z and known energy. (K.S.W.)

46462 POLARIZATION AT LARGE ANGLES. Yokosawa, A. (Argonne National Lab., Ill.). pp 117-43 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

Polarization parameters in the backward region and at large momentum transfer (2.0 to 5.0 GeV/c) are discussed. Previous experimental data obtained for the backward peaks in π^+p , K^+p , and pp and $\bar{p}p$ elastic scattering are successively examined and

compared with the theoretical baryon exchange models with or without Regge cuts. (France)

46463 COMPARISON OF $\bar{p}p \rightarrow \pi^-\pi^+(K^+K^-)$ ANNIHILATIONS AND $\pi^+p(K^+p)$ BACKWARD ELASTIC SCATTERING WITH CROSS-INGE SYMMETRY PREDICTIONS. Baglin, C. (CERN, Geneva). Ecole Polytechnique, Paris). pp 233-46 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

Recent data obtained for $\bar{p}p$, π^+p , K^+p elastic scattering, and $\bar{p}p$ annihilations into $\pi^-\pi^+$ and K^+K^- at 5 GeV/c allowed crossing relations to be experimentally checked. Quantitative comparisons between meson-nucleon backward elastic differential cross sections and the corresponding $\bar{p}p$ annihilations cross sections are presented, the correction factor R defined by Barger and Cline being taken into consideration. (France)

46464 ELASTIC SCATTERING AND ISOBAR PRODUCTION IN PROTON-PROTON COLLISIONS. Matthiae, G. (Istituto Superiore di Sanita, Rome). pp 373-96 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

Recent results obtained for elastic scattering and two-body inelastic processes in proton-proton collisions are given. A structure is shown in pp elastic scattering around $1.5(\text{GeV})^2$. This structure develops rapidly between 7 and 10 GeV/c and becomes more pronounced for further increases in energy. Evidence is given for the excitation of the nucleon resonances $N(1520)$, $N(1688)$, $N(2190)$ and $N(1400)$ in the two-body inelastic process. $N(1520)$ and $N(1688)$ cross sections exhibit a gradual change in slope inside the t region where the elastic scattering is structured. The momentum transfer dependence of these processes for $2 \leq t \leq 7(\text{GeV})^2$ is the same as for elastic scattering. (France)

46465 PROPERTIES OF MESON RESONANCES AS OBSERVED IN $\bar{p}$ - p ANNIHILATIONS AT REST. Montanet, L. (CERN, Geneva). pp 425-55 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

Experimental data obtained for $\bar{p}$ - p annihilation at rest are presented and analyzed in terms of the final-state interaction model. Indications of the presence of at least one scalar resonance, the epsilon, and perhaps a second resonance in the 1200-MeV region exist, the two resonances being separated by a zero of the $\pi^+\pi^-$ S-wave amplitude at 940 MeV. The $\omega\rho\pi^2$ enhancement observed at 1250 MeV is shown to be an impure state, two resonances probably being present, the B-meson $J^P = 1^+$, and the ρ $J^P = 1^-$ or a π $J^P = 0^-$. Veneziano amplitudes favor the second solution. The $\eta^0\pi^+$ enhancement also observed at 1300 MeV is shown to be associated to a 2^+ or a 0^+ resonance (1^- resonance being excluded). (France)

46466 NEUTRON MATTER. Vlasov, N. A. At. Energ. (USSR); 33: No. 1, 539-45 (Jul 1972). (In Russian).

The status of knowledge of nucleon structure and neutron stars is considered. Strong inelastic scattering of up to 20-GeV electrons by nucleons is evidence for the existence in nucleons of almost point components called partons. The discovery of pulsars attests to the existence of neutron stars having densities at their center exceeding nuclear densities, magnetic fields of the order of 10^{12} oersted, neutron superfluidity in the star mantle, and a variegated nuclear composition in the hard crust. (4 figures) (auth)

46467 DISCUSSION OF AN ENHANCEMENT AT ABOUT 2700 MeV/c² IN THE EFFECTIVE MASS OF THE $p\pi^-X^0$ SYSTEM IN THE REACTION $pp \rightarrow pp\pi^+\pi^-X^0$ AT 19 GeV/c INCIDENT-PROTON MOMENTUM. Haldorsen, J.; Jacobsen, T. (Univ. of Oslo). Nuovo Cim.; 10A: No. 3, 468-74 (1 Aug 1972).

In the reaction $pp \rightarrow pp\pi^+\pi^-X^0$ at 19 GeV/c an enhancement is seen in the effective mass of the $p\pi^-X^0$ system at about 2700 MeV/c², which probably is due to $\Delta(2850)$ production. A lower limit of $\frac{3}{2}$ is found for the spin of the $\Delta(2850)$. (auth)

46468 STUDY OF THE REACTIONS $\bar{p}p \rightarrow \bar{p}p$ AND $\bar{p}p \rightarrow \pi^-\pi^+X$ USING A POLARIZED PROTON TARGET. Etkin, Asher. New

Haven; Yale Univ. (1971). 208p. University Microfilms Order No. 72-16,319.

Thesis.

The results of measurements of asymmetry in $\bar{p}p \rightarrow \pi^+\pi^-$ (1.6 GeV/c incident $\bar{p}$ momentum) and $\bar{p}p \rightarrow \bar{p}p$ (1.6 and 2.5 GeV/c incident $\bar{p}$ momenta) are presented. The $\bar{p}p \rightarrow \pi^+\pi^-$ asymmetries were generally large and positive over the angular range covered. Asymmetries for $\bar{p}p \rightarrow \bar{p}p$ were also generally positive. The $\bar{p}p \rightarrow \pi^+\pi^-$ asymmetry data are the first such data obtained and have been fitted, along with all other $\bar{p}p \rightarrow \pi^+\pi^-$ data, to a partial wave expansion. The $\bar{p}p \rightarrow \bar{p}p$ data were parameterized by a spin-dependent diffraction model. An extensive description of a lanthanum magnesium nitrate polarized proton target is presented. Particular attention is paid to the nuclear magnetic resonance system used to measure the target polarization, and to non-linearity effects. Because the cryogenic system is somewhat unusual, an exhaustive description is given. The system described has several notable features: stable calibration, small non-linearity corrections, and automated measurement of polarization. The high resolution detection apparatus permitted the asymmetry measurements of the rare reaction $\bar{p}p \rightarrow \pi^+\pi^-$. This system also permitted a unique measurement to be made of the spatial distribution of the target polarization. (Diss. Abstr. Int., B)

46469 ANTIPROTON-PROTON INTERACTIONS FROM 1.51 TO 2.90 GeV/c. Parker, Donald Lester. East Lansing, Mich.; Michigan State Univ. (1971). 109p. University Microfilms Order No. 72-16,490.

Thesis.

In a bubble chamber experiment cross sections were obtained for zero, two, four, six and eight prong topologies, and the reaction cross sections and resonance production cross sections within each topology were obtained for nine antiproton momenta from 1.51 to 2.90 GeV/c. The extrapolation of small angle elastic scattering to $t = 0$ is discussed and compared with the results of other experiments. The differential cross sections are fitted to an adaptation of the Frahn-Venter optical model and also compared to Regge pole model predictions. All other channels are investigated with particular emphasis given to discovering the origin of the structure seen in the total cross section for antiproton-proton scattering. The structure cannot be explained as threshold effects, and viewed as direct channel effects, some channels can be ruled out as the origin of the structure. Pion multiplicity in the all-pionic annihilations was also investigated and comparisons of the predictions of a statistical model and a multiperipheral model are made. (Diss. Abstr. Int., B)

46470 FORWARD DIFFERENTIAL CROSS SECTION IN $\bar{p}p \rightarrow \bar{n}n$. Pittman, Roland Nathan. Stony Brook, N. Y.; State Univ. of New York (1972). 73p. University Microfilms Order No. 72-16,967.

Thesis.

The forward charge exchange reaction $\bar{p}p \rightarrow \bar{n}n$ was studied in an experiment at Brookhaven AGS. The incident antiprotons were identified by time of flight and a Cerenkov counter (at momenta above 2 GeV/c). The antineutrons and neutrons were detected with similar steel-scintillator sandwich hodoscope detectors. A liquid hydrogen target was used. Differential cross section data were taken at forty momenta from 1.00 to 3.17 GeV/c in 3% steps. The four-momentum transfer bite subtended was $|t| \leq .12(\text{GeV}/c)^2$. Results are given for the differential cross section $d\sigma/dt$ and $d\sigma/d\Omega$. Two features of the data are (1) the observation of a sharp forward peak for $|t| \leq .02(\text{GeV}/c)^2$ with a width of about one pion mass squared and (2) a dependent structure in the differential cross section which can be interpreted as direct channel resonance formation. The slope of $d\sigma/dt$ for $|t| \leq .02(\text{GeV}/c)^2$ is smaller than that predicted on the basis of a π -exchange Regge model; this, however, is consistent with the experimental t -resolution in the forward peak region. The angular dependence of the structure is consistent with a direct channel resonance of spin $3 \leq J_R \leq 5$. Its mass and width are consistent with those found in a previous study of the $\bar{p}p$ and $\bar{p}d$ total cross sections ($M = 2190$ MeV, $\Gamma = 85$ MeV). (Diss. Abstr. Int., B)

Strong (Meson-induced)

Refer also to abstracts: 45651, 46517, 46636, and 46869.

46471 (BNL-16954) EVIDENCE FOR EXISTENCE OF A NEW MESON IN THE 955 MeV MASS REGION. Eisner, Robert L. (Brookhaven National Lab., Upton, N. Y.). 5 Jun 1972. 17p. (CONF-720409-5). Dep. NTIS.

From third international conference on experimental meson spectroscopy; Philadelphia, PA. (28 Apr 1972).

Evidence is presented for the existence of a new meson in the

955 MeV mass region. The properties of the $M(953)$ are compared with those of the well-known $\eta'(958)$. (auth)

46472 (CONF-710816-12) MEASUREMENT OF THE RADIATIVE NEUTRAL DECAY MODES OF THE ω^0 AND ϕ^0 MESONS WITH THE ORSAY STORAGE RING. Benakkas, D.; Cosme, G.; Jean-Marie, B.; (and others) (Paris Univ., Orsay (France). Laboratoire de l'Accélérateur Lineaire). 1971. 18p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

The branching ratios for the decays $\phi^0 \rightarrow \eta^0(\pi^0) + \gamma$, $\omega^0 \rightarrow \pi^0 + \gamma$, $\omega^0 \rightarrow \pi^+ + \pi^- + \pi^0$, and $\omega^0 \rightarrow 2\pi^0 + \gamma$ are found using a large solid angle detector. (J.F.P.)

46473 (COO-1195-239) STUDY OF A_3 IN $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$. Ascoli, Giulio (Illinois Univ., Urbana). [nd]. Contract AT(11-1)-1195. 9p. (CONF-720409-4). Dep. NTIS.

From third international conference on experimental meson spectroscopy; Philadelphia, PA. (28 Apr 1972).

Analysis of bubble chamber and spark chamber data on the reaction $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$ at incident momenta from 11 to 40 GeV, shows that the A_3 enhancement in the $\pi^+\pi^-\pi^-$ system has $J^P = 2^-$, $M = 1.65$ GeV, $\Gamma = 0.27$ GeV, and decays entirely into $f^0\pi^-$. The phase of the A_3 does not show a resonant behavior. (auth)

46474 (COO-1428-306) FINAL STATES WITH THREE CHARGED PARTICLES AND A VISIBLE Λ FROM K^-d INTERACTIONS AT 4.5 GeV/c. Ammann, A. C.; Carmony, D. D.; Garfinkel, A. F.; Gutay, L. J.; Miller, D. H. (Purdue Univ., Lafayette, Ind. Dept. of Physics); Yen, W. L. (Indiana-Purdue Univ., Indianapolis, Dept. of Physics). 1972. Contract AT(11-1)-1428. 59p. Dep. NTIS.

A study is made of four or five body final states produced by the interaction of 4.48 GeV/c K^- mesons on neutrons. The data result from a 372,000 picture exposure in the Argonne National Laboratory 30 in. deuterium-filled bubble chamber. The reactions studied include $K^-n \rightarrow \Lambda\pi^+\pi^-\pi^-$, $K^-n \rightarrow \Sigma^0\pi^+\pi^-\pi^-$, $K^-n \rightarrow \Lambda\pi^+\pi^-\pi^0$, $K^-n \rightarrow \Lambda\pi^+\pi^-\pi^0\pi^0$, $K^-n \rightarrow \Lambda\pi^+\pi^-\pi^0\pi^0\pi^0$, $K^-n \rightarrow \Lambda\pi^+\pi^-\pi^0\pi^0\pi^0\pi^0$, where all the lambda particles decayed visibly. Evidence is presented for the existence of the $\Sigma(1640)$ whose mass and width are 1642 ± 12 MeV/c² and 55 ± 24 MeV/c², respectively. The branching ratios are found to be consistent with an octet assignment. The A_2 data are presented and are compatible with $J^{PC} = 2^{++}$. In addition, the productions of ρ^0 , ω , and ϕ^0 are discussed in conjunction with simultaneous $\Sigma^-(1385)$ production and the results are compared to quark model predictions. (auth)

46475 (IS-2886) OBSERVATION OF THE $M^0(940)$ AND $\delta^0(963)$. Cheshire, D. L.; Jacobel, R. W.; Lamb, R. C.; Peterson, F. C. (Iowa State Univ. of Science and Technology, Ames); Hoffman, E. W. (Argonne National Lab., Ill.); Garfinkel, A. F. (Purdue Univ., Lafayette, Ind.). 1 Jun 1972. 9p. (CONF-720409-7). Dep. NTIS.

From third international conference on experimental meson spectroscopy; Philadelphia, PA. (28 Apr 1972).

In a scintillation counter experiment two narrow, neutral mesons denoted $M^0(940)$ and $\delta^0(963)$ in the reaction $\pi^-p \rightarrow M^0n$ at 2.4 GeV/c were observed. Also the well established $\eta'(958)$ was observed. (auth)

46476 (JINR-P1-6408) INVESTIGATION OF GENERATION OF PARTICLES AND RESONANCE STATES IN INTERACTIONS OF 2.34 GeV/c, 5 GeV/c, AND 9 GeV/c π^- -MESONS WITH XENON NUCLEI. Slovinskii, B.; Strugal'skii, Z. S. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of High Energy). 1972. 13 p. (In Russian). Dep. NTIS (U. S. Sales Only).

The results of investigation of particle generation, mainly of π^0 -mesons and resonance states, in π^- -Xe interactions are described. (auth)

46477 (LBL-378) STUDY OF DIPION PRODUCTION IN THE REACTION $\pi^-p \rightarrow \pi^0\pi^0n$ BETWEEN 1.6 AND 2.4 GeV/c. Skuja, Andris (California Univ., Berkeley. Lawrence Berkeley Lab.). 25 Feb 1972. Contract W-7405-eng-48. 172p. Dep. NTIS.

Thesis.

An experiment was performed at the Berkeley Bevatron to study the $I = 0$, $J = 0$ π^-p system in the reaction $\pi^-p \rightarrow \pi^0\pi^0n$, at beam momenta of 1.60 to 2.40 GeV/c in 0.20-GeV/c intervals. The reaction occurred in a 20-cm liquid hydrogen target and the final state was identified by detecting the neutron and the γ -rays from the π^0 decays. The detection system consisted of a large cubic array of lead-plate optical spark chambers surrounding the target for converting γ -rays and a set of 20 plastic-scintillator neutron counters to measure time-of-flight and direction from the target, in the polar-lab-angle region of 12 to 72 degrees. The spark chambers were seven to eight radiation lengths thick and

covered five sides of a cube, with the sixth side (beam entrance face) being nearly closed by lead-scintillator sandwich gamma counters. A sample of 5000 four-gamma events was obtained that passed the kinematic fit to the hypothesis $\pi^-p \rightarrow \pi^0\pi^0n$ with a 1% confidence level. The total cross sections for the production of $\pi^-p \rightarrow \pi^0\pi^0n$ at each incident momentum were obtained. The dipion production distribution was parametrized using a modified one pion exchange model with form factors, and the off-mass-shell $\pi\pi$ scattering matrix element extracted. It was found that the $\pi^0\pi^0$ production distribution was more peripheral than the OPE model predictions and that above 850 MeV of the dipion mass, the matrix element varied rapidly, indicating a rapid variation of the $I = 0, J = 0$, phase shift in this region. It was also found that at low t , the reaction had substantial Δ^0 production ($\sim 50\%$). (auth)

46478 (ORNL-tr-2544) CROSS SECTION OF PION-NUCLEON INTERACTIONS. Barashenkov, V. S.; Okhlopova, V. A. (Joint Inst. for Nuclear Research, Dubna (USSR)). Translated for Oak Ridge National Lab., Tenn., from report JINR-P2-5924. 3p. Dep. NTIS.

Interpolation curves for total cross sections previously presented are refined and followed up to the high energy region for the reactions $\pi^- + p$ and $\pi^+ + p$. The total cross sections for elastic scattering, elastic scattering with charge exchange, and inelastic scattering for these reactions are given. (J.F.P.)

46479 (ORO-3065-24) PROGRESS REPORT [ON HIGH-ENERGY PHYSICS], JULY 1, 1971-JUNE 30, 1972. Walker, W. D.; Fortney, L. R. (Duke Univ., Durham, N. C. Dept. of Physics). 10 Jul 1972. Contract AT(40-1)-3065. 6p. Dep. NTIS.

A summary of research on the $\pi^+ + d$ experiment and pion reactions with neon is given. (J.F.P.)

46480 K^-p INTERACTIONS AT 10 GeV/c. Buschbeck, B.; Froehlich, A.; Markytan, M.; (and others) (Institut fuer Hohenenergiephysik, Vienna). Oesterr. Akad. Wiss., Math.-Naturwiss. Kl., Sitzungsber., Abt. II; 179: No. 8-10, 280-303(1971). (In German).

Presently available information about K^-p interactions at 10 GeV/c was reviewed. There were 3,600 bubble chamber events which yielded for the K^-p elastic scattering a cross section of 3.20 ± 0.14 mb, with a ratio of elastic to total cross section of 0.142 ± 0.006 . This value was somewhat lower than the 0.185 predicted as the upper limit by the Van Hove model for asymptotic energies. The dynamics of two-body reactions, quasi-two-body reactions, and multibody reactions were theoretically analyzed, and the data were compared with the experimental findings. Up to May 1969, seven Ω^- were discovered. Their decay was established as well as a new resonance demonstrated. (H.B.G.)

46481 PION EXCHANGE IN THE REACTION $\pi p \rightarrow pN$ AT 6 GeV/c. Scarr, J. Michael; Lai, Kwan-wu (Brookhaven National Lab., Upton, N. Y.). Phys. Rev. Lett.; 29: No. 5, 310-12 (31 Jul 1972).

A dip was observed at $-t \approx 0.5$ GeV² in $(\rho_{H0}^0 d\sigma/dt)(\pi p \rightarrow pN)$ for ρ^- , ρ^0 , and ρ^+ at 6 GeV/c. The dominant contribution to $(\rho_{H0}^0 d\sigma/dt)(\pi p \rightarrow pN)$ is from π exchange. (auth)

46482 CHARGE ASYMMETRY IN THE DECAY $\eta \rightarrow \pi^+\pi^-\gamma$. Thaler, J. J.; Appel, J. A.; Kotlewski, A.; (and others) (Columbia Univ., New York). Phys. Rev. Lett.; 29: No. 5, 313-16(31 Jul 1972).

The charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\gamma$ was measured. η mesons were produced in the reaction $\pi^-p \rightarrow \eta n$ at 730 and 810 MeV/c incident pion momentum. The asymmetry is 0.005 ± 0.006 for 36 155 events. No evidence for C-invariance violation in this decay was seen. (auth)

46483 MEASUREMENT OF THE CHARGE ASYMMETRY IN THE DECAY $\eta \rightarrow \pi^+\pi^-\pi^0$. Layter, J. G.; Appel, J. A.; Kotlewski, A.; (and others) (Columbia Univ., New York). Phys. Rev. Lett.; 29: No. 5, 316-19(31 Jul 1972).

The charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\pi^0$ was measured. η 's were produced in the reaction $\pi^-p \rightarrow \eta n$ at 730 and 810 MeV/c incident pion momentum. An asymmetry of -0.0005 ± 0.0022 was obtained for 220 659 events. No evidence for C-invariance violation in this decay was seen. (auth)

46484 EXPERIMENTAL SPECTRA FOR 12 GeV/c $k^+p \rightarrow pK^+\phi$ AND $pK^+\omega$. Davis, Philip J.; Alston-Garnjost, Margaret; Barbaro-Galtieri, Angela; (and others) (Univ. of California, Berkeley). Nucl. Phys.; B44: No. 2, 344-54(1972).

Experimental distributions are presented for the reactions $K^+p \rightarrow pK^+\phi$, $\phi \rightarrow K^+K^-$ and $K^+p \rightarrow pK^+\omega$, $\omega \rightarrow \pi^-\pi^+\pi^0$ at 12 GeV/c. The data were obtained from a 35 events/ μ b exposure of the SLAC

82-inch HBC. The cross sections for these reactions are (11 ± 1) μ b and (47 ± 3) μ b. $K\phi$, $K\omega$, and $p\omega$ low-mass enhancements were observed with no evidence for narrow structure. The momentum-transfer and angular-decay spectra of these systems are presented. (auth)

46485 REACTIONS $K^- + p \rightarrow \Lambda(\Sigma^0) + \text{VECTOR MESON AT 3.95 GeV/c}$. Rouge, A. (Ecole Polytechnique, Paris); Videau, H.; Videau, I.; (and others). Nucl. Phys.; B44: No. 2, 365-89(1972).

Complete results, concerning the five reactions $K^-p \rightarrow \Lambda\omega$, $K^-p \rightarrow \Lambda\phi$, $K^-p \rightarrow \Lambda\rho$, $K^-p \rightarrow \Sigma^0\rho$, and $K^-p \rightarrow \Sigma^0\rho$, are presented. The experimental data are well described by exchange mechanisms and the agreement with the SU(3) symmetry predictions is excellent. (auth)

46486 CROSS SECTIONS FOR (π^-, π^-n) REACTIONS ON ^{18}F AND ^{31}P IN THE 3 TO 3 RESONANCE REGION. Plendl, H. S. (Florida State Univ., Tallahassee); Burch, D.; Eberhard, K. A.; (and others). Nucl. Phys.; B44: No. 2, 413-21(1972).

Activation cross sections for the reactions $^{18}\text{F}(\pi^-, \pi^-n)^{18}\text{F}$ and $^{31}\text{P}(\pi^-, \pi^-n)^{30}\text{P}$ were measured at pion energies from 140 to 240 MeV and found to reach maxima of 39 ± 5 mb and 60 ± 11 mb, respectively, near 180 MeV. The results are compared with calculated cross sections and with the results of other measurements. (auth)

46487 INTERACTION OF K^- MESONS WITH NUCLEI. Burhop, E. H. S. (University Coll., London). Nucl. Phys.; B44: No. 2, 445-59(1972).

The experimental data for K^- capture by nuclei is interpreted in terms of K^- nucleon scattering amplitudes in a region of negative relative kinetic energy. It is shown that constant scattering length solutions which represent the data in the region of positive relative kinetic energy up to 100 MeV do not satisfactorily describe the interaction with bound nucleons and a modified scattering length solution is proposed. Measured values of the widths of K^- -mesic X-ray lines which require a strongly attractive K^- nucleons optical potential are successfully interpreted in terms of K^-N scattering lengths appropriate for bound nucleons. The information obtainable about relative neutron and proton density distributions in the nuclear periphery from studies of this kind is discussed. The need for more data on interactions of K^- in hydrogen at low kinetic energies and at rest and in other elements at rest is emphasized. (auth)

46488 FORWARD PRODUCTION OF DELTA(1236) IN PION-PROTON SCATTERING. Derado, I. (Max-Planck-Institut fuer Physik und Astrophysik, Munich). pp 39-47 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

The following reactions were studied: $\pi^+p \rightarrow \pi^+\pi^+n$ and $\pi^-p \rightarrow \pi^-\pi^0p$ around 4 GeV/c. Only events with $\cos \theta \geq 0$ were considered. Theta being the overall c.m.s. angle between the incident pion and the nucleon pion system. Effective masses were investigated for the $n\pi^-$, $p\pi^-$ and $p\pi^0$ systems. The only statistically significant structure was observed in the $n\pi^-$ effective mass. The angular distribution of π^- in $n\pi^-$ (c.m.s.) is presented. The differential cross sections for Δ^- , Δ^+ , Δ^0 and Δ^{++} are given as a function of U (momentum transfer). They were calculated for $U = 0$ and $U = U_{\max} = 0.2(\text{GeV}/c)^2$ in the case of Δ^{++} and Δ^- . Theoretical values and experimental results are compared. (France)

46489 EXPERIMENTAL SURVEY OF THE REACTIONS $\bar{K} N \rightarrow \pi Y$ AT MEDIUM AND HIGH ENERGY. Barloutaud, R. (CEN de Saclay, Gif-sur-Yvette, France). pp 49-63 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Recent results obtained for two body or quasi two body reactions of the $\bar{K} N \rightarrow \pi Y$ type, above 3 GeV/c, are surveyed as far as the simplest description seems to be a description in terms of meson or baryon exchange. Total cross sections are given. Forward scattering is discussed: upper limit of the $I = \frac{3}{2}$ channel amplitude; shape of $d\sigma/dt$ distribution; polarization; comparison with $\pi N \rightarrow KY$ reversed line reactions. Results obtained for backward scattering are also summarized. (France)

46490 POLARIZATION IN π^+-p BACKWARD ELASTIC SCATTERING AT HIGH ENERGY. Poulet, M. (Univ. Paris, Orsay, France). pp 97-104 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

The polarization parameter in backward elastic scattering $\pi p \rightarrow \pi p$ at 6 GeV/c was measured. The following two parameters were determined: the coplanarity of the three particles (incident, scattered, recoiled), and the angular correlations. Both must be zero for elastic backward events. Preliminary results are given. They are in disagreement with the predictions of pure Regge models and cuts based on the Carlitz-Kislinger method or the weak cut model, especially in the region of small w . (France)

46491 POLARIZATION IN CHARGE EXCHANGE REACTIONS. Guisan, O. (CEN de Saclay, Gif-sur-Yvette, France); Bonamy, P.; Borgeaud, P.; (and others). pp 105-16 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

New preliminary results concerning the polarization parameter $P(t)$ in the $\pi^+ p \rightarrow \pi^0 n$ reaction are presented. Measurements were achieved at 5.0 and 8.0 GeV/c from 0 to 2.0 (GeV/c)². The experimental lay-out is briefly described and the background problem encountered when using a polarized target is examined. Data obtained exclude a zero of $P(t)$ for $-t \leq 0.6$ (GeV/c)². Below this value a large positive polarization was found and above this value the polarization is compatible with zero within large errors, data obtained at 5.0 and 8.0 GeV/c being quite similar. Models involving double poles or double trajectories and most of the models involving cuts or absorption are not compatible with the results obtained. Models involving ρ' seem to work much better. (France)

46492 ELASTIC SCATTERING OF π^+ , K^+ , AND $\bar{p}$ ON PROTONS AT 5 GeV. Treille, D. (CERN, Geneva). pp 213-32 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

The complete angular distributions in $\pi^+ p$, $K^+ p$ and $\bar{p} p$ elastic scattering at 5 GeV were measured. A good sensitivity allowed the cross sections to be accurately measured at large angles for all the reactions, the cross sections being 10⁵ times smaller than in the case of forward scattering. Small backward peaks were unambiguously detected in the exotic channels: $K^- p \rightarrow p K^-$ and $\bar{p} p \rightarrow \bar{p} p$. (France)

46493 EVIDENCE FOR DOUBLE DIFFRACTIVE DISSOCIATION IN $\pi^+ p$ REACTIONS AT 5 GeV/c. Rinaudo, G. (Univ., Turin. Istituto Nazionale di Fisica Nucleare, Turin). Werbrouck, A. E.; Rost, M.; (and others). Nuovo Cim.; 10A: No. 1, 1-9(1 Jul 1972).

Evidence of double diffractive dissociation in 5 GeV/c $\pi^+ p$ reactions is found by analyzing angular distributions in selected longitudinal phase-space regions. The size of the observed effect is compatible with expectations from factorization. (auth)

46494 MEASUREMENT OF THE SLOPE PARAMETER IN THE DECAY $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$. Krenz, W. (Technische Hochschule, Aachen); Hopkins, H. W. K.; Evans, G. R.; Muir, J.; Peach, K. J. Lett. Nuovo Cim.; 4: No. 6, 213-19(10 Jun 1972).

The slope parameter in $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$ decay in a heavy-liquid bubble chamber was measured. The slope parameter was found to be -0.277 ± 0.018 . Also a test was made for CP violation in the charge asymmetry δ . It was found that $\delta = -0.0081 \pm 0.045$. (J.F.P.)

46495 PARTIAL WAVE ANALYSIS OF $\pi^+ p \rightarrow \pi^0 \Delta^{++}$ (1236) IN THE CENTER OF MASS ENERGY REGION 1820 TO 2090 MeV. Mehtani, Usha. Riverside, Calif.; Univ. of California (1971). 100p. University Microfilms Order No. 72-17,060.

Thesis.

The differential cross sections $d\sigma(\cos \theta_s)/d\Omega$ and Δ density matrix elements $\rho_{33}(\cos \theta_s)$, $\text{Re } \rho_{31}(\cos \theta_s)$ and $\text{Re } \rho_{3,-1}(\cos \theta_s)$ were measured for the reaction $\pi^+ p \rightarrow \pi^0 \Delta^{++}$ (1236) at seven incident momenta 1.28, 1.34, 1.42, 1.55, 1.67, 1.77 and 1.84 GeV/c. With these experimental data, an energy dependent partial wave analysis of this reaction was performed. It is found that the inelastic channel $\pi N \rightarrow \pi \Delta$ is dominated by F_{37} (1950) resonance with the branching fraction parameter $x_{\pi N \pi \Delta} = .25_{-0.10}^{+0.10}$. Other partial wave amplitudes which are strong in this energy interval are FF_5 , PP_1 , DD_5 , PP_3 and SD_1 . Phase shift analysis in the elastic channel shows the presence of the F_{37} (1950) resonance and suggests the existence of three other res-

onances F_{35} (1890), P_{31} (1910) and D_{35} (1960). Of the latter three resonances only P_{31} (1910) is indicated by the data with $x_{\pi N \pi \Delta} \approx .04$. The reaction $\pi^+ p \rightarrow \pi^0 \Delta^{++}$ goes by pure ρ exchange in the t -channel. All the angular distributions have features characteristic of Regge ρ trajectory exchange, i.e., they have dips at $t \approx 0$ and $t \approx -0.5$ (GeV/c)². The s -channel partial wave amplitudes were projected from the Regge ρ amplitude. The F_{37} amplitude, which is predominant in the s -channel, is negligible here. Also P_{31} is insignificant in the ρ amplitude. However, suggestive similarities between the partial wave fits and the s -channel projections of the Regge ρ amplitude are observed. (Diss. Abstr. Int., B)

46496 RHO-MESON PRODUCTION BY PIONS AT 15 GeV/c: A STUDY OF VECTOR DOMINANCE AND OF RHO-OMEGA INTERFERENCE. Ratcliff, Blair Norman. Stanford, Calif.; Stanford Univ. (1972). 170p. University Microfilms Order No. 72-16,778.

Thesis.

The results of a wire spark chamber experiment studying the reaction $\pi^- p \rightarrow \pi^+ \pi^- n$ at 15 GeV/c are presented. The differential cross section, density matrix elements, and dipion mass distributions are given in the rho region for momentum transfers (t) less than 0.30 (GeV/c)². The density matrix elements and differential cross section exhibit structure in the forward direction ($-t < m_\pi^2$). Using the vector dominance model, a comparison is made with polarized and unpolarized single-pion photoproduction data which also exhibit structure in the forward direction. Qualitative agreement with the vector dominance model is found and, in particular, the transverse rho differential cross section displays a pronounced forward peak, analogous to that observed in single-pion photoproduction. Quantitative agreement is found for the unnatural-parity-exchange cross section and for the asymmetry at small momentum transfers. Elsewhere there are disagreements in detail which may not be explained by changes in normalization or the value of the rho-photon coupling constant. The data show a significant (>4 standard deviations) rho-omega interference effect in the dipion mass distribution for $0.1 < |t| < 0.3$ (GeV/c)². The overall phase between the amplitudes in the observed data is dominated by transverse production and is shown to be -1.40 ± 0.45 (rad) which differs from the phase found at lower energies by $\sim \pi/2$. The lower limit on the branching ratio $\Gamma(\omega \rightarrow 2\pi)/\Gamma(\omega \rightarrow 3\pi)$ is 0.7% at the 95% confidence level. (Diss. Abstr. Int., B)

Particle Interactions and Properties (Theoretical)

Refer also to abstracts: 46126 and 46259.

General

Refer also to abstracts: 46691 and 46791.

46497 (IFVE-STF-71-36) ANALYTICAL PROPERTIES OF AMPLITUDES AND DIFFERENTIAL CROSS-SECTIONS FOR MULTIPARTICLE PRODUCTION PROCESSES OVER TWO ANGULAR VARIABLES. Ezhele, V. V. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoy Energii SSSR, Serpukhov. Institut Fiziki Vysokikh Energii). 1971. 11p. (In Russian). Dep. NTIS (U. S. Sales Only).

The amplitude of multiparticle production processes $T(s, n, \xi_1)$ and twice differential cross sections ($d^2\sigma_{\text{inel}}/d\Omega$) ($s, \vec{n}$) where n is a unit vector in the direction of the momentum of colliding particles considered as a function on a sphere $S^2(\vec{n}^2 = 1)$ are considered. It has been proved that both functions holomorphically continue into some domain on the complex sphere ($z_1^2 + z_2^2 + z_3^2 = 1$). It has also been proved that ($d\sigma_{\text{inel}}/d \cos \theta$) ($s, \cos \theta$) analytically continues in the Lehmann ellipse. (auth)

46498 SPIN SPECTRUM OF AN UNSTABLE PARTICLE. Fleming, Gordon N. (Pennsylvania State Univ., University Park). J. Math. Phys. (N. Y.); 13: No. 5, 626-37(May 1972).

A state vector (vector-valued distribution) of an unstable particle can be a 3-momentum eigenvector in at most one inertial frame as a consequence of the necessary width of the energy spectrum. This frame dependence of momentum eigenvectors of unstable particles is investigated and it is demonstrated that: (i) It is compatible with the principle of relativity. (ii) It leads to a distinction between two ways of defining the spin of the particle which are equivalent for stable particles. (iii) One definition, called kinematical spin, yields a precise value, and is determined by considerations of detailed balancing and other means of counting the degeneracy of the momentum eigenvectors. (iv) The second definition, called dynamical spin, need not yield a sharp value, is

determined by the dynamics of the decay process, and is measured by observing angular distributions of decay products. It is shown that an unstable particle with vanishing kinematical spin (pion) may have a small admixture of nonisotropic angular distribution of unpolarized decay products in the rest frame of the decay products. The order of magnitude of the effect is estimated and the coupling between the mass and spin spectrum is studied in the case of local interactions without derivative coupling. (auth)

46499 ONE-DIMENSIONAL CAUSAL THEORY OF TACHYONS. Antippa, A. F. (Univ. of Quebec, Trois-Rivieres). *Nuovo Cim.*; 10A: No. 3, 389-406(1 Aug 1972).

A one-dimensional theory of tachyons, which is noninvariant under space inversion and is based on extending the Lorentz transformations to inertial coordinate systems having relative velocities greater than that of light, is given. It is shown that the theory is fully causal and exhibits a reciprocity between tachyons and bradyons according to which a tachyon with velocity $\beta > 1$ behaves as a bradyon with velocity $1/\beta < 1$ and with the space and time components of its "two-vectors" interchanged. (auth)

46500 KINEMATICALLY CORRECTED CORRELATION FUNCTIONS FOR INCLUSIVE REACTIONS. Yaes, R. J. (Johannes-Gutenberg-Univ., Mainz). *Lett. Nuovo Cim.*; 4: No. 13, 611-15(29 Jul 1972).

A different set of correlation functions is defined in a phase-space model for inclusive interactions, which allows the dynamic effects to be separated from the kinematic effects. Sum rules for the inclusive cross sections are obtained. (L.B.S.)

46501 GRAVITATIONAL CORRECTION TO PCAC. Delbourgo, R.; Salam, A. (Imperial Coll. of Science and Tech., London). *Phys. Lett.*; 40B: No. 3, 381-2(10 Jul 1972).

A simple-minded perturbation calculation to order G (the Newtonian gravitational constant) of the two-graviton coupling to a pseudoscalar (P) or axial (A) current via a fermion loop gives an anomalous contribution which can be added to the Adler term in the form $\partial_\alpha A_\alpha = 2m P + e^2 \epsilon_{\kappa\lambda\mu\nu} F_{\kappa\lambda} F_{\mu\nu} / 16\pi^2 + \epsilon_{\kappa\lambda\mu\nu} R_{\kappa\lambda\rho\sigma} R_{\mu\rho\sigma} / 768\pi^2$. The anomalous terms can be interpreted as arising from an infinite-mass regulating loop. (auth)

46502 GENERAL PROPERTIES OF CORRELATION FUNCTIONS FOR INCLUSIVE PROCESSES. Ballestrero, A.; Nulman, R.; Predazzi, E. (Univ., Turin. Istituto Nazionale di Fisica Nucleare, Turin). *Nuovo Cim.*; 10A: No. 2, 311-24(21 Jul 1972).

General properties of correlation functions for inclusive cross sections are derived from sum rules obtained earlier. In particular, a recursion formula connecting correlation functions of contiguous orders is given which expresses the kinematical constraints of energy-momentum conservation. It follows that all correlation functions are nonzero. Zero and first moments of correlation functions are then given in terms of the multiplicities. (auth)

46503 ALGEBRAIC DERIVATION OF THE GENERALIZED TALMI EXPANSION. Domany, E.; Goshen, S.; Katriel, J. (Nuclear Research Center-Negev, Beer-Sheva, Israel). *Lett. Nuovo Cim.*; 4: No. 9, 389-90(1 Jul 1972).

The generalized Talmi expansion, in which the product of two harmonic-oscillator wave functions is expanded in terms of relative and generalized center-of-mass dependent functions, is derived by a strictly algebraic treatment. Use is made of the fact that transition to the relative and center-of-mass coordinates is obtained by a rotation in the x_1 - x_2 plane, and that the well parameter is affected by scaling of the corresponding coordinate. (L.B.S.)

46504 ENERGY-MOMENTUM CONSERVATION IN INCLUSIVE PROCESSES. Yang, T. C. (Univ. of California, San Diego, La Jolla). *Lett. Nuovo Cim.*; 4: No. 9, 352-6(1 Jul 1972).

The implications of energy-momentum conservation in single-particle and two-particle inclusive processes for correlation functions and cross sections are discussed. (L.B.S.)

46505 ESTIMATE OF THE QUASIPOTENTIAL PHASE SHIFT. Arkhipov, A. A. (Inst. of High Energy Physics, Serpukhov, USSR). *Teor. Mat. Fiz.*; 11: No. 3, 337-43(Jun 1972). (In Russian).

The quasipotential phase shift is calculated in the first and second orders with respect to the quasipotential. (auth)

46506 RELATIVISTIC WAVE FUNCTIONS FOR COMPOSITE PARTICLES. Peres, A. (Technion-Israel Inst. of Tech., Haifa). *Nuovo Cim.*; 10A: No. 2, 230-4(21 Jul 1972).

Clusters of free spinless particles are considered. It is shown that the relativistic wave functions of such clusters cannot be factorized into external and internal wave functions. However,

an approximate factorization is still possible, provided that the invariant mass of the cluster has a narrow distribution around some mean value. In that case the internal wave function of a cluster moving with relativistic speed is obtained from that of the same cluster at rest by means of a Lorentz contraction. (auth)

46507 IRREDUCIBLE REPRESENTATIONS OF PARA-BOSE OPERATORS. Alabiso, C.; Duimio, F. (Univ., Parma, Italy. Istituto Nazionale di Fisica Nucleare, Milan). *Nuovo Cim.*; 10A: No. 3, 499-512(1 Aug 1972).

A general procedure is given in order to evaluate the matrix elements of the para-Bose operators of any order. As a consequence, a class of irreducible representations of the $Sp_{2n,R}$ algebra is constructed. (auth)

46508 PHYSICS IN THE TWENTIETH CENTURY: SELECTED ESSAYS. Weisskopf, Victor F. Cambridge, Mass.; Massachusetts Institute of Technology Press (1972). 380p. \$7.95.

This collection of essays includes quality and quantity in quantum physics, the quantum ladder, Niels Bohr the quantum and the world, physics in the twentieth century, recent developments in the theory of the electron, the compound nucleus, problems of nuclear structure, quantum theory and elementary particles, nuclear models, the visual appearance of rapidly moving objects, light interactions with matter, the fall of parity, and the role of symmetry in nuclear, atomic, and complex structures. Also discussed are man and nature, Marie Curie and modern science, science and ethics, and the significance of science. (J.F.P.)

46509 PARTICLE PHYSICS. Lagarrigue, M. (Univ. Paris, Orsay, France). *Bull. Union Physiciens*; 66: No. 544, 783-853(Apr 1972). (In French).

The text of a Maitrise de Physique course given at Orsay is reproduced. The titles of the chapters are as follows: generalities about particles; the π meson; isotopic spin-strangeness-hypercharge; parity transformations and charge conjugations; weak interaction. (auth)

Electromagnetic

Refer also to abstracts: 46433, 46678, 46693, 46694, 46698, 46704, 46742, 46757, 46767, and 46867.

46510 (CALT-767-P-3-22) PERTURBATION TREATMENT OF HIGH ENERGY ELECTRON DIFFRACTION FROM IMPERFECT CRYSTALS. Lewis, Alan L.; Villagrana, Robert E. (California Inst. of Tech., Pasadena). 1970. 31p. Dep. NTIS.

A modified Bloch wave expansion is used to develop a theory of high energy electron diffraction from imperfect crystals. To compute these new Bloch waves one must solve a linear hyperbolic system in n unknowns. Scattering among the Bloch waves is controlled by the matrix elements of the perturbing potential, and various approximations to this scattering are discussed. The hyperbolic system is transformed to normal form; in making this transformation, the unknown functions become the plane wave amplitudes of the Darwin representation. The normal form reveals the region of determinacy of the system: a cone generated by the diffracted beams. The contraction of this cone to a line (the column approximation) is discussed in terms of the Bloch wave scattering. (auth)

46511 (CONF-710816-9) BEFORE APPLYING THE DISPERSION THEORY TO THE PHOTOPRODUCTION REACTIONS. Blazek, M. (Ceskoslovenska Akademie Ved, Prague. Fysikalni Ustav). [1971]. 21p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

The application of the dispersion relation techniques assumes the knowledge of the asymptotics together with the analytic properties of the amplitudes under consideration. A method is suggested which allows one to determine the locus where a partial wave amplitude for the photoreactions possesses an absorptive part in the complex s -plane. The present analysis yields the necessity of analytic continuation due to the appearance of the continua which give rise to the areas where a partial wave amplitude is not defined. Though this effect is relatively weak in the photoproduction of pions on nucleons, it is quite strong in the photoproduction of vector mesons. The complications encountered might be avoided by the (stable or unstable) particle exchanges. Then the necessity of the continuation does not appear. Suggestions for evaluation of the dispersion integrals are presented and some implications for full amplitude are mentioned. (auth)

46512 (CONF-710816-16) π^0 ELECTROPRODUCTION AT THE FIRST RESONANCE. Brasse, F. W.; Chazelas, E.; Fehrenbach, W.; (and others) (Deutsches Elektronen-Synchrotron, Ham-

burg (West Germany)). 1971. 29p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

Recent results on electroproduction of π^0 mesons in the region of the first resonance at four momentum transfers $q^2 = 0.6 \text{ GeV}^2$ and $q^2 = 1 \text{ GeV}^2$ are discussed and analyzed in terms of multipole amplitudes. Assuming that the angular distributions are dominated by the magnetic dipole M_{1+} , the square of the magnetic dipole amplitude and its interference terms with the multipoles E_{0+} , S_{0+} , E_{1+} , S_{1+} and M_{1-} are determined. The results are compared with theoretical models. The dependence of $|M_{1+}|^2$ on the invariant mass W at different q^2 is discussed by comparison with a standard Breit Wigner formula. (auth)

46513 (CONF-710816-17) IS THE PION FORM FACTOR DETERMINED BY ELECTROPRODUCTION EXPERIMENTS? Dombey, Norman; Read, B. J. (Sussex Univ., Brighton (England)). Aug 1971. 8p. Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

Experiments on pion electroproduction in the first resonance region which measure the pion form factor are reinterpreted using a very simple model in which the axial vector form factor appears explicitly. A significant correlation between F_π and G_A is found. It is concluded that the pion form factor has not yet been determined from electroproduction experiments since G_A is largely unknown. It is noted that a careful set of coincidence measurements in the region from threshold through the first resonance could determine both G_A and F_π . (J.F.P.)

46514 (CTH-ITP-71-30) NUCLEON FORM FACTORS IN A FIELD THEORY ON A HOMOGENEOUS SPACE OF THE POINCARÉ GROUP. Brink, Lars; Marnelius, Robert (Chalmers Tekniska Hogskola, Goteborg (Sweden). Institutionen for Teoretisk Fysik). 12 Nov 1971. 41p. Dep. NTIS (U. S. Sales Only).

The nucleon form factors are calculated to lowest order in a new field theory on an 8-dimensional homogeneous space of the Poincaré group. The simplest diagram does give both an electric and a magnetic form factor and a relation between them. It seems however, difficult to reproduce the experimental data, as e.g. any anomalous magnetic moment does not appear in a natural way. (auth) (Sweden)

46515 (DAMTP-71/24) DEEP INELASTIC ELECTROPRODUCTION AND RELATED PROCESSES. Landshoff, P. V.; Polkinghorne, J. C. (Cambridge Univ. (England). Dept. of Applied Mathematics and Theoretical Physics). 1971. 24p. (CONF-710816-18). Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

The parton approach, with the partons identified as quarks, is combined with a dual model to describe deep inelastic electron and neutrino scattering. This model has the important feature that it allows a complete calculation of the nondiffractive contributions to the structure functions, essentially without free parameters. The calculated difference $F_2^p(\omega) - F_2^n(\omega)$, which has no diffractive contribution, is in fair agreement with experiment. The calculated form of the nondiffractive contribution to $F_2^p(\omega)$ agrees well with the data for F_2^p for small ω . For large ω the difference between this calculated form and the data is quite small. If this difference is identified with the diffractive contribution, the neutrino structure functions are determined. These are in agreement with the rather preliminary data. The partons are introduced through a nonperturbative field-theory formalism. This reproduces most of the results of the more phenomenological approaches, except that in the case of heavy-muon-pair production it indicates that an extra Pomeron-exchange contribution is important. When this is allowed for, good agreement with the data for this process is obtained, including the "shoulder" in the differential cross section. The picture is extended to purely hadronic processes, and leads to the results of the additive quark model as an approximation, even though the structure of the hadrons is supposed to be considerably more complicated than in the naive quark model. The Wu-Yang model and s-channel helicity conservation also emerge as natural consequences. (auth)

46516 (DESY-72/21) TESTING TRIPLET MODELS. Suura, H. (Deutsches Elektronen-Synchrotron, Hamburg (West Germany); Walsh, T. F. (Hamburg Univ. (West Germany). Institut fuer Theoretische Physik (II); Young, Bing-Lin (Ames Lab., Iowa). May 1972. 11p. Dep. NTIS.

Radiative decays $\pi^0 \rightarrow 2\gamma$, $\omega \rightarrow \pi^0 + \gamma$, $\rho \rightarrow \pi^0 + \gamma$, $\eta \rightarrow 2\gamma$, $\epsilon(750) \rightarrow 2\gamma$, and $\eta^1 \rightarrow 2\gamma$ and $e^+ + e^-$ and $e^- + e^-$ interactions are used to distinguish the Han-Nambu and Gell-Mann-Zweig triplet quark models. (J.F.P.)

46517 (DESY-72/23) DIFFRACTION-DISSOCIATION MODEL FOR HIGH-ENERGY ELECTROPRODUCTION. Kramer, G.; Quinn, H. R. (Deutsches Elektronen-Synchrotron, Hamburg (West Germany)). May 1972. 54p. Dep. NTIS (U. S. Sales Only).

A model for two-pion electroproduction in the rho resonance region is constructed. The production proceeds by diffraction-dissociation of the virtual photon into a pion pair which interacts strongly in the p-wave state. The model is compared with recent experimental data from Cornell, DESY and SLAC. (auth)

46518 (ITEF-848) CONCERNING THE TWO-PHOTON MECHANISM OF PRODUCTION DURING COLLISION OF CHARGED PARTICLES. Terent'ev, M. V. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoi Energii SSSR, Moscow. Institut Teoreticheskoi i Eksperimental'noi Fiziki). 1971. 23p. (In Russian). Dep. NTIS (U. S. Sales Only).

The simple and invariant method of calculation with logarithmic precision of the asymptotic value of differential production cross section for charged particle collisions is formulated. The two-photon mechanism of production is considered. The main result is contained in given formulas. A number of examples is considered. The main features of the angular distributions and some numerical estimates are discussed considering the reaction of meson production as an example. (auth)

46519 (JINR-P2-6333) REACTION $\gamma + \gamma \rightarrow \pi + \pi$ AND ELECTROMAGNETIC DIFFERENCE FOR PION MASSES. Isaev, P. S.; Khleskov, V. I. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 26p. (In Russian). Dep. NTIS (U. S. Sales Only).

The reaction $\gamma + \gamma \rightarrow \pi + \pi$ was studied in the work by dispersion relations. The cross sections of the process were calculated for two parametrizations of s wave $\pi\pi$ phase shift. The dependence of cross sections on the threshold and high energy phase shift behavior and on inelasticity was investigated. The cross sections of the processes $e + e \rightarrow e + e + \pi + \pi$ were calculated. The pion electromagnetic mass was determined from dependence of the $\gamma + \gamma \rightarrow \pi^+ + \pi^-$ amplitude on photons "masses." (auth)

46520 (JINR-P2-6438) ELECTROMAGNETIC EFFECTS IN NUCLEON POLARIZATION AT MESON-NUCLEON SCATTERING. Vanzha, A. P.; Lapidus, L. I.; Tarasov, A. V. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Nuclear Problems). 1972. 24p. (In Russian). Dep. NTIS (U. S. Sales Only).

It is shown, that at high energy and small t the contribution of electromagnetic interaction of particles with spin is of importance for polarization effects in elastic (and some inelastic) processes of hadrons. The investigation of polarization $P_0(t)$ and also Wolfenstein parameters A and R at small t gives the possibilities for experimental determination of real and imaginary parts of hadronic invariant amplitudes A and B. The electromagnetic contribution for A and B is obtained. When the amplitude A' is involved, the expressions for the cross section, polarization P_0 , R and A are given. The expression for A'_{em} is given. When both hadronic and electromagnetic amplitudes are considered simultaneously the question of relative phases of amplitudes becomes of interest. The estimations for t_m give relativistic generalization of Schwinger slow neutron scattering results. The Coulomb interaction, which is dominant at $|t| < |t_c|$, where t_c is given strongly reduces polarization effects when compared with the case of meson-neutron scattering. Electromagnetic effects give essential deviations at small t. For the numerical estimation of the scale of electromagnetic effects, approximate hadronic amplitudes were taken from Rarita et al. The results, which were obtained in this way, are illustrated. (auth)

46521 (LBL-773) FINAL STATE HADRON DISTRIBUTIONS AND GENERALIZED SCALING LAWS IN DEEP-INELASTIC ELECTROPRODUCTION PROCESSES. Pak, Namik K. (California Univ., Berkeley. Lawrence Berkeley Lab.). Apr 1972. Contract W-7405-eng-48. 93p. Dep. NTIS.

Thesis.

Processes of the type "electron + nucleon $\rightarrow$ electron + hadron (pion) + Anything" at high energy and high electron momentum transfer are discussed, via the processes "virtual photon + nucleon $\rightarrow$ hadron (pion) + Anything." New general scaling laws are obtained for these processes which contain hadronic scaling laws (limiting fragmentation, and pionization) and Bjorken scaling as special cases. The derivation is model independent and is based on the expansion of the forward three-to-three amplitude, "virtual photon + nucleon + hadron $\rightarrow$ virtual photon + nucleon + hadron," into 0(1,2) harmonics. In getting the results it is assumed that the residue of pomeronchukon trajectory factorizes, and Bjorken scaling of the ordinary deep inelastic lepton scattering processes is used as an experimental fact. It is found that multiplicity of the produced particle grows with the energy logarithmically both

in the Regge and Bjorken scaling limits. Finally on the basis of PCAC hypothesis and the soft pion theorems of the current algebra the cross section of one soft pion production in deep inelastic electron scattering processes is calculated. (auth)

46522 (N-72-17763) CHARGED-PARTICLE MOTION IN THE VICINITY OF A NEUTRAL PLANE. Stern, David (National Aeronautics and Space Administration, Greenbelt, Md. Goddard Space Flight Center). Feb 1972. 30p. (NASA-TM-X-65821; X-641-72-22). NTIS.

The motion of a charged particle in a 2-dimensional neutral sheet with linear magnetic field variation is analyzed by Hamiltonian methods. From the basic Hamiltonian formulation, results of Sonnerup are recovered and given intuitive interpretation. The transformed Hamiltonian is used to derive the correct frequency of oscillation and serves as the basis of analytical treatment of perturbed versions of the motion, e.g. with electric field added. For 2-dimensional fields with slightly differing configurations—in particular, with added small field component orthogonal to the sheet, with an X-type null or with a chain of null points—two alternative methods are developed, reducing the problem either to motion in a 2-dimensional potential or to a pair of coupled oscillators. (auth) (STAR)

46523 (NYO-4076-17) INELASTIC e-p SCATTERING AND THE POLARIZATION CONTRIBUTION TO H HYPERFINE STRUCTURE. Jensen, D.; Kovess-Domokos, S.; Schonberg, E. (Johns Hopkins Univ., Baltimore, Md. Dept. of Physics). Jul 1971. Contract AT(30-1)-4076. 13p. Dep. NTIS.

The structure functions obtained from a specific resonance model of inelastic e-p scattering are used to evaluate the polarizability correction to the ground-state hyperfine splitting of hydrogen. By requiring that the spin-flip amplitude for real Compton scattering satisfy the Drell-Hearn sum rule, one can fix the only free parameter in the model. It is found that the polarizability corrections are approximately 1 ppm, in good agreement with the existing bounds on the uncertainty of Q. E. D. calculations of hyperfine structure. (auth)

46524 (NYO-4076-18) LIGHTLIKE QUARK MODEL AND CURRENT DENSITIES. Domokos, G. (Johns Hopkins Univ., Baltimore, Md. Dept. of Physics). Jul 1971. Contract AT(30-1)-4076. 23p. Dep. NTIS.

A field theory of quarks, with quark states being defined on space-time surfaces with lightlike normals, is formulated. The general structure of the field theory is discussed. It is found that single particle observables, and current densities in particular, have approximately a free field structure if the ground state is strongly degenerate. The structure functions of the nucleon in inelastic electron scattering exhibit a parton-like structure in any reference frame. (auth)

46525 (ORO-3992-80) LEPTONIC DECAY OF VECTOR MESONS IN AN ALGEBRAIC MODEL FOR HADRONS. Boehm, A. (Texas Univ., Austin. Center for Particle Theory). 4 Apr 1972. Contract AT(40-1)-3992. 25p. (CPT-134). Dep. NTIS.

The decay rates for $V \rightarrow e\bar{e}$ are calculated using an octet of transition operators that has been introduced previously in the framework of an algebraic model. The results are compared with experimental data and restrictions on the symmetry breaking relations for these transition operators are obtained, which are shown to be the analog of other well known symmetry breaking relations. (auth)

46526 (TPJU-14/71) COMPARISON OF THE ANGULAR DECAY DISTRIBUTIONS IN THREE QUARK MODELS OF PHOTOPRODUCTION. Gorczyca, Barbara (Jagellonian Univ., Krakow (Poland). Dept. of Theoretical Physics). Apr 1971. 11p. (CONF-710816-11). Dep. NTIS (U. S. Sales Only).

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

The constraints for angular decay distributions of resonances photoproduced in the reactions $\gamma B \rightarrow PB^*$, $\gamma B \rightarrow VB$, and $\gamma B \rightarrow VB^*$ in the "excited quark" model and "interacting quarks" model, where the quark structure is assumed only for baryons, are derived and compared with the corresponding predictions of the usual additive quark model of photoproduction including vector meson dominance and the additivity assumption. All the results of the "interacting quarks" model can be derived from the "excited quark" model and all the results of the "excited quark" model can be derived from the additive quark model. (auth)

46527 $\rho^0 - \omega$ INTERFERENCE IN THE REACTIONS $\gamma \rightarrow \pi^+\pi^-\Delta$. Achasov, N. N.; Shestakov, G. N. (Inst. of Mathematics, Novosibirsk, USSR). JETP Lett. (USSR) (Engl. Transl.); 15: No. 7, 302-5 (5 Apr 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 7, 427-30 (5 Apr 1972).

Interactions in which the cross sections for ω production are much larger than those for ρ^0 production are expected to exhibit a considerable enhancement of the $\rho^0 - \omega$ interference. It is shown theoretically that the interference for interactions of the type $\gamma + N \rightarrow \pi^+ + \pi^- + \Delta$ is enhanced by a factor of 9 over that for $\gamma + A \rightarrow \pi^+ + \pi^- + A$ interactions. The general question of separating the nonresonant background processes in the study of quasi-two-particle interactions is also considered. (R.W.R.)

46528 INFORMATION ON THE PION ELECTROMAGNETIC FORM FACTOR INSIDE ITS ANALYTICITY REGION PROVIDED BY BOUNDS ON ITS MODULUS ON THE CUT ($t \geq 4m^2$). Raz-zillier, I. (Inst. of Physics, Bucharest). Commun. Math. Phys.; 26: No. 2, 121-9 (1972).

A study is made of the mathematical aspects connected with the derivation of the model independent information that can be obtained on the pion electromagnetic form factor $F(t)$ inside the analyticity region (the cut t -plane) from the knowledge of upper and lower bounds of its modulus on the cut $t \geq 4m^2$ using analyticity, reality, and the normalization $F(0) = 1$. It turns out that (in a certain sense) this information depends only on the upper bound, whereas the lower one is irrelevant. (auth)

46529 MESON PHOTOPRODUCTION IN THE BACKWARD DIRECTION. Greenhut, G. K. (Daresbury Nuclear Physics Lab., Eng.). Nucl. Phys.; B44: No. 2, 422-36 (1972).

Two methods are discussed for predicting the cross sections for backward photoproduction of neutral mesons. One is an extension of the broken-SU(3) quark model plus vector dominance to the backward direction and yields cross sections for ρ^0 , ω , and ϕ production. The other uses factorization of Regge residues correlating the Regge fits to $\pi N \rightarrow N\pi$, $\pi N \rightarrow N\rho$, $\pi N \rightarrow N\eta$, and $\gamma N \rightarrow N\pi$ to obtain predictions for the cross sections for ρ^0 and η photoproduction. (auth)

46530 DANGEROUS SCALING LIMIT FOR VIRTUAL COMPTON SCATTERING OFF THE FORWARD DIRECTION. Min-kowski, P. (SIN, Zurich). Nucl. Phys.; B44: No. 2, 390-4 (1972).

Virtual Compton scattering off the forward direction is investigated in the scaling region. An example motivated by vector meson dominance is constructed in accordance with the Jost-Lehmann-Dyson representation which demonstrates that the light cone singularities of the current commutator fail to determine the amplitude if the square mass of one of the virtual photons remains finite. (auth)

46531 SOFT-PHOTON CROSS SECTIONS FOR $\pi^- d \rightarrow n n \gamma$. Picciotto, C. (Univ. of Victoria, B. C.). Can. J. Phys.; 50: No. 8, 798-802 (15 Apr 1972).

A phenomenological calculation of cross sections for the reaction $\pi^- d \rightarrow n n \gamma$ for pion lab energies of 150 to 300 MeV is presented. (auth)

46532 INTERACTION OF AN INTENSE QUASI-MONOCROMATIC ELECTROMAGNETIC BEAM WITH FREE ELECTRONS. Billoir, Pierre (Univ., Paris). Compt. Rend., Ser. B; 274: No. 15, 915-18 (10 Apr 1972). (In French).

The effect of a coherent photon beam with a high photon density on electrons at rest was studied using a simple quantum model. In a first approximation, this interaction can be described in terms of the scattering of the electrons at right angles to the beam direction by using a statistical property of bosons; the effect is quadratic with respect to the number of photons. (France)

46533 DOMINANCE OF ELECTROMAGNETIC EFFECTS IN HIGH ENERGY HADRON AND PHOTON INDUCED REACTIONS. Dar, A. (Technion-Israel Inst. of Tech., Haifa). pp 17-38 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris: Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Electromagnetic interactions in high energy hadron and photon induced reactions are examined. The one-photon exchange contribution to two-body final state reactions and also multibody final state reactions are discussed. The mechanism can be used for studying radiative decay rates of mesons, their elastic and inelastic form factors, and their structure functions and also for determining the phase of strong interaction amplitudes at high energy. Coulomb production in the nuclear field can explain the increase with energy observed in some recently reported inelastic cross sections. The dominance of the mechanism in the forward

direction can be used to measure radiative decay rates of mesons (Primakoff effect), and photon-meson and photon-photon cross sections. Interferences may lead to nonequality between particle and antiparticle cross sections. (France)

46534 K^+ PHOTOPRODUCTION. Capella, A. (Univ. Paris, Orsay, France). pp 289-97 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

K^+ photoproduction was studied with an incident momentum of the 5 to 15 GeV/c. Experimental data obtained exhibit the following features: a backward dip, but no dip in the $-0.4, -0.5$ (GeV)² region, below $2(\text{GeV})^2$, K and K^*-p contributions giving α_{eff} approximately zero, a $d\sigma/dt(K^0)/d\sigma/dt(KA)$ value approximately equal to 1 except in the forward direction. A model which considers only the K^* and K^{**} Regge poles and the K^*-p and $K^{**}-p$ leading cuts is presented. (France)

46535 CONNECTIONS BETWEEN LOW ENERGY AND HIGH ENERGY K^+ PHOTOPRODUCTION. Renard, F. M. (Univ., Montpellier, France). pp 293-307 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Characteristic features of the high energy K^+ photoproduction are shown to appear also in the low energy region (1 to 2 GeV range). The low energy integral is examined, using the finite energy sum rules (FESR), over the imaginary part of the scattering amplitude. FESR integrals were separately calculated for the s-channel part (N^*) and for the u-channel part (Y^*) in the case of the four t-channel helicity amplitudes. Experimental features are shown to be remarkably consistent with the FESR integrals in the low energy region. (France)

46536 QUARK MODEL PREDICTIONS FOR ANGULAR DECAY DISTRIBUTIONS OF PHOTOPRODUCED VECTOR MESONS IN INCLUSIVE PROCESSES. Gorczyca, B. (Jagellonian Univ., Krakow). Lett. Nuovo Cim.; 4: No. 8, 291-4(24 Jun 1972).

The quark model predictions on the decay distribution of vector mesons produced by polarized photons in inclusive interactions are presented. Relations for statistical tensors are given, which can be checked experimentally using linearly and circularly polarized photons. (L.B.S.)

46537 FIXED-POLE AND VIRTUAL COMPTON AMPLITUDE. Ezawa, Z. F. (Tokyo Univ.). Lett. Nuovo Cim.; 4: No. 8, 315-19 (24 Jun 1972).

Fixed poles in the Compton scattering amplitude are investigated in a field theory with the parton description for the virtual Compton scattering in the Bjorken limit. (L.B.S.)

46538 HIGH-ENERGY ELECTROMAGNETIC PROCESSES IN CONDENSED MEDIA. Ter-Mikaelyan, M. L. Interscience Tracts on Physics and Astronomy, No. 29. New York; John Wiley and Sons, Inc. (1972). 466p. \$27.50.

Coherent problems occurring in electromagnetic processes at high energies are examined from a unified viewpoint. The pseudo-photon method, coherent bremsstrahlung in a crystal, radiation processes at high energies in amorphous media, transition radiation, and radiation of particles passing uniformly through heterogeneous media are treated. (J.F.P.)

46539 DUALY CHARGED PARTICLES AND DUAL INVARIANCE. Schiff, H. (Univ. of Alberta, Edmonton). Can. J. Phys.; 50: No. 11, 1062-7(1 Jun 1972).

Dual charges are introduced which differ essentially from the conventional (Dirac) magnetic monopoles. A natural vehicle is shown to be a dual invariant Lagrangian to represent observable charges; the dual invariance can be broken in a simple way by a class of dually charged objects which exhibit no electromagnetic interaction with observable dually charged particles. A triplet formulation can also be obtained and may be of interest in providing a basic description of bosons and fermions. (auth) (Canada)

46540 e^+e^- ANNIHILATION INTO HADRONS: RESONANCES, REGGE POLES, FIXED POLES, SCALING, AND DUALITY. Layssac, J.; Renard, F. M. (Department de Physique Mathematique, Montpellier, France). Nuovo Cim.; 10A: No. 3, 407-44 (1 Aug 1972).

The behavior of the cross section for $e^+e^- \rightarrow \pi^0 + \text{hadrons}$ is discussed through the following properties of the structure functions $W_1(q^2, v)$: low-energy theorem at $q^2 = 0$, vector-meson

structures around $q^2 = m_\rho^2$, quasi-two-body contributions $\pi + R$ and $R + R'$ in the (1 to 2) GeV region, scaling Regge poles and fixed poles for high q^2 . Duality relations are proposed in order to connect these different properties. The shapes of the differential and total cross sections are discussed with respect to the preliminary experimental results. (auth)

46541 INDUCED RADIATION FROM NEUTRONS MOVING THROUGH A MEDIUM IN A MAGNETIC FIELD. Kul'kin, A. G.; Pavlenko, Yu. G. (Moscow State Univ.). Izv. Vyssh. Ucheb. Zaved., Fiz.; No. 12, 83-7(1971). (In Russian).

The conditions for the formation of a negative coefficient of reabsorption were investigated for the system neutron beam-medium in the presence of a magnetic field. A coherent generation mechanism of electromagnetic waves in a broad range of frequencies was realized. (K.S.W.)

46542 ARE VECTOR POTENTIALS MEASURABLE QUANTITIES IN ELECTROMAGNETIC THEORY? Vigier, J. P.; Marcilhacy, G. (Institut Henri Poincare, Paris). Lett. Nuovo Cim.; 4: No. 13, 616-18(29 Jul 1972).

Starting from a realistic description of electromagnetic sources, it is shown (a) that one can define the measurable vector potentials A^μ even in Maxwell's theory, and (b) that these A^μ are just the limit for $m_\gamma \rightarrow 0$ of the A^μ of the De Broglie-Proca theory (L.B.S.)

46543 ACOPLANARITY AND RADIATIVE CORRECTIONS IN BHABHA SCATTERING. Calva-Tellez, E. (College de France, Paris); Kessler, P.; Parisi, J. Lett. Nuovo Cim.; 4: No. 13, 619-23(29 Jul 1972).

A recent experiment performed at the ADONE colliding-beam facility showed the occurrence of an appreciable number of acoplanar events in Bhabha scattering ($e^+e^- \rightarrow e^+e^-$). Acoplanar was defined as those events where the angle $|\phi|$ between the two planes containing the beam axis and, respectively, the outgoing electron or positron was larger than 5° . These events were analyzed as being due to radiative corrections. Two distinct calculations are made in order to analyze the data in a more adequate way: an exact one, and a small-angle approximation, using the method of quasi-real processes. The acoplanarity distribution is shown to agree well with the experimental data. (L.B.S.)

46544 NONRELATIVISTIC HARMONIC-OSCILLATOR QUARK MODEL AND THE RATIO OF LONGITUDINAL TO TRANSVERSE CROSS-SECTIONS IN RESONANCE ELECTROPRODUCTION. Abdullah, T. (Dept. of Theoretical Physics, Oxford). Lett. Nuovo Cim.; 4: No. 13, 577-80(29 Jul 1972).

It is shown that the predictions of the nonrelativistic harmonic-oscillator quark model are essentially consistent with the results for the ratio of longitudinal to transverse cross sections in resonance electroproduction. It is pointed out that the predictions of the model are more sensitive to the choice of frame used for the physical interpretation of the matrix elements calculated than to the choice of parameters. The Breit frame is shown to be the most suitable frame for the nonrelativistic calculation. (L.B.S.)

46545 PHOTOPRODUCTION OF HYPERNUCLEI AND CURRENT ALGEBRA. Sidorov, A. A. (Tomsk Polytechnical Inst., USSR). Izv. Vyssh. Ucheb. Zaved., Fiz.; No. 12, 53-5(1971). (In Russian).

The differential cross section of the reaction $\gamma + {}^3\text{He} \rightarrow K^+ + \Lambda^3\text{H}$ was calculated within the limits of "soft" kaons, using the algebra of currents. In this reaction the nucleus ${}^3\text{He}$ and the hypernucleus $\Lambda^3\text{H}$ are considered as elementary particles. The amplitude of the photo-formation process for hypernuclei could be combined with the matrix element of a weak hadron current between the nucleus-hypernucleus states so that they could be expressed by corresponding nuclear form factors. The coupling constant of the axial current which characterizes β decay of the $\Lambda^3\text{H}$ may be equal to -0.67 , as in the case of the free Λ -particle. (tr-auth)

46546 EFFECT OF THE COLLECTIVE INTERACTION OF PARTICLES ON THE LONGITUDINAL DIMENSIONS OF AN ELECTRON BUNDLE IN AN ACCUMULATOR. Grigor'ev, Yu. N.; Grishaev, I. A.; Naugol'nyi, N. N. (Physical-Technical Inst., Kharkov). Ukr. Fiz. Zh. (Russ. Ed.); 16: No. 10, 1729-32(Oct 1971). (In Russian).

The effect of a force of collective interaction on the phase motion of relativistic electrons in a ring accumulator was studied. With an increase in the number of accumulated particles the decrement of synchrotron oscillations decreased, and the longitudinal dimensions of the bundle increased. The results of the theory are compared with experimental data on the latter phenomenon. (tr-auth)

46547 MOTION OF DIRAC PARTICLES POSSESSING AN ANOMALOUS MAGNETIC MOMENT IN A CENTRAL SYM-

METRIC ELECTRIC FIELD. Ternov, I. M.; Bagrov, V. G.; Bozrikov, P. V. (Tomsk Polytechnic Inst., USSR). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 38-45(1971). (In Russian).

States of a relativistic electron, having an anomalous magnetic moment, and moving in a central-symmetric field, were classified by a method which was very close to nonrelativistic classification. Spin quantum numbers are clearly distinguished. The motion of a neutron in a central field was also studied. New precise solutions for the Dirac-Pauli equations were found. A classical interpretation of the results is offered. (tr-auth)

46548 SPIN EFFECTS IN THEORY OF SYNCHROTRON RADIATION DUE TO MOTION OF AN ELECTRON IN A MEDIUM. Kul'kin, A. G.; Pavlenko, Yu. G. (Moscow State Univ.). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 7-15(1971). (In Russian).

The presence of a medium essentially changes the known spin effects during ultra-luminous motion of an electron in a magnetic field. In particular, the ground state became unstable and the first excited state lost its metastable character. An expression was obtained for the lifetime of the ground and first excited states. The intensity of radiation of the two chief components of polarization was investigated as a function of spin orientation. (tr-auth)

46549 ELECTROPRODUCTION OF π MESONS NEAR THRESHOLD AND CURRENT ALGEBRA. Kulish, Yu. V. (Physical-Technical Inst., Kharkov). *Ukr. Fiz. Zh. (Russ. Ed.)*; 16: No. 9, 1437-43(Sep 1971). (In Russian).

The amplitudes of electroproduction of π -mesons near a threshold having no kinematic singularities are obtained by means of current algebra and PCAC. The sensitivity of the electroproduction cross section to the nucleon axial-vector form factor is indicated. (auth)

46550 PECULIARITIES IN INTERACTION OF HIGH-ENERGY ELECTRONS AND POSITRONS WITH CRYSTALS. Grishaev, I. A.; Efimov, V. P.; Kasulov, V. I.; (and others) (Physical-Technical Inst., Kharkov). *Ukr. Fiz. Zh. (Russ. Ed.)*; 16: No. 9, 1548-50(Sep 1971). (In Russian).

The total yield of retarding photons was investigated as a function of crystal orientation during passage of electrons and protons with energies of 1 GeV through crystals of Si and Nb. The crystals were mounted so that the incident beam of electrons (positrons) penetrated in the direction of the axis (110) in the case of Si and (100) in the case of Nb. The crystal could be rotated around two perpendicular axes normal to the direction of the beam with a precision of 2×10^{-4} radians. The distinguishing parameter in the dependence of photon yield during the passage of electrons and protons through crystals was the width of the minimum in the region of small angles, which in the case of Si was 1.5 times smaller for positrons than for electrons; in the case of Nb was 1.6 times smaller for positrons than for electrons. (tr-auth)

46551 INDUCED EMISSION FROM AN ELECTRON IN A MAGNETIC FIELD AT THE LEVEL $n = 1$. Ternov, I. M.; Khalilov, V. R.; Zhuravlev, A. F.; Chizkov, G. A. (Moscow State Univ.). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 12, 97-100(1971). (In Russian).

The influence of an external magnetic field wave on the polarization of an electron in a magnetic field was examined. Allowing for an anomalous magnetic moment of the electron, it is possible to obtain complete polarization of an electron beam in the state $n = 0$. (tr-auth)

46552 e^+e^- ANNIHILATION INTO TWO HADRONS IN THE ENERGY INTERVAL 1400 TO 2400 MeV. Alles-Borelli, V.; Bernardini, M.; Bollini, D.; (and others) (CERN, Geneva.—Univ., Bologna. INFN, Bologna. INFN, Frascati, Italy). *Phys. Lett.*; 40B: No. 3, 433-6(10 Jul 1972).

The proof is given for the existence of the reaction $e^+e^- \rightarrow h^+h^-$ in the energy range 1400 to 2400 MeV, and its energy dependence is compared with that of $e^+e^- \rightarrow e^+e^-$, in the same experimental conditions of observation. The exponent of the s -dependence of the ratio $\alpha = (e^+e^- \rightarrow h^+h^-)/(e^+e^- \rightarrow e^+e^-)$ is measured to be $n = 2.08 \pm 0.45$, in the s -range (1.96-5.76) GeV², on the basis of 51 $e^+e^- \rightarrow h^+h^-$ events and 8918 $e^+e^- \rightarrow e^+e^-$ events observed. (auth)

46553 DRELL-HEARN SUM RULE AND THE LEPTON MAGNETIC MOMENT IN THE WEINBERG MODEL OF WEAK AND ELECTROMAGNETIC INTERACTIONS. Altarelli, G. (Univ., Rome); Cabibbo, N.; Maiani, L. *Phys. Lett.*; 40B: No. 3, 415-19(10 Jul 1972).

It is verified that the fourth-order Compton amplitude for the scattering of a photon off a charged lepton is finite in Weinberg's model of weak and electromagnetic interactions, and obeys the Drell-Hearn sum rule. The second-order weak corrections to

the anomalous magnetic moment of a charged lepton in the same model is computed. For the case of the muon, the following bounds are found: $+0.2 \times 10^{-8} \leq \kappa \leq +0.56 \times 10^{-8}$. (auth)

46554 NUCLEON FORM FACTORS IN A FIELD THEORY ON A HOMOGENEOUS SPACE OF THE POINCARÉ GROUPS. Brink, L.; Marnelius, R. (Inst. of Theoretical Physics, Goteborg). *Nuovo Cim.*; 10A: No. 3, 533-59(1 Aug 1972).

The nucleon form factors are calculated to lowest order in a new field theory on an 8-dimensional homogeneous space of the Poincaré group. The simplest diagram does give both an electric and a magnetic form factor and a relation between them. It seems, however, difficult to reproduce the experimental data, as, e.g., any anomalous magnetic moment does not appear in a natural way. (auth)

46555 PHOTOPRODUCTION OF RHO-MESONS ON DEUTERONS. Wilk, G. (International Centre for Theoretical Physics, Trieste, Italy). *Nuovo Cim.*; 10A: No. 2, 207-29(21 Jul 1972).

A formalism for $V_1 + D \rightarrow V_2 + D$, where $V_{1,2}$ are 1^- particles, is presented using the Glauber-model approach. The scattering amplitude is built up from $V_1 + N \rightarrow V_2 + N$ amplitudes treating the nucleons as spinless particles. Spin and D-state in the deuteron is taken into account in the usual manner. The formalism is applied to describe the photoproduction of rho-mesons on deuterons and comparison with experimental data is made. Some predictions on the density matrix elements of outgoing rho in terms of the rho-nucleon amplitude are given. It is shown that exact measurements of these can give information about spin structure of the rho-nucleon amplitude. (auth)

46556 CANONICAL ANOMALIES AND BROKEN SCALE INVARIANCE. Chanowitz, M. S.; Ellis, J. (Stanford Univ., Calif.). *Phys. Lett.*; 40B: No. 3, 397-400(10 Jul 1972).

Canonical behavior of strong interactions at short distances causes an anomaly in the trace identity involving two electromagnetic currents and the energy-momentum tensor. The anomaly is connected with the high energy behavior of $e^+e^- \rightarrow \gamma \rightarrow$ hadrons and the $\epsilon(700)\gamma\gamma$ coupling constant. (auth)

46557 t -DEPENDENCE OF THE FORM FACTORS OF THE NUCLEONS AND THE COMPOSITENESS OF THE ρ -MESON. Bordes, G. (College de France, Paris). *Lett. Nuovo Cim.*; 4: No. 9, 362-4(1 Jul 1972).

Magnetic and electric isovector form factors of the nucleon are calculated in the range of spacelike and low four-momentum transfer $0 < t < 1$ (GeV/c)². (L.B.S.)

46558 SCALING AND FINITE ELECTROMAGNETIC SELF-MASS OF HADRONS. Morita, K. (Dalhousie Univ., Halifax, Nova Scotia). *Lett. Nuovo Cim.*; 4: No. 9, 385-8(1 Jul 1972).

The general conditions to guarantee the finiteness of the electromagnetic self-mass of hadrons are derived under certain assumptions for scaling behavior, Regge behavior with scaling, and analyticity. (L.B.S.)

46559 DUALITY, SCALING, AND FIXED POLES IN VIRTUAL COMPTON SCATTERING. Miranskii, V. A.; Shelest, V. P.; Zinov'ev, G. M. (Inst. for Theoretical Physics, Kiev). *Nucl. Phys.*; B44: No. 2, 460-74(1972).

J-plane fixed poles are considered in the lepton-hadron scattering amplitudes. A relation between the scale-invariant behavior of the virtual Compton scattering structure W_2 and the $J = 1$ pole is established using the wrong signature FESR. A dual mechanism for the generation of fixed poles, leading to a number of interesting implications for a singular behavior of the pole residue in the variable q^2 (photon mass square), is indicated. (auth)

46560 KINEMATICS OF THE COUPLING OF REGGE POLES TO PHOTONS. Collins, P. D. B.; Gault, F. D. (Univ. of Durham, Eng.). *Nuovo Cim.*; 10A: No. 2, 189-206(21 Jul 1972).

Two problems which arise in understanding Regge-pole exchange in amplitudes which include photons are examined; namely how there can be a pion-exchange pole in pion photoproduction, and a finite pomeron coupling in forward Compton scattering. An attempt is made to resolve these problems by noting that the usual sense-nonsense zeros, which might seem to make the couplings vanish, can in fact be replaced in the partial-wave amplitudes by zeros in the kinematical factors. Since the zeros are absent from the total amplitudes, finite couplings remain. This replacement is a peculiarity of photon couplings. It has the disadvantage of changing the threshold behavior of the partial-wave amplitudes, but seems to be the only plausible way out of the dilemma. There are no other special requirements of gauge invariance, so that

these photon-induced processes may be Reggeized in exactly the same way as purely hadronic reactions. (auth)

46561 PION FORM FACTOR AND POSSIBLE HEAVY VECTOR MESON. Furuichi, Susumu (Rikkyo Univ., Tokyo); Kanada, Hiroyuki; Watanabe, Keiji. *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1790-2(May 1972).

The pion form factor is investigated for large t ($t > 1$ (GeV/c)²) for space-like and time-like region on the basis of a simple pole with ρ -meson dominance. It is shown that contributions from an additional pole ρ' greatly improve the fit to the experimental data for the space-like region. (L.B.S.)

46562 PARITY TESTS IN ELASTIC e - p SCATTERING AT LARGE ANGLES. Kuroda, Masaaki; Yamaguchi, Yoshio (Tokyo Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1676-81(May 1972).

The possibility of parity violation in elastic e - p scattering in large momentum transfer regions is discussed. (auth)

46563 PHOTOPRODUCTION OF VECTOR AND AXIAL VECTOR RESONANCES IN BROKEN NONLINEAR CHIRAL $SU_3 \times SU_3$ DYNAMICS. Kelemen, V. I.; Sabo, V. I.; Sosnivker, W. A.; Egorov, A. V. (Inst. of Theoretical Physics, Moscow). *Lett. Nuovo Cim.*; 4: No. 10, 401-6(8 Jul 1972).

The differential cross sections of photoproductions of all vector and all pseudoscalar resonances on baryons N and E are calculated within the framework of broken nonlinear chiral $SU(3) \times SU(3)$ dynamics, and are compared with the predictions of the dispersion approach and with the available experimental data. (L.B.S.)

46564 SIMPLE DIFFRACTION CONSIDERATIONS AND RELATIONS BETWEEN σ_T AND SLOPE OF VECTOR MESON ELECTROPRODUCTION. Eisenberg, Y. (Stanford Univ., Calif.). *Lett. Nuovo Cim.*; 4: No. 10, 425-9(8 Jul 1972).

The (ν, q^2) dependence of the total cross section for $\gamma_v + p \rightarrow \rho^0 + p$ is examined in the framework of the geometric diffraction picture. The behavior of the cross section in the Bjorken limit (ν and $q^2 \rightarrow \infty$ and finite ratio ν/q^2) is also examined. It is concluded that this picture relates in a simple way the observed decrease of the slope B with q^2 to the scaling mechanism and seems to agree with present experimental data. It is also predicted that in the Bjorken limit B would go to zero like $\sim 1/q^2$ if ϵ remains constant, or that $(1-\epsilon) \sim 1/q^2$ with $B \rightarrow \text{constant}$. (L.B.S.)

46565 DEEP INELASTIC SCATTERING, SCALING, AND REGGE BEHAVIOR IN A FEYNMAN DIAGRAM MODEL. Ahmed, M. A. (Univ. of Khartoum, Sudan). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1610-18(May 1972).

The Regge and scaling behavior of the scattering amplitude of massive photons on nucleons are investigated within the context of the dynamical Van Hove-type of model. The correct Regge behavior is obtained but the conventional formulation of the model does not exhibit scaling. The reason for this is pointed out and a cure that restores scaling is given. (auth)

46566 CHARGE DISTRIBUTIONS OF PARTONS AND SCALING IN DEEP INELASTIC SCATTERING. Chowdhury, A. Roy; Chowdhury, R. K. Roy (Jadavpur Univ., Calcutta). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1619-23(May 1972).

A charge distribution of partons inside the proton is conjectured. On this basis νW_2 is calculated in parton model for high energy inelastic scattering. Scaling behavior of νW_2 is reproduced and agrees fairly well with the experimental data. The predicted value of the mean square charge per parton is also lower than that obtained previously. (auth)

46567 SPECIAL METHOD IN S-MATRIX THEORY WITH THE p^4 MOMENTUM REPRESENTATION. Borelowski, Z.; Brzezowski, S. (Jagellonian Univ., Krakow). *Acta Phys. Pol.*; B3: No. 3, 363-73(1972).

A regularization method in the Kadyshevsky momentum representation for the boson-boson and boson-photon interactions is applied. (auth)

46568 DIRAC-HESTENES EQUATION AND THE MAXWELL EQUATIONS. Boudet, Roger (Univ., Aix-Marseille). *Compt. Rend., Ser. A*; 274: No. 16, 1278-80(17 Apr 1972). (In French).

The fusion method is applied to the Hestenes equation. Equations are obtained which are essentially real. (France)

46569 COMPARISON OF THE ANGULAR DECAY DISTRIBUTIONS IN THREE QUARK MODELS OF PHOTOPRODUCTION. Gorczyca, B. (Jagellonian Univ., Krakow). *Acta Phys. Pol.*; B3: No. 3, 391-6(1972).

The constraints for angular decay distributions of resonances photoproduced in the reactions $\gamma B \rightarrow PB^*$, $\gamma B \rightarrow VB$, and $\gamma B \rightarrow VB^*$

in the "excited quark" model and the "interacting quarks" model, where quark structure is assumed only for baryons, are derived and compared with the corresponding predictions of the usual additive quark model of photoproduction including vector meson dominance and the additivity assumption. All the results of the "interacting quarks" model can be derived from the "excited quark" model and all the results of the "excited quark" model can be derived from the additive quark model. (auth)

Weak

46570 (IC-72/54) VIRTUAL NEUTRINO EFFECTS. Budini, P. (International Centre for Theoretical Physics, Trieste (Italy)). Jun 1972. 14p. Dep. NTIS (U. S. Sales Only).

It is shown that the attractive forces due to exchange of neutrino pairs between leptons and antileptons are of such a nature that, if their long-range x^{-6} behavior continues up to distances of the order G_F^2 , they are apt to give massless composite lepton-antilepton states. These states could only have spin-one and they would presumably be medium strongly bound to free leptons. Possible consequences are discussed. (auth)

46571 (KFKI-72-15) SPONTANEOUS DECAY IN A TIME DEPENDENT EXTERNAL FIELD. Hrasco, P. (Magyar Tudományos Akadémia Kozponti Fizikai Kutató Intézete, Budapest). Indl. 11p. Dep. NTIS (U. S. Sales Only).

The parametric resonance in the decay of a system such as a positronium is discussed. It is shown, using Wigner-Weisskopf perturbation theory, that the decay constants depend on the frequency of the external perturbation. (auth)

46572 INTERPRETATION OF EXPERIMENTS ON THE SEARCHES FOR THE $K_S \rightarrow 2\mu$ DECAY. Dolgov, A. D.; Zakharov, V. I.; Okun, L. B. (Inst. of Theoretical and Experimental Physics, Moscow). *JETP Lett. (USSR) (Engl. Transl.)*; 15: No. y, 308-10(5 Apr 1972).

Translated from *Zh. Eksp. Teor. Fiz., Pis'ma Red.*; 15: No. 7, 434-7(5 Apr 1972).

In connection with the contradiction between the results of experiments on the decays $K_L \rightarrow 2\mu$ and $K_L \rightarrow 2\gamma$, it has been proposed that the suppression of the $K_L \rightarrow 2\mu$ decay is due to destructive interference of the $K_2 \rightarrow 2\mu$ and $K_1 \rightarrow 2\mu$ transitions. In this case the width for the $K_S \rightarrow 2\mu$ decay should be relatively large, the lower limit for the width being attainable only in the presence of a strong $2\pi-2\mu$ interaction. The possibility that this interaction is CP noninvariant is discussed, and the limitations imposed on the width of the $K_S \rightarrow 2\mu$ decay are analyzed under different hypotheses concerning the anomalous interactions of muons, photons, and pions. (R.W.R.)

46573 LIMITATIONS ON CURRENTS OF THE SECOND CLASS IN THE PROCESS $\nu_\mu + n \rightarrow p + \mu^-$. Lobov, G. A. *JETP Lett. (USSR) (Engl. Transl.)*; 15: No. 6, 253-5(20 Mar 1972).

Translated from *Zh. Eksp. Teor. Fiz., Pis'ma Red.*; 15: No. 6, 357-9(20 Mar 1972).

An upper bound is obtained for the tensor component of the second-class current in the process $\nu_\mu + n \rightarrow p + \mu^-$ from experimental data on the total cross section. (L.B.S.)

46574 PARTONS MAY HAVE INTEGRAL CHARGE. Budny, R.; Chang, T. H.; Choudhury, D. K. (Oxford Univ.). *Nucl. Phys.*; B44: No. 2, 618-28(1972).

Two parton models are presented for partons having the quantum number of the three triplet models proposed by Han and Nambu and by Cabibbo, Maiani and Preparata. Predictions of deep-inelastic lepton scattering are made and compared with experiment. The former model fares as well as the quark parton model, but the latter model is less successful. (auth)

46575 BOUNDS FOR K_{13} DECAY PARAMETERS. Micu, M. (Inst. for Atomic Physics, Bucharest. Joint Inst. for Nuclear Research, Dubna). *Nucl. Phys.*; B44: No. 2, 531-40(1972).

The extremum of the integral $\int_1^\infty f(x)|F(x)|^2 dx$ is determined for the case when the values taken by the function $F(x)$ at three points are known. The result is applied to the K_{13} problem, the function $F(x)$ being replaced by the form factor $f_+(q^2) + f_-(q^2)q^2/(m_K^2 - m_\pi^2)$ of the divergence of the strangeness changing vector current. (auth)

46576 RADIATIVE CORRECTIONS TO MUON DECAY. Ross, D. A. (Oxford Univ.). *Nuovo Cim.*; 10A: No. 3, 475-89(1 Aug 1972).

The radiative correction to the decay rate of the muon is considered. Particular attention is paid to the cancellation of the infrared divergences. The parts of the radiative correction which lead to these divergences are summed to all orders in the fine-

structure constant, whereas the rest of the correction is calculated to first order only. The result is convergent for all electron energies including the high-energy end point which diverged in previous first-order calculations. (auth)

46577 TESTS FOR A POSSIBLE CLASS OF CP INVARIANT SOLUTIONS TO THE $K_L \rightarrow \mu\bar{\mu}$ PROBLEM. Alles, W. (Univ., Bologna. Istituto Nazionale di Fisica Nucleare, Bologna); Pati, J. C. *Nuovo Cim.*; 10A: No. 2, 325-42(21 Jul 1972).

The possibility that the unitarity sum for the absorptive part of the $K_L \rightarrow \mu\bar{\mu}$ amplitude receives contribution in a CP-invariant manner from an on-mass-shell intermediate state involving a yet undetected light neutral boson (χ^0) is considered. It is suggested that the most relevant intermediate state is the ($\chi^0\gamma$) state with χ^0 having spin-parity 1^- . It is pointed out that the existence of such a particle, if it is not already excluded by present experiments, can be tested unambiguously and relatively easily by two experiments: (a) the measurement of the rate of the $K_L \rightarrow \chi^0 + \gamma$ decay (the χ^0 may be looked for easily via its decays to (e^-e^+) and ($\mu^-\mu^+$) systems), and (b) the production of the χ^0 by low-invariant-mass (e^-e^+) colliding-beam experiments. It is noted that the χ^0 , if it exists, receives a "natural" interpretation only in framework of hadron symmetries higher than the familiar SU_3 -group. The production of χ^0 by a host of production and decay processes is discussed subject to such an interpretation. (auth)

46578 LEPTON CHARGE AND WEAK NEUTRINO-LEPTON INTERACTION. Romanov, Yu. I. *Izv. Vyssh. Ucheb. Zaved.*, Fiz.; No. 12, 30-4(1971). (In Russian).

It was postulated that a system of leptons is characterized by one lepton charge with different magnitudes for μ -leptons and e -leptons. Within the framework of this scheme weak neutrino-lepton interactions were studied. "Charged" ($V - A$)-currents were introduced in order to describe $\bar{\nu}_e \rightleftharpoons \nu_\mu$ and $\bar{\nu}_\mu \rightleftharpoons \nu_e$ transitions. (tr-auth)

46579 INTERACTION PROCESSES OF NONZERO REST MASS ν AND $\bar{\nu}$ WITH LEPTONS. Rekalov, A. P. (Physical-Technical Inst., Kharkov). *Ukr. Fiz. Zh. (Russ. Ed.)*; 16: No. 10, 1585-92(Oct 1971). (In Russian).

Expressions for the differential and total cross sections of different processes of nonzero mass neutrino interaction with leptons are obtained in terms of invariant variables. The relations correct for $m_\nu = 0$ and $m_\nu \neq 0$, between the cross sections of lepton weak processes are found. The energy dependence of the energy loss and the scattering angle of ν and $\bar{\nu}$ in νe and $\bar{\nu} e$ scattering are analyzed in detail. (auth)

46580 $\Delta = 1/2$ RULE IN THE THREE-TRIPLET AND PARAQUARK MODELS. Kingsley, R. L. (Cambridge Univ., Eng.). *Phys. Lett.*; 40B: No. 3, 387-91(10 Jul 1972).

It is possible to propose a charged weak interaction current which will generate a non-leptonic weak Hamiltonian satisfying the $\Delta I = 1/2$ rule in the models of three triplets with integral charges, but not in the paraquark model. (auth)

46581 QUASI-ELASTIC NEUTRINO REACTION IN DEUTERIUM. Bernabeu, J.; Pascual, P. (Univ. of Valencia). *Nuovo Cim.*; 10A: No. 1, 61-75(1 Jul 1972).

The process $\nu + d \rightarrow \mu^- + p + p$ is studied in the framework of the impulse approximation, with or without using closure approximation. Final-state interactions are considered. (auth)

46582 ELECTROMAGNETIC DIVERGENCES IN WEAK PROCESSES. Bailin, D. (Univ. of Sussex, Eng.). *Nuovo Cim.*; 10A: No. 1, 159-72(1 Jul 1972).

Earlier calculations of the leading divergence in weak transitions between hadrons are generalized by including the electromagnetic interactions of the hadrons and of the charged W-boson. It is found that these can be removed by a self-mass counter term having the same form as that found earlier. However, there is an additional counter term in which the hadrons, in the form of the neutral Cabibbo current, are coupled directly to the electromagnetic field. (auth)

46583 K_{13} -DECAY PARAMETERS AND POSSIBLE SCALAR COUPLING IN WEAK INTERACTIONS OF URBARYONS. Kaneko, T. (Meijo Univ., Nagoya, Japan). *Lett. Nuovo Cim.*; 4: No. 9, 373-6(1 Jul 1972).

The K_{13} decay is discussed under the assumption of possible existence of scalar coupling in the weak interactions of urbaryon. It is pointed out that under this assumption, the relation of the scalar form factor to the Callan-Treiman relation can be understood. (L.B.S.)

46584 NEUTRINO REACTIONS AT ACCELERATOR ENERGIES. Smith, C. H. Llewellyn (Stanford Univ., Calif.). *Phys. Rep.*; 3C: No. 5, 261-379(Jun 1972).

Possible investigations of the properties of weak interactions using high energy neutrinos are reviewed. The use of neutrinos to probe hadron structure is discussed. (auth)

46585 MIXING PARAMETER ϵ IN NEUTRAL K DECAY. Ito, Yukio (Fukushima Univ., Japan). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1777-8(May 1972).

An explicit formula between the mixing parameter ϵ and $\gamma_1, \gamma_2, \Delta m (= m_1 - m_2)$ of K_1 and K_2 in superweak theory was previously shown by Kenny and Kabir (*Phys. Rev. D*; 2: 257(1970)). It is shown that the relation obtained by Kenny and Kabir holds in the general frame of Wu and Yang's theory without assuming $A_\alpha = \text{Re}A_\alpha$. (L.B.S.)

Strong (General)

Refer also to abstracts: 46462, 46683, and 46686.

46586 (JINR-P2-6501) FEYNMAN-WILSON-GAS MODEL AS A CONSEQUENCE OF JET APPROACH TO INELASTIC HADRON PROCESSES. Dushutin, N. K.; Mal'tsev, V. M.; Sheptil, V. I. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 14p. (In Russian). Dep. NTIS (U. S. Sales Only).

A possibility is shown to formulate the Feynman-gas model as a consequence of the jet approach to the process of deep-inelastic hadron interactions. The sense of the correlation functions, included into the model, is clarified. The further trends of the model generalization are shown. (auth)

46587 (RLO-2230-T-4-1) TESTS OF HIGH-ENERGY MODELS BY TWO-PARTICLE ANALYSES. Hwa, Rudolph C. (Oregon Univ., Eugene). 1972. Contract AT(45-1)-2230. 12p. Dep. NTIS.

Two-particle distribution functions are calculated in the diffractive-excitation model and compared with those of the multiperipheral model. Optimum regions for differentiating the predictions of the models are indicated. Decisive, yet feasible, experimental tests are suggested. (auth)

46588 POSSIBLE EXISTENCE OF HADRONIC ISOMERS. Pontecorvo, B. (Joint Inst. of Nuclear Research, Dubna, USSR). *JETP Lett. (USSR) (Engl. Transl.)*; 15: No. 6, 250-3(20 Mar 1972).

Translated from *Zh. Eksp. Teor. Fiz., Pis'ma Red.*; 15: No. 6, 353-6(20 Mar 1972).

It is pointed out that, owing to the uncertainty relation, an examination of the angular momenta reveals a unique similarity between quantum objects that differ strongly in their scales and in their mean energy distances between excitation levels. The question of possible metastability connected with large values of J in the field of hadron physics analogous to nuclear isomerism is considered. Experiments aimed at searching for hadronic isomers are discussed. (L.B.S.)

46589 WAVE PACKET REALIZATION OF LIGHTLIKE STATES. Domokos, G. (Johns Hopkins Univ., Baltimore). *Commun. Math. Phys.*; 26: No. 1, 15-23(1972).

A new realization of lightlike particle states is constructed. The realization considered corresponds to objects which form partially localized wave packets in the space dimensions transverse to the direction of the lightlike momentum. The states of the transverse motion are classified by the representations of an algebra $SU(1,1)$ constructed of the generators of the conformal group. It is conjectured that this realization is suitable for the formulation of the "parton"-model of hadrons. (auth)

46590 HIGH ENERGY PHENOMENOLOGY. Proceedings of the Second Session of the Sixth Rencontre de Moriond, Meribelles-Allues, France, March 7-9, 1971. Tran Thanh Van, J. (ed.). Paris: Laboratoire de Physique Theorique et Hautes Energies (1971). 559p. (In French and English). (CONF-710345-(Vol.2)).

There were 28 papers on high energy strong interaction phenomenology presented. Separate abstracts were prepared for 27 of these papers, the remaining paper having been published in abstract form. (France)

For abstracts of individual papers see: 46436-46438, 46462-46465, 46488-46492, 46533-46435, 46591, 46592, and 46627-46636.

46591 LARGE ANGLE SCATTERING OF HADRONS. Krzywicki, A. (Univ. Paris, Orsay, France). pp 187-211 of *High Energy Phenomenology*. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

Large angle elastic scattering of hadrons at high energy is discussed (a few GeV). Experimental data are briefly surveyed and the various theoretical models applying to the elastic case are presented: small angle scattering models extended to large angles, and the statistical model. (France)

46592 EXCHANGE DEGENERACY AND DIP MECHANISMS IN HIGH ENERGY COLLISIONS. Tran Thanh Van, J. (Univ. Paris, Orsay, France). pp 485-525 of *High Energy Phenomenology*. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Exchange degeneracy in high energy collisions is envisaged. It is shown to be verified for ρ - A_2 and K^*-K^{**} in the reactions with small Regge cuts. The two classes of high energy models which are essentially characterized by the presence or absence of non-sense wrong signature zeros are also discussed. Two crucial tests of the different models are presented: measurement of ρ_{00}^0 in $\pi^+n \rightarrow \omega p$ or $\pi^-p \rightarrow \omega n$, precise measurement of the $\pi^-p \rightarrow \eta n$ differential cross section at very small t as well as at large t . (France)

46593 ATTEMPT AT GENERALIZING DUAL-RESONANCE MODELS. A GROUP-THEORETICAL APPROACH. Matsumoto, H. (Tokyo Univ.). *Nuovo Cim.*; 10A: No. 3, 445-67 (1 Aug 1972).

Generalizations of dual-resonance models are discussed within the context of the group-theoretical formulation of the theory. Fields irreducible under $SL_{2,R}$ transformations are introduced, which are closely related to the $SL_{2,C}$ irreducible ones. The scheme has a simple factorization property and is convenient to see resonance structures. To satisfy the cyclic symmetry, three types of constructions of vertices are discussed. Gauge properties are also investigated in a model-independent way. It is shown that the Virasoro's general gauge is realized only for fields with the $SL_{2,R}$ spin 0, $-\frac{1}{2}$ and -1 . In the light of these results, it is concluded that the general gauge and realistic models ($\alpha_p(0) = \frac{1}{2}$) are incompatible in additive models. (auth)

46594 DYNAMIC INTERPRETATION OF BARYON AND MESON MASS FORMULAE. Libby, L. M. (Univ. of California, Los Angeles). *Phys. Lett.*; 40B: No. 3, 307-10 (10 Jul 1972).

A dynamical interpretation of baryon and meson mass formulae offers an explanation for associated production, for parity non-conservation, and for the $T = \frac{1}{2}$ rule. (auth)

46595 POLARIZATION IN INCLUSIVE REACTIONS. Kuroda, M.; Oyanagi, Y. (Tokyo Univ.). *Lett. Nuovo Cim.*; 4: No. 13, 627-30 (29 Jul 1972).

Polarization in hadronic inclusive interactions is examined and expressions are obtained for the asymmetry in the fragmentation region and in the triple-Regge region. The asymmetry of photo-induced inclusive interactions is briefly examined. (L.B.S.)

46596 DERIVATION OF THE CRITICAL TEMPERATURE FOR THE SUPERFLUID PHASE OF HADRONIC MATTER. Mann, A.; Weiner, R. (Indiana Univ., Bloomington). *Phys. Lett.*; 40B: No. 3, 383-6 (10 Jul 1972).

A quantitative derivation is given of the temperature at which superfluidity of hadronic matter ends. Using considerations similar to those used by Landau for ^4He , $T_c \leq 190$ MeV is found in close agreement with Hagedorn's "limiting" temperature. (auth)

46597 SOME CONSEQUENCES OF THE VANISHING OF THE TRIPLE-POMERONCHUKON-VERTEX. Rajaraman, R. (Cornell Univ., Ithaca, N. Y.). *Phys. Lett.*; 40B: No. 3, 392-6 (10 Jul 1972).

Starting with the assumption that the triple-pomeronchukon-vertex vanishes when all three pomeronchukons have zero mass, it is shown that in the absence of pionization, total cross sections would fall at least logarithmically with energy, while the mean multiplicity would approach a constant. Several reasons are presented to suggest that some pionization must occur and that, otherwise, limiting fragmentation itself would vanish. A secondary result, that the P - P' coupling will also vanish in certain regions, is also proved. (auth)

46598 POSSIBLE BOUNDS ON MUON PAIR AND W^+ PRODUCTION IN SUPER-HIGH-ENERGY HADRON COLLISIONS. Mikamo, S. (Tokyo Univ.). *Nuovo Cim.*; 10A: No. 2, 303-10 (21 Jul 1972).

Possible bounds on the muon pair and W^+ production cross sections in high-energy hadron collisions are discussed in connection with a theory predicting cross sections linearly rising with the incident-proton energy for both cases. It is pointed out that the cross sections do not seem to exceed the order of $10 \mu\text{b}$ up to several hundred TeV proton energies from the observation of the steeply falling spectrum of cosmic-ray muons where the energetic muons produced forward from such muon sources are expected to flatten the spectrum. (auth)

46599 PARTON MODEL AND THE CONCEPT OF DUALITY IN STRONG INTERACTIONS. Bandopadhyay, P. (Indian Statistical Inst., Calcutta); De, S. S. *Lett. Nuovo Cim.*; 4: No. 9, 377-80 (1 Jul 1972).

It is shown that the concept of duality can be incorporated into the framework of the parton model of hadrons from the dynamics of strong interactions. The various strongly interacting processes are generalized by arguing that all baryons as well as K -meson contain a π -meson in their structure and the specific interaction can be represented in terms of the pion-pion interaction and the corresponding rearrangement of partons. The $\pi N \rightarrow \pi N$ and $\Delta\Delta$ interactions are considered as examples. (L.B.S.)

46600 DUALITY PROPERTIES OF REGGE CUTS. Collins, P. D. B.; Hutt, P. K. (University Science Labs., Durham, Eng.). *Lett. Nuovo Cim.*; 4: No. 9, 347-51 (1 Jul 1972).

It is shown that quark-parton duality diagrams give a convenient mnemonic for the internal-symmetry and duality properties of Regge cuts. (L.B.S.)

46601 PHOTONLIKE PARTICLES, COMPTON SCATTERING, AND SUM RULES IN THE DUAL-RESONANCE MODEL. Friedman, K. A.; Rosenzweig, C. (Massachusetts Inst. of Tech., Cambridge). *Nuovo Cim.*; 10A: No. 1, 53-60 (1 Jul 1972).

An analogy is drawn between the real photons appearing in electrodynamics and the photonlike particle appearing in the dual-resonance model with unit intercept. Sum rules for the Compton scattering of these particles are derived and, by employing the analogy, interpreted in terms of low-energy theorems. In particular the "universal" property of magnetic moments and the lack of anomalous moments in the dual model are rederived. It is also shown that the Beg-type sum rule exists and is saturated by relatively few states. This is a direct result of the simplicity of the collinear algebra appearing in the dual model. (auth)

46602 SELF-CONSISTENT DIFFRACTION IN THE MULTIPERIPHERAL MODEL. Ball, J. S.; Zachariasen, F. (California Inst. of Tech., Pasadena). *Phys. Lett.*; 40B: No. 3, 411-14 (10 Jul 1972).

An exact self-consistent solution, for small t and very large s , of the multiperipheral bootstrap equations is presented. The solution predicts that total cross sections grow like $\log s$, that elastic total cross sections are constant, that elastic diffraction peaks shrink like $(\log s)^2$, and that multiplicities grow like $\log s$, all in the limit of infinite s . (auth)

46603 SPACE-TIME DESCRIPTION OF HADRONS. Mansouri, Freydoon. Baltimore; Johns Hopkins Univ. (1969).

134p. University Microfilms Order No. 72-16,839. Thesis.

The possibility of a consistent space-time description of hadrons is investigated. It is assumed that to zeroth approximation the amplitudes and propagators transform according to the irreducible representations of the proper complex Lorentz group, $L_+(C)$, or its covering group $SL(2,C) \otimes SL(2,C)$. To have both Lorentz covariance and Regge behavior, it becomes necessary to work in a realization of the representations of $SL(2,C)$ which is insensitive to the little group structure of the Poincare group. Amplitudes and propagators are constructed for free hadrons. The behavior of hadron amplitudes under C , P , and T is worked out. It is shown, moreover, that these amplitudes have definite transformation properties under fermion number and chiral gauge transformations, which determine the Toller quantum number, M . In particular, it is found that both fermions and mesons form chiral doublets. Explicit expressions for $M = 0$ and $M = \pm \frac{1}{2}$ propagators are given. They depend on one unknown parameter: The Lorentz trajectory of the corresponding hadron. To describe the interactions of hadrons, the structure of vertices is investigated. It is found that in general they are quasiloocal. The knowledge of wave functions, propagators, and vertex functions lead to a set of simple diagram rules for the construction of the invariant scattering amplitudes. The application of the diagram rules to meson-

meson scattering processes with a single exchange reproduces the well known contributions of the corresponding Lorentz poles with rapidly converging residues. (Diss. Abstr. Int., B)

46604 DOES THE SIMPLE QUARK LIGHT-CONE CURRENT ALGEBRA OF FRITZSCH AND GELL-MANN IMPLY THAT QUARKS DO EXIST. Haberler, P. L. F. (Institut fuer Hochenergiephysik der Oesterreich, Vienna). Lett. Nuovo Cim.; 4: No. 10, 393-7 (8 Jul 1972).

It is suggested that if the light-cone algebra of Fritzsche and Gell-Mann implies that the quark propagator behaves at high energies like a free propagator, then using standard assumptions, the existence of free quarks is implicitly postulated. (L.B.S.)

46605 INCLUSIVE REACTIONS IN THE REGGE POLE MODEL AND EXOTIC RESONANCES. Hara, Yasuo (Tokyo Univ. of Education). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1624-42 (May 1972).

The Regge pole model and duality are applied to processes of the type $a + b \rightarrow c + \text{anything}$. The exotic and nonexotic three-particle channels are defined. The existence of resonances in the $K^+\pi^+\pi^0$, $p\bar{n}\pi^+$, $K^0\pi^+\pi$ and $K^-\pi^-\pi$ channels is predicted in the model. (auth)

46606 URBARYON REARRANGEMENT AND INCLUSIVE PHENOMENOLOGY. Matsuoka, Takeo (Nagoya Univ., Japan). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1643-62 (May 1972).

From the standpoint of urbaryon rearrangement, the characteristic features of various inclusive processes are discussed. In the urbaryon rearrangement scheme, the condition for the early onset of limiting behavior of the inclusive process is predicted, which is different from those given by Chan et al. and Ellis et al. The inclusive processes are classified into six categories according to their distinctive features of the differential cross section. For the process in each category, definite predictions are obtained for the rate in which the differential cross section approaches a limit. It is shown that these predictions are in good agreement with available data. Sum rules among various inclusive processes are also derived. (auth)

46607 POSSIBLE EXPLANATION OF THE STRONG CORE PRESENT IN HADRON REACTIONS. Nakkagawa, Hisao (Kyoto Univ.). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1663-75 (May 1972).

Concerning the "strong core" in hadron scattering it is investigated, on the basis of the composite model, whether the physical relative wave function can have the inner oscillation when a redundant solution does not exist. The results are 1) even when there is no redundant solution, there exists an inner oscillation, or "strong core" and 2) as inelastic channels begin to open, this "strong core" gradually vanishes. These results strongly suggest the existence of fundamental entities in the hadron. (auth)

46608 SIZE OF HADRONIC CONSTITUENTS AND APPLICABILITY OF SCALING LAW. Matumoto, Ken-iti (Toyama Univ., Japan). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1795-6 (May 1972).

The effect of the size of the urbaryon, which appears together with the effect of the composite structure of hadrons is examined on the basis of transition amplitudes caused by electromagnetic or weak currents. Considering the deep inelastic process $e + p \rightarrow e + \text{anything}$, lower limits are obtained for the urbaryon and lepton. Information on the validity of the scaling behavior in this region is also obtained. (L.B.S.)

Strong (Baryon No. = 0)

Refer also to abstract: 46458.

46609 (ITF-72-13-P) ON THE DISTRIBUTION OF RESONANCE SPINS IN THE DUAL RESONANCE MODEL. Struminski, B. V. (Akademiya Nauk Ukrainskoi SSR, Kiev, Institut Teoreticheskoi Fiziki). 1972. 7p. (In Russian). Dep. NTIS (U. S. Sales Only).

The distribution of resonance spins in the dual resonance model is calculated by the statistical mechanics method. (auth)

46610 PARTIAL WAVE RELATIONS FROM FIXED- u DISPERSION RELATIONS. Baacke, J. (Freie Univ., Berlin). Nucl. Phys.; B44: No. 2, 594-604 (1972).

Partial wave relations are derived from fixed- u dispersion relations for meson-baryon and meson-meson scattering. General expressions for the integral kernels are given and some problems pertaining to the application of the relations are discussed. (auth)

46611 NONET STRUCTURE OF THE $J^{PC} = 3^{--}$ MESONS. Bramon, A.; Greco, M. (CNEN, Frascati, Italy). Nuovo Cim.; 10A: No. 3, 521-32 (1 Aug 1972).

The meson resonances $g(1660)$, $\varphi_N(1650)$, $\varphi_N(1830)$ and $K_N(1760)$ are tentatively grouped in an "ideal" SU_3 nonet having $J^{PC} = 3^{--}$. A discussion of their masses and their decay modes is presented showing general agreement with the available experimental data. The results are also discussed in the light of finite-energy sum rules for the reactions $\pi K \rightarrow \omega K$ and $\pi K \rightarrow \varphi K$. (auth)

46612 COLLINEAR DISPERSION RELATIONS AND THE $\rho-\pi-\pi$ AND $A_1-\rho-\pi$ COUPLINGS. Nasrallah, N. F. (Institut des Hautes Etudes Scientifiques, Bures-sur-Yvette, France). Nuovo Cim.; 10A: No. 2, 247-56 (21 Jul 1972).

It is shown that the use of collinear dispersion relations and of equal-time commutators obtained from current and field algebra, as well as pole dominance, allows the completion of the program of Riazuddin and Fayyazuddin and Geffen, i.e., the determination of the $\rho-\pi-\pi$ and the two $A_1-\rho-\pi$ coupling constants. It is found that, in this approach, the effect of the A_1 is to reduce considerably (by about 80%) the width of the ρ as given by the KSFR relation, and it is concluded that the assumptions of the previously mentioned authors on the on-mass-shell behavior of the amplitudes are too strong, as has already been suggested by Brown and West, who used on-mass-shell dispersion relations. The results yield the relation $f_A/f_\rho = \sqrt{M_A/M_\rho}$ between the normalizations of the A and ρ fields and, in conjunction with Weinberg's first sum rule a slightly modified version of the KSFR relation. (auth)

46613 LIMITING BEHAVIOR OF PARTIALLY INCLUSIVE REACTIONS. Satz, H. (Univ. of Bielefeld, Ger.). Lett. Nuovo Cim.; 4: No. 9, 357-61 (1 Jul 1972).

The limiting behavior of single-particle distributions in partially inclusive interactions is investigated for the case of $p\bar{p} \rightarrow \pi + \text{pions}$ for $S \rightarrow \infty$, where s is the squared total energy. The energy dependence of the cross section is also examined. (L.B.S.)

46614 PROCESS $\nu_\mu + p \rightarrow \nu_\mu + p + \pi^0$ IN WEINBERG'S MODEL OF WEAK INTERACTIONS. Lee, B. W. (National Accelerator Lab., Batavia, Ill.). Phys. Lett.; 40B: No. 3, 420-2 (10 Jul 1972).

The ratio $R = \sigma(\nu_\mu + p \rightarrow \nu_\mu + p + \pi^0) / \sigma(\nu_\mu + n \rightarrow \mu + p + \pi^0)$ in Weinberg's model of weak interactions is estimated on the basis of a static model for the neutrino production of the 3,3 resonance. For the Weinberg mixing angle α given by $\sin^2 \alpha \leq 0.35$ as deduced from the upper bound on $\sigma(\bar{\nu}_e + e \rightarrow \bar{\nu}_e + e)$, it is concluded that $R \gtrsim 0.4$, which is not consistent with the experimental bound $R \lesssim 0.14$. (auth)

46615 PION FORM FACTOR IN THE BETHE-SALPETER MODEL. Fleischer, J. (Univ., Karlsruhe, Ger.); Gutbrod, F. Nuovo Cim.; 10A: No. 2, 235-46 (21 Jul 1972).

A Bethe-Salpeter model is presented in which phenomenological potentials are used to reproduce the ρ peak in the $\pi\pi$ scattering amplitude. Then the off-shell amplitude is used to calculate the pion electromagnetic form factor. The Bethe-Salpeter equation is used in the ladder approximation and is solved by iteration and application of the Padé approximation method. Reasonable agreement with the vector-dominance model is achieved. (auth)

Strong (Baryon No. = 1)

Refer also to abstract: 46487.

46616 (BNL-16928) SCALING IN $\pi\pi^- \rightarrow pX^-$ AND AN ESTIMATE OF THE TRIPLE-POMERANCHUKON COUPLING. Paige, Frank E.; Wang, Ling-Lie (Brookhaven National Lab., Upton, N. Y.). Jun 1972. 6p. (CONF-720409-3). Dep. NTIS.

From third international conference on experimental meson spectroscopy; Philadelphia, Pa. (28 Apr 1972).

It is shown that the CERN-IHEP data on $p\pi^- \rightarrow pX^-$ at 25 and 40 GeV/c exhibit scaling in s'/s . An estimate of the triple-Pomeranchukon coupling is given, and the possibility of its measurement at NAL is discussed. (auth)

46617 (IFVE-STF-71-15) ESTIMATION OF INTERMEDIATE ENERGY CONTRIBUTION TO THE DISPERSION SUM RULES AND ROTATION OF POLARIZATION IN PION-NUCLEON SCATTERING AT HIGH ENERGIES. Aznauryan, I. G.; Solov'ev, L. D. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoi Energii SSSR, Serpukhov, Institut Fiziki Vysokikh Energii). 1971. 21p. (In Russian). Dep. NTIS (U. S. Sales Only).

The Serpukhov data on total cross sections show the violation of the Regge model at energies above 30 GeV. This fact increases the interest in the complete check of this model at lower energies. It is well known that the Regge pole model gives a good descrip-

tion of the cross sections of polarization for pion-nucleon elastic scattering at 4 to 25 GeV. It is shown that this model is also true in describing rotation of polarization in π^+p -scattering at 6 and 16 GeV. (auth)

46618 (IFVE-STF-71-71) TOTAL CROSS-SECTIONS FOR REACTION $\pi^+p \rightarrow \pi^0n$ AT HIGH ENERGIES. Bababev, Z. R. (Gosudarstvennyi Komitet po Ispol'zovaniyu Atomnoi Energii SSSR, Serpukhov. Institut Fiziki Vysokikh Energii). 1971. 8p. (In Russian). Dep. NTIS (U. S. Sales Only).

The total cross section for the reaction $\pi^+p \rightarrow \pi^0n$ at high energies was calculated and compared with experimental data. When calculating use was made of theoretical predictions on differential cross sections at 0° angle for this process with different parametrizations of the difference between the total cross sections for π^+ -scattering. (auth)

46619 (JINR-P1-6461) REMARK ON SPECTRUM SYMMETRY OF PION LONGITUDINAL MOMENTA IN π^+p INTERACTIONS AT $pc = 4$ GeV. Grishin, V. G.; Inogamov, Sh. V.; Yuldashev, B. S.; Yancho, G. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of High Energy). 1972. 7p. (In Russian). Dep. NTIS (U. S. Sales Only).

It is shown that the spectrum symmetry of the pion longitudinal momenta in π^+p interactions at $pc = 4$ GeV is observed in the system where $R = 1.69 \pm 0.15$. Possible interpretation of this effect is presented. (auth)

46620 (JINR-P4-6475) QUASI-THRESHOLD BEHAVIOR OF THE $\pi + N \rightarrow N + (\gamma\text{-VIRTUAL})$ PROCESS AMPLITUDE. Surovtsev, Yu. S.; Tkebuchava, F. G. (Joint Inst. for Nuclear Research, Dubna (USSR)). 1972. 8p. (In Russian). Dep. NTIS (U. S. Sales Only).

It is shown that the cross section of the process $\pi + N \rightarrow N + (\gamma\text{-virtual})$ at small values of the virtual photon momentum is determined, in general, by the Born terms of the amplitude. Thereby, the description of the process becomes nearly slowly independent when the virtual photon "mass" tends to its maximum value at fixed total energy. (auth)

46621 TESTS OF THE SPIN ASSIGNMENT AND EQUAL-PHASE ASSUMPTION FOR HIGH-SPIN BARYONS. Donohue, J. T. (Univ., Nice). Nucl. Phys.; B44: No. 2, 355-64(1972).

For baryon resonances with $\frac{1}{2} \leq J \leq \frac{3}{2}$ a set of tests are proposed for the spin assignment. The tests are relations between the sums of squares of the even-L statistical-tensors t_L^M and the target polarization for the reaction $M + B(P) \rightarrow M + B^*$, and are extensions of an earlier result. In addition it is shown how these relations can be used, when J is known, to test the dynamical assumption that all helicity amplitudes for the reaction have the same phase, this without a polarized target and without observing a parity-violating decay. (auth)

46622 PHENOMENOLOGICAL PRESCRIPTION FOR REGGE CUTS. Ringland, G. A. (Rutherford High Energy Lab., Chilton, Eng.); Roberts, R. G.; Roy, D. P.; Tran Thanh Van, J. Nucl. Phys.; B44: No. 2, 395-412(1972).

The failure of the old Reggeized absorption model is attributed to the wrong relative phase of pole and cut. A model for vector and tensor exchange is proposed in which the old absorption model is used to generate the cut coming from the component of the pole term with phase $e^{-i\pi\alpha(t)}$ [planar duality diagram]. However, the absorption model cut obtained from the constant phase part of the pole term (non-planar duality diagram) is rotated in phase by $\frac{1}{2}\pi$. This model may be applied over a wide range of energy, but not, by crossing symmetry, at infinitely high energies. This prescription is used to discuss $0^- + \frac{1}{2}^+ \rightarrow 0^- + \frac{1}{2}^+$ reactions using a dual pole input with wrong signature nonsense zeros and SU(3) constraints. The $\pi^+p \rightarrow \pi^0n$ polarization and also the cross section inequalities and polarizations of hypercharge exchange reactions are obtained. Further, this model gives a quantitatively better understanding of the crossover zeros. An additional bonus is that from a nonsense choosing P' input a P' amplitude is obtained qualitatively similar to that found by FESR and the no-compensation form required by pure pole fits. Predictions of polarization in $\bar{K}N$ and KN charge exchange and $\pi N \rightarrow \eta N$ are given. (auth)

46623 NATURE OF ABSORPTION. Barger, V. (CERN, Geneva); Geer, K.; Halzen, F. Contract AT(11-1)-881. Nucl. Phys.; B44: No. 2, 475-89(1972).

The imaginary helicity non-flip ω exchange amplitudes in $\bar{K}N$ and NN scattering are determined versus t and s from data on elastic differential cross section differences. The ω -universality prediction, $\text{Im } \omega_N(t) = 3 \text{ Im } \omega_K(t)$, is found to be satisfied for $0 \leq |t| \leq 0.5$, indicating that the absorptive cuts at small t preserve universality. The impact parameter distributions of the ω ampli-

tudes show: (i) a peripheral peak centered around $b \approx 1$ fm, with $\Delta b \approx 0.5$ fm in $\bar{K}N$ and $\Delta b \approx 0.75$ fm in NN ; (ii) a sizeable negative contribution at small b , more negative in NN than in $\bar{K}N$, (iii) a smaller tail of negative contributions around $b \approx 2$ fm. Strong absorption of the ω -pole imaginary amplitude with λ values around two is needed to reproduce this structure. A new version of the absorption model, suggested by πN amplitude analysis, is confirmed by our results. (auth)

46624 ANALYSIS OF KAON-NUCLEON SCATTERING AMPLITUDES IN THE INTERMEDIATE ENERGY REGION. Fukugita, M.; Inami, T. (Tokyo Univ.). Nucl. Phys.; B44: No. 2, 490-508(1972).

"Two component theory" of duality was tested with new $\bar{K}N$ and KN partial wave data in the intermediate energy region. A comprehensive study of the s -channel helicity amplitudes constructed separately from the background and resonant parts of partial wave amplitudes is presented. It is found that the background and resonant amplitudes have properties which are represented well by the pomeron and ordinary Regge exchanges, respectively. This approach also sheds new light on previous FESR calculations and the problem of exchange degeneracy. (auth)

46625 AMPLITUDE ANALYSIS OF πN COULOMB INTERFERENCE DATA. Krubasik, E. (Univ., Karlsruhe, Ger.). Nucl. Phys.; B44: No. 2, 558-72(1972).

πN scattering amplitudes near the forward direction were derived directly from the data in the 7 to 20 GeV/c region. A linear exponential behavior for the $I_t = 0$ part of the nonflip amplitude is found. The slopes of the diffraction peak yield a shrinkage in this energy range. Results for the real parts of the $I_t = 0$ and $I_t = 1$ components are compatible with the dispersion relation, but difficulties are encountered in a conclusive test of these relations. An energy independent crossover is found at $0.1 \leq -t \leq 0.15$ (GeV/c)² and its mechanism is explained. A non-Regge-pole contribution is seen in the $I_t = 1$ component of the nonflip amplitude. (auth)

46626 MASS RADIUS OF THE NUCLEON. Hare, M. G.; Papini, G. (Univ. of Saskatchewan, Regina). Can. J. Phys.; 50: No. 11, 1163-8(1 Jun 1972).

The mean radius of the mass distribution of the nucleon is determined to be $\langle r_1 \rangle = 0.7$ fm. The calculation makes use of sidewise, unsubtracted, threshold dominated dispersion relations for the form factors appearing in the matrix elements of the contracted energy-momentum tensor. It uses a π meson-nucleon intermediate state expansion. (auth) (Canada)

46627 DUALITY AND BACKWARD PEAKS. Halzen, F. (CERN, Geneva). pp 65-96 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

The implications of duality constraints on baryon exchange are studied using a factorizable Regge pole model for pseudoscalar meson baryon $\rightarrow$ pseudoscalar meson baryon scattering, and the patterns of baryon exchange degeneracy are derived. The experimental and phenomenological tests of the scheme are discussed: duality sum rule, effective residues, and Regge fits. Absence of parity doublets and the behavior of backward peaks at large momentum transfers are then examined. (France)

46628 BARYON SPIN DETERMINATIONS USING POLARIZED TARGETS. Donohue, J. T. (Univ., Nice). pp 181-5 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol. 2).

A new method for determining the spin (J) of the baryon resonances produced in quasi-two body reactions with polarized targets is discussed. Decays are analyzed in the framework of the spin-density matrix, using the concept of pure-state and the theorem of Eberhard and Good. The $a_{LM}(t, \phi)$ coefficients of the spherical harmonics expansion for even-L are measured for two or more values of the polarization (P); t_{LM} or ρ_P is then determined by assuming a definite value of J; $(d\sigma/d\Omega)^{\frac{1}{2}} (\text{Tr}(\rho_P^2))^{-\frac{1}{2}}$ is plotted as a function of $1-p^2$ and an attempt is made to discover whether it lies on a straight line passing through the origin. (France)

46629 LINE REVERSED REACTIONS. Michael, C. (CERN, Geneva). pp 247-57 of High Energy Phenomenology.

/Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Recent data on line reversed reactions are presented. Comparisons between $K^+n \rightarrow K^0p$ and $K^+p \rightarrow \bar{K}^0n$ at 2.3 and 3 GeV/c; $\pi^+\pi^- \rightarrow \pi^+\pi^+$ and $\pi^+\pi^- \rightarrow \pi^-\pi^-$ and $\pi^+\pi^- \rightarrow \pi^+\pi^-$, and $K^+p \rightarrow pK^+$ and $\pi^+p \rightarrow K^+K^+$ at 5 GeV/c; $\bar{K}N \rightarrow \pi Z$ and $\pi N \rightarrow KZ$, and $KN \rightarrow \pi A$ and $\pi N \rightarrow KA$ indicate the presence of low lying singularities explaining the energy dependence of the cross section variations in the 3 to 8 GeV/c region. The complexity of the angular momentum plane description for the scattering amplitude is thus emphasized. (France)

46630 PHENOMENOLOGICAL ANALYSIS OF πN AND KN CHARGE-EXCHANGE REACTIONS. Dronkers, J. (CEN de Saclay, Gif-sur-Yvette, France). pp 259-66 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971). (In French).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

$\pi^+p \rightarrow \pi^+n$, $\pi^+p \rightarrow \pi^0p$, $K^+p \rightarrow \bar{K}^0n$ and $K^+n \rightarrow K^0p$ reactions with $1^-(\rho)$ and $2^+(A_2)$ octet exchange in the crossed channel were studied. The $t = 0$ differential cross sections were extrapolated (independent of energy) from the most forward experimental points. Extrapolation errors are more important than experimental errors. PP, PC, and DP models were used, and an attempt was made to determine which parameters could be found by fitting. (France)

46631 COMPLEX CONJUGATE REGGE POLES. Roy, D. P. (Rutherford High Energy Lab., Chilton, Eng.). pp 309-23 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

The mechanism of pole-cut collision is studied using multi-peripheral unitarity. The general features of the resulting complex-conjugate poles are examined, the phenomenological implications of these complex conjugate poles being envisaged. Being on the unphysical sheet, these poles are demonstrated not to act directly on the scattering cross section but to control it indirectly through a bump of the Breit-Wigner type in the cut discontinuity. (France)

46632 K^+p PHASE SHIFT ANALYSIS AND EXOTIC RESONANCES. Yokosawa, A. (Argonne National Lab., Ill.). pp 325-36 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

New data concerning K^+p differential elastic cross sections and polarization (some at backward angles) were analyzed. The total number of data points is approximately twice that used in previous analysis, the incident momentum being in the 0.52 to 1.89 GeV/c range. A solution was obtained between 0.52 and 1.21 GeV/c and two other solutions were obtained between 1.37 and 1.89 GeV/c. Partial wave amplitudes are given for each of them. A resonant-like behavior in the P_1 partial wave was observed for each solution. (France)

46633 PHENOMENOLOGY OF TOTAL CROSS SECTIONS AND FORWARD SCATTERING AT HIGH ENERGY. Barger, V. (Univ. of Wisconsin, Madison). pp 337-55 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Recent σ_t experiments at Serpukhov indicate that, above 30 GeV, modifications in the Regge pole picture are necessary. The deviations of the Serpukhov $\sigma_t(\pi N)$ and $\sigma_t(KN)$ data from the Regge pole expectations can be ascribed to several radically different forms of asymptotic growth with energy: Pomeranchuk theorem violation, complex Regge poles, infinitely rising σ_t , vanishing asymptotic σ_t . The asymptotic forms selected by ρ and ω exchanges are tested using $K_L \rightarrow K_S$ regeneration, charge exchange scattering and the real part/imaginary part ratio for the elastic forward scattering amplitudes. (France)

46634 SOME IMPLICATIONS OF A VIOLATION OF THE POMERANCHUK THEOREM. Finkelstein, J. (CERN, Geneva).

pp 357-72 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

Theoretical consequences of the assumption of a violation of the Pomeranchuk theorem are discussed for nonforward scattering. K^+p and K^+n total cross sections are examined, and data on K^0 regeneration are shown not to favor the assumption of an asymptotic inequality for particle and antiparticle total cross sections at high energy. (France)

46635 EXPERIMENTAL LOW-ENERGY BEHAVIOR OF DIPS AND VENEZIANO DUALITY. Odorico, R. (CERN, Geneva). pp 397-424 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

The phenomenological convenience in considering dip behavior at low energy is illustrated. The K^+p scattering, πp scattering, and Λ -production processes are examined. Simple empirical relations between dips and resonances are demonstrated. Low-energy dip behavior is apparently simple only when resonances are regularly arranged according to a unique Regge sequence with a constant spacing in mass squared. (France)

46636 REVIEW OF NEAR-FORWARD πN SCATTERING AMPLITUDES AT HIGH ENERGIES. Hoehler, G. (Univ., Karlsruhe, Ger.). pp 457-84 of High Energy Phenomenology. /Tran Thanh Van, J. (ed.). Paris; Laboratoire de Physique Theorique et Hautes Energies (1971).

From Conference of Moriond meeting on the phenomenology of weak and electromagnetic interactions; Meribel-les-Allues, France (7 Mar 1971). See CONF-710345-(Vol.2).

An analysis of πN scattering experimental data at high energies and small momentum transfer is presented. Fixed- t dispersion relations and charge-independence are used for interpolating and determining the forward amplitudes $C^{(\rho,0)}$ (imaginary and real parts) and their derivatives. High energy models and phase shift sets are tested. The $A^+ \rightarrow 0$ condition is preferred to helicity conservation, provided that it is compatible with the data, since it involves a smaller number of adjustable functions. (France)

46637 s-CHANNEL HELICITY AMPLITUDES WITHOUT POMERON FOR ELASTIC K^+p AND π^+p CHARGE EXCHANGE SCATTERING. Karasuno, T.; Minami, S. (Osaka City Univ.). Lett. Nuovo Cim.; 4: No. 13, 647-50 (29 Jul 1972).

Two models are examined for the helicity-nonflip and helicity-flip amplitudes without pomeron for π^+p charge-exchange scattering. In the first model, only the imaginary part of the (nonpomeron) Regge amplitude is peripheral; in the second, both the imaginary and real parts on the (nonpomeron) Regge amplitudes without helicity-flip are peripheral. Using phase shift solutions, it is shown that the first model is more promising than the second. (L.B.S.)

46638 $A_{1\rho\pi}$ COUPLINGS AND t-CHANNEL HELICITY CONSERVATION. Roy, T.; Roychowdhury, A.; Banerjee, S. N. (Jadavpur Univ., Calcutta). Lett. Nuovo Cim.; 4: No. 13, 595-600 (29 Jul 1972).

It is shown from an analysis of the $\pi N \rightarrow A_1 N$ interaction that the helicity conservation in the t -channel is in no way connected with the S- and D-wave contributions to the $A_{1\rho\pi}$ coupling constant. (L.B.S.)

46639 POMERON AND f EXCHANGE AMPLITUDES IN ELASTIC π^+p SCATTERING. Davier, M. (Stanford Univ., Calif.). Phys. Lett.; 40B: No. 3, 369-72 (10 Jul 1972).

The dual absorptive model of Harari is shown to explain consistently the s and t dependence of the sum of π^+p and π^-p elastic differential cross sections at high energy. The Pomeron amplitude shrinks at a rate similar to that of K^+p elastic scattering while the f amplitude is shown to be dominated by peripheral partial waves at a radius ~ 1 fm. Finally, the shrinkage and s dependence of the f amplitude follow a canonical Regge behavior. (auth)

46640 REMOVAL OF AMBIGUITIES FROM INELASTIC PHASE-SHIFT ANALYSIS. Burkhardt, H. (Univ. of Birmingham, Eng.). Nuovo Cim.; 10A: No. 2, 379-88 (21 Jul 1972).

It is shown that the ambiguities in principle of phase-shift analysis are removed if, in addition to the differential cross section, the discontinuity of the amplitude across the nearby cut in $\cos \theta$ is given; unitarity is only used to resolve a twofold discrete ambiguity. The extension to pion-nucleon scattering shows that differential cross section and polarization measurements, together

with the discontinuity, are sufficient to leave only a fourfold ambiguity for unitarity, or an R measurement, to resolve. Questions of numerical analysis are not discussed. (auth)

46641 LOW-ENERGY THEOREM FOR PION-NUCLEON SCATTERING FROM PION SOURCE DYNAMICS. Pradhan, T.; Khare, A. V. (Saha Inst. of Nuclear Physics, Calcutta). *Lett. Nuovo Cim.*, 4: No. 9, 369-72 (1 Jul 1972).

A low-energy theorem is derived for the spin nonflip, isospin-flip πN scattering amplitude in terms of the πN coupling constant. The derivation of the theorem follows from the algebra of pion sources in which the pion sources and the isospin generators form a $SU(2) \times SU(2)$ algebra. It is suggested that the low-energy theorem may be used as a definition of πN coupling constant just as the Thirring theorem in quantum electrodynamics is used to define renormalized charge. (L.B.S.)

46642 DUALITY-BREAKING EFFECTS IN A MODEL FOR MESON-BARYON SCATTERING. Gage, D. W.; O'Donovan, P. J. (Arizona State Univ., Tempe). *Nuovo Cim.*, 10A: No. 3, 560-72 (1 Aug 1972).

Duality-violating effects are examined in the context of an exchange-degenerate multiple-scattering model incorporating strong EXD for poles, SU_3 at the vertices and s-u crossing. Ten parameters, including d/f ratios, are used to simultaneously fit cross sections for $K^+p \rightarrow \pi^+\Sigma^+$, $\pi^+p \rightarrow K^+\Sigma^+$, $K^+p \rightarrow \bar{K}^0n$, $K^+n \rightarrow K^0p$, $\pi^+p \rightarrow \pi^0n$, and $\pi^+p \rightarrow \eta n$. The qualitatively different energy dependences of the HCEX and CEX line-reversed cross section differences arise naturally from the fact that the model requires equal absorption in CEX line-reversed reactions and unequal absorption in the HCEX reactions. The cross section fits are excellent for the HCEX and reasonable for the CEX reactions; polarizations are correct in sign. The d/f ratios obtained are close to the quark-model SU_6 values. The model is compared with other multireaction fits and their differences and relative successes are discussed. (auth)

46643 PHOTON EXCHANGE IN $\pi + N \rightarrow \rho + N$ REACTIONS. Zima, V. G. (Donetsk Physical-Technical Inst., USSR). *Ukr. Fiz. Zh. (Russ. Ed.)*, 16: No. 9, 1518-23 (Sep 1971). (In Russian).

It is shown that interference of photon exchange in the t-channel of the reaction $\pi^+ + N \rightarrow \rho + N$ with nucleon exchange in the s- and u-channel is a remarkable quantity. It is necessary to take into consideration nucleon exchange at the same time with one-pion exchange if ρ -meson is coupled to a conserved source. In the reactions $\pi^+ + p \rightarrow \rho^+ + p$ photon exchange has independent importance for $|t| < 0.01 \text{ GeV}^2$. (auth)

46644 MANY-NEUTRON RESONANCES. Karmanov, V. A.; Kondratyuk, L. A.; Shapiro, I. S. (Inst. of Theoretical and Experimental Physics, Moscow). pp 496-511 of *Problems of Modern Nuclear Physics*. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Certain aspects of the theory of nuclear resonances are examined, for example, the lightest Δ -nuclei which contain the isobar $\Delta(1236)$. The scattering amplitude of an unstable particle by a stable N may be determined using the properties of the three-particle amplitude $\pi NN \rightarrow \pi NN$. Models for an induced nonexchange interaction ΔN and an exchange interaction ΔN are discussed. An enumeration of states participating in the isotriplet and the isosinglet is made for two-baryon Δ -nuclei and three baryon nuclei. (K.S.W.)

46645 VIOLATION OF POMERANCHUK THEOREM AND RELATIONS BETWEEN THE DIFFERENTIAL CROSS SECTIONS. Rekalov, M. P. (Physical-Technical Inst., Kharkov). *Ukr. Fiz. Zh. (Russ. Ed.)*, 16: No. 9, 1450-5 (Sep 1971). (In Russian).

Consequences of the Pomeranchuk theorem violation are analyzed, assuming that the amplitudes of elastic hadron scattering at a zero degree become real values at asymptotic high energies. It is shown that the differential cross sections of the charge-exchange processes $\pi^- + p \rightarrow \pi^0 + n$, $K^+ + n \rightarrow K^0 + p$ and of the elastic processes (πN) - and (KN) -scattering must be of one order in this energy region; while the differential cross sections of the processes $\pi^+p \rightarrow \eta + n$, $\pi^+p \rightarrow X + n$ must be essentially less. Consequences of the quark model for the processes $\gamma + N \rightarrow N + V^0$ must remain valid even if the Pomeranchuk theorem is not valid. (auth)

46646 HELICITY CONSERVATION IN $\pi N \rightarrow A_1 N$ SCATTERING. Biyajima, Minoru (Tokyo Univ. of Education). *Progr. Theor. Phys. (Kyoto)*, 47: No. 5, 1601-9 (May 1972).

The s- and t-channel helicity conservation hypotheses in $\pi N \rightarrow A_1 N$ reaction are studied. It is shown that the main properties of

density matrices for this reaction can be explained well by using the t-channel helicity conservation hypothesis and the hypothesis of the Pomeron dominance. Requiring the factorization theorem for s-channel helicity conservation in the nucleon vertex, the following scattering amplitude is obtained; $T_{\pi N \rightarrow A_1 N} = (e \cdot q_\pi) K_{\mu\nu} \gamma_\mu u$. However, this amplitude does not mean the s-channel helicity conservation. (auth)

46647 URBARYON REARRANGEMENT AND INELASTIC TWO-BODY PROCESSES. Kobayashi, Akizo (Nagoya Univ., Japan). *Progr. Theor. Phys. (Kyoto)*, 47: No. 5, 1779-81 (May 1972).

The nonvanishing polarization and the breaking of line reversal relation in the differential cross sections for inelastic meson-baryon two-body processes are investigated on the basis of urbaryon rearrangement diagrams. Qualitative results for elastic, charge-exchange, strangeness-exchange, and double charge-exchange interactions are found to reproduce rather well the experimental data. (L.B.S.)

46648 COMMENT ON THE BACKWARD $K^-n \rightarrow \Lambda \pi^-$ SCATTERING. Seth, V. P. (International Centre for Theoretical Physics, Trieste, Italy). *Progr. Theor. Phys. (Kyoto)*, 47: No. 5, 1774-6 (May 1972).

The contributions from N_α and N_γ trajectories for the backward scattering process $K^- + n \rightarrow \Lambda + \pi^-$ are taken with the trajectories' parameters obtained from the Chew-Frautschi plot to analyze the differential cross section and polarization with residue functions as the unknown constant parameters. The residue parameters thus obtained from the best fit are used to predict the values of some coupling constants. (L.B.S.)

Strong (Baryon No. ≥ 2)

Refer also to abstracts: 46457, 46787, and 47100.

46649 (JINR-P2-6395) ANALYSIS OF INELASTIC COLLISIONS OF PARTICLES WITH NUCLEI AT VERY HIGH ENERGIES. Barashenkov, V. S.; Eliseev, S. M. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 15p. (In Russian). Dep. NTIS (U. S. Sales Only).

Experimental data on pion- and nucleon-nucleus interactions at the energies much more than 10 GeV are analyzed on the basis of the cascade-evaporation model taking into account the decrease of density of number of intranuclear nucleons in the process of the cascade development. The experiment and theory can be co-ordinated right up to $T \approx 100 \text{ GeV}$ within the measurement errors. At large energies the cascade multiplicity of shower particles noticeably exceeds the experimental values. The influence of the many-particle interaction effect is discussed. (auth)

46650 (KFK-1547) SIMPLE THEORY OF PROTON-PROTON FINAL STATE INTERACTION IN THE REACTION $d + p \rightarrow p + p + n$. Matthaei, H. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Experimentelle Kernphysik); Zeh, H. D. (Heidelberg Univ. (West Germany). Institut fuer Theoretische Physik). Mar 1972. 20p. Dep. NTIS (U. S. Sales Only).

The amplitude for deuteron break-up in deuteron-proton collision including Coulomb final state interaction (in all partial waves) is derived analytically. The calculation is based on the assumption of a two-step reaction with break-up mechanism of zero range. Nuclear final state interaction is taken into account under the same approximations in s-waves. Comparison with experiments indicates a clear improvement compared to the Watson-Migdal theory. (auth)

46651 (RPP/T-18) AN ESTIMATE OF REGGE CUTS FROM INCLUSIVE FINITE MASS SUM RULES. Roy, D. P.; Roberts, R. G. (Rutherford High Energy Lab., Chilton (England)). 1972. 16p. Dep. NTIS (U. S. Sales Only).

Inclusive data on $K^+ \frac{p}{2} K^0$ and $K^- \frac{p}{2} \bar{K}^0$ are used to calculate the $\rho\rho$ and A_2A_2 Regge cuts in forward nucleon-nucleon scattering. Although the energies for which data exist are too low to allow a precise estimate, an upper bound for the cut magnitude can be obtained which is at the level given by the eikonal model. (auth)

46652 TWO-PION EXCHANGE AN POTENTIAL. Riska, D. O. (State Univ. of New York, Stony Brook). *Phys. Lett.*, 40B: No. 2, 177-80 (26 Jun 1972).

The two-pion exchange contribution to the ΛN potential is calculated from the field theoretic scattering amplitude. The effect of the Adler consistency condition for the $\pi\Lambda$ and πN amplitudes is taken into account, as well as the s-wave $\pi\pi$ interaction. (auth)

46653 RECOIL EFFECT IN QUASI-ELASTIC HADRON-DEUTERON SCATTERING. Alberi, G. (Univ., Trieste, Italy. Istituto Nazionale di Fisica Nucleare, Trieste, Italy); Gregorio, M. A.; Seth, V. P. *Nucl. Phys.*, B44: No. 2, 437-44 (1972).

The problem of hadron-deuteron quasi-elastic scattering is considered for large momentum transfers. The longitudinal part of the momentum transfer is included in the single and double scattering. It is shown that a simple enhancement factor relates the double scattering contribution to the differential cross section to the same quantity in Glauber theory. (auth)

46654 STRONG-INTERACTION FORM FACTOR OF PROTON IN HIGH-ENERGY PROTON-PROTON ELASTIC SCATTERING. Kato, S. (Hiroshima Univ.). *Nuovo Cim.*; 10A: No. 1, 48-52(1 Jul 1972).

An analysis of high-energy proton-proton elastic-scattering data is made in terms of one-particle exchange type amplitude. The results show that the proton form factor is constant at large momentum transfer. This may be interpreted as evidence for the existence of a pointlike core in the proton. (auth)

46655 SPIN STRUCTURE OF pd SCATTERING IN GLAUBER THEORY. Alberi, G. (Univ., Trieste, Italy. Istituto Nazionale di Fisica Nucleare, Trieste, Italy); Bertocchi, L.; Gregorio, M. A. *Nuovo Cim.*; 10A: No. 1, 37-47(1 Jul 1972).

The differential cross section, polarization and asymmetry, measured in recent proton-deuteron elastic scattering experiments, were computed using the Glauber multiple-scattering theory. The nucleon-nucleon amplitudes are obtained from recent phase-shift analyses; results from different sets of phase shifts are compared. (auth)

46656 FURTHER RESULTS FROM MULTI-REGGE MODELS APPLIED TO THE $pp \rightarrow pp\pi^+\pi^-$ REACTION AT 13.1 GeV/c. Le Guyader, J. (Institut de Physique Nucleaire, Paris); Sene, M.; Ginestet, J.; (and others). *Nuovo Cim.*; 10A: No. 3, 573-95(1 Aug 1972).

The $pp \rightarrow pp\pi^+\pi^-$ reaction at 13.1 GeV/c is analyzed in terms of the two multi-Regge models proposed by Chan et al., and by Berger et al. In the first model (CLA model) the introduction of the Δ trajectory exchange is discussed and a comparison is made with the experimental results. In the second model, which is a Reggeized pion-exchange model, the π trajectory has a hyperbolic form. The theoretical curves give a satisfactory representation of most of the distributions such as longitudinal momenta or effective-mass spectra. However, some discrepancies remain and are discussed. A comparison is made with the results previously obtained at 8 GeV/c. (auth)

46657 INCLUSIVE REACTIONS AND THE ASYMPTOTIC BEHAVIOR OF PROTON-PROTON COLLISIONS. Minakata, H. (Hiroshima Univ.). *Lett. Nuovo Cim.*; 4: No. 13, 593-7 (29 Jul 1972).

The asymptotic behavior of p-p inclusive interactions is studied, where the inclusive cross section is regarded as the totality of the inelastic processes. The inelastic cross section is written in terms of the one-particle distribution function (OPDF) and a dipole and a monopole fit to the OPDF are examined. (L.B.S.)

46658 EXPERIMENTAL SITUATION IN THE INVESTIGATION OF MANY-NUCLEON RESONANCES. Leksin, G. A. (Inst. of Theoretical and Experimental Physics, Moscow). pp 511-31 of *Problems of Modern Nuclear Physics*. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

A possible contradiction between the concepts of isonuclei and many-baryon resonances is explored. Well-known methods used in the study of boson and baryon resonances and in the analysis of di-baryon systems have led to the investigation of: energy dependence of total scattering cross sections; energy dependence of the differential cross section of a canal; the number of reaction occurrences as a function of the invariant mass of the system; and the number of reaction occurrences as a function of the "deficient mass," which is essentially the invariant mass of the system. Each of these problems is discussed. (K.S.W.)

46659 REGGE PHENOMENOLOGY OF INCLUSIVE SPECTRA IN THE CENTRAL REGION. Chan, Hong-mo (Rutherford High Energy Lab., Chilton, Eng.); Miettinen, H. I.; Roy, D. P.; Hoyer, P. *Phys. Lett.*; 40B: No. 3, 406-10(10 Jul 1972).

In the light of new data from the ISR, initial steps are taken towards the Regge analysis of inclusive spectra in the central region. It is shown that some features in the data necessitate the introduction of a further singularity in addition to the Pomeron and the meson trajectories ρ - f - ω - A_2 . The resulting model yields many predictions via duality and factorisation which can easily be tested at ISR and Batavia energies. (auth)

46660 SIMPLE DESCRIPTION OF HIGH ENERGY ELASTIC SCATTERING. Kane, G. L. (Rutherford High Energy Lab., Chilton, Eng.). *Phys. Lett.*; 40B: No. 3, 363-8(10 Jul 1972).

Motivated by ideas about the structure of high energy amplitudes in impact parameter space, a simple form (equation six) is suggested for the dominant elastic amplitude (Pomeron exchange in Regge language). It is imaginary at $t = 0$ but has a significant real part away from $t = 0$ even at very high energies. It is able to describe the experimental features of all elastic scattering to the highest ISR energies, including the increase in slope for $-t < 0.15$ and considerable shrinkage in pp scattering for $-t < 0.15$ with little shrinkage for $0.2 < -t < 0.3$; the small angle slope increase means that even elastic hadron interactions have an important peripheral contribution. The elastic amplitude has the phase variation with t needed in recent work to understand the πN scattering amplitude at 6 GeV/c in terms of the absorption model; this phase variation cannot be simply parametrized by a Regge pole. Using this Pomeron also solves the long-standing shrinkage problem for the absorption model. (auth)

46661 TRIPLE REGGE COUPLINGS AND THE CENTRAL REGION OF INCLUSIVE DISTRIBUTIONS. Humble, S. (Daresbury Nuclear Physics Lab., Warrington, Eng.). *Phys. Lett.*; 40B: No. 3, 373-7(10 Jul 1972).

A model for inclusive distributions near $x = 0$ is derived in terms of a product of triple Regge couplings. The model predicts a cusp structure in the high energy distributions, and explains the slow approach to a scaling limit for the exotic $\bar{p}$ and k^- production from pp collisions in terms of their masses. (auth)

46662 REMARKS ON AVERAGE PION MULTIPLICITIES AT HIGH ENERGIES. Charlton, G. R.; Thomas, G. H. (Argonne National Lab., Ill.). *Phys. Lett.*; 40B: No. 3, 378-80(10 Jul 1972).

The ISR data on $pp \rightarrow \gamma$ imply a π^0 transverse momentum spectrum which is consistent with the average of the π^+ and π^- spectra, at the same energy, both in shape and in normalization. Therefore, the average numbers of produced pions per inelastic pp collision at the ISR can be compared with data at lower energies, and with cosmic ray results. (auth)

46663 REGGEON CONTRIBUTIONS IN A TWO-FIRE-BALL MODEL. Margolis, B.; Meggs, W. J. (McGill Univ., Montreal). *Lett. Nuovo Cim.*; 4: No. 10, 413-16(8 Jul 1972).

The role of low-lying trajectories in inelastic pp interactions is investigated within the framework of two-fireball model. It is found that the proton spectrum will be dominated by protons from the decay of fireballs that are produced by pomeron exchange whereas the pion spectrum will be dominated at very high energies by pions from the decay of less probable but more massive fireballs produced by Regge exchange. (L.B.S.)

46664 INTRANUCLEAR CASCADES IN LIGHT NUCLEI. Abidinov, O. B.; Barashenkov, V. S. (Joint Inst. for Nuclear Research, Dubna, USSR). *Acta Phys. Pol.*; B3: No. 3, 385-90(1972). (In Russian).

The model of intranuclear cascades taking into account the changes in the density of a target-nucleus as the cascade develops, is employed for light nuclei of carbon type. In the energy range $T \leq 1$ GeV, the density affects slightly the properties of cascade particles (the number of low-energy particles being somewhat decreased) but it influences very strongly the yield of various isotopes already at the energies $T = 300$ to 400 MeV. The isotope yield also depends on the cluster structure of a target-nucleus. (auth)

Particle Invariance Principles and Symmetries

General

Refer also to abstracts: 46508 and 46714.

46665 (CTH-ITP-71-14) INTERNAL QUANTUM NUMBERS AND HOMOGENEOUS SPACES. Sears, Richard (Chalmers Tekniska Hogskola, Goteborg (Sweden). Institutionen for Teoretisk Fysik). 18 May 1971. 39p. Dep. NTIS (U. S. Sales Only).

A study of the possibility of finding explicit representations of the Poincaré group, which are labelled by additional discrete quantum numbers, was initiated. The main objective was to examine the extent to which a formalism developed by Kihlberg may be generalized to the case of a larger spectrum-generated group. An unsatisfactory aspect was the fact that it was impos-

sible to maintain the isospin invariance of the interactions. (Sweden)

46666 (IC-72/49) CONJUGACY AND STABILITY THEOREMS IN CONNECTION WITH CONTRACTION OF LIE ALGEBRAS. Shah, K. Tahir (International Centre for Theoretical Physics, Trieste (Italy)). Jun 1972. 20p. Dep. NTIS (U. S. Sales Only).

Let $a \in \Gamma_n(\bar{g})$, the n -dimensional Grassmannian variety, and $\bar{g}$ be a Lie algebra of the dimension $m \geq n$. Some properties of the set $B = \bar{G}(a) - \bar{G}(a)$ where $\bar{G}(a)$ is the Zariski closure of the orbit of $G = GL(\bar{g})$ at the point a are studied. The group $GL(\bar{g})$ is the group of automorphisms of $\bar{g}$. The set B is shown to be the set of Lie algebras which are contractions of $\bar{a}$. $\bar{a}$ is a subalgebra of $\bar{g}$. The main results are the theorems on the conjugacy of contracted algebras and the stability of Lie algebra $\bar{a}$ under contraction. The conjugacy theorem relates the algebras in the set B . (auth)

46667 (ITF-71-117-P) RELATIVISTIC TRANSFORMATIONS OF THE ONE-PARTICLE WAVE FUNCTIONS. Faustov, R. N. (Akademiya Nauk Ukrainskoi SSR, Kiev. Institut Teoreticheskoi Fiziki). 1971. 14p. (In Russian). Dep. NTIS (U. S. Sales Only).

The properties of the wave functions of free particles with arbitrary spins with respect to the transformations of the Poincaré group are considered. (auth)

46668 EIGENVALUES OF THE INVARIANT OPERATORS OF THE UNITARY UNIMODULAR GROUP $SU(n)$. Partensky, A. (Univ. Claude Bernard, Lyon). J. Math. Phys. (N. Y.); 13: No. 5, 621-5 (May 1972).

By a purely infinitesimal method the eigenvalues of the Biedenharn's invariant operators from I_2 to I_6 for the $SU(n)$ group are derived. It is shown that the general formula, quoted from Racah's work by Baird and Biedenharn, to obtain the eigenvalues of the invariant operators, is valid only for the two first I_2 and I_3 operators and not for the higher order invariants. The correct values are given for the first invariant till I_6 . (auth)

46669 STRUCTURE OF RELATIVITY GROUPS. Lugiato, Luigi A. (Univ., Milan); Gorini, Vittorio. J. Math. Phys. (N. Y.); 13: No. 5, 665-71 (May 1972).

An attempt is made to define the class of equivalent reference frames in special relativity in terms of a purely kinematical characterization of the notion of free material point and using homogeneity of space and time, isotropy of space, and unidirectional time flow. The problem of finding the possible forms of the relativity group and the significance of a "space-time homogeneity axiom" which was used in previous papers are discussed. (auth)

46670 NON-RELATIVISTIC DE SITTER SPACE-TIME AND THE HARMONIC OSCILLATOR. Sudbery, A. (Univ. of York, Heslington, Eng.). Nucl. Phys.; B44: No. 2, 520-30 (1972).

A nonrelativistic contraction of the de Sitter group $O(3,2)$ is shown to act as a dynamical group for the three-dimensional harmonic oscillator in a way which preserves many of the features of the relationship between the Galileo group and the free particle. (auth)

46671 ALGEBRAIC STRUCTURE OF A CLASS OF SOLVABLE QUANTUM PROBLEMS. Ghirardi, G. C. (Univ., Trieste, Italy). Nuovo Cim.; 10A: No. 1, 97-120 (1 Jul 1972).

A new algebraic approach to the treatment of quantum problems is introduced, which explains completely the algebraic structure of all those systems whose eigenvalue equations can be reduced, by means of proper functional transformations, to the hypergeometric equation. The results obtained constitute therefore the natural extension of the program started previously. In particular, it is proved that the eigenfunctions of the simple but physically interesting differential equation are associated to the states of the irreducible representations $D(\frac{1}{2}, \frac{1}{2}[1+2g]^h + 1)(1=0, 1, 2, \dots)$ of the Lie algebra $A_1 \approx B_1 \approx C_1$, characterized by the fact that the eigenvalue of the generator J_3 of the algebra takes the constant value $\frac{1}{2}[1+2g]^h$ for the considered states. When consideration is given to the S-wave Schrödinger equation for two particles interacting through the Hulthén potential, it is proved that the energy levels are associated with a properly selected submanifold of the space carrying the irreducible representations $D(\frac{1}{2}, -1)$ or $D(\frac{1}{2}, -1)$ of the same algebra. Finally, a general procedure is introduced which allows an analogous treatment of all problems whose eigenvalue equations can be reduced to the hypergeometric equation. An interesting three-body problem is also exhaustively investigated from the algebraic point of view. (auth)

46672 NONDEGENERATE COMPLEMENTARY SERIES OF REPRESENTATIONS OF THE GROUP $SU_{2,2}$. Macfadyen, N. W.

(Cambridge Univ., Eng.). Nuovo Cim.; 10A: No. 2, 268-76 (21 Jul 1972).

Two series of unitary irreducible representations of the non-degenerate complementary series of $SU_{2,2}$ are constructed. They are closely related to the first and second principal series of representations. A third series of irreducible representations, akin to the zeroth (discrete) principal series is also examined; an invariant Hermitian functional is presented, but its positivity could not be determined, and hence the unitarity of the representation. (auth)

46673 MANY-BODY REALIZATION OF THE SCHROEDINGER ALGEBRA. Burdet, G.; Perrin, M. (Centre National de la Recherche Scientifique, Marseille). Lett. Nuovo Cim.; 4: No. 13, 651-5 (29 Jul 1972).

It was previously shown that the largest group of space-time transformations which leaves invariant the free Schrödinger equation is a 12-parameter Lie group called Schrödinger group S , which contains the Galilei group G and one 11-dimensional subgroup of $Aut(G)$. The realization of the Schrödinger algebra is given in terms of canonical variables of N particles; this shows that the unitary irreducible representations of the extended Schrödinger group $\tilde{S}$ can describe the free system of these N interacting particles, but this extension of the Galilei group strongly limits the form of the interaction between particles compatible with the invariance property under $\tilde{S}$. (L.B.S.)

46674 REMARKS ON SOME SYMMETRY PROBLEMS IN COSMOLOGY AND IN THE THEORY OF PARTICLES AND FIELDS. Wataghin, G. (Univ., Turin). Lett. Nuovo Cim.; 4: No. 13, 608-10 (29 Jul 1972).

The first group of remarks concerns epochs in which the evolution of the Universe can be described as a quasi-adiabatic expansion. For the zero-mass constituent, like neutrinos and photons, the symmetry problem of the conformal group, and in the 3-dimensional space, of the $SO(4)$ group arises. The second part is concerned with the cut-off problems and the commutation relations. (L.B.S.)

46675 $SU(3)$ UNITARY REPRESENTATIONS. AN ALGEBRAIC METHOD WITH BOSE OPERATORS. Arisaka, Nakateru (Tohoku Univ., Sendai). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1758-71 (May 1972).

The unitary representations of the $SU(3)$ group are investigated in the Fock space defined by the Bose creation annihilation operators $a^k, a_1, b_1^{\dagger}$ and $b^{k\dagger}$ ($k=1, 2, 3$). The projection operators are obtained that pick up the vectors in the space of the arbitrary unitary representations of the $SU(3)$ group out of the Fock space. By the use of those projection operators, the canonical basis vectors $|T, Y(p, q)\rangle$ are explicitly obtained without the $SU(3)$ C.-G. coefficients. Each normalized isoscalar factor of the $SU(3)$ C.-G. coefficient in the coupling of $D(p, 0)$ and $D(0, q)$ is given in terms of the $SU(2)$ C.-G. coefficient. It is seen that all the possible $SU(3)$ transitions between the vectors of the arbitrary irreducible representation spaces occur by applying the irreducible tensor operators in the Fock space. The extension of the Bose operator method to the $SU(N)$ group is performed by introducing Bose creation annihilation operators with totally antisymmetric indices. (auth)

46676 BREAKDOWN OF SYMMETRY IN THE QUANTUM THEORY OF THE NONLINEAR REALIZATION OF A GROUP ON ITS SUBGROUP. Okabayashi, Takao (Tokyo Univ.). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1714-21 (May 1972).

The intuitively expected violation of symmetry in the quantum theory of preferred field is formulated definitely by constructing the transformation laws of operators in the interaction picture and of the Fock spaces. The components of the current belonging to the homogeneous space are not conserved, and their expectation values exhibit a PCAC-like behavior. The self-energy of the preferred field is explicitly given in the lowest order of perturbation. (auth)

Applications (Electromagnetic and Weak)

Refer also to abstract: 46494.

46677 OPTIMISATION OF POSSIBLE TESTS OF THE VALIDITY OF TIME-REVERSAL. Logan, B. A. (Univ. of Ottawa). Nucl. Instrum. Methods; 102: No. 2, 353-6 (1972).

Measurements of the angular correlation of cascade gamma rays from the decay of an oriented nuclear state can be used to test the validity of time-reversal invariance. Reactions induced by polarized proton beams can be used to orient suitable nuclear levels. The characteristics of some possible reactions are dis-

cussed and the relative merits of several suitable cascades are analyzed. (auth)

46678 ALGEBRAIC APPROACH TO THE SOLUTION OF QUANTUM-MECHANICAL PROBLEMS INVOLVING THE INTERACTION OF A CHARGED PARTICLE WITH ELECTROMAGNETIC FIELDS. Beers, B.; Nickle, H. H. (Southern Illinois Univ., Carbondale). *Lett. Nuovo Cim.*; 4: No. 8, 320-2 (24 Jun 1972).

An algebraic approach to the solution of quantum-mechanical problems involving the interaction of a charged particle with various types of electromagnetic fields is described. The method is essentially based on an investigation of the algebra satisfied by the generators of space-time translations, where the usual generators are regarded as being deformed by the presence of the electromagnetic field. (L.B.S.)

46679 LORENTZ TRANSFORMATIONS FOR ELECTROMAGNETIC FIELD VECTORS AND THE MOTION OF CHARGES IN CONSTANT AND HOMOGENEOUS ELECTROMAGNETIC FIELDS. Tarnovskii, A. S. (Kuibyshev State Univ., USSR). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 63-7 (1971). (In Russian).

A surface was established for the region of velocities corresponding to Lorentz transformations which minimize the values of vectors of a specified electromagnetic field. The connection with the Lorentz transformations which do not alter the values of vectors of a given electromagnetic field is revealed. The behavior of field vectors in a system associated with a charge moving in a constant and homogeneous electromagnetic field is examined. Such an electric charge moves in such a manner that the electromagnetic field vectors are decreased, and approach their direction. The transition time was found for a state for which the field vectors in the system associated with the charge are mutually parallel. (tr-auth)

46680 POSSIBLE CPT VIOLATION AND THE $K_L \rightarrow \mu^+ \mu^-$ DECAY. Achiman, Y. (Univ., Heidelberg). *Lett. Nuovo Cim.*; 4: No. 10, 437-42 (8 Jul 1972).

The puzzle between the value of the conventional unitary lower bound on the rate of $K_L \rightarrow \mu^+ \mu^-$ and a recent experimental value is examined. Possible CPT violation in the absorptive part of $K_L \rightarrow 2\mu$ is examined as a way to resolve this puzzle. (L.B.S.)

Applications (Strong)

Refer also to abstracts: 46482, 46483, and 46611.

46681 (NYO-4076-19) WAVE PACKET REALIZATION OF LIGHTLIKE STATES. Domokos, G. (Johns Hopkins Univ., Baltimore, Md. Dept. of Physics). Jul 1971. Contract AT(30-1)-4076. 14p. Dep. NTIS.

A new realization of lightlike particle states is constructed. The realization considered corresponds to objects which form partially localized wave packets in the space dimensions transverse to the direction of the lightlike momentum. The states of the transverse motion are classified by the representations of an algebra $SU(1,1)$ constructed of the generators of the conformal group. It is conjectured that this realization is suitable for the formulation of the "parton"-model of hadrons. (auth)

46682 MATHEMATICAL CHARACTERIZATION OF THE BREAKING OF HADRONIC INTERNAL SYMMETRY. Abellanas, Lorenzo (Institut des Hautes Etudes Scientifiques, Bures-sur-Yvette, France). *J. Math. Phys. (N. Y.)*; 13: No. 7, 1064-9 (Jul 1972).

The adjoint action of a semisimple complex Lie algebra G singles out some singular directions with special properties. The directions of breaking of the hadronic internal symmetry groups, e.g., $SU(3)$ or $SU(3) \times SU(3)$, by the electromagnetic, semileptonic, and nonleptonic weak interactions, are of this type. They are Hermitian (Q, Y) or nilpotent (Cabibbo direction) elements in the corresponding complexified Lie algebra. Their mathematical characterization is provided by the adjoint action of G . In particular, they satisfy a generalized extremal property for every G -invariant polynomial. (auth)

46683 ARE QUARKS POSSIBLE? Dowker, J. S. (Univ. of Manchester, Eng.). *Lett. Nuovo Cim.*; 4: No. 8, 301-4 (24 Jun 1972).

The existence of $SU(2)$ quarks, e.g., nucleons, and the nonobservation of $SU(3)$ quarks are considered. It is pointed out that there is a difference between $SU(2)$ and $SU(3)$ which could account for the absence of $SU(3)$ quarks. This difference is connected with the topological properties of the groups. It is concluded that if elementary particles are quantized kinks on $SU(3)$ they can have only zero triality, giving the eightfold way; thus quarks are excluded. (L.B.S.)

46684 TENSOR MESON PHOTOPRODUCTION AND $SU(6)$ SYMMETRY. Kulish, Yu. V. (Physical-Technical Inst., Kharkov). *Ukr. Fiz. Zh. (Russ. Ed.)*; 16: No. 9, 1444-9 (Sep 1971). (In Russian).

Within the framework of the $SU(6)$ symmetry the processes $\gamma N \rightarrow (1^+/2) + 2^+$ and $\gamma N \rightarrow (3^+/2) + 2^+$ with the magnetic dipole γ -quanta are considered. It is assumed that the mesons with $J^P = 2^+$ belong to the representation 189 or 405 the $SU(6)$ group. The comparison with the experiment is given. (auth)

46685 MODEL FOR THE MIXING AMONG THE 70 BARYON STATES BASED ON EXCHANGE DEGENERACY. Chiu, C. B.; Mitra, A. N. (Univ. of Texas, Austin). *Phys. Lett.*; 40B: No. 3, 401-5 (10 Jul 1972).

A model for the mixing between the 1^d , 8^d , 8^s , and 10^d states with $J^P = 3/2^-$ in the $SU(6)$ 70, $L = 1$ supermultiplet based on exchange degeneracy conditions of the pseudoscalar meson-baryon scattering is discussed. (auth)

46686 COMMENT ON THE VIRASORO GROUP IN DUAL-RESONANCE MODELS. Ramond, P. (Yale Univ., New Haven). *Lett. Nuovo Cim.*; 4: No. 10, 422-4 (8 Jul 1972).

The finite transformations of the Virasoro group are identified to be maps of periodic functions into periodic functions which leave their average unchanged. In analogy with the $SU(2)$ little group, it is expected that for each observable momentum P_μ , one representation of the Virasoro group exists which classifies the physical states. (L.B.S.)

46687 QUARTET SCHEME OF HADRONS IN CHIRAL $U(4) \otimes U(4)$. Maki, Ziro; Maskawa, Toshihide; Umemura, Isao (Kyoto Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1682-1703 (May 1972).

A general scheme of fundamental quartets is reformulated on the basis of strict lepton-baryon symmetry, and treated in a context of chiral $U(4) \otimes U(4)$ invariance. The structure of spontaneous breaking of it accompanied by sixteen Nambu-Goldstone bosons is studied in detail. One of the main predictions of the model is the existence of a new 0^- -meson, χ , with the mass heavier by ~ 0.5 GeV than the charmed partners, which may be identified with new heavy particles recently reported by Niu et al. It decays predominantly as $\chi \rightarrow K + K + (\pi, \eta \text{ or } \eta')$ with small decay widths ($\sim$ a few MeV). Such an indication of the existence of new heavy particles is encouraging enough to push forward our new treatment of the fundamental quartets (quartons) as a natural generalization of the Sakata-Nagoya model. (auth)

Field Theories

General

46688 WEYL AND LYRA MANIFOLDS. Sen, D. K.; Vanstone, J. R. (Univ. of Toronto). *J. Math. Phys. (N. Y.)*; 13: No. 7, 990-3 (Jul 1972).

It is shown that Weyl's geometry and an apparently similar geometry suggested by Lyra are special cases of manifolds with more general connections. The difference between the two geometries and their relationships with Riemannian geometry are clarified by giving a global formulation of Lyra's geometry. Finally the outline of a field theory based on the latter geometry is given. (auth)

46689 EXACT SOLUTIONS TO THE YANG-MILLS FIELD EQUATIONS. Rosen, Gerald (Drexel Univ., Philadelphia). *J. Math. Phys. (N. Y.)*; 13: No. 5, 595-7 (May 1972).

New spatially localized solutions to the Yang-Mills classical field equations are reported. (auth)

46690 HULTHEN AND SCHWARZSCHILD POTENTIALS IN THE KLEIN-GORDON EQUATION. Calucci, G. (Univ., Trieste, Italy). Istituto Nazionale di Fisica Nucleare, Trieste, Italy; Ghirardi, G. C. *Nuovo Cim.*; 10A: No. 1, 121-31 (1 Jul 1972).

The Klein-Gordon equation is discussed and solved for the S-wave in the presence of external fields, for two interesting cases. In the first case the external field corresponds to a short-range potential of the Hulthén type. The second case deals with the Klein-Gordon equation in a spherically symmetric gravitational field, within the framework of general relativity. Some interesting limiting cases of the considered equations as well as of the corresponding energy spectra and S-matrices are discussed. (auth)

46691 N. N. BOGOLYUBOV'S TRANSFORMATION IN THE STRONG COUPLING THEORY. II. Solodovnikova, E. P.; Tavkhelidze, A. N.; Khrustalev, O. A. (Inst. of High Energy

Physics, Serpukhov, USSR). *Teor. Mat. Fiz.*; 11: No. 3, 317-30 (Jun 1972). (In Russian).

N. N. Bogoliubov's transformation considered earlier in the works is generalized to the case of the strong particle-field interaction invariant under arbitrary continuous group of transformations. (auth)

46692 ESSENTIALLY NONLINEAR SCALAR FIELDS AND THE SPACE-TIME GEOMETRY. Nguyen Van Hieu. *Teor. Mat. Fiz.*; 11: No. 3, 293-300 (Jun 1972). (In Russian).

The curving of the space-time in essentially nonlinear fields is studied. The problem of the metrics determination in the field with a given Lagrangian is discussed. It is shown that if the Lagrangian is somehow known beforehand then the measurement of the velocity of signals in the known fields makes it possible to check the predictions of the theory. Another interesting consequence of the curving of space-time is also considered: the universal interaction of all fields with a given essentially nonlinear field. (auth)

46693 Σ^{Pq} TENSOR ASSOCIATED WITH THE POLARIZATION QUADRIVECTOR Σ_r AND THE MAXWELL Π^{Pq} TENSOR ASSOCIATED WITH THE ENERGY-MOMENTUM QUADRIVECTOR π_r IN COMPLEX FIELDS. Durand, Emile (Univ., Toulouse). *Compt. Rend.*, Ser. B; 274: No. 13, 843-5 (27 Mar 1972). (In French).

The Σ^{Pq} tensor associated with the polarization quadrivector, σ_r , is defined. It is shown to be a symmetrical universe tensor. The corpuscular character of the plane wave associated with the photon is demonstrated (helicity, mean spin density). (France)

46694 REVISED NONSYMMETRIC UNIFIED FIELD THEORY. Klotz, A. H. (Univ. of Sydney). *Acta Phys. Pol.*; B3: No. 3, 341-54 (1972).

A further development of the Einstein-Kaufman, nonsymmetric unified field theory is discussed. Static, spherically symmetric solutions of the field equations are considered. It is shown that there does not exist a solution corresponding to a magnetic monopole. In the purely electric case, one of Papapetrou's solutions is recovered and a new "cosmological" solution is found in which the space-time metric is that of a flat Minkowski world but a diverging electric field is present. It is pointed out that the theory may be significant as an account of charged matter. (auth)

46695 CYLINDRICAL WAVE SOLUTIONS OF BONNOR'S AND SCHRÖDINGER'S UNIFIED FIELD EQUATIONS. Lal, K. B.; Singh, T. (Univ. of Gorakhpur, India). *Tensor*; 23: No. 2, 151-6 (May 1972).

The solutions of Bonner and Schroedinger field equations are found for the case when the electromagnetic field is weak and the case when the electromagnetic field is not necessarily weak. (J.F.P.)

Quantum Field Theory

Refer also to abstracts: 46499, 46514, 46524, 46669, and 46676.

46696 (CONF-710816-19) THIRRING MODEL IN TERMS OF CURRENTS: SOLUTION AND LIGHT CONE EXPANSIONS. Dell'Antonio, G. F. (Naples Univ. (Italy). Istituto di Fisica Teorica); Frishman, Yitzhak (Weizmann Inst. of Science, Rehovoth (Israel)); Zwanziger, Daniel (New York Univ., N. Y.). 1971. 51p.

From International symposium on electron and photon interactions at high energies; Ithaca, N. Y. (23 Aug 1971).

Exact expansions of operator products, in terms of C-number functions singular on the light cone and regular operators, are exhibited explicitly in the Thirring model. For the products $\psi_1(x)\psi_1^*(x')$ and $\psi_2(x)\psi_2^*(x')$ of fermion fields the expansion reduces to one term only, with the C-number function having a singularity on the light cone which depends on the coupling constant, and the regular operator depending only on the currents, which are free. The resulting formula allows one to calculate all Wightman functions in terms of current matrix elements and thereby provides a simple and complete solution to the Thirring model. The different charge sectors are realized as inequivalent irreducible representation spaces of the canonical current commutation relations, on which the charged field ψ acts as an intertwining operator. (auth)

46697 (DESY-72/20) CURRENTS, STRESS TENSOR, AND GENERALIZED UNITARITY IN CONFORMAL INVARIANT QUANTUM FIELD THEORY. Mack, G.; Symanzik, K. (Deutsches Elektronen-Synchrotron, Hamburg (West Germany)). Apr 1972. 53p. Dep. NTIS (U. S. Sales Only).

Matrix elements of internal symmetry currents and energy momentum density tensor are constructed in Migdal Polyakov conformal invariant bootstrap field theory. Their 3-point func-

tions satisfy Bethe Salpeter equations which determine any free coefficients that may still occur in the conformal invariant Ansatz. Ward identities are verified for all n-point functions. They imply correct equal time current commutation relations. A proof of generalized unitarity is also given. Various equivalent forms of the propagator bootstrap are discussed. Algebraic techniques also yield an eigenvalue equation for first order correction to the exactly conformal invariant theory, assuming the latter is Gell-Mann Low large momentum asymptote of a renormalizable finite mass theory. (auth)

46698 (ITF-71-128-P) ON THE CHOICE OF A FORM FACTOR IN A NONLOCAL QUANTUM ELECTRODYNAMICS. Efimov, G. F.; Mogilevskii, O. A. (Akademiya Nauk Ukrainskoi SSR, Kiev, Institut Teoreticheskoi Fiziki). 1971. 28p. (In Russian). Dep. NTIS (U. S. Sales Only).

The photon Green function in the approximation of "principal logarithmic diagrams" is investigated in a local quantum electrodynamics. It is assumed that the result of the summation of "principal diagrams" leads to a nonlocal theory. The conditions, under which the Green function derived is unique, are stated. It turns out that any diagrams of perturbation series calculated by means of the partially summed photon Green function are convergent. Within the framework of the proposed procedure the value $l = (1/2m) \exp \{-(3\pi/2\alpha)\} \approx 10^{-268}$ cm is obtained for an elementary length. (auth)

46699 (JINR-P2-6394) STUDY OF CORRECTIONS TO EIKONAL REPRESENTATION OF AMPLITUDE IN SCALAR FIELD THEORY. Barbashov, B. M.; Nesterenko, V. V. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 16p. (In Russian). Dep. NTIS (U. S. Sales Only).

Based on the approach, called the straight-line paths approximation, the estimate of the high-energy behavior of the first correction term to the eikonal formula in scalar field theory $L_{\text{int}} = g\psi^2\phi$ is found. The loss of retardation in the eikonal representation is discussed as well as the connection of this effect with the corresponding approximation of the nucleon propagators. In conclusion, the modification of the eikonal representation, which is due to the introduction of additional interaction of the type $\lambda\phi^4$ is considered. (auth)

46700 (JINR-P2-6437) OPERATOR METHOD OF SOLVING THE QUASIPOTENTIAL EQUATIONS AND A CONCEPTION OF STRAIGHT-LINE PATHS AT HIGH ENERGIES. Kuleshov, S. P.; Matveev, V. A.; Sisakyan, A. N.; Smondyrev, M. A. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 13p. (In Russian). Dep. NTIS (U. S. Sales Only).

The operator method of solving the quasipotential equations is developed. It is shown that in high energy limit this method, just as the well-known straight-line path approximation in the framework of functional integration methods, is a realization of the general conception of straight-line paths. (auth)

46701 (JINR-P2-6454) QUANTUM FIELD THEORY: "DRESSING" CONTRA DIVERGENCIES. Shirokov, M. I. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 47p. (In Russian). Dep. NTIS (U. S. Sales Only).

The relativistic theory of charged ψ and neutral ϕ scalar fields is considered. The interaction $\lambda\psi^*\psi\phi$ is local, space-time is four-dimensional. Instead of "bare" creation-destruction operators the "dressed" ones are introduced. They have the following main property: "dressed" vacuum and one-particle states must be stable. This property requires in particular that the total Hamiltonian as a function of "dressed" operators must not contain trilinear terms, i.e. terms similar to the interaction ones (terms involving four, five... operators enter instead). The main equation is obtained by equating to zero the sum of all trilinear terms. A reason is presented in favor of the assertion: the equation solutions (if they exist) cannot entail the self-energy divergencies. A necessary condition of existence of the other solutions is obtained. It claims that some integrals involved in the equation must converge nonuniformly and must have some asymptotic property. It is shown that this property can be realized. (auth)

46702 (JINR-P2-6474) LOWER UNBOUNDEDNESS OF THE HAMILTONIAN WITH SCALAR ELECTRODYNAMICS. Zastavenko, L. G. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 11p. (In Russian). Dep. NTIS (U. S. Sales Only).

A proof of the inconsistency of the conditions, the s-matrix is finite, and the Hamiltonian is bounded from below, is given in the scalar electrodynamics for small values of coupling constant e^2 . It follows from this that the ground state for small e^2 and perturbation theory do not exist. (auth)

46703 (JINR-P2-6482) ASYMPTOTIC ESTIMATES OF QUASIPOTENTIAL PARAMETERS. Goloskokov, S. V.; Matveev, V. A. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 11p. (In Russian). Dep. NTIS (U. S. Sales Only).

The limitations on the quasipotential parameters which do not violate the absolute inequalities for the physical values proved in the local quantum field theory were obtained. (auth)

46704 (N-72-19766) LORENTZ-INVARIANT FORMULATION OF CHERENKOV RADIATION BY TACHYONS. Jones, Frank C. (National Aeronautics and Space Administration, Greenbelt, Md. Goddard Space Flight Center). Mar 1972. 34p. (NASA-TM-X-65851; X-641-72-73). NTIS.

Previous treatments of Cherenkov radiation, electromagnetic and gravitational, by tachyons were in error because the prescription employed to cut off the divergent integral over frequency is not a Lorentz invariant procedure. The resulting equation of motion for the tachyon is therefore not covariant. The proper procedure requires an extended, deformable distribution of charge or mass and yields a particularly simple form for the tachyon's world line, one that could be deduced from simple invariance considerations. It is shown that Cherenkov radiation by tachyons implies their ultimate annihilation with an antitachyon and demonstrates a disturbing property of tachyons, namely the impossibility of specifying arbitrary Cauchy data even in a purely classical theory. (auth) (STAR)

46705 ALGEBRAIC SOLUTION FOR THE KÄLLÉN-PAULI STATE VECTORS IN THE V_0 SECTOR BY A CONGRUENCE TRANSFORMATION. Nelson, C. A. (Louisiana State Univ., Baton Rouge). J. Math. Phys. (N. Y.); 13: No. 7, 1051-6 (Jul 1972).

From the representation of Bolsterli of the V_0 sector physical states in a basis having singular integral equations of the separable type, a congruence transformation carries the representation into the Källén-Pauli Heisenberg field components. This solves the Källén-Pauli singular integral equation algebraically. The resulting state vectors are proven explicitly to be both complete and orthonormal and furnish a new Möller wave matrix. (auth)

46706 HIGHER ORDER PERTURBATION THEORY FOR EXPONENTIAL LAGRANGIANS: THIRD ORDER. Pohlmeier, K. (Inst. for Advanced Study, Princeton, N. J.). Commun. Math. Phys.; 26: No. 2, 130-48 (1972).

The vacuum expectation value of the time-ordered product of three exponentials of free massless fields is defined as a continuous linear functional over a suitable test function space using minimal singularity as a criterion. (auth)

46707 CONNECTION BETWEEN THE SPECTRUM CONDITION AND THE LORENTZ INVARIANCE OF $P(\phi)$. Streater, R. F. (Bedford Coll., London). Commun. Math. Phys.; 26: No. 2, 109-20 (1972).

It is proved that, for the $P(\phi)$, quantum field theory, the Wightman functions are Lorentz invariant if the energy-momentum spectrum lies in the forward light-cone. The ingredients of the proof are the following facts, established by Glimm and Jaffe: the field satisfies local commutativity, and also the estimates $\phi_V(f, t) \leq \text{const } \|f\|_1 (H_V + 1)$ and $\pi_V(g, t) \leq \|g\|_2 (H_V + 1)$, where V is a space cut-off, uniformly in V . (auth)

46708 TEST OF QUANTUM-ELECTRODYNAMICAL CORRECTIONS IN MUONIC ATOMS. Walter, H. K.; Vuilleumier, J. H.; Backe, H.; (and others) (CERN, Geneva). Phys. Lett.; 40B: No. 2, 197-9 (26 Jun 1972).

Energies of the 5g-4f muonic X-rays were measured in ^{nat}Hg and ^{203}Tl with an accuracy of about 25 eV. They are found to be smaller than theoretical energies including recently revised calculations of higher-order vacuum polarization, by about two standard deviations. (auth)

46709 MULTIPOLE RADIATION SOLUTIONS FOR A CLASS OF MASSIVE PARTICLE WAVE EQUATIONS. Hajj, Jack P. (Suffolk Univ., Boston); Weaver, David L. J. Math. Phys. (N. Y.); 13: No. 5, 640-5 (May 1972).

The multipole radiation solutions are derived for the integer and half-integer spin massive particle wave equations of Weaver, Hammer, and Good. The rest system and massless limits of the solutions are found, the latter limit being shown to coincide with previous work by the present authors. Finally, as an application, the massive particle description is second quantized in terms of the multipole solutions and compared with some other formulations. (auth)

46710 RECENT DEVELOPMENTS IN THE COMPARISON BETWEEN THEORY AND EXPERIMENTS IN QUANTUM ELEC-

TRODYNAMICS. Lautrup, B. E. (CERN, Geneva); Peterman, A.; de Rafael, E. Phys. Rep.; 3C: No. 4, 193-259 (1972).

A survey is given of three main topics in quantum electrodynamics: fundamental bound systems; anomalous magnetic moments and high energy experiments. The emphasis lies particularly in recent developments concerning the electron and muon anomalous magnetic moments. (auth)

46711 CHOICE OF FORM FACTORS IN NON-LOCAL QUANTUM ELECTRODYNAMICS. Efimov, G. V.; Mogilevskii, O. A. (Inst. for Theoretical Physics, Kiev). Nucl. Phys.; B44: No. 2, 541-57 (1972).

The photon Green function in the approximation of principal logarithmic diagrams is investigated in local quantum electrodynamics. It is supposed that the result of the summation of principal diagrams leads to a non-local theory. Conditions under which the obtained Green function is unambiguous are stated. It turns out that any diagrams of the perturbation theory series, calculated by means of the partially summed photon Green function, are convergent. Within the frame of the procedure proposed, the value for elementary length is obtained: $1 = (1/2m) \exp \{-3\pi/2\alpha\} \approx 10^{-268}$ cm. (auth)

46712 ANALYTIC RENORMALIZATION OF THE EXPONENTIAL INTERACTIONS: THE THREE-POINT TIME-ORDERED PRODUCT WITH MINIMUM LIGHT CONE SINGULARITY.

Daniel, M. (Oxford Univ.); Mitter, P. K. J. Math. Phys. (N. Y.); 13: No. 7, 1026-41 (Jul 1972).

A method of analytic renormalization is developed to define the three-point time-ordered product of massless fields of exponential type as a strictly localizable distribution. The uniqueness property, known for the two-point T-product, is verified for the three-point product, for a special choice of fine renormalization. It is characterized by minimum singularity on the light cone: There are no delta function type singularities concentrated on the surface $x_1 = x_2 = x_3$. (auth)

46713 EQUIVALENCE OF DRESSING TRANSFORMATIONS. Federbush, Paul (Univ. of Michigan, Ann Arbor). J. Math. Phys. (N. Y.); 13: No. 7, 977-9 (Jul 1972).

The equivalence of the representations of the Weyl algebra for $\phi_{1/2}^4$ in a box induced by the dressing transformation of Glimm and the unitary dressing transformation is studied. Equivalence is shown for the simplified model in which only the most singular portion of the interaction is kept. (auth)

46714 POINCARÉ AND TCP INVARIANCE IN THE DETERMINATION OF WAVE EQUATIONS FOR PARTICLES OF ARBITRARY SPIN. Seetharaman, M.; Mathews, P. M. (Univ. of Madras). J. Math. Phys. (N. Y.); 13: No. 7, 938-43 (Jul 1972).

A brief outline is given of different types of approaches to the derivation of relativistic wave equations for free particles of arbitrary mass m and spin s . It is pointed out that all such equations can be subsumed in a standard form involving wavefunctions transforming according to the representation $D(0, s) \otimes D(s, 0)$ of the Lorentz group. Generalizing earlier work employing this standard form, all such equations (in a standard form) which are invariant under the Poincaré group and the combined operation $\Theta \equiv \text{TCP}$ are determined. It is shown that the resulting equations are automatically invariant under T, C, and P separately. (auth)

46715 REPRESENTATIONS OF THE CCR IN THE $(\phi^4)_3$ MODEL: INDEPENDENCE OF SPACE CUTOFF. Eckmann, Jean-Pierre (Univ., Geneva). Commun. Math. Phys.; 25: No. 1, 1-61 (1972).

The algebra of observables for the renormalized ϕ^4 interaction in 3-dimensional space-time is constructed. It is shown that the von Neumann algebras associated with observables in a bounded region B are independent of the space cutoff which is used in the construction of a Hamiltonian for this interaction. This result is shown to be useful in the construction of a translation invariant ϕ^4 theory in three dimensions. It also gives a physical criterion for the equivalence of non-Fock representations of the canonical commutation relations. (auth)

46716 SOME REMARKS ON THE NULL PLANE DEVELOPMENT OF A RELATIVISTIC QUANTUM FIELD THEORY. Schlieder, S.; Seiler, E. (Max-Planck-Institut fuer Physik und Astrophysik, Munich). Commun. Math. Phys.; 25: No. 1, 62-72 (1972).

Conditions are given for the existence of field operators on so-called null planes, and some consequences of the necessary restriction of the test function space concerning Haag's theorem and the possibility of unitary mappings intertwining between free fields of different masses are discussed. Conditions under which a unitary representation of the dilatations in the null plane gives

rise to a unitary representation of the dilatations in Minkowski space are discussed. (auth)

46717 EQUATION CURL $W_\mu(x) = 0$ IN QUANTUM FIELD THEORY. Pohlmeier, K. (Inst. for Advanced Study, Princeton, N. J.). Commun. Math. Phys.; 25: No. 1, 73-86(1972).

In one time and arbitrarily many space dimensions necessary and sufficient conditions for the existence of a local operator solution of the equation $\partial_\nu \omega = W_\nu$ are obtained. The given local fields W_ν satisfy $\partial_\mu W_\nu - \partial_\nu W_\mu = 0$, and the spectrum of the two point function $(\Omega, W_\mu(x) W_\nu(y) \Omega)$ is assumed to have a mass gap. (auth)

46718 $(\phi^{2n})_2$ FIELD HAMILTONIAN FOR COMPLEX COUPLING CONSTANT. Rosen, Lon; Simon, Barry (Univ. of California, Berkeley. Princeton Univ., N. J.). Trans. Amer. Math. Soc.; 165: 365-79(Mar 1972).

Hamiltonians $H_\beta = H_0 + \beta H_I(g)$, where H_0 is the hamiltonian of a free Bose field $\phi(x)$ of mass $m > 0$ in two-dimensional space-time, $H_I(g) = \int g(x): P(\phi(x)): dx$ where $g \geq 0$ is a spatial cutoff and P is an arbitrary polynomial which is bounded below, and the coupling constant β is in the cut plane, i.e., $\beta \neq$ negative real are considered. It is shown that H_β generates a semigroup with hypercontractive properties and satisfies higher order estimates of the form $\|H_\beta N^r R_\beta^s\| < \infty$, where N is the number operator, $R_\beta = (H_\beta + b)^{-1}$, r a positive integer, and β , s , and b are suitably chosen. For any $0 \leq \theta < \pi$, R_β converges in norm to R_0 as $|\beta| \rightarrow 0$ with $|\arg \beta| \leq \theta$. Finally applications of these results are discussed and asymptotic series and Borel summability for various objects in the real β theory are established. (auth)

46719 METHOD OF GAUGE-INVARIANT REGULARIZATION. Ashmore, J. F. (International Centre for Theoretical Physics, Trieste, Italy). Lett. Nuovo Cim.; 4: No. 8, 289-90 (24 Jun 1972).

It is suggested that by using the dimensionality of space-time as a complex parameter as in analytic regularization, a gauge-invariant renormalization could be achieved in a fairly simple manner. This idea is demonstrated in the example of second-order vacuum polarization in quantum electrodynamics. (L.B.S.)

46720 GAUGE PROBLEM IN QUANTUM FIELD THEORY. Mathews, P. M. (Univ. of Madras). Lett. Nuovo Cim.; 4: No. 8, 298-300(24 Jun 1972).

It was proved by Strocchi (Phys. Rev., 162: 1429 (1967)) from the general axioms of quantum field theory that the electromagnetic field tensor $F_{\mu\nu}$ will have to vanish identically if one insists that the potentials A_μ should a) satisfy Maxwell equation as an operator equation, and b) transform strictly as a four-vector under transformations of the Poincaré group. This work was criticized by Han and Biedenharn, who claimed that the conclusions are not valid. Comments are made on these works. It is suggested that the reason for the failure to get a nontrivial theory when a) and b) are demanded in the context of the conventional axioms of quantum field theory can be traced to the fact that the representation of the Poincaré group according to which the massless vector field transforms is an indecomposable one. (L.B.S.)

46721 BOUND STATES IN QUANTUM ELECTRODYNAMICS. Roy, T. (Jadavpur Univ., Calcutta). Indian J. Phys.; 45: No. 7, 328-36(Jul 1971).

Scattering of electrons, atoms and molecules including the creation-annihilation processes is considered from the point of view of quantum electrodynamics. The separation of the interaction Hamiltonian in the Coulomb part and the rest in the Schrödinger picture yields results which can be subjected to perturbation expansion, yet giving bound states wherever required. Because of the commutation relations proper symmetrizations are obtained even in a complicated case. (auth)

46722 POSSIBLE ORIGIN OF FINITE QUANTUM ELECTRODYNAMICS FROM WEAK INTERACTIONS. Budini, P. (International Centre for Theoretical Physics, Trieste, Italy); Furlan, P. Lett. Nuovo Cim.; 4: No. 8, 305-11(24 Jun 1972).

A method for regularizing quantum electrodynamics with the use of nonpolynomial weak Lagrangians and the concept of a regularizing coupling constant rather than a regularizing mass are examined. The photon constituents are taken to be lepton-antilepton pairs. It is shown that the value of the regularizing coupling constant to give massless vector bound states is of the order of magnitude of the Fermi weak-coupling constant. It is concluded that the idea that electrodynamical interactions are generated by the weak ones and that a finite quantum electrodynamics is originated in this way seems to be a plausible one. (L.B.S.)

46723 EXAMPLE OF A COVARIANT FIELD THEORY OF ELEMENTARY PARTICLES. Efinger, H. J. (Univ. of Otago, Dunedin, New Zealand). Lett. Nuovo Cim.; 4: No. 8, 312-14 (24 Jun 1972).

A brief review is first given of Fierz's model of a soluble field theory, that certain field equations that result from an interaction of a scalar gravitational field with a massless scalar field can be solved exactly. The possibility of generalizing this model to encompass elementary particles of nonzero mass is examined. (L.B.S.)

46724 CASUALITY GROUP IN FINITE SPACE-TIME. Blasi, A. A. (Univ. Genoa. Istituto Nazionale di Fisica Nucleare, Genoa); Gallone, F.; Zecca, A.; Gorini, V. Nuovo Cim.; 10A: No. 1, 19-36(1 Jul 1972).

An axiomatic definition of the relativity group for a finite four-dimensional space-time is given. Causality is represented as a relation on space-time and the relativity group is defined to be the casual group, namely the group of all permutations of space-time which preserve the causal relation. This group is found to be linear (modulo translations) and to contain, along with transformations which are strictly analogous to the usual Poincaré transformations, also transformations which interchange the inner with the outer part of light-cones. These transformations, which seem to set up a favorable framework for the introduction of tachyons, are possible in the real case only for a two-dimensional space-time and have to be introduced ad hoc, while in the scheme which is proposed here they arise naturally. It is further found that in this model the analogue of the orthochronous relativity group of the conventional real case is trivial, since it contains no boosts. (auth)

46725 REGGE POLES FOR LARGE COUPLING CONSTANTS IN THE WICK-CUTKOSKY MODEL. Mueller, H. J. W. (Univ., Trier-Kaiserslautern, Ger.). Nuovo Cim.; 10A: No. 1, 76-86(1 Jul 1972).

Following an initial investigation of Cheng and Wu the solutions of the Wick-Cutkosky model of the Bethe-Salpeter equation are examined in the case of unequal masses in the region of large coupling constants. Two branches of the solution are given covering the entire range of the independent variable. The tight-binding anomalies associated with the ladder approximation are discussed and the Regge trajectories are calculated. It is found that the pomeron trajectory—interpreted as the leading singularity of the "potential" rather than the amplitude—is located at $J > 1$ if the coupling is sufficiently strong. (auth)

46726 FACTORIZATION IN A MULTIPERIPHERAL CONFIGURATION: CAN ONE CONSTRUCT DUAL AMPLITUDES ON MORE GENERAL FIELD THEORY VERTICES AND PROPAGATORS. Lebrun, J. P. (CERN, Geneva). Nuovo Cim.; 10A: No. 1, 132-44 (1 Jul 1972).

Observing a simple connection between the zero-mode contribution in the conventional N-point dual Born term for unit intercept (Virasoro case) and the $\lambda\phi^3$ field theory, the possibility of constructing a new class of dual amplitudes based on more general field theory couplings and propagators is studied. The resulting amplitudes would have a number of particles suppressed on the leading trajectory, the spin-J ground state in a given channel being built on a two-dimensional operator formalism ("rotational" modes) that treats the integer and half-integer angular momenta in a unified way. The main part of this paper is devoted to a detailed study of factorization on a multiperipheral configuration in terms of "translational" and "rotational" modes. As a preliminary application in the context of dual models of the formalism developed the consequences of a simple factorization ansatz for the vertex into a pure "spin" part and a harmonic-oscillator part are studied. (auth)

46727 CROSSING AND UNITARITY IN THE CONTEXT OF THE BETHE-SALPETER EQUATION. Becker, W.; Grosser, D. (Univ., Tuebingen, Ger.). Nuovo Cim.; 10A: No. 2, 343-61 (21 Jul 1972).

Starting with the crossing-symmetric Bethe-Salpeter equation an approximation scheme for the T-matrix is discussed that consists of a recursive set of linear equations the solutions of which satisfy all symmetry properties of the exact T-matrix. The first of these equations is also derived by functional methods. It is established that in the frame-work of field theory essentially only the exact T-matrix satisfies both crossing symmetry and elastic unitarity, and the way in which the solutions of the equations violate unitarity is discussed. (auth)

46728 METHOD OF CHARACTERISTICS FOR FUNCTIONAL-DERIVATIVE EQUATIONS. Dahmen, H. D. (Institut fuer Theoretische Physik, Heidelberg. Kernforschungszentrum, Karlsruhe, Ger.); Jona-Lasinio, G.; Tarski, J. Nuovo Cim.; 10A: No. 3, 513-20(1 Aug 1972).

The method of characteristics is adapted to first-order functional-derivative equations. The general solution is then found to

three such equations in mathematical physics. Topological questions are not considered. (auth)

46729 ANALYTIC RENORMALIZATION VIA CONTINUOUS SPACE DIMENSION. Cicuta, G. M.; Montaldi, E. (Univ., Milan. Istituto Nazionale di Fisica Nucleare, Milan). *Lett. Nuovo Cim.*; 4: No. 9, 329-32(1 Jul 1972).

An analytic approach to renormalization is considered for the parametric integral representation for a Feynman integral in a scalar field theory in a space-time with d dimension. The parametric integral representation is extended by allowing the dimension d to take complex values. The renormalized integral is then obtained from an analytic continuation in d . (L.B.S.)

46730 THREE TYPES OF RELATIVISTIC EQUATIONS FOR PARTICLES WITH NONZERO MASS. Fushchich, V. I. (Inst. of Mathematics, Kiev). *Lett. Nuovo Cim.*; 4: No. 9, 344-6(1 Jul 1972).

It is shown that for free particles and antiparticles with mass > 0 and the arbitrary spin $s \geq \frac{1}{2}$ there exist three types of nonequivalent relativistic equations. (L.B.S.)

46731 CONFORMAL INVARIANCE OF ZERO-MASS KLEIN-GORDON-TYPE EQUATION. Mariwalla, K. H. (Inst. of Mathematical Science, Madras). *Lett. Nuovo Cim.*; 4: No. 8, 295-7 (24 Jun 1972).

It was recently shown by Bracken that $D_{M=0}^{m,0}$ representations of the Lorentz group that satisfy Klein-Gordon-type equations are not covariant under transformations of the conformal group if $mn \neq 0$. It is pointed out that the proof does not hold if one takes into account additional symmetries and constraints usually associated with these fields. (L.B.S.)

46732 LORENTZ TRANSFORMATIONS: DERIVATION OF LINEARITY AND SCALE FACTOR. Hegerfeldt, G. C. (Univ., Goettingen, Ger.). *Nuovo Cim.*; 10A: No. 2, 257-67(21 Jul 1972).

A transformation connecting two inertial frames S and S' must transform uniform motion of material bodies into uniform motion. However, since c is an upper bound on velocities, physics merely states that straight lines in the light-cone (and not necessarily all straight lines as for projective transformations) go over into straight lines. Nevertheless, it is shown in a quite elementary way that this weaker property also implies linearity. Constancy of light velocity then gives in a standard way the Lorentz transformations up to a "scale factor" λ which can, in principle, depend on the particular Lorentz transformation Λ and translation a . By a very simple group-theoretical argument it is shown that $\lambda \equiv 1$. (auth)

46733 MEASUREMENT OF AN OPTICAL FREQUENCY AND THE SPEED OF LIGHT. Bay, Z.; Luther, G. G.; White, J. A. (National Bureau of Standards, Washington, D. C.). *Phys. Rev. Lett.*; 29: No. 3, 189-92(17 Jul 1972).

The measurement of the frequency of the 633-nm red laser line is reported. This is the first measurement of an optical frequency in the visible range without reference to the speed of light or to a measured wavelength. Combination of the optical frequency with the known wavelength yields c to an accuracy higher than previously known. This method demonstrates the practicability of a single-standard time-length measurement system unified via a defined value of the speed of light. (auth)

46734 SUPERLUMINAL FRAMES AND THE GROUP OF GENERALIZED LORENTZ TRANSFORMATIONS IN FOUR DIMENSIONS. Mignani, R. (Univ., Catania, Italy. Centro Siciliano di Fisica Nucleare e di Struttura della Materia, Catania, Italy). *Istituto Nazionale di Fisica Nucleare, Catania, Italy*; Recami, E.; Lombardo, U. *Lett. Nuovo Cim.*; 4: No. 13, 624-6(29 Jul 1972).

A new group G of Lorentz transformations in four dimensions generalized for both subluminal ($\beta < 1$) and superluminal ($\beta > 1$) velocities was previously introduced. Comments are given to further clarify the new group, along with comments on the physical laws for tachyons the equivalence of inertial frames. (L.B.S.)

46735 DISPERSION RELATIONS FOR THE VERTEX FUNCTION FROM LOCAL COMMUTATIVITY. II. TWO-DIMENSIONAL DISPERSION RELATIONS. Andersson, B. (Univ. of Lund, Sweden). *Commun. Math. Phys.*; 25: No. 4, 308-35(1972).

Representation formulas for the vertex function are derived which are valid for arbitrary complex values of two of the scalar variables inside the corresponding domain of holomorphy, while the third variable is evaluated in a neighborhood of positive real values (the corresponding physical-region singularities). The analyticity domains always include at least the corresponding "axiomatic" Källén-Wightman domain. Two alternative versions

are given. In the first formula the occurring weight functions are the boundary values of the vertex function along the real axes of the scalar variables. The values of the arguments are such that the boundary values can be expressed in terms of the causal respectively time-ordered functions, which in a well-known way via reduction technique are related to the different form factors of the underlying field theory. In the second formula the weight functions are just the on-mass-shell matrix elements of the field operators which completely describe the dynamics of the theory and e.g., contain explicit information (in their support properties) on the mass spectrum of the field theory. The assumptions behind the results are the very general physical assumptions of Källén and Wightman, i.e., essentially Lorentz-covariance, translation-invariance, "reasonable" mass- and energy-spectrum, and local commutativity. Some moderate integrability- and boundedness-properties of the vertex function are needed. It is shown that the formalism can be extended to cover all cases of at most polynomial increase of the vertex function in asymptotic directions inside the Källén-Wightman holomorphy domain. The kernel functions in the representation formulas can be explicitly seen to result in possible singularities only along the physical-region-cuts, i.e., from the contributions of the corresponding physical states, as well as along the more complicated Källén-Wightman boundary-surfaces of the holomorphy domain, which correspond to the so-called "anomalous cuts." The kernel functions in the second version above are closely connected to simple perturbation theoretical functions. (auth)

46736 DISPERSION RELATIONS FOR THE VERTEX FUNCTION FROM LOCAL COMMUTATIVITY. I. ONE-DIMENSIONAL DISPERSION RELATIONS. Andersson, B. (Univ. of Lund, Sweden). *Commun. Math. Phys.*; 25: No. 4, 283-307(1972).

Dispersion relations for the vertex function are derived which are valid when two of the scalar variables are arbitrary complex inside certain domains of the product of the complex planes and the third scalar variable is evaluated just below or just above the physical region-cut. The domains of validity of the dispersion relations for the complex variables are domains with three real dimensions and can be described as neighborhoods of the boundaries of the "axiomatic" analyticity region of Källén and Wightman. The discontinuity of the vertex function across the cut-surface in the third variable for such values of the remaining variables is expressed only in terms of the dynamical on-mass-shell matrix elements of the locally commuting field operators. (auth)

46737 RIGGED HILBERT SPACES IN QUANTUM FIELD THEORY: A LESSON DRAWN FROM CHARGE OPERATORS. Antoine, J.-P. (Univ., Louvain, Belg. Univ., Geneva). *Helv. Phys. Acta*; 45: 77-92(1972).

Motivated by the problem of defining a charge operator, a systematic study is made of several classes of rigged Hilbert spaces suitable for quantum field theory. The result is that (essentially) only one of them satisfies all the requirements, namely $H_1 \subset H \subset (H_1)'$, where H_1 is the well-known space of quasi-local states. It is then shown that, even in the case of a nonconserved current, the charge operator always exists as a continuous operator from H_1 into $(H_1)'$. (auth)

46738 NONPOLYNOMIAL LAGRANGIANS (HIGHER PERTURBATION ORDERS). Volkov, M. K. (Joint Inst. for Nuclear Research, Dubna, USSR). *Teor. Mat. Fiz.*; 11: No. 3, 273-87 (Jun 1972). (In Russian).

It is proved that the theory with nonpolynomial Lagrangian remains finite in higher perturbation orders. It is proved also that the S -matrix is unitary in the third perturbation order. The power of increase of amplitudes in an arbitrary perturbation order is found. (auth)

46739 EQUATIONS FOR THE MATRIX ELEMENTS OF THE EUCLIDEAN QUANTUM ELECTRODYNAMICS. Rebenko, A. L. (Inst. of Theoretical Physics, Kiev). *Teor. Mat. Fiz.*; 11: No. 3, 301-16(Jun 1972). (In Russian).

The equations for the S -matrix elements of Euclidean quantum electrodynamics are obtained. The generating operator of these equations is investigated. On the basis of an analogy between the interaction Hamiltonian and the generating operator one succeeds in proving (under some restrictions put on the form factor) the convergence of the perturbation series for the arbitrary values of coupling constant. (auth)

46740 NOTE ON THE CONSTRUCTION OF THE BOGOLYUBOV SCATTERING OPERATOR IN THE $(\phi^4)_2$ THEORY. Wreszinski, W. F. (Eidgenössische Technische Hochschule, Zurich). *Teor. Mat. Fiz.*; 11: No. 3, 331-6(Jun 1972). (In Russian).

A Bogoliubov scattering operator $S(g)$ is constructed for the

(ϕ^4)₂ theory in the restricted case where the space-time cutoff has a product structure. Unitarity and causality are verified for this $S(g)$. Some problems in generalizing these results to a larger space of test functions are raised. (auth)

46741 QUANTUM THEORY OF PARTICLES WITH SPIN ZERO AND ONE HALF IN EXTERNAL FIELDS. Seiler, R. (Univ. of Pittsburgh). *Commun. Math. Phys.*; 25: No. 2, 127-51(1972).

The unitary (pseudo unitary) time-evolution operator for a particle with spin half (zero) in an external time-dependent electromagnetic (scalar) field is used to generate a Bogoliubov automorphism on the algebra of the free in field. For the case of an electric external field (scalar field) a finite expression for Ω_{out} is given and the S -matrix constructed. The latter is unitary and implements the Bogoliubov automorphism. Theorems by Shale and Stinespring are rederived. (auth)

46742 IS REGULARIZATION NECESSARY IN VACUUM POLARIZATION CALCULATIONS? Bui-Duy, Q. (Univ. of Montreal). *Lett. Nuovo Cim.*; 4: No. 10, 407-12(8 Jul 1972).

It is shown that, to all orders in perturbation theory, the vector current associated with the vacuum of a charged-particle motion in a prescribed electromagnetic field can be completely defined with renormalization prescription alone, and that regularization is not necessary. The technique used is the proper-time method developed by Schwinger. (L.B.S.)

46743 EQUAL-TIME AND EQUAL LIGHT-CONE COMMUTATOR BETWEEN THE DILATATION CHARGE AND THE FIELD IN MASSIVE $\lambda\phi^4$ -THEORY. Bandelloni, G. (Univ., Genoa). *Istituto Nazionale di Fisica Nucleare, Genoa. Lett. Nuovo Cim.*; 4: No. 10, 417-21(8 Jul 1972).

The existence of the commutation (D, ϕ) between the field and the dilatation charge in a field theory perturbative model is discussed in the equal-time and in the equal-light-cone limit. The massive $\lambda\phi^4$ model is used in the treatment. The results show that the equal-time commutator, as a limit of an unequal-time one, has no meaning due to the occurrence of the four dimensional term in the divergence of the dilatation current. (L.B.S.)

46744 RELATIVISTICALLY COVARIANT FORMULATION IN NONLINEAR LAGRANGIAN THEORIES AND FACTOR-ORDERING PROBLEMS. Suzuki, Tsuneo; Hattori, Chuichiro (Nagoya Univ., Japan. Aichi Inst. of Tech., Toyota, Japan). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1722-42(May 1972).

Factor ordering problems in non-linear Lagrangian theories are discussed. It is shown that symmetrization of the Hamiltonian is necessary in order to get the covariant Dyson formula by considering the factor ordering. As a result, there appear terms proportional to $\hbar^2(\delta^3(0))^2$ in the effective Hamiltonian density. In the evaluation of divergent integrals appearing in non-linear theories it is also necessary to take care of the order of integrations. Following the prescriptions thus obtained, Charap's discussions of the soft-pion theorems in chiral dynamics are reexamined in each order of the perturbation expansion. It is concluded that the soft-pion theorems are maintained, order by order, in perturbation theory also under the condition that the factor ordering is completely taken into account. (auth)

46745 TRANSFORMATION PROPERTIES OF OPERATORS IN LOCAL FIELD THEORY. Kikuchi, Kozo (Osaka Univ.); Taguchi, Yukio; Yamamoto, Kunio. *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1783-5(May 1972).

The necessity of the normal dependent Hamiltonian in addition to the ordinary Hamiltonian in quantum electrodynamics in the two-dimensional space-time was previously pointed out, which causes the modification of the energy-momentum tensor and hence the generators of the inhomogeneous Lorentz group. It is investigated how the notion of transformation should be changed on account of the modification of the generators. (auth)

46746 FUNCTIONAL DEFINITION AND CALCULATION OF GLOBAL OBSERVABLES IN NONLINEAR SPINOR FIELD QUANTUM THEORY. Stumpf, H. (Univ., Tuebingen, Ger.). *Z. Naturforsch.*; 27a: No. 7, 1058-72(Jul 1972).

Nonlinear spinor theory contains unobservable field operators which cannot be identified with free field operators. Therefore for the comparison with experiment a theory of observables for nonlinear spinor fields is required. This theory is developed for global observables by means of a map into functional space, and leads to a functional quantum theory of nonlinear spinor fields. (auth)

46747 REGGEON PERTURBATION THEORY IN A ZACHARIASEN-LIKE MODEL. Ebert, D. (Humboldt Univ., Berlin). *Acta Phys. Pol.*; B3: No. 3, 355-61(1972).

Gribov's Reggeon graph technique is used to construct a simple Zachariasen-like model with singularities in the j -plane. Decomposing the scattering amplitude into its one-Reggeon-irreducible and reducible parts yields a natural definition of the propagator, the vertex function, the form factor and the renormalization constants of an intermediate Reggeon. (auth)

Gravitation and General Relativity

Refer also to abstracts: 46192, 46193, 46261, 46501, and 46626.

46748 (JINR-P2-6270) BASIC STATE OF A MANY-FERMION SYSTEM, COUPLED BY GRAVITATIONAL LINES. Zastavenko, L. G. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 13p. (In Russian). Dep. NTIS (U. S. Sales Only).

A quasi-classical system of Fermi particles linked by lines of gravitation allows an asymptotic solution by means of the Thomas-Fermi method. The boundary-value problem for a differential equation is solved, and an expression for the energy of the basic state of a system of fermions is developed. (K.S.W.)

46749 (ORO-3992-94) ON A POSSIBLE REPULSIVE INTERACTION IN UNIVERSAL GRAVITATION. Chiang, C. C.; Dykla, John J.; Hwang, A. E. (Texas Univ., Austin. Center for Particle Theory). Jun 1972. Contract AT(40-1)-3992. 7p. Dep. NTIS.

A recent solution of the Einstein field equations suggests the possibility of a universal repulsive component in gravity. The consequences of such a possibility are examined by considering macroscopic systems, and it is concluded that such a repulsive component in gravity is compatible with the present observational data. (auth)

46750 APPROXIMATE SOLUTION FOR THE STATIC, SPHERICALLY SYMMETRIC METRIC DUE TO A POINT CHARGED MASS IN BRANS-DICKE THEORY. Mahanta, M. N.; Reddy, D. R. K. (Indian Inst. of Tech., New Delhi). *J. Math. Phys. (N. Y.)*; 13: No. 5, 708-9(May 1972).

An approximate solution of the field equations of Brans-Dicke theory is obtained for a static, spherically symmetric metric due to a charged point mass which can be considered to be an analog of the Reissner-Nordstrom metric in Einstein's theory. (auth)

46751 BUBBLE-TIME CANONICAL FORMALISM FOR GEOMETRODYNAMICS. Kuchar, Karel (Princeton Univ., N. J.). *J. Math. Phys. (N. Y.)*; 13: No. 5, 768-81(May 1972).

A functional differential version of the ADM canonical formalism is proposed. The existence of a canonical transformation separating canonical variables into internal coordinates, energy-momentum densities, and two pairs of true dynamical variables is assumed. The evolution of dynamical variables is governed by functional differential Hamilton's equations. They satisfy certain integrability conditions ensuring the internal path independence of dynamical evolution. The change of dynamical variables along any spacelike hypersurface is given by their Lie derivatives. This allows an elimination of $3\alpha^3$ components of the Hamiltonian equation, leading to a functional differential Hamilton equation based on a single bubble time. The Hamilton-Jacobi theory is built along the same lines. The formalism is illustrated in the mini-phase-space of the cylindrical Einstein-Rosen wave. (auth)

46752 STRUCTURE OF THE GRAVITATIONAL FIELD AT SPATIAL INFINITY. Geroch, Robert (Univ. of Chicago). *J. Math. Phys. (N. Y.)*; 13: No. 7, 956-68(Jul 1972).

A formalism is introduced for analyzing the structure of the gravitational field in the asymptotic limit at spatial infinity. Consider a three-dimensional surface S in the space-time such that the initial data on S is asymptotically flat in an appropriate sense. Using a conformal completion of S by a single point Λ "at spatial infinity," the asymptotic behavior of fields on S can be described in terms of local behavior at Λ . In particular, the asymptotic behavior of the initial data on S defines four scalars which depend on directions at Λ . Since there is no natural choice of a surface S in a space-time, the dependence of these scalars on S is essential. The asymptotic symmetry group at spatial infinity, whose elements represent transformations from S to other asymptotically flat surfaces, is introduced. It is found that this group, which emerges initially as an infinite-dimensional generalization of the Poincaré group, can be reduced to the Lorentz group. A set of evolution equations is obtained: These equations describe the behavior of the four scalars under the action of the asymptotic symmetry group. The four scalars can thus be considered as fields on a three-dimensional manifold consisting of all points at spatial infinity. The notion of a conserved quantity at spatial infinity is defined, and, as an example, the expression for the energy-momentum at spatial infinity is obtained. (auth)

46753 CANONICAL DYNAMICS OF SPINNING PARTICLES IN GRAVITATIONAL AND ELECTROMAGNETIC FIELDS.

Kuenzle, H. P. (Univ. of Alberta, Edmonton). *J. Math. Phys.* (N. Y.); 13: No. 5, 739-44 (May 1972).

In terms of the canonical form and the connection form on the bundle of Lorentz frames P over a space-time manifold V , a presymplectic form ω is defined on P , which induces a Poisson bracket on the set of real valued functions on the phase space of the system representing a spinning particle in an exterior gravitational and electromagnetic field. This structure coincides with the unique Poincaré invariant one for the free particle. Moreover, the projections into V of the integral manifolds of the kernel of ω on P yield precisely the world lines of a spinning particle as obtained for the dipole approximation of Dixon's equations of motion for extended test bodies in general relativity. (auth)

46754 TACHYON DUST METRIC IN GENERAL RELATIVITY.

Foster, J. C. Jr.; Ray, J. R. (Clemson Univ., S. C.). *J. Math. Phys.* (N. Y.); 13: No. 7, 979-82 (Jul 1972).

Special relativity allows the possibility of a class of particles, called tachyons, which travel with speeds greater than the speed of light in vacuum. These particles have spacelike 4-velocities. Since tachyons have energy and momentum, they will contribute to the gravitational field through the energy-momentum tensor. One question then is what types of solutions to the Einstein field equations will tachyons yield. A metric is considered which admits a four-parameter isometry group. When this metric is used in the field equations using a dust energy-momentum tensor, solutions exist only for spacelike 4-velocity of the dust. These are interpreted as solutions for a tachyon dust. Exact solutions to the field equations are obtained. (auth)

46755 GRAVITATIONAL FIELDS WITH GROUPS OF MOTIONS ON TWO-DIMENSIONAL TRANSITIVITY HYPERSURFACES IN A MODEL WITH MATTER AND A MAGNETIC FIELD.

Shikin, I. S. (Moscow State Univ.). *Commun. Math. Phys.*; 26: No. 1, 24-38 (1972).

For gravitational fields with metrics which admit of groups of motions multiply-transitive on 2-dimensional space-like invariant varieties, the exact solutions of the Einstein gravitational equations are given for the case when the sources of the gravitational field are dust-like matter and a magnetic field. A magnetic field is oriented along a direction orthogonal to transitivity hypersurface. The solutions contain arbitrary functions. In the case of transitivity hypersurface of positive curvature and in the absence of a magnetic field, the solution is reduced to the Tolman spherically symmetric solution for dust-like matter. The conditions are studied under which the solutions with a magnetic field become asymptotically isotropic and approach the flat and the open Friedmann models. The case of transitivity hypersurfaces with signature $(+,-)$ is also considered. (auth)

46756 ENERGY-MOMENTUM DETERMINATION IN THE GENERAL THEORY OF RELATIVITY.

Zhelnorovich, V. A. (Moscow State Univ.). *Dokl. Akad. Nauk SSSR*; 201: No. 5, 1078-81 (11 Dec 1971). (In Russian).

The assumption is made that the components of a pseudotensor or of an energy-momentum complex in a continuous medium (a field) are in a closed region, bounded by a piecewise continuous surface. The ensuing mathematical development is related to equations of Einstein, Meller-Mitskevich, and Lichnerowicz. (K.S.W.)

46757 NATURE OF GRAVITATIONAL SYNCHROTRON RADIATION.

Chitre, D. M.; Price, Richard H. (Univ. of Utah, Salt Lake City). *Phys. Rev. Lett.*; 29: No. 3, 185-8 (17 Jul 1972).

Ultra-relativistic geodesic particle orbits in the Schwarzschild geometry produce radiation (GSR) with angular distributions like that of synchrotron radiation. The spectra of high-frequency scalar, electromagnetic, and gravitational GSR are compared to each other and to the spectrum of synchrotron radiation. The differences among the spectra are explained in terms of the shape of the effective potential for GSR and the inapplicability of geometric optics. (auth)

46758 VARIATIONAL PRINCIPLES AND SPATIALLY-HOMOGENEOUS UNIVERSES, INCLUDING ROTATION.

MacCallum, M. A. H.; Taub, A. H. (Univ. of California, Berkeley). *Commun. Math. Phys.*; 25: No. 3, 173-89 (1972).

The validity of imposing spatial homogeneity on the variations in the usual action principle for Einstein's equations is studied. It is proved that with this procedure the standard and ADM Lagrangians give correct Einstein equations if and only if the space belongs to Class A of Ellis and MacCallum, i.e., the structure constants of the simply transitive group satisfy $C_{fg}^i = 0$. The possibility of overcoming the difficulty in the Class B spaces is examined. (auth)

46759 HOMOGENEOUS ANISOTROPIC SOLUTIONS OF EINSTEIN EQUATIONS WITH A COSMOLOGICAL TERM.

Aman, E. G. (Moscow State Univ.). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 12, 117-20 (1971). (In Russian).

A homogeneous, anisotropic model of the universe was obtained with planar, attendant space, filled with a powdery substance, taking into account the Λ -term. The presence of this term, independently positive or negative, accelerates the isotropisation of the solution. (tr-auth)

46760 GRAVITATIONAL ENERGY EMITTED BY AN ISOLATED MATERIAL SOURCE.

I. Silenko, P. N. (Moscow State Univ.). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 12, 101-4 (1971). (In Russian).

The solution of the Einstein field equation was examined for the case of axial symmetry, obtained by van der Brugge in 1966. Assuming that space-time everywhere is void outside of the time-like cosmic tube, which contains all field sources, the point of interest here is the empty realm of space-time. Using the tetrad formalism, the energy and momentum of the system were found, and it was shown that a gravitational radiation exists within the system. (tr-auth)

46761 COSMOLOGICAL TERM, SCREENING OF GRAVITATION, AND THE HYPOTHESIS OF NEGATIVE MASSES.

Baranov, A. A. *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 118-20 (1971). (In Russian).

A cosmological model is proposed which introduces the concepts of particles of both positive and negative masses, and assists in explaining the phenomenon of interacting galaxies and the screening effect caused by the presence of negative mass. The term $\Lambda = \rho_0/2 \times 8\pi G/c^2$ emerges from the theory, where G is Newton's gravitational constant. The density of positive matter ρ_0 is assigned two different values for purposes of discussion: 5×10^{-29} and 5×10^{-31} g/cm³. Other constants derived from these values indicate that with density changes of positive matter within the generally accepted limits, the value of the masses of gas from minus-particles with a temperature -3°K varies from the mass of the electron to the mass of the muon. (K.S.W.)

46762 SOLUTION OF THE EINSTEIN EQUATIONS WITH A NULL BUNDLE.

III. Dozmorov, I. M. (Moscow Higher Technical School). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 76-81 (1971). (In Russian).

The transformation of a Kerr-Schild type of null bundle was employed for obtaining new, accurate solutions to Einstein-Maxwell equations from known stationary solutions in a vacuum. Among these is the Kerr solution which describes the gravitational field of a homogeneous, rotating ring. (tr-auth)

46763 SOLUTION OF THE EINSTEIN EQUATIONS WITH A NULL BUNDLE.

II. Dozmorov, I. M. (Moscow Higher Technical School). *Izv. Vyssh. Ucheb. Zaved., Fiz.*; No. 11, 68-76 (1971). (In Russian).

The possibility of obtaining new accurate solutions to Einstein equations in a vacuum, and to the Einstein-Maxwell equations already known solutions, using a transformation with a null type bundle, was explored. The change in algebraic properties was analyzed for such a transformation in space permitting a congruence of isotropic, geodesic, with null distortion and rotation. Concrete examples illustrating the new, accurate solutions are presented. (tr-auth)

46764 CHARACTERISTIC DEVELOPMENT OF TRAPPED SURFACES IN EINSTEIN-SCALAR SPACETIMES.

Demmie, Paul Norman. Pittsburgh; Univ. of Pittsburgh (1971). 164p. University Microfilms Order No. 72-18,105.

Thesis.

The concept of a trapped surface provides for a precise characterization of gravitational collapse which has proceeded beyond the point of no return. Since there is no compelling reason to assume that trapped surfaces always develop during gravitational collapse, an important problem arises concerning the determination of the data required to be given on a null hypersurface in order that they develop in certain spacetimes. An investigation based on the general theory of relativity of this problem of the characteristic development of trapped surfaces was undertaken in this thesis for Einstein-scalar spacetimes. The main results of this investigation are the conditions on the characteristic data sufficient for trapped surfaces to develop in both spherically symmetric Einstein-scalar spacetimes which either contain or do not contain a non-diverging null hypersurface. An important further result of this investigation is the evidence presented supporting a relationship between a certain piece of the characteristic data and angular momentum. (Diss. Abstr. Int., B)

46765 MANY-BODY FORCES AND THE EFFECT OF THE MATTER DISTRIBUTION IN THE UNIVERSE TO THE GRAVITATIONAL CONSTANT. Hida, K. (Tokyo Univ.). *Lett. Nuovo Cim.*; 4: No. 10, 398-400 (8 Jul 1972).

The importance of the effects due to gravitational many-body forces on the observed gravitational constant is pointed out. The three-body problem is considered and the effects are discussed within the framework of the Minkowski space, considering curvature effects as interactions. The contributions from the Earth, the Sun, the Galaxy, and the Universe for the test bodies on the surface of the earth are estimated. Implications for a model of the expanding Universe are indicated. (L.B.S.)

46766 GAUGE TRANSFORMATION AND GRAVITATIONAL POTENTIALS. Hida, Kichiro; Okamura, Hiroshi (Tokyo Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1743-57 (May 1972).

It is believed in particle physics that the velocity-dependent part of potential is, in general, ambiguous as far as it is derived from S-matrix. The most general form of the graviton propagator is considered under an arbitrary q-number gauge transformation for the graviton field. The propagator depends on twelve arbitrary functions of k^2 , k being the space part of the momentum of the graviton. The arbitrariness of the gauge functions can be used to remove the ambiguity of one-graviton exchange potential up to the order of $(G/r)(v/c)^4$, G being the gravitational constant. The potential thus obtained depends only on one gauge parameter, say x . The perihelion motion of two-body system is gauge independent, although the potentials in the order of $(G/r)(v/c)^2$ and G^2/r^2 depend on the gauge parameter x . The potentials are derived from those in a fixed gauge parameter x by a coordinate transformation. (auth)

46767 MODEL FOR PARTICLES WITH SPIN IN THE GRAVITATIONAL AND ELECTROMAGNETIC FIELD. Duval, Christian; Fliche, Henri-Hugues; Souriau, Jean-Marie (Univ., Aix-Marseille). *Compt. Rend., Ser. A*; 274: No. 13, 1082-4 (27 Mar 1972). (In French).

A model is constructed for particles with spin. The model is consistent with experimental results and the principles of symplectic mechanics and general relativity. It contains an arbitrary function $m(\omega)$ which may be experimentally determined. (France)

46768 ELASTIC EXCITATIONS EXCITED BY GRAVITATIONAL WAVES. Papapetrou, A. (Univ., Paris). *Ann. Inst. Henri Poincaré, Sect. A*; 16: No. 1, 63-78 (1972). (In French).

The general relativistic equation of the motion of a continuous material distribution combined with the relativistic equation of elasticity leads to the equation of elastic waves which determines the vibrations of an elastic body induced by a gravitational wave. (France)

46769 INTEGRATION OF THE NEWMAN-PENROSE EQUATIONS IN A MINKOWSKI SPACE. Pons, Roland (Univ., Paris). *Ann. Inst. Henri Poincaré, Sect. A*; 16: No. 1, 51-62 (1972). (In French).

The Newman-Penrose equations are integrated in Minkowski space. The Bondi-Metzner transformations are studied in a Minkowski space having axial symmetry and the radial Newman-Penrose coordinate. (France)

46770 COVARIANT CANONICAL EQUATIONS OF THE TEST-PARTICLE MOTION IN GENERAL RELATIVITY. Pyragas, Kazimieras A. (Inst. for Theoretical Physics, Kiev). *Tensor*; 23: No. 2, 241-9 (May 1972).

The canonical equations of motion are derived in a covariant form by a direct generalization of the Hamiltonian variational principle. Also the notion of correspondence of the points on a set of curves is discussed. (J.F.P.)

46771 SPECIAL CLASS OF STATES OF TOTAL PURE RADIATION IN GENERAL RELATIVITY. Zund, J. D. (New Mexico State Univ., Las Cruces). *Tensor*; 23: No. 2, 144-6 (May 1972).

A special class of solutions of the Lichnerowicz radiation equations and the Maxwell equations for a singular source-free electromagnetic field are shown for a nonspecial recurrent space-time. (J.F.P.)

46772 NEUTRINOS IN SPHERICALLY SYMMETRIC GRAVITATIONAL FIELDS. Audretsch, J. (Queen Elizabeth Coll., London). *Lett. Nuovo Cim.*; 4: No. 9, 339-43 (1 Jul 1972).

The solutions ψ of the Weyl equation for neutrino interactions with gravitational fields are studied in the general spherically symmetric space-time without restrictions on the metric. It is shown that in any spherically symmetric space-time, the Weyl equation describing neutrinos has no solution apart from the trivial one ($\psi \equiv 0$) under certain constraints. (L.B.S.)

Scattering Theory

Refer also to abstracts: 46699, 46727, 46738-46741, 46747, 47068, and 47284.

46773 (CTH-ITP-71-17) BREAKDOWN OF SIMPLE CAN-CELLATION IN THE OPERATOR MULTIPLE SCATTERING SERIES. Karlsson, B. R.; Namyslowski, J. M. (Chalmers Tekniska Hogskola, Goteborg (Sweden). Institutionen for Teoretisk Fysik). 19 Jul 1971. 6p. Dep. NTIS (U. S. Sales Only).

The Watson multiple scattering series is considered for a three-body problem in the eikonal approximation. The sum of the series, i.e. the off-shell Faddeev-Lovelace transition amplitude, is expressed entirely in terms of off-shell two-body transition amplitudes. The result shows that the simple cancellation that occurs in the eikonal multiple scattering series for the on-shell amplitude breaks down for the off-shell amplitude. (auth) (Sweden)

46774 (ITF-72-44-E) THRESHOLD BEHAVIOUR OF DAMA. Bugrii, A. I.; Enkovskii, L. L.; Kobylinskii, N. A. (Akademiya Nauk Ukrainskoi SSR, Kiev. Institut Teoreticheskoi Fiziki). 1972. 11p. Dep. NTIS (U. S. Sales Only).

The threshold behavior of a dual amplitude with Mandelstam analyticity (DAMA) is investigated. The behavior of the trajectory associated with the amplitude is consistent with the threshold behavior following from the potential scattering and also from unitarity and the Mandelstam representation. (auth)

46775 (JINR-P2-6393) S-MATRIX UNITARITY IN THEORIES WITH NONPOLYNOMIAL LAGRANGIANS. Volkov, M. K. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 18p. (In Russian). Dep. NTIS (U. S. Sales Only).

Unitarity of S matrix in the third order of perturbation theory in G in nonpolynomial theories is proved. The role of anomalous singularities is cleared up. The scattering amplitude in the fourth order in G is also considered. (auth)

46776 (JINR-P2-6442) INVESTIGATION OF LARGE ANGLE SCATTERING IN QUASIPOTENTIAL APPROACH. Goloskokov, S. V. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 15p. (In Russian). Dep. NTIS (U. S. Sales Only).

The solution of the quasipotential equation in the quasipotential approximation was found. The estimates of the large-angle scattering amplitude at high energies for the exponential-type quasipotential were obtained. (auth)

46777 PROBABILISTIC INTERPRETATION OF THE QUANTUM SCATTERING CROSS SECTION. Babbitt, Donald G. (Univ. of California, Los Angeles). *J. Math. Phys. (N. Y.)*; 13: No. 5, 637-40 (May 1972).

The probabilistic interpretation of the quantum scattering cross section in the case of potential scattering is discussed in terms of Poisson random measures on the impact parameter plane and the sphere of outgoing directions. (auth)

46778 SOME FORMAL DEVELOPMENTS ON THE EIKONAL TREATMENT OF PRODUCTION PROCESSES. Calucci, G. (Univ., Trieste, Italy. Istituto Nazionale di Fisica Nucleare, Trieste, Italy). *Nucl. Phys.*; B44: No. 2, 629-38 (1972).

A treatment of the eikonal formalism for production processes is presented in which a formal solution for the collision problem, analogous to the S-matrix construction in the interaction picture, is obtained. Examples in which the formal solution allows a non-perturbative calculation of the transition amplitudes are shown. (auth)

46779 PRACTICAL RELATIVISTIC THREE-PARTICLE THEORY WITH ARBITRARY PARTICLE INTERACTIONS. Weiss, U. (Univ., Stuttgart). *Nucl. Phys.*; B44: No. 2, 573-93 (1972).

Relativistic three-particle equations are established by the rules of Blankenbecler and Sugar, which put the intermediate particles on the mass shell. By use of the Wightman-Garding momenta the correct cluster properties and time-reversal invariance are preserved and a spurious $s = 0$ singularity is avoided. A reduction to effective multichannel two-particle equations which is not restricted to separable potentials is discussed. The approach admits the systematic application of perturbation theory to relativistic composite particle scattering problems. (auth)

46780 TIME-DEPENDENT AND TIME-INDEPENDENT POTENTIAL SCATTERING FOR ASYMPTOTICALLY COULOMB POTENTIALS. Marchesin, Dan; O'Carroll, Michael L. (Pontificia Univ. Catolica, Rio de Janeiro). *J. Math. Phys. (N. Y.)*; 13: No. 7, 982-90 (Jul 1972).

The three-dimensional quantum mechanical problem of non-relativistic potential scattering is considered, where the Hamiltonian is $H_2 = H_1 + V = H_0 + V_C + V$; H_0 is the free particle Hamiltonian, V_C is the Coulomb potential, and $V(x)$ is a real-valued potential function defined for $x \in \mathbb{R}^3$. The existence of the wave operators $W_{\pm}(H_2, H_D) = s\text{-}\lim_{t \rightarrow \pm\infty} e^{iH_2 t} e^{-iH_D t}$ on $L^2(\mathbb{R}^3)$ for $V = V_2 + V'$ is shown, where $V_2 \in L^2(\mathbb{R}^3)$, $V' \in L^\infty(\mathbb{R}^3) \cap L^p(\mathbb{R}^3)$ ($p < 3$), and $e^{-iH_D t}$ is the family of unitary operators used by Dollard to show the existence of the wave operators $W_{\pm}(H_1, H_D)$ appropriate for the pure Coulomb case. If V is spherically symmetric then $W_{\pm}(H_2, H_D)$ are shown to be absolutely continuous complete. If in addition $V(r)$ is continuous in $(0, \infty)$, then $W_{\pm}(H_2, H_D)$ are continuous complete. In both cases $S = W_{+}^{*} W_{-}$ is unitary. The connection between the more physical, time-dependent wave operator approach and the traditional time-independent method is made, and phase shift formulas are obtained for the wave and scattering operators. (auth)

46781 NONRELATIVISTIC TIME-DEPENDENT SCATTERING THEORY AND VON NEUMANN ALGEBRAS. I. SINGLE CHANNEL SCATTERING. Zachary, W. W. (Naval Research Lab., Washington, D. C.). *J. Math. Phys. (N. Y.)*; 13: No. 5, 609-15 (May 1972).

For single-channel nonrelativistic time-dependent scattering an analysis is made of the von Neumann algebras $M_{\pm}$ generated by the wave operators $\Omega_{\pm}$ and the algebra B generated by the spectral projections of both the total and free Hamiltonians. Each of the algebras $M_{\pm}$ is the direct sum of an Abelian algebra and a factor of type I_{∞} , but both summands need not occur. For a large class of central potentials B is the direct sum of a countable number of factors of type I_{∞} and its commutant the direct sum of a countable number of finite factors of type I . In the latter case the type of each direct summand is uniquely determined by the angular momentum of the corresponding partial wave. Similar decompositions are obtained for algebras related to B and B' in a natural way that have certain connections with the absolutely continuous and point spectra of the total Hamiltonian. The structure of some algebras related to $M_{\pm}$ is connected with certain properties of the bound states. (auth)

46782 SOLUTION OF FADDEEV EQUATIONS FOR A ONE-DIMENSIONAL SYSTEM. Majumdar, Chanchal K. (Tata Inst. of Fundamental Research, Bombay). *J. Math. Phys. (N. Y.)*; 13: No. 5, 705-8 (May 1972).

Exact solutions are derived for the Faddeev equations for a one-dimensional system of three distinguishable particles of identical mass, interacting pairwise through a delta function potential. The method employed here uses scattering boundary conditions directly on the system of equations. (auth)

46783 SCATTERING THEORY IN FOCK SPACE. Prugovecki, Eduard (Univ. of Toronto). *J. Math. Phys. (N. Y.)*; 13: No. 7, 969-76 (Jul 1972).

It is shown that the wave operators which can be introduced in quantum field theories with cutoff interactions can be represented by wave transformers acting on states represented by statistical operators. Integral representations can be found for these wave transformers, which then form the basis of a time-independent scattering theory for such quantum field theories. It is pointed out that this leads to a time-independent perturbation theory in Fock space. (auth)

46784 COMPLETE SOLUTION OF THE INVERSE SCATTERING PROBLEM AT FIXED ENERGY. Sabatier, Pierre C. (Univ., Montpellier, France. Indiana Univ., Bloomington). *J. Math. Phys. (N. Y.)*; 13: No. 5, 675-99 (May 1972).

Let D be the class of functions which are bounded by $Cr^{-1+\epsilon}$ and $Cr^{-3-\epsilon}$, E the class of potentials $V(r)$ such that $V(r)$, $rV'(r)$, and $r^2V''(r)$ belong to D . E is dense in the class of potentials γ with finite norm $\int_0^\infty \rho |V(\rho)| d\rho$ in which almost all the results of potential scattering are derived. A complete solution of the inverse scattering problem at fixed energy is given in a class $\tilde{E}$ of potentials which contains E . This means that given any set of phase shifts bounded by $CI^{-1-\epsilon}$, all the potentials of $\tilde{E}$ which fit this set of phase shifts are constructed. They depend on an arbitrary function. The fundamental tool in the solution is the "scattering structure function." The method is derived in such a way that an approximation theory and numerical computations are feasible. These, together with various studies of the solutions, are the object of forthcoming papers. (auth)

46785 FREE RIEMANN-GREEN'S FUNCTION OF THE INVERSE PROBLEM OF SCATTERING THEORY. Gugushvili, E. I.; Mentkovskii, Yu. L. (Inst. of Theoretical Physics, Kiev). *Nuovo Cim.*; 10A: No. 2, 292-302 (21 Jul 1972).

A closed expression for the free Riemann-Green's function of the inverse problem of scattering theory is obtained which is

rigorously valid for arbitrary complex angular momentum l . An estimate (from above) for this function is given. The spectral representation of the free Riemann-Green's function which was formally obtained in a previous paper is rigorously derived. (auth)

46786 SOLUTION OF THE STANDING-WAVE INTEGRAL EQUATION. Minelli, T. A.; Zardi, F. (Univ., Padua). *Nuovo Cim.*; 10A: No. 2, 362-78 (21 Jul 1972).

The standing-wave integral equation is considered in the complex momentum plane. This equation is solved, and the solution is used to inquire about some properties of the reactance matrix. As an illustrative application, the pole expansion of the reactance matrix and the two-potential problem are then investigated. (auth)

46787 NUMERICAL ASPECTS OF MOMENTUM-SPACE SCATTERING EQUATIONS FOR PEAKED POTENTIALS. Adelberg, Michael L.; Saperstein, Alvin M. (Wayne State Univ., Detroit). *J. Comput. Phys.*; 9: No. 3, 398-411 (Jun 1972).

Several programming techniques are suggested for solving partial wave scattering equations in momentum space when the potential matrix, as a function of momentum variables, is strongly peaked about the scattering singularity. In particular, an algorithm is described that makes use of the second Born approximation to estimate error in performing principal value quadratures. For a given required accuracy, one obtains a region of integration about the singularity and a Gauss-Legendre quadrature on this region whose truncation error equals the error due to discarding the region away from the singularity and whose order is much lower than that required by full region quadrature to get the same accuracy. Considerable saving in computer time is achieved. (auth)

46788 INVERSE PROBLEM OF SCATTERING THEORY AND RIEMANN'S METHOD. Gugushvili, E. I.; Mentkovskii, Yu. L. (Inst. of Theoretical Physics, Kiev). *Nuovo Cim.*; 10A: No. 2, 277-91 (21 Jul 1972).

By extending Riemann's method to the case of unbounded domains, a new rigorous approach to the inverse problem of scattering theory is developed. The proposed approach allows one to deal with scattering of both charged and neutral particles in the framework of a universal scheme which is also suitable for treating both real- and complex-angular-momentum cases. (auth)

46789 S-MATRIX POLE TRAJECTORIES IN THE N/D APPROACH. Kok, L. P.; van der Ploeg, F. A.; Greben, J. M. (Univ., Groningen, Netherlands). *Phys. Lett.*; 40B: No. 3, 300-4 (10 Jul 1972).

Solutions of the N/D equations in potential scattering are described for some specific local interactions for which the left hand singularities are poles (exponential, Hulthén potential). The quality and the anomalies of the resulting approximate solutions are discussed in detail. (auth)

46790 EXACT STATIONARY SOLUTIONS FOR THE CLASSICAL RELATIVISTIC THREE-PARTICLE PROBLEM. Chern, De-ching. Philadelphia; Temple Univ. (1972). 150p. University Microfilms Order No. 72-17,683.

Thesis.

The existence of certain exact solutions of the nonrelativistic three-body problem was recently shown by Havas. The solutions describe 3 particles revolving around a common axis with the same angular velocity; the configuration is fixed in a plane which also contains the axis of rotation. In this thesis, the corresponding relativistic problem is investigated for electromagnetic, scalar and vector meson interactions. Radiation is excluded by assuming that the interactions are time-symmetric. Equations of motion derived from both field theory and the theory of action at a distance are considered. The existence of exact solutions of the above type is established. These include the relativistic generalization of Havas' solutions, as well as some highly relativistic motions without nonrelativistic counterpart. However, the latter are shown to be not verifiable experimentally with known particles and interactions. (Diss. Abstr. Int., B)

46791 HIGH ENERGY PRODUCTION PROCESSES IN THE RELATIVISTIC EIKONAL APPROXIMATION. Tanaka, Syu. Philadelphia; Univ. of Pennsylvania (1971). 79p. University Microfilms Order No. 72-17,429.

Thesis.

For high energy production processes, the factorization properties of inelastic effects on the elastic amplitude are studied, first by the eikonal approximation method in a perturbative approach (without assuming the momentum of a produced particle is small), and second by functional methods for many- but soft-particle production. The latter corresponds to the situation in the Bloch-Nordsieck model. (Diss. Abstr. Int., B)

46792 MULTIPLE-SCATTERING EXPANSION AND THE GLAUBER APPROXIMATION. Yamazaki, Masakatsu (Tokyo

Inst. of Tech.). Progr. Theor. Phys. (Kyoto); 47: No. 5, 1593-1600(May 1972).

In potential theory a general derivation of the Glauber approximation from the multiple scattering expansion is given by making the high-energy small-angle approximation in Green's function. The problem is treated in coordinate representation. This treatment makes the derivation easier than that of Harrington does. It is found that each sub-reaction matrix which consists of a different number of connected parts of the total reaction matrix is reduced to the corresponding Glauber term at high energies. (auth)

46793 RELATIVISTIC SCATTERING OF PARTICLES WITH SPIN. Barthelemy, M. C. (Univ., Angers, France); Jolivet, Andre. Compt. Rend., Ser. A; 274: No. 13, 1078-81(27 Mar 1972). (In French).

The Faxen-Holtzmark method is generalized in the framework of a relativistic theory for particles with spin. The system of partial differential equations describing the particles is irreducible, linear of the first order, of finite rank and has a covariant form with respect of the Poincare group. A general formula is given for the scattering amplitude. (France)

PHYSICS (LOW-TEMPERATURE)

Superfluidity

Refer also to abstract: 47299.

46794 ENERGY FLUX BETWEEN TWO SUPERFLUID HELIUM BATHS SEPARATED BY A THIN LAYER. Seiden, Joseph (Univ. Paris, Orsay, France). *Compt. Rend., Ser. B*; 274: No. 18, 1049-52 (3 May 1972). (In French).

The energy flux crossing a thin metal layer separating two superfluid helium baths at different temperatures was calculated. The results were applied to experiments performed by Challis and Sherlock. (France)

46795 PHONON DISPERSION AND CONTRIBUTION TO THE HEAT CAPACITY IN LIQUID ^4He . Yamada, Kazuo; Ishikawa, Koki (Nagoya Univ., Japan). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1455-63 (May 1972).

The temperature-dependent shift of the phonon excitation spectrum in liquid ^4He , based on the phonon interactions in quantum hydrodynamics, is calculated over a wide range of momentum. It is found that in a small-momentum limit, the result is in agreement with previous papers, and in a large-momentum limit, the shift decreases logarithmically with increasing momentum and turns into a negative value beyond a critical momentum. Further, a correction term to the specific heat, necessary because of this temperature-dependent shift of the spectrum and giving a deviation from the T^3 law, is evaluated from the lowest-order thermodynamical potential. The predicted contribution is proportional to T^7 and results in negligible magnitude at low temperature and pressure compared with the term γT^5 , which is dominant in the analysis of the experiments for the heat capacity by Phillips et al. As a result, the positive dispersion ($\gamma < 0$) at zero temperature was not altered by the evaluation obtained here. (auth)

46796 CONSEQUENCES OF MICROSCOPIC SYMMETRIES AND LINEAR RESPONSE THEORY FOR THE HYDRODYNAMICS OF SUPERFLUID HELIUM. Fryer, Michael Owen, Seattle; Univ. of Washington (1972). 94p. University Microfilms Order No. 72-20,865. Thesis.

The hydrodynamic properties of liquid He II are analyzed. It is noted that these properties may be described by two-fluid hydrodynamics and by quantum statistical mechanics. By insisting that the linearized versions of these theories produce the same results in the hydrodynamic regime, it is possible to derive information about both types of theories. How the two-fluid hydrodynamic equations are modified by the presence of external fields is discovered. Once the hydrodynamic equations in the presence of the fields described above are known, the hydrodynamic limit of several microscopic correlation functions are determined. Further information about the viscosities of He II is derived in terms of the correlation functions. Finally, the hydrodynamic properties of the order-parameter magnitude are analyzed. A conservation law for it is derived that has the correct high- and low-frequency form, and a single relaxation-time model of it for all frequencies is constructed. (TSS) (Diss. Abstr. Int., B)

46797 ENERGY TRANSFER FROM A PULSED THERMAL SOURCE TO He II BELOW 0.3 K. Pfeifer, Charles Dean. St. Louis; Washington Univ. (1971). 170p. University Microfilms Order No. 72-9363. Thesis.

Measurements are presented of the angular distribution of the energy flux generated by each of two small ($\sim 1 \text{ mm}^2$) pulsed heaters with different surface structures. The heaters are immersed in a large ($\sim 100 \text{ cm}^3$) volume of He II at a temperature near 230 mK. Heater pulse widths are varied from 50 to 1000 μs and heater power densities (W_H) from 100 to 6 w/cm^2 . The amplitude of the radiated energy flux is monitored by six small carbon detectors oriented normal to the flux. They are located at distances (d) of 1 and 2 cm from the heaters and at angles (ϕ) varying from 7 to 89° with respect to the heater normals. For small W_H , the flux travels at $238 \pm 10 \text{ m/s}$ with no spatial attenuation or dispersion. At the critical heater flux density $W_{SE} = 0.7 \text{ w/cm}^2$ for one heater and $W_{SE} = 1.5 \text{ w/cm}^2$ for the other the radiated flux develops a slower component (second edge) traveling at the velocity, $v_{SE} = 130 \pm 40 \text{ m/s}$. At lower heater power densities the angular distribution of flux amplitudes is characteristic of the diffuse radiation of a noninteracting flux of phonons from the heater surface. When W_H exceeds 0.5 w/cm^2 , the angular distribution of flux amplitudes is altered, indicating

appreciable interactions among the excitations in the radiated flux near the heater surface. A semiempirical model for the growth of a helium black-body source from the heater surface is described that accounts for the angular distribution of flux amplitudes in the heater power density range $0.5 \leq W_H \leq 1.2 \text{ w/cm}^2$. The second edge flux component is identified as the roton flux emitted from a helium black-body source at a temperature, $T_s > 0.6 \text{ K}$. A second-sound pulse model is proposed that predicts qualitatively the angular distribution of the second edge flux amplitudes as well as the difference in the critical values W_{SE} for the two heaters. A spectral analysis of the energy flux is attempted to determine the extent to which the helium source is in thermal equilibrium at different values of W_H . (Diss. Abstr. Int., B)

46798 SURFACE-BOUND ELECTRONS ON LIQUID HELIUM: A NEGATIVE RESULT. Ostermeier, R. M.; Schwarz, K. W. (Univ. of Chicago). *Phys. Rev. Lett.*; 29: No. 1, 25-8 (3 Jul 1972).

An experiment is described which was designed to measure both the lifetime and the mobility of image-potential-bound electrons on liquid helium. No evidence for bound states with a lifetime greater than a few microseconds was observed, in disagreement with the results of two different experiments recently presented by other authors. (auth)

46799 ABSENCE OF A QUADRATIC TERM IN THE ^4He EXCITATION SPECTRUM. Roach, Pat R.; Abraham, B. M.; Ketterson, J. B.; Kuchnir, M. (Argonne National Lab., Ill.). *Phys. Rev. Lett.*; 29: No. 1, 32-4 (3 Jul 1972).

By directly measuring the difference in velocity of 30 and 90 MHz sound waves in liquid ^4He below 100 mK, it was determined that the elementary excitation spectrum cannot be described by an expression $\epsilon(k) = \hbar k(1 + \alpha_1 k + \alpha_2 k^2 + \dots)$ which has a value for α_1 that is greater than 0.01 Å. (auth)

46800 QUANTIZED VORTEX RING DYNAMICS AND THE VORTEX CORE RADIUS IN He II. Steingart, M.; Glaberson, W. I. (Rutgers-The State Univ., New Brunswick, N. J.). *J. Low Temp. Phys.*; 8: No. 1-2, 61-77 (Jul 1972).

Using a newly developed time-of-flight technique, the temperature dependence of the vortex core radius in the range $0.35 \text{ K} \leq T \leq 0.60$ was determined by means of precise measurements of vortex ring dynamics. This work was extended to a measurement of the pressure dependence of the vortex core radius. The results are in agreement with the earlier work of Rayfield and Reif at $T = 0.28 \text{ K}$ and zero pressure and with a model of the vortex core proposed by Glaberson, Strayer, and Donnelly. The precision of the measurements enabled us to observe differences in the dynamics of oppositely charged vortex rings in the same electric field configurations. It was possible to quantitatively account for these differences in a simple way by incorporating the frictional drag associated with the ion bound on the vortex core into the equations of motion. The results suggest that the negative ion shape is not greatly distorted by its presence on the vortex core, nor is the negative ion strongly shielded by the vortex velocity field. (auth)

46801 PERSISTENT CURRENT STATES IN ROTATING SUPERFLUID HELIUM. Kojima, H.; Veith, W.; Guyon, E.; Rudnick, I. (Univ. of California, Los Angeles). *J. Low Temp. Phys.*; 8: No. 3-4, 187-93 (Aug 1972).

Persistent current states in superfluid helium contained in a rotating superleak are studied for various velocities and initial conditions. The change in the superfluid-component velocity as the normal-component velocity changes is interpreted in terms of a circulation-free potential flow and the entrance of vortices. The appearance of vortices of opposite sign to that of pre-existing ones in the superleak is discussed, and this is related to the history-dependent nature of persistent currents. An analogy between a superfluid helium system and a highly irreversible Type-II superconductor is made. (auth)

46802 SPIN-WAVE PHENOMENA IN LIQUID ^3He SYSTEMS. Corruccini, L. R.; Osheroff, D. D.; Lee, D. M.; Richardson, R. C. (Cornell Univ., Ithaca, N. Y.). *J. Low Temp. Phys.*; 8: No. 3-4, 229-54 (Aug 1972).

The magnetic effects predicted by Leggett and Rice for Fermi liquids in the collisionless regime were verified. Spin-echo measurements are reported that demonstrate the existence of spin waves and undamped spin currents in liquid ^3He and dilute ^3He - ^4He solutions. The experimental results are compared with the

theoretical expressions of Leggett and Rice, and values of the Fermi liquid interaction parameter $\lambda = [1/(1 + F_0^2)] - \{1/[1 + F_0^2/3]\}$ are derived for liquid ^3He at $P = 0$ and 27 atm and a 6.4% ^3He - ^4He solution. In addition, some techniques for direct observation of individual spin-wave modes are explored and the results reported. (auth)

46803 MICROSCOPIC THEORY OF IMPURITIES IN LIQUID HELIUM. II. Bowley, R. M. (Univ. of Nottingham, Eng.). *J. Low Temp. Phys.*; 8: No. 3-4, 261-75 (Aug 1972).

A microscopic theory of impurities in liquid helium is developed from the theory of Sunakawa and coworkers. A canonical transformation is used to renormalize the phonons. The theory is applied to ^3He impurities, to positive ions, to the interaction between ^3He impurities, and to the interaction between a ^3He quasi particle and a positive ion. It is found that these interactions are zero in the long-wavelength limit, if the phonons are not renormalized. The coupling between the ^3He and the phonons due to the change in the binding energy of the ^3He with density variations is also considered. (auth)

46804 HIGH-RESOLUTION HEAT-CAPACITY STUDY OF ^4He ADSORBED ON BARE COPPER. Princehouse, D. W. (Univ. of Washington, Seattle). *J. Low Temp. Phys.*; 8: No. 3-4, 287-303 (Aug 1972).

Detailed heat-capacity measurements were made over the range $x = 0.03$ to 1.15 monolayers and the temperature range $T = 1$ to 10 K. A small, on-line computer acquired and reduced the data in a very reproducible manner. Vapor-pressure measurements indicate that the substrate is strongly heterogeneous. The heat capacity, while qualitatively similar to previous data on Ar-plated copper, is best explained by a model based on extreme surface heterogeneity. (auth)

Superconductivity

Refer also to abstracts: 46021, 46031, 47269, and 47299.

46805 (BNL-16913) EFFECT OF SURFACE CONDITIONS ON SOME SUPERCONDUCTING PROPERTIES OF NIOBIUM. Strongin, M.; Farrell, H. H.; Varmazis, C.; Halama, H. J.; Kammerer, O. F.; Varma, M. N. (Brookhaven National Lab., Upton, N. Y.); Dickey, J. M. (Queens Coll., New York). 1971. 6p. (CONF-720513-8). Dep. NTIS.

From Applied superconductivity meeting; Annapolis, MD. (1 May 1972).

Measurements of gaseous impurities on the niobium surface are discussed which indicate higher oxygen levels than the bulk concentrations. Superconducting penetration depth measurements with various bulk oxygen concentrations are discussed, and they also indicate higher amounts of oxygen in the surface regime. Finally, the properties of thin films of Nb_3Sn grown on niobium, which can be used as a high T_c protective layer for the delicate niobium surface, are discussed. (M.H.R.)

46806 SINUSOIDAL CURRENT-PHASE RELATION FOR SUPERCONDUCTING THIN-FILM MICROBRIDGES. Gregers-Hansen, P. E.; Levinsen, M. T.; Pedersen, G. Fog (Univ. of Copenhagen). *J. Low Temp. Phys.*; 7: No. 1-2, 99-117 (Apr 1972).

Superconducting systems separated by a very short weak link are treated using the one-dimensional Ginzburg-Landau theory. It is shown that, in the limit of extremely narrow and/or dirty links, $I \sim \sin \Delta\phi$. In the dirty limit the well-known expression for the Josephson tunnel current is obtained. However, the normal tunnel resistance is replaced by the resistance of the link in the normal state. The theoretical results are compared with our experimental investigations on thin-film microbridges of the Dayem type. Deviations from a sinusoidal current-phase relation are discussed in connection with microwave-induced subharmonic steps sometimes observed in the current-voltage characteristics. (auth)

46807 MAGNETIC PROPERTIES OF A SUPERCONDUCTOR WITH HEXAGONAL SYMMETRY. Takanaka, Kenji; Ebisawa, Hiromichi (Tohoku Univ., Sendai). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1781-3 (May 1972).

The anisotropic behavior of H_{c2} , K_2 , and the distortion of the lattice structure of flux lines of technetium are discussed. The discussion is restricted to the isotropic BCS model Hamiltonian for the pairing interaction, in order to elucidate only the effects of the anisotropy of the velocity on the Fermi surface. (M.H.R.)

46808 THERMODYNAMICAL PROPERTIES OF A JOSEPHSON JUNCTION WITH A NORMAL-METAL BARRIER. Ishii, Chikara (Tokyo Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1464-81 (May 1972).

Thermodynamical properties are investigated for the current-carrying metastable states of a model of Josephson junction with a normal-metal barrier. The one-particle density of states and the thermodynamic potential are derived, including the effects of multiple particle-hole reflection of the electrons inside the barrier. The canonical relation between the phase-coupling energy and the barrier supercurrent is confirmed and expressed in a general form in terms of the phase-dependent density of states. (auth)

46809 LOCALIZED MAGNETIC STATES IN SUPERCONDUCTOR. Machida, Kazushige; Inoue, Chuya; Shibata, Fumiaki (To-kyo Univ. of Education). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1482-90 (May 1972).

The effect of a localized magnetic moment on superconductivity is investigated in the s-d exchange model with the generalized Hartree-Fock approximation. A bound state is shown to exist within the energy gap of a superconductor when the coupling is antiferromagnetic and the coupling constant J is greater than the pairing interaction g . When J is much greater than g , the system is shown to have the singlet ground state with the doublet excited state; in other cases, there is no bound state in the energy gap. Depression of superconducting temperature due to magnetic impurities is also calculated by using the t-matrix obtained in this approximation. The result obtained is rather different from those of other theories. (auth)

46810 STUDY OF LOSSES IN AN ASSEMBLY OF SUPERCONDUCTING FILAMENTS TRANPOSED IN A CONDUCTING MATRIX AND SUBJECTED TO A PULSED MAGNETIC FIELD. Hlasnik, I.; Pouillange, J.-P. (CEN de Saclay, Gif-sur-Yvette, France). *Rev. Gen. Elec.*; 81: No. 2, 119-28 (Feb 1972). (In French).

A quantitative study was made of the losses that can occur in a superconductor formed from an assembly of noninsulated superconducting filaments transposed in a normal-metal matrix when exposed to a transverse magnetic field. Expressions for the electric field demonstrate the existence of a critical length, l_c in a direction normal to the magnetic field and the thickness of the assembly, for which the current circulating between the superconducting plates is equal to their individual critical currents. It is demonstrated that for $l \leq 0.21l_c$ a negligible increase in losses in the coil results from the circulating currents. These results are extrapolated to deal with the real case of filament conductors. (France)

46811 TUNNELING GENERATION, RELAXATION, AND TUNNELING DETECTION OF HOLE-ELECTRON IMBALANCE IN SUPERCONDUCTORS. Tinkham, M. (Cambridge Univ., Eng.). *Phys. Rev., B*; 6: No. 5, 1747-56 (1 Sep 1972).

The theory of electron tunneling is extended to treat the generation and detection of an imbalance Q between electronlike and holelike quasiparticle populations in superconductors. The equilibration of injected high-energy quasiparticles owing to inelastic phonon processes is discussed, and it is shown that in tin, Q relaxes in $\tau_Q \approx [\Delta(0)/\Delta(T)](2 \times 10^{-10})$ sec. It is also shown that even a small ($\sim 1\%$) anisotropy of the gap can lead to significant Q relaxation by elastic scattering processes. This discussion extends the previous brief interpretation of the pair-quasiparticle potential difference measured by Clarke. (auth)

46812 NUMERICAL CONSTANTS FOR ISOLATED VORTICES IN SUPERCONDUCTORS. Hu, Chia-ren (Univ. of Southern California, Los Angeles). *Phys. Rev., B*; 6: No. 5, 1756-60 (1 Sep 1972).

For isolated vortex lines in high- κ , type-II superconductors, Abrikosov derived the expressions $B(0) = \kappa^{-2}(\ln \kappa + C_0)H_{c2}$ and $H_{c1} = \frac{1}{2}\kappa^{-2}(\ln \kappa + C_1)H_{c2}$, but the numerical values he provided for the constants C_0 and C_1 were previously found to violate an identity $C_1 - C_0 - \frac{1}{2} = C_\gamma > 0$. The constants are reevaluated, giving $C_0 = -0.282$, $C_1 = 0.497$, $C_\gamma = 0.279$. Furthermore, for superconductors containing a high concentration of magnetic impurities, it was previously shown that the electric field generated at the center of an isolated vortex in flux-flow situations is proportional to H_{c2} and the flux-flow velocity v . The proportionality constant C_E in the high- κ limit is numerically evaluated here to be 0.951, which, together with the value for C_γ , determines the flux-flow resistivity $\rho_f = 0.381 \rho_n(B)/H_{c2}$ in the low-applied-field limit when vortices are very far apart. (auth)

46813 THERMODYNAMIC PROPERTIES OF SMALL SUPERCONDUCTING PARTICLES. Muehlschlegel, B. (Univ., Cologne); Scalapino, D. J.; Denton, R. *Phys. Rev., B*; 6: No. 5, 1767-77 (1 Sep 1972).

In bulk superconducting systems the large intrinsic range of the pair coherence length implies an extremely narrow critical region. However, as is well known, the size of the critical region

becomes larger as the dimensions of the system decrease below the coherence length. An interesting limiting case corresponds to particles with all dimensions less than a coherence length which form essentially zero-dimensional systems. Here, the thermodynamic properties of small superconducting particles are calculated using a static approximation. Within this approximation, both the order parameter and the quasiparticle contribution are explicitly studied. (auth)

46814 DYNAMIC STRUCTURE OF VORTICES IN SUPERCONDUCTORS. III. GRAPHS FOR $B \approx H_{c2}$. Thompson, R. S.; Hu, C.-R.; Smith, T. L. (Univ. of Southern California, Los Angeles). Phys. Rev., B; 6: No. 5, 2044-7 (1 Sep 1972).

Graphs are presented illustrating the backflow generated by vortices moving in a superconductor as driven by a transport current. (auth)

46815 STUDY OF FLUX-FLOW NOISE IN TYPE-II SUPERCONDUCTORS. Shen, Shu-wei. Minneapolis; Univ. of Minnesota (1972). 95p. University Microfilms Order No. 72-20,145.

Thesis.

Flux-flow noise in type-II superconductors is investigated theoretically and experimentally. The observed noise turns out to be shot noise as expected from the assumption that the flux-flow voltage is generated by independently drifting flux bundles. Bundle sizes as large as 10^2 to 10^3 are found in vanadium foils, dependent on the biased condition. The experiment shows qualitative agreement with the theory. (Diss. Abstr. Int., B)

46816 ULTRASONIC ATTENUATION IN SUPERCONDUCTING INDIUM AND INDIUM-TIN ALLOYS. Binnie, David Earl. University Park, Pa.; Pennsylvania State Univ. (1971). 99p. University Microfilms Order No. 72-19,274.

Thesis.

The ultrasonic attenuation coefficients in the normal and superconducting states were measured as a function of temperature and orientation in single-crystal indium and three single-crystal indium-tin alloys with 0.01, 0.03, and 0.1 at. % tin. The measurements were made at temperatures between 1.0 and 4.2°K for ultrasonic waves propagating along the [001], [100], and [110] directions in the indium face-centered tetragonal lattice. The superconducting energy gap for each orientation and alloy was calculated from the ultrasonic data. (TSS) (Diss. Abstr. Int., B)

46817 RESISTIVE TRANSITION OF A SUPERCONDUCTOR. Thomas, Gordon Albert. Rochester, N. Y.; Univ. of Rochester (1972). 173p. University Microfilms Order No. 72-15,341.

Thesis.

Experiments testing the limits of applicability and the systematic breakdown of the mean field theory of superconductivity are presented. The electrical resistance is used as a probe of the theory in a study dealing with four interrelated aspects of the general problem. First, the linearized, time-dependent Ginzburg-Landau (GL) theory is investigated through experiments on the fluctuation conductivity of Al thin films in large electric fields above the transition temperature T_c . Second, one-dimensional microstrips are studied, and it is found that, although the simple linearized theory breaks down close to T_c , the theory can be extended into the region near T_c by the self-consistent inclusion of the fourth-order term in the GL free energy. The existence of a characteristic current for nonlinear phase slippage and a fluctuation resistance in the zero-current limit below T_c in one-dimensional microstrips are demonstrated in experiments. The inadequacy of the simple mean field treatment of the fluctuation conductivity in clean samples above T_c is elucidated in the final section, where three studies of the effects of ephemeral Cooper pairs on the normal electrons are presented: two-dimensional $\text{Sn}_{1-x}\text{In}_x$ films in parallel magnetic fields, one-dimensional Al microstrips in parallel magnetic fields, and two-dimensional Al films with the addition of magnetic impurities. In all cases a quenching of the normal electron paraconductivity was observed as predicted by the theory. Experimental evidence is presented in this dissertation that the resistive transition of small superconducting specimens is strongly modified by the presence of thermodynamic fluctuations. These studies have yielded new information about the nature of phase transformations in solids and the time-dependent theory of superconductivity. (TSS) (Diss. Abstr. Int., B)

46818 ULTRASONIC ATTENUATION IN PURE NIOBIUM AT LOW TEMPERATURES. Alexander, Edward Michael. College Park, Md.; Univ. of Maryland (1972). 130p. University Microfilms Order No. 72-15,340.

Thesis.

Ultrasonic attenuation measurements were made on very pure niobium (RR ~ 50%) with q values up to 11 at T_c . These mea-

surements were made using a pulse-transmission method at frequencies up to 270 MHz and thin-sample handling techniques. The high value of q was determined by fitting the data to a normal-state electronic-attenuation model and independently using magneto-acoustic criteria. Even though q was much greater than 1, no rapid fall in the attenuation ratio was observed below T_c . This is the first such observation in the literature. Further, a low-temperature maximum in the normal-state attenuation as a function of temperature was observed. This phenomenon has thus far been reported for only one other metal, rhenium, and not previously from niobium. The relation of the maximum to the absence of rapid fall is also discussed. The data are compared with the data of all other groups who have measured shear wave attenuation in niobium. It is noted that for $q < 1$ the attenuation ratio vs. temperature data follows BCS predictions, but that as sample purity increases there is increasing deviation from BCS. For $q \sim 1$ the deviation from BCS predictions appears to be maximized. As frequency is increased in a high-purity specimen, the deviation from BCS becomes smaller, suggesting a return to the BCS prediction as q is increased from $q \sim 1$ to $q \gg 1$. The deviations from BCS theory are examined in relation to recent models which will account for phonon scattering and interband effects. For the first time in niobium, magneto-acoustic oscillations (at least four) in the high-purity sample were also observed. From these oscillations it was possible to measure calipers of the niobium Fermi surface for directions in the (100) plane. These calipers are in good agreement with the theoretically predicted calipers for the so-called distorted ellipsoids of the niobium Fermi surface. (Diss. Abstr. Int., B)

46819 FLUX FLOW IN NONIDEAL TYPE-II SUPERCONDUCTORS. Kusayanagi, Eiichirou; Kawashima, Teruko; Yamafuji, Kaoru (Kyushu Univ., Fukuoka). J. Phys. Soc. Jap.; 33: No. 2, 551-60 (Aug 1972).

The resistive behavior of the mixed state is studied theoretically in nonideal Type-II superconductors containing many defects. The fundamental equations are the time-dependent Ginzburg-Landau equations generalized from the conventional ones so as to describe elastic interactions of flux lines with defects. The solutions are obtained for the case of weak and localized defects by perturbation calculations in the vicinity of the upper critical field. It is shown that the flux-line lattice in nonideal samples can carry a finite amount of nondissipative transport current, which leads to the appearance of the depinning current term in the relation of the induced electric field to the transport current. A detailed expression of the critical current density is derived for a simple type of defect. An expression of the flow conductivity is shown to be also modified by the presence of defects. (auth)

46820 INELASTIC NEUTRON SCATTERING FROM VORTEX LATTICE IN TYPE-II SUPERCONDUCTORS. II. THE LONDON MODEL. Maki, Kazumi (Nuclear Research Center, Juelich, Ger.). J. Low Temp. Phys.; 8: No. 1-2, 47-59 (Jul 1972).

The inelastic neutron scattering from the vortex lattice in type-II superconductors is considered. Use is made of the London model to describe the vibration of the vortex lattice. It is found that the differential cross section has sharp peaks besides the Bragg peaks when the transferred momentum q approaches the reciprocal lattice vector G . These peaks are due to excitation of the vibrational modes and diverge as $|q - G|^{-2}$. The Debye-Waller factor is also calculated, and is extremely small.

46821 TIME-DEPENDENT SUPERCONDUCTIVITY AND QUANTUM DISSIPATION. Rieger, T. J. (Univ. of California, Santa Barbara); Scalapino, D. J.; Mercereau, J. E. Phys. Rev., B; 6: No. 5, 1734-43 (1 Sep 1972).

A time-dependent Ginzburg-Landau theory is used to calculate the time-varying currents and potentials in a one-dimensional superconductor undergoing quasi-continuous phase slip. To account for the irreversible nature of this process, the concept of superconducting dissipation is introduced. This analysis is consistent with many observations on weak links and provides an adequate model for at least their low-frequency behavior. (auth)

46822 PROBING MANY-BODY EFFECTS WITH SUPERCONDUCTIVITY. Chaikin, Paul Michael. Philadelphia; Univ. of Pennsylvania (1971). 251p. University Microfilms Order No. 72-17,336.

Thesis.

The principle forming the basis of this thesis is that superconductivity is the most (maybe the only) well-understood many-body problem in solids. The point emphasized is that other single- and many-body interactions can be probed with superconductivity and a great deal of information gained. To illustrate the usefulness of superconductivity in studying many-body effects, experiments dealing with magnetic impurities and the Kondo effect

were carried out. Two sets of experiments were undertaken: one involved the critical-field behavior of superconductors containing magnetic impurities and another involved tunneling into proximity-effect sandwiches with magnetic impurities in the normal-film side. Using multiple pairbreaking theory, the H_{C2} data for pure La and for La-Ce and La-Gd alloys were analyzed. The tunneling experiments entail inducing a normal metal film (which contains magnetic impurities) superconducting by the proximity effect. This allows study of the density of states characteristic on the normal-metal side. The systems studied were AuFe/Pb and PdCo/Pb. (TSS) (Diss. Abstr. Int., B)

46823 RESISTIVITY OF THE INTERMEDIATE STATE OF TYPE-I SUPERCONDUCTORS. Lowell, J.; Parkinson, S.; Rose-Innes, A. C. (Univ. of Manchester, Eng.). *J. Low Temp. Phys.*; 8: No. 3-4, 277-85(Aug 1972).

The interaction between transport current and flux spots in the intermediate state of a superconductor is investigated. It is shown that the force on the flux spots depends on factors such as current lead configuration, and it is therefore not possible to define a unique flow resistivity for type-I superconductors. The conclusions are supported by experimental evidence. (auth)

46824 EFFECT OF STRAIN ON ULTRASONIC ATTENUATION IN SUPERCONDUCTORS. Das, H. L.; Chaudhuri, K. D. (Univ. of Delhi). *Physica*; 59: No. 4, 693-6(15 Jun 1972).

Ultrasonic-attenuation measurements in single crystals of 99.9999% pure tin were taken both in the superconducting and in the normal-conducting states with the help of the standard double-ended pulse-echo technique. Strain in the specimen has a pronounced effect on the attenuation in the superconducting state while attenuation in the normal conducting state remains uninfluenced by the amount of strain in the specimen. Normal conducting attenuations are also seen to be independent of amplitude of the ultrasonic pulses. For sufficiently strained samples, attenuation in the superconducting state becomes identical with that in the normal-conducting state. (auth)

46825 KINETIC EQUATION FOR SUPERCONDUCTING ALLOYS LOCATED IN WEAK STATIC FIELDS. Melik-Barkhudarov, T. K. (Inst. of Physical Research, Erevan). *Dokl. Akad. Nauk Arm. SSR*; 53: No. 1, 21-5(1971). (In Russian).

A simple case of a superconducting alloy positioned in a weak static field is mathematically developed. Certain matrix elements are encountered, which are assumed to be significantly different from zero in a narrow band of energy close to the Fermi surface of the order of the critical temperature. (K.S.W.)

46826 LOCAL MAGNETIC FIELD DISTRIBUTIONS IN SUPERCONDUCTING NIOBIUM AT 4.2 K BY NEUTRON DIFFRACTION. Schelten, J.; Ullmaier, H.; Lippmann, G. (Kernforschungsanlage, Juelich, Ger.). *Z. Phys.*; 253: No. 3, 219-31(1972).

Neutron-diffraction experiments on flux line lattices in superconducting Nb are reported. From the integral reflectivities of the (10), (11), (20), (21), and (31) reflections the absolute values of the form factors were obtained as a function of the flux density. Considering certain properties of the flux line lattice it is also possible to determine the phases of the form factors. From this the local field distribution in the mixed state could be calculated by Fourier transformation. The dependences of the resulting maximum, minimum, and saddle-point fields on the flux density and on the impurity parameter are discussed and compared with theory. (auth)

46827 TIME-DEPENDENT PHENOMENA IN WEAKLY-COUPLED SUPERCONDUCTORS. Lepae, L.; Mancini, F.; Umezawa, H. (Univ. of Wisconsin, Milwaukee). *Nuovo Cim.*; 10B: No. 1, 267-82(11 Jul 1972).

Two superconducting systems joined by a weak link present a great variety of phenomena which are not totally explained by the Josephson formulation. A general theory is given of the phenomena observed in these systems when time-dependent effects are taken into account. The formulation, presented as an application of the boson method in superconductivity, is derived from first principles in the framework of a microscopic theory. The results show the appearance of harmonic and subharmonic steps in the I-V characteristic of the system; the presence or not of subharmonic steps is essentially determined by the geometry of the system. The existence of subharmonic terms in ac effects is shown to be strongly related to the diffraction pattern presented by the I_c-H curves in dc experiments performed on the same sample. A comparison of the usual theory and the present formulation is discussed. (auth)

46828 IDENTIFICATION OF THE SUPERCONDUCTING PHASES IN THE MOLYBDENUM-PLATINUM SYSTEM. Flueckiger, R.; Paoli, A.; Roggen, R.; Yvon, K.; Muller, J. (Univ., Geneva). *Solid State Commun.*; 11: No. 1, 61-3(1 Jul 1972).

Structural-phase analysis was combined with an extensive study of the superconducting transitions, as observed magnetically and calorimetrically. The results show that the highest critical temperatures occurring in this system have been erroneously ascribed to the A-15-type structure. (auth)

46829 CRITICAL CURRENTS IN Nb₃Al. Martin, P. J.; Campbell, A. M.; Evetts, J. E. (Cambridge Univ., Eng.). *Solid State Commun.*; 11: No. 1, 123-5(1 Jul 1972).

Alternating current measurements on Nb₃Al showed that large amounts of the sample are driven normal at relatively low fields. Small variations of T_c are associated with very large variations of H_{C2}, caused, most probably, by varying degrees of disorder. These regions of low H_{C2} material may explain the low critical current densities measured in Nb₃Al and other A15 superconductors. (auth)

46830 SURFACE EXCITATIONS OF TYPE-I SUPERCONDUCTORS IN A STATIC MAGNETIC FIELD. Rogani, A.; Tabet, E. (Istituto Superiore di Sanita, Rome). *Solid State Commun.*; 11: No. 1, 127-30(1 Jul 1972).

Surface excitations of type-I superconductors in a static magnetic field are studied assuming diffuse scattering at the superconductor surface. The energy spectrum and the density of states for these excitations are obtained in the self-consistent field approach. Some remarks are made about the relevance of the surface states for the microwave impedance evaluation. (auth)

46831 DISCONTINUOUS FLUX FLOW IN SUPERCONDUCTING NbSe₂. Frindt, R. F.; Huntley, D. J.; Kopp, J. (Simon Fraser Univ., Burnaby, British Columbia). *Solid State Commun.*; 11: No. 1, 135-7(1 Jul 1972).

The flux flow resistance region was observed to end by a series of discontinuous steps to zero as the applied magnetic field or specimen current is increased. This supports Pippard's model of the flux line lattice deforming to match the pinning centres. A new interpretation of the peak effect in the critical current curve is also given. (auth)

46832 ANOMALOUS CRITICAL FIELD CURVES OF THE KONDO SUPERCONDUCTOR (La_{1-x}Ce_x)Al₂. Riblet, G.; Winzer, K. (Univ., Cologne). *Solid State Commun.*; 11: No. 1, 175-7(1 Jul 1972).

The critical field curves of (La_{1-x}Ce_x)Al₂ alloys were investigated and a maximum in H_{C2}(T) for finite temperatures between 0.4 and 0.6°K for Ce concentrations x > 0.003 was observed. The results are in rough agreement with the theory of additive pair-breaking as applied to a Kondo superconductor. (auth)

46833 MAGNETIC FIELD DISTRIBUTIONS IN TYPE II SUPERCONDUCTORS. Inhaber, H. (Univ. of Oklahoma, Norman). *Solid State Commun.*; 11: No. 1, 275-8(1 Jul 1972).

The internal induction distribution of Type-II superconductors is calculated using experimental data on a Nb ellipsoid of Cline et al., (*J. Appl. Phys.*, 37: 1(1966)) and a theory of Inhaber (1972). Qualitative agreement is found between theory and experiment; however, the distortion of fluxoid lattice due to shape, possible inhomogeneities in the distribution of pinning centers, and lack of probe resolution may be factors in preventing quantitative agreement. (auth)

46834 MODEL OF THE JOSEPHSON EFFECT. Turski, L. A. (Univ., Warsaw). *Nuovo Cim.*; 10B: No. 1, 102-10(11 Jul 1972).

A simple quantum-mechanical model of two coupled Bose systems exhibiting phase coupling, i.e., the Josephson effect, is proposed. The model is constructed in the spirit of the Bogolyubov's approach to the many-boson problem. The coherent-state technique is used. (auth)

46835 VOLTAGES INDUCED IN SUPERCONDUCTORS IN THE ABSENCE OF TRANSPORT CURRENTS. Ohmer, Melvin C.; Joiner, W. C. H. (Univ. of Cincinnati). *Phil. Mag.*; 26: No. 1, 15-32(Jul 1972).

When a superconducting sample is subjected to superimposed dc and oscillatory magnetic fields, under certain circumstances, voltages are found to exist in the sample even though no external currents are supplied. A dc voltage component is produced which reverses sign with a reversal in polarity of the dc magnetic field. Except for this sign reversal, the dc voltage versus dc field trace is symmetric with respect to field reversals. Voltage components are also present which are harmonics of the oscillatory field. The effects are quite complex and depend on the values of the many

parameters in the experiment. The results are discussed in terms of the existence of asymmetric flux flow through the sample during each cycle of the oscillatory field. (auth)

46836 EFFECT OF DISLOCATION TANGLES ON SUPERCONDUCTING PROPERTIES. Dew-Hughes, D.; Witcomb, M. J. (Univ. of Lancaster, Eng.). *Phil. Mag.*; 26: No. 1, 73-96 (Jul 1972).

The microstructure of deformed Mo-34 at.% Re alloys was correlated with normal-state resistivity and superconducting properties. A model based on the dislocation cell structure is able to account for the variation of normal-state resistivity, upper and lower critical fields, and magnetic hysteresis, in the superconducting state. The superconducting properties are shown to be dominated by the presence of high-resistivity cell walls. In particular, the movement of quantized flux vortices is hampered by an interaction with the interfaces of this cell structure. Calculated parameters are in reasonable agreement with observed values. (auth)

46837 THEORY OF ELECTRON TUNNELING IN NIS SYSTEMS WITH PARAMAGNETIC IMPURITIES. Cheishvili, O. D. (Inst. of Physics, Tbilisi, USSR). *J. Low Temp. Phys.*; 8: No. 3-4, 213-18 (Aug 1972).

Current-voltage characteristics of NIS junctions were studied when there are paramagnetic impurities in a narrow layer of the superconductor close to the insulator. It is shown that paramagnetic impurities in such systems considerably change the current-voltage characteristic in the narrow range of the applied potential $eV \sim \Delta$ and $V \sim 0$. (auth)

46838 SUPERCONDUCTIVITY AND LOCALIZED SPIN FLUCTUATIONS IN CONCENTRATED ALUMINUM-3d-TRANSITION-METAL ALLOYS. Babic, E.; Ford, P. J.; Rizzuto, C.; Salamoni, E. (CNR, Genoa). *J. Low Temp. Phys.*; 8: No. 3-4, 219-28 (Aug 1972).

Measurements are presented of the superconducting transition temperatures of concentrated alloys of aluminum containing Ti, V, Cr, Fe, and Co, prepared using an ultrarapid quenching technique. These results extend the concentration range of existing measurements by one or two orders of magnitude. The data were analyzed using the theories of Kaiser and of Rivier and MacLaughlin, and it was found that both theories can adequately describe the results for the initial slopes $(dT_c/dn)_0$ and concentration dependence of T_c . In particular, from the theory of Rivier and MacLaughlin, characteristic temperatures were found for these alloys that are in agreement with those obtained from measurements on other physical parameters. (auth)

46839 CRYOGENIC PROBLEMS OF FORCED COOLED SUPERCONDUCTING SYSTEMS. Keilin, V. E.; Klimentenko, E. Yu. (Inst. of Atomic Energy, Moscow). *Cryogenics*; 12: No. 4, 12: No. 4, 292-6 (Aug 1972).

Methods of producing cold compressed helium, various ways of refrigerant distribution, and ways of reducing cool-down time being applied to forced-cooled superconducting systems (FCSS) are discussed. The conditions obtained are for preventing normal-zone propagation from current leads inwards into the winding. The analysis shows that main cryogenic problems of FCSS can be solved relatively simply. (auth)

46840 PHASE SENSITIVE METHOD FOR DETECTING SUPERCONDUCTIVE TRANSITIONS IN THE MICROWAVE REGION. di Crescenzo, Enrico; Flamini, Mario; Indovina, Pietro L.; Onori, Sandro; Rosati, Alfredo. *Ann. Ist. Super. Sanita*; 7: 560-84 (1971). (In Italian).

The measurement method described detects superconductive transitions as a function of the magnetic field applied by means of a phase-sensitive technique. A detailed description of the modulation parameters is given according to the criteria followed in EPR experiments. In the description of the experimental equipment, particular attention is given to the stabilization system of

the microwave generator and to the detection method. Furthermore, the resonant cavity associated with the cryogenic system is fully explained. (auth)

Equipment and Devices

Refer also to abstract: 45679.

46841 INTRODUCERE IN FIZICA TEMPERATURILOR JOASE. (Introduction to Low Temperature Physics). Balla, D.; Deutsch, R. V. Bucharest; Editura Academiei Republicii Socialiste Romania (1970). 494p. Lei 31.

Proceeding from a study of the solution properties of the Schroedinger equation, problems related to the isoenergetic and Fermi surfaces are examined. The Brillouin zones and energy bands are introduced, conclusions arising from the dispersion equations are formulated, and theoretical methods used for the study of the isoenergetic and Fermi surfaces are presented. A concise description of the principles of phenomenological thermodynamics is given in a form corresponding to the presence of electric and magnetic fields. A systematic exposition of the statistical physics methods used is given. Practical methods used for realization of low temperatures are described. Temperature measurement methods are covered. Magnetic properties of substances at low temperatures are presented. The theory of specific heats of solids at low temperature is dealt with. Transport phenomena in solids at low temperature are covered. Liquid helium properties are described. Experimental and theoretical results referring to superconductivity are presented. (M.H.R.)

46842 METHOD AND MEANS FOR CHARGING OR DISCHARGING SUPERCONDUCTING WINDINGS. Bergmann, Winfried H. U. S. Patent 3,667,029. 30 May 1972. Filed 15 Apr 1970.

A winding of superconducting material is charged by electrically shorting the winding with a hard Type-II series superconducting material and cooling the winding and shorting material to a superconducting temperature. A magnetic flux is generated to penetrate the shorting material to establish therein fluxoids and a temperature gradient is established across the shorting material to cause motion of the fluxoids across the shorting material into the magnetic circuit of the winding to effect charging or discharging thereof. (Official Gazette)

46843 SUPERCONDUCTIVE MAGNETS. Donaldson, Gordon W.; Graham, Charles D. Jr.; Hart, Howard R. Jr. (to General Electric Co.). U. S. Patent 3,665,351. 23 May 1972. Filed 9 Jul 1970.

Superconductive tape-wound solenoids used for the production of high magnetic fields are susceptible to being driven into the normal resistance state by flux jumping occurring in the end coils. This is caused by the high concentration of magnetic flux components perpendicular to the planes of the tapes in these coils. The performance of such solenoids may be substantially improved by providing an alternative path for such flux components through a body having a relatively high saturation magnetization such as iron or mild steel closely adjacent the end coils but electrically isolated therefrom. (Official Gazette)

46844 TUBULAR NIOBIUM/COPPER CONDUCTORS FOR AC SUPERCONDUCTIVE POWER TRANSMISSION. Graeme-Barber, C. (Central Electricity Research Lab., Leatherhead, Eng.); Maddock, B. J.; Popley, R. A.; Smedley, P. N. *Cryogenics*; 12: No. 4, 317-8 (Aug 1972).

Composite niobium/copper tubes with diameters of the order of those that would be required for superconducting cables (eg 63.5 and 100 mm) were made with power ratings of a few GVA. Although the detailed electrical performance of these tubes has still to be determined, the results of measurements on small samples suggest that power losses will be acceptably low. Areas in which problems remain are outlined. (UK)

PHYSICS (NUCLEAR)

Neutron Interactions with Matter

Refer also to abstracts: 47220-47224 and 47251.

46845 (BNL-16935) FOURIER TRANSFORM OF INELASTIC NEUTRON SCATTERED INTENSITY FOR SOLID ^4He . Gilbert, Ernest (Brookhaven National Lab., Upton, N. Y.). 1972. 47p. Dep. NTIS.

The inelastic scattering of neutrons was used to study the properties of solid ^4He . Fourier transforms were used to determine the probability density of the helium atoms. (J.W.R.)

46846 (JEN-221) GREEN FUNCTION IN THE SECONDARY MODEL OF THERMALIZATION. Guasp, Jose (Junta de Energia Nuclear, Madrid (Spain)). 1972. 170p. (In Spanish). Dep. NTIS (U. S. Sales Only).

The Green function of the thermalization problem is studied in the secondary model case through the spatial Fourier transform. A relation between singularities and eigenvalues allows the determination of the analyticity dominion. The eigenvalues spectrum has a purely discrete part, lying on an interval of the imaginary axis of the k complex plane (the Fourier parameter), and another part, purely continuous, lying in the remainder of the imaginary axis. A correspondence between discrete eigenvalues and "exponential modes" of the Green Function, extremal properties for the eigenvalues and some remarkable properties of the eigenfunctions are established. The numerical methods used, results about the possibility of disappearance of the fundamental eigenvalue when the absorption is increased, and the apparition of a "pseudo-fundamental" mode in the Green Function are discussed. (auth)

46847 TIME-DEPENDENT NEUTRON TRANSPORT EQUATION WITH DELAYED NEUTRONS. Yoo, Kun Joong; Pac, Pong Youl (Seoul National Univ.). J. Korean Nucl. Soc.; 4: No. 2, 102-8 (Jun 1972).

The time-dependent neutron transport equation with delayed neutrons is analytically solved for the case of isotropic scattering with constant cross sections. The equations in the two divided time regions are obtained from the original equation by the asymptotic method. It is shown that the approximate solutions in each time region are uniformly valid in time to the order of the inverse magnitude of the velocity. (auth)

46848 FAST NEUTRON TIME-OF-FLIGHT SPECTROMETER WITH HIGH RESOLUTION. Cho, Mann (Atomic Energy Research Inst., Seoul). J. Korean Nucl. Soc.; 4: No. 2, 116-31 (Jun 1972).

A fast neutron time-of-flight spectrometer was constructed (with suitable choice of target thickness and proton bombarding energy in the $^7\text{Li}(p,n)^7\text{Be}$ nuclear reaction) for a continuous keV spectrum of neutrons at 0 degrees in 1-ns pulses from a Van de Graaff. A time-pick-up fast neutron detector with a 5-mm-thick, 92% enriched ^{10}B slab and four heavily shielded $4'' \times 3''$ NaI scintillation detectors was assembled. The energy resolution of this spectrometer is better than 0.3% at 50 keV, and the signal-to-background ratio is also improved. Total cross section measurements of several separated single isotopes were carried out with this spectrometer and analyzed by an R-matrix multi-level computer code. The spin values and resonance parameters of each individual resonance are given. (auth)

46849 NEW RESULTS IN THEORY OF PULSED THERMAL NEUTRONS. Protopopescu, V. (Inst. of Atomic Physics, Bucharest). Stud. Cercet. Fiz.; 24: No. 3, 347-67 (1972). (In Romanian).

Topics discussed include the analytical structure of the solution in the continuous sections case, an experimental situation, the analytical structure of the solution in the discontinuous sections case, and the analytical extension of the dispersion law. (J.N.R.)

46850 GREEN'S FUNCTION OF THE NEUTRON TRANSPORT EQUATION FOR A MOVING MEDIUM. Garusov, E. A.; Petrov, Yu. V. This paper was summarized in At. Energ. (USSR); 30: No. 5, 444-5 (May 1971) as paper No. 485/5933. (In Russian).

Full copy of this paper is held by TIC.

The system of eigenfunctions of the stationary neutron transport equation is generalized to the case of nuclei of infinite mass that move with the same velocity u in a direction perpendicular to the plane of the source. It is assumed that the cross sections are independent of the energy and that the neutrons are scattered isotropically in the center of mass system. (auth)

46851 SOLUTION OF MULTI-DIMENSIONAL NEUTRON TRANSPORT EQUATION BY FINITE ELEMENT METHOD. Ukai, Seiji (Kyoto Univ.). J. Nucl. Sci. Technol. (Tokyo); 9: No. 6, 366-73 (Jun 1972).

A scheme of treatment with the finite element Galerkin method is proposed for the approximation of solutions of multi-dimensional steady state neutron transport equations, and it is proved that the approximate solutions yielded by the treatment converge to the solutions of the transport equations under reasonable hypotheses. These approximate solutions are used also to show the existence of weak solutions of transport equations. Finally, based on the Galerkin method, the conditions for convergence are discussed for the multi-group approximation and for the method of spherical harmonics. (auth)

46852 PROTECTION FROM THE NEUTRON EMISSION OF ^{238}Pu DIOXIDE. Zharkov, V. A.; Chudotvorov, A. A.; Moiseenko, B. A. This paper was summarized in At. Energ. (USSR); 31: No. 1, 50-1 (Jul 1971) as paper No. 505/6061. (In Russian).

The neutron spectrum, dose multiplicity of the attenuation afforded by water shielding, and cross section for neutron removal by iron were investigated for a specimen of $^{238}\text{PuO}_2$. The neutron yield was measured to be $(2.0 \pm 0.2) \times 10^4$ n/s-g of ^{238}Pu . Calculations were made to determine the specific contributions made by the neutrons from the (α, n) reaction with oxygen and by spontaneously generated neutrons to the total yield; the calculated spectrum agreed well with the experimental one. The flux-to-dose conversion coefficient was found to be $(3.70 \pm 0.25) \times 10^{-5}$ mrem/n/cm² for the specimen examined. The experimentally obtained curve for the dose multiplicity of the attenuation of neutron radiation from $^{238}\text{PuO}_2$ by water shielding in semi-infinite geometry was compared with the corresponding calculated curve. They differed somewhat; this difference was attributed to the barrier effect. The removal cross section was found to be 1.80 ± 0.07 b. It was established that the neutron removal cross section is independent of the thickness of the removing materials (iron) and the thickness of the water shielding, provided that this is between 30 and 45 cm. (4 figures) (R.W.R.)

46853 INTERFERENCE IN THE ULTRACOLD NEUTRON TRANSMISSION THROUGH THIN LAYERS. Steyerl, A. (Technische Univ., Munich). Z. Phys.; 252: No. 5, 371-82 (1972).

At the time-of-flight spectrometer for ultracold neutrons (UCN) with energies below 10^{-4} eV at the FRM, Munich, the perpendicular neutron transmission through carbon and gold films of thickness 450 to 800 Å was measured as a function of neutron velocity in the range $11 \geq v_z \geq 3.7$ m/s. The transmission curves obtained show the expected interference patterns due to interference between the partial waves transmitted and reflected at the two surfaces. From these curves absolute values for the thickness and density could be obtained. From the UCN transmission through a $4.71\text{-}\mu\text{m}$ -thick copper foil a value $a_{\text{coh}} = 7.5 \pm 0.15$ F was derived for the nuclear scattering length of Cu. (5 figures) (auth)

46854 TWO-DIMENSIONAL TRANSPORT PROBLEMS IN A QUARTER SPACE. Lam, Siu Kee. Stanford, Calif.; Stanford Univ. (1971). 108p. University Microfilms Order No. 72-11,593.

Thesis.

Neutral particle transport phenomena in a quarter space are investigated by considering the one-speed integral transport equation. Both the homogeneous (Milne's problem) and inhomogeneous (constant-source problem) form of the transport equation are examined. For both problems, the method of attack is double Fourier transformation and use of the one-dimensional Wiener-Hopf technique. All contributions to the density asymptotically away from the corner are determined explicitly by considering two appropriate half-space problems and by extensive use of contour integration in two complex variables. The remaining contributions near the corner are obtained by iterating the transport equation. Numerical results are given for a particular symmetric Milne's problem and a particular constant-source problem. Diffusion theory results for these problems are given for comparison. Extension to the case of a quarter space surrounded by another medium with the same total cross section but different mean number of particles emitted per collision is also given. (Diss. Abstr. Int., B)

46855 SUBCRITICAL MULTIPLICATION OF ^{252}Cf NEUTRONS AND ITS APPLICATIONS. Hansen, L. E.; Wogman, N. A.; Perkins, R. W.; Clayton, E. D. (Battelle-Northwest, Richland, Wash.). Nucl. Technol.; 15: No. 3, 422-30 (Sep 1972).

Calculations were performed which indicate that thermal flux

increases of a factor of 20 should be possible through the optimal use of fissile material regions surrounding ^{252}Cf sources. For demonstration purposes, a subcritical system was assembled and used for trace element determination in marine life, terrestrial biota, and mineral specimens. This multiplying assembly consisted of PuO_2 admixed with polystyrene and reflected with Plexiglas. Source neutrons were provided by a 200 microgram ^{252}Cf source. A neutron flux amplification of about fivefold was achieved at the ^{252}Cf source, while an amplification of some 40-fold was obtained 16 cm from the source where the majority of source exposures would be made. (auth)

46856 REMARK ON NEUTRON SLOWING DOWN. Dvorak, S.; Kropac, B. (Tesla, Roznov, Czech.). Czech. J. Phys.; 22: No. 6, 459-67(1972).

The density, $W_n(U)$, of the neutron lethargy probability distribution after the n^{th} collision with nuclei of a moderator was found by using the integral representation of the Dirac δ function. It was assumed that the collisions are elastic and lethargy increments in individual collisions are independent. This procedure leads to expression of $W_n(U)$ by an improper integral, which is in turn converted to a finite sum. In addition, the asymptotic behavior of $W_n(U)$ for a great number of collisions n was determined. (auth)

46857 DISAPPEARANCE OF NEUTRON DIFFUSION LENGTH OF BERYLLIUM AND GRAPHITE ACCORDING TO TEMPERATURE. Yamagishi, Tomejiro; Deguchi, Ichiro; Sekiya, Tamotsu (Osaka Univ.). J. Nucl. Sci. Technol. (Tokyo); 9: No. 6, 375-8(Jun 1972).

A modified synthetic scattering kernel was used to examine the temperature dependence of the neutron diffusion length in beryllium and graphite. The critical temperature for the disappearance of the diffusion length was determined for both cases. (J.W.R.)

46858 SLOW NEUTRON SPECTRA IN THE LIQUID AND SOLID METHANE. Inoue, Kazuhiko (Hokkaido Univ., Sapporo); Otomo, Norio; Utsuro, Masahiko; Fujita, Yoshiaki. J. Nucl. Sci. Technol. (Tokyo); 9: No. 6, 374-5(Jun 1972).

Neutron spectra in cold methane were measured at four temperatures from 13 to 112°K. A pulsed source and the time-of-flight technique were used. The results show that very cold solid methane should be an effective moderator for cold pulsed neutron sources. (J.W.R.)

46859 PARALLEL SHOOTING METHOD FOR BOUNDARY-VALUE PROBLEMS: APPLICATION TO THE NEUTRON TRANSPORT EQUATION. Canosa, Jose; Penafiel, H. R. (IBM Data Processing Division Scientific Center, Palo Alto, Calif.). IBM J. Res. Develop.; 16: No. 4, 354-64(Jul 1972).

A direct method is given for the solution of the spherical harmonics approximation to the Boltzmann equation for neutron transport in slab geometry. The roundoff instability of the problem is eliminated by performing linear transformations of the matrices involved, which ensure that the matrix columns are linearly independent. The novelty of the method lies in that only a minimum number of matrix transformations is performed, the precise number being determined dynamically, efficiently, and in a new way by the program itself in the course of the computation. (auth)

Nuclear Properties and Reactions, A ≤ 5

Refer also to abstract: 46887.

46860 (LYCEN-7186) FINAL STATE INTERACTION BETWEEN THE PROTON AND THE NEUTRON FROM THE $^1\text{H}(d, pn)^1\text{H}$, $^2\text{H}(d, pn)^2\text{H}$, $^3\text{He}(d, pn)^3\text{He}$, AND $^4\text{He}(d, pn)^4\text{He}$ REACTIONS. Ille, Bernard (Lyon Univ. (France). Institut de Physique Nucleaire). 1971. 55p. (In French). Dep. NTIS (U. S. Sales Only).

Thesis.

The following reactions were studied at 27.5 MeV: $d(p, pn)p$, $d(d, pn)d^*$, $d(^3\text{He}, pn)^3\text{He}$ and $d(^4\text{He}, pn)^4\text{He}$. The experimental results were analyzed using the Watson-Migdal final interaction model and the Phillips-Griffy-Biedenharn sequential model to establish the production of the d^* singlet. Both models satisfactorily explain the final state interaction between the proton and the neutron in $d + p$, $d + d$, and $d + ^4\text{He}$ reactions. $^4\text{He}^*$ and $^4\text{Li}^*$ resonances complicate the $d + ^3\text{He}$ reaction. From the point of view of the deuteron singlet, these different reactions establish with certainty a peak in the final state interaction corresponding to the production of d^* considered as a pseudo-particle with isospin $T = 1$. (France)

46861 $^2\text{H}(n, p)2n$ REACTION AT 50 MeV AND THE NEUTRON-NEUTRON SCATTERING LENGTH. Stricker, A. (Tokyo Univ.); Saji, Y.; Ishizaki, Y.; (and others). Nucl. Phys.; A190: No. 2, 284-92(1972).

The proton energy spectrum from the deuteron break-up induced by 50 MeV neutrons has been measured with a broad-range magnetic analyzer followed by a 200-proportional-counter array and a plastic scintillation counter at the mean proton angle of 3.8° in the lab system. The measured spectrum was compared with the theoretical one calculated as a function of the neutron-neutron scattering length a_{nn} on the basis of the impulse approximation. From this comparison, $a_{nn} = -21.7 \pm 1.2$ fm was extracted at the effective range of 2.60 fm. (auth)

46862 NON-RESONANT ATTRACTION BETWEEN NEUTRONS IN THE $\pi^- + ^3\text{He} \rightarrow \pi^+ + 3n$ REACTION. Phillips, A. C. (Univ. of Manchester, Eng. Daresbury Nuclear Physics Lab., Eng.). Phys. Lett.; 33B: No. 4, 260-2(26 Oct 1970).

It is shown that the tendency for a low energy three-neutron system to be emitted in the $\pi^- + ^3\text{He} \rightarrow \pi^+ + 3n$ reaction does not imply the existence of a three-neutron resonance. (auth)

46863 ELASTIC SCATTERING OF PROTONS BY DEUTERONS AT ENERGIES OF 14, 16, AND 17.6 MeV. Kiseleva, Z. P.; Skorodumov, B. G.; Radyuk, G. A.; Rasulov, E. N.; Talanin, Yu. N. (Inst. of Nuclear Physics, Tashkent). Izv. Akad. Nauk Uzb. SSR, Ser. Fiz.-Mat. Nauk; No. 4, 83-5(1971). (In Russian).

The angular distributions of protons scattered by deuterated polyethylene were measured from 20 to 150° . The 17.6-MeV protons were generated in a cyclotron; energy reduction to 16 or 14 MeV as desired was accomplished by aluminum foils. The results obtained may be used in phase analysis. (1 figure, 1 table) (K.S.W.)

46864 ANALYSIS OF NEUTRON-DEUTERON ELASTIC SCATTERING AT LOW ENERGY. Viennet, R. (Univ., Neuchatel, Switzerland). Nucl. Phys.; A189: No. 2, 424-38(1972). (In French).

The results of an effective-range analysis of elastic $n-d$ scattering below the deuteron break-up threshold are given. Total spin and orbital angular momentum are supposed to be conserved, and J degeneracy is admitted. The S -, P -, and D -states only are taken into account. No confirmation of the pole of Van Oers and Seagrave in the effective-range function of the ^2S state was obtained. (9 figures, 4 tables) (auth)

46865 DEUTERON-DEUTERON SCATTERING IN THE CASE OF NONCONSERVATION OF CHANNEL SPIN. Pham Dinh Lien (Institut des Sciences Nucleaires, Grenoble, France). Nucl. Phys.; A189: No. 3, 665-72(1972).

Deuteron-deuteron elastic scattering is studied in case of non-conservation of the channel spin, that is, the possibility of transitions between singlet and quintet channel-spin states. The mixing parameter, which measures the coupling between two different orbital angular momenta, is also taken into account in the calculation of the transition-matrix amplitudes. Expressions are given for the cross section and the polarization in terms of the phase shifts. (auth)

46866 CHARGE-DEPENDENT EFFECTS IN THE PHOTO-DISINTEGRATION OF ^4He . Londergan, J. T. (Indiana Univ., Bloomington. Case Western Univ., Cleveland); Shakin, C. M. Phys. Rev. Lett.; 28: No. 26, 1729-32(26 Jun 1972).

$^4\text{He}(\gamma, p)^3\text{H}$ and $^4\text{He}(\gamma, n)^3\text{He}$ cross sections have been calculated for photon energies from the (γ, n) threshold to 34 MeV. Effects of Coulomb interactions, channel spin mixing, and mass differences have been taken into account using a coupled-channel continuum-shell-model calculation of $p\text{-}^3\text{H}$ and $n\text{-}^3\text{He}$ elastic and charge-exchange scattering. The ratio $\sigma(\gamma, p)/\sigma(\gamma, n)$ is predicted to be close to 1, except at energies very close to the (γ, n) threshold. (auth)

46867 MAGNETIC MOMENTS OF ^3He AND ^3H . Gerstenberger, R. V.; Nogami, Y. (McMaster Univ., Hamilton, Ont.). Phys. Rev. Lett.; 29: No. 4, 233-7(24 Jul 1972).

The electromagnetic structure of a nucleon in a nucleus is different from that of a free nucleon. This difference is examined for the two-pion part of the form factor by considering interactions with other nucleons in the course of pion emission and absorption, but without exchanging charge. For ^3He and ^3H , the isovector part of the magnetic moment increases by an amount which is comparable to the outstanding discrepancy between the experimental and theoretical values. (auth)

46868 EVIDENCE FOR AN ANOMALY IN THE EXCITATION FUNCTION OF THE REACTION $^2\text{H}(p, \gamma)^3\text{He}$. Chang, C. C.; Diener,

E. M.; Ventura, E. (Stanford Univ., Calif.). *Phys. Rev. Lett.*; 29: No. 5, 307-9(31 Jul 1972).

The excitation function for the capture reaction ${}^2\text{H}(\text{p},\gamma){}^3\text{He}$ was measured in the range $E_p = 10$ to 17.5 MeV. An anomaly was observed and is interpreted as a broad resonance in ${}^3\text{He}$ at an excitation energy of 14.5 ± 0.5 MeV with a width of about 2 MeV. (auth)

46869 EXPERIMENTAL INVESTIGATION OF THE REACTION $(\pi^-, \pi^+ p)$ ON ${}^6\text{Li}$ AND ${}^{12}\text{C}$ AT A MOMENTUM OF 1 GeV/c. Bayukov, Yu. D.; Vorob'ev, L. S.; Grigor'ev, E. A.; (and others) (Inst. of Theoretical and Experimental Physics, Moscow). pp 410-22 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The reactions $\pi^- + {}^{12}\text{C} \rightarrow \pi^- + \text{p} + {}^{11}\text{B}$ and $\pi^- + {}^6\text{Li} \rightarrow \pi^- + \text{p} + {}^5\text{He}$ were investigated with a 1 GeV beam of π^- -mesons from a 7 GeV proton synchrotron. A magnetic spectrometer with four spark chambers was employed in analyzing the momentum of the scattered π^- -mesons. The reactions may be adequately described with a polar diagram; however, at large imparted momenta other diagrams provide a significant contribution. (K.S.W.)

46870 NEW RESULTS ON $(\pi^+, 2p)$ REACTIONS IN LIGHT NUCLEI. Bressani, T.; Charpak, G.; Favier, J.; Meyerhof, W. E.; Zupanic, C. (CERN, Geneva). pp 444-56 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Some results on $(\pi^+, 2p)$ reactions on O, ${}^4\text{He}$, ${}^6\text{Li}$, ${}^{12}\text{C}$, ${}^{14}\text{N}$, and ${}^{16}\text{O}$ at 76 MeV are presented. The protons were detected in coincidence at a mean angle of 76° with respect to the incident pion beam. The excitation energy of the residual system was determined. Cross sections were derived assuming a peripheral mechanism. The results are compared with other data and with theory. (J.W.R.)

46871 STUDY OF ${}^2\text{H} + {}^3\text{He}$ ELASTIC SCATTERING AND THE REACTION ${}^2\text{H}({}^3\text{He}, {}^4\text{He}){}^1\text{H}$ BETWEEN 17.5 AND 44.1 MeV. King, Thomas Richard. Boulder, Colo.; Univ. of Colorado (1971). 198p. University Microfilms Order No. 72-17,276.

Thesis.

An investigation of ${}^2\text{H} + {}^3\text{He}$ elastic scattering and the reaction ${}^2\text{H}({}^3\text{He}, {}^4\text{He}){}^1\text{H}$ has been conducted at bombarding energies between 17.5 and 44.1 MeV to determine whether any states in ${}^5\text{Li}$, with appreciable widths for decay into two-body final states, exist in the region of excitation from 23.4 to 34 MeV in ${}^5\text{Li}$. An excitation function for ${}^2\text{H} + {}^3\text{He}$ elastic scattering was measured at three laboratory scattering angles, and for ${}^2\text{H}({}^3\text{He}, {}^4\text{He}){}^1\text{H}$ at four laboratory scattering angles. No evidence was found for virtual states of the nucleus ${}^5\text{Li}$ having appreciable two-body decay widths between 23.4 and 30.4 MeV excitation. Any such states have widths less than 200 keV between 23.4 and 25 MeV, less than 1.4 MeV between 25 and 30.4 MeV, less than 100 keV between 30.4 and 31 MeV, and less than 1 MeV between 31 and 34 MeV excitation in ${}^5\text{Li}$. Complete angular distributions for ${}^2\text{H} + {}^3\text{He}$ elastic scattering and the reaction ${}^2\text{H}({}^3\text{He}, {}^4\text{He}){}^1\text{H}$ were taken at several bombarding energies between 18.5 and 44.1 MeV. A partial-wave decomposition of the elastic scattering angular distributions has been performed. No anomalous behavior is observed. The angular distributions were also fitted with a linear Legendre polynomial series. Finally, the angular distributions have been compared to the predictions of two simple direct-reaction theories: the Butler stripping theory and the diffraction theory. The fits to the Butler theory are satisfactory, and the fits to the diffraction theory are remarkably good. Absolute differential cross sections were measured to an accuracy of 3%. At bombarding energies below 21.6 MeV they supplement existing data of comparable accuracy. Between 34.8 and 40.5 MeV bombarding energy, existing data are sparse, and of only moderate (14%) accuracy. The present work extends the data in this region, and improves the absolute accuracy by a factor of five. The differential cross sections measured above 40 MeV in the present work are the only data in this energy region. (Diss. Abstr. Int., B)

46872 DIFFERENTIAL CROSS SECTIONS FOR SMALL ANGLE n - p SCATTERING AT 24 MeV. Masterson, Thomas G. Madison, Wis.; Univ. of Wisconsin (1972). 163p. University Microfilms Order No. 72-13,984.

Thesis.

The differential cross section for n - p elastic scattering was measured at 19.5° and 25.1° (lab) for an incident neutron energy of 24 MeV. The neutron source was the $\text{T}(\text{d}, \text{n}){}^4\text{He}$ reaction at 0° . Both target and detector were liquid scintillators and neutrons were detected in each by the recoil protons produced. A time-of-flight coincidence requirement between target and detector

pulses reduced background. A neutron-gamma pulse shape discrimination system eliminated gammas from the detecting scintillator. An on-line computer stored and analyzed the recoil proton pulse height spectra. Normalization of the scattering data to the 0° neutron flux yielded an absolute differential cross section. A change in the flight time requirement specified the accidental coincidence background. The insertion of a shadow bar between target and detector determined room scattered backgrounds. Corrections were made for finite target and detector thicknesses, finite solid angles, wall effects and multiple scattering. The total cross sections calculated by Hopkins and Breit were used to determine detector efficiencies at different energies and relativistic kinematics to convert angles and cross sections from the lab to the CM system. The cross section measured at a CM scattering angle of 39.3° is 31.20 ± 0.45 mb/sr. This value agrees with the cross section predicted by Hopkins and Breit which is 31.3 mb/sr using the LRL phase shifts or 31.0 mb/sr using the Yale phase shifts. The cross section measured at a CM scattering angle of 50.5° is 30.7 ± 0.6 mb/sr. This value also agrees with the cross section predicted by Hopkins and Breit which is 31.0 mb/sr using the LRL phase shifts or 30.7 mb/sr using the Yale phase shifts. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, $6 \leq A \leq 19$

Refer also to abstracts: 46486, 46664, 46870, 46903, 46907, 46912, 46915, 46929, 47074, 47075, 47077, and 47083.

46873 (CEA-N-1522) NUCLEAR PHYSICS DEPARTMENT. Progress Report, October 1, 1970–October 15, 1971. (Commisariat à l'Energie Atomique, Saclay (France). Centre d'Etudes Nucleaires). 358p. (In French). Dep. NTIS (U. S. Sales Only).

This progress report covers the whole set of experiments and technological studies undertaken at the Nuclear Physics Department during the past year. The experimental activity at the 600-MeV electron linear accelerator (ALS) is now well in progress with beams of electrons, photons (bremsstrahlung) and pions giving interesting and sometimes unexpected results. Activity continues at the cyclotron, particularly with the study of fourth-order deformations in heavy nuclei. The tandem Van de Graaff is mainly used for systematic research on many-nucleon-transfer reactions with heavy-ion beams. The first experiments of high-resolution spectroscopy at the Saturne Synchrotron (1-GeV protons) are expected for the beginning of 1972. Giant resonance experiments and fission studies carried out by a branch recently connected with the Nuclear Physics Department, the activity of which is centered around a 60-MeV electron linac (AL 60) and reactors (EL 3 and Triton), are presented in this report for the first time. (auth)

46874 (JUL-834-RG) ABSOLUTE MEASUREMENT OF THE ${}^9\text{Be}(n, 2n)$ CROSS SECTION IN THE ENERGY REGION OF 2.37 TO 3.24 MeV. Blosser, Manfred (Kernforschungsanlage, Juelich (West Germany). Institut fuer Reaktorentwicklung). Mar 1972. 101p. (In German). Dep. NTIS (U. S. Sales Only).

Thesis. Submitted to Technische Hochschule, Aachen (West Germany).

Since the energy of DD neutrons is angle dependent, quasi-monoenergetic neutrons of 2.37, 2.53, 2.77, 2.99, 3.21, and 3.34 MeV of energy could be produced; their energy distribution was measured by a ${}^9\text{Be}$ semiconductor spectrometer. The direct determination of the $(n, 2n)$ reaction probabilities by means of a 4π Gd liquid scintillator with a double-pulse technique for identification of $(n, 2n)$ neutrons was done absolutely by determining the efficiency of the Gd detector. The applied scattering block-method for the measurement of the neutron intensity was not primarily based on the knowledge of other cross sections. After taking into account the measured energy distribution of primary neutrons and multiple scattering processes in the Be target by a simple Monte Carlo process, the ${}^9\text{Be}(n, 2n)$ cross sections are given, and possible reaction mechanisms are discussed. (17 figures, 6 tables) (auth)

46875 (KFK-EXT-10/72-1) MEAN EXCITATION ENERGIES OF THE DAUGHTER NUCLEI ON β^- -DECAY. Keller, K. A.; Muenzel, H. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Radiochemie). Feb 1972. 51p. (In German). Dep. NTIS (U. S. Sales Only).

For the prediction of unknown half-lives of β^- -decay the knowledge of mean excitation energies of the daughter nuclei is of great importance. Therefore a compilation of mean excitation energies is given in tables and figures. The accuracy of the half-lives taking into account these mean excitation energies is discussed. (auth)

46876 (ORNL-TM-3859) REVIEW OF THE CROSS SECTIONS OF THE ISOTOPES ^6Li AND ^7Li AS TABULATED ON THE ENDF/B VERSION III FILES. Tobias, Melvin (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 60p. Dep. NTIS.

The following report reviews the cross section information contained in the ENDF/B files concerning the isotopes ^6Li and ^7Li . The purpose of this study is to seek out those areas of uncertainty which may be of concern to fusion reactor technology. Other nuclides will be similarly examined and the information obtained will be used in cross section sensitivity studies for fusion reactor design. (auth)

46877 CIRCULAR POLARIZATION OF γ RAYS IN ^{18}F . A TEST FOR WEAK INTERACTION THEORIES. Wambach, U. M.; Gari, M.; Kuemmel, H. (Ruhr Univ., Bochum, Ger.). Phys. Lett.; 33B: No. 4, 253-6 (26 Oct 1970).

The circular polarization from the transition of the $(0^-, 0; 1.08 \text{ MeV})$ level to the $(1^+, 0)$ ground state in ^{18}F has been computed with two-body matrix elements and short range correlations via the Bethe-Goldstone equation for finite nuclei. Results have been obtained for many different weak interaction theories. (auth)

46878 CHARGE FORM FACTOR AND QUADRUPOLE MOMENT OF ^6Li . Jain, A. K.; Sarma, N. (Bhabha Atomic Research Center, Bombay). Phys. Lett.; 33B: No. 4, 271-3 (26 Oct 1970).

The charge density of ^6Li is calculated using antisymmetrized and modified cluster model wave functions. Satisfactory agreement with the measured charge form factor and quadrupole moment has been obtained. Exchange terms have been found to be important in both calculations. (auth)

46879 BETA-DELAYED PROTON DECAY OF ^{13}O : A VIOLATION OF MIRROR SYMMETRY. Esterl, J. E.; Hardy, J. C.; Sextro, R. G.; Cerny, J. (Univ. of California, Berkeley). Phys. Lett.; 33B: No. 4, 287-90 (26 Oct 1970).

Beta-delayed protons, $E_{1ab} > 1.2 \text{ MeV}$, were observed following the decay of ^{13}O , $\tau_{1/2} = 8.95 \pm 0.20 \text{ ms}$. Comparison of ^{13}O decay rates with those of the mirror nuclide ^{13}B indicates a 15% deviation from mirror symmetry. (auth)

46880 SPECTRA OF NEUTRONS EMITTED AFTER MUON CAPTURE IN ^{16}O , ^{28}Si , ^{32}S AND ^{40}Ca . Hill, Lemuel L. (Naval Ordnance Lab., Silver Spring, Md.); Ueberall, H. Nucl. Phys.; A190: No. 2, 341-61 (1972).

On the basis of the particle-hole model for the closed-(sub)shell nuclei ^{16}O , ^{28}Si , ^{32}S and ^{40}Ca , a calculation of the partial muon-capture rates to the $T = 1$ states of the model (which mainly represent the giant resonance), as well as of the spectra of neutrons emitted in the decay of these states to the ground and excited states of the corresponding daughter nuclei is presented. Improved positions and empirical total widths for the resonant states have been used as obtained from photonuclear and electron scattering studies of the levels. The theoretical total muon-capture rates are in good agreement with measured rates. The calculated neutron spectra are compared to recently measured spectra following muon capture in ^{16}O , ^{32}S and ^{40}Ca . (auth)

46881 TEST OF TIME REVERSAL INVARIANCE THROUGH DETAILED BALANCE OF THE REACTION $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$. Von Wimmersperg, U.; Kernel, G.; Allardice, B. W.; Mason, W. M.; Tanner, N. W. (Nuclear Physics Lab., Oxford). Phys. Lett.; 33B: No. 4, 291-3 (26 Oct 1970).

The angular distributions of $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ and its inverse $^{16}\text{O}(\gamma, \alpha)^{12}\text{C}$ have been measured at an excitation of 13.1 MeV where there is a strong asymmetry favoring the forward direction due to $E1/E2$ interference. The asymmetries for (α, γ) and (γ, α) were the same within one standard deviation, and reveal no evidence of a failure of time reversal invariance. (auth)

46882 ENERGY LEVELS OF LIGHT NUCLEI $A = 18$ TO 20. Ajzenberg-Selove, F. (Univ. of Pennsylvania, Philadelphia). Nucl. Phys.; A190: No. 1, 1-196 (1972).

A compilation of energy levels of $A = 18, 19$ and 20 nuclei, with emphasis on material leading to information about the structure of the $A = 18-20$ systems, is presented. (auth)

46883 ELASTIC SCATTERING OF ^6Li IONS. Davydov, V. V.; Novatskii, B. G.; Ogloblin, A. A.; (and others). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2399-2405 (Nov 1971). (In Russian).

The elastic scattering of ^6Li ions with an energy of 30 MeV by the nuclei ^{12}C , ^{16}O , ^{28}Si , ^{40}Ca , ^{56}Ni , ^{120}Sn , ^{197}Au , and ^{208}Pb was studied. The maximum possible angles (170 to 175°) were attained in all cases. The scattering of 36.6 MeV α particles by ^6Li nuclei

was similarly measured. The parameters of the Saxon-Woods potential are tabulated for each of the eight nuclei. (K.S.W.)

46884 EVIDENCE FOR SHAPE OR QUASIMOLECULAR RESONANCES IN THE $^{16}\text{O} + ^{16}\text{O}$ INTERACTION. Singh, P. P. (Indiana Univ., Bloomington); Sink, D. A.; Schwandt, P.; Malm, R. E.; Siemssen, R. H. Phys. Rev. Lett.; 28: No. 26, 1714-17 (26 Jun 1972).

Differential cross sections have been measured for the reaction $^{16}\text{O}(^{16}\text{O}, ^{12}\text{C})^{20}\text{Ne}$ leading to the first three states of ^{20}Ne at c.m. energies from 17.5 to 30 MeV and c.m. angles of approximately 62°, 76°, and 90°. The yield curves are characterized by (1) gross structures and narrow structures, and (2) strong angular and cross correlations. It is suggested that these results present direct evidence for shape or quasimolecular resonances in the $^{16}\text{O} + ^{16}\text{O}$ system. (auth)

46885 MECHANISM OF THE $^9\text{Be}(p, p_1)^9\text{Be}$ 1.68 MeV REACTION. Vasil'ev, S. S.; Galakhmatova, B. S.; Dem'yanova, A. S.; (and others). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2364-7 (Nov 1971). (In Russian).

Measurements of the angular distributions of elastic and inelastic scattering of protons by ^9Be nuclei were conducted at energies of 6.36 and 6.48 MeV and in the angular range 15 to 150°. The results of the measurements according to the excitation of the 1.68 MeV, $1/2^+$, level were compared with calculated values based on the mechanism characterized by a quadrangular diagram. (tr-auth)

46886 ORBITAL REARRANGEMENT IN LIGHT NUCLEI. Muether, H.; Goeke, K.; Faessler, Amand (Kernforschungsanlage, Juelich, Ger.). Z. Phys.; 253: No. 1, 61-70 (1972).

Orbital rearrangement effects due to the removal of a proton are studied. Calculations for the nuclei ^{16}O , ^{20}Ne , and ^{40}Ca are performed using the Yale potential and the Brink-Boeker force B1. The odd-mass nuclei are not restricted to time reversal symmetry—in this way a large overestimation of the rearrangement energies is avoided. Although the orbital rearrangement energies are small ($<1.5 \text{ MeV}$), by utilizing density-independent forces one can find larger changes in the wave functions due to the rearrangement, as exhibited in the discussion of the quadrupole moments and the mass distributions. (1 figure, 4 tables) (auth)

46887 INVESTIGATION OF TEMPERATURE DEPENDENCE OF CROSS SECTION OF COLD- AND THERMAL-NEUTRON SCATTERING BY O_2 AND D_2 MOLECULES. Vertebnyi, V. P.; Gul'ko, V. A.; Maistrenko, A. N.; Matviishina, N. N.; Razbudei, V. F. (Inst. of Nuclear Research, Kiev). Ukr. Fiz. Zh. (Russ. Ed.); 16: No. 12, 1967-74 (Dec 1971). (In Russian).

The total cross sections for cold- and thermal-neutron scattering by diatomic gases (oxygen and deuterium) at temperatures of -100 to $+100^\circ\text{C}$ are reported. The measured cross sections are compared with those calculated by the theories of Krieger-Nelkin and Young-Koppel. (4 figures, 2 tables) (auth)

46888 ANGULAR AND ENERGY DISTRIBUTIONS OF PROTONS IN (γ, p) AND (e, p) REACTIONS ON ^{12}C NUCLEUS FOR 130 MeV EXCITATION ENERGY. Egiyan, K. Sh.; Bohek, G. L.; Kulibaba, V. I.; Grishaev, I. A. (Inst. of Physics, Erevan). Physical-Technical Inst., Kharkov). Izv. Akad. Nauk Arm. SSR, Ser. Fiz.; 6: No. 3, 161-7 (1971). (In Russian).

The angular and the energy distributions of the secondary protons from (γ, p) and (e, p) reactions on ^{12}C at 130 MeV excitation energy are studied. The experimental results are compared with the quasi-deuteron and one-particle absorption models of γ quanta in the nucleus. For both cases the parameters of the proton wave function in the nucleus are estimated. (4-figures, 1 table) (auth)

46889 SOME REMARKS ON THE TRANSPARENCY OF COMPLEX NUCLEI TO PHOTOPRODUCED CHARGED PIONS AT INTERMEDIATE ENERGIES. de Carvalho, H. G. (Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro); Martins, J. B.; Tavares, O. A. P.; Nazareth, R. A. M. S.; di Napoli, V. Lett. Nuovo Cim.; 4: No. 9, 365-8 (1 Jul 1972).

Nuclear transparencies of carbon-12, aluminum-27, manganese-55, ruthenium-100, and uranium-238 to photoproduced π^+ and π^- are calculated as functions of the energies of the respective particles. (L.B.S.)

46890 NEW EXPERIMENTAL DATA ON THE FINE STRUCTURE OF DIPOLE RESONANCE. Goryachev, B. I.; Ishkhanov, B. S.; Shevchenko, V. G. pp 362-78 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Results are reported on cross section measurements on (γ, n) -

and (γ, p) -reactions obtained with a betatron for $E_{\gamma, \max} = 35$ MeV, in energy steps of 20, 50, and 100 keV. Light, average, and heavy nuclei were investigated with the above reactions: ^{12}C , ^{16}O , ^{24}Mg , ^{26}Mg , ^{27}Al , ^{28}Si , ^{31}P , ^{32}S , ^{40}Ca , ^{51}V , ^{52}Cr , ^{58}Ni , ^{60}Ni , ^{63}Cu , ^{89}Y , ^{90}Zr , ^{93}Nb , $^{92,98}\text{Mo}$, $^{110-116}\text{Cd}$, ^{163}Ho , ^{166}Er , ^{181}Ta , ^{208}Pb . Cross sections were also measured for (γ, f) -reactions on ^{232}Th , ^{235}U , and ^{238}U . (K.S.W.)

46891 DIRECT REACTIONS INDUCED BY HIGH ENERGY PROTONS. Palevsky, Harry (Brookhaven National Lab., Upton, N. Y.). pp 423-43 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Some experiments on direct nuclear reactions induced by high energy protons are surveyed. The type of information obtained in such experiments is discussed. Some conclusions regarding nucleon-nucleon correlations in nuclei which were drawn from the experiments are presented. (J.W.R.)

46892 QUADRUPOLE MOMENTS OF LIGHT NUCLIDES. Ruehl, Franklin R. Jr. (Univ. of California, Los Angeles). Indian J. Phys.; 45: No. 4, 149-54 (Apr 1971).

As predicted by a trio of effective nucleon-nucleon interaction potentials, the quadrupole moments of seven light axially symmetric nuclei are analyzed. Based on agreement with experiment, the ground state shapes favored for these nucleides are then indicated. (auth)

46893 DEVELOPMENT OF A TWO-SECTION Ge(Li) COMPTON POLARIMETER WITH APPLICATIONS IN THE $2s-1d$ SHELL. Lumpkin, Alexander Henderson, Baltimore; Johns Hopkins Univ. (1971). 121p. University Microfilms Order No. 72-16,910.

Thesis.

A two-section Ge(Li) Compton polarimeter was developed, and its sensitivity to polarization was determined by an asymmetry calculation and a calibration with highly polarized gamma rays. This capability for absolute measurement of the linear polarization of gamma rays with high energy resolution was utilized experimentally with direct angular correlation measurements to study reaction gamma rays. Theoretical gamma ray linear polarization and angular distribution formulae in terms of the phase-defined reduced matrix elements of Rose and Brink are used in the analysis to determine, independently of the reaction mechanism, multipole mixing ratios of the gamma rays. These mixing ratios are compared in both magnitude and sign to the predictions of nuclear models. In particular gamma ray transitions of ^{19}F and ^{27}Al were studied. In ^{19}F the 1.459 MeV level was the major concern and the three transitions from this level which was populated by inelastic proton scattering were studied. The mixing ratios were determined to be $-0.1 < \delta_{M_2/E_1} < 0.0$ for the $1.459 \rightarrow 0.197$ MeV transition, $0.30 < \delta_{E_2/M_1} < 0.38$ for the $1.459 \rightarrow 0.110$ MeV transition, and $|\delta_{M_2/E_1}| < 0.06$ for the $1.459 \rightarrow 0.0$ MeV transition. The results indicate the level in question cannot be explained as a $p_{3/2}$ proton hole coupled to the pure ground state rotational band as has been accepted in the past. In ^{27}Al the transitions to the ground state from the 2.21, 2.73, 2.98, and 3.00 MeV levels were studied. Mixing ratios δ_{E_2/M_1} for the transitions were determined to be $\delta(2.21) = -0.57 \pm 0.08$, $\delta(2.73) = 0.22 \pm 0.09$, and $\delta(2.98) = 0.05 \pm 0.03$. The 3.00 MeV transition is apparently a pure E2 transition. These results are compared to the prediction of the rotational-vibrational model of Röpke. These applications demonstrate the usefulness of the two-part Ge(Li) polarimeter in the study of reaction gamma rays. (Diss. Abstr. Int., B)

46894 SMALL ANGLE INELASTIC SCATTERING OF 40 MeV PROTONS BY ^{12}C , ^{40}Ca , ^{56}Fe , AND ^{58}Ni . Ingham, David Ransford. Santa Barbara, Calif.; Univ. of California (1971). 93p. University Microfilms Order No. 72-18,240.

Thesis.

Angular distributions have been measured for inelastic scattering of 40 MeV protons to several states in ^{12}C , ^{40}Ca , ^{56}Fe and ^{58}Ni in the angular range from 6° to 20° . The scattered protons were stopped in a "stack" of three Si(Li) detectors and a computer was used to reject false events which had abnormal distributions of energy loss. The results are compared to DWBA calculations, to previous Coupled Channels calculations and to the Parity Rule. Discrepancies are noted between the data and the DWBA, but the Coupled Channels formalism appears capable of resolving these at least in part. (Diss. Abstr. Int., B)

46895 INVESTIGATION OF TWO NUCLEON TRANSFER REACTIONS IN ^{19}F USING 45 MeV PROTONS. Huber, Kent Alvin. Los Angeles; Univ. of Southern California (1972). 149p. University Microfilms Order No. 72-17,474.

Thesis.

The $^{19}\text{F}(p, ^3\text{He})$ and $^{19}\text{F}(p, t)$ reactions leading to the mirror nuclei ^{17}O and ^{17}F , respectively, were studied at a bombarding energy of 45 MeV. Absolute experimental differential cross sections were obtained for transitions to the ground, 0.871, 3.06, and 3.85 MeV levels of ^{17}O and to the ground, 0.500, 3.10, and 3.85 MeV levels of ^{17}F . Targets consisted of commercially available Teflon (CF_2) foils and target deterioration was carefully monitored. Data were collected for center-of-mass scattering angles between 10° and 110° . All distributions exhibit the strong forward peaking indicative of a direct transfer of two nucleons from the target to the incident probe. Total cross sections for the $^{19}\text{F}(p, ^3\text{He})$ reaction to the ground and 0.871 MeV states of ^{17}O were found to be 1.855 ± 0.005 millibarn and 0.660 ± 0.006 millibarn, respectively. Total cross sections for the $^{19}\text{F}(p, t)$ ground and 0.500 MeV transitions were measured as 2.58 ± 0.02 millibarn and 1.56 ± 0.09 millibarn. The experimental data are compared to the results of a DWBA analysis in which the initial nucleus is assumed to consist of an inert ^{16}O core with three nucleons in the $1d_{5/2}$ and $2s_{1/2}$ orbitals. The configuration of the ground state for both ^{17}F and ^{17}O assumed the ^{16}O core plus one nucleon in the $1d_{5/2}$ orbital. The first excited states assumed the inert core with one nucleon in the $2s_{1/2}$ orbital. Angular distributions are determined assuming the transfer of a heavy cluster of two nucleons. These calculations utilized the DWBA code JULIE and used optical model parameters from the literature to describe the elastic scattering in the entrance and exit channels. The predictions for the $^{19}\text{F}(p, t)$ ^{17}F $L = 0$ transfer to the 0.500 level exhibits the same oscillations as the data but fails to predict the correct magnitude. Results for the analogous $^{19}\text{F}(p, ^3\text{He})$ reaction to the 0.871 MeV levels also display the correct frequency but are displaced approximately 5° toward higher angles. Angular distributions predicted for the ground state $L = 2$ transitions are not as representative of the experimental results. (Diss. Abstr. Int., B)

46896 STUDY OF THE CONTINUUM STATES OF THE ^7Be SYSTEM BELOW 10 MeV. Gintner, Henry. Cleveland; Case Western Reserve Univ. (1972). 103p. University Microfilms Order No. 72-18,692.

Thesis.

A distorted wave Born approximation K-matrix formalism is employed to study the scattering states of the ^7Be system below 10 MeV. The ^7Be system is considered to be composed of three elementary particles: an α particle, a deuteron and a proton. Optical potentials that depend on angular momentum are used to describe the elastic scattering cross sections for $^6\text{Li} + p$ and $^4\text{He} + ^3\text{He}$. A small D-state admixture in the ground state of ^6Li leads to a large resonant structure in the reaction $^4\text{He}(^3\text{He}, p)^6\text{Li}$ in the energy region corresponding to the $^4\text{P}_{3/2}$ level of ^7Be . The fit to the experimental cross sections in the region around the $^4\text{P}_{3/2}$ level is good. The K-matrix method is compared with the standard DWBA T-matrix method. The K-matrix method and the T-matrix method give essentially the same results outside the energy region of the $^4\text{P}_{3/2}$ level. In the energy region of the $^4\text{P}_{3/2}$ level the K-matrix method is superior since it takes into account the loss of flux from the incident channel to the outgoing channel. Inclusion of other channels and more cluster structures in ^6Li and ^7Be should lead to improved results. (Diss. Abstr. Int., B)

46897 $^3\text{He}(^6\text{Li}, \alpha p)^4\text{He}$ REACTION AT 1.54 MeV. Thompson, Daniel Thomas. Pullman, Wash.; Washington State Univ. (1972). 106p. University Microfilms Order No. 72-18,493.

Thesis.

Asymmetry in the proton yield about the ^5Li direction in the sequential reaction ($^3\text{He} + ^6\text{Li} \rightarrow \alpha_1 + ^5\text{Li} \rightarrow \alpha_1 + \alpha_2 + p$) has been measured for the ground and first excited states of ^5Li at a beam energy of 1.54 MeV. The initial reaction mechanism has been confirmed to be a pickup reaction in which the incoming ^3He picks up a neutron from the target ^6Li nucleus. A model based upon the physical aspects of the short lifetime of the ^5Li nucleus is capable of explaining the asymmetry for both the ground and first excited states of ^5Li . This model assumes that the ^5Li retains a "memory" of its formation and then shows that protons will be preferentially emitted forward of the direction of recoil of the ^5Li . The possibility that distorted wave Born approximation (DWBA) stripping theory may explain the asymmetric results has also been investigated. This is significant since the DWBA theory makes no assumptions about the lifetime of the ^5Li nucleus. The effects of Coulomb final state interactions (FSI) among the final state particles has been studied. Coulomb FSI do not contribute to the observed asymmetry, but they can significantly affect observed spectra for many reactions. An experimental technique has been developed which makes it possible to separate kinematically interfering events for three body final state reactions in which two or more of the final state particles are nonidentical. (Diss. Abstr. Int., B)

46898 EXPERIMENTAL INVESTIGATION OF THE $^{12}\text{C}+^{12}\text{C}$ REACTION AT LOW ENERGIES. Mazarakis, Michael Geras. Philadelphia; Univ. of Pennsylvania (1971). 203p. University Microfilms Order No. 72-17,396.

Thesis.

The reactions $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ and $^{12}\text{C}(^{12}\text{C}, p)^{23}\text{Na}$ have been studied in order to extend the available excitation curves down to lower energies and to provide reliable predictions of the reaction cross sections in the energy region of interest for stellar thermonuclear carbon burning. A carbon foil target of about $30 \mu\text{g}/\text{cm}^2$ ($65 \mu\text{g}/\text{cm}^2$ for measurements of $E_{\text{c.m.}} < 2.8 \text{ MeV}$) was bombarded with a beam of $^{12}\text{C}^{++}$ of $0.5 - 1 \mu\text{Amp.}$ from the University of Pennsylvania Tandem Accelerator. The measurements covered a range of 2.4 to 5 MeV in the center-of-mass in steps of 125 keV. Many excited state transitions have been observed for the protons and alpha particles. The spectra of protons and alphas were obtained at eight forward laboratory angles: 20° , 30° , 40° , ..., 90° . Four silicon solid state detectors were used in two successive runs for each energy studied. The obtained angular distributions were transformed separately to the center-of-mass system and integrated to give the total cross sections for the alpha particles and protons. The present measurements give cross sections in agreement with those obtained by J. R. Patterson, H. Winkler and C. S. Zaidins in the overlapping region and extend the available information to the actual static astrophysical region (hydrostatic carbon burning). The obtained cross sections, corrected for the transmission probability of the Coulomb barrier (nuclear factor $\bar{S}$), show a continuation of the prominent "structure" observed near 6 MeV by E. Almquist, D. A. Bromley and J. A. Kuehner. Below 3 MeV an increase of the factor $\bar{S}$ was observed with a possible maximum near 2.4 MeV and a corresponding total reaction cross section of the order of 10^{-8} barns. This coincides with sharp resonances separately observed in the compound ^{24}Mg nucleus in the $^{16}\text{O}(^{12}\text{C}, \alpha)^{24}\text{Mg}$ reaction. The comparison of the branching ratios measured in the present work with those of the above resonances does not preclude a similar origin. The excitation functions of the different reaction products were analyzed in detail and two intermediate resonances were found below 5 MeV at 4.17 and 4.9 MeV with the same properties as those found near 6 MeV. A spin assignment was carried out and a possible $J^+ = 4^+$ has been suggested. A phenomenological analysis of the present results (reaction cross sections and branching ratios) has recently been carried out by G. J. Michaud and E. W. Vogt (M6) based on an α - α model interaction. The main conclusions of this paper are discussed with the presentation of the experimental results. (Diss. Abstr. Int., B)

46899 STUDY OF THE $^{16}\text{O}(\text{d}, \text{n})$ REACTION FROM 3 TO 4 MeV. Anderson, Bryon Don. Cleveland; Case Western Reserve Univ. (1972). 109p. University Microfilms Order No. 72-18,666.

Thesis.

The cross sections and polarizations of the neutron groups from the $^{16}\text{O}(\text{d}, \text{n})^{17}\text{F}$ reaction leading to the ground and first excited states of ^{17}F have been measured in the deuteron energy range from 3 to 4 MeV. Angular distributions were determined at three energies. Excitation functions of cross sections at two angles and of polarizations at one angle were performed. Time-of-flight techniques were employed throughout. The neutron polarimeter for the polarization measurements consisted of a gas helium-xenon scintillator with two plastic-scintillator side detectors. Calculations were performed to determine to what extent the DWBA theory could describe the experimental results. The cross-section angular distributions and the highest energy ground state polarization angular distribution could all be reproduced simultaneously with only small modifications made to optical-model parameters taken from recent elastic-scattering data analyses. The rest of the polarization data appears to be affected by large compound-nucleus contributions and could not be reproduced simultaneously. Spectroscopic factors were extracted and are in satisfactory agreement with those obtained by previous workers. (Diss. Abstr. Int., B)

46900 SEARCH FOR PARITY AND ISOBARIC SPIN VIOLATING DECAY OF THE 3.56 MeV LEVEL IN ^6Li . Kosiara, Andrzej. Cleveland; Case Western Reserve Univ. (1972). 59p. University Microfilms Order No. 72-18,703.

Thesis.

The self interacting current-current theory of weak interactions predicts the existence of a parity violating force between nucleons. A technique utilizing time-of-flight in addition to the usual energy and particle identification has been applied to the investigation of the decay of the 3.56 MeV, O^+ , $T = 1$ level in ^6Li . This level is forbidden by conservation of parity and isobaric spin to break up into a deuteron and an alpha particle. It was populated via the $^9\text{Be}(p, \alpha)^6\text{Li}$ (3.56 MeV) reaction. The proton beam of 2.56 MeV energy was supplied by the CWRU Van de Graaff accelerator at a 1 Mhz

repetition rate. This beam when reaching the target has a further time structure which consisted of 1 ns long groups of particles separated in time by 100 nsec. Most of the protons are contained in one group, called the main pulse. This group would be the most desirable to populate the 3.56 MeV in ^6Li . The subsequent parity-forbidden breakup would have to be investigated in the presence of the intense background due to elastically scattered protons. The high intensity of the elastically scattered protons from the main pulse introduces nearly 100% dead time in the electronic analysis and makes detection of a parity-forbidden event difficult. This problem could be circumvented by operating the accelerator at low current but this introduces beam instability which makes it impossible to interrelate data. The experiment was performed by electronically gating out the elastically scattered protons due to the main pulse. The products of the bombardment of the beryllium target by the subsequent 10 Mhz proton groups were observed with a scintillation detector. The low intensity of the elastically scattered protons due to those groups permitted keeping the dead time of the multichannel analyzer less than 2%. The data consist of time-of-flight spectra with the part due to the main pulse gated out. They were analyzed by the least-squares technique for the presence of deuterons from the forbidden breakup. The background was obtained by observing gated time-of-flight spectra of the same reaction at the energy where the population of the 3.56 MeV level is unlikely. No evidence of the forbidden breakup has been observed. The limit on joint parity and isobaric spin conservation is $^2F \sim 8.1 \times 10^{-6}$. (Diss. Abstr. Int., B)

46901 NUCLEAR SPINS AS DETERMINED BY THE METHOD OF GAMMA-GAMMA ANGULAR CORRELATIONS. Spyr, David Jon. Iowa City, Iowa; Univ. of Iowa (1972). 110p. University Microfilms Order No. 72-17,606.

Thesis.

The method of gamma-gamma angular correlations (Litherland and Ferguson, method D) was used to make spin determinations for levels in ^{10}B , ^{23}Na , and ^{42}K . The quadrupole/dipole amplitude mixing ratios for each transition were also measured as well as the magnetic substate populations for the initial decaying level. A new and flexible computer-based data taking system was employed to make the first measurement for the ^{42}K levels and to confirm the rest. The levels investigated together with the most probable or accepted spins were: $^{10}\text{B}_{1.740 \rightarrow 0.717 \rightarrow 0.0}$ (0, 1, 3); $^{23}\text{Na}_{2.390 \rightarrow 0.439 \rightarrow 0.0}$ ($1/2$, $5/2$, $3/2$); $^{42}\text{K}_{0.252 \rightarrow 0.104 \rightarrow 0.0}$ (4, 3, 2). (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, $20 \leq A \leq 38$

Refer also to abstracts: 46486, 46664, 46873, 46875, 46880, 46882, 46883, 46886, 46889, 46890, 46892, 46893, 46901, 46924, 46929, 47066, and 47072.

46902 (BARC-598) STUDIES ON ANALOG STATES IN ^{33}Cl BY ISOPIN-FORBIDDEN RESONANCES IN THE REACTION $^{32}\text{S}(p, \gamma)^{33}\text{Cl}$. Eswaran, M. A.; Ismail, M.; Ragoowansi, N. L. (Bhabha Atomic Research Centre, Bombay (India)). 1972. 26p. Dep. NTIS (U. S. Sales Only).

The residual activity between bursts of a mechanically chopped beam was used to measure the yield of the reaction $^{32}\text{S}(p, \gamma)^{33}\text{Cl}$ systematically in the bombarding energy range $E_p = 3.36$ to 5.41 MeV. Two $T = 3/2$ states in ^{33}Cl at $E_p = 3.371 \pm 0.005 \text{ MeV}$, $E_x = 5.550 \pm .007 \text{ MeV}$ and at $E_p = 5.282 \pm .006 \text{ MeV}$, $E_x = 7.402 \pm .008 \text{ MeV}$ were located with the resonance strengths $(2J + 1)\Gamma_p/\Gamma = 0.76 \pm 0.18 \text{ eV}$ and $1.50 \pm 0.37 \text{ eV}$, respectively. Each of these resonances was narrower than the estimated 2 keV spread in the proton beam. These two states are interpreted as the analogs of the ground state and the second excited state of ^{33}P with J^π values of $1/2^+$ and $5/2^+$, respectively. γ decay of the lower resonance, investigated with a Ge(Li) detector, shows $>88\%$ and $<12\%$ branchings to the first excited state and ground state of ^{33}Cl , respectively. The M1 strengths of these transitions are compared with those obtained from beta analog transitions and with the theoretical predictions based on many-particle shell-model calculations. (5 figures, 4 tables) (auth)

46903 (COO-1120-149) FINAL REPORT. (Kansas Univ., Lawrence, Nuclear Structure Lab.). Jun 1972. Contract AT-(11-1)-1120. 91p. Dep. NTIS.

MAGNESIUM-24—energy levels from sodium-23 (p, γ) reactions, widths and spin-parity assignments for, (E)
NITROGEN-14—proton reactions (p, γ) at 3.1 to 3.5 MeV, γ spectra and excitation function for, (E)
—proton inelastic scattering at 3.1 to 3.5 MeV, excitation function for, (E)

- OXYGEN-15—energy levels at 10.275, 10.459, and 10.466 MeV from nitrogen-14 (p, γ) reactions, spin-parity assignments and decay properties of, (E)
- PROTONS—scattering by sodium-23 at 1.4 to 3 MeV, resonance structure in excitation functions for elastic and inelastic, (E)
- SODIUM-23—proton reactions (p, γ) at 1.8 to 2.8 MeV, γ yields for, (E)
- proton reactions (p, α) at 1.4 to 3 MeV, resonance structure in excitation functions for, (E)
- proton elastic and inelastic scattering at 1.4 to 3 MeV, resonance structure in excitation functions for, (E) (J.W.R.)

46904 ENERGY LEVELS OF ^{36}K . Dzubay, T. G. (Duke Univ., Durham, N. C. Triangle Universities Nuclear Lab., Durham, N. C.); Jaffe, A. A.; Ludwig, E. J.; (and others). *Phys. Lett.*; 33B: No. 4, 302-4 (26 Oct 1970).

The $^{36}\text{Ar}(^3\text{He}, t)^{36}\text{K}$ reaction has been observed at 22.17 MeV, and a Q value of -12.93 ± 0.04 MeV deduced. Excited states in ^{36}K were seen at 0.800, 1.115, 1.59, 1.67, 1.89, 2.27, 2.41, 2.56, 2.85 and 3.35 MeV. A comparison is made with levels in ^{36}Cl . (auth)

46905 LIFETIME OF THE 1.013-MeV STATE IN ^{27}Al . Kalpov, D. K.; Kosyak, Yu. G.; Smirin, L. N.; Shubnyi, Yu. K. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2323-4 (Nov 1971). (In Russian).

In an investigation of the γ resonance scattering by ^{27}Al the lifetime of the 1.013 MeV level was determined as $\tau = 1.22 \times 10^{-2}$ sec. Radioactive ^{27}Mg was used as the source of the γ radiation. (tr-auth)

46906 LOW-LYING POSITIVE-PARITY STATES OF ^{34}Cl AND ^{36}Cl FROM (p, d) REACTIONS AND SHELL-MODEL INTERPRETATIONS. Vignon, B. (Institut des Sciences Nucleaires, Grenoble, France); Longueue, J. P.; Towner, I. S. *Nucl. Phys.*; A189: No. 3, 513-25 (1972).

Low-lying positive-parity states of ^{34}Cl and ^{36}Cl nuclei have been studied from the $^{35,37}\text{Cl}(p, d)$ reactions. A DWBA analysis of the angular distributions for the strong transitions has been performed and spectroscopic factors deduced. The experimental results have been compared with shell-model calculations in the $2s_{1/2} - 1d_{3/2}$ vector space, and with recent calculations involving the full $s-d$ shell. Most of the excitation energies and spectroscopic factors were successfully reproduced. (auth)

46907 STUDY OF EXCITED STATES OF ^{21}Ne . Rolfs, C. (Univ., Freiburg i. B. Univ. of Toronto); Trautvetter, H. P.; Kuhlmann, E.; Riess, F. *Nucl. Phys.*; A189: No. 3, 641-64 (1972).

The 2.79, 3.66, 3.73 and 3.88 MeV states in ^{21}Ne have been studied with the $^{18}\text{O}(\alpha, n\gamma)^{21}\text{Ne}$ reaction at $E_\alpha = 4.9$ to 6.6 MeV. Both neutrons and γ -rays were observed. The investigation involved measurements of γ -ray angular distributions as well as $n\gamma$ and $\gamma\gamma$ angular correlations for procuring spin and multipole mixing information. The results of this investigation combined with previous work lead to respective J^π assignments of $1/2^-, 3/2^-, 5/2^+$ and $5/2^-$. The states above 4 MeV excitation energy have been studied at $E_\alpha = 7.5$ to 11.5 MeV with a $\text{Ge(Li)}-\text{NaI(Tl)}$ pair spectrometer. Two new states at $E_x = 5525.0 \pm 1.5$ and 5682 ± 2 keV have been found. Evidence is also presented for a doublet comprising the states with excitation energies 5821 ± 3 and 5823 ± 3 keV. Excitation energies, γ -ray decay schemes, mean-lives and limits on spin assignments to all the states are reported. Some of the results are compared with the available data for the mirror nucleus ^{21}Na and discussed in terms of the rotational model. (auth)

46908 LIFETIME OF THE 975-keV, $3/2^+$ STATE IN ^{25}Mg . Alexander, T. K.; Haeusser, O.; MacDonald, A. B.; Yang, J. T. (Chalk River Nuclear Labs., Ont.). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2287-91 (Nov 1971). (In Russian).

Measurements were made using the reaction $^{24}\text{Mg}(p, \gamma)^{25}\text{Mg}$ in which the velocity of the nucleus of recoil ^{25}Mg was 5.5% of the velocity of light. From the results of direct measurements of the lifetimes pertaining to the decays $975 \rightarrow 585$ and $975 \rightarrow 0$ the averaged lifetime of the 975 state was 14.6 ± 1.5 psec. (K.S.W.)

46909 GAMMA-SPECTRA AND CROSS SECTION OF THE ($n, n'\gamma$) REACTION IN THE ISOTOPES ^{27}Al , ^{51}V , ^{52}Cr , ^{55}Mn , ^{56}Fe , ^{93}Nb , ^{208}Pb , AND ^{209}Bi . Degtyarev, Yu. G.; Protopopov, V. N. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2341-4 (Nov 1971). (In Russian).

The excitation of the low-lying levels of the eight isotopes ^{27}Al , ^{51}V , ^{52}Cr , ^{55}Mn , ^{56}Fe , ^{93}Nb , ^{208}Pb , and ^{209}Bi was investigated during inelastic scattering of neutrons with energies of from ~ 0.6 to ~ 3.4 MeV. The measurements were conducted at standard ring geometry in an electrostatic accelerator. The γ spectra

from the ($n, n'\gamma$) reactions were detected with a Ge(Li) -spectrometer using a 2.5 cm^3 crystal. The absolute cross sections of excitation of all γ transitions were determined through a comparison of their yield with the intensity of the 846 keV line from ^{56}Fe . (K.S.W.)

46910 INVESTIGATION OF THE GAMMA DECAY OF EXCITED STATES IN ^{26}Al . Piiparinen, M. (Univ. of Helsinki). *Z. Phys.*; 252: No. 3, 206-18 (1972).

The gamma decay of the low-lying energy levels ($E_x < 4$ MeV) in ^{26}Al was studied through the $^{24}\text{Mg}(p, \gamma)^{26}\text{Al}$ reaction by using seven resonances in the proton energy range $E_p = 200$ to 1700 keV. Ge(Li) detectors and isotopically separated targets were used. Energies and branching ratios of several excited states were measured. The Q value of the reaction was determined to be 2271.3 ± 0.8 keV. Lifetimes, investigated by the Doppler-shift attenuation method, and γ -ray angular distributions were determined using the $E_p = 823, 1201, 1490$, and 1660 keV resonances of the $^{24}\text{Mg}(p, \gamma)^{26}\text{Al}$ reaction. (4 figures, 5 tables) (auth)

46911 BETA DECAY ASYMMETRY AND NUCLEAR MAGNETIC RESONANCE OF ^{28}Al , ^{38}Cl , AND ^{116}In PRODUCED BY CAPTURE OF POLARIZED NEUTRONS. Lades, Heinrich (Technische Univ., Munich). *Z. Phys.*; 252: No. 3, 242-8 (1972). (In German).

Polarized ^{28}Al , ^{38}Cl , and ^{116}In nuclei were produced by capture of polarized thermal neutrons. The nuclear magnetic resonance was observed by measuring the asymmetry of the β decay. The nuclear magnetic moments were found to be $\mu(^{28}\text{Al}) = 2.789(1) \text{ nm}$ and $\mu(^{38}\text{Cl}) = 2.05(2) \text{ nm}$ (uncorrected). The β -decay asymmetry of ^{116}In was observed in superconducting In metal. (5 figures) (auth)

46912 K^- -NUCLEUS SCATTERING LENGTHS. Seki, Ryo-ichi (California State Univ., Northridge). *Phys. Rev. Lett.*; 29: No. 4, 240-3 (24 Jul 1972).

K^- -nucleus scattering lengths are extracted from all K^- -mesonic atom data currently available. The scattering lengths obtained are shown to be model independent as long as the K^- -nucleus optical potential is local. (auth)

46913 ($^{16}\text{O}, ^{12}\text{C}$) STUDIES IN THE 2s-1d SHELL. Maher, J. V.; Erb, K. A.; Wedberg, G. H.; Ricci, J. L.; Miller, R. W. (Univ. of Pittsburgh). *Phys. Rev. Lett.*; 29: No. 5, 291-5 (31 Jul 1972).

Angular distributions were measured for the ($^{16}\text{O}, ^{12}\text{C}$) reactions with $^{24,26}\text{Mg}$, ^{27}Al , and ^{28}Si at 42 MeV. States in the " α -particle" nuclei ^{28}Si and ^{32}S were populated strongly for $0 \leq E_x \leq 11$ MeV, while cross sections for transitions to states in ^{30}Si and ^{31}P were small. The ^{28}Si and ^{32}S ground-state angular distributions are sharply oscillatory in nature. The 4.43-MeV state of ^{12}C is strongly excited in the reaction $^{24}\text{Mg}(^{16}\text{O}, ^{12}\text{C})^{28}\text{Si}$, but not in the others. (auth)

46914 NEW RESULTS ON THE FINE STRUCTURE OF ANALOG RESONANCES. Bilpuch, E. G. (Duke Univ., Durham, N. C. Triangle Universities Nuclear Lab., Durham, N. C.). pp 318-49 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

High-resolution experiments on proton elastic scattering by $A = 20$ to 66 nuclei have been performed at proton energies of 1 to 3.3 MeV. The fine structure of isobaric analog resonances observed in these experiments is discussed. (J.W.R.)

46915 NUCLEAR STRUCTURE OF ^{23}Na : CHARGED PARTICLE REACTION STUDIES. Powers, John Richard. Philadelphia; Univ. of Pennsylvania (1971). 265p. University Microfilms Order No. 72-17,412.

Thesis.

The nuclear structure of ^{23}Na has been studied using single-proton, two-nucleon, three-nucleon, and four-nucleon stripping in the $^{22}\text{Ne}(^3\text{He}, d)^{23}\text{Na}$ ($E_{^3\text{He}} = 15$ MeV), $^{21}\text{Ne}(^4\text{He}, p)^{23}\text{Na}$ ($E_{^4\text{He}} = 18$ MeV), $^{20}\text{Ne}(^7\text{Li}, \alpha)^{23}\text{Na}$ ($E_{^7\text{Li}} = 22$ MeV), and $^{19}\text{F}(^8\text{Li}, d)^{23}\text{Na}$ ($E_{^8\text{Li}} = 16$ MeV) reactions and using proton pickup in the $^{24}\text{Mg}(t, \alpha)^{23}\text{Na}$ ($E_t = 15$ MeV) reaction. The $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ ($E_d = 11.57, 11.82$, and 12.07 MeV) and $^{12}\text{C}(^{12}\text{C}, p)^{23}\text{Na}$ ($E_{^{12}\text{C}} = 30$ MeV) reactions were also studied. Angular-momentum transfer values were obtained from the systematics of the ($^3\text{He}, d$), ($^3\text{He}, p$), (t, α), ($^6\text{Li}, d$), and ($^7\text{Li}, \alpha$) angular distributions. The cross sections of states of known spin obtained from the (d, α) reaction were shown to vary in accord with the $(2J + 1)$ rule. Restrictions on the spin of unknown levels were then determined using this rule. By combining results several new spin-parity assignments were made. Transition strengths were obtained by a comparison of the experimental angular distributions with DWBA predictions for the ($^3\text{He}, d$),

(t, α), and ($^6\text{Li}, d$) reactions and with PWBA predictions for the ($^7\text{Li}, \alpha$) reaction. The pickup and stripping strengths obtained from these studies enabled a detailed analysis of the configuration of many states as predicted by the Nilsson model and shell model. Significant revision of previous understanding of the negative-parity state was proposed. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, $39 \leq A \leq 58$

Refer also to abstracts: 46848, 46873, 46875, 46883, 46886, 46889, 46890, 46894, 46901, 46909, 46914, 46932, 46934, 46936, 46940, and 47069.

46916 (AE-452) FAST NEUTRON ELASTIC SCATTERING EXPERIMENTS. Salama, M. (Aktiebolaget Atomenergi, Studsvik (Sweden)). 1972. 98p. Dep. NTIS (U. S. Sales Only).

Thesis. Submitted to Cairo Univ. (Egypt).

In the present study measurements were carried out on fast neutron elastic scattering angular distributions of the elements Cr, Fe and Ni in the energy interval 1.77 to 2.76 MeV and of As, Nb, Mo, Cd, Sb, Hf, Au, Pb, (radiogenic lead) and Pb at 8.05 MeV, using the time-of-flight technique. The fast neutrons were produced with the $T(p,n)^3\text{He}$ and the $D(d,n)^3\text{He}$ reactions using gas targets. The neutrons were scattered by hollow right cylindrical samples of the natural elements. The scattered neutrons were detected with a fast plastic scintillator discriminated against gamma rays to reduce background. The elastic scattering effect was separated from the inelastic scattering processes by use of a time-of-flight spectrometer. The spectrometer time resolution was about 3 ns. The angular distribution measurements were performed in the angular interval ranging from 20° to 160° with 3 meters flight path. The experimental data were collected in steps of 10° while in the extreme forward directions the data were taken every 5° . The experimental data were normalized and were corrected for background. Various corrections due to neutron attenuation in the sample, neutron multiple scattering and the finite geometry of the source-sample-detector system were applied to the measured distributions in order to derive accurate absolute cross sections. The corrected measured angular distributions were compared with the distributions calculated using a nuclear optical model assuming a central local potential consisting of Saxon-Woods and derivative Saxon-Woods and Thomas form factors describing the real, imaginary and spin orbit potentials. The numerical calculations were made using the ABACUS II computer code. From the nuclear structure point of view, the elastic scattering data were examined to obtain information upon the distribution of neutrons and protons inside nuclei. The extension of the matter and charge distributions have been calculated for Sb and Au from neutron and electron root-mean-square radii confirming other observations that the mass distribution extends beyond that of the nuclear charge distribution. It has also been confirmed that the difference between the root-mean-square matter and charge radii increases somewhat with the neutron excess in nuclei. (auth)

46917 (BNL-tr-477) γ -EMISSION IN RADIATIVE CAPTURE OF INTERMEDIATE NEUTRONS. Gamalii, A. F.; Zemtsev, B. V.; Ivanov, V. B.; Netserov, B. V.; Kham'yanov, L. P. Translated by S. J. Amoretty (Brookhaven National Lab., Upton, N. Y.), from Yadern. Fiz.; 15: No. 1, 3-9(1972). 12p. Dep. NTIS.

The spectra of the capture gamma rays from radiative capture of thermal neutrons and of neutrons with energies 2 and 25 keV by ^{95}Mo , ^{62}Ni , ^{48}Ti nuclei were studied using a γ spectrometer with a Ge(Li) detector. The neutrons with average energies of 2 and 25 keV were selected from the solid reactor spectrum with the help of scandium and ferroaluminum filters. (3 figures, 1 table) (auth)

46918 (JEN-230-DMe/I-25) TABLE OF CROSS SECTION (n, p), (n, α), AND ($n, 2n$) REACTIONS IN STEEL COMPONENTS AND OTHER NUCLEAR MATERIALS. Lopez Jimenez, Jose (Junta de Energia Nuclear, Madrid (Spain)). Division de Metalurgia. 1972. 35p. (In Spanish). Dep. NTIS (U. S. Sales Only).

Reactions (n, p) and (n, α) produce in the materials large amounts of hydrogen and helium atoms. The presence, especially of helium, changes the physical properties of materials and particularly reduces the ductility of irradiated stainless steel cladding above 500°C . Cross sections of all isotopes which constitute the S. S. and other clad materials, have been compiled. Experimental available data were obtained from BNL (1956, 64 and 68), and the rest, from J. C. Roy and J. J. Hawton calculations in a fission neutron spectrum (1960). By the Fortran IV program COFRE, the

fission-spectrum-averaged cross section of experimental data and the average values in lethargy constant intervals were obtained. The estimated cross section of ROY has been transformed to threshold "equivalent" cross sections using the effective energy E_{eff} . These data will be the in-put data of PROTALFA program, which determine the amount of H and He gas produced by the interactions of thermal and fast neutron flux in S. S. and other clad alloys. (auth)

46919 FURTHER RESULTS ON THE PROTON RADIOACTIVITY OF ^{53}Co . Cerny, Joseph; Gough, R. A.; Sextro, R. G.; Esterl, John E. (Univ. of California, Berkeley). Nucl. Phys.; A188: No. 3, 666-72(1972).

Additional studies on the proton radioactivity of the 247 ± 12 msec isomer ^{53m}Co lead to an improved estimate of ≈ 17 sec for its partial half-life for emission of 1.59 ± 0.03 MeV protons to the ^{52}Fe ground state. An upper limit of 1/250 for the ratio of direct proton decay to the $^{52}\text{Fe}^*(2^+, 0.84 \text{ MeV})$ state relative to decay to the $^{52}\text{Fe}(\text{g.s.})$ can be set. (auth)

46920 ENERGY OF LEVELS OF ^{45}Sc UP TO 3 MeV FROM THE $^{44}\text{Ca}(\text{p}, \gamma)^{45}\text{Sc}$ REACTION. Ekstroem, P.; Erlandsson, B.; Marcinkowski, A.; Tillman, J. (Univ. of Lund, Sweden). Z. Phys.; 252: No. 3, 189-205(1972).

The gamma-ray decay of the 1658-keV resonance, which is a member of the isobaric analogs of the 1904-keV $\frac{3}{2}^-$ level in ^{44}Ca , was studied with a Ge(Li) detector. A number of gamma-ray transitions are described and associated with the decay of 15 levels between 1.50 and 3.05 MeV. These observations made it possible to deduce spin and parity assignments for some of the levels. By comparing excited levels of ^{43}Sc and ^{45}Sc it was possible to assign certain levels as members of one negative-parity band and two positive-parity bands with $K^\pi = \frac{1}{2}^+$ and $\frac{3}{2}^+$. (9 figures, 2 tables) (auth)

46921 THEORETICAL COEFFICIENTS OF INTERNAL CONVERSION FOR ^{51}V . Trusov, V. F. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2338-40(Nov 1971). (In Russian).

Internal conversion coefficients were found for ^{51}V in K -, L -, M -, and M_1 -subshells of the first two multipole orders or electrical and magnetic transitions in the energy range 0.05 to 1 m.e.c.^2 . Screening was taken into account using the Gashpar analytical potential in a variant of non-linear deformation of effective charge. The transition $M1-E2$ with an energy of 320 keV was examined, for which experimental data exist. (tr-auth)

46922 ELASTIC AND INELASTIC SCATTERING OF 6-MeV PROTONS BY THE EVEN ISOTOPES OF CHROMIUM. Antropov, A. E.; Zarubin, P. P.; Ioannu, P. D.; Orlov, B. N.; Romanov, V. S. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2355-8 (Nov 1971). (In Russian).

The energy and angular distribution of elastic and inelastic scattering of 6 MeV protons by the isotopes $^{50,52,54}\text{Cr}$ were measured. Peculiarities were discovered which were successfully explained by the different contribution to the scattering of indirect processes depending on the magnitude of the threshold of the (p, n) reaction. (tr-auth)

46923 EFFECT OF DIFFUSION OF THE BOUNDARY OF A NUCLEUS AND THE COULOMB INTERACTION ON THE DIFFRACTION DISINTEGRATION OF DEUTERONS. Berezhnoi, Yu. A.; Evlanov, M. V. (Physical-Technical Inst., Kharkov. Inst. of Nuclear Research, Kiev). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2406-9(Nov 1971). (In Russian).

A mathematical development led to the calculation of the differential cross section of elastic scattering of 100 MeV deuterons by ^{58}Ni nuclei under three different assumptions: the possession by the nucleus of a sharp boundary but without allowing for the Coulomb interaction of the deuteron with the nucleus, the existence of a diffuse boundary for the nucleus but without allowing for deuteron-nucleus interaction, and the existence of a diffuse nuclear boundary, and making allowance for the Coulomb interaction. The third case significantly changed the behavior of the differential cross sections of interaction processes between deuterons and nuclei, and this fact may prove to be important in the comparison of theoretical results with experimental data. (K.S.W.)

46924 ELASTIC SCATTERING OF POLARIZED PROTONS BY ^{28}Si AND ^{40}Ca . Klucharev, A. P.; Slabospitskii, R. P.; Andreev, G. G.; (and others) (Physical-Technical Inst., Kharkov, USSR). Ann. Phys. (N. Y.); 71: No. 2, 438-42(Jun 1972).

Proton polarization was measured for elastic scattering of polarized protons in the reaction $^{28}\text{Si}(\text{p}, \text{p})^{28}\text{Si}$ in the energy region 1.5 to 3.0 MeV and in $^{40}\text{Ca}(\text{p}, \text{p})^{40}\text{Ca}$ in the energy region 2.3 to 2.8 MeV at angles of 60° and 90° c.m. As a result of the investigation, the total angular momentum for excited states of ^{29}P , $J^\pi = \frac{3}{2}^-, \frac{1}{2}^+, \frac{1}{2}^-$, was determined. For states of ^{41}Sc with

energies of 3.729 and 3.769 MeV, $J^\pi = 1/2^-$. But instead of the state at 3.467 MeV, two states with energy excitation 3.460 and 3.475 MeV with $J^\pi = 1/2^-$ were discovered. All four investigated levels of ^{41}Sc obviously are terms of multiplet of one-particle excited state. Phase-shift analyses made it possible to obtain widths for the levels from the energy dependence of the polarization for the levels investigated. (3 figures) (auth)

46925 ^{53}mCo : A PROTON-UNSTABLE ISOMER. Jackson, K. P. (Oxford Univ.); Cardinal, C. U.; Evans, H. C.; Jelley, N. A.; Cerny, J. Phys. Lett.; 33B: No. 4, 281-3 (26 Oct 1970).

A 1.53 ± 0.04 MeV proton activity with a 245 ± 20 ms half-life has been observed in the reaction of ^{16}O on ^{40}Ca . The most plausible origin of this activity is the proton radioactivity of ^{53}mCo , although the decay of this isomer by beta-delayed proton emission remains a possibility. (auth)

46926 CONFIRMED PROTON RADIOACTIVITY OF ^{53}mCo . Cerny, J.; Esterl, J. E.; Gough, R. A.; Sextro, R. G. (Univ. of California, Berkeley). Phys. Lett.; 33B: No. 4, 284-6 (26 Oct 1970).

Proton-induced reactions on ^{54}Fe produce a proton activity [1.57 ± 0.03 MeV; 242 ± 15 ms] with a threshold of 26.3 ± 0.4 MeV which can only arise from ^{53}mCo . Failure to detect positron-proton coincidences in the decay of this isomer establishes its direct proton radioactivity. (auth)

46927 $^{58}\text{Ni}(\alpha, \alpha')^{58}\text{Ni}$ AND THE NATURE OF HIGH-LYING STATES IN ^{58}Ni . Bruge, G.; Chaumeaux, A.; DeVries, R.; Morrison, G. C. (CEN de Saclay, Gif-sur-Yvette, France). Phys. Rev. Lett.; 29: No. 5, 295-8 (31 Jul 1972).

Levels in ^{58}Ni up to 10 MeV excitation energy were studied through the (α, α') reaction at 30 MeV; surprising overall agreement between the observed spectra and those from α -transfer reactions was found. (auth)

46928 MISSING LOW-LYING STATES IN ^{39}K . Durell, J. L. Max-Planck Institut fuer Kernphysik, Heidelberg; Metag, V.; Repnow, R.; (and others). Phys. Rev. Lett.; 28: No. 26, 1723-5 (26 Jun 1972).

The previously known levels of ^{39}K at 3.94 and 4.52 MeV have been shown to be doublets. Two of these levels are candidates for the missing low-lying negative-parity states predicted by shell-model and phonon-hole calculations. (auth)

46929 ANGULAR DISTRIBUTION OF NEUTRONS FROM (p,n) REACTIONS ON ^{18}O , ^{37}Cl , ^{41}K , ^{53}Cr , AND ^{59}Co NUCLEI. Dryapachenko, I. P.; Nemets, O. F.; Pilipchenko, V. A. (Inst. of Nuclear Research, Kiev). Ukr. Fiz. Zh. (Russ. Ed.); 16: No. 12, 1947-52 (Dec 1971). (In Russian).

Angular distributions of neutrons from (p,n) reactions with ^{59}Co , ^{53}Cr , ^{18}O nuclei and total cross sections for the reactions $^{59}\text{Co}(p,n)^{59}\text{Ni}$ and $^{53}\text{Cr}(p,n)^{53}\text{Mn}$ at energies from 2.8 MeV to 4 MeV were measured by means of the long counter and the single-crystal scintillation spectrometer. Angular distributions of neutrons from (p,n) reactions with ^{37}Cl , ^{59}Co , and ^{41}K to ground states of the final nuclei at proton energies of 3.7 MeV and 3.9 MeV were also measured with the single-crystal scintillation spectrometer. The influence of the interference effects on the angular distributions from the reactions $^{37}\text{Cl}(p,n)^{37}\text{Ar}$ and $^{18}\text{O}(p,n)^{18}\text{F}$ was observed. For the remaining nuclei the angular distributions slightly differ from isotropic ones. The results of the measurements are compared with the data of other authors. (7 figures) (auth)

46930 NUCLEAR STRUCTURE AND REACTION MECHANISM STUDIES USING THE $(^3\text{He}, t)$ REACTION. Fadner, Willard Lee. Boulder, Colo.; Univ. of Colorado (1971). 214p. University Microfilms Order No. 72-17,256.

Thesis.

Transitions to the isobaric analog ground states of ^{46}Ti , ^{50}Cr , ^{54}Fe , ^{62}Ni , ^{64}Ni and ^{90}Zr , and to several excited states of the particle-hole mirror nuclei ^{46}V and ^{50}Mn were studied with the $(^3\text{He}, t)$ reaction over the energy range of 21.4 to 37.5 MeV. For the transitions to the isobaric analog ground states, unexpected and complex variations in the shapes of the experimental angular distributions were observed. The variations in the shape as a function of energy had distinct features for each of the nuclei examined. These effects were poorly described by DWBA calculations based on a simplified shell-model theory (microscopic model). Calculations based on a generalized optical model theory (macroscopic model), using different interaction terms for each nucleus, gave a somewhat better description. Interaction strengths, normalized to fit the data, required a significant decrease with increasing incident energy for both theoretical models. This effect, documented herein, has been a major factor in variations in interaction strengths previously noted between independent studies. It was observed, however, that the interaction strengths were approximately constant as a function of target nucleus at the incident

energy of 37.5 MeV, whereas this constancy was not seen at lower energies. The poor description, both in shape and in strength, of the energy dependence of the data is evidence of basic inadequacies in the theories as they are presently used. On the basis of a simple shell-model, it is expected that the energy levels of excited states of the nucleus ^{46}V should be essentially the same as those for ^{50}Mn . This is because the two nuclei can be considered to be particle-hole conjugates. Particles and holes in a particular shell should be equivalent, according to the shell-model. To investigate this relationship, angular distributions were obtained for 16 levels in ^{46}V , ranging up to 4.01 MeV excitation, and 8 levels in ^{50}Mn , ranging up to 2.00 MeV excitation. The data were compared to both macroscopic and microscopic distorted-wave calculations, using a tensor force when appropriate. Several J, π assignments were made. The resulting level schemes for the two nuclei were compared with shell-model calculations. Significant discrepancies with the theory, and between the levels for the two nuclei, were noted. Presumably the experimental results now made available could aid in further theoretical refinements. The excited states of ^{50}Mn were also studied as a function of energy. Macroscopic asymmetry potential strengths and microscopic interaction strengths decreased significantly with energy, similar to the effect observed for the isobaric analog ground states. This points out the generality of the energy dependence of the strengths using the present theoretical methods. Values of the neutron excess deformation parameters, β_2 and β_4 were only mildly energy dependent for the energies examined in this study. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, $59 \leq A \leq 89$

Refer also to abstracts: 46853, 46873, 46875, 46890, 46914, 46916, 46929, 46930, 46985, and 47066.

46931 NEUTRON DEFICIENT EVEN-EVEN ISOTOPES OF Kr, Se, Ge, AND PSEUDO SU_3 COUPLING. Nolte, E.; Kutschera, W.; Shida, Y.; Morinaga, H. (Technical Univ., Munich). Phys. Lett.; 33B: No. 4, 294-6 (26 Oct 1970).

Gamma-ray spectra from the reactions $^{58,60,61,62}\text{Ni}(^{16}\text{O}, \text{xn yp } \gamma\gamma)$ and $^{58}\text{Ni}(^{12}\text{C}, \text{xn yp } \gamma\gamma)$ were studied and low-lying levels of neutron deficient even-even isotopes of Kr, Se and Ge were obtained. The results are compared with the recently proposed pseudo-LS and pseudo- SU_3 coupling schemes. (auth)

46932 LEVEL DENSITIES FROM SPECTRA OF DIFFERENT REACTIONS POPULATING THE SAME NUCLEUS. Lu, C. C.; Vaz, L. C.; Huizenga, J. R. (Univ. of Rochester, N. Y.). Nucl. Phys.; A190: No. 2, 229-60 (1972).

Level densities of ^{56}Fe , ^{59}Co , ^{60}Ni , ^{62}Ni , ^{63}Cu and ^{65}Cu are determined from analyses of several charged-particle spectra populating the same residual nucleus. Experimental spectra are reported for the following reactions: $^{56}\text{Fe}(\alpha, \alpha')^{56}\text{Fe}$, $^{59}\text{Co}(p, \alpha')^{56}\text{Fe}$, $^{59}\text{Co}(p, p')^{59}\text{Co}$, $^{58}\text{Co}(\alpha, \alpha')^{58}\text{Co}$, $^{56}\text{Fe}(\alpha, p)^{59}\text{Co}$, $^{62}\text{Ni}(p, \alpha)^{59}\text{Co}$, $^{60}\text{Ni}(\alpha, \alpha')^{60}\text{Ni}$, $^{63}\text{Cu}(p, \alpha)^{60}\text{Ni}$, $^{62}\text{Ni}(p, p')^{62}\text{Ni}$, $^{62}\text{Ni}(\alpha, \alpha')^{62}\text{Ni}$, $^{59}\text{Co}(\alpha, p)^{62}\text{Ni}$, $^{65}\text{Cu}(p, \alpha)^{62}\text{Ni}$, $^{63}\text{Cu}(p, p')^{63}\text{Cu}$, $^{63}\text{Cu}(\alpha, \alpha')^{63}\text{Cu}$, $^{60}\text{Ni}(\alpha, p)^{63}\text{Cu}$, $^{65}\text{Cu}(p, p')^{65}\text{Cu}$, $^{65}\text{Cu}(\alpha, \alpha')^{65}\text{Cu}$ and $^{62}\text{Ni}(\alpha, p)^{65}\text{Cu}$. Each of these experimental spectra is analyzed with an exact theory giving the angular and energy-dependent differential cross sections for compound-nucleus reactions including explicitly the angular momentum. From these analyses the Fermi-gas level-density parameters were determined: ^{56}Fe , $a = 5.7$ MeV $^{-1}$, $\Delta = 0.7$ MeV; ^{59}Co , $a = 6.2$ MeV $^{-1}$, $\Delta = -0.8$ MeV; ^{60}Ni , $a = 6.4$ MeV $^{-1}$, $\Delta = 1.3$ MeV; ^{62}Ni , $a = 6.4$ MeV $^{-1}$, $\Delta = 0.5$ MeV; ^{63}Cu , $a = 6.8$ MeV $^{-1}$, $\Delta = -0.5$ MeV; and ^{65}Cu , $a = 6.6$ MeV $^{-1}$ and $\Delta = -0.5$ MeV. All of the spectra were also analyzed with the conventional slope technique which utilizes the Weisskopf expression for the differential cross sections. The results from such an analysis depend on the form of the pre-exponential term in the level-density formula and may give sizeable errors in the Fermi-gas level-density parameter a . A number of theoretical spectra are computed with the exact theory which explicitly includes angular momentum and these spectra are analyzed with the conventional methods to illustrate the errors introduced when angular momentum is not treated properly for such reactions. An estimate of precompound proton emission is made for several (p,p') reactions. (auth)

46933 INTENSITY RATIO OF 2158 keV AND 2505 keV GAMMA-RAYS IN THE DECAY OF ^{60}Co . Fujishiro, Masatoshi; Furuta, Junichiro (Radiation Center of Osaka Prefecture, Sakai, Japan). J. Nucl. Sci. Technol. (Tokyo); 9: No. 6, 348-52 (Jun 1972).

The intensity ratio of the 2,158 and 2,505 keV γ -rays in the decay of ^{60}Co was measured by applying the photonuclear reactions $^9\text{Be}(\gamma, n)$ and $^2\text{D}(\gamma, n)$. The result obtained was an intensity ratio of $\{I(2,158)/I(2,505)\} \sim 237$, which exceeded all the values reported in available literature on previous measurements. (auth)

46934 (α, p) REACTION ON EVEN NICKEL ISOTOPES. Bucurescu, D.; Ivascu, M.; Semencescu, G.; Titirici, M. (Inst. of Atomic Physics, Bucharest). Nucl. Phys., A189: No. 3, 577-91(1972).

The (α, p) reaction on $^{58,60,62,64}\text{Ni}$ has been studied at an incident energy of 19.3 MeV. Zero-range DWBA calculations based on a triton-transfer mechanism describe well the whole set of experimental data. The strong J-dependence of the $L = 1$ transitions is described by using the spin-orbit coupling in the proton channel. The systematic behavior of the transitions for each L and J and the good agreement with the DWBA curves enable one to make spin assignments for levels in ^{67}Cu . The relative intensities of the ground state transitions agree with calculated values. The relative strengths for transitions to excited states are compared with those determined from proton-transfer reactions. (auth)

46935 DEPENDENCE OF THE DEORIENTATION OF NUCLEI OF RECOIL ATOMS (^{103}Rh AND ^{74}Ge) WHILE FREELY FLYING IN A VACUUM ON TIME. Ward, D.; Geiger, J. S.; Graham, R. L. (Chalk River Nuclear Labs., Ont.). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2325-30(Nov 1971). (In Russian).

Nuclear spin alignment may be strongly disturbed in the absence of a decelerating medium. This type of deorientation was investigated using a plunger type of apparatus (Alexander and Bell, 1970), in which a beam of heavy ions impinged on a thin target of ^{74}Ge . Inelastic scattering of particles of the beam was recorded, as Coulomb excitation of the target atoms occurred. Similar measurements were made of the Coulomb excitation of the 295 keV $3/2^-$ level in ^{103}Rh . (K.S.W.)

46936 EXCITATION CROSS SECTION OF THE FIRST 2+ LEVELS OF Ni AND Zn NUCLEI DURING INELASTIC SCATTERING OF NEUTRONS IN THE NEAR-THRESHOLD ENERGY REGION. Konobeevskii, E. S.; Musaelyan, R. M.; Popov, V. I.; Surkova, I. V. (Lebedev Inst. of Physics, Moscow). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2345-8(Nov 1971). (In Russian).

The energy dependence of the (n, n γ) reaction cross section with excitation of the first 2+ levels of $^{58,60}\text{Ni}$ and $^{64,66,68}\text{Zn}$ was investigated. The experimental data for Ni is satisfactorily described by the optical model without allowing for the correlation of the resonance widths. The cross section for Zn exceeded the calculated value. (tr-auth)

46937 EXCITATION OF INPUT STATES DURING THERMAL NEUTRON CAPTURE BY ^{68}Zn , ^{70}Ge , AND ^{80}Se . Murzin, A. V.; Kolomiets, V. M. (Inst. of Nuclear Research, Kiev). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2349-54(Nov 1971). (In Russian).

Possible types of input states were examined which were excited during capture of thermal neutrons by ^{68}Zn , ^{70}Ge , and ^{80}Se nuclei. Special features of the γ spectra of capture are explained qualitatively for the isotopes ^{68}Zn , ^{71}Ge , and ^{81}Se , and their low excited levels are depicted schematically. (tr-auth)

46938 DECAY OF ^{76}Br . Dzhelepov, B. S.; Dmitriev, A. G.; Zhukovskii, N. N. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2249-54(Nov 1971). (In Russian).

Two β^+ components were derived from the ^{76}Br spectrum: 3,980 and 3,440 keV. The ratio of their intensities was 0.22. The ratio of the intensities of the transitions γ 559 and β^+ 3,980 was 7.6. Six new lines were detected in the ^{76}Br γ spectrum; their relative intensities were measured. The energy values and relative intensities of the ^{76}As γ lines were also determined. The decay scheme of ^{76}Br was refined. High values for log ft of the β^+ -transitions on the basis of the excited states (ground, 0^+ , and first-excited, 2^+) of ^{76}Se are evidence of $J^\pi = 1^-$ for the ground state of ^{76}Br . (tr-auth)

46939 IZOBARNYE YADRA S MASSOVYM CHISLOM $A = 72$. (Isobaric Nuclei with Mass Number $A = 72$). Vitman, V. D.; Dzhelepov, B. S.; Medvedev, A. I. Svoistva Atomnykh Yader, Vypusk 13. Leningrad: Izdatel'stvo Nauka Leningradskoe Ot-delenie (1972). 120p.

This monograph summarizes the voluminous material on the nuclei ^{72}Zn , ^{72}Ga , ^{72}Ge , ^{72}As , ^{72}Se , ^{72}Br , and ^{72}Kr which has appeared in print up to September 1, 1971. The information compiled includes energies and intensities of radioactive emissions from these isotopes, detailed decay schemes, quantum characteristics

of excited levels, and similar properties. The reliability of the data is evaluated, and additional investigation is recommended for certain work. (tr-auth)

46940 NUCLEAR γ RAYS ASSOCIATED WITH STOPPED KAONS. Barnes, P. D. (Carnegie-Mellon Univ., Pittsburgh); Eisenstein, R. A.; Lam, W. C.; (and others). Phys. Rev. Lett.; 29: No. 4, 230-3(24 Jul 1972).

A large number of nuclear γ rays associated with negative kaons stopped in Ni, Cu, Si, and Al were identified. Prominent among these for both the Cu and the Ni targets are transitions that take place in even-even nuclei lighter than the target by up to about twelve nucleons. In both cases, the total intensity of all of the identified nuclear γ rays amounts to about twice that of the strongest x ray. (auth)

46941 SEARCH FOR THE REACTION $\mu^- + \text{Cu} \rightarrow e^+ + \text{Co}$. Bryman, Douglas Andrew. Blacksburg, Va.; Virginia Polytechnic Inst. and State Univ. (1972). 62p. University Microfilms Order No. 72-18,005.

Thesis.

A search was performed for the reaction $\mu^- + \text{Cu} \rightarrow e^+ + \text{Co}$, which is allowed by the lepton conservation scheme that assigns the same lepton number to the μ^- , e^+ , ν_μ , and $\bar{\nu}_e$. Momentum analysis of particles emitted following muon capture was performed using a magnetic spectrometer with magnetostriuctive readout wire spark chambers. The time of flight through the spectrometer and energy loss in a scintillation counter were measured for each event. One event was observed, at momentum 89.9 ± 3.5 MeV/c, which is consistent with the above process. Background from radiative muon capture was expected to produce ~ 0.01 events that could stimulate the search for reaction. However, there exists uncertainty in the relative yield and cutoff for high energy photons from radiative muon capture. Although it is not certain whether or not the reaction $\mu^- + \text{Cu} \rightarrow e^+ + \text{Co}$ has been detected, an upper limit was set on the branching ratio relative to ordinary muon capture of $R \leq 2.6 \times 10^{-8}$ (90% confidence level). According to a calculation by Kisslinger, this can be related to an upper limit on the coupling constant of this type of interaction relative to the vector coupling constant of (G_A/G_V) $\leq 0.10 \pm 0.03$. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, $90 \leq A \leq 149$

Refer also to abstracts: 46873, 46875, 46883, 46889, 46890, 46909, 46911, 46916, 46917, 46930, 46935, 47066, 47072, 47073, 47075, and 47209.

46942 (JINR-P6-6512) STUDY OF THE DECAY ^{148}Tb ($T_{1/2} = 66$ min) AND ^{150}Tb ($T_{1/2} = 3.48 \pm 0.16$ h). Vylov, Ts.; Gromov, K. Ya.; Gromova, I. I.; Iskhakov, G. I.; Kuznetsov, V. V.; Kuznetsova, M. Ya.; Potempa, A. V.; Fominykh, M. I. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Nuclear Problems). 1972. 28p. (In Russian). Dep. NTIS (U. S. Sales Only).

Gamma-ray, conversion electron, and γ - γ coincidence spectra in the decay of ^{148}Tb and ^{150}Tb were studied. The half-life was measured for ^{150}Tb ($T_{1/2} = 3.48 \pm 0.16$ h). The decay schemes $^{148}\text{Tb} \rightarrow ^{148}\text{Gd}$ and $^{150}\text{Tb} \rightarrow ^{150}\text{Gd}$ are proposed. The comparison is made for the excited states of neighboring nuclei ^{148}Gd , ^{148}Sm , ^{144}Nd and ^{150}Gd , ^{148}Sm , ^{146}Nd . (auth)

46943 (N-72-18705) PARTICLE-HOLE STATES IN ^{120}Sn . Leonard, Regis F. (National Aeronautics and Space Administration, Cleveland, Ohio, Lewis Research Center). 31 Jan 1972. 9p. (CONF-720119-1; NASA-TM-X-68004; E-6772). NTIS. From American Physical Society meeting; San Francisco, CA. (31 Jan 1972).

The 45 MeV deuteron beam from the Berkeley 88-inch cyclotron was used to study the particle-hole states in ^{120}Sn . The present work resolved a number of individual levels and compared their strength and excitation energy with that observed in the $\text{Sn}-102(\text{d}, \text{He}-3)$ reaction, which was also remeasured. A schematic diagram of the scattering system is given along with a description. (auth) (STAR)

46944 DECAY OF ISOMERS $^{126,128,130}\text{Sb}$ AND QUASIPARTICLE LEVELS $^{126,128,130}\text{Te}$ WITH LARGE SPIN. Kiselev, V. G.; Levkovskii, V. N.; Artem'ev, O. I.; Peker, L. K. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2300-5(Nov 1971). (In Russian).

The gamma spectra of $^{126,128,130}\text{Sb}$ were measured, and the resulting data were utilized in the construction of a decay scheme.

Analysis of the data showed two-particle levels with $I^\pi = 7^-$ and four particle levels with $I^\pi = 9^-$ for the isotopes $^{126,128,130}\text{Te}$. (tr-auth)

46945 ^{135}Ba LEVELS. Begzhanov, R. B.; Glad'shev, D. A.; Kobilov, O. Sh.; Radzhapov, P. S.; Sabirov, Kh. S. (Inst. of Nuclear Physics, Tashkent). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2312-15 (Nov 1971). (In Russian).

The angular correlations (γ 373), (γ 481) and (γ 633), (γ 218) were measured in the ^{135}Ba nucleus. Spin and parity of $\frac{1}{2}^+$, $\frac{5}{2}^+$, and $\frac{3}{2}^+$ are uniquely described by the levels 218, 481, and 852 keV, respectively. The lifetime of the 481 keV level was evaluated by delayed $\gamma\gamma$ coincidence: $T_{1/2} \leq 10$ psec. (tr-auth)

46946 NEUTRON SPECTROMETRY OF THE Sb NUCLEUS. Mateciuc, V. A. (Inst. of Atomic Physics, Bucharest). *Stud. Cercet. Fiz.*; 24: No. 2, 165-97 (1972). (In Rumanian).

A pulsed neutron source and a time-of-flight spectrometer were used to study neutron reactions with Sb. Measured quantities included the Sb neutron cross section at 0.05 to 150 eV, radiative capture widths, reduced neutron widths, average level spacings, the neutron strength function, Doppler effect and crystal binding effect on neutron resonances, and ^{121}Sb resonance parameters. The experimental facilities and equipment are described. (J.W.R.)

46947 DETERMINATION OF THE TOTAL COEFFICIENT OF INTERNAL GAMMA-RADIATION CONVERSION OF ^{139}La . Aristov, E. A.; Bazhenov, V. A. *Meas. Tech. (USSR) (Engl. Transl.)*; 14: No. 12, 1883-5 (Dec 1971).

Translated from *Izmer. Tekh.*; 14: No. 12, 61-2 (Dec 1971).

The decay of ^{139}Ce was studied using a proportional counter-NaI(Tl) scintillation detector coincidence setup. Measurements were made on γ -x-ray coincidences for six carefully prepared sources. The results were used to determine the total internal conversion coefficient for the 166-keV transition in ^{139}La . (J.W.R.)

46948 ^{132}Ba LEVELS EXCITED DURING THE DECAY OF ^{132}La AND $^{132*}\text{La}$. Amov, B.; Bylov, Ts.; Enikova, M.; (and others). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2266-71 (Nov 1971). (In Russian).

The isotope ^{132}La was obtained by irradiating a gadolinium target with 660 MeV protons in a synchrotron. Gamma spectra, electron conversion spectra, and $\gamma\gamma$ coincidence were studied. A decay scheme was constructed for excited levels of ^{132}Ba , and the intensities of some γ -transitions which arose during the decay of ^{132}La and $^{132*}\text{La}$ were measured. (K.S.W.)

46949 ISOMERIC TRANSITION $8^- \rightarrow 6^+$ IN THE ^{130}Ba NUCLEUS. Arsen'ev, D. A.; Dzholos, R. V. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2281-6 (Nov 1971). (In Russian).

A model was proposed for describing the properties of a transient nucleus, based on energy calculations of the deformation of a nucleus. Using this model the isomeric state 8^- in the ^{130}Ba nucleus was studied. (tr-auth)

46950 OPTICAL DETERMINATION OF THE QUADRUPOLE MOMENT OF ^{138}La . Fischer, W. (Univ., Marburg, Ger.); Huehnermann, H.; Mandrek, K.; Ihle, H. *Phys. Lett.*; 40B: No. 1, 87-8 (12 Jun 1972).

The rare lanthanum isotope ^{138}La was highly enriched with the aid of an electromagnetic isotope separator. The hyperfine structure of two La lines was investigated by means of optical interference spectroscopy, which allowed the determination of the nuclear electric quadrupole moment $Q(^{138}\text{La}) = 0.51 \pm 0.09$ b. (auth)

46951 EXCITATION OF THE ^{99}Tc NUCLEUS. Begzhanov, R. B.; Sabirov, Kh. S. (Inst. of Nuclear Physics, Tashkent). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2316-22 (Nov 1971). (In Russian).

The $\gamma\gamma$ -angular correlation of cascades 740-180 and 740-40-140 keV in ^{99}Tc was investigated. The lifetimes of the 922 and 180 keV states were measured ($T_{1/2} = 0.2 \times 10^{-8}$ sec and 3.36×10^{-9} sec, respectively). Conclusions were drawn regarding the quantum characteristics of several levels. (tr-auth)

46952 SUB-BARRIER (p,n) REACTION IN THE REGION OF SINGLE-PARTICLE RESONANCE. Sirotkin, V. K.; Chesnokova, V. D. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2359-63 (Nov 1971). (In Russian).

The energy dependence of a sub-barrier (p,n) reaction of Sn isotopes in the region of single-particle 3p-resonance was analyzed on the basis of the shell approach to the theory of nuclear reactions. A significant difference in the results of the shell approach and of the optical model was detected. The results of numerical calculations were compared with experimental data. (tr-auth)

46953 CONVERSION OF BREMSSTRAHLUNG PHOTO NEUTRON MEAN ENERGIES TO MONOCHROMATIC PHOTO NEUTRON MEAN ENERGIES. Thies, H. H.; Thomas, B. W.; Lee, F.; Kochi, R.; Crawford, D. M. (Univ. of Western Australia, Netherlands). *Nucl. Instrum. Methods*; 102: No. 2, 219-22 (1972).

Using a method described by Barrett and Thies one can measure readily the mean energies of neutrons produced in bremsstrahlung experiments. It is shown here how one can calculate from data on bremsstrahlung photo neutron mean energies the neutron mean energies that correspond to an idealized experiment using monochromatic gammas. Simple formulae are derived to estimate the accuracy of calculated mean energies. The practical limits of experiments on photo neutron mean energies are discussed. (auth)

46954 MULTIPOLE ORDER OF TRANSITIONS AND CHARACTERISTICS OF THE EXCITED STATE OF ^{147}Eu BY THE METHOD OF ANGULAR $\gamma\gamma$ CORRELATION. Vyvryshchuk, Ya.; Zhuk, V.; Krupa, E.; (and others). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2331-4 (Nov 1971). (In Russian).

Measurements of the angular correlation of γ quanta with energies of 396, 765, 893, and 1,130 keV were made, coincident with the most intense γ transition of 229 keV in ^{147}Eu . For the 1,359 keV level, $I^\pi = \frac{9}{2}^+$, and for the 1,122.3 keV level the most probable value of $I^\pi = \frac{5}{2}^+$. (tr-auth)

46955 HINDERED $\frac{1}{2}^+ \rightarrow \frac{3}{2}^-$ β -DECAY OF ^{143}Pr . Griffin, H. C. (Univ. of Michigan, Ann Arbor); Rengan, K. *Nucl. Phys.*; A190: No. 2, 293-6 (1972).

A low-intensity 742 keV γ -ray has been found in the decay of ^{143}Pr . This γ -ray is interpreted as evidence for a weak β -transition ($1.2 \times 10^{-8}\%$, $\log f_{\beta} = 13.2$, $\log f_{\beta} t = 11.6$) from ^{143}Pr ($J^\pi = \frac{1}{2}^+$) to the 742 keV first excited state of ^{143}Nd ($\frac{3}{2}^-$). (auth)

46956 $^{90}\text{Zr}(d,n)^{91}\text{Nb}$ AND $^{96}\text{Zr}(d,n)^{97}\text{Nb}$ REACTIONS. Horton, J. L. (Univ. of Texas, Austin); Hollas, C. L.; Riley, P. J.; (and others). *Nucl. Phys.*; A190: No. 2, 362-72 (1972).

The (d,n) reaction on ^{90}Zr and ^{96}Zr has been studied at 12 MeV deuteron bombarding energy using the neutron time-of-flight technique with an overall neutron time resolution of 1.9 ns. Angular distributions of neutron groups leading to states in ^{91}Nb and ^{97}Nb were measured in the angular range between 15° and 60° . The measured cross sections were analyzed in the framework of the distorted-wave theory of stripping reactions to deduce l-values and proton spectroscopic factors of states in the residual nuclei. The results are compared with the corresponding data available from ($^3\text{He},d$) studies. The fractionation of the single-particle proton states and their centroid energies are determined. (auth)

46957 GAMMA-GAMMA ANGULAR CORRELATION OF THE 171- TO 427-keV CASCADE AND THE g-FACTOR OF THE 462-keV, $5/2^+$ LEVEL OF ^{126}Te . Nigam, A. K.; Bhattacharyya, R. (Saha Inst. of Nuclear Physics, Calcutta). *Indian J. Phys.*; 45: No. 5, 214-24 (May 1971).

The gyromagnetic ratio of the 462 keV $\frac{5}{2}^+$ state in ^{126}Te has been measured by IRF method using perturbed angular correlation technique and the g-factor value obtained as $g = -0.56 \pm 0.70$. The angular correlation coefficients of the 171-427 keV cascade, used in the g-factor measurement of the 462 keV state, have also been measured. The magnetic moment of the 462 keV, $\frac{5}{2}^+$ level is computed on the basis of Kisslinger-Sorensen model. (auth)

46958 GYROMAGNETIC RATIO OF THE 2083-keV, 4^+ STATE OF ^{140}Ce . Nigam, A. K.; Bhattacharyya, R. (Saha Inst. of Nuclear Physics, Calcutta). *Indian J. Phys.*; 45: No. 5, 225-9 (May 1971).

The g-factor of the 2083 keV level ($T_{1/2} = 3.41$ ns) of ^{140}Ce was measured using differential perturbed angular correlation method at a magnetic field of about 19 kilo-oersteds. The measurements were performed using γ_1 - γ_2 and γ_1 - γ_3 radiation cascades of the 329-487-1596 keV triple cascade, separately. The values of the g-factor obtained by both measurements are in good agreement. The average value of the g-factor comes out to be $g = 1.16 \pm 0.08$. Besides this, the angular correlation measurements using γ_1 - γ_2 and γ_1 - γ_3 cascades were also performed. A differential angular correlation measurement, however, determined the attenuation of the angular correlation coefficients. (auth)

46959 MAGNETIC MOMENT OF THE 2761 keV 8^+ STATE IN ^{92}Mo . Cochavi, S.; McDonald, J. M.; Fossan, D. B. (State Univ. of New York, Stony Brook). *Phys. Lett.*; 33B: No. 4, 297-8 (26 Oct 1970).

The g-factor for the 2761 keV 8^+ state in ^{92}Mo has been measured by a time-differential technique with the $^{90}\text{Zr}(\alpha,2n)^{92}\text{Mo}$ reaction. The result is $g = 1.40 \pm 0.07$ ($\mu = 11.2 \pm 0.6$ nm) which

is consistent with a pure $(1g_7)^2, 8^+$ wave function. Evidence for an enhanced g_1 is discussed. (auth)

46960 MEASUREMENT OF THE QUADRUPOLE MOMENT OF A SHAPE ISOMER. Gerschel, C.; Perrin, N.; Valentin, L. (Institut de Physique Nucleaire, Orsay, France). *Phys. Lett.*; 33B: No. 4, 299-301(26 Oct 1970). (In French).

A perturbed differential angular correlation measurement on lanthanum magnesium double nitrate yields the quadrupole moment $Q = 1.6 \pm 0.2$ b, and the deformation $\beta = 0.47$ for the 535 keV level of ^{135}La . Discussion of these values and that of the magnetic moment confirms the existence of shape isomerism. (auth)

46961 COULOMB EXCITATION OF ^{117}Sn AND ^{118}Sn . Stelson, P. H.; Milner, W. T.; McGowan, F. K.; Robinson, R. L.; Raman, S. (Oak Ridge National Lab., Tenn.). *Nucl. Phys.*; A190: No. 2, 197-217(1972).

A total of 5 states in ^{117}Sn and 4 states in ^{118}Sn were observed to be Coulomb excited with α -particle and ^{16}O projectiles. Level energies, spins, B(E2) and B(M1) transition probabilities and mean-lives were obtained. For ^{117}Sn , spin $\frac{3}{2}^+$ states were observed at 158.7 and 1005 keV and spin $\frac{5}{2}^+$ states were observed at 1020, 1180 and 1447 keV. For ^{118}Sn , spin $\frac{3}{2}^+$ states were observed at 23.9 and 920.5 keV and spin $\frac{5}{2}^+$ states were observed at 921.4, 1090 and 1354 keV. A total of 20 B(E2) values were obtained and they varied in magnitude from $\frac{1}{10}$ to 10 times the single-particle estimate. About 12 B(M1) values were obtained and they range from 10^{-1} to 10^{-4} of the single-particle estimate. Evidence was found for the existence of quadrupole collective states built on low-lying excited quasiparticle states. (auth)

46962 POSSIBLE EVIDENCE FOR A CONTRIBUTION FROM DIRECT COLLISIONS IN K-SHELL INTERNAL IONIZATION ACCOMPANYING β DECAY. Isozumi, Y.; Mukoyama, T.; Shimizu, S. (Kyoto Univ.). *Phys. Rev. Lett.*; 29: No. 5, 298-301(31 Jul 1972).

Rigorous calculations of the energy-dependent K-shell internal ionization probability were performed by using an improved one-step theory of electron shakeoff. The measured probability for ^{147}Pm considerably exceeds the theoretical curve at low electron energies. This suggests that the direct collision process may play an important role when both a β particle and a K electron are emitted with low kinetic energies. (auth)

46963 GAMMA GAMMA DIRECTIONAL CORRELATION MEASUREMENTS IN ^{131}Cs . Sturm, R.; Thiele, F. W.; Thielmann, R.; Gruenberg, E. (Univ., Marburg, Ger.). *Z. Phys.*; 252: No. 4, 312-23(1972). (In German).

^{131}Cs decay was studied by angular correlation measurements of the (404-216), (832-216), and (247-373) keV γ cascades using NaI(Tl) detectors in connection with a two-parameter multichannel analyzer. Unique spin assignments were obtained for the 216 ($J^\pi = \frac{3}{2}^+$) and 373 ($J^\pi = \frac{3}{2}^+$) keV levels. A previously discussed value of $J^\pi = \frac{5}{2}^+$ for the 1048-keV level was ruled out. The multipole character of the 404-keV transition was found to be E2 with less than 2% M1. (2 figures, 2 tables) (auth)

46964 LOW-LYING STATES AND CONFIGURATIONS IN ^{138}La . Lasiyo, Romedon (Florida State Univ., Tallahassee); Sheline, Raymond K.; Griffioen, Roger D.; DuBard, James L. *Z. Phys.*; 253: No. 1, 17-27(1972).

Tritons from the reaction $^{138}\text{La}(d,t)^{138}\text{La}$ at $E_d = 16$ MeV were analyzed at eleven reaction angles from 22° to 90° with a broad-range magnetic spectrograph. The Q value of the reaction is -2522 ± 5 keV. The nine lowest-lying states in ^{138}La are interpreted in terms of the shell-model configurations $(\pi g_7)^{-1}(\nu d_{3/2})^{-1}$, $(\pi g_7)^{-1}(\nu s_{1/2})^{-1}$, and $(\pi g_7)^{-2}(\nu d_{3/2})^{-1}(\nu d_{5/2})^{-1}$. Seven levels in the energy range of 700 to 1300 keV are populated by $l = 5$ transitions and are interpreted as coming from the $(\pi g_7)^{-1}(\nu h_{9/2})^{-1}$ configuration. The ground state of ^{138}La is shown to have $J^\pi = 5^+$; therefore, beta decay by unique, second-forbidden transitions to the 2^+ one-phonon states of ^{138}Ce and ^{138}Ba must be inferred in spite of the unusually high log-ft values of 19.2 and 18.5, respectively. (5 figures, 2 tables) (auth)

46965 COLLECTIVE STATES OF EVEN-EVEN NUCLEI OF Mo AND Ru INCLUDING UNPAIRED NUCLEONS. Peker, L. K. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2390-2 (Nov 1971). (In Russian).

In even-even nuclei, collective states, including unpaired nucleons, should possess considerable energy and therefore are located in the region of spectra with a large density of levels. It is shown that in even-even Mo and Ru nuclei there are manifest collective levels, including unpaired nucleons. (K.S.W.)

46966 EXPERIMENTAL NEUTRON RESONANCE SPECTROSCOPY. Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970). 544p. \$29.50.

Separate abstracts were prepared for each of the five chapters in this book. (R.W.R.)

For abstracts of individual chapters see: 45641, 46967-46969, and 47036.

46967 TOTAL NEUTRON CROSS SECTION MEASUREMENTS. Firk, F. W. K. (Yale Univ., New Haven); Melkonian, E. pp 101-54 of *Experimental Neutron Resonance Spectroscopy*. /Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970).

A brief introduction is followed by sections on experimental techniques, analysis of neutron spectrometer transmission data, the self indication method, and data obtained from total cross section measurements. An extensive list of references is included. (J.W.R.)

46968 NEUTRON SCATTERING AND CAPTURE CROSS-SECTION MEASUREMENTS. Rae, Ernest R. (Atomic Energy Research Establishment, Harwell, Eng.); Block, Robert C. pp 155-233 of *Experimental Neutron Resonance Spectroscopy*. /Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970).

Measurements in the resonance region, including determination of partial widths and spins, and average cross section measurements are discussed in the introduction. General considerations and experimental measurements on neutron capture and scattering are discussed. Shell transmission and activation measurements on neutron capture are covered. Some results are reviewed. An extensive list of references is included. (J.W.R.)

46969 GAMMA RAYS FROM NEUTRON CAPTURE IN RESONANCES. Bollinger, Lowell M. (Argonne National Lab., Ill.). pp 235-345 of *Experimental Neutron Resonance Spectroscopy*. /Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970).

Sections are included on general characteristics of the spectra, apparatus, analysis of spectra, and experimental results. An extensive list of references is also included. (J.W.R.)

46970 LOG-F TABLES FOR BETA DECAY. Gove, N. B.; Martin, M. J. (Oak Ridge National Lab., Tenn.). *Nucl. Data, Sect. A*; 10: No. 3, 205-317(Nov 1971).

An extensive tabulation of log-f values is presented for use in the calculation of the comparative half-lives for beta transitions. Values of capture-to-positron ratios are also given. These log-f and ϵ/β^+ values incorporate screening and finite nuclear size corrections. They are tabulated for allowed and first-forbidden unique transitions for Z from 6 to 95 as a function of transition energy. The range of transition energies covered is 0.010 MeV to 10 MeV with energy increments chosen so that intermediate values can be obtained from the tables by means of linear extrapolation. The values for first-forbidden unique transitions are larger than those given in older tables by 1.079, the log of 12. (auth)

46971 DIRECT MEASUREMENT OF gamma-RAY ENERGIES AND HALF-LIFE OF ^{141}Cs SEPARATED BY PAPER ELECTROPHORESIS. Tamai, Tadaharu; Takada, Jitsuya; Matsushita, Rokuji; Kiso, Yoshiyuki (Kyoto Univ.). *J. Nucl. Sci. Technol. (Tokyo)*; 9: No. 6, 378-80(Jun 1972).

Cesium-141 was separated from fission products using paper electrophoresis. The γ spectra from the decay of ^{141}Cs were studied using a Ge(Li) detector and a multichannel analyzer. The observed γ energies and intensities were compared with previous data. A half-life of 25 sec was determined. (J.W.R.)

46972 GAMMA RAY SPECTROSCOPIC STUDIES OF MEMBERS OF THE A = 141 DECAY CHAIN. Todd, Richard Ross. East Lansing, Mich.; Michigan State Univ. (1971). 199p. University Microfilms Order No. 72-16,527.

Thesis.

The techniques of beta and gamma ray spectroscopy have been applied to a study of the decay schemes of ^{141m}Sm and ^{141}Pm . The goal of this study was to increase our understanding of nuclear structure and to obtain information about energy level systematics in these members of the A = 141 decay chain below the N = 82 closed shell. Ge(Li) singles, Ge(Li)-Ge(Li) two parameter coincidence and Ge(Li)-NaI(Tl) (anti) coincidence spectrometers were used to identify and establish the sequence of gamma ray transitions. The decay schemes were determined and comparisons are made with previous studies as well as the results of recent reaction work. Forty-seven transitions have been identified with the beta decay of ^{141m}Sm and 40 of these have been incorporated in a decay scheme involving 16 levels in the daughter ^{141}Pm . These levels are at 0, 196.6, 628.6, 804.5, 837.1, 974.0, 1108.1, 1167.2, 1313.2, 1414.8, 1834.0, 1983.1, 2063.5, 2091.6, 2119.0, and 2702.4 keV. The lifetime of the state at 628.6 keV in ^{141}Pm was measured to be 700 ± 20 ns and the E3 transition from this level to the ground state was found to exhibit an enhancement of 1.6 spu. The decay of ^{141}Sm was observed to populate levels at

403.9, 438.4, 728.0, 1292.7, 1495.7, 2004.5, and 2037.7 keV in ^{141}Pm . Of the 50 transitions observed in the decay of ^{141}Pm , 43 of these have been placed in a level scheme containing 23 levels. The levels identified are at 0, 193.8, 1223.3, 1345.8, 1564.8, 1596.8, 1820.4, 1897.1, 1968.0, 2066.4, 2073.7, 2109.6, 2145.2, 2246.6, 2265.3, 2303.4, 2354.4, 2388.4, 2430, 2506, 2620, 2805, and 2986 keV. Limits on the possible spin assignments of the levels identified have been made from calculated log ft values and observation of transitions to levels of known spins. A three-quasiparticle multiplet of high-spin, odd parity states was identified between 1.4 and 2.7 MeV in ^{141}Pm . The decay characteristics of the multiplet suggest the possibility of obtaining information about its structure from shell model calculations using a limited basis. A survey has been made of the energy level systematics and properties of low-lying states in the $N = 80, 81$, and 82 odd mass isotones. These observations have led to some predictions and suggestions for future experiments. (Diss. Abstr. Int., B)

46973 ISOTOPE SHIFT STUDIES IN THE RARE EARTHS AND LEAD. Lee, Paul Lung Sang. Pasadena, Calif.; California Inst. of Tech. (1972). 88p. University Microfilms Order No. 72-17,644.

Thesis.

Isotope shifts of $K_{\alpha 1}$ x-ray transitions were measured for the Neodymium isotopes Nd 142, 143, 144, 145, 146, 148 and 150, the Samarium isotopes Sm 147, 148, 149, 150, 152 and 154, the Gadolinium isotopes Gd 154, 155, 156, 157, 158 and 160, the Dysprosium isotopes Dy 162 and 164, the Erbium isotopes Er 166, 168 and 170, the Hafnium isotopes Hf 178 and 180 and the Lead isotopes Pb 204, 206, 207 and 208. A curved crystal Cauchois spectrometer was used. The analysis of the measurement furnished the variation of the mean square charge radius of the nucleus, $\delta \langle r^2 \rangle$, for 23 isotope pairs. The experimental results were compared with theoretical values from nuclear models. Combining the x-ray shifts and the optical shifts in Nd and Sm yielded the optical mass shifts. An anomaly was observed in the odd-even shifts when the optical and the x-ray shifts were plotted against each other. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, 150 ≤ A ≤ 189

Refer also to abstracts: 46149, 46875, 46916, 46953, 46962, 46973, 47008, and 47066.

46974 (FRNC-TH-207) DE-EXCITATION OF THE COMPOUND NUCLEI FORMED IN $^{181}\text{Ta} + ^{11}\text{B}$ REACTIONS. Delagrè, Hugues (Bordeaux Univ. (France)). 1972. 87p. (In French). Dep. NTIS (U. S. Sales Only).

Thesis.

Twelve isotopes were identified after the irradiation of thick-recoil thick type stacks of Ta targets with ^{10}B and ^{11}B ions having energies in the 10 MeV per nucleon range. Very complicated spectra were obtained from nonchemically treated targets using high resolution α analysis equipment. Statistical analysis techniques were used to resolve the decay curves and to accurately determine the efficiency of the Ge(Li) junction used. The production cross sections and the mean paths of eight of the isotopes were measured. The form of the excitation functions and the variation of the mean path with energy demonstrate that ^{188}Pt , ^{186}Ir and ^{185}Os result from the formation of the compound nucleus. Calculation of the production cross sections made for these three isotopes, using the statistical evaporation model, explain the experimental results, the two principal parameters being the moment of inertia of the nuclei and the γ transition strength parameters. The magnitude of the energy carried off by the first neutrons confirms the existence of γ emission at the end of the evaporation chain. (France)

46975 (JINR-P4-6462) ON PARAMETRIZATION OF B(E2)-TRANSITION IN DEFORMED EVEN-EVEN NUCLEI WITHIN THE SU(3) SCHEME. Ralchev, P. P. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 18p. (In Russian). Dep. NTIS (U. S. Sales Only).

A problem is discussed concerning possible applications of the SU(3) scheme to deformed even-even nuclei. The transition probabilities are parametrized using the free parameters arising in the SU(3) scheme. The results obtained are compared with experimental data. (3 figures, 1 table) (auth)

46976 (JINR-P6-6492) STUDY OF THE ^{179}Re DECAY. THREE-QUASIPARTICLE STATES IN ^{179}W . Arlt, R.; Bayar, B.; Vylov, Ts.; (and others) (Joint Inst. for Nuclear Research,

Dubna (USSR). Lab. of Nuclear Problems). 1972. 20p. (In Russian). Dep. NTIS (U. S. Sales Only).

The decay scheme of ^{179}Re has been studied. The following rotational bands in the decay of ^{179}Re were identified: $7/2^-$ (514), $1/2^-$ (512), $9/2^-$ (624), $5/2^-$ (512), $7/2^+$ (633) and three-quasiparticle states: $3/2^+$ ($1/2^-$ (514), $5/2^+$ (402), $1/2^-$ (514)] and $7/2^+$ ($1/2^-$ (514), $5/2^+$ (402), $1/2^-$ (514)]. (auth)

46977 NEW ISOTOPES OF IRIDIUM. Nadzhakov, E.; Bochev, B.; Venkova, Ts.; (and others). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2202-6 (Nov 1971). (In Russian).

In the irradiation of Ho by the ions ^{22}Ne and ^{20}Ne there are obtained highly-neutron-deficient isotopes of iridium. They were identified by means of their genetic association with daughter isotopes of osmium and rhenium. The new isotopes discovered were ^{181}Ir (10 ± 2 min), ^{180}Ir (6.5 ± 1.5 min), and ^{179}Ir (4 ± 1 min). (tr-auth)

46978 ELECTRON CONVERSION SPECTRUM OF ^{182}Re . Gavriljuk, V. I.; Kupryashkin, V. T.; Latyshev, G. D.; (and others). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2232-48 (Nov 1971). (In Russian).

The electron conversion spectrum of isomers of ^{182}Re with $T_{1/2} = 12.7$ hr and 64 hr was investigated with a resolution of 0.03 to 0.04% with respect to H/p. Many unknown transitions were discovered. The internal conversion ratio and the multipole order of the majority of γ transitions were determined. A mixture of multipole orders was established according to the ratios of intensities of conversion lines in the L- and M-subshells. The decay scheme was constructed. (tr-auth)

46979 ^{175}Hf STATES EXCITED DURING THE DECAY OF ^{175}Ta . Gadzhokov, V.; Zvol'ski, I.; Molnar, I.; Nenov, N. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2255-65 (Nov 1971). (In Russian).

The excited levels of ^{175}Hf which arose from the decay of ^{175}Ta were studied, and the electron conversion spectrum, γ rays, and $\gamma\gamma$ -coincidence were evaluated. On the basis of experimental data a decay scheme for the process $^{175}\text{Ta} \rightarrow ^{175}\text{Hf}$ is proposed with 23 new levels. (tr-auth)

46980 ROTATIONAL LEVELS WITH HIGH SPINS IN ^{171}Lu . Barneoud, D.; Foin, C.; Baudry, A.; (and others) (Univ., Grenoble, France). J. Phys. (Paris); 33: No. 1, 15-22 (Jan 1972). (In French).

Energies and intensities of the gamma-rays following ^{189}Tm ($\alpha, 2n$) ^{171}Lu reaction were measured with Ge(Li) detectors. Rotational bands involving the $1/2^+$ [404] (ground state) $1/2^-$ [541], $1/2^+$ [411], $5/2^+$ [402], $5/2^-$ [514] intrinsic states were identified using the two-dimensional γ - γ coincidence technique and analyzed with classical methods. The large perturbations observed for $1/2^-$ [541] band are studied in terms of a Coriolis mixing in the $3/2^-$ [532] band. The best fit is obtained when the energy of the $3/2^-$ [532] band head is comprised between 600 and 700 keV. A tentative identification of this state at 674 keV is suggested. (France)

46981 MAGNETIC PROPERTIES OF ^{153}Eu , ^{155}Eu , AND ^{189}Os . Begzhanov, R. B.; Gaffarov, D. G.; Muminov, A. I.; Radzhapov, P. S. (Inst. of Nuclear Physics, Tashkent). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2295-9 (Nov 1971). (In Russian).

The half-lives of the levels 69, 95, and 219 keV of the ^{189}Os nucleus: (2.12 ± 0.10), ≤ 0.3 and ≤ 0.3 nsec, were measured using a delayed $\gamma\gamma$ coincidence method. The angular correlation of six cascades of γ radiation from the ^{189}Os nucleus was investigated. The g-factors of the excited states 103.2 keV for ^{153}Eu , 105 keV for ^{155}Eu , and 69 and 95 keV for ^{189}Os were measured. (tr-auth)

46982 PARITY NONCONSERVING EFFECTS IN NUCLEI ^{175}Lu AND ^{181}Ta . Grabowski, J. (Inst. of Nuclear Physics, Krakow). Phys. Lett.; 33B: No. 4, 268-70 (26 Oct 1970).

The nuclear part R of the parity nonconserving circular polarization in 343 keV and 396 keV gamma transitions in ^{175}Lu and 482 keV gamma transition in ^{181}Ta have been calculated. (auth)

46983 STRUCTURE OF GROUND STATES OF ODD-ODD NUCLEI ^{160}Tb , ^{182}Ta , ^{182}Re , AND OF ISOMER ^{182m}Re . Galan, P. (Inst. of Physics, Bratislava). Czech. J. Phys.; 22: No. 6, 468-75 (1972).

The ground states of the odd-odd nuclei ^{160}Tb , ^{182}Ta , and ^{182}Re and the isomeric state ^{182m}Re are interpreted in terms of the Nilsson nuclear model. The conclusions about the structure of these states were drawn from the experimental results obtained in the study of their decay to the neighboring even-even nuclei as well as from the systematics of single-particle proton and neutron states in neighboring odd-mass nuclei. The agreement between experimental and calculated values of the moments of inertia of these states is demonstrated. (auth)

46984 LEVEL STRUCTURE OF ^{187}Os FROM THE DECAY OF ^{187}Ir . Ahlgren, Karin; Daly, P. J. (Purdue Univ., Lafayette, Ind.). Nucl. Phys.; A189: No. 2, 368-84(1972).

The decay of ^{187}Ir was investigated using Ge(Li) detectors in singles and coincidence γ -ray measurements. While the measured γ -ray energies and intensities were in good agreement with the values obtained in a recent study, the coincidence results were invaluable in correctly placing all the strong transitions and in establishing an extensive ^{187}Os level spectrum accommodating more than 98% of the decay intensity. An interpretation of the ^{187}Os level structure in terms of the Nilsson model is presented which includes assignments of levels in the 500 to 1000 keV excitation energy range to the Nilsson orbitals $3/2^+ [624]$, $3/2^- [501]$, and $5/2^- [503]$. Evidence for considerable phonon-quasiparticle mixing in ^{187}Os is discussed. (3 figures, 3 tables) (auth)

46985 ISOBARS FROM THE SPALLATION REACTION FOR NUCLEAR SPECTROSCOPY. Arlt, R.; Bystrov, V. A.; Habenicht, W.; (and others) (Joint Inst. for Nuclear Research, Dubna, USSR). Nucl. Instrum. Methods; 102: No. 2, 253-6(1972).

A method for isolation of radioactive nuclei of the isobars of some elements with a first ionization potential less than 6.5 eV from a target irradiated by 660 MeV protons is described. For this purpose an electromagnetic mass-separator and ion-source with surface ionization have been used. An estimation of the activities of the nuclei obtained on the collector of the mass-separator is done and methods for their identification are described. As an illustration, the experimental results of the Rb and Sr and some rare earth nuclei production are given. A new isotope ^{71}Rb ($T_{1/2} = 3.9$ min) has been identified. (auth)

46986 STUDY OF DECAY OF STATES OF A NUCLEUS OF 89 NEUTRONS: ^{155}Dy . Torres, J. P. (Centre de Spectrometrie Nucleaire et de Spectrometrie de Masse, Orsay, France); Paris, P.; Lecouturier, D.; Kilcher, P. Nucl. Phys., A189: No. 3, 609-31(1972). (In French).

The decay of ^{155}Ho to ^{155}Dy ($T_{1/2} = 48$ min) was investigated by means of Ge(Li) and Si(Li) detectors and a β -spectrograph having a homogeneous field. With the help of a multidimensional coincidence device, a scheme with 21 levels explained in terms of single-particle Nilsson-model and vibrational levels is proposed. To describe the sequence of the levels an important Coriolis coupling has to be introduced. The configurations involved and their band-head energies are for negative-parity levels: $3/2^- [521]$ (0 keV), $5/2^- [523]$ (136.3 keV), $3/2^- [532]$ (202.4 keV), and for positive-parity levels: $3/2^+ [651]$ (240.3 keV), $1/2^+ [660]$ (320.0 keV), $5/2^+ [642]$ (408.5 keV). (auth)

46987 EXPERIMENTAL PROOF OF THE EXISTENCE OF TWO BRANCHINGS IN THE ROTATIONAL BAND $1/2^- [541]$ IN ^{171}Lu . Barneoud, D.; Foin, C.; Baudry, A.; Gizon, A.; Valentin, J. (Inst. of Nuclear Physics, Grenoble, France). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2291-4(Nov 1971). (In Russian).

The reaction $^{168}\text{Tm}(\alpha, 2n)^{171}\text{Lu}$ was used to determine the rotational band and to detect branching for which $I + 1/2$ was an even, whole number. The experimental data also established the spacing of this $1/2^- [541]$ rotational band in ^{171}Lu . (K.S.W.)

46988 CORIOLIS COUPLING OF THE BANDS $1/2^- [541]$ AND $3/2^- [532]$ IN ^{171}Lu . Piepenbring, R. (Inst. of Nuclear Research, Grenoble, France). Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2309-11(Nov 1971). (In Russian).

States of ^{171}Lu with a negative parity which arise from $h_{9/2}$ -states in nuclei in the rare-earth region with odd proton number are discussed. Good development of bands have been observed for ^{171}Lu ; accordingly this isotope was used to detect Coriolis pairing. The bands $1/2^- [541]$ and $3/2^- [532]$ were investigated, and the best agreement between experimental data and calculated data is obtained if the $3/2^-$ -level is assumed to lie between 600 and 700 keV. (K.S.W.)

46989 HYPERFINE STRUCTURE OF THE GROUND STATE OF ^{165}Ho . Merzyn, G.; Penselin, S.; Wolber, G. (Univ., Bonn). Z. Phys.; 252: No. 5, 412-20(1972).

Four hyperfine structure frequencies were measured in the ground state $4I_{9/2}$ of ^{165}Ho with high precision by using the atomic-beam magnetic resonance method at magnetic fields of up to 800 oersted. In the analysis of these measurements a possible electrical sedecimpole (16-pole) interaction was included. The following multipole interaction constants could be calculated: $A = 800.582 \pm 8$ (14) MHz, $B = -1668.100$ (91) MHz, $C = -2224$ (7520) Hz, $D = -398$ (790) Hz. Thus, within the limits of this experiment no octopole or sedecimpole interaction could be established. (4 tables) (auth)

46990 DIRECT MEASUREMENT OF THE NUCLEAR MAGNETIC DIPOLE MOMENT OF ^{165}Ho WITH THE ATOMIC BEAM

MAGNETIC RESONANCE METHOD. Haberstroh, R. A.; Moran, T. I.; Penselin, S. (Univ., Bonn). Z. Phys.; 252: No. 5, 421-7(1972).

Four rf transitions between different Zeeman levels of the $4I_{9/2}$ ground state of ^{165}Ho were measured in an external magnetic field of about 300 gauss with an atomic-beam magnetic resonance apparatus. The interaction between the nuclear magnetic dipole moment and the external field could be deduced from these measurements. Because the magnetic field was measured by calibration transitions in ^{39}K , ^{85}Rb , and ^{87}Rb , the following value could be determined for the nuclear magnetic dipole moment: $\mu_I(^{165}\text{Ho}) = 4.094(44) \mu_N$ (uncorrected for diamagnetic shielding). The g_I factor of the ground state of ^{165}Ho was measured to be $g_I(^{165}\text{Ho}) = 1.1951445(40)$. (1 figure, 2 tables) (auth)

46991 $2^+ \rightarrow 2^+$ TRANSITION OF 879 keV IN ^{160}Dy INVESTIGATED BY ELECTRON-GAMMA DIRECTIONAL CORRELATION. Zupancic, M.; Bikit, I.; Cvjeticanin, D.; Marinkov, L. (Boris Kidric Inst. of Nuclear Science, Belgrade). Z. Phys.; 252: No. 3, 237-41(1972).

$e-\gamma$ directional correlation measurements of the $2^+(879 \text{ keV}) \rightarrow 2^+(87 \text{ keV}) \rightarrow 0^+$ cascade in ^{160}Dy were made, and the magnitude of the electric monopole to quadrupole mixing ratio for the 879-keV transition was found to be $q_K = -0.03(10)$ for $\lambda = 0$ (λ according to the Hager-Saltzer definition). (2 figures, 1 table) (auth)

46992 MIXING OF TWO-PROTON BANDS WITH TWO-NEUTRON BANDS IN ^{176}Hf . Khoo, T. L.; Waddington, J. C.; O'Neil, R. A.; (and others) (McMaster Univ., Hamilton, Ont.). Phys. Rev. Lett.; 28: No. 26, 1717-20(26 Jun 1972).

Evidence for the interaction of high-spin two-quasiproton bands with two-quasineutron bands in ^{176}Hf is presented. The two $K^\pi = 6^+$ bands at 1333.1 and 1761.5 keV are shown to be of highly mixed character. A $K^\pi = 8^-$ band at 1559.4 keV, which is largely of two-proton character, is shown to be mixed with a predominantly two-neutron $K = 8^-$ band at 1860.3 keV. (auth)

46993 DEFORMATION PARAMETERS OF ^{152}Sm BY ELECTRON SCATTERING. Bertozzi, W. (Massachusetts Inst. of Tech., Cambridge); Cooper, T.; Esselin, N.; (and others). Phys. Rev. Lett.; 28: No. 26, 1711-13(26 Jun 1972).

Cross sections for excitation of the ground-state rotational band in ^{152}Sm by high-energy electrons were measured. The ground-state charge distribution has been determined including the deformation parameters β_2 and β_4 of the nuclear surface. (auth)

46994 EFFECTIVE g_s FACTOR FOR THE ODD PROTON ^{153}Sm NUCLEUS DETERMINED BY THE INTERNAL CONVERSION PROCESSES. Krpic, D.; Milanovic, J.; Bikit, I.; Stepic, R. (Boris Kidric Inst. of Nuclear Sciences, Belgrade). Z. Phys.; 253: No. 1, 71-7(1972).

The K/L_3 , L_1/L_3 , L_1/L_2 , and L_2/L_3 conversion ratios for the 103-keV transition in ^{153}Sm were measured by means of a high-resolution $\pi\sqrt{2}$ electron spectrometer. The values of the M1 conversion penetration parameter $\lambda = 5.0 \pm 0.4$, and the mixing ratio $\delta^2 = 0.144 \pm 0.006$ are deduced. By comparing the experimental penetration parameter with the value obtained from Nilsson-model calculations the effective spin gyromagnetic ratio is found to be $g_s^{\text{eff}} = (3.7 \pm 0.4) \mu_N$. (1 figure, 1 table) (auth)

46995 VALIDITY OF THE ROTOR MODEL IN THE ODD-Z NUCLIDE ^{171}Lu . Hjorth, S. A.; Ryde, H.; Skaaneberg, B. (Univ. of Stockholm). J. Phys. (Paris); 33: No. 1, 23-6(Jan 1972).

A metallic thulium target was irradiated with 28-MeV α -particles to produce the $^{169}\text{Tm}(\alpha, 2n)^{171}\text{Lu}$ reaction. The excitation functions for the observed γ -rays and the angular distributions of these γ -rays were studied. A search was made for isomeric transitions with half-lives longer than 10 ns. The 137.05 keV γ -ray depopulating the $1/2^+ [411]$ state into the $1/2^- [541]$ level was found to have a half-life of 33 ± 8 ns. No other delayed transitions were observed. Rotational bands built on the $1/2^+ [411]$, $5/2^+ [402]$, $1/2^+ [404]$ (ground-state), $1/2^- [541]$ and $3/2^- [514]$ intrinsic states were obtained. The properties of these bands are examined with respect to deviations from the rotor model. Evidence is obtained of coupling term with $\Delta K = 2$, a variable moment of inertia explaining the available data. As usual the assumption of independent quasiparticle motion overestimates the effect of the Coriolis force. (France)

46996 BAND MIXING IN ^{156}Gd AND ^{158}Gd . Kluk, A. F. (Oak Ridge National Lab., Tenn.); Johnson, Noah R.; Hamilton, J. H. Z. Phys.; 253: No. 1, 1-12(1972).

A detailed investigation of the energies and intensities of the γ -rays that depopulate the low-spin levels of the β - and γ -vibrational bands of ^{156}Gd and the γ -vibrational band of ^{158}Gd was conducted. Both singles and γ - γ coincidence measurements were made on sources of 15-d ^{156}Eu and 46-min ^{158}Eu by use of large-

volume, high-resolution Ge(Li) detectors. In addition to the γ band at 1154.09 keV, two $K^\pi = 0^+$ bands were observed in ^{156}Gd with band heads at 1049.45 and 1168.11 keV. The 2^+ and 3^+ members of the γ -vibrational band in ^{156}Gd were observed at 1187.12 and 1265.43 keV, respectively; a new $K^\pi = 0^+$ band was seen at 1195.98 keV. A first-order perturbational treatment of the branching ratios was applied to both nuclei. In addition, the mixing between the ground state, the β^- , and the γ -vibrational bands of ^{156}Gd was considered from two approaches, but neither satisfactorily explains all the experimental B(E2) ratios. (3 figures, 2 tables) (auth)

46997 DECAY OF 1.4 Min ^{173}Er . Pursiheimo, V.; Tuurnala, T.; Raumamaa, T. (Univ. of Helsinki). *Z. Phys.*; 252: No. 4, 283-99(1972).

A (1.4 ± 0.1) -min activity, which is assigned to β^- decay of ^{173}Er , was produced with 14- to 15-MeV neutrons through the reaction $^{178}\text{Yb}(n, \alpha)^{173}\text{Er}$. Its decay was studied with Ge(Li) detectors and several γ - γ as well as β - γ coincidence arrangements. Eight gamma rays with energies 94.2, 116.14, 118.6, 122.40, 192.8, 199.2, 800.8, and 895.2 keV were assigned to the decay of ^{173}Er . The proposed level scheme of the daughter nuclide ^{173}Tm contains levels at 0, 2.5, 118.6, 124.9, 317.7, 411.9, and 1212.8 keV. The 317.7-keV level is an isomeric state with a measured half-life of 10 ± 3 μs . This 10- μs isomer and the 411.9-keV level are thought to be the band head and $5/2^-$ rotational member of $1/2^-$ [523] Nilsson state, respectively. The levels at 0, 2.5, 118.6, and 124.9 keV are interpreted as $1/2^+$, $3/2^+$, $5/2^+$, and $7/2^+$ members of the $1/2^+$ [411] band; and the level at 1212.8 keV is interpreted as the $5/2^-$ [514] Nilsson state. Some systematic considerations and theoretical transition probability calculations are also included. (7 figures, 4 tables) (auth)

46998 GAMMA- AND X-RAYS IN THE DECAY OF ^{164}Ho . von Droste, G. Frh.; Pessara, W. (Physikalisch-Technische Bundesanstalt, Brunswick). *Z. Phys.*; 252: No. 4, 300-11(1972). (In German).

The photon spectrum in the decay of ^{164}Ho was investigated using a high-resolution solid-state detector separating the K x rays of Dy, Ho, and Er. The two γ rays depopulating the 3^+ and 2^+ levels in ^{164}Ho were found to have energies of 56.6 and 37.3 keV. The 91.4-keV radiation of Er from the sample seems to have a nonresolved high-energy component with about 2.5% intensity. It may be due to the cross-over transition $3^+ \rightarrow 1^+$ or a sum line. The half-life of the isomeric state of ^{164}Ho was found to be 36.7 ± 1 min. The total transition rates between different levels were calculated from the measured photon intensities. The fraction 0.27 found in the branching of the e^- -capture feeding the 2^+ state of Dy has nearly the same value as the analog branching in the β^- decay. By combining the measured complex ground-state decay curve and the transition intensities of the isomeric and the ground state, the ground-state half-life was deduced; it was found to be (28.1 ± 1.0) min. The corresponding ratio of the ground-state and the isomeric-state production rates is $2.84:1$. A ratio of 0.04 was found for the cross sections of the $^{165}\text{Ho}(\gamma, n)^{162}\text{Ho}$ and $^{165}\text{Ho}(\gamma, n)^{164}\text{Ho}$ reactions for a 90-MeV bremsstrahlung spectrum. (13 figures, 2 tables) (auth)

46999 NEW ISOTOPE ^{189}Tl . Vandlik, T. B.; Vandlik, Ya.; Zaitseva, N. G.; (and others) (Joint Inst. for Nuclear Research, Dubna, USSR). *JETP Lett. (USSR) (Engl. Transl.)*; 15: No. 7, 271-4(5 Apr 1972).

Translated from *Zh. Eksp. Teor. Fiz., Pis'ma Red.*; 15: No. 7, 386-9(5 Apr 1972).

A PbF_2 target was bombarded by 660-MeV protons, the Tl was separated on-line, and the Tl fraction was separated with a mass separator. Gamma spectra were taken with a Ge(Li) detector having a resolution of 3.5 keV at 600 keV. After all known nuclides were taken into account, three gamma rays (of 216.3, 229.0, and 335 keV) were left; they were assigned to the decay of isotope ^{189}Tl with a half-life of 1.4 ± 0.4 min. (R.W.R.)

47000 MEAN LIVES OF SOME HIGH ENERGY LEVELS IN ^{160}Dy . Abou-Leila, Hussein; Darwish, Saher; Abdel-Halim, A. F. *Compt. Rend., Ser. B*; 274: No. 14, 896-9(5 Apr 1972). (In French).

The mean life of the 1358.7 keV level of ^{160}Dy was determined using a double magnetic lens spectrometer and time-to-amplitude converters. An upper limit was determined for the 966.1 and 1264.7 keV excited levels. The following results were obtained: tau (1358.7 keV) = 3.9 ± 0.2 nsec, tau (966.1 keV) ≤ 20 psec, and tau (1264.7 keV) ≤ 15 psec. (France)

47001 NEW ISOTOPES OF TUNGSTEN. Nadzhakov, E.; Bochev, B.; Venkova, Ts.; (and others). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2207-12(Nov 1971). (In Russian).

In reactions of the composite nucleus with neutrons during the irradiation of ^{155}Gd and ^{156}Gd with heavy ions ^{22}Ne and ^{20}Ne there

is produced neutron-deficient isotopes of tungsten. They were identified through their genetic association with the daughter isotopes of tantalum and hafnium. The new isotopes were identified as ^{171}W (9.0 ± 1.5 min) and ^{170}W (4 ± 1 min). (tr-auth)

Nuclear Properties and Reactions, $190 \leq A \leq 219$

Refer also to abstracts: 46461, 46883, 46909, 46916, 46973, and 47073.

47002 g-FACTOR OF THE 12^+ ISOMERIC STATE IN ^{206}Pb AND THE EFFECTIVE ORBITAL g-FACTOR OF THE NEUTRON. Nakai, K. (Univ. of Copenhagen); Herskind, B.; Blomqvist, J.; (and others). *Nucl. Phys.*; A189: No. 3, 526-38(1972).

The g-factor of the 12^+ two-quasiparticle isomeric state, $[(v1\frac{1}{2})^-]_{12^+}$, in ^{206}Pb has been measured by the time-differential perturbed angular correlation method. The result obtained is $g(12^+, ^{206}\text{Pb}) = -0.155 \pm 0.004$. Three different methods are applied to deduce the orbital g-factor of the neutron around the ^{208}Pb core, namely (i) the M1 core polarization model, (ii) the configuration mixing model, and (iii) a combination of the M4 transition rate in ^{207}Pb with the present result. The g_1^{eff} (neutron) deduced in the three different ways are in good internal agreement: (i) -0.036 ± 0.010 , (ii) -0.040 ± 0.03 and (iii) -0.034 ± 0.010 . Although it is hard to estimate the overall uncertainty, it is probably significant that the anomaly in the orbital g-factor of the neutron has opposite sign from that of the proton, and that the size is smaller. From known g-factors in the lead region the polarization part δg_s and $\delta g'_s$ of the g-factor of the neutron can be derived. These values are $\delta g_s = 2.77 \pm 0.15$ and $\delta g'_s = -3.30 \pm 0.22$. By comparing the experimental g-factors for the $i_{1/2}$ proton and neutron we find that the part of the g-factor deviating from the Schmidt line has an isovector character. This means that δg_s and $\delta g'_s$ should have the same magnitude but opposite sign for the proton. Calculated g-factors based on this conclusion agree well with the experimental g-factors for the $h_{9/2}$ and $i_{13/2}$ protons. (auth)

47003 DECAY OF ^{194}Au . Vishnevskii, I. N.; Gavriluk, V. I.; Kupryashkin, V. T.; (and others). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2213-31(Nov 1971). (In Russian).

The e^- spectrum of ^{194}Au ($T_{1/2} = 39.5$ hr) was investigated on the iron β -spectrometer $\pi\sqrt{2}$, $r_0 = 50$ cm, with a resolution of 0.03 to 0.04%, and the accompanying γ spectrum was determined on the Ge(Li) detector with a resolution of 4 keV. The multipole order of the majority of transitions was established; a mixture of multipole orders were confirmed according to the ratio of intensities of the conversion lines in the L- and M-subshells for many transitions. Many new transitions were detected in the e^- spectrum. A decay scheme was constructed. (tr-auth)

47004 α -DECAY OF NEUTRON-DEFICIENT ISOTOPES OF RADON. Golovkov, N. A.; Ivanov, R. B.; Kolachkovski, A.; Norseev, Yu. V.; Chumin, V. G. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2272-80(Nov 1971). (In Russian).

The α -spectra of Rn isotopes having masses of $206 \leq A \leq 212$ were investigated with a magnetic α -spectrograph. New α -transitions were detected, and their relative intensities were measured. Earlier α -group energies and a portion of the α -decay scheme were revised to provide better accuracy. The dependence of $T_{1/2}$ or E_α was studied for isotopes with neutron numbers $N > 128$ and $N \leq 128$. (tr-auth)

47005 K CONVERSION COEFFICIENT NEAR THRESHOLD: $^{191\text{m}}\text{Os}$. Kirschner, H. (Univ., Goettingen, Ger.). *Z. Phys.*; 252: No. 3, 249-54(1972). (In German).

The K-conversion coefficient of the predominantly M3 transition $^{191\text{m}}\text{Os} - ^{191}\text{gOs}$ $E_\gamma = 74.37$ keV was measured to be $\alpha_K = 131 \pm 12$. This differs from other measurements and from theoretical values. (3 figures) (auth)

47006 DECAY OF ^{210}At TO LEVELS IN ^{210}Po . Jardine, L. J.; Prussin, S. G.; Hollander, J. J. (Univ. of California, Berkeley). *Nucl. Phys.*; A190: No. 2, 261-83(1972).

The electron-capture decay of ^{210}At has been reinvestigated and present data define twenty-three levels in ^{210}Po . The multipolarity of thirty-six transitions has been measured and combined with data from recent reaction studies to assign spins and parities to these levels. Evidence is presented which locates the 3^- collective level in ^{210}Po at 2387 keV above the ground state. The electron-capture transition rates to odd-parity levels are discussed with reference to the particle-hole excitations in ^{208}Pb . Transition probabilities for γ -decay of members of the $\pi(h_{9/2})$ multiplet are compared to predictions from recent theoretical calculations. (auth)

47007 FIRST FORBIDDEN BETA DECAY OF ^{209}Pb . Behrens, H.; Kobelt, M.; Thies, W. G.; Appel, H. (Univ., Karlsruhe, Ger. Kernforschungszentrum, Karlsruhe, Ger.). *Z. Phys.*; 252: No. 4, 349-61 (1972).

The $\frac{3}{2}^+ \rightarrow \frac{3}{2}^-$ first-forbidden β transition of $^{209}\text{Pb} \rightarrow ^{209}\text{Bi}$ was investigated. The end-point energy was determined to be (644.6 ± 1.2) keV; the half-life to be (3.253 ± 0.014) h. For the shape factor no deviation from the allowed form was found within an accuracy of 1%. The shape factor and ft-value can be excellently interpreted by the shell model in jj coupling. (auth)

47008 VERY INELASTIC SCATTERING OF ALPHA PARTICLES FROM HEAVY NUCLEI. Halpern, I. (Univ. of Washington, Seattle. Univ. of Copenhagen). pp 255-62 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Energy spectra and cross sections were measured for (α, α') , $(\alpha, ^3\text{He})$, $(^3\text{He}, ^3\text{He}')$, and $(^3\text{He}, \alpha)$ reactions on ^{181}Ta , ^{197}Au , and ^{208}Pb for bombarding energies of about 50 up to 80 MeV. The observed structure in the spectra and cross sections is discussed. (J.W.R.)

47009 NUCLEAR ENERGY LEVELS IN EVEN ISOTOPES OF LEAD FROM (α, xn) REACTIONS ON SEPARATED MERCURY ISOTOPES. Burns, Erskine John Thomas. Davis, Calif.; Univ. of California (1971). 355p. University Microfilms Order No. 72-17,899.

Thesis.

Levels in ^{206}Pb , ^{204}Pb , ^{202}Pb , ^{200}Pb and ^{198}Pb are identified with 30 to 55 MeV. (α, xn) reactions on enriched Hg targets with $x = 2$ and 4. The data includes in-beam, beam-swept and off-beam gamma-ray spectra corresponding to time distributions of 3 to 125 nanoseconds, 100 to 625 nanoseconds and greater than .1 seconds; along with in-beam conversion electrons. Assignments of levels are based on relative intensities, both prompt (< 3 ns.) and delayed (> 3 ns.), relative excitation functions, conversion coefficients, and energy level systematics. New (7 $^-$) isomers (> 3 nsec.) are assigned to ^{202}Pb , ^{200}Pb and ^{198}Pb with half-lives of 80 nsec., 310 nsec., and greater than 1 μ sec; respectively. These isomers are believed to be two-particle (quasi-particle) configuration mixed states made up of the $(i_{1/2}, p_{3/2})$, $(i_{3/2}, p_{3/2})$ and $(i_{1/2}, i_{3/2})$. Energy level systematics and transition probabilities tested the following microscopic models: (1) strict shell model; (2) pairing plus quadrupole model; (3) surface delta residual interaction; and (4) extended shell model with a realistic residual interaction. States populated via (α, xn) on lead are shown to be relatively pure two-particle (quasi-particle) levels. (Diss. Abstr. Int., B)

Nuclear Properties and Reactions, A $\geq$ 220

Refer also to abstracts: 46148, 46461, 46889, 46966, 47042, and 47955.

47010 (ANCR-1060) TOTAL NEUTRON CROSS SECTION OF ^{243}Am FROM 0.5 TO 1000 eV. Simpson, O. D.; Simpson, F. B.; Harvey, J. A.; Slaughter, G. G.; Benjamin, R. W.; Ahlfeld, C. E. (Aerojet Nuclear Co., Idaho Falls, Idaho). Mar 1972. Contract AT(40-1)-1375. 95p. Dep. NTIS.

Transmission measurements were made on two high-purity samples of ^{243}Am having inverse thicknesses of 1288.2 and 279.3 b/a. Data were collected from 0.5 to 1000 eV using the Oak Ridge Electron Linear Accelerator. High-resolution data were taken using 10-ns, 140-MeV electron bursts, 10-ns channel widths, and a flight path of 18.576 meters. Single-level Breit-Wigner resonance parameters were obtained from area analysis below 250 eV. The resonance parameters were also determined below 18 eV from shape analysis. From an analysis of 24 resonances an average value of $\Gamma = 39 \pm 1$ MeV was determined. The average observed level spacing between resonances was found to be 0.67 ± 0.06 eV. A neutron S-wave strength function of $(0.96 \pm 0.10) \times 10^{-4}$ was determined from the resonance parameters. The resonance-absorption integral for neutrons with energies above 0.625 eV was calculated to be 1813 ± 68 b. (22 figures, 6 tables) (auth)

47011 (BNWL-1648) PHOTON INTENSITIES AND THEIR IMPORTANCE IN COUNTING TRANSURANIC MATERIALS. Swinsh, K. L. (Battelle Pacific Northwest Labs., Richland, Wash.) May 1972. Contract AT(45-1)-1830. 28p. Dep. NTIS.

Ratios of x-ray to alpha intensities are presented for a number of transuranic materials, and how these ratios can vary with

changing isotopic composition is illustrated. Plutonium is used as an example. Also shown is how the biological hazard as measured by the maximum permissible body burden (MPBB) can vary with isotopic composition for plutonium. Plutonium from spent power reactor fuels presents a greater hazard on a unit weight basis than high-grade fissile material. The x-ray intensity is initially smaller for the power reactor material but increases as the ^{241}Am grows into the material. In fact, the x-ray intensity will eventually be greater than that from the high-grade material. Although the isotopic composition is an important facet in determining the biological hazard and in interpreting the photon intensities from transuranic materials, it is not the only complication in interpretation of counting data. The low penetrating power (absorption) and, because of this, the manner in which the material is deposited will also affect the interpretation of counting data. (15 figures, 1 table) (auth)

47012 (CEA-N-1526) LIBRARY OF DATA FOR FISSION PRODUCTS (THIRD EDITION). Blachot, Jean; de Tourrell, Roland (Commissariat a l'Energie Atomique, Grenoble (France). Centre d'Etudes Nucleaires). Mar 1972. 272p. (In French). Dep. NTIS (U. S. Sales Only).

The third version of the library of fission product data is presented. Two tables give a description of the nucleides and gamma transitions. (France)

47013 (COO-3058-23) MeV TOTAL NEUTRON CROSS SECTIONS OF ^{238}U . Hayes, S. H.; Stoler, P.; Clement, J. M.; Goulding, C. A. (Rensselaer Polytechnic Inst., Troy, N. Y. Dept. of Physics). 1972. Contract AT(11-1)-3058. 16p. Dep. NTIS.

A measurement of the total neutron cross sections of ^{238}U was made using the RPI electron LINAC. The operating electron energy was about 60 MeV, and the beam burst width was 20 ns. The neutron energies were determined using the time-of-flight technique with a resolution of .08 ns/m. The cross sections are presented from 0.8 to 30.0 MeV. The cross section curve is smooth in this region, and the averaged data have a statistical precision of 1%. The data reported here have been compared with other recent measurements and with the ENDF-B file. There is generally good agreement with a few percent deviations in certain energy regions. In addition, the experimental cross sections were fit using coupled channel code JUPITOR to obtain optical model and deformation parameters. (auth)

47014 (FRNC-TH-184) CHEMICAL SEPARATION AND HIGH RESOLUTION NUCLEAR SPECTROSCOPY IN THE $^{228}\text{Ra} \rightarrow ^{228}\text{Ac} \rightarrow ^{228}\text{Th}$ RADIOACTIVE CHAIN. Herment, Micheline (Grenoble Univ. (France). Faculte des Sciences). 1971. 70p. (In French). Dep. NTIS (U. S. Sales Only).

Thesis.

The $^{228}\text{Ra} \rightarrow ^{228}\text{Ac} \rightarrow ^{228}\text{Th}$ decay scheme was studied in order to precise the nuclear properties of the odd-odd Ac nucleus. The results were obtained by γ spectrometry, γ - γ coincidence spectrometry, electron spectrometry, and e- γ coincidence spectrometry. 190 transitions were examined in the framework of this study, which enabled the modes of excitation, and the intrinsic states of the deformed nuclei to be investigated. Examination of the exact ratios of the transition probabilities allowed a comparison to be made with theoretical predictions and the confirmation or proposition of certain levels. A level scheme is presented for ^{228}Th and ^{228}Ac . The chemical separation methods used to obtain ^{228}Ra from ^{232}Th , and the preparation of ^{228}Ac and ^{228}Ra sources are presented. (France)

47015 (INDC-7-U) PROCEEDINGS OF THE TOPICAL CONFERENCE ON "NEUTRON INDUCED FISSION" HELD DURING THE FOURTH INDC MEETING, BHABHA ATOMIC RESEARCH CENTER, BOMBAY, WEDNESDAY, JULY 14, 1971. (International Nuclear Data Committee. International Atomic Energy Agency, Vienna (Austria)). 162p. (CONF-710725).

Separate abstracts were prepared for 10 of the 11 papers included. The remaining paper is a brief summary containing no data. (J.W.R.)

For abstracts of individual papers see: 47016-47021 and 47043-47046.

47016 (INDC-7-U, pp 23-5) DID SUPERHEAVY NUCLEI EXIST IN OUR SOLAR SYSTEM? Bhandari, Narendra (Tata Inst. of Fundamental Research, Bombay (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

The available evidence on the existence of superheavy elements is evaluated and the results are discussed in terms of their possible physical and chemical properties. There are two lines of evidences, both obtained from a study of primitive objects of our solar systems, the meteorites and lunar dust grains which indicate that superheavy elements were extant at the time these objects solidified. The rare gas evidence based on the presence of an

anomalous component of fissiogenic xenon has been discussed in detail by Anders and Heymann (1969) and by Rao (1970). Recently Bhandari et al (1971) have observed fission fragment tracks characteristic of nuclei $Z > 110$. These tracks are present in a number of meteorites. The estimated number of tracks due to ^{244}Pu and element $Z > 110$ relative to ^{238}U as observed in certain enriched regions of the silicate crystals are given. Based on these observations arguments are developed for possible chemical nature of elements responsible for the characteristic long (>15 micron) tracks. The fact that the meteorite Angra Dos Reis is highly depleted in such tracks but unusually enriched in U, Pu, Ba etc. lead us to conclude that the element(s) may not be a chemical homologue of these groups. However its exact chemical properties can not be ascertained from the available data. Near absence of the superheavy tracks in lunar rocks which are younger by about 1 billion years indicates that they must have a half life $\lesssim 10^8$ years and must now be extinct. (auth)

47017 (INDC-7-U, pp 31-59) MASS DISTRIBUTIONS IN FISSION AND SOME RELATED PHENOMENA. Felvinci, J.; Melkonian, E.; Havens, W. W. Jr. (Columbia Univ., New York).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

Some experiments on mass distribution and related phenomena in fission are discussed. The systematics of mass distributions are reviewed. Topics discussed include the fission fragment mass distribution, variation with mass of the fissioning nucleus, mass distribution as a function of fragment kinetic energy, primary and post-neutron mass distributions and prompt neutron emission, effect of energy on the mass distribution, variations of mass distribution with spin and/or K quantum number, and effects of mass distribution on cross measurements. (J.W.R.)

47018 (INDC-7-U, pp 61-4) FISSION ISOMERIC STATE IN ^{236}U . Derengowski, M. A.; Felvinci, J. P.; Melkonian, E. (Columbia Univ., New York).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

A delayed fission activity with a half-life of 7 ± 1 nsec was found in the neutron fission of ^{235}U at neutron energies of a few hundred eV up to about 3 keV. A lower limit of 2% was set for the delayed fission relative to prompt fission. (J.W.R.)

47019 (INDC-7-U, pp 89-112) PROMPT RADIATIONS EMITTED IN LOW ENERGY FISSION. Nadkarni, D. M. (Bhabha Atomic Research Centre, Bombay (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

The emission of prompt radiation in fission, including neutrons, γ -rays, conversion electrons, x-rays, and long range charged particles is discussed. Some experimental results are presented and some previous work is reviewed. (J.W.R.)

47020 (INDC-7-U, pp 113-29) RECENT STUDIES ON PROMPT NEUTRON AND GAMMA RAYS FROM FISSION. Conde, H. (Forsvarets Forskningsanstalt, Stockholm (Sweden)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

Some recent experiments on prompt neutron and γ -ray emission in fission are reviewed. Special emphasis is given to results on average prompt neutron yields and fission neutron spectra. (J.W.R.)

47021 (INDC-7-U, pp 131-44) RADIATIONS EMITTED IN FISSION. Ribon, Pierre.

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

The properties of the various particles which are emitted during or after the fission process are briefly examined. Their emission can occur at three different stages: 1- at the scission time. They are mainly α particles and other light particles; the results for these emissions are numerous and coherent, and allow to obtain informations on the scission configuration. On the contrary, the neutron emission is not well enough known. 2- by the fission fragments. These deexcited mainly by neutron emission; nevertheless, if they have high spin values ($I \sim 15$), γ -ray emission is expected to compete strongly with the neutron emission. But some very recent results give lower spin values ($I \sim 7$). 3- by the fission products. The delayed neutron precursors are now quite well known, and the interest is turning more and more upon their energy spectra. The calculations which have been done to fit the previously known spectra does not describe the structure of the most recent results. (auth)

47022 (JINR-P7-6466) ANGULAR DISTRIBUTION OF FRAGMENTS OF SPONTANEOUSLY FISSIONING ISOMERS. Gangrskii, Yu. P.; Nguyen Cong Khan; Pulatov, D. D.; Fam Zui Hein (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Nuclear

Reactions). 1972. 15p. (In Russian). Dep. NTIS (U. S. Sales Only).

Measurements were made of the angular distributions of the fragments from the spontaneously fissioning isomers ^{236}Pu , ^{237}Pu and ^{243}Cm , produced in α -particle reactions. Anisotropy of the angular distribution was observed for ^{236}Pu and ^{237}Pu isomers. The effects of the neutrons and of γ quanta, emitted from the compound nucleus, as well as of extranuclear magnetic fields, on the angular anisotropy are discussed. (5 figures, 1 table) (auth)

47023 (KFK-1537) PREPARATION AND INVESTIGATION OF THE NEPTUNIUM ISOTOPES 231, 232, AND 233. Weiss-Reuter, R. (Kernforschungszentrum, Karlsruhe (West Germany)). Institut fuer Radiochemie). Dec 1971. 54p. (In German). Dep. NTIS (U. S. Sales Only).

The decay of ^{231}Np , ^{232}Np , and ^{233}Np was studied by γ -ray spectroscopy. All three isotopes decay predominantly by electron capture. For ^{233}Np 24 γ -ray transitions were detected. An extended decay scheme is given. For ^{233}Np and ^{231}Np 21 and 15 γ -ray transitions, respectively, were observed; decay schemes for these isotopes are proposed. (16 figures, 18 tables) (auth)

47024 (KFK-1563) MEASURE OF THE RATIO OF CAPTURE-TO-FISSION CROSS SECTION FOR ^{235}U AND ^{239}Pu BETWEEN 8 AND 60 keV NEUTRON ENERGY. Bandl, Rolf-Eckart (Kernforschungszentrum, Karlsruhe (West Germany)). Institut fuer Angewandte Kernphysik. Mar 1972. 71p. (In German). Dep. NTIS (U. S. Sales Only).

The ratio of capture to fission cross section, α , has been measured for ^{235}U and ^{239}Pu in the neutron energy range from 8 to 60 keV. A 3 MV Van-de-Graaff accelerator generated a pulsed proton-beam. The $^7\text{Li}(p,n)^7\text{Be}$ reaction was used as the neutron-source. The absorption was found by a comparison of the scattered neutrons from a fissile sample with those from a non-absorbing lead sample. Simultaneously the fast fission neutrons were detected with an organic liquid scintillator. The neutron energy was determined by the time-of-flight method, the time resolution was about 20 ns/m. The measurements were normalized between 40 and 50 keV to the results of recent evaluations. The error in α amounts to 15% between 10 and 40 keV neutron energy. (auth)

47025 PROMPT GAMMA DECAY OF FISSION FRAGMENTS. Sarkar, R. (Saha Inst. of Nuclear Physics, Calcutta); Chatterjee, A. Phys. Lett.; 33B: No. 4, 263-7(26 Oct 1970).

Prompt γ -ray energies and yields from fission fragments for the thermal neutron fission of ^{235}U and spontaneous fission of ^{252}Cf are calculated from an improved renormalized Fermi gas model and the collective potential energy surface concept. The predictions agree fairly well with the experimental observations. (auth)

47026 FISSION OF ALIGNED ^{233}U NUCLEI BY NEUTRONS FROM 0.4 TO 2000 eV. Kuiken, R. (Kamerlingh Onnes Laboratorium, Leiden); Pattenden, N. J.; Postma, H. Nucl. Phys.; A190: No. 2, 401-18(1972).

The anisotropy in the angular distribution of fission fragments from neutron-induced fission of aligned ^{233}U has been measured as a function of neutron energy from 0.4 to 2000 eV. The ^{233}U nuclei were aligned in thin layers of monocrystalline rubidium uranyl nitrate, effectively cooled to about 0.3 K. The variation in the anisotropy over the resonances in the energy region from 0.4 to 60 eV and also over the unresolved region from 60 to 2000 eV was found to be rather small: Comparing the experimental distribution of anisotropy values with theoretical distributions showed, that for the $J = 3$ resonances, channels with $K = 3$ are not open; K being the projection of the compound spin J on the deformation axis. Channels with $(J^\pi, K) = (2^+, 2)$ are at most partially open. Several channels must be assumed to be open in order to explain the experimental distribution. A slight systematic variation observed in the anisotropy values averaged over 100 eV intervals may indicate the presence of intermediate structure. (auth)

47027 MASS YIELDS FROM THERMAL-NEUTRON INDUCED FISSION OF ^{245}Cm . Harbour, R. M.; MacMurdo, K. W. (E. I. du Pont de Nemours and Co., Aiken, S. C.). J. Inorg. Nucl. Chem.; 34: No. 7, 2109-14(Jul 1972).

The fission yields of 21 mass chains were measured for the thermal-neutron induced fission of ^{245}Cm . Fission product gamma activities from ^{245}Cm were determined relative to those from ^{235}U with a high resolution lithium-drifted germanium detector. The measured errors ranged from 1 to 10% over a mass number range of 85 to 147. No significant fine structure in the mass yield curve was apparent. (auth) (UK)

47028 CONVERSION TRANSITIONS IN ^{226}Ra . Tret'yakov, E. F.; Tret'yakova, N. I.; Konyaev, V. F. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2306-8(Nov 1971). (In Russian).

Internal conversion electron spectra which accompanied the α decay of ^{230}Th were measured. The E1 transitions with energies 186 and 253 keV in ^{226}Ra were determined. (tr-auth)

47029 ENERGY LEVELS IN ^{231}Th AND ^{233}Th . Grottdal, T.; Limstrand, J.; Nybo, K.; Skar, K.; Thorsteinsen, T. F. (Univ. of Bergen. Univ. of Copenhagen). Nucl. Phys.; A189: No. 3, 592-608(1972).

The energy levels of ^{231}Th and ^{233}Th have been studied by the (d,t) and (d,p) reactions at deuteron bombarding energies of 12 and 13 MeV. The scattered particles were analysed in a magnetic spectrograph at angles of observation of 60°, 90° and 125°. A total of 9 Nilsson orbitals in ^{231}Th and 7 Nilsson orbitals in ^{233}Th were identified on the basis of the total cross sections and the intensity patterns for the rotational states. The influence of Coriolis mixing was taken into account. (auth)

47030 ON-LINE MEASUREMENTS OF RARE-GAS FISSION YIELDS IN 14 MeV NEUTRON FISSION. Bocquet, J. P.; Brisot, R.; Crancon, J.; Moussa, A. (CEN, Grenoble, France). Nucl. Phys.; A189: No. 3, 556-76(1972).

The independent and cumulative fission yields of rare-gas isotopes produced from the fission of ^{233}U , ^{235}U , ^{238}U and ^{232}Th by 14 MeV neutrons have been measured by means of on-line isotopic separation. Chain yields can be deduced from the cumulative yields and Gaussian curves fitted to the independent yield distributions. Values of $Z_p - Z_{\text{UCD}}$ and σ_t have been compared with those obtained from other fissioning systems. The total number of neutrons, ν_{tot} , emitted in ^{232}Th fission has been estimated. (auth)

47031 STUDY OF X-RAYS AND NUCLEAR γ -RAYS IN MUONIC THALLIUM. Backe, H.; Engler, R.; Jahnke, U.; (and others). (CERN, Geneva). Nucl. Phys.; A189: No. 3, 472-512(1972).

Energies and intensities of muonic x-rays, nuclear γ -rays and μ -capture γ -rays were measured in natural muonic thallium with Ge(Li) detectors. The absolute intensities of higher μ x-rays were reproduced by a cascade calculation starting with a statistical population at $n = 20$ including K-, L- and M-conversion. The electron screening effect was deduced from energies of higher μ x-rays. Eight prompt nuclear γ -rays were found, due to an excitation of $(4.1 \pm 1.0)\%$ and $(14.6 \pm 3.0)\%$ of the first $3/2^+$ level in μ ^{203}Tl and μ ^{205}Tl , respectively, and of $(8.8 \pm 0.9)\%$ of the $5/2^-$ level at 2.63 MeV in ^{205}Tl . This excitation explains the anomalous intensity ratios of the 2p-1s and 3d-2p fine structure components. From the nuclear γ -rays of the first excited states were deduced: the magnetic hf splittings $\epsilon^{203}(1/2) = 2.57 \pm 0.15$ keV, $\epsilon^{205}(1/2) = 2.30 \pm 0.02$ keV, muonic isomer shifts $\Delta E_{\text{is}}(203) = 0.26 \pm 0.15$ keV, $\Delta E_{\text{is}}(205) = -(0.31 \pm 0.11)$ keV; E2/M1 mixing ratios $\delta_{\text{E}}^2(203) = 0.89 \pm 0.27$, $\delta_{\text{E}}^2(205) = 1.00 \pm 0.05$, and the half-life in the presence of the muon $T_{\mu}(3/2) = 1.4 \pm 0.1$ nsec in ^{205}Tl . The values of $\epsilon(1/2)$ and ΔE_{is} were compared with shell-model calculations. From δ_{E}^2 and $T_{\mu}(3/2)$ evidence for a magnetic nuclear polarization was found. From the 2p-1s, 3d-2p, 3p-2s and 2s-2p transitions the size and shape of the nuclear charge distribution and a nuclear polarization due to the $1s_{\mu}$ muon of $\Delta E_{\text{np}} = 8.3 \pm 1.2$ keV and 9.5 ± 1.2 keV for ^{203}Tl and ^{205}Tl respectively were determined in agreement with theoretical estimates. An isotope shift of $\Delta E = 10.35 \pm 0.25$ keV was measured for the $1s_{\mu}$ state which is compared with data from optical spectroscopy. From an analysis of the time distribution of delayed γ -rays from μ -capture, an isotopic effect of $(5.4 \pm 2.5)\%$ was found for the capture rates, in fair agreement with predictions from Primakoff's μ -capture formula. (auth)

47032 HARD GAMMA RADIATION FROM ^{238}Pu . Makarenko, A. I.; Ostretsov, L. A.; Forafontov, N. V. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2335-7(Nov 1971). (In Russian).

The γ radiation from three sources, prepared from metallic Pu with activities of 5.7, 8.8, and 17.0 Ci, was investigated. During a determination of the relative intensity, the varying effectiveness of the detector at different γ quantum energies and varying degrees of absorption in the lead filter were taken into account. The results are compared with the γ spectra of ^{234}Pa and ^{234}Np . Besides the known γ lines with energies at 743.0, 767.0, and 786.0 keV, there were also observed γ lines at 707, 807, 852, 882, 926, 945, 1,002, 1,039, and 1,083 keV, all of which accompany the α decay of Pu. A revised decay scheme for ^{238}Pu is presented. (K.S.W.)

47033 SUBBARRIER FISSION RESONANCES IN Th ISOTOPES. Back, B. B.; Britt, H. C.; Garrett, J. D.; Hansen, Ole (Los Alamos Scientific Lab., N. Mex.). Phys. Rev. Lett.; 28: No. 26, 1707-10 (26 Jun 1972).

Subbarrier fission resonances have been found in $^{230}\text{Th}(t,pf)$ and $^{232}\text{Th}(d,pf)$ reactions. This is in disagreement with the fission barriers predicted by the Strutinskii theory. (auth)

47034 GAMMA BRANCHING AND TOTAL CONVERSION COEFFICIENTS OF INTRABAND TRANSITIONS IN THE GROUND-

STATE ROTATIONAL BAND OF ^{246}Cm . Schmidt-Ott, W.-D.; Fink, R. W. (Georgia Inst. of Tech., Atlanta). Z. Phys.; 253: No. 1, 13-16(1972).

With calibrated Ge(Li) detectors, γ - γ coincidences were observed in the decay of ^{246}Cf . The total conversion coefficients of the 54.7- and 66.7-keV transitions in the first rotational band were determined to be 237 ± 30 and 69 ± 15 , respectively. The branching ratio of the 121.5-keV crossover transition is $(22 \pm 7)\%$. The errors are the 2σ statistical uncertainties, linearly added to the errors in the detector efficiencies. (auth)

47035 TRANSFER REACTIONS OF A LARGE NUMBER OF NUCLEONS INDUCED BY ^{84}Kr ON ^{232}Th : STUDY OF THE REACTION THRESHOLDS. Bimbot, R.; Gauvin, H.; Le Beyec, Y.; (and others) (Institut Physique Nucleaire, Orsay, France). Nucl. Phys.; A189: No. 3, 539-44(1972). (In French).

Excitation functions have been measured for the formation of ^{225}Th , ^{226}Th , ^{223}Ac , and ^{224}Ac , when ^{232}Th was bombarded by ^{84}Kr ions at energies ranging between 450 and 494 MeV. The thresholds of all the reactions appear at 450 MeV (330 MeV c.m.), a higher energy than expected. Cross sections of the order of 1 mb are reached for these transfer reactions corresponding to a large number of neutrons. The results are compared with the same reactions induced with lighter projectiles. (auth)

47036 MEASUREMENTS ON FISSIONLE NUCLIDES. Moore, M. S. (Idaho Nuclear Corp., Idaho Falls). pp 347-508 of Experimental Neutron Resonance Spectroscopy. /Harvey, J. A. (ed.). New York; Academic Press, Inc. (1970).

After an introduction on the development of fission theory and fission as a multistage process sections are included on experimental methods in low-energy neutron spectroscopy of fissionable nuclides, cross-section measurements on common fissionable isotopes in the resonance region, fragment angular distributions and fission near threshold, the scission point and nuclear structure of the fragments, and the question of spin dependence of the mass distribution. The relationships among the various multilevel approaches to resonance structure are treated in an appendix. In addition, an extensive list of references is included. (J.W.R.)

47037 LONG RANGE ALPHA PARTICLE EMISSION IN THE FISSION OF HEAVY ELEMENTS. Rajagopalan, Mahalingam. Princeton, N. J.; Princeton Univ. (1971). 103p. University Microfilms Order No. 72-14,163.

Thesis.

Emission of alpha particles in the spontaneous fission of ^{252}Cf and in the proton induced fission of ^{238}U was investigated. The angular distribution of alpha particles emitted in coincidence with light as well as heavy fragments were determined in the spontaneous fission of ^{252}Cf . The FWHM (Full width at half maximum) for the angular distribution of alpha particles emitted in coincidence with light fragments was found to be 23 degrees (distinctly narrower than has been previously reported). The FWHM for the angular distribution of alpha particles emitted in coincidence with all the fragments was found to be $37\frac{1}{2}$ degrees. Trajectory calculations were done to find the best set of initial parameters that give rise to the angular distribution of alpha particles in agreement with experiment. The results are consistent with the sudden snap mechanism of alpha particle emission proposed by Halpern. They are also compatible with the adiabatic model of fission and contradictory to predictions by the statistical model. The dependence of the probability of alpha particle emission in fission on excitation energy was investigated by measuring the emission rates in the fission of ^{238}U induced by 16-MeV and 42-MeV protons. It was found that the probability of alpha particle emission in 42-MeV proton induced fission was $20 \pm 5\%$ higher than in the case of 16-MeV proton induced fission. A least square analysis was done with the available data on the alpha particle emission rates in fission induced by intermediate energy projectiles and the best fit was given by $Y_{\alpha} = (1.658 \pm 0.03) + (0.0217 \pm 0.0015) E^*$ where Y_{α} is the alpha particle emission probability per thousand binary fissions and E^* is the excitation energy of the fissioning system. The observed increase in the alpha particle emission rate with increase in excitation energy is consistent with expectations from the two-mode fission hypothesis. (Diss. Abstr. Int., B)

Nuclear Theory

Refer also to abstracts: 46466, 46508, 46644, 46658, 46787, 47015, and 47209.

47038 (BNL-tr-487) CHANGE IN THE MECHANISM OF INELASTIC INTERACTIONS BETWEEN PARTICLES AND NUCLEI IN THE ENERGY REGION $T \approx 5$ GeV. Barashenkov, V. S.; Gudima, K. K.; Toonev, V. D. (Joint Inst. for Nuclear Research,

Dubna (USSR)). Translated for Oak Ridge National Lab., Tenn., from report JINR-P2-4302. 9p. Dep. NTIS.

The analysis of the available experimental data on inelastic nucleon- and meson-nucleus interactions shows that the energy dependence of different characteristics connected with the cascade and evaporation particle production is changed in the energy region $T \approx 5$ GeV. This change is of "saturation" character, and an essential modification of the conventional intranuclear cascade model is required for its interpretation. One of the possible explanations is based on the progressive decrease of the number of nucleons in the nucleus as the intranuclear cascade develops. (4 figures) (auth)

47039 (BNL-tr-488) SELF-CONSISTENT FIELD METHOD AS APPLIED TO THE NUCLEUS. Kompaneets, A. Translated by S. J. Amoretti (Brookhaven National Lab., Upton, N. Y.), from Dokl. Akad. Nauk SSSR; 85: No. 2, 301-4(1952). 8p. Dep. NTIS.

The self-consistent field method is used to explain, in general terms, certain regularities in nuclear structure; the approximate constant energy and volume per nucleon. Many constants of the nuclear force being unknown, the work is formulated in a way that minimizes the use of specific numerical values. (R.W.R.)

47040 (BNL-tr-489) TOWARD THE THEORY OF NUCLEAR SHELLS. Ivanenko, D. D.; Sokolov, A. A. Translated for Brookhaven National Lab., Upton, N. Y., from Dokl. Akad. Nauk SSSR; 74: No. 1, 33-6(1950). 7p. Dep. NTIS.

An approximate general formula is derived for the "critical" numbers N of nucleons completing successive nuclear shells as a function of the nucleonic density ρ . The statistical method of Fermi, which had already been employed for the approximate deduction of critical numbers of electrons in an atom, was followed. The statistical model leads directly to the formula $N = (21 + 1)^3/24 \pi^2 r^3 \rho$. Assuming a certain distribution of ρ , the N 's are obtained successively for $l = 0, 1, 2, 3$, etc. By taking $\rho = \text{constant}$, for heavy nuclei, one gets $N = 0, 1, 2, 6, 12, 24, 38, 60, 86, 122 \dots$ If, for light nuclei, ρ is assumed to vary continuously in space, the numbers obtained are $N = 2, 10, 28, 60, 110 \dots$ The assumptions having an approximate, preliminary value only, the results should be regarded as giving only a certain general answer to the problem. Corrections, such as that for the Coulomb interaction between protons, or that for the effect of the surface layer, could be introduced. Valente [Phys. Rev. 78: 77(1950)] is an analysis of the empirically known series of "magic" numbers, obtained the formula $N = (l^3 + 51)/3$ which is related to that given above. (TSS)

47041 (CEA-N-1529) GEOMETRICAL PROPERTIES OF NUCLEAR REACTIONS. Raynal, Jacques (Commissariat à l'Énergie Atomique, Saclay (France). Centre d'Études Nucleaires). Mar 1972. 118p. (In French). Dep. NTIS (U. S. Sales Only).

The angular dependence of reaction products with a polarized beam and a polarized target exhibits some characteristics which do not depend upon the reaction mechanism but upon the particle and target spin values. A general description of a polarized beam is given. The properties of the reaction amplitudes with respect to the spins in the initial and the final states are studied, taking into account parity and time reversal invariances. Certain properties of the azimuthal distribution with respect to the incident beam, for the cross section and the polarizations of the outgoing particles are presented. The number of angular distributions which can be defined is a function of the spins. (France)

47042 (COO-3001-12) IMPORTANCE OF THE QUADRUPOLE PAIRING FIELD IN THE $J^\pi = 0^+$ VIBRATIONS OF SHAPE DEFORMED NUCLEI. Bes, D. R. (Comision Nacional de Energía Atómica, Buenos Aires (Argentina). Los Alamos Scientific Lab., N. Mex.); Broglia, R. A. (State Univ. of New York, Stony Brook. Inst. for Theoretical Physics. Los Alamos Scientific Lab., N. Mex.); Nilsson, B. (State Univ. of New York, Stony Brook. Dept. of Physics). 1972. Contract AT(11-1)-3001. 16p. Dep. NTIS.

The monopole and quadrupole pairing residual interactions acting simultaneously on a system of particles which move in a set of Nilsson single-particle levels and which display a non-uniform distribution of positive and negative quadrupole moments around the Fermi surface give rise to BCS state dependent gaps, which can produce effects which are similar to those appearing when there is an energy gap in the single-particle spectrum. The competition between monopole and quadrupole pairing correlations allows for the existence of $J^\pi = 0^+$ pairing vibrations which carry large (p,t) cross sections and weak (t,p) cross sections or vice versa. This model is used in the analysis of the observed two-nucleon transfer data on the U-isotopes. (auth)

47043 (INDC-7-U, pp 3-19) EXCITATION ENERGY DEPENDENCE OF SHELL EFFECTS IN NUCLEAR FISSION. Kapoor, S. S. (Bhabha Atomic Research Centre, Bombay (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

Some aspects of nuclear fission which are related to nuclear shell effects and their dependence on excitation energy are discussed. Properties of the fission barrier and the ground state deformation are analyzed. (J.W.R.)

47044 (INDC-7-U, pp 65-79) THEORY OF FRAGMENT MASS AND CHARGE DISTRIBUTION IN FISSION. Ramamurthy, V. S. (Bhabha Atomic Research Centre, Bombay (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

A model for fission which assumes the preformation of fragments in the last stages of the division process is discussed. It is shown that with certain simplifying assumptions regarding the structure of single-particle levels in the nascent fragments, the observed mass and charge distributions for fission fragments as well as their dependence on the excitation energy of the fissioning nucleus may be explained. (J.W.R.)

47045 (INDC-7-U, pp 81-6) DISTRIBUTION OF THE TOTAL FISSION WIDTH. Ullah, Nazakat (Tata Inst. of Fundamental Research, Bombay (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

Some recent developments in the distribution of the total fission width are presented. General expressions for the correlation coefficients of the resonance parameters which take into account the effects due to unitarity are given. A two-resonance two-channel model is used to derive an expression for the distribution of the total width. (auth)

47046 (INDC-7-U, pp 145-60) FISSION BARRIER HEIGHTS, ENERGY RELEASE, AND FRAGMENT ENERGY PARTITION PROCESSES FROM THE VIEWPOINTS OF THE NUCLEAR STRUCTURAL MODELS. Chatterjee, A. (Calcutta Univ. (India). Saha Inst. of Nuclear Physics, Calcutta (India)).

From fourth meeting on neutron induced fission; Bombay, India (14 Jul 1971).

Some recent theoretical work on various aspects of nuclear fission phenomena is briefly summarized. The work is mainly on the development of microscopic models to explain fission barrier heights, energy release, and fragment energy partition processes. (J.W.R.)

47047 (JINR-P2-6503) ACCOUNT OF NUCLEAR BOUNDARY DIFFUSIVITY IN THE INTRANUCLEAR CASCADE MODEL. Barashenkov, V. S.; Gudima, K. K.; Zhereg, F.; Toneev, V. D. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 9p. (In Russian). Dep. NTIS (U. S. Sales Only).

Taking into account the nuclear potential diffusivity and the density distribution of nuclear constituents in Monte-Carlo calculations is considered. This method does not use the division of the nucleus into spherical zones with constant density. The developed method corresponds to more realistic nuclear model and, at the same time, simplifies essentially the intranuclear cascade calculations, especially in the case of interactions of high-energy ions with nuclei. (auth)

47048 (JINR-P4-6452) POSSIBLE EXISTENCE OF SU(3) MULTIPLETS IN SPECTRA OF DEFORMED EVEN-EVEN NUCLEI. Raichev, P. P. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 15p. (In Russian). Dep. NTIS (U. S. Sales Only).

An assumption is made that the excited states of deformed even-even nuclei are combined into some SU_3 multiplets. The problem on the origin of these multiplets is discussed. (auth)

47049 (JINR-P4-6458) $SU(3)$ PHYSICAL BASIS REALIZATION AND PROBABILITIES OF $B(E2)$ -TRANSITIONS IN $SU(3)$ SCHEME. II. Afanas'ev, G. N.; Avramov, S. R.; Raichev, P. P. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Theoretical Physics). 1972. 16p. (In Russian). Dep. NTIS (U. S. Sales Only).

The probabilities of transitions inside the rotational bands and between them are calculated in the framework of the $SU(3)$ scheme. It is shown that there can be chosen the operators Ω_1 and Ω_2 separating sufficiently well the rotational bands. (auth)

47050 β, γ TRANSITIONS AND STRUCTURE OF ANTI-ANALOG STATES. Aleshin, V. P.; Kolomiets, V. M. Izv. Akad. Nauk SSSR, Ser. Fiz.; 35: No. 11, 2368-73(Nov 1971). (In Russian).

The possibility of excitation and the structure of anti-analog configuration states and states with a polarized core are discussed in connection with $\beta\gamma$ processes. Electromagnetic transitions between anti-analog states and concomitant T_c -states in odd nuclei ($N, Z + 1$) are examined in shell models. (K.S.W.)

47051 REGULARITY IN MULTIPLET PROPERTIES OF ODD-ODD NUCLEI. Vorob'ev, V. D.; Peker, L. K.; Kharitonov, Yu. I. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2374-9 (Nov 1971). (In Russian).

A shell model was used to explain regularity in multiplet properties relative to a deformation of the multiplet with change in the number of particles in the shell. A large number of theoretical investigations which contained calculations of levels of odd-odd nuclei lead to the conclusion that the dominating influence in multiplet formation was the p-n pair interaction. Three isotopes with $N = 127$ were examined in the ^{208}Pb region: ^{212}At , ^{214}Fr , and ^{216}Ac , in addition to a number of Bi and Nb isotopes. Study of the ^{90}Zr region was confined to the isotopes $^{94,96}\text{Nb}$. (K.S.W.)

47052 SPIN CORRELATION DURING BREMSSTRAHLUNG RADIATION OF LONGITUDINALLY POLARIZED ELECTRONS, ALLOWING FOR MULTIPOLE MOMENTS OF THE NUCLEI. Kerimov, B. K.; Sokolov, A. A.; El Gavkhar, A.; Ablakulov, Kh. A. (Moscow State Univ.). *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2380-5 (Nov 1971). (In Russian).

The influence of a magnetic dipole moment of a nucleus with spin I on the transfer of spin of a longitudinally polarized electron during emission of circularly polarized bremsstrahlung had previously been studied. That result is here generalized for the case in which the electric quadrupole and magnetic octupole moments of the nucleus-target are considered. A formula was obtained which determined the effect of longitudinal polarization of a scattered electron and of multipole moments of the target-nucleus on the extent of circular polarization of decelerating protons. (K.S.W.)

47053 ISOTOPIC STRUCTURE OF THE GIANT DIPOLE RESONANCE AND THE ISOSPIN FREQUENCY IN THE GROUND STATE OF THE NUCLEUS. Kuriyandskii, A. S. *Izv. Akad. Nauk SSSR, Ser. Fiz.*; 35: No. 11, 2393-8 (Nov 1971). (In Russian).

The excitation of analog states in the process of photoabsorption was studied. The particle-hole interaction in a channel leads to the formation of the giant dipole resonance in the cross section of photoabsorption. This process may be interpreted as the excitation of a "collective state" of a given particle-hole type. The isotopic structure of giant dipole resonance, and the isospin frequency are examined. (K.S.W.)

47054 VALIDITY OF DWBA FOR DEUTERON STRIPPING. I. THE MITRA THREE-BODY MODEL. Bouldin, D. P.; Levin, F. S. (Brown Univ., Providence). *Nucl. Phys.*; A189: No. 3, 449-71 (1972).

Previous work showing that there exists an exact formulation of the DWBA for stripping in the S-wave, separable potential, three-body model of Mitra is discussed and extended. The one-body equation obeyed by the c.m. wave function used in the reformulated DWBA is derived and compared with the equation obeyed by the wave function used in the standard formulation of DWBA, viz., the deuteron elastic scattering wave function. Results obtained by other workers on application of three-body methods to direct reactions are discussed in light of the fact that an exact DWBA exists for the separable potential model. (auth)

47055 CHARGE-SYMMETRY BREAKING AND N-N SCATTERING LENGTHS. Henley, E. M.; Keliher, T. E. (Univ. of Washington, Seattle). *Nucl. Phys.*; A189: No. 3, 632-40 (1972).

Recent experimental and theoretical investigations of ρ - ω and η - π mixing have caused us to re-examine the effects of these mixings on the charge-symmetry breaking of nuclear forces. The charge asymmetry of the nucleon-nucleon force in the singlet S-state is used to predict the n-n scattering length in terms of the measured p-p one. The extra attraction of the n-n potential in the $^1S_0$ state gives rise to a change of roughly -0.8 fm in the corresponding scattering length. The magnitude (and sign) of the charge asymmetry is approximately that required to understand the binding-energy differences of the mirror nuclei ^3He - ^3H and ^{41}Sc - ^{41}Ca . (auth)

47056 HEAVY ION NUCLEAR REACTIONS. PHENOMENA INTERMEDIATE BETWEEN THE PHENOMENON OF THE COMPOUND NUCLEUS AND DIRECT TRANSFER EFFECTS. Basile, R. (Univ., Orleans, France); Galin, J.; Guerreau, D.; Lefort, M.; Tarrago, X. *J. Phys. (Paris)*; 33: No. 1, 9-14 (Jan 1972). (In French).

An exchange of nucleons is observed in heavy ion induced nuclear reactions in the very forward direction. It involves a deep nuclear interaction in which the nuclear matter is compressed. An attempt is given to explain this process using Coulomb potential, centrifugal barrier, Bruckner's attractive and repulsive nuclear potentials in the region (high nuclear density) where the projectile enters into the target nucleus. Transitory compound

systems are formed when l is lower than a given value of h which depends on the energy (43h at 78 MeV, 54h at 100 MeV). By considering the number of excitons created in the area of interaction, it is found that for l lower than 37h, the number of excitons is large enough to carry a true compound nucleus process. A range of l value between 37 and 42h for 78 MeV (and between 46 and 53h for 100 MeV) yields a very short lived state which might explain the very inelastic cases of nucleon transfers. (France)

47057 CALCULATION OF THE NUCLEAR QUADRUPOLE MOMENT. Stamp, A. P. (Univ. of Auckland, New Zealand). *Phys. Lett.*; 33B: No. 4, 257-9 (26 Oct 1970).

The single-particle configurations which must be taken into account to give reliable results for the calculation of quadrupole operators are determined. (auth)

47058 THEORY OF ANOMALIES OBSERVED IN (d,p) EXCITATION CURVES AT THRESHOLDS FOR NEUTRON ANALOGUE CHANNELS. Lane, A. M. (United Kingdom Atomic Energy Authority, Harwell, Eng.). *Phys. Lett.*; 33B: No. 4, 274-8 (26 Oct 1970).

It is shown that several essential features of the (d,p) threshold anomalies observed with targets $A \approx 90$ may be understood physically without elaborate calculations. In particular, their narrow width and their occurrence on threshold over the mass range 80-100 results from a remarkable energy compression effect operating on the single particle 3p-neutron state which is basically responsible for the anomaly. It is remarked that the wide variety of observed anomalies are classified very neatly by the Q-values of the (d,p) reactions. (auth)

47059 ISOSPIN DEPENDENCE OF OPTICAL MODEL POTENTIALS USING A REALISTIC EFFECTIVE INTERACTION. Sinha, B. C. (King's Coll., London). *Phys. Lett.*; 33B: No. 4, 279-80 (26 Oct 1970).

A linear iso-spin dependence of the optical model potential is found using a strong density dependent effective interaction. A correction, which depends on nuclear matter distribution, is suggested to the volume integral of the optical potential. (auth)

47060 SINGLE-PARTICLE SPIN-ORBIT POTENTIAL FOLLOWING FROM THE NUCLEON-NUCLEON SPIN-ORBIT INTERACTION. Penzel, R.; Stocker, W. (Univ., Munich). *Nucl. Phys.*; A190: No. 2, 218-28 (1972).

The single-particle spin-orbit potential following from the nucleon-nucleon spin-orbit interaction is investigated for the case of a plane nuclear surface. Scheerbaum's result, that in the case when the triplet-even part of the spin-orbit interaction vanishes the exchange term is equal in first inhomogeneity order to the direct term, is verified. The second-order contribution to the exchange part is given explicitly. For a Gaussian form of the two-nucleon spin-orbit interaction it compensates partly for the first-order contribution, the ratio being about $-1/3$. The spin-orbit splittings following from the two-nucleon interaction of Green are calculated. (auth)

47061 SPECTROSCOPY WITH TWO-NUCLEON TRANSFER REACTIONS IN DEFORMED NUCLEI. I. FORM FACTORS. Ascuitto, R. J. (Yale Univ., New Haven); Soerensen, Bent. *Nucl. Phys.*; A190: No. 2, 297-308 (1972).

A method for the evaluation of two-nucleon transfer form-factors is devised, which is suitable for the treatment of reactions between deformed nuclei. The nuclear-structure-dependent parts are evaluated in a rotating (intrinsic) frame. (auth)

47062 SPECTROSCOPY WITH TWO-NUCLEON TRANSFER REACTIONS IN DEFORMED NUCLEI. II. RARE-EARTH $K = 0$ BANDS STUDIES WITH THE (p,t) REACTION. Ascuitto, R. J. (Yale Univ., New Haven); Soerensen, Bent. *Nucl. Phys.*; A190: No. 2, 309-27 (1972).

The ratio of the (p,t) strength to excited $K = 0$ bands relative to the ground band varies from a few to about 40% across the rare-earth region. This behavior is reproduced by a calculation that uses conventional pairing plus quadrupole interactions for the structure calculation and includes inelastic excitations in the treatment of the reaction. (auth)

47063 VERTEX PARAMETERS AND COUPLING CONSTANTS FOR THE VIRTUAL DECAY OF LIGHT NUCLEI. Rinat, A. S. (Weizmann Inst. of Science, Rehovot, Israel); Kok, L. P.; Stingl, M. *Nucl. Phys.*; A190: No. 2, 328-40 (1972).

Various sources of information on vertex functions are compared. Explicit wave functions, scattering and reaction data, and dispersion relations are used to extract range parameters and spectroscopic factors of parametrized vertex functions describing the virtual decay of nuclei with $A \leq 6$. Mutual agreement is found in general, with the exception of the ^6Li - ^4He -d vertex parameters. (auth)

47064 NUCLEON-NUCLEUS POTENTIAL IN THE HARTREE-FOCK APPROXIMATION WITH SKYRME'S INTERACTION. Dover, C. B. (Brookhaven National Lab., Upton, N. Y.); Nguyen Van Giai. Nucl. Phys.; A190: No. 2, 373-400(1972).

The connection between the self-consistent nuclear field in the Hartree-Fock (HF) approximation with the Skyrme interaction and the real part of the phenomenological nucleon-nucleus optical potential was investigated. It is shown that a simple local transformation $u(r) = (m^*(r)/(m))^{\frac{1}{2}} u^L(r)$ of the radial wave function $u(r)$ leads to an ordinary Schroedinger equation for $u^L(r)$. The energy-dependent potential $V_L(r, E)$ which appears in this equation is identified with the real part of the nucleon optical potential. $V_L(r, E)$ is decomposed into a central part V_{av} , a symmetry part V_{sym} and a spin-orbit part $V_{s.o.}$. The radial form factors which characterize V_{av} , V_{sym} and $V_{s.o.}$ are expressed in closed form in terms of the total nuclear density $\rho(r)$ and the neutron excess density $\Delta\rho(r)$. Using force parameters obtained by fitting bound-state properties of closed-shell nuclei, V_L is calculated numerically and compared to empirical potentials. V_{av} strongly resembles a Woods-Saxon form, with depth $V_0 \approx -43$ MeV, radius $R_0 = 1.10A^{\frac{1}{3}} + 0.75$ fm and diffuseness $a \approx 0.55$ fm; V_{av} also contains an energy-dependent part of depth 0.43 E and radius $R_0^E = R_0 - 0.25$ fm. It is shown that V_{sym} is proportional to $\Delta\rho(r)$ times a smooth function of the density. The calculated symmetry potential changes from a surface form for medium nuclei (Ca) to a surface plus volume for heavier nuclei (Zr, Sn, Pb). The form factor for $V_{s.o.}$ is close to a Thomas form, but peaks at a radius inside that of the central potential. (auth)

47065 METHOD FOR SOLVING FADDEEV EQUATIONS. Benayoun, J. J. Gignoux, C. (Institut des Sciences Nucleaires, Grenoble, France). Nucl. Phys.; A190: No. 2, 419-36(1972).

Faddeev equations, written as coupled integro-differential equations, are solved by using a relaxation method. The initial guess solution for this method is obtained by solving Faddeev equations by means of separable expansions for two-body Green functions. This method is applied to the problem of the bound states for three bosons interacting via two potentials, one of them having a repulsive part. Preliminary results on the triton problem are given. (auth)

47066 SYSTEMATICS OF EXCITATION FUNCTIONS FOR FAST NEUTRON INDUCED (n,p)- AND (n, α)-REACTIONS. Krivan, V.; Muenzel, H. (Nuclear Research Centre, Karlsruhe, Ger.). J. Inorg. Nucl. Chem.; 34: No. 7, 2093-2108(Jul 1972).

The systematics of excitation functions for (n,p)- and (n, α)-reactions were investigated. The dependence of the position and value of the maximum cross-section, the half-width and an "asymmetry factor" on the charge number of the target material were evaluated from published data. Using these relationships, a procedure was developed for the construction of unknown excitation functions. To check this procedure, the activities produced by (n,p)-reactions with ^{27}Al , ^{51}V , ^{59}Co , ^{75}As and ^{103}Rh , and by (n, α)-reactions with ^{51}V , ^{75}As , ^{133}Cs and ^{165}Ho , were experimentally determined. The neutrons were obtained by bombarding a thick Be converter with 33 MeV and 53 MeV deuterons. Only for As do the determined activities deviate appreciably (by about a factor of two) from the values calculated with the aid of the calculated excitation functions. In all other cases the agreement was within 35%. (auth) (UK)

47067 TWO-PARTICLE ORBITAL MATRIX OF DENSITY FOR ONE-QUANTA OSCILLATORY-EXCITED SU_3 -CONFIGURATIONS FOR p-NUCLEI. Kalinauskas, R. K.; Vanagas, V. V. (Inst. of Physics and Mathematics, Vilnius, USSR). Liet. Fiz. Rinkiny; 11: No. 4, 581-93(1971). (In Russian).

In the case of one-quanta oscillatory-excited states of p-shell nuclei the decomposition of the unitary scheme wave function in terms of wave functions characterized by SU_3 -configurations has been found. The explicit expressions for the spin-isospin independent two-particle matrix of density in space of SU_3 -configurations for the maximum oscillatory energy for the relative motion of a pair of nuclei have been obtained. (auth)

47068 RESONANCE SCATTERING THEORY AND THE STRUCTURE OF COMPOUND STATES. Dumitrescu, O.; Kuemmel, H. (Ruhr Univ., Bochum, Ger.). Ann. Phys. (N. Y.); 71: No. 2, 556-68(Jun 1972).

A rigorous equation for compound-state wave functions (with only one channel open) is obtained. It is shown that the compound states may have a complex structure; however, this difficulty can be overcome using some solvable models. One such solvable model is proposed for the case of α decay. (auth)

47069 GENERATOR-COORDINATE FORM FACTORS FOR TRANSFER REACTIONS BETWEEN HEAVY IONS. Bonche, P.; Giraud, B. (CEN de Saclay, Gif-sur-Yvette, France). Phys. Rev. Lett.; 28: No. 26, 1720-3(26 Jun 1972).

In the framework of the distorted-wave Born approximation, a generator-coordinate formalism is used for optical waves. This results in a microscopic prescription for the calculation of transfer reaction form factors. A preliminary analysis of the reaction $^{54}\text{Fe}(^{16}\text{O}, ^{12}\text{C})^{58}\text{Ni}$ gives a reasonable fit to the angular distribution. (auth)

47070 SINGLE-PARTICLE EFFECTS ON FISSION PROBABILITIES FOR THE LIGHTER ELEMENTS. Vandenbosch, R.; Mosel, U. (Univ. of Washington, Seattle). Phys. Rev. Lett.; 28: No. 26, 1726-9(26 Jun 1972).

The excitation energy dependence of the competition between neutron emission and fission has been calculated for nuclei in the vicinity of the ^{208}Pb closed shell. A two-center model has been used to calculate single-particle effects on the nuclear level density as well as on the barrier heights. Comparison with experiment demonstrates the importance of including the dependence of the single-particle level density on deformation. (auth)

47071 NUCLEAR LEVEL SEQUENCE OF SINGLE-PARTICLE ENERGIES IN A MOMENTUM-DEPENDENT POTENTIAL. Lodhi, M. A. K.; Waak, B. T. (Texas Tech Univ., Lubbock). Phys. Rev. Lett.; 29: No. 5, 301-3(31 Jul 1972).

Single-particle energy calculations were made for nuclei from boron through beyond lead, a form of momentum-dependent nucleon-nucleus potential being used. The results are in good agreement with the observed data for energies of single-particle levels. (auth)

47072 LIMITS ON ANGULAR MOMENTUM IN HEAVY-ION-COMPOUND-NUCLEUS REACTIONS. Blann, M. (Univ. of Rochester, N. Y.); Plasil, F. Phys. Rev. Lett.; 29: No. 5, 303-6(31 Jul 1972).

Experimentally measured fusion excitation functions for the reactions induced on ^{27}Al with ^{16}O and on ^{107}Ag with ^{20}Ne are interpreted in terms of an equilibrium model with fission competition during de-excitation of the compound nucleus. The results of the calculation are in excellent agreement with experimental results—limits to angular momenta for nuclei surviving de-excitation of the compound nucleus are thereby predicted. (auth)

47073 COUPLED-CHANNEL CALCULATIONS OF THE ENERGY DEPENDENCE OF THE (p,n) CHARGE-EXCHANGE REACTION. Hoffmann, G. W.; Coker, W. R. (Univ. of Texas, Austin). Phys. Rev. Lett.; 29: No. 4, 227-9(24 Jul 1972).

Excitation functions of the (p,n) reaction on ^{91}Zr , ^{119}Sn , ^{208}Pb , and ^{209}Bi are compared with the predictions of exact solutions of the coupled Lane equations, and it is found that the distorted-wave Born-approximation and coupled-channel predictions have essentially the same overall shape if the same optical and symmetry potentials are used. A linear energy dependence of the complex symmetry potential, suggested earlier from distorted-wave Born-approximation analyses, as well as modification of the neutron optical potential are required to describe the available data from threshold to ~ 45 MeV if global optical parameters are used. (auth)

47074 SOLUTION OF THE BETHE-GOLDSTONE EQUATION IN FINITE NUCLEI FROM N-N PHASE SHIFT DATA. Hodgson, R. J. W. (Univ. of Ottawa). Can. J. Phys.; 50: No. 9, 940-6(1 May 1972).

An approach is outlined wherein the Bethe-Goldstone wave function may be computed in a shell-model basis from a knowledge of the harmonic-oscillator matrix elements. These in turn may be derived from the N-N phase shifts by a number of different methods. The Pauli operator is treated exactly in the space of two-particle shell-model states. The method is applied to a calculation of the levels of ^{18}O and ^{18}F and results obtained using different two-body matrix elements are compared. (auth) (Canada)

47075 TRANSFER REACTIONS TO UNBOUND STATES BY THE DIFFRACTION MODEL. Sharaf, M. A. (International Centre for Theoretical Physics, Trieste, Italy). Z. Phys.; 253: No. 1, 28-37(1972).

Transfer reactions leading to unbound states are studied by means of the diffraction model. The unbound state is described by a resonance scattering wave function. The radial integrals are evaluated using some reasonable approximations. Comparison with the experimental results is very satisfactory. (auth)

47076 QUASI-CLASSICAL TREATMENT OF MULTINUCLEON TRANSFER REACTIONS. Toepffer, Christian (Univ., Frankfurt am Main). Z. Phys.; 253: No. 1, 78-88(1972).

The surface nature of the transfer reaction and the quasi-classical relative motion of the nuclei imply strong conditions on optimal scattering angles and Q values. These are evaluated using the method of stationary phases. Recoil effects are considered. The agreement with experiment is good. (3 figures, 1 table) (auth)

47077 CONSIDERATION ON DIRECT REACTION MECHANISMS IN THE UNIFIED MICROSCOPIC NUCLEAR THEORY. Hub, R.; Clement, D.; Wildermuth, K. (Univ., Tuebingen, Ger.). *Z. Phys.*; 252: No. 4, 324-48(1972).

In the framework of a unified microscopic nuclear theory the starting equations for the discussion of the different phenomenological direct reaction theories are derived. Subsequently the $^6\text{Li}(p,^3\text{He})^4\text{He}$ reaction is discussed as a realistic example to obtain a better physical insight into the different kinds of reaction mechanisms. (10 figures, 1 table) (auth)

47078 REPULSIVE CORE EFFECTS IN (t,p) STRIPPING REACTIONS. Osman, A.; Zimanyi, J. (Central Research Inst. for Physics, Budapest). *Lett. Nuovo Cim.*; 4: No. 7, 246-50 (17 Jun 1972).

The effect of the repulsive core in the nucleon-nucleon interaction on (t,p) stripping reactions was analyzed. The effect was found to be a small constant multiplicative factor relative to the nonrepulsive case. (J.W.R.)

47079 ALPHA INTENSITIES TO ROTATIONAL BANDS OF ODD-A NUCLEI. Sharma, R. C. (Sir George Williams Univ., Montreal). *Lett. Nuovo Cim.*; 4: No. 7, 257-61(17 Jun 1972).

The total probability for exciting a rotational level in a daughter nucleus with spin J, after the emission of an alpha particle with angular momentum 1, was calculated for 22 odd-A nuclei from A = 227 to 257 using the formula of Nosov. Deviations between the calculated and experimental values are discussed. (J.W.R.)

47080 ANALYSIS OF EXPERIMENTAL ENERGIES OF EXCITED STATES OF EVEN-EVEN DEFORMED NUCLEI USING ROTATIONAL FORMULAS. Abdurakhmanov, S. R.; Belen'kii, V. M. (Inst. of Nuclear Physics, Tashkent). *Izv. Akad. Nauk Uzb. SSR, Ser. Fiz.-Mat. Nauk*; No. 5, 66-9(1971). (In Russian).

The empirical formula of Y. P. Varshni (1968) was used to analyze known experimental values for energies of excited states of rotational bands from even-even deformed nuclei in the region $150 < A < 190$ and $A > 220$. The results from over 40 nuclei were compared with those of an analogous analysis using the formula of Bohr and Mottelson (1963), and it was concluded that Varshni's formula represented the experimental values more accurately. A simple criterion was obtained for excluding incorrect experimental energies of excited states with a given spin. (K.S.W.)

47081 COLLECTIVE-SINGLE-PARTICLE STATES OF DEFORMED ODD NUCLEI. Begzhanov, R. B.; Salakhitdinov, A. N.; Sharipov, Sh. (Inst. of Nuclear Physics, Tashkent). *Izv. Akad. Nauk Uzb. SSR, Ser. Fiz.-Mat. Nauk*; No. 5, 76-9(1971). (In Russian).

The same authors had previously examined excited states of odd, nonspherical atomic nuclei on the basis of a phenomenological model of the odd nucleus. New calculations have now been made on odd nuclei, with a basic state spin of $\frac{1}{2}$, and allowing for nuclear deformation. The theory yielded results for the isotopes ^{165}Ho and ^{253}Cf which agreed very well with experimental values. (K.S.W.)

47082 ROLE OF THE EXIT STATES OF A COMPOUND NUCLEUS IN THE EMISSION OF HIGH-ENERGY γ -RAYS. Brzosko, J. S.; Piotrowski, J.; Soltan, A. Jr.; Szefflinski, Z. (Univ., Warsaw). *Nucl. Phys.*; A189: No. 3, 545-55(1972).

An explanation is proposed for the appearance of the bump in γ -spectra emitted by nuclei excited to 10 MeV in the (n, γ), (p, γ) and (γ,γ') reactions. The study is based on a previous analysis which indicated that decay of the compound nucleus is responsible for this phenomenon. It has been assumed that the main mode of the compound-nucleus decay is an exit state consisting of a highly excited nucleon and a low excited core. The estimates carried out reproduce the characteristic features of the phenomenon studied. (auth)

47083 ALPHA-CLUSTER STRUCTURE AND WEAK COUPLING FEATURE IN LIGHT sd-SHELL NUCLEI. GENERAL DISCUSSION AND ^{12}F PROBLEM. Nemoto, Fumiki; Bando, Hiroharu (Kyoto Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 4, 1210-34(Apr 1972).

With the aim to investigate the core excited states in nuclei from ^{20}Ne to $^{16}\text{O}^*$, an α -cluster model is presented which can be applied to non- α -nuclei. Both alpha-clustering and overall antisymmetrization are found to be important to reproduce the weak α -hole repulsion, which is responsible for weak coupling features and other trends of the core excited states observed in these nuclei. The nucleus ^{19}F (^{18}Ne) is treated especially in detail and it is shown that the observed weak coupling behavior is understood consistently within this model. (auth)

47084 (44) AND (60) + (06) INTERACTIONS IN THE s-d SHELL. Taruishi, Koji (Tokyo Metropolitan Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 4, 1422-3(Apr 1972).

The expansion of Slater integrals in terms of Talmi integrals is discussed. Two equations are given as an example of direct integrals reducing to a sum of several integrals with different Legendre polynomial orders. Expansion coefficients are calculated for the (44) and (60) + (06) interactions. (J.W.R.)

47085 (N-Z) DEPENDENCE OF (n, γ) CROSS SECTIONS. Matsuoka, Kazuo (Osaka Univ.); Isomoto, Yukuo; Sano, Mituo; Igarasi, Sin-iti; Nishimura, Kazuaki *Progr. Theor. Phys. (Kyoto)*; 47: No. 4, 1428-30(Apr 1972).

Cross sections for (n, γ) reactions above resonance energies have been observed to decrease gradually with increasing neutron excess. Exceptions were found for magic neutron numbers and for nuclei in the transition region. Origins for these characteristics are discussed. The N-Z dependence of the neutron separation energy and the shell effect for the level density are two reasons for these characteristics. (J.W.R.)

47086 PROBLEMS OF MODERN NUCLEAR PHYSICS. Proceedings of the Second Problem Symposium on Nuclear Physics, Novosibirsk, USSR, June 12-19, 1970. Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). 544p. (In Russian and English). (CONF-700635).

Separate abstracts were prepared for 23 of the 30 complete papers included. Abstracts for the remaining 7 papers have previously appeared in NSA. (J.W.R.)

For abstracts of individual papers see: 46644, 46658, 46869, 46870, 46890, 46891, 46914, 47008, and 47087-47101.

47087 INTEGRAL EQUATIONS OF SCATTERING THEORY AND FEW-NUCLEON SYSTEMS. Faddeev, L. D. pp 5-19 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From 2. symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Two circumstances simplify the application of accurate integral equations of scattering theory to few-nucleon systems: the small radius of inter-nucleon interaction and the fact that only one value exceeds unity for two or three particles. After introducing the desired integral equations, three methods are given for solving them (Gilbert-Smith, Bateman, and the unitary polar approximation, UPA). The review continues with characteristics of three spinless bosons, three-nucleon systems, and a discussion of theoretical results by selected authors. (55 references) (K.S.W.)

47088 BOUND STATES OF FEW-NUCLEON SYSTEMS. Badalyan, A. M. (Inst. of Theoretical and Experimental Physics, Moscow). pp 20-30 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Calculations of various characteristics for bound states of few-nucleon systems were made under different conditions: pair potentials at small distances, without tensor forces, etc. A K-harmonic method was used for 3- and 4-particle systems Darevich-Green potentials, and Woods-Saxon potentials were also used in treating the data. (K.S.W.)

47089 RECENT EXPERIMENTS ON THE FOUR AND FIVE NUCLEON SYSTEMS. Meyerhof, W. E. (Stanford Univ., Calif.). pp 30-49 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Some recent experiments on four and five-nucleon systems are reviewed. Theory and experiment are compared for some of the data. (J.W.R.)

47090 HYPERSPHERICAL HARMONICS METHOD IN THE THEORY OF NUCLEAR BOUND STATES. Simonov, Yu. A. (Inst. of Theoretical and Experimental Physics, Moscow). pp 51-78 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The method of multidimensional harmonics is described, and basic equations and physical premises of the method are given with some properties of their solution. A method is given for calculating matrix elements, and for extending the calculations to wave functions with $K = K_{\min} + 1$ and $K_{\min} + 2$. The results allow an insight into the spectrum of excited states, and the addition of higher harmonics to the basic state. The problem of saturation is discussed in the framework of the present method, and the latter is related to other approaches in the theory of the nucleus. (K.S.W.)

47091 USE OF THE K-HARMONICS METHOD TO CALCULATE NUCLEAR PROPERTIES. Baz, A. I. (Inst. of Atomic Energy, Moscow). pp 79-102 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

A number of theoretical schemes resulted from the application of the K-harmonics method to the calculation of some basic properties of the nucleus. The general formula of the K-harmonics method is elucidated, and a fundamental approximation is given to assist in its practical use. The calculation of matrix elements between the K-harmonics is explained, and a shell model is introduced which was utilized in the calculation of nuclear spectra. (K.S.W.)

47092 COLLECTIVE LOW-LYING NUCLEAR STATES. Belyaev, S. T. (Inst. of Nuclear Physics, Novosibirsk, USSR). pp 129-41 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Low excited states of even-even nuclei (<2.5 to 3 MeV) generally possess a collective nature. Phenomenological and microscopic models are briefly discussed, and two models for coupling nucleons at two levels were employed in a theoretical calculation of spectra and energies of excited states for comparison with experimental curves. A method for examining (quasi) rotational excitation is divided into several steps: selection of groups of states interfering strongly among themselves but relatively weakly connected with other states, the establishment of symmetry and quantum numbers of collective states, and the solution of equations for density matrices in generalized space. (K.S.W.)

47093 COMPARISON OF VARIOUS APPROACHES TO NUCLEAR THEORY. Migdal, A. B. (Inst. of Atomic Energy, Moscow). pp 157-66 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The nucleus is first treated as a multi-body problem with excitation theory and the so-called gaseous approximation outlined as the simpler approximative approaches. The "microscopic" approach to nuclear theory, beginning with the interaction of two nucleons in a vacuum, is developed, and the basic ideas relating to the quasiparticle method are presented. (K.S.W.)

47094 CRITICAL EXPERIMENTS FOR DETERMINING PARAMETERS OF EFFECTIVE PARTICLE INTERACTION IN A NUCLEUS. Sliv, L. A. (Ioffe Inst. of Physics and Tech., Leningrad). pp 188-200 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

It is assumed that the effective interaction in the nucleus is analogous to the interaction between free nucleons, with basic members of the effective interaction being central, tensor, spin-orbital, and quadratic in L . Theoretical treatment is accorded to each of these topics. (K.S.W.)

47095 THEORY OF LOW-LYING NUCLEAR STATES IN THE SCHEME $U_{3(A-1)}$. Vanagas, V. V. (Inst. of Physics and Mathematics, Vilnius, USSR). pp 201-18 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The properties of low-lying levels of nuclei were investigated with the help of a model representation based solely on the kinematic properties of the nuclear wave function. The fundamental development of the model which was based on the scheme of the unitary group $U_{3(A-1)}$ is reviewed. Following this development the unitary scheme is related to properties of real nuclei. The operator for potential energy of the nucleus is expanded through group operators, which leads to the general properties of the spectrum of the operator for the potential energy of the nucleus. (K.S.W.)

47096 PROGRESS IN THE THEORY OF NUCLEAR MATTER AND ITS APPLICATION TO FINITE NUCLEI. Negele, J. W. (Univ. of Copenhagen). pp 218-43 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The physical foundations of nuclear matter theory are discussed. It is shown how the present knowledge of nuclear matter enables one to understand the ground-state properties of finite nuclei. Some of the criticisms of nuclear matter theory are answered. (J.W.R.)

47097 COULOMB ENERGIES AND NUCLEAR RADII. Schiffer, John P. (Argonne National Lab., Ill. Univ. of Chicago). pp 246-55 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Empirical Coulomb energy differences and charge radii were used to extract a mean-square radius for the neutron excess in a number of nuclei. The calculations were restricted to closed proton shells in order to avoid complications arising from uncertainties in configurations. The results are discussed and compared with other results. (J.W.R.)

47098 OVERLAPPING LEVELS AND GIANT RESONANCES. Shapiro, I. S. (Inst. of Theoretical and Experimental Physics, Moscow). pp 273-84 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The concept of the input state is formulated, not being based on model representation. The phenomenological theory of giant nuclear resonances is subsequently developed, and related to the dynamics of the formation of a composite nucleus. (tr-auth)

47099 SPECTROSCOPIC STUDIES WITH ISOBARIC ANALOG RESONANCES. von Brentano, Peter (Max-Planck-Institut fuer Kernphysik, Heidelberg). pp 285-318 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The spectroscopic information obtainable from isobaric analog resonances is described. Methods for analyzing experimental data on these resonances are reviewed. The S -matrix for isobaric analog resonances is discussed in detail. Some experimental data is surveyed. (J.W.R.)

47100 DESCRIPTION OF THE INTERACTION OF HIGH ENERGY PARTICLES WITH NUCLEI. Feshbach, H. (Massachusetts Inst. of Tech., Cambridge). pp 379-91 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

Some quantitative reasons for errors in the diffraction model due to correlations are given. An alternate formalism for the description of the scattering of high energy particles by nuclei is offered. (J.W.R.)

47101 MECHANISM OF DIRECT PROCESSES AT HIGH ENERGIES. Kolybasov, V. M. (Inst. of Theoretical and Experimental Physics, Moscow). pp 391-410 of Problems of Modern Nuclear Physics. /Kolybasov, V. M. (ed.). Moscow; Izdatel'stvo Nauka (1971). (In Russian).

From second symposium on the problems of nuclear physics; Novosibirsk, USSR (12 Jun 1970).

The wide variety of nuclear reactions are discussed with respect to possible mechanisms: inelastic collisions $A(x,x')A^*$, knock-out reactions $A(x,x')B$, the polar mechanism $A + x \rightarrow B + y + z$, the decay reaction $A \rightarrow B + i$, etc. The discussion touches upon polarization effects, verification of isotopic ratios, differential cross sections, and angular distribution of proton paths in reactions emitting protons. (K.S.W.)

47102 THEORETICAL INVESTIGATION OF COLLECTIVE EXCITATIONS IN MEDIUM AND HEAVY NUCLEI. Das, Tapan Kumar. Philadelphia; Univ. of Pennsylvania (1971). 163p. University Microfilms Order No. 72-17,342.

Thesis.

The collective excitations in medium and heavy doubly even nuclei have been investigated by examining a few phenomenological models. Attempts have also been made to understand the nuclear rotations and vibrations on a uniform footing by providing theoretical foundations of some of the phenomenological models. A theoretical foundation as well as a higher order generalization for the highly successful Variable Moment of Inertia (VMI) model have been provided utilizing the variation after projection (VAP) method. The derivation gives a theoretical justification for the applicability of the model outside the strongly deformed region. By analogy, a variable frequency (VF) model is obtained for one dimensional vibrations. Extending the ideas, a multi-band VMI-VF model has been proposed and successfully applied to the available

data. The multi-band VMI model has been shown to reduce to the form of the original VMI model for the excitation energy with respect to the band head of a given band. This indicates that the effects of band mixing are already incorporated in the original VMI model. The ground state bands have also been investigated in terms of the Harris or generalized cranking model. It has been proved that this model is exactly equivalent to the generalized VMI model, when all orders are taken. As an independent foundation of the model, a derivation of the cranking variational equation (CVE) for a two dimensional case was performed. Next an approach to the problem starting from the vibrational end is made by proposing a model of anharmonic vibrations (AVM). The AVM has been applied to the ground state bands. This gives a power series in angular momentum quantum number for the excitation energy of members of the band. It has been shown that this series converges in the rotational limit, when successive pairs of terms of the series are considered. This model has been applied to the available data for the excitation energies and $B(E2)$ values of members of the ground state band. Finally an attempt has been made to generalize the AVM description to include excited bands. A complete set of computing operators has been found for complete classification of the states in the vibrational description. To apply the generalization to the excited rotational bands, one needs to discover a correlation between the two sets of quantum numbers specifying the vibrational and rotational descriptions. An attempt in this direction met with no success indicating that no such simple correlation exists. (Diss. Abstr. Int., B)

47103 NUCLEAR THREE-BODY PROBLEM AND GENERALIZED SPHERICAL HARMONICS. Schneider, Thomas Richard.

Philadelphia; Univ. of Pennsylvania (1971). 193p. University Microfilms Order No. 72-17,421.

Thesis.

A scheme for solving the three-body bound state problem by expanding the wavefunction in a complete set of generalized spherical harmonics is presented. This reduces the three-body Schroedinger equation to an infinite set of coupled ordinary differential equations in one variable. Restricting the problem to a finite subspace the convergence properties of this approximation as the number of states increases is dealt with. These states are labeled by the eigenvalues of the operator Λ^2 . The corresponding classical operator is zero only when all three particles are at the same point in space at the same time, and thus is a measure of the "compactness" of the system. The convergence problems associated with the expansion are related to the Fourier problem. A theorem about the convergence rate of generalized Fourier series is proved. Combining this with theorems on the convergence of the approximate energies and wavefunctions gives one the asymptotic convergence rate of the binding energies. Examining the properties of the Schrödinger equation, one can relate the convergence rate to the properties of the two-body potential. The approximate energies converge as inverse powers of the eigenvalues of Λ^2 , with the power depending upon the exact form of the two-body potential. Some calculations of the binding energies of three-boson systems are presented. These calculations are performed, with potentials of both Yukawa and square well form, by numerically integrating the coupled differential equations. Even potentials with repulsive Yukawa cores can be easily solved using generalized spherical harmonics. (Diss. Abstr. Int., B)

PHYSICS (PLASMA AND THERMONUCLEAR)

47104 PLASMA: THE FOURTH STATE OF MATTER. Frank-Kamenetskii, D. A. New York; Plenum Publishing Corporation (1972). 165p. \$12.50.

Translation of *Plazma: Chetvertoe Sostoyanie Veshchestva* by Joseph Norwood Jr.

The fundamental concepts of plasma physics are discussed without relying on intricate mathematics. The magnetohydrodynamics, as well as the physical kinetics of plasma, are described. Plasma oscillations, instability, compression, and confinement by magnetic fields are discussed in a very elementary manner. (M.O.W.)

Confinement and Heating

47105 VISCOSITY AND THERMAL CONDUCTIVITY EFFECTS ON THE CONTAINMENT OF A TOROIDALLY CONFINED RESISTIVE PLASMA WITH FLOW. Grimm, R. C.; Johnson, J. L. (United Kingdom Atomic Energy Authority, Abingdon, Eng.). *Plasma Phys.*; 14: No. 6, 617-34 (Jun 1972).

Plasma containment in toroidal systems is studied using a fluid model containing the effects of plasma inertia, resistivity (electron-ion momentum transfer), viscosity (ion-ion collisions), and thermal conductivity. Asymptotic expansion techniques are used to separate the problem into two distinct parts—determination of steady-state configurations where time variations occur only on a diffusion time scale, and investigation of the stability of these equilibria. The previously known steady-state solutions are shown to be stable. However, terms in the stress tensor associated with the curvature of the magnetic field lines, which introduce parallel viscosity into the components of the momentum equation perpendicular to the field, are shown to restrict the possible solutions. (auth)

47106 NONADIABATIC ENERGY LIMIT VERSUS MIRROR RATIO IN MAGNETIC-WELL GEOMETRY. Foote, James H. (Univ. of California, Livermore). *Plasma Phys.*; 14: No. 5, 543-52 (May 1972).

The energy at which ions trapped in a static quadrupole magnetic well begin to exhibit nonadiabatic behavior in their single-particle motion was determined for mirror ratios from 1.4 to 8.1. The analyzed data are derived partly from digital-computer orbit calculations and partly from experiment. The investigated particles either reflect at closed contours of constant magnetic-field magnitude or are constrained by the field lines to reflect in regions where the contours approximate sections of closed magnetic-field surfaces. The calculated trajectories pass through the central region of the well and represent the first trapped particles in a closed-contour system to become nonadiabatic as the energy is increased. (UK)

47107 COLLISIONLESS HEATING OF A PLASMA BY MEANS OF OSCILLATING EXTERNAL CURRENTS. Lashmore-Davies, C. N. (United Kingdom Atomic Energy Authority, Abingdon, Eng.). *Plasma Phys.*; 14: No. 4, 357-66 (Apr 1972).

The transfer of power between an external circuit and a collisionless plasma is described. A very simple model is used that enables heating for arbitrary values of the frequency to be analyzed. Transit-time magnetic pumping and ion-cyclotron heating are described in detail. The simple model used enables the mechanism of magnetic pumping to be described much more precisely than has been done previously. (UK)

47108 HEATING OF TOROIDAL PLASMAS BY NEUTRAL INJECTION. Stix, Thomas H. (Princeton Univ., N. J.). *Plasma Phys.*; 14: No. 4, 367-84 (Apr 1972).

A brief review of the physics of ion acceleration, charge exchange and ionization, trajectories for fast ions in toroidal magnetic fields, and fast-ion thermalization is presented. The injection of fast atoms is found to be a highly competitive method both for heating present-day experimental toroidal plasmas and for bringing full-scale toroidal CTR plasmas to low-density ignition. (auth)

47109 PLASMA HEATING NEAR THE ION-CYCLOTRON FREQUENCY AND ITS HARMONICS. Clark, Reginald Wayne. Austin, Tex.; Univ. of Texas (1971). 134p. University Microfilms Order No. 72-19,570.

Thesis.

Additions are made to the theory of the quasi-ion Bernstein modes that propagate for wave frequencies near the harmonics of the ion cyclotron frequency, and experience ion cyclotron damping as the harmonic resonances are approached. The effects of

collisions and electron Landau damping are included, and group-velocity calculations are performed. When the effects of collisions are included, the theory predicts strong collisional damping in high-density, low-temperature collision-dominated plasmas, whereas the original theory predicted wave propagation in the absence of collisions. The inclusion of electron Landau damping effects in the theory modifies the propagating bands so that the waves propagate undamped only in narrow bands near cutoff, i.e., where the component of the propagation vector parallel to the magnetic field is small. The ion acoustic-electrostatic ion cyclotron (IA-EIC) wave also propagates near the fundamental and in bands near harmonics of the ion cyclotron frequency but requires for weak ion and electron Landau damping that $T_e > T_i$, which is an undesirable temperature condition for plasma heating. However, due to nonlinear effects, the constraints imposed by electron Landau damping considerations may not be important for large-amplitude waves. The constraints imposed by ion Landau damping, however, remain valid over much longer times and distances than those imposed by the electrons and hence represent much stronger constraints. An experimental investigation of harmonic ion cyclotron wave propagation in high-density, low-temperature collisionally dominated plasma indicates that the waves are very heavily damped. These results indicate that a high-density plasma must be preheated to a temperature where collisions become unimportant before harmonic ion cyclotron waves can be used for plasma heating. Subsequent experimental investigation of heating in the turbulently preheated plasma in the Cornell University THM-2 plasma machine indicates plasma heating at densities as high as $\sim 7 \times 10^{18}/\text{m}^3$. Ion energy measurements indicate ion temperatures the order of 100 to 110 eV and a nonMaxwellian energy distribution with a higher-energy tail, while diamagnetic loop measurements indicate that the sum of the ion and electron temperatures is roughly between 100 and 200 eV. Interferometer measurements indicate that the plasma is contained for the duration of the 600- μsec rf pulse. Fluctuations were observed that are attributed to particles bunching and executing longitudinal oscillatory motion between the magnetic mirrors of the THM-2. These observations raise the possibility that the heating resulted from the nonlocal resonance that may arise when particles oscillate in an inhomogeneous magnetic field. The observed heating was proportional to the net power input, and no saturation effects were seen. The overall energy-transfer efficiencies are estimated to be at least 20 to 40%. (Diss. Abstr. Int., B)

47110 PLASMA INJECTION AND COLLISION PHENOMENA IN AN OCTUPOLE MAGNETIC FIELD. Ligon, Jerry Lloyd. Austin, Tex.; Univ. of Texas (1971). 182p. University Microfilms Order No. 72-19,624.

Thesis.

This work investigates the transverse plasma injection into a linear octupole magnetic field by using two colliding plasmas for injection instead of the usual single plasma. The two plasmas enter the octupole field from opposite sides and collide at selected locations in the field region. The theory that applies to plasma injection and collision in a transverse magnetic field is presented, and the possible effects of Coulomb collisions, collective instabilities, and polarization electric fields are discussed. Expected results from a typical experiment are also presented. A computer simulation of two-plasma injection into a magnetic field similar to the octupole field is carried out. The program is a modification of a program written for analysis of single gun injection. The results of the program agree with the experimental results in describing some aspects of the injection and interaction. (TSS) (Diss. Abstr. Int., B)

47111 PARALLEL MAGNETIC MULTIPOLE CONFINEMENT OF A MAGNETIC-FIELD-FREE PLASMA. Limpaechee, Rudolf. Los Angeles; Univ. of California (1972). 105p. University Microfilms Order No. 72-20,454.

Thesis.

A new multimagnetic mirror plasma device is described. The walls of this cylindrical 86-liter "box" are lined with 1252 ($1/2'' \times 1/2'' \times 1.5''$) ALNICO V permanent magnets to form a series of parallel broken-line cusp geometry. Since the magnetic field falls off exponentially ($\delta \approx 1.2$ cm) the interior is essentially magnetic field free, even though B_{max} (at the walls) is 1800 gauss. The argon, helium, hydrogen, or deuterium plasma is produced by electron emission from 12 short tungsten filaments with an emission current of $0 < I_{\text{ee}} \leq 5$ A over a neutral density range of $5 \times 10^{-6} \leq n_0 \leq 2 \times 10^{-3}$ torr and a plasma density of over 2×10^{11} cm^{-3} torr. For argon and helium the empirical relation $n \propto$

$n_0 = 3.3 \pm 0.05 \times 10^{18}$ is found over the neutral density and emission current range specified. A plasma nonuniformity of less than 1% is recorded in the magnetic field-free region with a plasma noise level of $\delta n/n = 3 \times 10^{-4}$. A steady-state plasma lifetime of over 20 msec is obtained. The system ionization efficiency is particularly impressive. With a total input power of 500 watts (heater and emission) the device has a volume of 10^{13} Debye spheres, where 4×10^{13} particles per watt of input power are maintained in the steady-state condition. At 5×10^{-6} torr, a fractional ionization of 35% is obtained. Strong evidence exists that the ions are contained magnetically as well as electrostatically. The electrostatic fields, which are maintained by the plasma electron and emission electron interaction with the magnetic multipole field, drastically reduce the ion cusp losses. With this multipole structure, high density quiescent collisionless plasmas can be produced with any size and any geometry. An obvious extension is the application of a uniform magnetic field for high-beta experiments. A new system is proposed to contain high energetic ions by replacing internal emission electron sources by electron beams. These electrons have to be injected through the central cusp regions of the electromagnets. (Diss. Abstr. Int., B)

47112 TURBULENT HEATING OF THE IONS IN A TOROIDAL DISCHARGE. Mah, Seck Qui. Saskatoon, Saskatchewan; Univ. of Saskatchewan (1971). 173p. University Microfilms Order No. 72-18,654. Thesis.

Studies of plasma heating were carried out in a toroidal, high-voltage discharge. The argon plasma of density $\sim 10^{18}/\text{m}^3$ is formed initially by rf preionization at low pressure (≈ 0.3 mtorr). The electric field (~ 5 kV/m) of the main discharge is applied parallel to a steady, toroidal magnetic field (≤ 0.2 Wb/m²). Rapid ion heating, to $\sim 10^5$ K, is observed in the time interval from 1 to 2 μsec after initiation of the electric field. The ion temperature is determined from the Doppler broadening of the Ar II 4806-Å emission using a time-resolved, single-channel Fabry-Perot interferometer with piezoelectric scanning. Ion acoustic waves induced by the electron current during the discharge are shown to be the probable cause of the rapid heating. This conclusion was reached by measuring, in addition to the ion temperature, the electron temperature, electron density, plasma current and electric field, all as a function of time. The measurements show that, when rapid ion heating is observed, the electron drift velocity is above the critical value for growth of ion acoustic waves and, more significant, the rapid heating ceases when the mean electron velocity falls below this critical value. Anomalous low electrical conductivity is observed in the interval when the ions are rapidly heated. The observed conductivity is about two orders of magnitude lower than classical, indicating a large collision frequency between electrons and the waves excited in the plasma. Detailed comparison with the conductivities predicted by theories of plasma turbulence due to the ion acoustic instability show that these cannot provide a satisfactory explanation. Partial agreement is found between experiment and the theory of Dupree. At the onset of the high-voltage discharge, electrons are heated rapidly, up to several times 10^5 K, within a fraction of a microsecond. Only a small portion of the input energy is converted into particle drift; the rest goes into electron heating. Energy accounting is not possible at later times in the discharge, where rapid electron cooling due to some unknown process is observed. (Diss. Abstr. Int., B)

47113 ELECTROSTATIC CONTAINMENT IN FUSION RE-ACTORS. Hirsch, Robert L. (to International Telephone and Telegraph Corp.). U. S. Patent 3,664,920. 23 May 1972. Filed 21 Jun 1968.

Apparatus is described for producing controlled nuclear-fusion reactions. A generally circular cathode has concentrically positioned therein a generally circular anode having multiple openings for the free flow of electrons. The anode encloses a volume free of tangible structure. An electron-emissive electrode in the interelectrode space between the anode and cathode emits electrons which penetrate the anode and enter the central volume, establishing a negative space charge. Ions of a fusion reacting gas are injected at a location within the volume, at a potential less than that applied to the anode, and are focused toward a common cross-over point in the central portion thereof. The magnitude of the negative space charge as well as the energies of the ions introduced into the volume are so controlled that collisions of the ions at or near the cross-over point result in the fusing thereof. Electrons introduced in the interelectrode space traverse the cathodic space, thereby making a number of trips through the anode. In this invention, the anode structure is effectively shielded such that electron interception thereby with consequent high energy electron losses can be made to approach zero. This is accomplished by creating a magnetic field around

the portions of the anode which define the openings, this field serving to shield the conducting portions thereof. (Official Gazette)

47114 PLASMA CONFINEMENT BY EXTERNAL RF SIGNAL. Chiplonkar, V. T.; Sapre, A. V. (Inst. of Science, Bombay). Curr. Sci. (India); 41: No. 15, 559-60 (5 Aug 1972).

Observations are reported on the current density distribution on the anode surface of a cylindrical gas discharge plasma when an rf signal is applied to a pair of ring electrodes placed at a distance of 1.3 cm from the anode on the inner and outer walls of the discharge tube. The data were used to obtain a quantitative estimate for the confinement effect. (M.O.W.)

47115 GENERALIZED NONLINEAR EXPRESSION FOR THE PLASMA DISPLACEMENT IN HARMONICA 2 AND TOKAMAK. Misguich, J. H.; Mercier, C. (EURATOM-CEA, Fontenay-aux-Roses, France). Plasma Phys.; 14: No. 4, 418-20 (Apr 1972).

An expression for the displacement of plasma in a toroidal device is modified by taking into account nonlinear effects, varying plasma radius, and nonvanishing surface currents, and related to recent results obtained with the Harmonica 2 device. (UK)

Diagnostics

47116 (IPP-IV-12) SIMULTANEOUS SPECTROSCOPIC DETERMINATION OF ELECTRON DENSITY AND TEMPERATURE AS A PROOF OF RELAXATION EFFECTS IN A STREAMING ARGON-POTASSIUM-PLASMA. Dodel, G. (Max-Planck-Institut fuer Plasmaphysik, Garching (West Germany)). Jun 1971. 128p. (In German). Dep. NTIS (U. S. Sales Only).

Simultaneous measurements of electron temperature and density distributions prove that significant relaxation effects of the electron density occur in an MHD-generator-type argon-potassium plasma streaming through a channel with segmented-electrode array. Both parameters are determined spectroscopically, the electron temperature by line reversal of the potassium resonance doublet $4S_{1/2}-4P_{3/2}$, and the electron density by absolute intensity measurement of the $4P_{3/2}-10D_{5/2}$ transition. The applicability of these two spectroscopic methods, which is by no way trivial in a transient plasma, is investigated on the basis of the quasi-steady-state theory of the relaxation process by Bates et al. The principal problem of spectroscopy in transient plasmas is substantiated in terms of this theory in a rather general way. The numerical calculations of the theory are based on an eighteen-level model of the potassium atom, the semiclassical THOMSON-GRYZINSKI collision cross sections being used. They yield population densities of the excited levels, as well as ionization and recombination coefficients as functions of the electron density and temperature. Thus, density and temperature ranges can be specified in which the spectroscopic methods mentioned above are justified. Furthermore, the specific relaxation problems occurring in the experiment can be calculated and compared to the measurements. Good qualitative agreement is found. Quantitatively, the measured electron density relaxation occurs faster than is predicted by the calculations. Finally, the numerical calculations yield the equilibrium electron density as a function of the electron temperature. Due to radiation escape, this equilibrium density appreciably deviates from the Saha density. In the experiment, however, such a significant deviation could not be verified. The discrepancy between calculated and measured equilibrium electron densities can be reduced by taking wall radiation into account. This seems to indicate that wall radiation plays a role in establishing equilibrium. (auth)

47117 SPECTRAL INTENSITY MEASUREMENTS FROM HIGH-PRESSURE NITROGEN PLASMAS. Cooper, David M. (Ames Research Center, Moffett Field, Calif.). J. Quant. Spectrosc. Radiative Transfer; 12: No. 8, 1175-89 (Aug 1972).

Spectral measurements of the equilibrium radiation from nitrogen at high pressures were obtained. Experiments were performed in a ballistic range and an explosively driven shock tube. The tests covered a temperature range from 9000 to 17,200 K and a pressure range from 1.3 to 63.0 atm. Averaged electronic absorption f-numbers for $N_2^+B^2\Sigma_u^+ - x^2\Sigma_g^+$, $N_2C^3\Pi_u - B^3\Pi_g$, and $N_2B^3\Pi_g - A^3\Sigma_u^+$ bands are reported. N^+ continuum radiation was not observed. (auth)

47118 REFLECTION OF MICROWAVE THROUGH LABORATORY PLASMA. Gwal, A. K.; Singh, K. P.; Shukla, P. K.; Misra, K. D. (Banaras Hindu Univ., India). Int. J. Electron.; 33: No. 1, 91-5 (Jul 1972).

Propagation of microwaves through the laboratory plasma was studied. Assuming the linear electron density variation, an expression for the envelope of the reflected wave was obtained. It was observed that the propagating wave through the microwave

reflection probe is distorted when reflected back from the other end of the probe having plasma boundary. By analyzing the distorted signal thus received, the plasma parameter is estimated. (UK)

47119 TECHNIQUE FOR RECORDING LANGMUIR PROBE CHARACTERISTICS IN AFTERGLOW PLASMAS. Dean, A. G.; Smith, D.; Plumb, I. C. (Univ. of Birmingham, Eng.). *J. Phys.*, E (London); 5: No. 8, 776-8 (Aug 1972).

A technique is described for obtaining Langmuir probe current-voltage characteristics in afterglow plasmas under conditions where the ratio of the signal-to-noise levels is small. This has enabled accurate measurement to be taken of very low electron temperatures and electron densities under conditions for which it is essential to use very small probes in order to ensure that significant depletion of the plasma does not occur. The technique involves the successive accumulation of data from repetitive afterglows using multichannel scaling equipment. Sample data are presented to illustrate the value of the technique. (auth)

47120 METHOD FOR DETERMINING THE DIRECTION OF THE MAGNETIC FIELD IN A TOKAMAK PLASMA BY LASER LIGHT SCATTERING. Sheffield, John (Univ. of Texas, Austin). *Plasma Phys.*; 14: No. 4, 385-95 (Apr 1972).

It is shown that laser light passing through a magnetized plasma undergoes significant modulation when the scattering wave number is within about one degree of the magnetic field direction. By using a Fabry-Perot etalon of free spectral frequency range equal to the electron cyclotron frequency, the resulting modulation can be detected and the effect of the magnetic field determined. (UK)

47121 HIGH-SPEED PHOTOGRAPHY OF A PLASMA FOCUS. Lee, S.; Chen, Y. H.; Chow, S. P.; (and others) (Univ. of Malaya, Kuala Lumpur). *Int. J. Electron.*; 33: No. 1, 85-90 (Jul 1972).

A coaxial plasma gun is operated with an outer glass cylinder in place of the usual copper cylinder. This enables streak and framing photographs to be readily taken of the shock-propagation region, the transition to the focus region, and the focus itself. (auth)

47122 ELECTRON TEMPERATURE DETERMINATION FROM THE RATE OF IONIZATION. Mah, S. Q.; Skarsgard, H. M.; Strilchuk, A. R. (Univ. of Saskatchewan, Saskatoon). *Plasma Phys.*; 14: No. 4, 422-5 (Apr 1972).

The use of a technique to measure the electron temperature of a low-pressure, high-electric-field, toroidal discharge is reported. The method utilizes the observation of an increase in plasma density, and relates ionizing rate to electron energy. (UK)

47123 CALCULATION AND MEASUREMENT OF THE CHARACTERISTIC FLOW PARAMETERS OF A STATIONARY SUPERSONIC PLASMA FLOW IN A MAGNETIC DE LAVAL NOZZLE. Bitter, M.; Bartoli, C. (European Space Research Inst., Frascati, Italy). *Plasma Phys.*; 14: No. 6, 575-82 (Jun 1972).

A Q device modified into a magnetic de Laval nozzle is used to produce a supersonic cesium plasma. It is shown that the plasma in such an experimental setup can be described by a simple adiabatic theory. The characteristic parameters of a stationary supersonic flow are calculated both for collisional and the collisionless case. The ion saturation currents measured with Langmuir probes at various positions along the magnetic nozzle are found to be in agreement with the theoretically expected thermal ion fluxes. (auth)

47124 OPTIMIZATION OF MAGNETIC PROBE MEASUREMENTS IN TRANSIENT PLASMAS. Pachner, Jaroslav Jr. Vancouver, B. C.; Univ. of British Columbia (1971). National Library of Canada, Ottawa.

Thesis.

A significant improvement in magnetic probe measurements was achieved by developing a 3-coil magnetic probe that partially corrects for the boundary error, i.e., for the error caused by an exclusion of the plasma current from the space occupied by the probe. The spatial resolution l of the three-coil probe is roughly one-half of the probe radius, a . For current distributions that vary slowly with distance (scale length, λ) it is shown that the fractional error in the magnetic field is $0.2 (l/\lambda)^2$. For a conventional probe, the error is at least four times as large. Also, it was shown that spurious signals arising from poor probe geometry (and which often obscure the signals produced by the measured magnetic fields themselves) can be eliminated by making use of the symmetry of the discharge fields. Measurements were made on a z-pinch discharge that confirm the claimed 3-coil probe performance by revealing a "fine" structure of the current distribution in helium in the filling pressure range 0.5 to 4 torr. The measurements are presented in the form of a catalogue of the

spatial distribution of magnetic field and the current density for a z-pinch discharge in He and Ar. Using the probe measurements, a qualitative model of the collapse of the current sheet for 4 torr in He is developed that differs from previous models. (Diss. Abstr. Int., B)

47125 STUDY OF OPTICAL-SCATTERING METHODS IN LABORATORY PLASMA DIAGNOSTICS. Phipps, Claude Raymond Jr. Stanford, Calif.; Stanford Univ. (1972). 127p. University Microfilms Order No. 72-20,725.

Thesis.

Electron velocity distributions are deduced along axes parallel and perpendicular to the magnetic field in a pulsed, linear Penning discharge in hydrogen by means of a laser Thomson-scattering experiment. Results obtained are numerical averages of many individual measurements made at specific spacetime points in the plasma evolution, and take advantage of the high level of reproducibility of density and velocity parameters possible in this type of discharge. Observation solid angles are made quite small to maximize resolution between the two distributions. Because of the high resolution in k space and the relatively low maximum electron density $2 \times 10^{15} \text{ cm}^{-3}$, special techniques were required to obtain measurable scattering signals. These techniques are discussed and experimental results are presented. The Thomson scattering measurements show identical behavior of the longitudinal and azimuthal distributions, within experimental error. For magnetic fields below the critical value for onset of strong instability in the Penning discharge, the results show impressively good fit to a calculated Gaussian distribution, for electron energies up to 6 times the thermal energy. However, measurements above this critical field show a distinct excess of high-energy electrons which cannot be accounted for by a single Gaussian, within experimental error. Equipment-scattered light levels are the lowest ever attained in an experiment of this kind. (Diss. Abstr. Int., B)

47126 INTERPRETATION OF SIGNALS FROM A CYLINDRICAL LANGMUIR PROBE IN A STREAMING LASER-PRODUCED PLASMA. Segall, Stephen Barrett. College Park, Md.; Univ. of Maryland (1972). 235p. University Microfilms Order No. 72-20,261.

Thesis.

The applicability of the cylindrical Langmuir probe to the diagnosis of the properties of streaming, laser-produced plasmas is investigated. The regime in which probe measurements can be made is defined, and the relation of the electrostatic probe to other means of plasma diagnostics for laser-produced plasmas is explained. A theory to explain the current characteristic for a probe in a laser-produced plasma is developed. First, the theory for a probe in a stationary plasma is analyzed to develop a stationary analog to the flowing-plasma case. In the process, it is found that the general theory for a cylindrical probe in a stationary plasma, as developed by Bernstein and Rabinowitz and Laframboise, can be considerably simplified both theoretically and computationally. This simplified general theory is then used to reexamine several fundamental concepts of probe theory and the behavior of the probe for various changes of plasmas and probe parameters. Relations applicable to the case of a probe in a high-velocity flowing plasma are developed and used to modify the Langmuir theory for a probe in a collisionless flowing plasma. Experimental procedures for the use of cylindrical Langmuir probes to diagnose streaming, laser-produced plasmas are described, and experimental current-voltage characteristic curves are analyzed using the theory for a probe in a flowing plasma which has been developed. Density and temperature values are obtained for copper and aluminum plasmas at various times during the flow. Reasons for deviations from theoretical behavior are discussed. (Diss. Abstr. Int., B)

47127 TIME-RESOLVED DIAGNOSTIC METHOD FOR HYDROGEN PLASMAS. Edwards, David F. (Colorado State Univ., Ft. Collins); Bacon, M. E.; Litvak, M. M. *Int. J. Electron.*; 33: No. 1, 105-12 (Jul 1972).

A time-resolved plasma diagnostic method is described as it applies to laser-produced sparks in hydrogen gas. Electron densities and temperatures in the ranges of 10^{19} to 10^{17} cm^{-3} and 10^6 to 10^4 K , respectively, can be determined with nanosecond time resolution. The diagnostic method is not necessarily restricted to hydrogen plasma and suggestions are given as to possible extensions to other types of plasma. (auth)

47128 PLASMA-SAMPLING DIAGNOSTICS USING ION KINETIC ENERGY DISTRIBUTIONS. Seguin, J. G.; Dugan, C. H.; Goodings, J. M. (York Univ., Toronto). *J. Mass Spectrom. Ion Phys.*; 9: No. 2, 203-13 (Jul 1972).

Kinetic energy distributions of ions extracted from a microwave discharge in nitrogen are used to diagnose the influence of the

plasma sheath on the sampling process. Nitrogen plasmas having $n_e \approx 10^9$ to 10^{10} cm $^{-3}$ and $kT_e \approx 1$ eV were examined over the pressure range 0.02 $_2$ to 0.42 torr. Over this range N_2^+ and N^+ are the major ion species with N_2^+ predominant by approximately one order of magnitude. The ions are sampled from the bulk phase through the plasma sheath which forms over a tiny platinum leak. The leak itself acts as a circular planar Langmuir probe. After passage through the leak, the ions are analyzed for kinetic energy by a series of planar retarding grids and then mass-analyzed using a quadrupole mass spectrometer. Analysis of the ion kinetic energy distributions indicates that a model of the plasma sheath dominated by mobility considerations is applicable for pressures >0.42 torr. On the other hand, a collisionless free fall sheath model is applicable at pressures $<0.02_2$ torr. However, the sampling is complicated by the occurrence of ion-molecule reactions in the sheath. The evidence suggests that the cross section for dissociative charge transfer $N_2^+ + N_2 = N^+ + N + N_2$ exceeds that for symmetric charge transfer $N_2^+ + N_2 = N_2 + N_2^+$ at low pressure; the reaction $N_2^+ + N_2 = N^+ + N_3$, if present at all, does not play a major role. (auth)

47129 SPECTROGRAPHIC MEASUREMENT OF THE TEMPERATURE OF A PLASMA ANALYZED BY STATISTICAL METHODS. Murino, Pasquale (Univ., Naples). Aerotec.-Messili Spazio; 50: No. 6, 352-9 (Dec 1971). (In Italian). Thesis.

The local temperature of an ionized gas can be indirectly determined by measuring the emitted radiation intensity per unit volume. For an axisymmetric and optically thin plasma the lateral intensity $I(x)$ of the emitted radiations can be easily measured through spectroscopic methods; therefore, it is necessary to solve Abel's integral equation to get the radiation intensity per unit volume. The numerical evaluation of Abel's integral was obtained through a method based on the approximation of $I(x)$ with a sum of polynomials orthogonal with respect to a suitable weight function. The method seems to be interesting as it is not very sensitive to the random errors of the measurements while it shows a fairly good numerical accuracy. Spectrographic measurements were made of the radiations emitted by a laboratory plasma in several experimental conditions. Abel's inversion was done by using this method and the results obtained were satisfactory. (auth)

Kinetics (General)

Refer also to abstracts: 46361 and 46425.

47130 PLASMA CURRENT CONDUCTION TO A COLD ELECTRODE. Kern, Robert Allyn. Cleveland; Case Western Reserve Univ. (1972). 180p. University Microfilms Order No. 72-18,698.

Thesis.

The sheath region adjacent to a cold electrode in a partially ionized gas is investigated in detail for the case of the Debye length smaller than an ion-atom mean free path. From the physics of ion-atom collisions, two limiting collision models are developed for the case of argon. Utilizing these models, a procedure is developed for joining the kinetic theory and continuum descriptions at the point where the two expressions give the same value for the electron-temperature gradient. This matching procedure effectively utilizes a generalized Bohm criterion as a boundary condition. This matching procedure is compared with sheath models for local variations in physical quantities and for the effects on the solution for an ionization dominated flowing plasma under shock tube conditions. This latter comparison is chosen because it provides a good test of the sheath models, it is typical of MHD applications, and the results can be compared to shock-tube experiments. It is shown that the various sheath models differ considerably in their description of the region adjacent to the electrode. Nevertheless, in some cases the ion current drawn to the electrode is insensitive to these discrepancies because there is little ionization near the electrode. By comparison with experiments and estimates of convection it is determined that for the lower range of shock tube pressures for argon, convection is the dominant current-producing mechanism. At the higher range of shock tube pressures, ionization is probably dominant. Although the matching procedure is not strictly valid at the higher shock tube pressures, there are ionization-dominated applications where the matching procedure would be valid. (TSS) (Diss. Abstr. Int., B)

47131 OBSERVATIONS OF PRECURSOR IONIZATION IN ELECTROMAGNETIC TUBES. McRae, Thomas G. Gainesville, Fla.; Univ. of Florida (1971). 65p. University Microfilms Order No. 72-16,631.

Thesis.

An experiment was designed which measures the photoioniza-

tion of a room-temperature gas by the radiation emitted from a hot plasma. The radiation source is a Mach 5 shock wave in xenon which is separated from an absorption chamber by a lithium-fluoride window. The shock was produced by a curved electromagnetic T-tube. The emission just prior to shock-wave reflection was collimated and projected through the absorption chamber to an electrostatic double probe 232 cm away. Electron number densities were recorded in Xe, Ar, Kr, and residual gas at various absorption chamber pressures. A qualitative test of the photoelectric effect on the system boundaries was conducted and both axial and transverse shock tube spectra were recorded. The results of this experiment show that, in electromagnetically driven shock tubes, precursor ionization is the result primarily of radiation, not of ionizing potential waves as recently hypothesized. Furthermore, this ionization is produced by wavelengths greater than 1200 Å but less than 3000 Å and it is predominantly the impurities in the system which become ionized. The photoelectric effect on the system walls was present but found to be negligible at normal shock-tube pressures. (Diss. Abstr. Int., B)

47132 PRELIMINARY RESULTS OBTAINED FROM THE STUDY OF AN ARGON PLASMA CONTINUUM IN THE NEAR ULTRAVIOLET AND VISIBLE RANGES AT HIGH PRESSURES. Goldbach, Claudine; Nollez, Gerard (Centre National de la Recherche Scientifique, Paris). Compt. Rend., Ser. B; 274: No. 16, 970-3 (17 Apr 1972). (In French).

The continuum produced by thermal plasmas in the 10 ^{4} K temperature region is essentially due to recombination, and may be theoretically described using the Biberman-Schluter $\xi(\lambda, T)$ function. The values of $\xi(\lambda, T)$ calculated by Schluter explain most of the experimental results; very little work has been done in comparing theory and experiment for pressures higher than atmospheric pressure. Measurement of the K_λ absorption coefficient of the plasma, produced in a variable-length stabilized arc, allowed the $\xi(\lambda, T)$ function to be determined in the $1 \leq p(\text{atm}) \leq 30$ pressure range. (France)

47133 ELECTRON ENERGY DISTRIBUTION AND STRIATIONS IN THE LANGMUIR-PARADOX REGION. Kondrat'ev, A. A.; Milenin, V. M.; Tsendin, L. D. (Leningrad State Univ.). Sov. Phys.-Tech. Phys. (Engl. Transl.); 17: No. 1, 71-5 (Jul 1972).

Translated from Zh. Tekh. Fiz.; 42: No. 1, 94-9 (Jan 1972).

A quantitative study is made of striations in a helium discharge under the conditions of the Langmuir paradox. The effective frequency of randomizing collisions is calculated; the pressure dependence of this frequency is consistent with a wall mechanism for electron randomization. The hydrodynamic approximation is used to analyze striations at low pressures; the theoretical and experimental results show agreement. (auth)

47134 NOTE ON PLASMA ACCELERATION CAUSED BY PLASMA TURBULENCE. Motz, H. (Oxford Univ.); Tsytovich, V. N. Plasma Phys.; 14: No. 6, 583-9 (Jun 1972).

Momentum exchange between two plasmas in relative motion, one more dense than the other, is investigated. It is concluded that the dominant turbulent process is the scattering of Langmuir waves by ions of the denser plasma. The length over which the relative motion ceases increases with the fourth power of the ratio of the relative speed to the thermal velocity. When the relative velocity exceeds a critical value, the generation of Langmuir turbulence is stabilized and eventually the order motion is no longer dispersed. Application to a laboratory experiment is discussed and it is also suggested that the process investigated might be of importance in the case of galaxies expanding into intergalactic gas. (auth)

47135 IMPEDANCE OF A PARTIALLY IONIZED CESIUM PLASMA. Yamada, Jun; Okuda, Takayoshi (Nagoya Univ., Japan). Jap. J. Appl. Phys.; 11: No. 7, 1032-8 (Jul 1972).

The impedance of a partially ionized cesium plasma is measured over a wide range of signal frequency. A theory is developed, taking into account the perturbations of the density, velocity, and temperature of the electron and also of the population of the excited states of the atom for the response of such a plasma to an ac electric field of small amplitude. The impedance formula derived from the above is confirmed by the experiment. (auth)

47136 CATHODE HEATING IN AN ASYMMETRICALLY BURNING PENNING DISCHARGE. Hopfgarten, N. (Royal Inst. of Tech., Stockholm). Plasma Phys.; 14: No. 4, 397-401 (Apr 1972).

Ions and electrons bombarding the cathodes in a Penning discharge in a magnetic mirror field heat the cathodes and release secondary electrons. From measurements of cathode heating, currents, and burning voltages, it is possible to estimate the density and energy of ions leaving the discharge to the cathodes, as

well as the currents of electrons arriving at the cathodes and the currents of electrons leaving them. (UK)

47137 MASS, MOMENTUM, AND ENERGY FLOW FROM AN MPD ACCELERATOR. Cory, John Slane. Princeton, N. J.; Princeton Univ. (1971). 112p. University Microfilms Order No. 72-18,799.

Thesis.

The mass, momentum, and energy flows through a quasi-steady, self-field MPD accelerator are measured over a current range of 8 to 50 kA and inlet mass flows of 2 to 36 g/s of argon. Time-resolved arc chamber pressure and impact pressure in the exhaust jet, measured with a piezoelectric pressure probe capable of operation with a flat response from 1 to 140 kc, are used to calculate the momentum flux profiles, the gasdynamic contribution to thrust, and the total thrust. The momentum flux profile indicates that the accelerator produces a uniform, 2-inch-diameter axial jet at the anode which expands into a Gaussian profile at an axial station 11 inches from the anode. The electromagnetic component of the thrust is found to follow the familiar quadratic dependence on arc current, while a more complex empirical relation is needed to correlate the gasdynamic contribution with the current and mass flow rate. Using available time-of-flight velocity profiles at a current of 16 kA and a mass flow of 5.9 g/s, calculated flux profiles of mass and kinetic energy exhibit a tendency for some fraction of the inlet mass flow to leak out at a low velocity around the central high-velocity core. Discharge chamber geometry changes to eliminate the leakage will doubtless improve the arc performance. (Diss. Abstr. Int., B)

47138 PROCESSES AND MEASUREMENTS IN AN ABNORMAL NEGATIVE GLOW PLASMA. Kaul, Ashok Kumar. College Park, Md.; Univ. of Maryland (1972). 174p. University Microfilms Order No. 72-20,251.

Thesis.

A stable dc discharge system using a brush cathode, and capable of generating a negative glow of useful dimensions, is described. Measurements of electron densities and temperatures were made under steady-state conditions along the axis of the glow, using a 1-mil cylindrical Langmuir probe. A preliminary investigation of the "hysteresis" encountered in the Langmuir-probe characteristics showed that this effect is probably caused by the variation in surface contact potential arising from the neutralization of certain types of impurity ions and their subsequent adsorption by the probe surface. Electron and ion densities are found to satisfy quasi neutrality for the negative glow plasma, and range from about 2.4×10^8 to $3 \times 10^{12} \text{ cm}^{-3}$, with the electron temperature first decreasing with increasing pressure from about 1500 to 650°K, and then increasing again to about 2000°K at a pressure of approximately 1 torr. The plasma is found to be fairly homogeneous axially, and the effect of the discharge current was also investigated. A theoretical model is presented to explain the observed densities and temperatures, and in particular, the behavior of the electron temperature. This model assumes the existence of the metastable-metastable ionizing collisions and the superelastic collisions in addition to the impact ionization heating mechanism. The loss or cooling mechanisms assumed are ambipolar diffusion, recombination, and elastic collisions with ions and neutral gas atoms, while the effect of electron-electron collisions is also included. This yields reasonable agreement with measured electron temperatures and gives the correct temperature variation. The observed electron densities are well explained on the basis of a recombination-dominated plasma over most of the range of the experiment. The recombination-rate coefficients obtained are in the vicinity of the collisional-radiative rates of Kingston, Bates, and McWhirter. (Diss. Abstr. Int., B)

47139 OBSERVATION OF SECOND COMPRESSION IN THE FINAL STAGE OF A DISCHARGE OF THE PLASMA FOCUS TYPE. Gribkov, V. A.; Korzhavin, V. M.; Krokhn, O. N.; (and others) (Lebedev Inst. of Physics, Moscow). JETP Lett. (USSR) (Engl. Transl.); 15: No. 6, 232-4 (20 Mar 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 6, 329-32 (20 Mar 1972).

Results are presented of a high-speed interferometry study of the final plasma focus (PF) stage. The results show that the PF current sheath becomes ruptured at the instant of maximum of the hard x-ray pulse, that the remaining parts of the sheath are additionally compressed, and turbulence develops at the rupture place. The results also establish an additional mechanism whereby the plasma flows out of the sheath. (M.O.W.)

47140 MODEL FOR THE HOLLOW CATHODE DISCHARGE. Lorente-Arcas, A. (Paris Univ., Orsay, France). Plasma Phys.; 14: No. 6, 651-9 (Jun 1972).

The existence of a local overpressure in a hollow-cathode discharge (HCD) has been detected. In order to explain this, it is

supposed that the plasma density in the active zone is essentially determined by the equilibrium between the ionization produced by the electron current and the electron-ion recombination. In the same way, the total density of atoms plus ions in the active zone is determined by the gas flow. The up-stream limit of the active zone is found to be a thin sheath created by ambipolar diffusion and recombination between electrons and ions (ADAR sheath); one criterion for such a sheath to exist has been found. The variation of pressure up-stream from the active zone in terms of the discharge current has been found. By comparing the theory with experiment, a total recombination coefficient is found whose order of magnitude is in agreement with the values given in the literature. (auth)

Kinetics (Experimental)

47141 (LA-DC-72-703) PLASMA EXPERIMENTS IN THE SCYLLAC 5-METER, LINEAR θ PINCH. Thomas, K. S.; Harris, H. W.; Jahoda, F. C.; Sawyer, G. A.; Siemon, R. E. (Los Alamos Scientific Lab., N. Mex.). 1971. 4p. (CONF-720710-5). Dep. NTIS.

From second topical conference on pulsed high-beta plasmas; Garching, Germany (3 Jul 1972).

Experiments are reported on the behavior of the plasma column in a straight high-voltage θ pinch which may be operated both with and without strong magnetic mirrors. Operation without mirror fields produced a high temperature (2 to 3 keV) plasma column which lasted about 15 μsec , showed considerable "wobble," and also showed effects due to plasma contamination of the discharge tube by material "boiled" off the quartz discharge tube by the previous discharge. Enlarging the discharge tube outside the coils reduced this problem. In the data so far, when mirror fields which give a mirror ratio of 2 to 3 are applied at the same time as the main field, the plasma column shows evidence of a $m = 1$ instability. (auth)

47142 NEUTRAL-POINT DISCHARGE EXPERIMENT. Ohyaabu, Nobuyoshi; Kawashima, Nobuki (Tokyo Univ.). J. Phys. Soc. Jap.; 33: No. 2, 496-501 (Aug 1972).

The behavior of a plasma current through a magnetic neutral point is investigated experimentally. A quasi-steady magnetic field (either cusp or quadrupole magnetic field) is created by a slow capacitor discharge, and the plasma sheet current is excited by a slow capacitor discharge. The plasma current sheet thus formed is collapsed into small filamentary pinches, and they are ejected outward on the current sheet plane along the magnetic field lines. This seems to correspond to the ejection of plasma and energetic particles associated with solar flares. The visual observation together with double probe and Rogowski coil measurements show that the collapse of the plasma current occurs earlier as the plasma current and the external magnetic field are increased and as the initial gas pressure is decreased, and this is explained qualitatively by the resistive tearing-mode instability. (auth)

47143 ENHANCED FAR-INFRARED RADIATION FROM A THETA-PINCH PLASMA. Chin-Fatt, Cecil A. College Park, Md.; Univ. of Maryland (1972). 95p. University Microfilms Order No. 72-20,785.

Thesis.

A nonthermal spectrum was observed of far infrared radiation, covering a wavelength range of approximately 0.4 mm to 2 mm emitted by the turbulent helium plasma of a theta pinch with a strongly diffusive magnetic piston, at a peak intensity that is at least three orders of magnitude greater than that expected of an equivalent thermal plasma. Line emission and magnetic probe studies in addition to the temporal development and the observed two-banded structure of the enhanced nonthermal spectrum, support the interpretation of this far infrared emission to be direct evidence of two nonlinear radiation phenomena, involving the interactions of longitudinal electron waves with plasma inhomogeneities and with themselves and resulting in bremsstrahlung radiation at the plasma frequency, ω_{pe} , and its harmonic $2\omega_{pe}$; these radiation processes were theoretically proposed to explain types II and III radio bursts associated with solar flares. A theoretical estimate, based on a fluid plasma model, gave fair agreement with the observed intensity of the far infrared radiation. (Diss. Abstr. Int., B)

47144 SYNCHROTRON RADIATION FROM A MAGNETICALLY CONFINED HOT-ELECTRON PLASMA. Nee, Soe-mie Foeng. College Park, Md.; Univ. of Maryland (1972). 189p. University Microfilms Order No. 72-20,260.

Thesis.

The spectrum of the synchrotron radiation emitted by the elec-

tron cloud produced in the University of Maryland pulsed magnetic-mirror compression experiment was measured in a wavelength region from 3.0 cm to 0.3 mm using three independent detector systems. The measured spectra agree with the theoretical spectra calculated using a distribution function derived from a relativistic Maxwellian distribution function that conserves the magnetic moments of the trapped electrons during magnetic compression of the electron cloud. This agreement demonstrates that the electron cloud is compressed adiabatically and that electrons are not lost during the decay of magnetic field. The analytical expressions of this distribution function and its generalization to the three-dimensional case are derived. Single-particle effects on the measured spectrum of synchrotron radiation are also investigated. The effects of a finite aperture to broaden the cyclotron lines and to change the intensity by $(1 - \beta_{\parallel} \cos \theta)$ such that the observed synchrotron radiation, including the aperture effect and the retardation effect, is just equal to the radiation emitted by an electron drifting freely along the magnetic field. The correction for the electron passing through the observing window is to split the broadened spectrum due to the finite aperture into many lines separated by the longitudinal frequency. The spectrum of the synchrotron radiation for an electron confined in a static inhomogeneous magnetic mirror field covers a wide continuous frequency range approximately from $\omega_c/(1 + \beta_{\parallel} \cos \theta_0)$ to $\omega_c/(1 - \beta_{\parallel} \cos \theta_0)$, and its shape depends entirely on the field configuration. This correction is important for observation along the magnetic axis and is negligible for observation perpendicular to the magnetic axis. (Diss. Abstr. Int., B)

Kinetics (Theoretical)

Refer also to abstracts: 47163, 47164, and 47182.

47145 (COO-3086-15) RESEARCH IN COMPUTATIONAL PLASMA PHYSICS. Progress Report, March 1, 1971-March 1, 1972. Chu, C. K.; Taussig, R. T. (Columbia Univ., New York). 1 Mar 1972. Contract AT(11-1)-3086. 10p. Dep. NTIS.

Research progress on the following four areas is briefly described: (1) numerical simulation of the shock tube; two-dimensional one-fluid MHD code, (2) heat transfer studies for the shock tube, (3) Vlasov equation and ion acoustic phenomena, and (4) simulation of slightly collisional plasmas. (M.O.W.)

47146 KINETIC THEORY OF A TWO-DIMENSIONAL MAGNETIZED PLASMA. Vahala, George Martin. Iowa City, Iowa; Univ. of Iowa (1972). 199p. University Microfilms Order No. 72-17,613.

Thesis.

Several features of the equilibrium and nonequilibrium statistical mechanics of a two-dimensional plasma in a uniform dc magnetic field are investigated. The calculations have been motivated by the recent derivation of Bohm's diffusion coefficient given by Taylor and McNamara for this system. The charges interact only through electrostatic (logarithmic) potentials. The problem is considered both with and without the guiding-center approximation. With the guiding-center approximation, an appropriate Liouville equation and BBGKY hierarchy predict no approach to thermal equilibrium for the spatially uniform case. For the spatially nonuniform situation, a guiding-center "Vlasov" equation is discussed and is solved in special cases. The guiding-center model is shown to be equivalent to the fluid mechanics problem of the motion of rectilinear vortices in an inviscid, incompressible fluid. The most interesting features of thermal equilibrium theory (with and without the guiding-center approximation) are (1) a collapse of the system above a critical value of the plasma parameter (May 1967); and (2) a divergence in the electric field fluctuation spectrum (minus the self-energy terms) for small plasma parameter and very large systems. For the nonequilibrium, nonguiding-center case, a Boltzmann equation and a Fokker-Planck equation are derived in the appropriate limits. The Fokker-Planck equation is more tractable than the Boltzmann equation, but contains a divergent integral which must be cut off on physical grounds. This cutoff is compared to the standard cutoff introduced in the two-dimensional unmagnetized Fokker-Planck equation. In the low plasma parameter it is shown, under the assumption that for large (integer) n $(\gamma n / \gamma_{n+1}) = O(n^p)$, $p < 2$ where $\gamma_n \equiv \omega_n - n\Omega$, with ω_n the n^{th} Bernstein mode and Ω the electron gyrofrequency, that the Balescu-Lenard collision term is zero in the long-time average limit if one considers only two-body interactions. The energy transfer from a test particle to an equilibrium plasma is discussed and also shown to be zero in the long-time-average limit. This supports the result of zero Balescu-Lenard collision term. (Diss. Abstr. Int., B)

47147 NUMERICAL SIMULATION OF RELATIVISTIC ELECTRONS CONFINED IN AN AXISYMMETRIC MIRROR FIELD. Brettschneider, Marcel. Davis, Calif.; Univ. of California (1971). 145p. University Microfilms Order No. 72-17,898.

Thesis.

In the controlled fusion experiment, Astron, a beam of relativistic electrons is injected into a partially evacuated cylindrical region in which an externally applied magnetic field has been established. The cylindrical region is bounded by two concentric aluminum shells. The applied field, in most cases, is solenoidal. It is generated by a large number of coils located outside of and concentric with the cylindrical region. Generally, the currents in the coils are programmed to create a shallow magnetic mirror near one end of the region and a steep mirror field at the other end. The electrons are injected near that end of the region which contains the shallow mirror. Their subsequent trajectories are helical. To make it easier to trap the electrons, a cylinder of azimuthally oriented resistor wires, concentric with and located somewhat inside of the outer cylinder, was installed. The electrons trapped form a cylindrical shell known as the E-layer. If enough electrons are trapped, the self-field of the E-layer can exceed the applied field, resulting in a magnetic field configuration which includes a region which is minimum $|B|$ and has no loss cones. A mathematical model that is capable of simulating the formation of the E-layer was developed and is described here. The model assumes axial symmetry. It includes all the physics described above, and also the effects of the neutralization of the electric field of the injected electrons due to their interaction with the background gas. This model was used to study the formation of an E-layer when one pulse of electrons was injected. It was also used to study the effect of injecting a number of pulses in succession. The results are in good qualitative agreement with the observed experimental results. (TSS) (Diss. Abstr. Int., B)

47148 EFFECTS OF ION MOTION ON SPECTRAL LINE BROADENING IN PLASMAS. Roszman, Larry Joe. Gainesville, Fla.; Univ. of Florida (1971). 219p. University Microfilms Order No. 72-16,652.

Thesis.

The effect of ion motion on the broadening of atomic spectral lines by plasmas is examined. A review and discussion of the quasi-static ion theory of plasma line broadening and previous work on the problem of ion motion are included. It is concluded that two important aspects of the ion perturbations of the radiator must be taken into account when examining ion motion. First, many ions interact simultaneously with the radiator, i.e., multiple collisions between the ions and the radiator are overwhelmingly significant. Second, neither the quasi-static nor the time-dependent ion-radiator interaction can be well treated by finite-order time-dependent perturbation theory. Forms of the time-dependent ion microfield distribution suitable for numerical computation are derived first for an ion gas neglecting ion-ion correlations, and then for an ion gas including ion-ion correlations. The derivations of the expressions for the second case are based upon time-dependent generalizations of the cluster-expansion and collective-coordinate methods employed by Hooper for the quasi-static microfield distribution. Some numerical results are given for the case which neglects ion-ion correlations. It is found that the effects of ion motion are significant for frequency separations from the line center less than the reciprocal of the time it takes an ion traveling with the average thermal velocity to cross the ion sphere radius. (TSS) (Diss. Abstr. Int., B)

47149 TOTAL ENERGY LOSS OF A TEST CHARGE IN A COLLISIONAL MAGNETOPLASMA. Shukla, P. K. (Umeaa Univ., Sweden); Singh, R. N. Phys. Scr.; 4: No. 6, 281-9 (1971).

A general theory of energy attenuation of a charged test particle in a cold, collisional plasma is developed. The effect of collisions on the plasma properties is investigated. The radiation characteristics of a spiralling test particle are discussed. In the presence of collisions it is found that absorption is important when the collision frequency is comparable to the wave frequency. Estimates of radiated and absorbed power by test charges of different energies are made, and the role of collisions on the power spectrum is studied in detail. (auth) (Sweden)

47150 SELF-CONSISTENT PAIR CORRELATIONS AND DIELECTRIC RESPONSE IN A PLASMA. Abramo, M. C.; Tosi, M. P. (Univ., Messina, Italy. CNR, Messina, Italy). Nuovo Cim.; 10B: No. 1, 21-35 (11 Jul 1972).

The self-consistent approach to the determination of the dielectric response function and of pair correlations in an electron liquid, developed by Singwi et al., is systematically examined in the classical limit. The self-consistent equation for the pair correlation function is obtained from the second equation of the BBGKY hierarchy, by a decoupling approximation on the triplet

correlation function which takes account of the limiting behaviors at long wavelengths and at short distances. In the same order of approximation an effective two-body potential is derived by an alternative decoupling scheme, whose use allows one to improve the self-consistent pair correlation at short distances. A collective potential, inclusive of local field corrections, is also proposed for the evaluation of the wave vector and frequency-dependent dielectric function in both the classical and the degenerate electron liquid. The decoupling approximation is shown to be consistent with the second-order perturbative treatments of the triplet function which separately apply in the long-wavelength domain and in the short-distance domain, and the systematic extension of the decoupling scheme to higher orders is indicated and illustrated. The present results are expected to be of practical usefulness in the treatment of correlations in the degenerate electron gas at metallic densities and in the high-density, low-temperature plasma. (auth)

47151 NON-LINEAR THEORY OF INSTABILITY OF A MONO-ENERGETIC ELECTRON BEAM IN PLASMA. Matsuboro, N. G.; Onishchenko, I. N.; Shapiro, V. D.; Shevchenko, V. I. (Physical Technical Inst., Kharkov). *Plasma Phys.*; 14: No. 6, 591-600 (Jun 1972).

The excitation of a monochromatic plasma wave by a "cold" electron beam is investigated. The solution obtained describes the transition from the exponential growth of the wave amplitude in the linear stage to the amplitude oscillations associated with trapping of the beam in the potential well created by the wave. The maximum amplitude and its oscillation period are found. Both nonrelativistic and relativistic beams are investigated. (auth)

47152 BREMSSTRAHLUNG PHOTON EMISSION RATE FROM MAXWELLIAN PLASMAS. Halverson, W. (Massachusetts Inst. of Tech., Cambridge). *Plasma Phys.*; 14: No. 6, 601-4 (Jun 1972).

The bremsstrahlung photon emission rate from a Maxwellian plasma for photon energies greater than a given threshold value is expressed in terms of a rapidly converging series. The results are used to determine the applicability of soft x-ray diagnostic techniques for laser-produced plasmas, as a function of electron temperature and detector threshold energy. (auth)

Production

47153 (NP-19364) PRODUCTION AND MEASUREMENT OF INTENSE ION BEAMS AND BEAM-PRODUCED PLASMA IN A CUSPED MAGNETIC FIELD. Huizenga, W. (Stichting voor Fundamenteel Onderzoek der Materie, Jutphaas (Netherlands). Instituut voor Plasma-Fysica). Mar 1972. 86p. Dep. NTIS (U. S. Sales Only).

Experimental work on the production and current measurement of intense hydrogen-ion beams in pulsed condition (60 mA at 20 kV) is described. In order to produce an intense ion beam, use is made of a duoplasmatron provided with a plasma-expansion cup at anode potential. Results obtained with various configurations of grids mounted in the ion-extraction region were compared. The use of concave grids appears to be a good expedient for the production of ion beams with sufficient initial focusing to balance space-charge repulsion. Beam-current measurements were made with a Faraday collector provided with a secondary-electron suppressor. Observations show that the voltage needed to suppress the escape of secondary electrons from the collector is approximately proportional to the space-charge potential of the beam in the center of the collector entrance. Special attention was paid to the beam composition, as was proved necessary in these experiments. Next, the formation of plasma by injection of a 20-keV hydrogen-ion beam into a quadrupole magnetic-containment device was studied. Electron-density measurements were made with a 9-mm microwave interferometer and with resonance-shift methods using the metallic vessel as a cavity. Also, current measurements were made with collectors located in the mirror regions. The plasma density in the steady state appears to be proportional to the gas pressure in the range up to 10^{-4} torr. Under normal conditions (injected current 50 mA, gas pressure 2×10^{-5} torr), the average density is about 5×10^8 el cm^{-3} . The observed plasma-ion temperature is about 150 eV at low gas pressure, whereas the electron temperature does not exceed a few eV. The energy transfer from the hot-ion component to the plasma is in the order of one part per thousand. The energy transfer is thought to be caused by collective effects (beam-plasma interaction). Considerations on the plasma-particle balance lead to the existence of a positive plasma potential in the center during the steady state (about 70 V at low gas pressure). The existence of this potential was confirmed experimentally. (auth)

47154 FORMATION OF THE ENERGY SPECTRUM OF THE IONS OF A LASER PLASMA. Bykovskii, Yu. A.; Vasil'ev, N. N.; Degtyarenko, N. N.; (and others) (Moscow Engineering Physics Inst.). *JETP Lett. (USSR) (Engl. Transl.)*; 15: No. 6, 217-20 (20 Mar 1972).

Translated from *Zh. Eksp. Teor. Fiz., Pis'ma Red.*; 15: No. 6, 308-11 (20 Mar 1972).

The effects of acceleration and recombination on the ion energy spectra were investigated. A detailed study was made of the energy distributions of hydrogen and zirconium ions and deuterium and lithium ions (LiD) at different flux densities. Energy distribution diagrams are given. (M.O.W.)

47155 MOMENTUM TRANSFER AND PLASMA FORMATION ABOVE A SURFACE WITH A HIGH-POWER CO₂ LASER. Pirri, A. N.; Schlier, R.; Northam, D. (Avco Everett Research Lab., Everett, Mass.). *Appl. Phys. Lett.*; 21: No. 3, 79-81 (1 Aug 1972).

An experiment was performed to measure the total impulse transferred to a surface by a high-power pulsed 10.6- μ laser under atmospheric conditions, and also to determine the degree to which the observed plasma decouples the laser energy from the surface. Specific interest was directed toward conditions under which a "radiation-driven detonation wave" may form due to the presence of a surface. The momentum transferred in dyn sec/J decreases with an increase in the ratio of incident energy to spot area since a larger fraction of the energy is decoupled from the surface. (auth)

Stability

47156 (LA-DC-72-706) PLASMA EXPERIMENTS ON $l = 1, 0$ HELICAL EQUILIBRIA IN THE SCYLLAC 5-METER θ -PINCH TOROIDAL SECTOR. Ellis, W. R.; Hammer, C. F.; Jahoda, F. C.; Quinn, W. E.; Ribe, F. L.; Siemon, R. E. (Los Alamos Scientific Lab., N. Mex.). 1972. 4p. (CONF-720710-3). Dep. NTIS.

From second topical conference on pulsed high-beta plasmas; Garching, Germany (3 Jul 1972).

Experiments are reported on a sector comprising one third of the 4.8-m-diameter Scyllac torus. The toroidal force and outward plasma drift are compensated by an interference force produced by a combination of $l = 1$ helical fields and $l = 0$ bumpy fields. The toroidal equilibrium is observed, followed by an $m = 1$ (sideward) motion of the plasma column, which is nearly the same all along the torus, independent of the $l = 0$ periodicity, with plasma containment times as large as 11 μsec , comparable to times for plasma end loss. The $m = 1$ motions, which occur predominantly in the horizontal plane of the torus, suggest either an imbalance between the $l = 1, 0$ and toroidal forces at later times or a long wavelength $m = 1$ instability. Measurements of the applied magnetic fields show that the product of $l = 1$ and $l = 0$ fields for plasma equilibrium agrees with sharp-boundary MHD theory. Time-resolved measurements of the plasma diamagnetism, electron density distribution, neutron emission, and beta are reported. (auth)

47157 (LA-DC-72-707) FEEDBACK STABILIZATION ON AN $l = 1$ THETA-PINCH PLASMA COLUMN. Gribble, R. F.; Burnett, S. C.; Harder, C. R. (Los Alamos Scientific Lab., N. Mex.). 1971. 4p. (CONF-720710-4). Dep. NTIS.

From second topical conference on pulsed high-beta plasmas; Garching, Germany (3 Jul 1972).

An $l = 0$ MHD feedback experiment to control the $m = 1$ instability on a plasma column subject to $l = 1$ helical fields is described. The feedback system has three basic components: a position detector, a signal processor, and a power amplifier. Feedback stabilization is being tried first on a linear θ pinch (Scylla IV-3). Horizontal and vertical motions of the plasma column are optically sensed with fast-rise-time silicon bi-cell detectors. The signal processor provides amplitude and derivative control and normalization of the detector signals. Ten power-amplifier modules in push-pull configurations are used to drive the $l = 0$ windings. The $l = 1$ coils which drive the instability are driven by a crowbarred capacitor bank. The module output power of 30 MW rises in 1.3 μsec and may be sustained for as much as 20 μsec . From detector output to power-module output the power gain of the feedback system is 7×10^{14} . (auth)

47158 (LA-DC-72-726) RECENT RESULTS FROM THE SHOCK-HEATED TOROIDAL Z-PINCH EXPERIMENT ZT-1. Burkhardt, L. C.; Di Marco, J. N.; Forman, P. R.; Haberstich, A.; Karr, H. J.; Phillips, J. A. (Los Alamos Scientific Lab., N. Mex.). 1971. 4p. (CONF-720710-1). Dep. NTIS.

From second topical conference on pulsed high-beta plasmas; Garching, Germany (3 Jul 1972).

Preionization in this experiment is produced by a ≈ 20 kA z-current of period ~ 30 μ sec. The degree of ionization and radial distribution of current density are functions of the filling pressure and strength of the preionization current. After the start of the main z current, with initial B_θ (wall) of ~ 60 kG/ μ sec and peak currents of ~ 200 kA, plasma pressure profiles show a positive pressure gradient on axis, as required for MHD stability, existing for times of at least 8 μ sec (current zero at ~ 11 μ sec). The plasma β_θ is ~ 0.4 at 5-mtorr filling pressure and increases with the filling pressure. Plasma temperatures ($T_i + T_e$) of 1 to 4 keV are inferred from the pressure profiles and initial gas-filling densities with the higher temperatures corresponding to the lower pressures (2 mtorr). When the B_z field outside the pinch column is reversed, streak photography and MHD stability analysis show significant improvement towards stability. High plasma conductivity prevents the zero in B_z from moving away from the wall more than ~ 0.5 cm. (auth)

47159 (UCRL-Trans-10630) STABILITY OF A FINITE-PRESSURE PLASMA WITH AN ANISOTROPIC ELECTRON-VELOCITY DISTRIBUTION. Egorenkov, V. D.; Stepanov, K. N. Translated for Univ. of California Lawrence Livermore Lab., Livermore, from Ukr. Fiz. Zh. (Russ. Ed.); 15: No. 2, 325-30 (1970). 12p. Dep. NTIS.

A homogeneous finite-pressure plasma with hot electrons and cold ions and with an anisotropic electron distribution is considered. It is shown that, in the high-frequency range $\omega_{He} \gg |\omega| \gg \omega_{Hi}$, the instability region for electromagnetic waves analogous to whistlers and ion-acoustic waves coincides with that for diamagnetic and firehose instabilities. Since the growth rates of the unstable oscillations considered are appreciably greater than the ion cyclotron frequency, these instabilities are more dangerous than the diamagnetic and firehose ones, for which the growth rates are considerably smaller than ω_{Hi} . (auth)

47160 CONVECTION OF A PLASMA IN A GRAVITATIONAL FIELD. Wobig, Horst (Princeton Univ., N. J.). Plasma Phys.; 14: No. 4, 403-16 (Apr 1972).

It is shown that the interchange instability of a magnetoplasma in a gravitational field can be stabilized by the combined effects of viscosity and resistivity. (UK)

47161 ABSOLUTE AND CONVECTIVE INSTABILITIES. Bijlsma, S. (FOM-Instituut voor Atoom-en Molecuulfysica, Amsterdam). Plasma Phys.; 14: No. 4, 420-2 (Apr 1972).

It is shown that the use of a saddle-point method can lead to the determination of criteria for distinguishing between absolute and convective instabilities in a one-dimensional linear infinite system. (UK)

47162 FLUTE MODES IN A PLASMA IN THE PRESENCE OF NONUNIFORM ELECTRIC FIELDS. Carenza, L.; Engelman, F. (EURATOM-CNEN, Rome). Nuovo Cim.; 10B: No. 1, 127-43 (11 Jul 1972).

The effects of the presence of a nonuniform electric field on the stability of flute modes in a plasma are studied in the approximation of small Larmor radius. Most of the work is concerned with discussing and applying to the corresponding eigenvalue problem a step approximation for the profiles of the electric and diamagnetic drifts. Furthermore, a new sufficient stability criterion for flute modes is formulated. The influence of the drifts on the stability properties of the flute modes is discussed, and, in particular, the stabilizing effect of a density gradient is displayed. (auth)

47163 STATISTICAL THERMODYNAMICS OF COLLISIONLESS PLASMA. I. THE THERMODYNAMICAL POTENTIALS. Minardi, E. (EURATOM-FOM, Jutphaas, Netherlands). Plasma Phys.; 14: No. 5, 427-41 (May 1972).

It is shown that the equilibria and the stability of nonMaxwellian inhomogeneous plasma can be described by a theory with a thermodynamical structure, provided that, in the linearly unstable cases, the perturbation modes under consideration are not too far from marginal stability. On the one hand, variational functionals prove to exist which can characterize the equilibrium and its stability by means of their extremum properties; on the other hand, a statistical model is proposed from which the same functionals can be derived in an independent way. (UK)

47164 STATISTICAL THERMODYNAMICS OF COLLISIONLESS PLASMA. II. STABILITY THEORY OF INHOMOGENEOUS SYSTEMS. Minardi, E. (EURATOM-FOM, Jutphaas, Netherlands). Plasma Phys.; 14: No. 5, 443-8 (May 1972).

The statistical model developed in Part I is applied to the discussion of the stability of inhomogeneous Vlasov equilibria. The existence of instability is expressed by a general condition provided a certain class of variations compatible with the physical constraints imposed on the system does exist. It is also shown, by means of the entropy principle, that the isolated inhomogeneous

collisionless systems evolve towards more homogeneous equilibria, associated with a higher entropy, by transforming their collective energy into the energy of the random individual fluctuations. Moreover, nearly homogeneous but electrostatically unstable collisionless plasmas prove to evolve towards the marginal state of the stable equilibrium (which is a state of maximum entropy for the collisionless systems), while an electrostatic quasi-static marginally unstable mode with long wavelength appears in the system. (auth)

47165 MICRO-INSTABILITIES OF HIGH PRESSURE PLASMA ASSOCIATED WITH DEPENDENCE OF THE ION VISCOSITY ON THE DERIVATIVES OF THE LONGITUDINAL THERMAL FLUX. Mikhailovskii, A. B.; Tsypin, V. S. (Inst. of Atomic Energy, Moscow). Plasma Phys.; 14: No. 5, 449-59 (May 1972).

Microinstabilities of an inhomogeneous, collisional, high-pressure plasma are investigated. Attention is paid mainly to perturbations with large longitudinal wave numbers, for which the longitudinal ion viscosity is significant. It is shown that for large plasma pressure, the derivatives of the longitudinal thermal flux must be included in the expressions for ion viscosity. Effects of the ion viscosity lead to instability of high-pressure plasma, both in the absence of a magnetic field gradient and when it is oriented in the same direction as the density gradient. (auth)

47166 DRIFT-KINETIC EQUATION AND ELECTROSTATIC INSTABILITY THEORY IN TOKAMAK PLASMAS. Cheung, Lung. Austin, Tex.; Univ. of Texas (1971). 139p. University Microfilms Order No. 72-19,569.

Thesis.

Approximate solutions are obtained to the Vlasov equation that are valid under the conditions that the particle gyro-motion is on a small scale compared to the macroscopic scale. Stability theory of drift modes in a nonuniform, collisionless plasma in the Tokamak confinement geometry is discussed. The stability theory developed substantially generalizes previous systematic treatments given for the plane-slab geometry and the closed-field-line multipole geometry. In particular, the stability of a Maxwellian distribution with a density gradient and a toroidal current is considered in detail. The modified stability criteria and growth rates for the modes are found. (TSS) (Diss. Abstr. Int., B)

47167 EQUILIBRIUM AND STABILITY THEORY OF NON-NEUTRAL PLASMAS. Bogema, Bernard LeRoy. College Park, Md.; Univ. of Maryland (1972). 123p. University Microfilms Order No. 72-18,882.

Thesis.

Equilibrium and stability properties of magnetically confined nonneutral plasmas are studied theoretically. Cold-fluid equilibria are constructed in the relativistic and nonrelativistic regimes, exhibiting a variety of density profiles. Electromagnetic perturbations about nonrelativistic equilibria are considered for the ordinary and extraordinary modes of polarization. For constant-density plasmas, the perturbed fields are determined. Electrostatic surface waves on finite-radius columns are studied. Special attention is directed to the case that the plasma has two components rotating at different velocities. In this case, instability may develop for sufficiently high density. Finite-temperature effects are included through a Vlasov description of nonneutral plasmas. The distribution function is assumed to depend on perpendicular energy $H_\perp$ and angular momentum L through the linear combination $H_\perp - \omega_R L$, where ω_R is a constant. The corresponding equilibria rotate rigidly in the mean about the axis of symmetry with angular velocity ω_R . In the nonrelativistic limit, a necessary and sufficient condition for radial confinement is derived, and it is shown that the perpendicular distribution function $\int dp_\parallel^2 (H_\perp - \omega_R L, p_\parallel)$ may be obtained from the equilibrium density profile. The equilibrium equations appropriate to relativistic rigid rotors are also presented. Examples of equilibria are given in both the nonrelativistic and relativistic regimes. Stability of small amplitude electrostatic perturbations about nonrelativistic Vlasov equilibria is considered, emphasizing finite radius effects. Body waves are derived for small temperature and temperature corrections are given for surface waves. A dispersion relation is obtained in the case that the plasma is composed of two equilibrium electron beams counterstreaming along the external magnetic field $B_0 \hat{z}$. An unstable region of k_z is found that exists only for finite column radius. Finally, an analogue of the Rostoker equation is derived for stable nonneutral plasmas, and quasi-linear equations are developed for weakly unstable systems. (Diss. Abstr. Int., B)

47168 EFFECTS OF PARALLEL WAVELENGTH ON THE COLLISIONAL DRIFT INSTABILITY. Schlitt, Leland Glenn. Princeton, N. J.; Princeton Univ. (1972). 88p. University Microfilms Order No. 72-18,787.

Thesis.

An extensive investigation of the dependence of drift-wave

characteristics upon parallel wavelength in collisional potassium Q-machine plasmas is presented. The experiment was performed on the Princeton Q-1 device modified to permit variation of plasma column length ($L = 15$ to 135 cm) and thus parallel wavelength ($\lambda_{||} \approx 2L$ under "electron-rich" conditions; $n_0 = 10^{10}$ to 2×10^{11} cm $^{-3}$, $T_e = 2650$ K). Results are compared with a two-fluid slab-model theory which includes the effects of parallel resistivity and thermal conductivity for electrons and Larmor radius, collisional viscosity, and motion parallel to the magnetic field for ions. The general-stability characteristics of the drift wave are confirmed: stability at low magnetic field ($B/k_{\perp} \approx 0.4$ kG-cm) due to ion viscosity, stability at both long and short parallel wavelength ($\lambda_{||} \gtrsim 200$ and $\lambda_{||} \lesssim 100$ cm, respectively) due to the combined effects of ion viscosity and parallel electron fluid expansion, and stability of individual modes at high magnetic field ($B/k_{\perp} \approx 1$ kG-cm, $\lambda_{||} \approx 100$ cm) due to ion parallel motion. In addition, quantitative agreement between experiment and theory to within a factor of 2 is demonstrated for the detailed parametric dependences of drift-wave properties (instability onset, complex frequency, and oscillation amplitude) on parallel wavelength, magnetic field, and density. (Diss. Abstr. Int., B)

47169 THRESHOLD OF IONIZATION INSTABILITY IN AN UNBOUNDED LOW TEMPERATURE PLASMA. Zampaglione, V. (CNEN, Frascati, Italy). Plasma Phys.; 14: No. 6, 664-6 (Jun 1972).

The complete dispersion relation for ionization waves propagating in an unbounded low-temperature plasma is analytically solved by neglecting the electron pressure term in the Ohm's Law solution, together with the heat convection in the electron energy equation. Within this limit it is found that the threshold of ionization instability does not depend on the relaxation rates within a broad range of operating conditions for the MHD energy conversion method. (UK)

Thermonuclear Processes

Refer also to abstract: 45433.

47170 (IPP-IV-32) POWER PRODUCTION COSTS AND POSSIBLE APPLICATIONS OF SHORT-PERIOD COMBUSTION MHD GENERATORS. Buende, R.; Helm, S.; Muntenbruch, H. (Max-Planck-Institut fuer Plasmaphysik, Garching (West Germany)). Jan 1972. 58p. (In German). Dep. NTIS (U. S. Sales Only).

Possibilities of applying kerosene-oxygen fired MHD plants for fusion experiment power supply and for reserve power production in industrial and public grids were investigated. Costs of power generation by a plant that can reach its full load only 0.3 to 0.5 sec after being started, and that can vary its load with a time constant of about 10 ms, were determined. As to the power supply of fusion experiments (tokamaks and stellarators), this type of MHD generator will cover quite well present and future requirements, and it may lead to an important reduction of energy costs. Concerning public and industrial grids, this MHD plant is suited to short-period reserve power production required to prevent breakdown of the grid in the event of a sudden shutdown of a unit. When this plant is only operated up to about 100 hours per year, the total power generation costs per year are considerably lower than those incurred by other reserve-power-supply methods. (auth)

47171 EXPERIMENTS ON A SUPERPOSED FIELDS THETA PINCH. Bono, R.; Costa, S.; Mondino, P. L.; Rostagni, G. (Univ. Padua). Plasma Phys.; 14: No. 5, 553-70 (May 1972).

The so-called "superposed fields" or "sustained-field" technique, as applied to a theta pinch is described, and a series of experiments performed in order to compare its characteristics with those of a simple theta-pinch are discussed. The desired magnetic field waveform is obtained by means of two coaxial coils, one fed by a steady, or quasi-steady current, the other by a pulsed one. The geometrical field shape is a function of the current ratio and maps for different conditions are shown. Discharges in hydrogen between 10 and 200 mtorr and in helium between 50 and 400 mtorr have been studied and the main results are reported. The influence of a preionizing axial discharge on plasma behavior is also discussed. (UK)

47172 THERMONUCLEAR COMBUSTION WAVES IN A BOUNDED PLASMA. Nastoyashchii, A. F.; Shevchenko, L. P. At. Energ. (USSR); 32: No. 6, 451-5 (Jun 1972). (In Russian).

Existence conditions and propagation velocity of a thermonuclear combustion wave in an infinite layer of a plasma confined by walls of heavy material are studied. Two methods are considered: 1) finding approximate analytical solutions for "cold" and "hot"

regions and joining the solutions; 2) the phase plane method. It is shown that a combustion wave occurs only in a region of densities and temperatures where the temperature derivative of the effective heat source, including the thermal effect of the fusion reaction thermal losses on the walls and plasma radiation, is negative. (auth)

47173 BREEDING RATIO, INVENTORY, AND DOUBLING TIME IN A D-T FUSION REACTOR. Vogelsang, W. F. (Univ. of Wisconsin, Madison). Nucl. Technol.; 15: No. 3, 470-4 (Sep 1972).

A simple model is proposed to investigate the relations between breeding, inventory, and doubling time in the blanket of a fusion reactor using the $T(D,n)He^4$ reaction. It is assumed that the amount of tritium removed per unit time is proportional to the amount present, and the effects of radioactive decay are included. A series of numerical calculations were made using parameters appropriate for a 5000 MW $_h$ reactor. From these calculations it appears that, to keep the blanket inventory and the initial inventory for startup to reasonable values, the tritium removal system must be designed to provide a mean tritium residence time in the blanket less than ~ 1 day, preferably with the mean residence time approaching 0.1 day. Breeding ratios in the range of 1.02 to 1.1 give acceptable doubling times and, in general, higher breeding ratios are not desirable, especially when considered from a hazards viewpoint. (auth)

47174 CONTROLLED-FUSION REACTOR. Stix, Thomas H. (to United States Atomic Energy Commission). U. S. Patent 3,663,362. 16 May 1972. Filed 22 Dec 1970.

Method and apparatus for plasma confinement and/or heating in a toroid-like magnetic system is described, wherein a dense electron-rich sheath is maintained between the surface of a toroid-like body of quasi-neutral plasma and an adjacent conducting wall or grid in order to create a deep electrostatic potential well for the ions of said quasi-neutral plasma and thus to enhance the confinement of ions in said system and/or to accelerate injected ions for the heating of said plasma. (Official Gazette)

47175 PROTON E-LAYER ASTRON FOR PRODUCING CONTROLLED FUSION REACTIONS. Christofilos, Nicholas C. (to United States Atomic Energy Commission). U. S. Patent 3,664,921. 23 May 1972. Filed 16 Oct 1969.

High-energy charged particles (molecular ions) are injected and trapped in a linear axial magnetic mirror containment field to produce a cylindrical E-layer of relativistic ions, e.g., protons rotating about the axis of said field. The magnetic field of the E-layer interacts with the mirror field to produce a pattern of closed magnetic field lines defining a charged-particle containment zone into which neutral fuel materials may be introduced to be ionized by interaction with the E-layer particles forming a high-temperature gas or plasma therein. (Official Gazette)

47176 NEUTRONICS CALCULATIONS FOR TOROIDAL THERMONUCLEAR REACTOR BLANKETS CONTAINING IRON. Smith, Jay Edward. Pittsburgh; Carnegie-Mellon Univ. (1972). 266p. University Microfilms Order No. 72-17,884.

Thesis. A method is developed for determining the true angular dependence of source neutrons entering the blanket region of a thermonuclear reactor of arbitrary shape. Application of this method is made to various specific geometries including the toroidal tokamak for which the effect of exact angular source distribution on blanket neutronics calculations is estimated by means of a synthesis technique. The feasibility and effects of including discrete amounts of iron within the blanket region are investigated for several possible reasons, including the reduction of required lithium pumping power. The neutron source treatment and synthesis technique mentioned above are used to estimate the effects of iron addition to the blanket of a tokamak reactor. (Diss. Abstr. Int., B)

47177 NUCLEAR FUSION DEVICE OF THE AIR-CORE TOKAMAK TYPE. Yoshikawa, Shoichi (to United States Atomic Energy Commission). U. S. Patent 3,663,361. 16 May 1972. Filed 17 Feb 1970.

An air-core tokamak, having high symmetry, is described for providing an increased ratio of plasma radius over the major radius, stability, high-B, and minimizing field error, as well as a low aspect ratio, high accessibility to the plasma, and separation of the plasma column from the proximity of the material of the vacuum container wall and limiters. The toroidal confinement device has an air-core toroidal solenoid disposed along the inside diameter of the plasma column for producing a toroidal electric field, and spaced apart, toroidal conductors disposed along the outside diameter of the plasma column for compensating for the

return flux of said solenoid and providing a transverse magnetic field for stably confining the plasma. (Official Gazette)

47178 ALL-METAL DISCHARGE TUBE. Phillips, James A.; Schofield, Aldred E.; Briscoe, William L. (to United States Atomic Energy Commission). U. S. Patent 3,667,059. 30 May 1972. Filed 9 Mar 1971.

In a fast linear or toroidal zeta-pinch device, a discharge tube that fits precisely into the primary winding of the device is described. The tube is a series of anodized aluminum rings stacked upon each other and held together by a nonconducting epoxy layer, which is coated with a slightly conducting layer, and then an outer structural layer. (Official Gazette)

47179 FUSION REACTORS. Hancox, R. (United Kingdom Atomic Energy Authority, Abingdon, Eng.). Aust. Elec. World; 37: No. 5, 14-20(10 May 1972).

Technological advances towards fusion reactors are described and some conceptions of future fusion systems are forwarded. The engineering of large and complex magnetic systems with average fields of 75 kG peaking to 150 kG with improvement in critical current densities may be required. Structural materials capable of withstanding high neutron doses without damage require development, and other advances that may be crucial for the success of fusion reactors are: refueling of a magnetically contained plasma, control of a reacting plasma, and removal of escaping plasma and reaction products. First-generation reactors are likely to be fueled with deuterium and tritium, which is bred in the nuclear blanket by lithium disintegration. At any time the amount of fuel present in the plasma would be sufficient for only a few seconds operation. (UK)

Wave Phenomena

Refer also to abstracts: 46313-46315, 46318, and 47118.

47180 SPONTANEOUS EMISSION OF PLASMA WAVES IN THE PRESENCE OF A FINITE-AMPLITUDE WAVE. Katayama, Yoshishige (Kyoto Univ.). J. Phys. Soc. Jap.; 33: No. 2, 483-95(Aug 1972).

The spontaneous Cherenkov emission of longitudinal waves in a plasma with small but finite amplitude wave (hereafter referred to as the carrier wave) is investigated. Two effects of the carrier wave are taken into account; one is the modification of the test-particle orbit and the other that of the dielectric character of the plasma. It is shown that particles trapped in the potential-energy troughs of the carrier wave emit a discrete spectrum of sideband waves (in addition to the carrier wave itself) whose frequencies are different from the carrier-wave frequency by approximately integral multiples of the bounce frequency. An untrapped particle also emits waves at several different frequencies, including the one which satisfies the ordinary Cherenkov condition, but these frequencies depend on the particle velocity. The results are compared with those reported in a previous paper where only the effect of the orbit modification is taken into consideration. (auth)

47181 NONLOCAL DRIFT WAVES IN A CYLINDRICAL PLASMA. Midzuno, Yukio (Nagoya Univ., Japan). J. Phys. Soc. Jap.; 33: No. 2, 502-10(Aug 1972).

Propagation characteristics of nonlocal drift waves in a cylindrical plasma are investigated on the basis of two-fluid model composed of cold ions and warm electrons. An equation is derived for describing the radial behavior of the wave amplitude. The dispersion relation is obtained from solutions of an eigenvalue problem, and the latter is proved to have a unique solution if the radial density is moderate and the mode numbers are prescribed. The eigenvalue problem is solved numerically for a density profile of a Gaussian type, for which the drift frequency ω_* can be defined unambiguously. It is found that the dispersion relation obtained in the conventional slab approximation is a good approximation to the one of a cylindrical plasma, provided that the drift frequency ω_* is much smaller than the ion cyclotron frequency. (auth)

47182 NUMERICAL SIMULATION OF THE EVOLUTION OF AN EXCITED MONOCHROMATIC WAVE IN A ONE-DIMENSIONAL MAXWELLIAN PLASMA. Lorente, Ambrosio (Univ. Paris, Orsay, France). Compt. Rend., Ser. B; 274: No. 18, 1057-60 (3 May 1972). (In French).

A one-dimensional Maxwellian electron plasma was simulated using electron packets (whose electric charge and dimension were defined from the density of the initial plasma) and a uniform cloud of stationary ions. The existence of a Landau damping of waves initially excited was established together with the nonlinear effects due to trapping. (France)

47183 KINETIC ANALYSIS OF COLLISION EFFECTS ON NONLINEARITIES RELATED TO ELECTROMAGNETIC WAVE PROPAGATION IN A WEAK DISSIPATIVE-IONIZED MEDIUM. Jancel, Raymond (Univ., Paris). Compt. Rend., Ser. B; 274: No. 18, 1061-4(3 May 1972). (In French).

A kinetic method for studying the effect of collisions on nonlinearities in weakly or partially ionized plasmas is proposed. Effects related to electromagnetic-wave propagation such as generation of harmonics and mode coupling are investigated in particular. (France)

47184 CYCLOTRON HARMONIC WAVES IN A REFLEX DISCHARGE. Brand, G. F.; Cohen, D. D. (Univ. of Sydney). Plasma Phys.; 14: No. 5, 571-4(May 1972).

An experiment to examine the dispersion relation for cyclotron harmonic waves in a reflex discharge revealed a large effect due to the nonMaxwellian electron velocity distribution. The wavelengths were several times shorter than those predicted by Bernstein's dispersion equation. (auth)

47185 NONLINEAR FOCUSING OF TM WAVES IN ISOTROPIC PLASMA. Demchenko, V. V.; Hussein, A. M. (Atomic Energy Establishment, Cairo). Plasma Phys.; 14: No. 4, 417-18(Apr 1972).

The self-focusing properties of small-amplitude waves in isotropic plasma are considered. It is shown that, if the dielectric constant varies with the wave amplitude, a waveguide channel is formed and maintained by the electromagnetic wave pressure. (UK)

47186 GENERATION OF SHOCK WAVES BY EXPLODING CURRENT SHEATH. Komel'kov, V. S.; Modzolevskii, V. I. JETP Lett. (USSR) (Engl. Transl.); 15: No. 6, 210-12(20 Mar 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 6, 299-301(20 Mar 1972).

A dense plasma was accelerated in strong magnetic fields and in short nozzles. When the magnetic piston reaches the end of the nozzle, a unique explosion-like rapid expansion of the current sheath occurs and strong shock waves are produced. Some experimental results are given. (M.O.W.)

47187 PARAMETRIC AMPLIFICATION OF ALFVEN WAVES. Lehane, J. A.; Paoloni, F. J. (Univ. of Sydney). Plasma Phys.; 14: No. 5, 461-71(May 1972).

Observation of the amplification of Alfvén waves launched into a perturbation region is reported. The effect was found to depend on the relative frequencies of the wave and the perturbation field, on the strength of the perturbation field, and on the phase of the wave entering the perturbation region. (UK)

47188 CYCLOTRON HARMONIC WAVES IN AN INHOMOGENEOUS PLASMA. I. THEORY. Derfler, H.; Leuterer, F. (Max-Planck-Institut fuer Plasmaphysik, Garching, Ger.). Plasma Phys.; 14: No. 5, 473-97(May 1972).

The Buchsbaum-Hasegawa equation describing the rf electric field within a one-dimensional magnetoplasma slab is derived from first principles. In a laboratory plasma the electron collisions are sufficiently numerous, in which case one can ignore the multiple reflection of the waves between the hybrid layers, and describe the electric field by two propagating waves alone. (UK)

47189 CYCLOTRON HARMONIC WAVES IN AN INHOMOGENEOUS PLASMA. II. EXPERIMENT. Leuterer, F. (Max-Planck-Institut fuer Plasmaphysik, Garching, Ger.). Plasma Phys.; 14: No. 5, 499-521(May 1972).

Electrostatic cyclotron harmonic waves were excited in an inhomogeneous plasma column with axial magnetic field by means of an externally applied rf capacitor field. Amplitude and phase of the rf potential inside the plasma were measured with a probe. For frequencies just above the harmonics of the electron cyclotron frequency, forward waves were found in the low density region of the plasma where the local density is less than the hybrid density. For frequencies just below the harmonics, backward waves were found in the dense core of the column. Time-of-flight measurements show that both the forward and the backward waves are excited around the hybrid layers. Multiple reflection of backward wave trains at the hybrid layers was observed. (UK)

47190 PARAMETRIC COUPLING BETWEEN ELECTRONIC AND IONIC WAVE MODES IN AN AUTO-RESONANT PLASMA. Leprince, P. (Laboratoire de Physique des Plasmas, Orsay, France). Plasma Phys.; 14: No. 5, 523-41(May 1972). (In French).

The experimental results in a low-pressure rf self-sustained plasma discharge are discussed. A parametric coupling between several electronic modes with different azimuthal wavenumbers and an ionic mode are seen to explain one of the emissions whose

frequency lies close to ionic plasma frequency. This oscillation is also observed when the exciting frequency lies close to the thermal Tonks-Datner resonances (secondary resonances) frequencies in a low-pressure glow discharge. (UK)

47191 LONGITUDINAL WAVES IN PLASMAS AND PHYSICAL MECHANISM OF LANDAU DAMPING. Lacina, J. (Inst. of Plasma Physics, Prague). *Plasma Phys.*; 14: No. 6, 605-16 (Jun 1972).

Expressions for the kinetic energy and the kinetic momentum of longitudinal waves are derived including the nonlinear motion up to second-order in the perturbation. The Landau dispersion equation is then derived from the energy-momentum conservation laws. It is shown that the effect of the electric field of a wave on the motion of particles results both in quasi-harmonic oscillations (damped or growing) and in the changed value of uniform motion. (UK)

47192 MODIFIED ORDINARY WAVE. Arumi, F. N.; Oakes, M. E. (Univ. of Texas, Austin). *Plasma Phys.*; 14: No. 6, 635-49 (Jun 1972).

The modified ordinary wave in the presence of contraststreaming beams is studied using a macroscopic model which, through the use of a pressure tensor, includes perpendicular temperature of the beams. Regions of instability are determined and growth rates given. (auth)

47193 EXPERIMENTAL STUDY OF WAVE-WAVE COUPLING PROCESSES IN A BEAM-PLASMA SYSTEM. Chang, James Shih-tsih. Urbana, Ill.; Univ. of Illinois (1972). 92p. University Microfilms Order No. 72-19,806.

Thesis.

The wave-wave coupling processes in a cold-beam cold-plasma system were investigated under conditions such that nonlinear wave-particle effects are negligible. In this experiment, a low-density mono-energetic beam in the energy range of 15 to 16 keV was premodulated at two frequencies, $f_0 = 9.5$ GHz and $f_1 = 9.4$ GHz and injected into the afterglow of a discharge in neon. The interaction between these two launched waves and their difference frequency, f_2 , was investigated and at low wave amplitudes was found to be in general agreement with the theory of the three-wave decay instability. At larger wave amplitudes, a strong remixing of the three waves occurred, giving rise to "satellite waves" at frequencies $f_0 \pm nf_2$ and $f_1 \pm nf_2$, where n is an integer. Measurements of the spatial evolution of these satellite waves and their dependence on the modulation power and difference frequency are presented. The results are discussed in the light of the theory of coupling among spatially growing waves. (Diss. Abstr. Int., B)

47194 INVESTIGATION OF LOW-DENSITY PLASMA SHOCK WAVES IN AN EXPANDING DISCHARGE. Davis, Harold Alan. San Diego, Calif.; Univ. of California (1972). 166p. University Microfilms Order No. 72-19,718.

Thesis.

Plasma waves launched into a low-density, photoionized, magnetized, argon plasma were studied. The shock waves were generated by an inverse pinch discharge, an expanding discharge as opposed to the more conventional contracting discharges normally used to study shock waves. The driving current of this device initially flows in the region near the insulator, which is transiently filled with a relatively high density (30 m torr) of neutral gas just prior to the initiation of the discharge. As this current layer expands, it acts as a piston, driving shocks into the low-density region ahead of it. The low-density region contains a uniform 2-m torr fill of neutral argon which is 30% photoionized (2×10^{13} electrons/cm³) by the intense radiation from the plasma associated with the piston. Initially, the experimental volume is filled with a uniform axial magnetic field of 500 G. The diagnostic techniques employed included Kerr-cell photography of the plasma luminosity, magnetic and floating electric probing, Thomson scattering of ruby laser light at low α to obtain both electron temperature and density, and the measurement of spectroscopic line-intensity profiles to infer ion velocities. Magnetic probe data indicated that a conventional Mach 2 shock, well separated from the driving field having a compression ratio of 2, was generated. However, the scattering measurements showed that the electron temperature was too low and the electron density rise across the shock was too high for the observed propagating magnetic field jump to be a conventional plasma shock. Subsequent line-intensity profile measurements showed that a component of the plasma was moving at shock speed. This led to the development of a new model. In this model, a beam of twice-ionized argon ions moves ahead of the piston and carries some electrons with it in order to preserve space-charge neutrality. This produces the observed electron density rise. Since the electrons are tightly coupled to the magnetic field, magnetic

flux is carried with the ion beam producing the observed rise in the magnetic field. The origin of the ion beam was investigated by calculating the trajectories of ions, starting at various radii, using the measured radial electric field. The calculation demonstrated that the fast ions originated near the insulator and were not the result of specular reflection from the piston field, whose current sheet boundary moved at exactly half shock speed. In addition, the stability of the ion beam moving through the ionized stationary background was investigated (the ion mean free path due to Coulomb collisions was very large). The beam was found to be stable. This result was compared with the theory of interpenetrating ion beams in a magnetic field, and was found to agree for the conditions of this experiment. (Diss. Abstr. Int., B)

47195 STRUCTURE OF THE PLASMA-WAVE ECHO. Goforth, Robert Ronda. Austin, Tex.; Univ. of Texas (1971). 86p. University Microfilms Order No. 72-19,591.

Thesis.

Electron plasma wave echo experiments are described in which the observed structure of the echo is in good agreement with the theoretical description from the Vlasov equation. The theory is extended to third order for comparison with the experiment. Measurements of the spatial periodicity and envelope of the third-order echo are in good agreement with the extended theory. Electron plasma wave echoes of second order were experimentally studied in a two-density plasma, and the observed structure compares favorably with theory. Because the observed structure contains information about the primary waves, these experiments demonstrate the reversibility of Landau damping. (Diss. Abstr. Int., B)

47196 STRUCTURE OF A LARGE-AMPLITUDE STATIONARY HYDROMAGNETIC DISTURBANCE IN A COLLISIONLESS PLASMA. Mahdavi, Mohamad Ali. San Diego, Calif.; Univ. of California (1972). 157p. University Microfilms Order No. 72-19,720.

Thesis.

Structure of a shock-like hydromagnetic disturbance in a collisionless plasma was studied. The current sheet of an Inverse Pinch Discharge driven into a low-density magnetized plasma was used to generate the shock wave. Argon at a static pressure of 2×10^{13} torr was used as a filling gas. A certain amount of argon gas was puffed into the discharge region prior to firing of the energy banks to ensure uniform breakdown of the gas. Preionization of the low-density ambient gas was achieved by intense uv radiation from the piston plasma (current sheet of the pinch). Diagnostic techniques used were electric and magnetic field probes, light scattering and spectroscopy. Electron densities and temperatures in the post and pre-shock regions were measured by light scattering. Streaming velocities of the ions were determined by the measurement of Doppler shift of spectral lines. Electron temperature in the post-shock plasma was also determined by the ratio of spectral line intensities. A stationary shock-like structure of velocity of 10^7 cm/sec propagates into undisturbed plasma of 2×10^{13} el/cm³ and ambient transverse magnetic field of 500 G. The experimentally measured plasma parameters do not fit the De Hoffman-Teller jump conditions. The explanation offered for this violation is presence of a beam of fast ions emerging from the magnetic piston in the upstream direction. Doppler shift of spectral lines showed presence of ions traveling at the shock velocity. The plasma associated with the piston and the ion beam was primarily double ionized. Measured electron temperature in the post-shock plasma is ≈ 3 eV. An upper bound set on the ion temperature is ≈ 300 eV. (Diss. Abstr. Int., B)

47197 ELECTRICAL SHOCK-TUBE PRECURSOR. Mills, James Ignatius Jr. Norman, Okla.; Univ. of Oklahoma (1972). 104p. University Microfilms Order No. 72-19,020.

Thesis.

Observations of luminosity and ionization far in advance of the plasma-acoustic expansion in electrically driven shock tubes have been made by many investigators, and the conflicting results and explanations of these studies have led to much confusion. The results described in this dissertation clarify the situation. Precursor waves were created and studied in helium by means of an electrically driven shock tube employing two different discharge-circuit configurations. The component causes of precursive effects were separated and identified and studied individually in detail. Parameters of basic interest such as wave speed and electron temperatures and densities were monitored, using nuclear data handling techniques and photomultipliers for the wave speeds and a revised spectral-line-ratio technique for the temperatures and densities. Velocities of the order of 10^8 to 10^9 cm/sec were recorded. Electron temperatures of 10 eV and electron densities ranging from 10^9 to 10^{10} el/cm³ were observed. Exten-

sive studies employing the use of a magnetic field were conducted in order to gain insight into the thermal conduction processes acting to transfer energy to the wave front. Many investigations of the electrical properties of the wave including the use of external calibrated probes to measure the electric field strengths associated with the front were undertaken. A discussion of the bearing this new knowledge has on explaining past studies as well as planning future studies with the electrical shock tube is included. (Diss. Abstr. Int., B)

47198 STUDY OF THE STRUCTURE OF COLLISIONLESS SHOCK WAVES PROPAGATING PERPENDICULAR TO AN INITIAL FIELD. Phillips, Perry Edward, Austin, Tex.; Univ. of Texas (1971). 137p. University Microfilms Order No. 72-19,648.

Thesis.

An experimental study of the structure of collisionless shock waves was carried out on the oblique shock apparatus at the University of Texas. A brief review of the structure of shock waves is presented, starting from ordinary gasdynamic shocks and progressing to the structure of collisionless shock waves. Calculations of the effects of various dissipative mechanisms on the shock thickness were performed. A comprehensive experimental study of the structure of low- to intermediate-strength shocks was conducted. Special emphasis was placed on understanding the double structure previously observed in shocks for low- β plasmas. The apparatus produced shocks with Alfvén Mach numbers from 2 to 6, which propagated into a preheated plasma with a density of $\sim 5 \times 10^{14} \text{ cm}^{-3}$ and equal electron and ion temperatures of 1 eV. The initial plasma is collision-dominated; however, due to heating in the shock frame, collisionless conditions are rapidly produced. A novel magnetic probe was used to examine in detail the magnetic structure of the shock waves moving perpendicular to the field. The familiar double structure, or foot, was observed above the so-called critical Alfvén Mach number $M_A^* \approx 3$. A double structure, involving a pronounced change in slope, was also observed below M_A^* . Using a two-fluid MHD model it is shown that the latter can be explained in terms of a transition from classical to collisionless resistivity within the shock front. The observed foot structure above the critical Mach number was consistent with the concept of ions being reflected from the shock front and executing gyro orbits out ahead of the shock. For verification of this hypothesis two methods of detecting the presence of reflected ions were attempted. First spectroscopic measurements of the H α line profiles were performed. For shocks with Mach numbers above M_A^* , they exhibited a significant amount of radiation out at a Doppler-shifted frequency corresponding to ions traveling at twice the shock velocities. This excess radiation was not observed for Mach numbers below M_A^* . A direct measurement of the reflected ion current was obtained by a modified ion probe. These measurements were compared with a model proposing the link between the reflected ions and the observed foot structure. Excellent agreement was achieved. A model for the reflecting mechanism is also proposed which can explain the importance of the critical Mach number and is not

in contradiction to the experimental data. From this work it can be concluded that ions are reflected from the shock front and that they are the cause of the double structure observed in shock waves for $M_A > M_A^*$. (Diss. Abstr. Int., B)

47199 INFLUENCE OF REFLECTED IONS ON THE MAGNETIC STRUCTURE OF A COLLISIONLESS SHOCK FRONT. Phillips, P. E.; Robson, A. E. (Univ. of Texas, Austin). Phys. Rev. Lett.; 29: No. 3, 154-7 (17 Jul 1972).

A special probe was used to detect ions reflected from a collisionless shock wave propagating perpendicular to the field in a magnetized, low- β plasma. Reflected ions are observed only above a critical Alfvén number $M_A \approx 2.7$, and their orbits coincide with the familiar "foot" on the magnetic structure of supercritical shocks. The origin of the reflected ions is discussed. (auth)

47200 EFFECT OF THERMAL CONDUCTIVITY ON THE STRUCTURE AND CRITICAL PARAMETERS OF SHOCK WAVES IN A PLASMA. Berezin, Yu. A.; Dudnikova, G. I. Zh. Prikl. Mekh. Tekh. Fiz.; No. 2, 8-14 (1972). (In Russian).

Results are given from a numerical solution to the problem of propagation of shock waves in a cold, rarefied plasma across a magnetic field in the presence of a finite conductivity and electronic thermal conductivity. The results are compared with those obtained neglecting thermal conductivity. At small Mach numbers ($M < 2.5$) it is possible to disregard the thermal conductivity. At Mach numbers of $2.8 \lesssim M \lesssim 3.3$, an isomagnetic discontinuity in density is obtained. An increase in the amplitude of the magnetic field at the plasma boundary leads to a dissolution of the isomagnetic discontinuity. The critical Mach number characterizing the shock wave is $M^* > 3.4$. (tr-auth)

47201 EXCITATION OF A REGULAR PLASMA WAVE BY A MODULATED BEAM WITH HIGH ENERGY DENSITY. Krasovitskii, V. B. (All-Union Research, Design, and Technological Inst., Moscow). JETP Lett. (USSR) (Engl. Transl.); 15: No. 6, 244-7 (20 Mar 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 6, 346-9 (20 Mar 1972).

Investigations were made of the interaction of a relativistic beam with a nonlinear plasma, which point to the possibility of synchronism between the beam and the wave during the nonlinear stage of instability development if the beam and plasma parameters are so chosen that the beam velocity and the phase velocity of the wave decrease with time in accordance with an identical law. In this case the energy transferred from the beam to the field greatly exceeds the value obtained earlier. (M.O.W.)

47202 SOLITONS IN A TWO STREAM PLASMA. Alikhanov, S. G.; Klichigin, G. N.; Konkashbaev, I. K. (Inst. of Nuclear Physics, Novosibirsk, USSR). Plasma Phys.; 14: No. 6, 661-4 (Jun 1972).

The propagation of solitons in a nonisothermal collisionless plasma in the absence of a magnetic field was investigated, in a system consisting of several inter-penetrating ion streams. (UK)

PHYSICS (RADIATION AND SHIELDING)

Radiation Physics

Refer also to abstracts: 45604, 46359, 46365, 46448, 46854, and 47238.

47203 (N-72-18797) RADIATION DOSIMETRY FOR THE GEMINI PROGRAM. Richmond, Robert G. (National Aeronautics and Space Administration, Moffett Field, Calif. Ames Research Center). Mar 1972. 16p. (NASA-TN-D-6695; MSC-S-222). NTIS.

The principal source of radiation for low-earth-orbit, low inclination space flights is in the area of the South Atlantic magnetic anomaly. None of the Gemini dose measurements reported are of high-enough intensity to be considered hazardous. There is a trend toward larger doses as missions are flown higher and longer. Extended orbital operations between 1400 and 4400 km would encounter high interior radiation levels. Pronounced spacecraft geometry effects were measured in manned spacecraft. Instrumentation for radiation measurements on Gemini spacecraft is described. (auth) (STAR)

47204 INVESTIGATIONS ON ABSOLUTE DIFFERENTIAL SCATTERING CROSS SECTIONS OF 0.662 MeV PHOTONS. Thein Aung (Arts and Science Univ., Rangoon); Thein Htut, Hla Pe; (and others). Union Burma J. Sci. Technol.; 3: No. 1, 245-50 (Apr 1970).

Absolute values of the differential scattering cross sections for ^{137}Cs (0.662-MeV) photons were measured in the angular range of 15° to 120° . A uniform-sensitivity photon counter suitable for low-energy photons was used to measure the relative values of the cross sections. The cross sections were then converted to absolute values by measuring incomplete scattering cross section. The experimental values, when corrected for coherent scattering, show good agreement with the Klein-Nishina formula. (auth)

47205 TRANSMISSION OF LOW-ENERGY X-RAY THROUGH VARIOUS LOCALLY AVAILABLE MATERIALS. Lee, H. (Ateneo Univ., Manila); Terrado, J.; Petel, R.; Bustamante, N. Philipp. Nucl. J.; 2: No. 2, 215-22 (1970).

The transmission of low-energy x rays through various materials was determined using a Victoreen condenser roentgen meter. Initial calibration of the x-ray unit was done to determine the characteristics of the equipment. The method of least squares was used to obtain a good curve fit on the transmission measurement data. The linear attenuation coefficient of the various materials as a function of the energy of the x-ray beams were determined. (auth)

47206 EFFECTIVE ATOMIC NUMBERS FOR PHOTOELECTRIC PROCESS IN VARIOUS SHELLS IN ALLOYS. Narasimha Raju, M. L.; Chandran, C. R.; Parthasaradhi, K.; Ramamurty, S. (Andhra Univ., Waltair, India). Indian J. Phys.; 45: No. 2, 92-3 (Feb 1971).

The effective atomic numbers for various shell effects in three typical alloys: solder (Pb, 67%; Sn, 33%), plumber solder (Pb, 50%; Sn, 50%), and soft solder (Pb, 40%; Sn, 60%) are estimated using procedures described previously and making use of theoretical shell cross sections. It was shown that effective atomic numbers, within the range of errors, can be defined uniquely irrespective of shells and incident photon energy in all the alloys. (M.H.R.)

47207 EFFECTIVE ATOMIC NUMBERS FOR DIFFERENTIAL PROCESSES OF GAMMA-RAY INTERACTIONS. Siddappa, K.; Parthasaradhi, K.; Rama Rao, J. (Andhra Univ., Waltair, India). Indian J. Phys.; 45: No. 4, 184-6 (Apr 1971).

Systematic investigations were made on differential effects in photoelectric incoherent and coherent scattering processes. The available theoretical cross sections were used, since there is no adequate accurate experimental data on the differential effects. (M.H.R.)

47208 RANDOM METHOD FOR BOLTZMANN'S EQUATION. Patraulea, N. N. Stud. Cercet. Mec. Apl.; 30: No. 6, 1281-5 (1971). (In Rumanian).

A random method is presented for the solution of the basic equation of rarefied gases. The method is equally applicable to elastic collisions and to any other models of collision laws. Random evaluations of the collision are made over the 8-dimensional interval $(\bar{v}_1, \bar{v}_2, \bar{P})$ connected to each point of a finite network in

the physical space. Satisfactory convergence is mainly based on the lower dispersion in consequence of the selection of the random velocities $\bar{v}_1, \bar{v}_2$ of the waves under consideration with probability densities $t/\int f d\bar{v}$. (auth) (Rom. Sci. Abstr.)

47209 FLUORESCENCE PHOTON AND AUGER ELECTRON SPECTRA. Charlton, D. E. (Sir George Williams Univ., Montreal). Radiat. Res.; 50: No. 3, 455-63 (Jun 1972).

Empirical formulas giving the fluorescence photon and Auger electron spectra following electron capture and internal conversion as a function of atomic number are presented. Processes originating in the K and L shells only are considered, and these account for approximately 95% of the total energy escaping from the atom. It is suggested that the remaining 5% is probably carried by Auger electrons originating in the M and N shells. The fluorescence photon and Auger electron spectra of tellurium atoms produced by the decay of ^{125}I are evaluated and compared to those of other authors. (auth)

47210 LOW-ENERGY ELECTRON TRANSPORT BY THE METHOD OF DISCRETE ORDINATES. Bartine, David Elliott. Rolla, Mo.; Univ. of Missouri (1971). 108p. University Microfilms Order No. 72-18,186.

Thesis.

The one-dimensional discrete-ordinates code ANISN was adapted to simulate the transport of low-energy (on the order of a few MeV) electrons. Two different calculational techniques were utilized for the treatment of electron-electron collisions that result in a small energy transfer. One method treats such collisions by a continuous slowing-down approximation, while the other method treats these collisions by the use of a very approximate cross section. Calculated results obtained with ANISN are compared with experimental data for the transmitted energy and angular distributions for 1-, 2.5-, 4-, and 8-MeV electrons normally incident on aluminum slabs of various thicknesses and for 1-MeV electrons normally incident on a gold slab. The calculated and experimental results are in reasonably good agreement for the aluminum slabs but are in poor agreement for the gold slab. Calculated results obtained with ANISN are also compared with calculated results obtained with Monte Carlo methods. (Diss. Abstr. Int., B)

47211 DISCRETE-ORDINATES METHODS FOR THE NUMERICAL SOLUTION OF THE TRANSPORT EQUATION. Lathrop, K. D. (Los Alamos Scientific Lab., N. Mex.). Reactor Technol.; 15: No. 2, 107-35 (1972).

The theory, advantages, disadvantages, and capabilities of the discrete-ordinates method for numerical solution of the Boltzmann transport equation are reviewed. To illustrate the simplicity of the method, the author derives the theory directly for the specific case of one-dimensional spherical geometry and thereby provides a clearer picture of the method than is available in more general expositions. On this foundation the essential generalizations of the techniques are made. The major limitations of the method, including orthogonal geometries, negative fluxes, ray effects, convergence requirements, and data storage are discussed in detail, and current work to remedy these limitations is reviewed. The capabilities of the discrete-ordinates method are surveyed at length, and numerical programs applying the discrete-ordinates techniques are tabulated and compared. An ancestral chart of current discrete-ordinates programs is provided. A general guide to the usage of discrete-ordinates programs is given, and the application of the method is illustrated in examples including a shielding bench-mark problem, a reactor core-design problem, a control-rod-worth and radiation heating study, a pulsed-neutron problem, and a shielding parameter survey. (66 references) (auth)

47212 CALCULATION OF THE SPECTRA OF ELECTRON ENERGY LOSSES IN GOLD FROM 1 TO 12 eV. COMPARISON WITH THE SPECTRA OBSERVED AT 35 keV. Wehenkel, Claude (Univ., Paris). Compt. Rend., Ser. B; 274: No. 15, 939-42 (10 Apr 1972). (In French).

The electron energy loss spectra were calculated from experimental optical constants taking into account propagation effects. The effects of the zero energy loss line and other parameters are discussed. The calculation was compared with experimental spectra obtained on the transmission by a gold layer of known thickness. The agreement is better for the positions of the lines than for their relative intensities. (France)

47213 METHOD OF DETERMINING THE MEAN ENERGY OF SECONDARY ELECTRONS IN ELECTRON EQUILIBRIUM

REGIONS USING THEORETICAL X-RAY SPECTRA (10 TO 20 MV) AND IONIZATION MEASUREMENTS IN DIFFERENT MEDIA. Guille, B.; Martin, P. N.; Bernard, M. (Univ., Limoges, France). *RadioProtection*; 7: No. 1, 45-51(1972). (In French).

In an x or γ radiation field, the absorbed dose at every point in matter is linked to the ionization produced in an air cavity and to the stopping power relative to the secondary electrons. When particular electron equilibrium conditions are set up, this dose depends also upon the spectral distribution of the incident radiation, the attenuation of the various media traversed, and the coefficient of energy transfer. These two relations enable the mean energy of the secondary electrons resulting from ionization phenomena to be determined. (France)

47214 RADIATIONNAYA FIZIKA KRISTALLOV I p-n-PEREKHODOV. (Radiation Physics of Crystals and p-n Junctions). Sirota, N. N. (ed.). Minsk: Izdatel'stvo Nauka i Tekhnika (1972). 168p.

Problems in radiation physics of solids, associated with channeling of charged particles of relatively high energy, in crystals of semiconductors are examined in this collection of 21 articles by different authors. Eight papers are concerned with the interaction of radiation with Si and Ge p-n junctions, and the effects of temperature, annealing behavior, and the spectral composition of neutrons. Seven additional papers focus attention on the effects of radiations (fast neutrons, ^{60}Co gamma rays, Si^+ ions, x rays) on chemical compounds such as GaSb, CdS, Si, ferrites, alkali halides, KMnO_4 , α -alanine, glycine, and triglycinesulfate. (K.S.W.)

47215 GAMMA QUANTA DISTRIBUTION AT LARGE DISTANCES FROM THE SOURCES. Nikolaishvili, Sh. S.; Gabrashvili, A. I.; Dzhgarkava, N. N.; Iordaniashvili, E. A. *At. Energ. (USSR)*; 32: No. 6, 504-5(Jun 1972). (In Russian).

A generalized method is outlined for calculating the distribution of γ quanta and neutrons at arbitrarily large distances from the sources of the original radiation. The generalization touches upon that portion of the method in which the problem is solved with the help of the final number or original momenta. (K.S.W.)

Radiation Protection Standards

47216 NATIONAL STANDARDS FOR PHOTON DOSIMETRY. Guitho, J. P. (CEN de Saclay, Gif-sur-Yvette, France). *Radio-protection*; 7: No. 1, 3-12(1972). (In French).

A coherent photon dosimetry system of national references was established. Gamma sources (^{137}Cs and ^{60}Co) were used to produce reference photon beams, their decay/time characteristics being determined. The dimensions of the radiation source, the characteristics of the collimator, and the environment were studied. In particular, the effect of the various geometric parameters of the collimator on the spectral distribution of the incident photons was investigated. The exposure due to the photon beams was determined; their energy fluence is currently being investigated. The references for the absorbed dose rates will be defined. (France)

Radiation Source Calibration and Standardization

Refer also to abstract: 45799.

47217 STANDARD BLACK BODY AT 10,000°K (HIGH PRESSURE PLASMA ARC). Peyturaux, R.; Goldbach, C.; Nollez, G. (CNRS, Paris). pp 65-75 of *Proceedings of the Meeting on Metrology*. Paris: Bureau National de la Metrologie (1971). (In French). (CONF-701151-1).

From Meeting on metrology; Paris, France (17 Nov 1970).

Research was conducted to produce a radiation standard corresponding to temperatures in excess of 5000°K. Theoretical studies postulate that a 10-cm argon plasma subjected to a pressure of 100 bars and brought to a temperature of 10,000°K would radiate as a black body. In order to achieve such conditions, a stabilized arc was developed involving a 5-cm-long plasma operating under a pressure of 50 atm. At atmospheric pressure, the electronic temperature is 1600°K (for dissipated 16 kW). Although the determination of the electron temperature poses problems at high pressures (reabsorption of radiation), it appears that the electron temperature decreases with pressure (for constant dissipated power) while the brightness temperature increases and tends to remain constant over a large spectral range. (France)

47218 EXPOSURE MEASUREMENTS AND SPECTROMETRY IN THE MEDIUM ENERGY X-RAY REGION. Amat, G.; Am-

broisi, M. (Laboratoire Central de l'Armement, Arcueil, France). pp 181-8 of *Proceedings of the Meeting on Metrology*. Paris: Bureau National de la Metrologie (1971). (In French). (CONF-701151-2).

From Meeting on metrology; Paris, France (17 Nov 1970).

Exposure measurements consist of measuring the total charge (of a particular sign) collected as a result of the secondary particles liberated from interactions between photons and a given mass of air. This ionization is measured using a free-air ionization chamber which defines, perfectly, an interaction volume for the photons. In the usual x-ray apparatus, this chamber is the primary instrument employed to measure exposure rates, cavity chambers being used for the transfer of standards. The spectrometry of beams differs from the γ spectrometry of sources in three aspects: the energy distribution of the photon fluence is no longer discrete, but continuous and does not require very high resolution. A very large number of photons penetrate into the spectrometer and the possibility of deforming the initial spectrum must be reduced by diaphragms. The photons penetrate the detector as narrow parallel beams leading to a high detection efficiency. (France)

47219 METROLOGY OF IONIZING RADIATIONS. Le Gallic, Y. (CEN de Saclay, Gif-sur-Yvette, France). pp 175-80 of *Proceedings of the Meeting on Metrology*. Paris: Bureau National de la Metrologie (1971). (In French). (CONF-701151-3).

From Meeting on metrology; Paris, France (17 Nov 1970).

The metrology of ionizing radiations utilizes three reference units: the Curie, the Roentgen, and the Rad, which define two types of measurement: the metrology of activities, and the dosimetry of radiations. The accuracy obtained in this type of metrology is distinct from that obtained in other metrological disciplines, an accuracy of 10^{-4} being currently sought. It is recalled that the essential factors necessary for an improvement are: purity of the radioisotope; the development of new measurement techniques; knowledge of fundamental parameters; development of national reference standards. Although considerable work has yet to be accomplished, the results already obtained are very encouraging. (France)

Shielding Calculations and Theory

47220 (ORNL-TM-3867) SDT1. IRON BROOMSTICK EXPERIMENT. Maerker, R. E. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 20p. (ENDF-166). Dep. NTIS.

The experimental and calculational details for a CSEWG integral data testing shielding experiment are presented. The experiment was designed to test a given set of neutron total cross sections for iron in the range 0.8 to 11 MeV. A FORTRAN package is available to perform all the manipulation described. (M.H.R.)

47221 (ORNL-TM-3868) SDT2. OXYGEN BROOMSTICK EXPERIMENT. Maerker, R. E. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 18p. (ENDF-167). Dep. NTIS.

The experimental and calculational details for a CSEWG integral data testing shielding experiment are presented. The experiment was designed to test a given set of neutron total cross sections for oxygen in the range 1.9 to 8.6 MeV. A FORTRAN package is available to perform all the manipulation described. (M.H.R.)

47222 (ORNL-TM-3869) SDT3. NITROGEN BROOMSTICK EXPERIMENT. Maerker, R. E. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 18p. (ENDF-168). Dep. NTIS.

The experimental and calculational details for a CSEWG integral data testing shielding experiment are presented. The experiment was designed to test a given set of neutron total cross sections for nitrogen in the range 0.8 to 10 MeV. A FORTRAN package is available to perform all the manipulation described. (M.H.R.)

47223 (ORNL-TM-3870) SDT4. SODIUM BROOMSTICK EXPERIMENT. Maerker, R. E. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 18p. (ENDF-169). Dep. NTIS.

The experimental and calculational details for a CSEWG integral data testing shielding experiment are presented. The experiment was designed to test a given set of neutron total cross sections for sodium in the range 0.8 to 11 MeV. A FORTRAN package is available to perform all the manipulation described. (M.H.R.)

47224 (ORNL-TM-3871) SDT5. STAINLESS-STEEL BROOMSTICK EXPERIMENT. Maerker, R. E. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 26p. (ENDF-170). Dep. NTIS.

The experimental and calculational details for a CSEWG inte-

gral data testing shielding experiment are presented. The experiment was designed to test a given set of neutron total cross sections for a type-310 stainless steel in the range 1.2 to 11 MeV. A FORTRAN package is available to perform much of the manipulation described. (M.H.R.)

PHYSICS (SOLID-STATE)

Radiation Effects

Refer also to abstracts: 45664, 45668, 45671, 46076, 46550, and 47214.

47225 (AWRE-O-31/72) SOME MEASUREMENTS OF THE ELECTRICAL CONDUCTIVITY INDUCED IN POLYMERS BY PULSED X-RAY IRRADIATION. Barry, J. F.; Culliford, E. R. (Atomic Weapons Research Establishment, Aldermaston (England)). May 1972. 22p. Dep. NTIS (U. S. Sales Only). UK 50 pence.

The electrical conductivity induced by pulsed x-ray irradiation was measured for the materials Mylar, polythene, and PTFE. Dose rates in the range 10^7 to 10^{11} rad/sec at x-ray energies up to 4.5 MeV were used. For a given material, the results exhibit considerable scatter from sample to sample. This is attributed to minor variations in the electrode structure and in the composition of the materials, their effects being enhanced at the low conductivity levels involved. The results indicate that the conductivity varies linearly with dose rate, no strong evidence being found for saturation effects up to the maximum dose rate of 10^{11} rad/sec. (auth)

47226 RADIATION-INDUCED CENTERS OF PARAMAGNETIC AND OPTICAL ABSORPTION IN GLASSES OF THE SYSTEMS SrO-SiO_2 AND BaO-SiO_2 . Tyul'kin, V. A. Inorg. Mater. (USSR) (Engl. Transl.); 8: No. 1, 131-4 (Jan 1972).

Translated from *Izv. Akad. Nauk SSSR, Neorg. Mater.*; 8: No. 1, 152-5 (Jan 1972).

The EPR and electron absorption spectra of gamma-irradiated glasses from the systems SrO-SiO_2 and BaO-SiO_2 were studied. The data are compared with the results of previous investigations of glasses of the systems MgO-SiO_2 and CaO-SiO_2 . Regardless of the type of MO oxide, identical radiation centers arose in glasses in the system in question; however, the oxide concentration range in which a given variety of center predominates depends on the type of MO. The influence of concentration and M^{2+} cation type on the EPR line half-widths is discussed. Broadening of the lines was mainly due to dipole-dipole interaction between holes (or electrons) and the magnetic moment of the nucleus of a M^{2+} ion in the first coordination sphere. (auth)

47227 PROPERTIES OF RUBY LASER RADIATION UNDER THE ACTION OF ^{60}Co GAMMA RADIATION. Arifov, U. A.; Bedilov, M. R.; Khaidarov, K.; Egamov, U. (Inst. of Nuclear Physics, Tashkent). Dokl. Akad. Nauk SSSR; 203: No. 1, 68-70 (1 Mar 1972). (In Russian).

It is known that small doses ($<10^4$ rad) for γ radiation increases the efficiency of a ruby laser by 300%. A study was made of the effect of ^{60}Co γ radiation in doses of 0 to 10^4 rad on the properties of the stimulated radiation of a ruby laser, utilizing photoelectric, calorimetric, spectral, and photographic methods. It was found that doses 10^3 rad and greater decreased the output of the laser, altered the absorption spectrum of the ruby itself, and changed the distribution of radiation with respect to the ruby cross section and the duration of pulse. (K.S.W.)

47228 ION CHANNELING. Dearnaley, Geoffrey (United Kingdom Atomic Energy Research Group, Harwell, Eng.). Sciences (Paris); No. 76, 39-47 (Jan 1972). (In French).

Developments that have occurred since the experimental discovery of the channeling phenomenon are reviewed. The phenomenon is briefly defined and its consequences are examined (ion implantation and decreases in the yield of nuclear reactions). Possible applications are envisaged. (France)

47229 ELECTRON IRRADIATION EFFECTS IN ALUMINUM SINGLE CRYSTALS. Kirkland, Leland Rose Jr. Clemson, S. C.; Clemson Univ. (1971). 122p. University Microfilms Order No. 72-20,781.

The work on Al single crystals was undertaken to determine the relationship between the I_B recovery and the (100) directions. Information about this relationship was obtained by observing how the I_B recovery depends upon irradiation energy, sample thickness, and sample crystallographic direction parallel to the irradiation beam. Experimentally, the defects were detected by changes in the electrical resistivity of the samples. The dependence of the residual electrical resistivity upon sample size is determined from preirradiation measurements on the samples. Between measurements, the sample thicknesses were reduced by using a chemical

polish. The theory of Dingle is adapted to analyze the data. The results give good agreement with the theory of the size effect and show that the electron mean free path in the present samples is 0.0134 cm. The product of the bulk resistivity and this mean free path is $7.6 \times 10^{-12} \Omega \text{ cm}^2$. Damage rate studies give results that are in agreement with previous work. The damage rate is not sensitive to the crystallographic direction parallel to the irradiation beam for recoil energies below 25 eV. At higher recoil energies, the damage rate does have a slight crystallographic dependence. The analysis of the I_B recovery confirms the assignment of the I_B substage to the recovery of close-pair defects. Comparison of experiment to first-order recovery theory shows that I_B defects have an activation energy of 0.042 eV. Annealing data show that the magnitude of the I_B recovery is sensitive to irradiation energy, sample thickness, and sample crystallographic direction parallel to the irradiation beam. A theory is derived for the creation of the I_B defects based on the fact that these defects are caused by atoms recoiling with low energy along (100) directions. An essential assumption of this theory is that the number of I_B defects is proportional to the electron fluence capable of causing low energy atomic recoils along a (100) direction. Comparison of theory to experimental data indicates that I_B defects are created when atoms are displaced with energies in the interval from 16 to 21 eV and recoil within 16° of a (100) direction. (TSS) (Diss. Abstr. Int., B)

47230 RECOMBINATION DOMAIN INSTABILITY IN HIGH-ENERGY ELECTRON-IRRADIATED GERMANIUM. Wada, Takao (Mie Univ., Tsu, Japan); Fukuoka, Yutaka; Arizumi, Tetsuya. Jap. J. Appl. Phys.; 11: No. 7, 1009-15 (Jul 1972).

When a bar of n-type germanium (1 to $2 \Omega\text{-cm}$) is irradiated with a total flux of about 10^{15} to 10^{16} electrons/ cm^2 at 8.5 MeV, domains of high electric field in the irradiated crystal appear at a high field above 350 V/cm and 77 K. At sufficiently high electric fields, current-voltage characteristics, estimated by observing an instantaneous current density and a time constant associated with the trapping of hot electrons at defect centers, have a voltage-controlled differential negative resistance region; thus, the high-field domain is formed by the buildup of the space-charge boundaries. The velocity of the moving domain ranged from 10^5 cm/sec to 2×10^6 cm/sec, depending upon the light intensity. (8 figures, 1 table) (auth)

47231 COMPARISON OF LATTICE PARAMETER AND RESISTIVITY CHANGE IN ELECTRON-IRRADIATED COPPER. Dworschak, F.; Wagner, H.; Wombacher, P. (Kernforschungsanlage, Juelich, Ger.). Phys. Status Solidi (b); 52: No. 1, 103-9 (1 Jul 1972).

Lattice parameter and residual resistivity measurements were performed simultaneously on high-purity copper samples irradiated with 2.6-MeV electrons at 6°K up to defect concentrations of about 0.1%. Lattice-parameter changes could be measured with a resolution of $\Delta a/a = 5 \times 10^{-6}$. In six irradiations, the ratio $\eta = (\Delta a/a)/\Delta \rho$ was determined as a function of defect concentration and thermal recovery. During defect production, the ratio η amounts to $(1.75 \pm 0.10) \times 10^3 (\Omega\text{cm})^{-1}$ and is independent of the defect concentration. Furthermore, the ratio η stays constant throughout the recovery stages I, II, and III. (auth)

47232 OPTICAL WAVEGUIDING IN PROTON-IMPLANTED GaAs. Garmire, E. (California Inst. of Tech., Pasadena); Stoll, H.; Yariv, A.; Hunsperger, R. G. Appl. Phys. Lett.; 21: No. 3, 87-8 (1 Aug 1972).

Optical waveguides were produced in n-type GaAs by implantation with 300-keV protons. The guiding is shown to be due to the elimination of charge carriers from the implanted region. Annealing of the waveguide leads to very large reductions in the $1.15\text{-}\mu$ guided-wave absorption. (auth)

47233 PARAMAGNETIC HOLE CENTERS IN CeO_2 . Wagner, G. R.; Murphy, J. (Westinghouse Research Labs., Pittsburgh). Phys. Rev., B; 6: No. 5, 1638-44 (1 Sep 1972).

ESR spectra of three hole centers were observed in nominally pure crystals of CeO_2 following uv irradiation at 77°K . Designated V_I , V_{II} , and V_{III} , the centers exhibit orthorhombic symmetry with principal axes (x, y, z) parallel to [110], [001], and [110], respectively. The g values are: $g_x(I) = 2.0311$, $g_y(I) = 2.0065$, $g_z(I) = 2.0330$; $g_x(II) = 2.0175$, $g_y(II) = 2.0054$, $g_z(II) = 2.0317$; and $g_x(III) = 2.0268$, $g_y(III) = 2.0050$, $g_z(III) = 2.0415$. The V_{III} center exhibits a hyperfine interaction with a nuclear spin $I = 1/2$ for which $A_x = 3.1$, $A_y = 2.3$, and $A_z = 2.7$ in units of 10^{-4} cm^{-1} . The thermal bleaching of the centers was correlated with the thermolumines-

cence of the same crystals measured by Land and Rodine. Models proposed for these centers involve an O_2^- molecular ion near a stabilizing impurity ion. (auth)

47234 CHANNELING OF PROTONS IN THIN $BaTiO_3$ CRYSTALS AT TEMPERATURES ABOVE AND BELOW THE FERRO-ELECTRIC CURIE POINT. Gemmell, D. S.; Mikkelsen, R. C. (Argonne National Lab., Ill.). Phys. Rev., B; 6: No. 5, 1613-35 (1 Sep 1972).

The channeling of 3.8-MeV protons in thin (3 to 20- μ -thick) $BaTiO_3$ crystals was studied at 140°C (i.e., in the cubic paraelectric phase) and at room temperature (i.e., in the tetragonal ferroelectrically polarized phase). The influence of channeling on backscattering yields, on characteristic x-ray production, and on the transmitted protons, was determined. For the purpose of fitting the data for the cubic phase, the formulas that Barrett found to fit a large range of data on channeling in monatomic crystals were extended to the polyatomic case. The quality of the fit thus obtained is fair, but not as good as those obtained in monatomic lattices. Data are also presented on dechanneling and on the influence of radiation damage on the channeling properties of $BaTiO_3$. In the polarized tetragonal phase, it is found that the channeling properties are strongly affected by the relative orientations of the crystal planes and the polarization of the crystal. The effect of polarization is attributed to the associated high internal electric fields. (auth)

47235 DISTRIBUTION OF ALKALI IONS IN NICKEL BY IONIC BOMBARDMENT. Kasymov, A. Kh.; Pugacheva, T. S.; Safarov, A. (Tashkent Polytechnic Inst.). Izv. Akad. Nauk Uzb. SSR, Ser. Fiz.-Mat Nauk; No. 6, 41-6(1971). (In Russian).

The technique of alloying a material by the use of ion bombardment was further investigated by calculations with beams of Li, Na, K, Rb, Cs, and Ba ions having energies of from 1 to 10 keV incident on polycrystalline nickel. A threshold displacement energy of 25 eV was assumed. The extent of penetration of light ions was, in general, greater than that of heavy ions. The coefficient of reflection weakly decreases with increase in the energy of the incident ions, and increases with smaller ion masses. (K.S.W.)

47236 MASS DEPENDENCE OF CRITICAL AMORPHIZING DOSE IN ION IMPLANTATION. Furukawa, Seiji; Ishiura, Hiroshi (Tokyo Inst. of Tech.). Jap. J. Appl. Phys.; 11: No. 7, 1062-3(Jul 1972).

Equations are derived for the critical dose to produce an amorphous layer: $\phi_c = 4(\Delta R_p)NE_d/\xi E_c$. Calculations were performed for Si targets and compared with the experimental mass dependence for ions from H to Bi. (D.L.C.)

47237 ELECTRICAL PROPERTIES OF n-TYPE SI LAYERS DOPED WITH PROTON BOMBARDMENT INDUCED SHALLOW DONORS. Ohmura, Y.; Zohta, Y.; Kanazawa, M. (Tokyo Shibaura Electric Co. Ltd., Kawasaki, Japan). Solid State Commun.; 11: No. 1, 263-6(1 Jul 1972).

Sheet Hall coefficients and resistivities of n-type Si layers doped with shallow donors produced by proton bombardment were measured between 35 and 300°K. The donor ionization energy is (26 ± 1) MeV. The donor concentration profile was determined. (auth)

47238 ANGULAR DEPENDENCE OF THE EROSION RATE AND OF SPECIFIC ENERGY LOSSES OF 5.6 keV NOBLE GAS IONS ON GLASSES. Bach, H. (Jenaer Glaswerk Schott and Gen., Mainz). J. Mass Spectrom. Ion Phys.; 9: No. 3, 247-52 (Aug 1972). (In German).

With an ion-beam apparatus, the angular dependence of the sputtering yields, specific energy losses, and luminescence were measured for glasses and thin oxide layers. According to the results obtained, the angular dependence of the sputtering yield is not always mainly due to the orientation of the impulse vector. It is also influenced by the elastic reflection of a considerable number of those ions that do not contribute to the yield. This number increases for lower angles of incidence if the ratio $\mu = M_2/M_1$ of the effective mass numbers of the glass components M_2 and the ions M_1 is increased. (auth)

47239 FIELD-ION MICROSCOPE STUDY OF ION-IMPLANTATION IN IRIIDIUM. I. PHILOSOPHY AND PRELIMINARY CONSIDERATIONS. O'Connor, G. P.; Ralph, B. (Cambridge Univ., Eng.). Phil. Mag.; 26: No. 1, 113-28(Jul 1972).

The field-ion microscope was used to investigate some of the processes occurring during ion implantation, in particular, the fate of the implanted species. Attention is therefore paid to point defect contrast in field-ion images to establish the validity of counting such defects. The importance of bright spots is emphasized

since they may represent the only type of contrast which could lead to unambiguous detection of isolated implanted atoms. Since self-interstitial atoms released from traps by the imaging process contribute to the bright spot concentration, consideration is given to the imaging mechanism and to means of distinguishing between bright spots caused by different defects. It is shown that, in the case of oxygen implants, such a distinction is possible, and further, that the implanted species is trapped, possibly as part of a "complex." Control experiments to ensure that real effects are observed, and errors in defect counts are briefly mentioned and it is concluded that significant data can be extracted from field-ion micrographs provided care is taken in selecting a suitable implantation. (auth)

47240 FIELD-ION MICROSCOPE STUDY OF ION-IMPLANTATION IN IRIIDIUM. II. RESULTS AND DISCUSSION. O'Connor, G. P.; Ralph, B. (Cambridge Univ., Eng.). Phil. Mag.; 26: No. 1, 129-42(Jul 1972).

The methods of data analysis described in Part I are applied to a determination of the damage and implant profiles for iridium specimens bombarded with keV ions of carbon, oxygen, xenon, silver, and iron. It is shown that bright spot counting is a valid technique and can, in suitable circumstances, lead to a direct derivation of the implant profile. The damage profile in all cases is positioned deeper into the specimen than the implant profile and, since the displacement between them increases with decreasing temperature of irradiation, it is concluded that focused replacement sequences provide the main method for the transport of energy. The effects of recoil implantation of surface contaminant species are observed directly, and it is shown that the concentration of impurity species has an important effect on the annealing behaviour of ion-implanted specimens. (auth)

47241 DIRECT OBSERVATION OF GRAIN BOUNDARY DISLOCATION CLIMB IN ION-IRRADIATED GOLD BICRYSTALS. Komem, Y.; Petroff, P.; Balluffi, R. W. (Cornell Univ., Ithaca, N. Y.). Phil. Mag.; 26: No. 1, 239-52(Jul 1972).

Thin-film gold bicrystals containing (001) low-angle twist boundaries, with twist angles in the range $\theta = 7$ to 14° , and high-angle twist boundaries with $\theta \approx 37^\circ$, were prepared by welding evaporated films together face-to-face. All the boundaries contained regular orthogonal arrays of intrinsic screw grain boundary dislocations (GBD's) as well as more randomly occurring extrinsic edge GBD's having Burgers vectors essentially perpendicular to the boundary. The specimens were irradiated on one face with 20-keV gold ions producing excess self-interstitial atoms which diffused to the extrinsic GBD's, causing them to climb in the boundary. The extrinsic GBD climb, in turn, caused the boundaries to act as overall interstitial sinks. The climb motion was observed directly by transmission electron microscopy. The intrinsic screw GBD's in the high-angle boundaries (and probably also in the low-angle boundaries) remained unaffected during the process, but possibly they acted as short circuiting pipes which delivered interstitials to the extrinsic GBD's. It is concluded generally that both low- and high-angle boundaries may act as point defect sinks (sources) by means of the climb of GBD's having Burgers vectors with components normal to the boundary plane. This process is actually a form of internal crystal growth (dissolution) with the GBD's acting as "internal crystal ledges." (auth)

47242 ELECTRON STIMULATED DESORPTION OF CARBON MONOXIDE ADSORBED ON NIOBIUM (111). Davis, Paul Richard; Pullman, Wash.; Washington State Univ. (1972). 56p. University Microfilms Order No. 72-18,510.

Thesis.

Electron-stimulated desorption of carbon monoxide adsorbed on the (111) crystal face of niobium at 150°K was found to produce both O^+ and CO^+ ions. The CO^+ -producing state was desorbed completely at temperatures above 250°K. At temperatures above 700°K, the O^+ ion current completely disappeared, due to dissociation of the adsorbed molecules. Threshold energies for formation of O^+ and CO^+ ions were measured to be 19.0 ± 0.5 eV and 10.0 ± 0.5 eV respectively. The cross section for CO^+ production by 100-eV electrons was found to be $(9 \pm 2) \times 10^{-20}$ cm². (Diss. Abstr. Int., B)

47243 DETERMINATION OF THE SPATIAL DISTRIBUTION OF DEEP CENTERS FROM CAPACITANCE MEASUREMENTS OF PN JUNCTIONS. Zohta, Yasuhito; Ohmura, Yamichi (Tokyo Shibaura Electric Co. Ltd., Kawasaki, Japan). Appl. Phys. Lett.; 21: No. 3, 117-19(1 Aug 1972).

Effects of a spatial distribution of deep centers on the junction capacitance are reported for the first time. It is shown that the spatial distribution of deep centers can be calculated from capacitance-voltage method and the Copeland method, if their energy

level is known or assumed. The theory is examined with the experiment on silicon p-n junctions irradiated with 300-kV protons. The analysis yields a value 0.40 eV below conduction band for the energy level of radiation-induced defects, as well as a distribution of the defects with depth. (auth)

47244 INTERACTION OF A RUBY WITH IONIZING RADIATION. Ermolaev, E. A.; Shelekhin, Yu. L.; Votinov, M. P. Tr. Leningrad. Politekh. Inst.; No. 325, 78-80(1971). (In Russian).

A rose ruby (0.05 percent Cr_2O_3 in burden) was irradiated by ^{60}Co γ rays (10^8 R) and thermal neutrons (10^{18} to 10^{17} n/cm² at room temperature, and single crystals of corundum (and ruby) were irradiated by x rays (10^4 roentgens) at liquid-nitrogen temperature. The irradiated crystals acquired an orange tint. Two additional absorption bands with centers at 370 and 460 nm appeared in the optical spectrum, the Cr^{3+} content decreased by 11 to 18%, and the nonuniformity of the intracrystalline field increased by 30 to 40%. It was confirmed that the induced coloring is associated with a change in the valence state of the chromium and the formation of color centers in the matrix defects. (JPRS)

47245 VTORICHNO-EMISSIONNYE I STRUKTURNYE SVOISTVA TVERDYKH TEL. (Secondary Emission and Structural Properties of Solid Bodies). Arifov, U. A. (ed.). Tashkent; Izdatel'stvo Fan Uzbekskoi SSR (1970). 212p.

This publication contains 36 papers by different authors on the interaction of ion, electron, and positron beams with matter. The phenomena of sputtering, secondary ion and electron emission, and the changes in the physico-chemical properties of substances are discussed. Adsorption during ion bombardment and the annihilation of positrons in solids are reviewed. (tr-auth)

47246 ACCUMULATION OF F CENTERS IN NaCl CRYSTALS AS A FUNCTION OF THE PROTON-BEAM INTENSITY. Vaisburd, D. I.; Alekseev, P. D. (Tomsk Polytechnic Inst., USSR). Izv. Vyssh. Ucheb. Zaved., Fiz.; No. 11, 27-31(1971). (In Russian).

The dependence of the yield of radiation defects on the radiation intensity is discussed. The form of these relations is determined by the ratio between the relaxation time of processes in particle tracks and the overlapping time of tracks. The existence of an optimum intensity was demonstrated at which the yield of defects was maximum. In the accumulation of F centers at a slow rate at 295°K, an essential role is played by processes having a relaxation time of the order of 10 seconds. (tr-auth)

Physical Properties

Refer also to abstracts: 45992 and 46023.

47247 (BNWL-1511) HIGH-TEMPERATURE ELECTRICAL CONDUCTIVITY OF CeO_{2-x} . Bates, J. Lambert (Battelle Pacific Northwest Labs., Richland, Wash.). Jun 1972. Contract AT(45-1)-1830. 21p. Dep. NTIS.

The electrical conductivity of CeO_{2-x} was measured from 16 to 1998°C in argon containing a low partial pressure of oxygen. The electrical conductivity of CeO_2 increased during five heating cycles as the O/Ce ratio decreased to a final value of 1.82. The energy for conduction below 1900°C decreased during the heating cycles as the conductivity increased. Three discontinuities in the log σ versus $1/T$ curves were observed at approximately 1000, 1150, and 1900°C. Above 1900°C, the conductivity decreased sharply with increasing temperature. After heating to 1900°C, no discontinuities were observed in the log σ versus $1/T$ curve for $\text{CeO}_{1.82}$ below 1900°C. (auth)

47248 ELECTRICAL AND THERMAL CONDUCTIVITIES OF RHENIUM, OSMIUM, AND IRIUM AND SOME OF THEIR ALLOYS WITH THORIUM AND LANTHANUM IN A WIDE TEMPERATURE RANGE. Gugin, A. A.; L'rov, S. N.; Mal'ko, P. I.; Nemchenko, V. F.; Starodubov, I. P. (Kherson Pedagogical Inst., USSR). Sov. Powder Met. Metal Ceram. (Engl. Transl.); 11: No. 1, 51-5(Jan 1972).

Translated from Porosh. Met., Akad. Nauk Ukr. SSR; 11: No. 1, 65-70(Jan 1972).

Studies were made on the effects of temperature on the electric conductivity, ρ , and thermal conductivity, λ , of Re-Th, Os-Th, and Ir-La alloys produced from powders of their starting components. Results are given for measurements of ρ and λ for pressed and fused Re, Os, and Ir specimens and the above alloys in the ranges 100 to 1700°C and 100 to 1200°C, respectively. The electric conductivity in the temperature range investigated was much higher than that of their base metals. The thermal conductivity in the temperature range investigated was much less than that of

the pure base metals. This, together with the increase in the electric conductivity of the alloys, was evidence that the Wiedemann-Franz law is qualitatively satisfied. With rise in temperature, the thermal conductivity of the alloys fell, and, at high temperatures, quite distinctly tended to a constant value. On this basis, it was concluded that the thermal conductivity of these alloys at such temperatures is mainly electronic in character. (P.C.H.)

47249 ELECTRICAL CONDUCTIVITY, THERMO-EMF, AND THERMIONIC EMISSION OF BaO AND SrO CRYSTALS. Kovalev, N. N.; Sorokin, O. V. (Inst. of Semiconductors, Leningrad). Inorg. Mater. (USSR) (Engl. Transl.); 8: No. 1, 94-9(Jan 1972).

Translated from Izv. Akad. Nauk SSSR, Neorg. Mater.; 8: No. 1, 111-16(Jan 1972).

The thermo-emf was measured for single crystals of BaO and SrO in order to calculate the position of the Fermi level relative to the bottom of the conductivity zone. From measurements of the thermoemission current, the thermodynamic work function was obtained. Comparison of the electron affinity with its value for the surface, obtained from photoelectric measurements, showed that in vacuum the bending of the zones at the surface must be small in comparison with ϕ and χ . Estimation of the positions of the donor levels in the forbidden zone led to the conclusion that, in all probability, these are F centers. The electron mobilities in the conductivity zones of BaO and SrO were estimated. Comparison with theoretical data showed that they are mainly determined by interaction with polar optical vibrations. It was found that when $E \geq 10^4$ V/cm, the thermionic emission follows the normal Schottky law. (auth)

47250 CARBIDE RESISTORS. Grebenkina, V. G. (Inst. of Problems in Materials Science, Kiev). Inorg. Mater. (USSR) (Engl. Transl.); 8: No. 1, 161-2(Jan 1972).

Translated from Izv. Akad. Nauk SSSR, Neorg. Mater.; 8: No. 1, 181-2(Jan 1972).

The temperature dependence of the electric conductivity of niobium carbides was investigated between room temperature and 1000°C. In the region of homogeneity, the electric conductivity of niobium monocarbide increased with the defect density of the carbon sublattice. This was due to an increase in the concentration of additional scattering centers in the form of vacancies in the carbon sublattice. The temperature coefficient of the resistance (TCR), on the other hand, decreased as the lower limit of the region of homogeneity was approached. Test results for resistors with $\text{NbC}_{0.74}$ as current-carrying phase revealed that resistors prepared from this substance have the following principal characteristics: rated value, 1 Ω ; TCR, -5×10^{-4} deg⁻¹; moisture retention 1%. (P.C.H.)

47251 NEUTRON SCATTERING. Jacrot, Bernard (Institut Max von Laue-Paul Langevin, Grenoble, France). Recherche (Paris); 3: No. 22, 325-33(Apr 1972). (In French).

The usefulness of thermal neutrons from high-flux nuclear reactors for studying the condensed structure of matter (positions and motion of atoms in matter) is emphasized. Neutron- and x-ray-diffraction techniques are compared (magnetic structures and crystalline structures). The use of inelastic neutron scattering in the study of the dynamics of condensed matter is presented. (France)

47252 ELECTRON SPIN RESONANCE IN SINGLE CRYSTAL YTTRIA STABILIZED ZIRCONIA. Thorp, J. S.; Aypar, A.; Ross, J. S. (Durham Univ., Eng.). J. Mater. Sci.; 7: No. 7, 729-34(Jul 1972).

Electron spin resonance was observed at 35 GHz in 8 and 12 mol % yttria-stabilized zirconia single crystals before and after blackening by current passage. Measurements were made between 293 and 77°K. In both as-grown and blackened crystals, the spectra showed anisotropic lines (type A) characterized by $g_{\parallel} = 2.003 \pm 0.002$ and $g_{\perp} = 1.880 \pm 0.002$ with respect to a [111] symmetry axis and a broader, slightly anisotropic line (type B) centered near $g = 1.993$. In current-blackened crystals, a weak isotropic line (type C) was also found near $g = 1.986$. The type-A lines are attributed to a charged complex formed by an electron trapped at an oxygen vacancy and associated with an yttrium ion. Tentative models are suggested to explain the type-B and type-C lines. (auth)

47253 ANISOTROPY IN THE PARAMAGNETIC SUSCEPTIBILITY OF SOME RARE EARTH ELEMENTS. Aleonard, Roland (Centre National de la Recherche Scientifique, Grenoble, France). Compt. Rend., Ser. B; 274: No. 16, 967-9(17 Apr 1972). (In French).

The changes in the paramagnetic susceptibilities of Gd, Tb, Dy,

Ho, and Er monocrystals as a function of temperature were measured (5 to 750°C) for various orientations of the field with respect to the crystallographic axes. (France)

47254 ELECTRON DENSITY IN TUNGSTEN CRYSTAL ACCORDING TO NEUTRON-ELECTRON INTERACTION DATA. Aleksandrov, Yu. A.; Ozerov, R. P.; Ranev, N. V. (Inst. of Physics and Chemical Research, Moscow). JETP Lett. (USSR) (Engl. Transl.); 15: No. 7, 280-2 (5 Apr 1972).

Translated from Zh. Eksp. Teor. Fiz., Pis'ma Red.; 15: No. 7, 398-400 (5 Apr 1972).

Measurements of the electron density distribution on the (001) plane of W were made using neutron diffraction and x-ray scattering. The two results are identical. (D.L.C.)

47255 ENERGY SPECTRUM AND CRYSTALLINE STRUCTURE OF WURTZITE-LIKE MODIFICATION OF BORON NITRIDE. Frantsevich, I. N.; Zhurakovskii, E. A.; Kurdyumov, A. V.; Vasilenko, N. N. (Inst. of Problems of Materials, Kiev). Dokl. Akad. Nauk SSSR; 203: No. 1, 87-90 (1 Mar 1972). (In Russian).

The four known crystal modifications of BN are BN_{hex} , BN_{rhomb} , BN_{cube} , and BN_{hex-w} , where the last is a hexagonal structure resembling wurtzite. A spectral and x-ray-structure study was made of BN_{hex-w} prepared by pulsed compression of BN_{hex} . B-N bond lengths, interlayer distances, and emission $K\alpha$ bands in the various BN modifications were compared. (K.S.W.)

47256 MAGNETIC SURFACE STATES IN ZINC AND CADMIUM. Rahn, John Phillip. Seattle; Univ. of Washington (1972). 101p. University Microfilms Order No. 72-20,886.

Thesis.

Magnetically induced surface electron state resonances as a function of sample orientation were studied for the first time in high-purity single crystals of zinc and cadmium. The transition magnetic fields are recorded for all sample orientations in three principal crystallographic planes. The experimentally observed magnetic fields of the surface resistance derivative maxima agree with the Nee-Prange theory. The angular behavior of these magnetic fields is compared to those predicted by a semi-empirical OPW pseudopotential band model. Conversely, an iteration scheme is used to obtain the point Fermi velocities from the measured data points for the third-band electron lens. The agreement between the predicted and observed magnetic field is excellent for the lens. From the point velocities, it is concluded that the electron-phonon renormalization of the one-electron energy bands is not constant even on the spherical portion of the lens. (Diss. Abstr. Int., B)

47257 TEMPERATURE DEPENDENCE OF THE WORK FUNCTION OF THE Ba- W_{10} SYSTEM. Shul'man, A. R.; Kirsanova, T. S.; Belova, V. M. Radio Eng. Electron. Phys. (USSR) (Engl. Transl.); 16: No. 10, 1792-3 (Oct 1971).

Translated from Radiotekh. Elektron.; 16: No. 10, 2003-4 (Oct 1971).

The work function of W(110) covered with Ba (coverage varied from $\theta = 0.5$ to 2) was measured at 600 to 900°K. The coefficient $\Delta\phi/\Delta T$ has different values, being $\sim 4 \times 10^{-4}$ eV/deg at 580 to 700°K and 1.2×10^{-3} eV/deg at 700 to 900°K. The results are compared with these for the BaO-W system. Dips are visible in the $\phi(T)$ curves, especially for $\theta = 0.8$. (D.D.C.)

47258 dc ANISOTROPIC MAGNETIZATION AT 20.4 K AND THE IDEAL CRYSTAL-FIELD-ONLY BEHAVIOR OF TmSb. Foner, S. (Massachusetts Inst. of Tech., Cambridge); Cooper, Bernard R.; Vogt, Oscar. Phys. Rev., B; 6: No. 5, 2040-4 (1 Sep 1972).

The anisotropic magnetization of TmSb at 20.4°K was measured in applied dc fields up to 144 kOe. The experimental results are in excellent agreement with isothermal crystal-field-only theory for parameters deduced from earlier pulsed-field experiments at 1.5°K. This resolves an ambiguity posed by earlier pulsed-field magnetization experiments at 20.4°K, the results of which fell between the adiabatic and isothermal crystal-field-only theoretical curves. High-resolution dc measurements of the magnetization at 1.5°K were also made up to 52 kOe and are in good agreement with the earlier pulsed-field data. (auth)

47259 PARAMAGNETIC SUSCEPTIBILITIES OF METALLIC SAMARIUM COMPOUNDS. Stewart, A. M. (Univ., Genova. Imperial Coll. of Science and Tech., London). Phys. Rev., B; 6: No. 5, 1985-98 (1 Sep 1972).

It is shown that the influence of conduction-electron polarization effects upon the susceptibilities of metals containing the tripositive samarium ion is very much greater than upon metals containing normal rare earths. A theory of the susceptibility of metallic samarium materials is developed that takes account of these polarization effects and of interionic Heisenberg exchange cou-

plings, the admixture of the $J = 7/2$ state into the $J = 5/2$ ground state, which is assumed to be the only one to be thermally populated, but which, however, does not take account of crystal-field splittings. The susceptibility is found to be of the unexpectedly simple form $\chi(T) = \chi_0 + D/(T - \theta)$, in which the only dependence on temperature T is that explicitly shown. This expression is found to fit the published data for the susceptibility of dhcp samarium to an accuracy of 1% in the temperature region 110 to 230°K, and the parameters extracted from the fit are found to be in excellent agreement with those obtained for the other light rare-earth metals. An expression is also derived for the susceptibilities of metals containing normal rare earths, which takes account of both conduction-electron polarization and crystal field effects. (auth)

47260 SINGLE-ELECTRON ENERGIES, MANY-ELECTRON EFFECTS, AND THE RENORMALIZED-ATOM SCHEME AS APPLIED TO RARE-EARTH METALS. Herbst, J. F. (Cornell Univ., Ithaca, N. Y.); Lowy, D. N.; Watson, R. E. Phys. Rev., B; 6: No. 5, 1913-24 (1 Sep 1972).

A systematic investigation of certain electronic properties of the rare-earth metals is reported. Calculations are performed within the framework of the renormalized-atom method in which Hartree-Fock free-atom solutions, with electronic configurations appropriate to the metal, are initially computed; the wave functions are then renormalized to the Wigner-Seitz sphere and used to construct l-dependent Hartree-Fock-Wigner-Seitz crystal potentials. The following results are obtained: (i) Recent spectral information together with the free-atom solutions permits one to estimate the change in neutral-atom correlation energy associated with changing the 4f electron count; contrary to expectation, it is found that correlation effects are more significant in a configuration with one fewer 4f and one more 5d electron. (ii) Band extrema and Fermi levels are placed. (iii) The positions of occupied and unoccupied 4f levels are estimated in both a one-electron approach and a multielectron method taking screening and relaxation effects into account in a definite way. The one-electron approximation for the 4f levels fails badly in reproducing the results of recent photoemission experiments, while the multielectron calculations are in surprisingly good accord with experiment. The effective Coulomb-interaction energy between two 4f electrons at the same site, the familiar U , is reduced from the single-particle value of approximately 27 eV to about 7 eV with the inclusion of multielectron effects. (iv) Hartree-Fock values for the 4s- and 5s-shell exchange splittings are compared with soft-x-ray photoemission studies of the rare-earth fluorides and oxides; the calculated 4s splittings are roughly twice as large as experiment while, unexpectedly, the 5s results are in almost precise agreement. (auth)

47261 MAGNETIC PHASE DIAGRAM OF THE UAs-US SYSTEM. Lander, G. H.; Mueller, M. H.; Reddy, J. F. (Argonne National Lab., Ill.). Phys. Rev., B; 6: No. 5, 1880-8 (1 Sep 1972).

The magnetic phase diagram of the $UAs_{1-x}S_x$ system ($0 \leq x \leq 1$) was investigated by neutron diffraction with powder samples. The solid solutions are face-centered cubic (NaCl crystal structure), and the anion atoms are distributed randomly over the anion sublattice. For $x = 0$ (UAs), the type-I antiferromagnetic ordering ($T_N = 127^\circ K$), consisting of ferromagnetic sheets stacked $+-+-$, changes at $0.5T_N$ to the type-IA ordering, in which the sheets are stacked $++--$. This transition is accompanied by an abrupt increase in the magnetic moment from 1.9 to $2.2 \mu_B$ per uranium atom. As x increases from zero, the stability of the type-IA structure increases until for $x = 0.20$ this phase is present at all temperatures below T_N . For x between 0.25 and 0.34, the magnetic moments are not arranged in a simple commensurate structure, but their modulation from layer to layer follows a sinusoidal form. Near T_N , a single harmonic is present: The repeat distance in real space varies from ~ 2.8 unit cells at $x = 0.25$ to ~ 4.2 unit cells at $x = 0.34$. A third harmonic, implying some squaring of the sine wave, appears at about $0.5T_N$. The periodicity is a function of concentration (x) only and does not vary with temperature for a given sample. Although at the ends of this intermediate range the stable phases at low temperature tend to be either type IA (low x) or ferromagnetic (high x), the longitudinal-wave structure is stable to at least $2.2^\circ K$ over a narrow composition range. (auth)

47262 DETERMINATION OF THE ELECTRON-PHONON ENHANCEMENT FACTOR FROM SPECIFIC-HEAT DATA. Knapp, G. S. (Argonne National Lab., Ill.); Jones, R. W. Phys. Rev., B; 6: No. 5, 1761-7 (1 Sep 1972).

The electron-phonon enhancement factor is determined for scandium, yttrium, vanadium, platinum, and palladium solely from specific-heat data. The key to the method is that at high temperatures the electron-phonon enhancement factor is zero. Therefore, if the electronic specific-heat coefficient at high tem-

peratures is calculated and compared with the low-temperature value, both the band-structure density of states and the electron-phonon enhancement factor can be determined. (auth)

47263 STUDY OF HYDROGEN IN VANADIUM. Wanagel, John, Ithaca, N. Y.; Cornell Univ. (1971). 70p. University Microfilms Order No. 72-15,421.

Thesis.

The manner in which the hydride precipitate manifests itself on a polished (100) surface of a single crystal was examined by optical methods. Two different surface features were associated with the formation of a vanadium hydride precipitate. Orientation relationships between co-existing phases were determined using x-ray and electron-microscopy techniques. The axes of the β (bct) hydride phase were shown to be slightly tilted away from the cube axes of the α (bcc) phase. The relationship between the β (bct) phase and the γ (fcc) phase was shown to be such that $(100)\beta \parallel (110)\gamma$ and $[001]\beta \parallel [001]\gamma$. In the single-phase β (bct) region, frequent examples of twinning were observed on (101) planes. Electron-microscope specimens of vanadium, thinned by techniques known to add hydrogen, showed evidence for ordering in β -vanadium hydride. Extra half-order reflections were observed and concluded to be due to scattering from hydrogen. The transformation between the ordered and disordered phases occurs reversibly at a specific temperature in the range 160 to 190°C depending on the specimen. The reciprocal lattice as determined by electron diffraction was shown to be in the monoclinic Laue group. Indirect evidence places the composition of the ordered phase in the vicinity of V_2H . The possible monoclinic structures consistent with the reciprocal lattice in all cases place the hydrogens in tetrahedral sites on alternate (101)_{bct} planes. Application of tensile stress at 200°K along the [100] direction of a single crystal of vanadium containing about 1.2 at. % hydrogen caused the disappearance of hydride plates and the creation of a large number of slip lines. Deformation twinning was suppressed by the precipitate and fracture was by brittle cleavage on a (100) plane. (Diss. Abstr. Int., B)

47264 LOW-TEMPERATURE ELECTRON-PHONON UMKLAPP SCATTERING IN SIMPLE METALS. Ekin, John W. Ithaca, N. Y.; Cornell Univ. (1971). 121p. University Microfilms Order No. 72-18,552.

Thesis.

The first conclusive evidence for the long postulated low-temperature disappearance of the electron-phonon umklapp scattering contribution to the electrical resistivity of potassium is reported. Accurate low temperatures (1 to 25°K) resistivity data, obtained from potassium samples having residual resistance ratios $[RRR = \rho(20^\circ K)/\rho(0^\circ K)]$ in the range from 100 to 8000, are interpreted in terms of a detailed theoretical analysis of umklapp scattering effects. Deviations from Matthiessen's rule, while present, are comparatively small and do not alter appreciably the temperature dependence of the resistivity for even the least pure sample ($RRR = 136$). In addition, a theoretical analysis of umklapp effects in potassium is performed for the ultrasonic attenuation (in the local limit) and the thermal resistivity. Using various model pseudopotentials and a realistic phonon spectrum, both these quantities are evaluated at temperatures from 1 to 20°K. The rapid decrease of the umklapp component at very low temperatures is emphasized in each case. Accurate ($\pm 0.01\%$ relative accuracy) low-temperature (1 to 100°K) data on the purity and temperature dependence of the electrical resistivity of the simple polyvalent metals, aluminum and cadmium, are also presented and interpreted in terms of electron-phonon umklapp interaction. In contrast to potassium, the resistivity of aluminum exhibits relatively large deviations from Matthiessen's rule and it is suggested that anisotropic umklapp relaxation of the electron distribution may be playing an important role in determining the magnitude of such deviations. (Diss. Abstr. Int., B)

47265 ION-ION EMISSION: A NEW TOOL FOR MASS-SPECTROMETRIC INVESTIGATIONS OF PROCESSES ON THE SURFACE AND IN THE BULK OF SOLIDS. Fogel, Ya. M. (Gorky State Univ., Kharkov). J. Mass Spectrom. Ion Phys.; 9: No. 2, 109-25 (Jul 1972).

A brief account of the properties of the ion-ion emission phenomenon is given. The possibilities of applying this phenomenon to the study of processes on the surface and in the bulk of solids are discussed. The foundations of the new method are considered. (auth)

47266 MAGNETIC SUSCEPTIBILITY OF CUBIC NIOBIUM AND TANTALUM CARBIDES IN THE LOW TEMPERATURE RANGE. Borukhovich, A. S.; Dubrovskaya, L. B.; Matveenko, I. I.; Gel'd, P. V. (Inst. of Chemistry, Sverdlovsk, USSR). Tr. Inst. Khim., Akad. Nauk SSSR, Ural. Filial; No. 20, 71-3 (Sep 1970). (In Russian).

The magnetic susceptibility of the cubic carbides of Nb and Ta

was measured in their regions of homogeneity at low temperatures. The Curie-Weiss function χ for the composition $NbC_{0.80}$ was detected below 30°K. The results were considered in terms of the available concept of the electron structure of these compounds. (tr-auth)

47267 DYNAMICAL THEORY FOR THE CONTRAST OF PERFECT AND IMPERFECT CRYSTALS IN THE SCANNING ELECTRON MICROSCOPE USING BACKSCATTERED ELECTRONS. Spencer, J. P.; Humphreys, C. J.; Hirsch, P. B. (Oxford Univ.). Phil. Mag.; 26: No. 1, 193-213 (Jul 1972).

A many-beam dynamical theory, which approximately takes into account multiple inelastic scattering, is formulated and used to calculate the intensity of electrons backscattered from perfect and imperfect crystals. The contrast of channeling pattern lines is discussed and the occurrence of "missing" lines explained. The backscattered intensity due to an inclined stacking fault is calculated. The image consists of damped fringes of periodicity ξ_g ; the nature of the first fringe depends on the sign of $g \cdot R$ and no contrast occurs for integral values of $g \cdot R$. Similar calculations were made for dislocations. Reversing the sign of $g \cdot b$ reverses the image contrast for dislocations near the surface, and the usual TEM invisibility criteria hold, namely invisibility for $g \cdot b = g \cdot b_{\text{u}} = 0$. The experimental conditions necessary for the imaging of defects are discussed. Imaging defects in solid specimens should be possible using cold field-emission electron sources. (auth)

47268 STUDY OF DOPPLER-BROADENED ANNIHILATION RADIATION IN METALS WITH A Ge(Li) DETECTOR. Alfano, F.; Reynolds, T.; Ramaswamy, M. K. (Fordham Univ., Bronx, N. Y.). Nuovo Cim.; 10B: No. 1, 334-8 (11 Jul 1972).

Doppler-broadened positron annihilation radiation spectra were investigated in twelve metals. A plot of αX^2 vs. E_F for these metals, when grouped according to the periodic table, results in straight lines. (α is the effective-electron-mass ratio m^*/m_0 , X is a measure of the observed broadening, and E_F is the Fermi energy.) This feature is consistent with a simple theoretical model. In addition, it suggests a possible method for determining the electron effective mass in other metals. This was utilized to estimate the effective-mass ratio for B as 2.8. (auth)

47269 SOME CRYSTALLINE FIELD EFFECTS IN METALS. Fulde, Peter; Peschel, Ingo (Institut Max von Laue-Paul Langevin, Garching, Ger.). Advan. Phys.; 21: No. 89, 1-67 (1972).

Different physical effects are discussed which are caused by the crystalline electric field splitting of rare earth ions in metals. Thus, the rare earth ions may be impurities dissolved in a metallic matrix or they may form a regular lattice. In the former case, the cases of a normal conducting and a superconducting matrix are distinguished between. The influence of the crystalline field splitting on the properties of the conduction electrons is calculated. In the case of a normal matrix, anomalous behavior of the thermoelectric power is found due to the impurity levels. If the matrix is superconducting, large deviations result from the theory of Abrikosov and Gorkov (1960) which describes the influence of non-split magnetic impurities on superconductivity. A comparison of the theory with available experiments is presented. For the case that the rare earth ions form a regular lattice, various aspects of the collective excitations in the paramagnetic state are discussed. Special attention is paid to the soft mode problem of exchange-induced ferromagnets. Furthermore, the influence of impurities on the excitation spectrum (localized modes, resonant modes, etc.) and on the magnetic ordering temperature is discussed. (auth)

47270 GADOLINIUM g-SHIFTS AND BAND STRUCTURE. Gomes, A. A.; Guimaraes, A. P.; Iannarella, L. (Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro). Notas Fis., Cent. Brasil. Pesqui. Fis.; 18: No. 14, 191-8 (1972).

A simple s-, d-band model for the rare earths is used to describe the g shifts in metallic systems containing gadolinium. The temperature dependence of the d-electron susceptibility $\chi^{(d)}$ is suggested to explain the apparent discrepancies between the exchange parameters J_{eff} measured using different techniques (EPR, bulk magnetization measurements, hf fields). A numerical calculation shows that a model band with a narrow (~ 0.1 eV wide) peak near the Fermi level could account for the required variation of $\chi^{(d)}$ in gadolinium metal. (auth)

47271 REMARKS ON Gd g-SHIFTS. Gomes, A. A.; Guimaraes, A. P.; Iannarella, L. (Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro). Notas Fis., Cent. Brasil. Pesqui. Fis.; 18: No. 15, 201-7 (1972).

The s-d hybridization is given a simple formulation in order to discuss its relevance to the Gd g shifts; it is apparent that the

effective exchange parameters J_{eff} can be negative even with negligible interband mixing. A possible explanation for the signs of the g shift and band magnetization in Gd metal is also suggested. (auth)

47272 MAGNETIC PROPERTIES OF SOME ACTINIDE HALIDE COMPOUNDS. Hendricks, Melvin Elton. Columbia, S. C.; Univ. of South Carolina (1971). 195p. University Microfilms Order No. 72-12,007. Thesis.

Magnetic susceptibility measurements on 13 actinide compounds were made with a vibrating-sample magnetometer, most in the temperature range 2.5 to 70°K at 10,000 Oe. These compounds, classed according to site symmetry of the actinide ion, are: (1) octahedral: $\text{Cs}_2\text{NaMCl}_6$ where M is a trivalent actinide, U, Np, Pu, Am, Cm, Bk, the latter two doped into $\text{Cs}_2\text{NaLuCl}_6$; (2) C_{3h} : UCl_3 , NpCl_3 , PuCl_3 ; (3) D_{2d} : PaCl_4 , UCl_4 ; (4) low-symmetry: PuF_4 , UBr_4 . Theoretical calculations, including intermediate coupling and limited J-mixing, were performed for several cases for comparison with experimental magnetic properties. Results for the octahedral compounds include determination of ground levels, observation of small distortion effects, and measurements of limits on crystal-field parameters. In the $\text{Cs}_2\text{NaMCl}_6$ matrix, the following ground levels were determined: U^{3+} , Γ_8 ; Np^{3+} , Γ_6 ; Pu^{3+} , Γ_6 ; Am^{3+} , Γ_1 ; Bk^{3+} , Γ_1 . The U, Np, and Pu compounds, at the lowest temperatures, gave deviations from Curie-Weiss behavior that are attributed to small distortions from octahedral symmetry. Crystal-field calculations, including J-mixing with the first-excited state, were made for U^{3+} and Np^{3+} for ratios of the octahedral crystal-field parameters, B_6^0/B_4^0 , between 0 and +1, varying B_6^0 from 0 to 20,000 cm^{-1} . Assuming an approximately constant B_6^0/B_4^0 ratio in the $\text{Cs}_2\text{NaMCl}_6$ series, the limits $+0.07 \leq (B_6^0/B_4^0) \leq +0.77$ were established from analysis of U^{3+} and Np^{3+} data. $\text{Cs}_2\text{NaAmCl}_6$ below 70°K exhibits temperature-independent paramagnetism, in good agreement with the theoretical value. Bk^{3+} in $\text{Cs}_2\text{NaLuCl}_6$ also gives temperature-independent paramagnetism. In the C_{3h} trichloride series, several types of magnetic phenomena were observed. Antiferromagnetic ordering was found for UCl_3 , $T_N = 22 \pm 1^\circ\text{K}$, and for PuCl_3 , $T_N = 4.5 \pm 1^\circ\text{K}$. NpCl_3 exhibited temperature-independent paramagnetism below 50°K. Each of the C_{3h} compounds displays Curie-Weiss behavior at relatively low temperatures, with measured moments in close agreement with the free-ion values. Using previously reported wave functions and energy levels for PuCl_3 , calculated and experimental susceptibilities were in excellent agreement from 10 to 500°K. The D_{2d} compound PaCl_4 was found to have a ferromagnetic transition, with $T_C = 182 \pm 2^\circ\text{K}$, an unusually high transition temperature for a ferromagnetic insulator. Below 182°K, the magnetization was unsaturated at fields up to 12,000 Oe; hysteresis was not observed above 3.29°K. In the paramagnetic region, PaCl_4 obeys the Curie-Weiss law from 182 to 219°K; crystal-field calculations for this region indicate that J-mixing is significant. The D_{2d} compound UCl_4 has a magnetic moment with complex behavior below 190°K, with temperature-independent paramagnetism below 35°K; the susceptibility in this region was reproduced by a simple theoretical model. The magnetic behavior of both PuF_4 and UBr_4 is consistent with low-symmetry crystal fields. PuF_4 showed temperature-independent paramagnetism below 70°K, indicating a fairly large crystal field. UBr_4 was paramagnetic at the lowest temperatures measured, consistent with a small C_{2v} crystal field. Freshly prepared PuF_4 below 15°K gave an anomalously large, temperature-dependent diamagnetism, which disappeared upon aging. (Diss. Abstr. Int., B)

47273 MAGNETIC COMPENSATION IN IRON-GADOLINIUM AND IRON-TERBIUM FILMS. Buravikhin, V. A.; Bochkarev, V. F.; Egorov, V. A.; (and others) (Irkutsk Pedagogical Inst., USSR). Izv. Vyssh. Ucheb. Zaved., Fiz.; No. 11, 88-92(1971). (In Russian).

Physical properties and phase composition of Fe-Gd and Fe-Tb films possessing magnetic compensation were investigated. The effect of phase compensation on galvanomagnetic and magnetic properties was studied. Experiments were conducted for the purpose of elucidating the nature of magnetic compensation which was observed during the initial annealing. Experimental and calculated data were obtained on the phase composition of films after annealing, involving the compounds GdFe_5 , Gd, Gd_2Fe_7 , and $\text{Gd}_2\text{Fe}_{17}$. (tr-auth)

Moessbauer Effects and NMR

Refer also to abstract: 46911.

47274 OPTIMUM ABSORBER THICKNESS FOR SINGLE LINE MOESSBAUER EFFECT IN THE PRESENCE OF STRONG NON-

RESONANT ABSORPTION. Pelzl, J. (CEN, Grenoble, France). Nucl. Instrum. Methods; 102: No. 2, 349-51(1972).

Estimates of the optimum absorber thickness for a single-line Mössbauer effect measurement are derived. Numerical results for different magnitudes of the background noise are presented. (auth)

47275 ANALYSIS OF FIVE LINE HYPERFINE SPLIT MOESSBAUER ABSORBER SPECTRA. Hershkowitz, N.; Ruth, R. D.; Wender, S. A.; Carpenter, A. B. (Univ. of Iowa, Iowa City). Nucl. Instrum. Methods; 102: No. 2, 205-17(1972).

Sums of Lorentzians are fit to numerical solutions of the transmission integral for five-line hyperfine-split absorber spectra. The general dependences of fit parameters are discussed. Procedures are given to determine actual absorber parameters. (auth)

47276 PRESSURE DEPENDENCE OF MOESSBAUER RECOILLESS FRACTION AND SECOND-ORDER DOPPLER SHIFT. Srivastava, B. N.; Tyagi, R. N. (Government Coll., Naini Tal, India). Indian J. Pure Appl. Phys.; 10: No. 2, 89-92(Feb 1972).

The effect of pressure on the recoilless fraction f for ^{57}Fe (14.4 keV) gamma transition in some host lattices was studied theoretically through the variation of rigidity and density with pressure in low- and high-temperature ranges. The calculated values of f agree closely with the available experimental values. The pressure dependence of second-order Doppler shift for a number of lattices in the pressure range of 0 to 300 kbar was investigated through the variation of rigidity and density. The calculated values of temperature and pressure coefficients for ^{57}Fe compare reasonably well with the values reported by earlier workers. (auth)

47277 PULSE FOURIER TRANSFORM NUCLEAR MAGNETIC RESONANCE SPECTROSCOPY. Netzel, Daniel A. (Northwestern Univ., Evanston, Ill.). Appl. Spectrosc.; 26: No. 4, 430-42(1972).

Recent advances in NMR spectroscopy allow one to study extremely dilute systems or nuclei which occur in low natural abundance and/or have poor sensitivity for NMR detection. This paper describes various pulse and pulse Fourier transform NMR techniques. Included in this introduction are: (1) a brief review of the nuclear relaxation phenomenon, (2) sensitivity enhancement using the pulse technique, (3) pulse instrumentation, and (4) the analytical applications of pulse Fourier transform NMR. 31 references. (auth)

47278 SOME RECENT DEVELOPMENTS IN HIGH RESOLUTION NUCLEAR MAGNETIC RESONANCE. Becker, Edwin D. (National Institutes of Health, Bethesda, Md.). Appl. Spectrosc.; 26: No. 4, 421-30(1972).

Techniques for studying high-resolution nuclear magnetic resonance spectra have been considerably broadened in recent years. The most far reaching development—pulse Fourier transform (FT) methods—is discussed in detail. Applications of FT techniques to measurement of relaxation times and to enhancement of weak signals, especially from natural-abundance ^{13}C , are reviewed. Double resonance methods, particularly the nuclear Overhauser effect, and the use of lanthanide shift reagents are also covered in this "mini-review." 39 references. (auth)

47279 POWDER ENDOR LINE SHAPES: NUCLEAR RELAXATION INDUCED BY MOTION OF NEARBY ELECTRON SPINS. Leniart, Daniel S.; Hyde, James S.; Vedrine, Jacques C. (Varian Associates, Palo Alto, Calif.). J. Phys. Chem.; 76: No. 15, 2079-87(20 Jul 1972)

The line shape and intensity of electron-nuclear double resonance (ENDOR) spectra in powders is determined, in part, by the specific form of the mechanism(s) that produce nuclear spin-lattice relaxation. In all cases investigated, nuclear relaxation is assumed to stem from modulation of the electron-nuclear dipolar (END) interaction. The initial model considered assumes that the unpaired electron is trapped in a cage and undergoes a series of jumps between at least two equally probable sites. The dimensions of the cage are such that any fluctuations in angle are severely restricted, and this jumping motion results only in a modulation of the interparticle distance between the trapped H atom and surrounding nuclei outside the cage. It is shown that the dominant relaxation component of the surrounding nuclei is proportional to $\sin^4 \theta$ and leads to a doubly peaked ENDOR spectrum. The second model assumes that the trapped electron resides in a cage whose walls are at infinite potential, and the system is treated quantum mechanically as a particle in a three-dimensional box. The thermal lattice motions of the cage modulate the dipolar interaction in some random manner to produce a spectral density which differs from that of the jump model by the

appearance of a Boltzmann factor describing the particle populations and a frequency term corresponding to the quantized transitions induced between the energy levels of the box. It is shown that for particle transition frequencies, $\omega_{\nu\nu'}$, which are less than or equal to spin transition frequencies, $\omega_e \pm \omega_n$, the quantum mechanical spectral densities reduce to those of the semiclassical jump model. As the temperature is reduced below 20°K the ENDOR line shape expected from the "box" model is shown to resemble that of a matrix ENDOR signal. At very low temperatures a change in both the ENDOR line shapes and intensities is predicted. (auth)

47280 MOESSBAUER EFFECT STUDIES OF ^{57}Co AND ^{57}Fe IN β -RHOMBOHEDRAL BORON. Stanke, F.; Parak, F. (Technische Univ., Munich). Phys. Status Solidi (b); 52: No. 1, 69-77(1 Jul 1972).

The behavior of ^{57}Fe - and ^{57}Co -doped β -rhombohedral boron was investigated by means of Moessbauer spectroscopy. Samples prepared by diffusing ^{57}Fe at temperatures above 1000°C and ^{57}Co at temperatures of 1500°C or higher showed a reproducible quadrupole interaction and a strongly anisotropic f-factor. Using combinations of the different parameters obtained from the spectra, it was attempted to determine the position of the impurities. There is some evidence for an incorporation within the 28-atom subgroup. It was found that the sites occupied by Co and Fe do not have a C_{5v} point symmetry. (auth)

47281 NUCLEAR MAGNETIC RESONANCE IN THE LaX_3 INTERMETALLIC COMPOUNDS. Welsh, L. B. (Northwestern Univ., Evanston, Ill.); Toxen, A. M.; Gambino, R. J. Phys. Rev., B; 6: No. 5, 1677-85(1 Sep 1972).

The ^{139}La NMR Knight shift and spin-lattice relaxation time (T_1) in LaPb_3 were measured from 1.5 to 300°K. The isotropic and anisotropic ^{207}Pb Knight shifts were measured at 1.5, 4.2, and 77°K and T_1 was measured at 1.55°K. Also presented are measurements of the $\text{La } T_1$ in LaIn_3 from 1.5 to 300°K which are more complete than previously reported. These measurements

are compared with previously reported measurements for LaSn_3 . The ^{139}La Knight shift (+0.010% at 4.2°K) in LaPb_3 has a temperature dependence roughly five times that found in LaSn_3 and LaIn_3 even though the bulk susceptibility is considerably smaller. For LaPb_3 , the $\text{La } T_1 T$ product is nearly temperature independent ($T_1 T = 0.59 \text{ sec}^\circ\text{K}$ at 4.2°K) and is about half that found in LaSn_3 . For LaIn_3 , $T_1 T$ increases by 50% between 1.5 and 300°K and has a temperature dependence similar to LaSn_3 . The isotropic and anisotropic Pb Knight shifts ($K_{\text{iso}} = +0.70\%$ and $3K_{\text{ax}} = +0.43\%$ at 4.2°K) are nearly temperature independent. The Pb relaxation time in LaPb_3 ($T_1 T = 63 \times 10^{-4} \text{ sec}^\circ\text{K}$) is about four times faster than in Pb metal, indicating a larger s-contact contribution than in Pb metal and a large negative Knight-shift component. The La NMR and bulk-susceptibility data are partitioned using the customary approximation for transition metals. The results for LaIn_3 and LaSn_3 are quite similar, suggesting that in both these materials the La d-orbital susceptibility is much larger than the d-spin susceptibility and that exchange-enhancement effects are probably not important. For LaPb_3 , d-orbital effects are less important than the d-spin contributions. In all three compounds, the La d-spin contributions to the NMR properties and the bulk susceptibility are of similar magnitude. (auth)

47282 DETERMINATION OF THE BACKGROUND OF MOESSBAUER TRANSMISSION EXPERIMENTS USING IRON ALLOYS WITH 3d-TRANSITION ELEMENTS. Semenkin, V. A.; Shevchenko, Yu. A. (Ural Polytechnic Inst., Sverdlovsk, USSR). Instrum. Exp. Tech. (USSR) (Engl. Transl.); 14: No. 6, 1659-61 (1971).

Translated from Prib. Tekh. Eksp.; 14: No. 6, 67-9(1971).

The background development mechanism and the principal sources of errors in the absolute measurement of Moessbauer-effect probability are described, and estimates of their magnitudes are presented. A procedure is described for simultaneous measurement of (a) multiple Compton scattering in the absorber, and (b) the fraction of high-energy radiation which falls in the "window" that cuts out the resonance peak. (auth)

PHYSICS (THEORETICAL)

Refer also to abstract: 46671.

47283 (CERN-72-8) METHODS OF NONLINEAR ANALYSIS. Brown, Clifford C.; Wehrl, Alfred (European Organization for Nuclear Research, Geneva (Switzerland)). 24 May 1972. 47p. Dep. NTIS (U. S. Sales Only).

The topics discussed are the Banach and Brouwer-Schauder fixed-point theorems, fixed points of groups, semigroups of transformations, and the local degree of a continuous mapping including the basic concepts, applications, and infinite-dimensional spaces. (J.F.P.)

47284 (IC-72/45) ASYMPTOTIC NUMBERS: ONE GENERALIZATION OF THE NOTION OF NUMBER. Christov, Chr.; Todorov, T. (International Centre for Theoretical Physics, Trieste (Italy)). Mar 1972. 14p. Dep. NTIS (U. S. Sales Only).

An algebra is proposed whose elements are called power asymptotic numbers. Every number is represented by one class of sequences—functions $a(s)$ with power expansion in the neighborhood of the point $s = 0$, which have fixed coefficients and arbitrary residual terms. A function with power expansion is a function of the form $a(s) = s^\lambda \varphi(s)$ where $\varphi(s)$ has derivatives up to a given order λ ($\lambda = 0, 1, 2, \dots$). (According to Taylor's theorem, functions of that kind can be expanded in power series.) Several properties of these numbers are established. The power asymptotic numbers are objects interesting in themselves, but they are also auxiliary quantities useful for the introduction of the asymptotic functions—a new variant of generalized functions, quite analogous to the distributions of Sobolev-Schwartz but allowing the operation of multiplication. The final aim is to develop an elaborate method for operations with nonnormalizable wave functions in quantum mechanics, necessary, for example, for the investigation of the connection between scattering matrix and cross section. (auth)

47285 (ITF-71-101-P) IMPOSSIBILITY OF INTRODUCING THE "HIDDEN" PARAMETERS INTO QUANTUM MECHANICS. Akhiezer, A. I.; Polovin, R. V. (Akademiya Nauk Ukrainiskoi SSR, Kiev. Institut Teoreticheskoi Fiziki). 1971. 44p. (In Russian). Dep. NTIS (U. S. Sales Only).

Increased interest has recently been displayed in the gnosological foundations of quantum mechanics, in the first place for von Neiman's theorem about the impossibility of introducing the so-called "hidden" parameters into the quantum mechanics. The most simple and transparent proofs of this fundamental theorem are given. The above is preceded by a brief account of basic quantum mechanics, including the measurement process. (auth)

47286 (LYCEN-7202) GENERATING FUNCTION METHOD AND DEVELOPMENTS IN QUASI-BOSON. Hage-Hassan, M.; Lambert, M. (Lyon Univ. (France). Institut de Physique Nucleaire). Jan 1972. 27p. (In French). Dep. NTIS (U. S. Sales Only).

An isomorphism between the H' subspace of physically bound states and a normalized F functional space is established. This is accomplished by introducing a generating function of the H' states. By adopting a Bargmann space for the F space, an isomorphism is obtained between the H_F space of the fermion states and the H_B space of the boson states. An expansion of any quasi-boson operator of H_B is thus feasible. This is particularly interesting for the Hamiltonian operator. In this case an expansion is obtained which is similar to the Beliaev-Zelevinski and Marumori expansions, but more simple and more convergent. (France)

47287 (ORNL-4778) KWIC INDEX FOR NUMERICAL ALGEBRA. Householder, A. S. (Oak Ridge National Lab., Tenn.). Jun 1972. Contract W-7405-eng-26. 538p. Dep. NTIS.

This listing combines, updates, and otherwise augments those issued previously in three separate volumes, two dealing with linear algebra and one on nonlinear algebraic and transcendental equations. The intention was to list recent literature and selected older classics that have a possible interest for those working in this branch of numerical analysis. (auth)

47288 USE OF A QUASI-CLASSICAL APPROXIMATION FOR THE SOLUTION OF A SYSTEM OF HARTREE-FOCK EQUATION. Gataullin, R.; Arifov, P. U. (Inst. of Electronics, Tashkent). Izv. Akad. Nauk Uzb. SSR, Ser. Fiz.-Mat. Nauk; No. 4, 58-62(1971). (In Russian).

With the integro-differential Hartree-Fock equations as a basis, it was shown that valid quasi-classical conditions of quantization with high accuracy are obtained so that numerical solutions may be derived with an electronic computer for the

plotting of λ_{quant} . The value λ_{quant} thus obtained may be used for subsequent refinement by Ridley's formula. (3 figures, 1 table) (K.S.W.)

47289 GENERAL FORMULA FOR OBTAINING EXPECTATION VALUES FROM THE HYPERCHARACTERISTIC FUNCTION. Rosen, G. (Drexel Univ., Philadelphia). Lett. Nuovo Cim.; 4: No. 9, 391-2(1 Jul 1972).

Comments are made on a previous derivation of quantum-mechanical expectation values from the hypercharacteristic function (Shankara: Lett. Nuovo Cimento; 4; 867(1970)). (auth)

47290 IMPROVEMENTS ON A NEW METHOD FOR THE NUMERICAL SOLUTION OF THE SCHROEDINGER EQUATION. Grimm, R. C.; Storer, R. G.; Davies, B. (Flinders Univ. of South Australia, Bedford Park). J. Comput. Phys.; 9: No. 3, 538-46(Jun 1972).

Two procedures are described for improving the accuracy and efficiency of a method recently proposed for the numerical solution of the bound state Schrödinger equation. Firstly, an analysis of the errors in the method dictates the most efficient way for obtaining accurate results and also allows for the use of extrapolation techniques. Secondly, an expansion around the solution for a harmonic oscillator potential yields a more rapidly convergent procedure. Illustrations for a simple one-dimensional potential are given. (auth)

47291 UNITARY OPERATOR FORM OF THE GENERAL BOGOLYUBOV-TYABLIKOV TRANSFORMATION. Zietek, W. J. (Inst. for Low Temperature and Structure Research, Wroclaw). Phys. Lett.; 40A: No. 2, 139-40(3 Jul 1972).

The unitary operators corresponding to the general Bogolyubov-Tyablikov transformation for Bose as well as Fermi operators are determined and some special cases are discussed. (auth)

47292 INCOMPATIBILITY OF GAUGE INVARIANCE AND NONRELATIVISTIC LOCALITY IN PATH INTEGRAL FORMULATION. Patera, J. (Univ. of Montreal). J. Statist. Phys.; 5: No. 3, 185-8(1972).

It is shown that a modification of the usual gauge transformation is essential to the path integral formulation of nonrelativistic quantum mechanics as a consequence of defining the locality condition as follows: The contribution from each path comes entirely from the points on the path. Arguments are based on the similarity between Wiener and Feynman functional (path) integrals. (auth)

47293 FINITE DIFFERENCE SCHROEDINGER EQUATION. Meessen, A. (Louvain Univ., Belgium). Ann. Soc. Sci. Brux. Ser. I: 86: No. 1, 89-100(1972).

To avoid any arbitrariness, one should consider the ultimate limit for the smallest measurable distance as an unknown quantity, whose value could be finite as well as zero. This implies that the usual differential equations of quantum mechanics should be considered as limiting cases of some adequate finite difference equations. The "correspondence principle" which allows one to set up these finite difference equations, is justified by checking the internal consistency of the theory, in the case where the light velocity $c = \infty$. (auth)

47294 COVERING LAW IN QUANTAL PROPOSITION SYSTEMS. Ochs, Wilhelm (Univ., Munich). Commun. Math. Phys.; 25: No. 3, 245-52(1972).

A recent justification of the covering law by Jauch and Piron founded on the existence of ideal measurements of the first kind is shown to contain a gap. This gap is closed by an additional assumption about such measurements. (auth)

47295 AXIOMS OF QUANTUM MECHANICS. Bugajska, K.; Bugajski, S. (Silesian Univ., Katowice, Poland). Bull. Acad. Pol. Sci., Ser. Sci. Math., Astron. Phys.; 20: No. 3, 231-4(1972).

The lattice postulate is eliminated from Zierler's axiomatic scheme by proving that the postulate is a consequence of some other axioms. The result may serve as justification of the "critical" lattice assumption of the logical approach. (auth)

47296 REMARKS ON THE THREE BODY PROBLEM IN QUANTUM MECHANICS. Makai, E. (Mathematical Inst., Budapest). Stud. Sci. Math. Hung.; 6: No. 3-4, 467-80(1971).

Some solutions of the Schroedinger equation of the quantum mechanical three body problem are found avoiding group-theoretical methods. The solutions have the form of power series expansions. (J.F.P.)

47297 POSSIBLE GENERALIZATION OF NONRELATIVISTIC QUANTUM MECHANICS. Kuryshkin, V. V. (Institut de France, Paris). *Compt. Rend., Ser. B*; 274: No. 20, 1107-10 (15 May 1972). (In French).

The different attempts made to construct a new theory which would provide a deterministic description of quantum processes and have statistical properties that coincide with the usual quantum mechanics are reviewed and discussed. (France)

47298 VARIATIONAL PRINCIPLE FOR NONLINEAR LAGRANGIAN IN QUANTUM MECHANICS. Ohtani, Teruya; Sugano, Reiji (Okaka City Univ.). *Progr. Theor. Phys. (Kyoto)*; 47: No. 5, 1704-13 (May 1972).

The variation principle has not been well defined in quantum mechanics for a Lagrangian containing a velocity dependent potential. The method of variation is investigated to derive an Euler-Lagrange equation in connection with Noether's theorem when the Lagrangian is invariant under some point transformation. Obtained are the Euler-Lagrange equation retaining the above invariance by means of the variation method and the Hamiltonian

with which the canonical equation of motion leads to the Euler-Lagrange equation. (auth)

47299 LECTURES ON QUANTUM STATISTICS. VOLUME 2. QUASI-AVERAGES. Bogolyubov, N. N. New York; Gordon and Breach, Science Publishers, Inc. (1970). 237p.

Translated from the Russian by L. Klein and S. Glass.

The concept is introduced of a quasi average and of thermodynamic Green's functions defined with quasi averages. These quasi averages are thermal averages with a symmetry breaking term introduced into the Hamiltonian. This results in a violation of a conservation law and allows nonzero results to be obtained for the quantities of interest. This method is applied to problems of the spontaneous magnetization of the Heisenberg ferromagnet, the crystal lattice, and, in an extensive treatment, the superconductor. The model Hamiltonian for superconductivity theory is examined and bounds for the eigenvalues are obtained. The hydrodynamic equations for superfluids are derived and the corresponding Green's functions are examined. The self consistent field method is related to the method of compensation of dangerous diagrams. (M.H.R.)

REACTOR TECHNOLOGY

Power Reactor Development

Refer also to abstracts: 45181, 45288, and 45439.

47300 ATOMIC ENERGY IN INDIA. Grenon, Michel. Rev. Fr. Energ.; 23: No. 239, 246-57 (Feb 1972). (In French).

The structure of the energy sources of India is reviewed. The various aspects of the Indian electronuclear program are then studied. Although India possesses modest U reserves, she possesses the world's largest Th reserves. In spite of this, the preliminary phase of the nuclear program was upon the Canadian natural U, heavy-water, reactor series (Candu), a type of reactor having a very high Pu yield. An interest exists in fast reactors using Th fuel as well as in breeder reactors using this fuel. Mining, fuel manufacture industries, and heavy water production and irradiated fuel reprocessing plants were developed independently of the reactor construction program. (France)

47301 REACTOR TECHNOLOGY. VOLUME 15, NUMBER 2. One of Four Technical Progress Reviews Published for the Technical Information Center, USAEC. Washington, D. C.; Atomic Energy Commission (1972). 98p. \$0.75.

Four review articles are presented on various aspects of reactor design, fuels, and components. A separate abstract was prepared for each article. (H.D.R.)

For abstracts of individual papers see: 45960, 47211, 47302, and 47303.

47302 DEVELOPMENT OF MATERIALS AND SYSTEMS FOR THE MOLTEN-SALT REACTOR CONCEPT. MacPherson, H. G. (Univ. of Tennessee, Knoxville). Reactor Technol.; 15: No. 2, 136-55 (1972).

Interest in molten-salt-fueled reactors stems largely from the ease and relatively low cost of purifying and reconstituting the fuel. Molten-salt thermal breeder reactors—graphite moderated and with salt constituents that absorb few neutrons—will have low enough inventories and doubling times to greatly ease the future requirements for uranium mining and for enrichment facilities. Among the reactor designs that have been studied, the favorite is a single-fluid breeder fueled with a mixture of ${}^7\text{LiF}$, BeF_2 , ThF_4 , and ${}^{233}\text{UF}_4$. The chemical characteristics of such salts and the behavior of fission products in them are quite well understood from many years of research and from the operation of the Molten Salt Reactor Experiment (MSRE). A number of processes for removing the soluble fission products (and protactinium) from the salt have been explored. Of these, reductive extraction into bismuth is being developed most vigorously. A bubble generator and centrifugal gas stripper are being developed to ensure the speedy removal of the noble-gas fission products, which are only sparingly soluble. Unresolved problems receiving attention include the control of tritium arising from neutron reactions with lithium, surface fissures in Hastelloy N probably arising from fission-product attack, the retention of the radiation resistance of Hastelloy N to high temperatures, and the development of a steam generator. (101 references) (auth)

47303 EFFECTS OF METAL SWELLING AND CREEP ON FAST REACTOR DESIGN AND PERFORMANCE. Huebotter, Paul R. (Argonne National Lab., Ill.). Reactor Technol.; 15: No. 2, 156-83 (1972).

Dimensional instability of stainless steel, due to radiation-induced swelling and radiation-enhanced creep, profoundly affects the design and performance of fast reactors. The design literature is reviewed in an attempt to clarify the impact of swelling and creep and to scope the options available for accommodating these fast-neutron irradiation effects. The problems of swelling and creep are stated for selected reactor components, and contemporary-design thinking on each component is presented. The subject is then discussed on a reactor-by-reactor basis with either a summary description of the design strategy to accommodate swelling and creep or, in the case of reactors designed prior to discovery of these phenomena, the impact of swelling and creep that has been experienced or is foreseen. Finally the probable effect of swelling and creep on fast reactor economics is conjectured. This effect is translated into an incentive for research and development focused toward a metallurgical solution to the problem (14 references) (auth)

47304 STUDY OF THE MECHANICAL AND THERMAL BEHAVIOR OF A HIGH TEMPERATURE REACTOR CORE. Felten, P.; Gros, G.; Malherbe, J. (CEN de Saclay, Gif-sur-Yvette, France). Bull. Inform. Sci. Tech. (Paris); No. 167, 7-19 (Feb 1972). (In French).

The various thermal and mechanical problems relating to the core of a prismatic element HTR are analyzed (assembling and handling of the blocks, dimensional changes induced by graphite irradiation, fuel performance, gas leaks through the stack). A possible core is presented. The behavior of two different types of fuel elements in this core is examined. (France)

Physics and Core Engineering

Refer also to abstracts: 47581 and 47585.

47305 (ORNL-TM-3832) DESIGN STUDIES OF A MOLTEN-SALT REACTOR DEMONSTRATION PLANT. Bettis, E. S.; Alexander, L. G.; Watts, H. L. (Oak Ridge National Lab., Tenn.). Jun 1972. Contract W-7405-eng-26, 69p. Dep. NTIS.

The MSDR, a 350-MW(e) Molten-Salt Reactor Demonstration Reactor, is based on technology much of which was demonstrated by the MSRE. The cylindrical vessel (26 ft diam by 26 ft high) houses a matrix of graphite slabs forming salt passages having a volume fraction in the core of 10%. The flow of fuel salt is distributed so that the temperature rise along any path is the same—from 1050 to 1250°F. In the primary exchanger, heat is transferred to barren carrier salt (in at 900°F, out at 1100°F). In the secondary exchanger, heat is transferred to a stream of Hitec salt (in at 800°F, out at 1000°F). The Hitec oxidizes tritium to tritiated water which is removed and disposed of. The Hitec generates steam at 900°F, 2400 psi in a boiler, superheater, and reheater. Electricity is produced at an overall efficiency of 36.6%. Soluble fission products are removed by discarding the carrier salt every 8 years after recovery of the uranium by fluorination. Volatile fission products are removed by sparging the fuel salt with helium bubbles in the reactor primary system. The fuel cycle cost was estimated to 0.7 mill/kWhr for inventory, 0.3 mill/kWhr for replacement, and 0.1 mill/kWhr for processing, giving a total of 1.1 mills/kWhr. (auth)

47306 BURNUP AND POWER SHIFT EFFECTS IN LARGE FAST BREEDER REACTORS. Philbin, J. S. (Sandia Labs., Albuquerque, N. Mex.); Axford, R. A. Nucl. Technol.; 15: No. 3, 327-42 (Sep 1972).

A method for analyzing the changes in the neutronic characteristics of large fast reactors during long-term operation is developed. The non-uniform buildup and depletion of the fuel nuclides is taken into account by using eigenfunction expansions in the solution of the one-dimensional, multigroup diffusion equations. A refueling scheme is introduced in order to study the behavior over many reloading cycles as well as between two loading events. The depletion of fissile nuclides in the core and the buildup of fissile nuclides in the blanket cause a power shift into the blanket. The effect of this power shift upon the breeding ratio, the power density, and the reactivity coefficients of a large mixed oxide fast reactor is explored in detail. The power shift into the blanket during the approach to equilibrium is 1.55% of the total reactor power. The breeding ratio increases by only 0.56% during reactor operation. The sodium void coefficient increases by 11.2% from startup to the end of an equilibrium period. The Doppler coefficient, which changes only slightly during the approach to equilibrium, becomes more negative. The total change in the Doppler coefficient is less than 2%. 35 references. (auth)

47307 ON-LINE REACTIVITY FEEDBACK ANALYSIS OF EBR-II BY ROD DROP. Larson, H. A. (Argonne National Lab., Idaho Falls, Idaho); Engen, I. A. Nucl. Technol.; 15: No. 3, 462-4 (Sep 1972).

Reactivity feedback is an important indicator in the analysis of the dynamic behavior of a nuclear reactor. The rod-drop perturbation method is used at EBR-II to produce time-dependent power data which is processed with an on-line digital data reduction procedure to produce time-dependent reactivity feedback; reactivity feedback models are defined in the time and frequency domains. This streamlined procedure has resulted in decreased data reduction time and manpower requirements along with increased data integrity. 12 references. (auth)

Control and Instrumentation

47308 REGULATION OF THE WATER-STEAM CIRCUIT OF A NUCLEAR POWER STATION. A ONCE-THROUGH STEAM GENERATOR WITHOUT BOILER. Barber, P. (CEN de Saclay, Gif-sur-Yvette, France). Bull. Inform. Sci. Tech. (Paris); No. 167, 21-7 (Feb 1972). (In French).

The logical process leading to the choice of control system for the water-steam circuit of a power station is exposed. Particular attention is given to the problems of dimensions which arise in the case of the steam generator of a high-temperature reactor. (France)

Coolant System

47309 (CISE-R-257) DIGITAL PROGRAMS FOR THE STUDY OF THE THERMOHYDRAULIC BEHAVIOR OF THE PRIMARY CIRCUIT OF THE REACTOR CIRENE (PECVAR, CRIP). Muscettola, M. (Centro Informazioni Studi Esperienze, Milan (Italy)). Sep 1968. 221p. (In Italian). Dep. NTIS (U. S. Sales Only).

The characteristics and differences of the various digital programs set up at CISE for the study of the thermo-hydraulic behavior of the primary circuit of the CIRENE nuclear reactor are summarized. The program PECVAR and the program CRIP are thoroughly described. The program PECVAR computes a detailed profile of pressure along the portion of circuit including the feeders, the channel with the bottom and top end-fittings, the risers and the steam-water separators. It also computes mean densities and mass and energy content in the channels and the safety margin with respect to the heat transfer crisis. The program CRIP computes the water and steam mass flow rate distribution among the various channels when, for a given geometry, the thermal power, the drum pressure, the steam feeder header pressure and the total mass flow rate of the recirculated water are imposed. In this calculation the overall energy and mass balance of the primary circuit is taken into account. 11 references. (auth)

47310 STUDY OF PRESSURE FLUCTUATIONS GENERATED BY AERODYNAMIC SINGULARITIES IN A CIRCUIT. Gibert, R. J. (CEN de Saclay, Gif-sur-Yvette, France). Bull. I Inform. Sci. Tech. (Paris); No. 167, 33-9 (Feb 1972). (In French).

Incidents that have occurred in various nuclear power reactors emphasize the problems that can arise from pressure and speed fluctuations in the primary and secondary circuits of the reactor and the corresponding induced vibrations. Certain singularities giving rise to fluctuations (local instabilities and the corresponding acoustic phenomena) were studied experimentally. A calculation on the transmission of such vibrations throughout the circuit is presented. (France)

47311 REINFORCED CONCRETE VESSELS FOR NUCLEAR POWER REACTORS: PRESENT STATUS AND FUTURE TRENDS. Balanescu, M. (Inst. of Atomic Physics, Bucharest). Stud. Cercet. Fiz.; 24: No. 4, 401-23 (1972). (In Rumanian).

French and English developments for the graphite-gas reactor line have demonstrated that the design and erection of a concrete calandria of the size required for power reactors do not raise any basic difficulty. Given the already known techniques and methods of building, it is considered that for high temperature or fast neutron commercial reactors, steel pressure vessels should no more be adopted, the future belonging to the prestressed reinforced concrete ones. The development status of the prestressed reinforced concrete calandria for water reactors, the difficulties pertaining to this line, as well as the proposed solutions, are reviewed. (auth)

Fuels and Fuel Elements

47312 (CONF-720802-1) HEAT TRANSFER PERFORMANCE OF SHORT TWISTED TAPES. Klepper, O. H. (Oak Ridge National Lab., Tenn.). [1971]. 23p. Dep. NTIS.

For Presentation at 13, national heat transfer conference; Denver, CO. (6 Aug 1972).

The use of a short twisted tape extending over a portion of a tube's heated length to enhance the heat transfer to a gas flowing within the tube was investigated with emphasis on the vortex decay region downstream from the tape. This problem is of particular significance to the cooling of high-performance gas-cooled nuclear reactors. Heat-transfer and pressure-drop data were obtained using nitrogen gas to cool tubes heated by an electric current passing axially through the tube wall. Tape twists (measured as the ratio of the tube length per 180-degree tape twist to the tube diameter) varied from 2.38 to 8.05; Reynolds moduli ranged from 20,000 to 380,000; and wall-to-gas absolute temperature ratios from 1.0 to 2.1. Results are presented. 13 references. (auth)

47313 SPACE-ENERGY-DEPENDENT PHYSICS PROBLEMS OF PLUTONIUM FUELS IN HTGRs. Asmussen, Keith E.; Waelti, Paul (Gulf General Atomic, San Diego, Calif.). Nucl. Technol.; 15: No. 3, 359-65 (Sep 1972).

The plutonium loadings for a test element for irradiation in the Peach Bottom HTGR were determined. The test element is a

graphite circular cylinder with an array of eight fuel rods in a circular pattern. The fuel rods contain dense spherical kernels of PuO_2 or $\text{PuO}_2/3 \text{ ThO}_2$. Because of the strong Pu resonances, the heterogeneity effects in the test element are strongly marked. A code for treating this rather complicated problem rigorously was not available. Thus the problem was simulated within a one-dimensional transport code by equating three collision probabilities. The heterogeneities were seen to result in significant grain self-shielding, rod self-shielding and mutual shielding of the rods. The Pu loading for the central fuel body of the test element was determined to be 19.0 grams. 12 references. (auth)

47314 STUDY OF VIBRATIONS OF THE CENTERING PARTS OF FUEL ELEMENT CARTRIDGES. Livolant, M.; Hoffmann, A. (CEN de Saclay, Gif-sur-Yvette, France). Bull. Inform. Sci. Tech. (Paris); No. 167, 29-32 (Feb 1972). (In French).

In a nuclear reactor of the natural uranium-graphite-gas type, the highly pressurized cooling fluid flows at great speed along the fuel cartridges which are held in the center of the channel by light alloy elements: the centering parts. The pressure fluctuations generated by the flow instability produce random stresses in these centering parts. Under particular conditions this has resulted in rupture due to metal fatigue. The extensive program undertaken by the CEA to study these phenomena is reviewed. (France)

Operation and Economics

47315 (CISE-R-178) ACTUAL VALUE METHOD EVALUATION OF THE EFFECTS OF FUEL COSTS ON THE COST OF POWER PRODUCED BY A NUCLEAR REACTOR FOR THE CASE OF BATCH REFUELING AND FOR THE CASE OF CONTINUOUS REFUELING. Astolfi, A. (Centro Informazioni Studi Esperienze, Milan (Italy)). Oct 1966. 59p. (In Italian). Dep. NTIS (U. S. Sales Only).

Using the present-worth method, the derivation of fuel cost expressions accounting for all fuel expenditures during the entire life of the power station in both cases of batch and continuous loading is shown. The resulting expressions are applied for calculating the fuel cost for the metallic and oxide uranium fuelled reactors of the CIRENE program. In the case of continuous loading a simple approximate procedure to estimate the capital invested in the fuel cycle and its interest charges is also presented. (auth)

47316 FAST BREEDER: SCHEDULE LENGTHENS, COST ESCALATES. Olds, F. C. Power Eng.; 76: No. 6, 33-5 (Jun 1972).

The timetable for the U. S. LMFBR program is examined. The effects of delays on estimated costs are discussed. (H.D.R.)

Safety and Siting

47317 (BNWL-1635) TECHNOLOGICAL CONSIDERATIONS IN EMERGENCY INSTRUMENTATION PREPAREDNESS. Phase II-A. Emergency Radiological and Meteorological Instrumentation Criteria for Reactors. Selby, J. M.; Andersen, B. V.; Carter, L. A.; Corley, J. P.; Elderkin, C. E.; Kathren, R. L.; Schwendiman, L. C.; Unruh, C. M.; Watson, E. C.; Woodruff, R. K. (Battelle Pacific Northwest Labs., Richland, Wash.). May 1972. Contract AT(45-1)-1830. 125p. Dep. NTIS.

Criteria for several instrument and instrumentation systems are presented. These include: (1) Liquid and gaseous radiological effluent monitoring instrumentation, both within the facility and in the environs. (2) Ambient radiation monitoring instrumentation. (3) Portable radiological monitoring instrumentation. (4) Meteorological instrumentation, both for synoptic and predictive purposes. The criteria are for instrumentation for determining the nature and extent of a radiological incident immediately after its occurrence. Emergency instrumentation criteria are examined for the contemporary reactor, in consonance with what is needed, reasonable, and achievable with the state-of-the-art, while at the same time identifying areas in which additional study is indicated. Before criteria for reactor emergency instrumentation could be developed, it was necessary to establish the maximum fission product inventory which could be involved in an accident, and also to make certain assumptions about the fraction of the inventory which might escape from the core. Once the fission products were characterized, instrumentation requirements for quickly assessing the release and predicting the trajectory and airborne concentration downwind were defined. Based on the quantity and characterization of airborne fission products which could escape, guides to instrument selection are given along with criteria for performance and placement, and some techniques presented for estimating the magnitude of the release. Instrumentation for meteorological parameter measurement is considered in detail because of the important role such measure-

ments will play in determining the direction of flow and dispersion in the atmosphere, and in the initial response by monitoring personnel. 115 references. (auth)

47318 (RT/PROT-72-4) DECAT: COMPUTER CODE FOR ANALYSIS OF THE LONG TERM PRESSURE AND TEMPERATURE TRANSIENT AFTER A LOSS OF COOLANT ACCIDENT IN A G.E. TYPE PRESSURE SUPPRESSION CONTAINMENT. Morici, A.; Santarossa, G. (Comitato Nazionale per l'Energia Nucleare, Rome (Italy)). 23 Feb 1972. 33p. (In Italian). Dep. NTIS (U. S. Sales Only).

The necessary input data are given and the use of the code for the calculation of the maximum values of pressure and temperature is illustrated with the aid of a test case. The analytical model is given. A listing is included and the nomenclature is given with the symbols used in the computer language. (auth)

Commercial Power Reactors

Refer also to abstract: 47579.

Physics and Core Engineering

Refer also to abstract: 47595.

47319 (GULF-GA-B-10959) COMPUTER SIMULATION OF PULSED-NEUTRON EXPERIMENTS FOR THE FORT ST. VRAIN HTGR. Pfeiffer, Wayne (Gulf General Atomic Co., San Diego, Calif.). 29 Mar 1972. Contract AT(04-3)-633. 40p. Dep. NTIS.

Prior to startup of the 330 MW Fort St. Vrain HTGR, pulsed-neutron experiments will be performed to determine the reactivity for various subcritical, control rod configurations. To determine the reactivity, the inhour method will be used. This method requires that the time response following the pulse exhibit the die-away of the fundamental prompt mode. For such a large reactor, it is important to carefully position the source and detector in order to minimize the contribution of the higher prompt modes to the time response. Moreover, for a power reactor, economics dictates that the optimum source and detector locations be determined before the experiments. This can be done by simulating the experiments on a computer. Such computer simulations have been performed for all ten control rod configurations for which pulsing is planned. The optimum source and detector locations have been determined as well as the expected time responses. 11 references. (auth)

Control and Instrumentation

47320 (DOCKET-50003-118) INDIAN POINT STATION, UNIT 1. Measures Taken to Repair Control Rod and Safety Implications of the Incident. (Consolidated Edison Co. of New York, Inc., New York). 10 Jul 1972. 9p. Dep. NTIS.

47321 (DOCKET-50244-109) ROBERT EMMETT GINNA NUCLEAR POWER PLANT, UNIT 1. Loss of Rod Control Logic System Power Supply. (Rochester Gas and Electric Corp., N. Y.). 7 Aug 1972. 2p. Dep. NTIS.

47322 (DOCKET-50247-77) INDIAN POINT NUCLEAR GENERATING UNIT 2. Summary Report of Control Rod Drive Line Malfunction. (Consolidated Edison Co. of New York, Inc., New York). 2 Aug 1972. 9p. Dep. NTIS.

47323 (DOCKET-50263-141) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. Supplemental Letter to the May 25, 1972 Report of Low APRM Gain Setting. (Northern States Power Co., Minneapolis, Minn.). 4 Aug 1972. 2p. Dep. NTIS.

47324 ANALOG STUDIES OF THE DYNAMICS OF A POWER PLANT CONTROL SYSTEM INCORPORATING A 440 MW(e) REACTOR. Golyand, L. N.; Dunaevskii, S. Ya.; Ostashenko, V. F.; Afanas'eva, T. M. This paper was summarized in *At. Energ. (USSR)*; 30: No. 1, 44 (Jan 1971) as paper No. 453/5856. (In Russian).

Results of analog simulation studies of the dynamics of the control system for a power plant unit incorporating the V-440 reactor are presented. The reliability of the results obtained in the analog simulation are confirmed by experiments conducted on real equipment (the first and second power units of the Novaya Voronezh' nuclear power plant). Transients in the control system of the first and second power units are compared to the simulation results, with the power output brought down to 70 and to 120 MW respectively. The simulation accuracy was estimated in neutron flux at 3-5% rating, in terms of pressure to about 1 atm, in terms of coolant temperature to within 1-2°C. The static characteristics

of the third power unit of the Novaya Voronezh' nuclear power plant (with the V-440 reactor) are presented along with the dynamic characteristics of the unit, the latter including the speed at which the reactor is brought up to full power, the extent to which reactor parameters are smoothed out automatically, and delay times, in response to reactivity surges and turbogenerator loads. The characteristics are reported for different power output levels and different reactivity coefficients with respect to coolant temperature. The dynamic characteristics of the third power unit in response to a stepwise drop in station power output from 100% to 50%, with a relay type controller acting on the reactor control rods in a constant-speed mode, are discussed. (auth)

Containment Systems, Waste Processing, and Radiation Protection

47325 (DOCKET-50265-101) QUAD-CITIES STATION, UNIT 2. Timer Found to be Inoperable. (Commonwealth Edison Co., Chicago, Ill.). 13 Aug 1972. 1p. Dep. NTIS.

47326 (DOCKET-50346-30) DAVIS-BESSE NUCLEAR POWER STATION. Shipping Damage to Containment Vessel Equipment Hatch. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 9 Aug 1972. 4p. Dep. NTIS.

Coolant System

47327 (DOCKET-50016-160) ENRICO FERMI ATOMIC POWER PLANT, UNIT 1. Execution of Periodic Test Activity Delayed by Piping Installation. (Power Reactor Development Co., Detroit, Mich.). 7 Jul 1972. 1p. Dep. NTIS.

47328 (DOCKET-50280-78) SURRY POWER STATION, UNIT 1. Authorization Request to Drain Reactor Coolant System Water and Lower Water Level in Reactor Vessel. (Virginia Electric and Power Co., Richmond). 4 Aug 1972. 2p. Dep. NTIS.

47329 (DOCKET-50341-44) ENRICO FERMI ATOMIC POWER PLANT, UNIT 2. Reactor Pressure Vessel Ultrasonic Indications. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 3 Aug 1972. 3p. Dep. NTIS.

47330 (DOCKET-50409-7) LA CROSSE BOILING WATER REACTOR. Forced Circulation System Behavior. (Dairyland Power Cooperative, La Crosse, Wis.). 14 Jul 1972. 2p. Dep. NTIS.

Auxiliary Systems and Engineered Safeguards

47331 (DOCKET-50237-250) DRESDEN NUCLEAR POWER STATION, UNIT 2. Failure of No. 2 Diesel Generator. (Commonwealth Edison Co., Morris, Ill. (USA)). 25 Jul 1972. 2p. Dep. NTIS.

47332 (DOCKET-50249-148) DRESDEN NUCLEAR POWER STATION, UNIT 3. Reactor Pressure Scram Switch Drifted. (Commonwealth Edison Co., Chicago, Ill.). 1 Aug 1972. 2p. Dep. NTIS.

47333 (DOCKET-50263-142) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. HPCI Turbine Check Valve Leakage. (Northern States Power Co., Minneapolis, Minn.). 4 Aug 1972. 1p. Dep. NTIS.

47334 (DOCKET-50263-143) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. Fast Start Failure of the High Pressure Coolant Injection System. (Northern States Power Co., Minneapolis, Minn.). 3 Aug 1972. 1p. Dep. NTIS.

47335 (DOCKET-50265-100) QUAD-CITIES STATION, UNIT 2. Failure of Inboard Primary Containment Isolation Valve in the Reactor Core Isolation Cooling System. (Commonwealth Edison Co., Cordova, Ill.). 3 Aug 1972. 2p. Dep. NTIS.

Operation and Economics

47336 (DOCKET-1151-57) ELK RIVER REACTOR. Operating Authorization DPRA-3, Amendment 1. (United Power Association, Elk River, Minn. (USA)). 14 Jul 1972. 32p. Dep. NTIS.

The operating authorization for the Elk River Power Reactor is requested to be transferred from Rural Cooperative Power Association to United Power Association by virtue of a company merger. (H.D.R.)

47337 (DOCKET-50003-119) INDIAN POINT STATION, UNIT 1. Secondary Coolant Activity in Excess of 1.5×10^{-2}

$\mu\text{Ci/ml.}$ (Consolidated Edison Co. of New York, Inc., New York). 19 Jul 1972. 3p. Dep. NTIS.

47338 (DOCKET-50010-114) DRESDEN NUCLEAR POWER STATION, UNIT 1. Analysis of Minimum Hydro Test Requirements and Fluence Calculations. (Commonwealth Edison Co., Chicago, Ill.). 3 Aug 1972. 8p. Dep. NTIS.

Information is presented concerning the minimum hydro test temperature and the beltline neutron flux calculations for the pressure vessel. (D.C.C.)

47339 (DOCKET-50016-164) ENRICO FERMI ATOMIC POWER PLANT, UNIT 1. Report for April 1972. (Power Reactor Development Co., Detroit, Mich.). 21 Jul 1972. 12p. (PRDC-EF-104). Dep. NTIS.

47340 (DOCKET-50016-165) ENRICO FERMI ATOMIC POWER PLANT, UNIT 1. Failure to Measure Radioactivity of Sodium at Specified Times. (Power Reactor Development Co., Detroit, Mich.). 2 Aug 1972. 1p. Dep. NTIS.

47341 (DOCKET-50029-189) YANKEE NUCLEAR POWER STATION. Operation Report 138 for Month of June 1972. (Yankee Atomic Electric Co., Westboro, Mass.). 21 Jul 1972. 9p. Dep. NTIS.

47342 (DOCKET-50133-91) HUMBOLDT BAY POWER PLANT, UNIT 3. Transmission Tripped out of Service by Forest Fire. (Pacific Gas and Electric Co., San Francisco, Calif.). 14 Aug 1972. 2p. Dep. NTIS.

47343 (DOCKET-50213-154) HADDAM NECK PLANT, UNIT 1. Operation Report No. 72-6 for the Month of June 1972. (Connecticut Yankee Atomic Power Co., Haddam). 27 Jul 1972. Contract AT(30-1)-3144. 20p. (NYO-3144-72). Dep. NTIS.

47344 (DOCKET-50219-196) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Relay Failure. (Jersey Central Power and Light Co., Morristown, N. J.). 26 Jun 1972. 1p. Dep. NTIS.

47345 (DOCKET-50219-197) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Surveillance Test Failures. (Jersey Central Power and Light Co., Morristown, N. J.). 26 Jun 1972. 2p. Dep. NTIS.

The failure of the rubber expansion joint between the emergency service water pump and the discharge piping is described. The maintenance required for the main steam isolation valve is discussed. (D.C.C.)

47346 (DOCKET-50219-199) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Experience With the Rod Worth Minimizer. (Jersey Central Power and Light Co., Morristown, N. J.). 14 Jul 1972. 3p. Dep. NTIS.

47347 (DOCKET-50219-203) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Surveillance Failure. (Jersey Central Power and Light Co., Morristown, N. J.). 11 Aug 1972. 2p. Dep. NTIS.

The failure of an isolation condenser steam line high flow differential pressure instrument to trip at the normal trip point is described. (D.C.C.)

47348 (DOCKET-50219-204) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Containment Spray Pump Failure. (Jersey Central Power and Light Co., Morristown, N. J.). 11 Aug 1972. 2p. Dep. NTIS.

47349 (DOCKET-50220-100) NINE MILE POINT NUCLEAR STATION, UNIT 1. Malfunctions Which Occurred During the Periodic Testing of the Core Spray Systems. (Niagara Mohawk Power Corp., Lycoming, N. Y.). 24 Jul 1972. 4p. Dep. NTIS.

47350 (DOCKET-50220-101) NINE MILE POINT NUCLEAR STATION, UNIT 1. Delay in Delivery of Nitrogen Gas. (Niagara Mohawk Power Corp., Syracuse, N. Y.). 1 Aug 1972. 2p. Dep. NTIS.

47351 (DOCKET-50220-103) NINE MILE POINT NUCLEAR STATION, UNIT 1. Technical Specifications, Proposed Changes. (Niagara Mohawk Power Corp., Syracuse, N. Y.). 10 Aug 1972. 18p. Dep. NTIS.

Revisions are proposed in the Nine Mile Point Power Reactor, Unit 1 technical specifications by the incorporation of changes in the plant operating organization. (H.D.R.)

47352 (DOCKET-50237-245) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Drift in Calibration of Some of EHC Oil Pressure Scram Switches. (Commonwealth Edison Co., Morris, Ill. (USA)). 25 Jul 1972. 2p. (DOCKET-50249-143). Dep. NTIS.

47353 (DOCKET-50237-247) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Failure to Monitor Plant Chimney Effluents July 26, 1972. (Commonwealth Edison Co., Morris, Ill. (USA)). 26 Jul 1972. 1p. (DOCKET-50249-145). Dep. NTIS.

47354 (DOCKET-50237-248) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Oil Low Pressure Scram Switches Found to Have Drifted. (Commonwealth Edison Co., Chicago, Ill.). 27 Jul 1972. 2p. (DOCKET-50249-146). Dep. NTIS.

47355 (DOCKET-50237-251) DRESDEN NUCLEAR POWER STATION, UNIT 2. Proposed Change 10, Supplement. (Commonwealth Edison Co., Chicago, Ill.). 31 Jul 1972. 6p. Dep. NTIS.

The results of the Dresden Power Reactor Unit 2 inservice inspection performed during the second partial refueling outage are presented. (D.C.C.)

47356 (DOCKET-50237-252) DRESDEN NUCLEAR POWER STATION, UNIT 2. Sample Pump Tripped. (Commonwealth Edison Co., Chicago, Ill.). 1 Aug 1972. 2p. Dep. NTIS.

Information is presented concerning the malfunction of the stack monitoring system sample pump. (D.C.C.)

47357 (DOCKET-50237-253) DRESDEN NUCLEAR POWER STATION, UNIT 2. Diesel Generator Failed to Operate. (Commonwealth Edison Co., Chicago, Ill.). 1 Aug 1972. 4p. Dep. NTIS.

47358 (DOCKET-50237-254) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. QUAD-CITIES STATION, UNITS 1 AND 2. Operating History of the Rod Worth Minimizers. (Commonwealth Edison Co., Chicago, Ill.). 3 Aug 1972. 5p. (DOCKET-50249-149; DOCKET-50254-111; DOCKET-50265-98). Dep. NTIS.

47359 (DOCKET-50244-106) ROBERT EMMETT GINNA NUCLEAR POWER PLANT, UNIT 1. Report of Exposure of an Individual to Radiation. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 25 Jul 1972. 2p. Dep. NTIS.

47360 (DOCKET-50245-143) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Abnormal Occurrence AO-72-18. (Millstone Point Co., Hartford, Conn.). 20 Jul 1972. 1p. Dep. NTIS.

Information is presented concerning the incorrect trip times for delay relays of the average power range monitoring system. (D.C.C.)

47361 (DOCKET-50245-146) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Semiannual Report, January 1, 1972-June 30, 1972. (Millstone Point Co., Hartford, Conn.). 9 Aug 1972. 63p. Dep. NTIS.

47362 (DOCKET-50249-151) DRESDEN NUCLEAR POWER STATION, UNIT 3. During Monthly Surveillance Testing the Inboard Isolation Valve Failed to Close. (Commonwealth Edison Co., Chicago, Ill.). 10 Aug 1972. 2p. Dep. NTIS.

47363 (DOCKET-50254-105) QUAD-CITIES STATION, UNIT 1. Failure of RCIC Motor Operated Valve MO-1-1301-49 to Open on Signal. (Commonwealth Edison Co., Cordova, Ill.). 10 Jul 1972. 2p. Dep. NTIS.

47364 (DOCKET-50254-108) QUAD-CITIES STATION, UNITS 1 AND 2. Interim Report on Condenser Pump Room Modifications, Change. (Commonwealth Edison Co., Chicago, Ill.). 24 Jul 1972. 2p. (DOCKET-50265-93). Dep. NTIS.

47365 (DOCKET-50261-101) H. B. ROBINSON, UNIT 2. Malfunction of "B" Steam Line Isolation Valve. (Carolina Power and Light Co., Raleigh, N. C.). 18 Jul 1972. 1p. Dep. NTIS.

47366 (DOCKET-50261-102) H. B. ROBINSON UNIT 2. Boron Injection Tank Dilution. (Carolina Power and Light Co., Raleigh, N. C.). 24 Jul 1972. 1p. Dep. NTIS.

47367 (DOCKET-50263-137) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. Turbine Trip at 100 percent Power and Subsequent Events. (Northern States Power Co., Minneapolis, Minn.). 20 Jul 1972. 7p. Dep. NTIS.

47368 (DOCKET-50263-139) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. High Pressure Coolant Injection System Failure. (Northern States Power Co., Minneapolis, Minn.). 27 Jul 1972. 2p. Dep. NTIS.

47369 (DOCKET-50263-140) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. Malfunction of the "D" Steam-line Safety/Relief Valve and Inoperability of Bellows Leakage

Monitoring Systems. (Northern States Power Co., Minneapolis, Minn.). 28 Jul 1972. 4p. Dep. NTIS.

47370 (DOCKET-50265-95) QUAD-CITIES STATION, UNIT 2. Fire in Two Electrical Cable Trays. (Commonwealth Edison Co., Cordova, Ill.). 24 Jul 1972. 6p. Dep. NTIS.

47371 (DOCKET-50265-96) QUAD-CITIES STATION, UNIT 2. Reactor Core Isolation Cooling System Inboard Steam Supply Valve Failed in Open Position. (Commonwealth Edison Co., Cordova, Ill.). 28 Jul 1972. 1p. Dep. NTIS.

47372 (DOCKET-50266-73) POINT BEACH NUCLEAR PLANT, UNITS 1 AND 2. Boric Acid Concentration in "A" Boric Acid Storage Tank Below Technical Specifications Limit. (Wisconsin Electric Power Co., Milwaukee). 26 Jul 1972. 3p. (DOCKET-50301-65). Dep. NTIS.

47373 (DOCKET-50266-75) POINT BEACH NUCLEAR PLANT, UNIT 1. Steam Generator Cladding Repair. (Wisconsin Electric Power Co., Milwaukee). 19 Jul 1972. 58p. Dep. NTIS.

47374 (DOCKET-50271-90) VERMONT YANKEE NUCLEAR POWER STATION. License Application, Amendment 31. (Vermont Yankee Nuclear Power Corp., Rutland). 19 Jul 1972. 195p. Dep. NTIS.

Financial data on the owners of the Vermont Yankee Power Reactor are presented. The 1971 Annual Reports for the Central Vermont Public Service Corporation, Green Mountain Power Corporation, New England Power Company, Connecticut Light and Power Company, Central Maine Power Company, Public Service Company of New Hampshire, Hartford Electric Light Company, Cambridge Electric Light Company, Eastern Utilities Associates, and the Western Massachusetts Electric Company are included. (D.C.C.)

47375 (DOCKET-50277-51) PEACH BOTTOM ATOMIC POWER STATION, UNITS 2 AND 3. Amendment 17, Substitution of a New Exhibit "U." (Philadelphia Electric Co., Pa.). 14 Jul 1972. 4p. (Docket-50278-51). Dep. NTIS.

Additional information is presented concerning estimated operating and permanent shutdown costs. (D.C.C.)

47376 (DOCKET-50293-80) PILGRIM NUCLEAR POWER STATION. Abnormal Occurrence AO-72-3. (Boston Edison Co., Mass.). 6 Jul 1972. 3p. Dep. NTIS.

An unplanned actuation of isolation valves for the reactor core isolation cooling system is described. (D.C.C.)

47377 (DOCKET-50293-82) PILGRIM NUCLEAR POWER STATION. Abnormal Occurrence AO-72-4. (Boston Edison Co., Mass.). 17 Jul 1972. 3p. Dep. NTIS.

Incorrect operating procedures resulting in the inoperability of the reactor water cleanup system high flow isolation differential pressure switches are described. (D.C.C.)

47378 (DOCKET-50293-83) PILGRIM NUCLEAR POWER STATION. Abnormal Occurrence AO-72-5. (Boston Edison Co., Mass.). 24 Jul 1972. 3p. Dep. NTIS.

Information is presented concerning the unplanned restart of recirculation pumps during a loss of power startup test. (D.C.C.)

47379 (DOCKET-50304-59) ZION STATION NUCLEAR POWER PLANT, UNIT 2. Leaks in Canopies of Two Control Rod Drive Assemblies. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 3 Aug 1972. 2p. Dep. NTIS.

47380 (DOCKET-50305-46) KEWAUNEE NUCLEAR POWER PLANT. Application for Construction Permit and Operating License, Amendment 20. (Wisconsin Public Service Corp., Green Bay). 18 Jul 1972. 5p. Dep. NTIS.

Additional information is presented concerning estimated operating and permanent shutdown costs. (D.C.C.)

47381 (DOCKET-50317-53) CALVERT CLIFFS NUCLEAR POWER PLANT, UNITS 1 AND 2. Failure of Polar Crane Upper Limit Switch. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 8 Aug 1972. 6p. (DOCKET-50318-51). Dep. NTIS.

47382 (DOCKET-50409-6) LA CROSSE BOILING WATER REACTOR. Incident Reports. (Dairyland Power Cooperative, La Crosse, Wis.). 20 Jul 1972. 9p. (DPC-72-5; DPC-72-6). Dep. NTIS.

Information is presented concerning revised bolting requirements for shutdown condenser partition, and fire in the forced circulation pump casing insulation. (D.C.C.)

47383 (DOCKET-50409-8) LA CROSSE BOILING WATER REACTOR. Incident Report, Hydrogen Detonation in Main Condenser Off-Gas System. (Dairyland Power Cooperative, La Crosse, Wis.). 2 Aug 1972. 5p. (DPC-72-7). Dep. NTIS.

47384 (DOCKET-50409-9) LA CROSSE BOILING WATER REACTOR. Monthly Operating Report, June 1972. (Dairyland Power Cooperative, La Crosse, Wis.). Jun 1972. Contract AT(11-1)-851. 31p. (DPC-851-44). Dep. NTIS.

47385 TRENDS IN CAPITAL COSTS OF GENERATING PLANTS. Roe, Kenneth A.; Young, William H. (Burns and Roe, Inc., New York). Power Eng.; 76: No. 6, 40-3(Jun 1972).

The rising trend of construction and capital costs for new electric power generating plants is examined. Some possible cost reduction solutions are discussed. (H.D.R.)

47386 ENVIRONMENTAL BENEFIT-COST ANALYSIS FOR NUCLEAR POWER GENERATION. Morrison, D. L. (Battelle Columbus Labs., Ohio); Males, R. H.; Duke, K. M.; Sharp, V. L.; Ritzman, R. L. Nucl. News; 15: No. 6, 50-7(Jun 1972).

The problems and procedures in preparing a cost-benefit analysis for nuclear power plants are discussed. The environmental considerations for the LaSalle County Station are used as an example of the practical limits and application of cost-benefit analyses to power reactors. (13 references) (H.D.R.)

47387 COMPARATIVE STUDY OF THE METHODOLOGY EMPLOYED IN ECONOMIC CALCULATIONS RELATING TO NUCLEAR POWER STATIONS IN THE SOVIET UNION AND WESTERN COUNTRIES. Koryakin, Yu. I. (Inst. of Atomic Energy, Moscow). At. Energy Rev.; 10: No. 2, 233-49(1972).

The methodological principles applied in Western countries and the Soviet Union in studying the economics of nuclear power stations are compared. It is shown that the mathematical formation to which these principles give rise coincides in many instances, although there are a number of differences in methodological approach, particularly as regards the use of mathematical operations, their conception and justification. 19 references. (auth)

Safety and Siting

Refer also to abstracts: 45535 and 45536.

47388 (DOCKET-50003-120) INDIAN POINT STATION, UNIT 1. Topical Report on General Electric Induction Motors; Inside Containment Low Pressure Core Cooling. (Consolidated Edison Co. of New York, Inc., New York). 3 Aug 1972. 177p. Dep. NTIS.

47389 (DOCKET-50010-113) DRESDEN NUCLEAR POWER STATION, UNIT 1. Proposed Change 17 to Operating License DPR-2. (Commonwealth Edison Co., Chicago, Ill.). 10 Jul 1972. 8p. Dep. NTIS.

The extent and consequences of hydrogen production in the containment of the Dresden Power Reactor, Unit 1 following a loss of coolant accident are evaluated. (H.D.R.)

47390 (DOCKET-50016-163) ENRICO FERMI ATOMIC POWER PLANT, UNIT 1. Answer to Show Cause Order and Amendment to Request for Extension of Expiration Date of Provisional Operating License. (Power Reactor Development Co., Detroit, Mich.). 20 Jul 1972. 43p. Dep. NTIS.

47391 (DOCKET-50029-190) YANKEE NUCLEAR POWER STATION. Withdrawal of Proposed Change 101. (Yankee Atomic Electric Co., Westboro, Mass.). 31 Jul 1972. 1p. Dep. NTIS.

47392 (DOCKET-50133-89) HUMBOLDT BAY POWER PLANT, UNIT 3. Technical Specifications, Proposed Change 40. (Pacific Gas and Electric Co., San Francisco, Calif.). 19 Jul 1972. 13p. Dep. NTIS.

Revised technical specifications are presented concerning the design characteristics of Type 3 Batch 4 fuel assemblies. (D.C.C.)

47393 (DOCKET-50155-117) BIG ROCK POINT NUCLEAR PLANT. Technical Specification Change Request 31, Errata. (Consumers Power Co., Jackson, Mich.). 27 Jun 1972. 2p. Dep. NTIS.

An illustration of a Reload G fuel rod bundle is presented. (D.C.C.)

47394 (DOCKET-50155-119) BIG ROCK POINT NUCLEAR PLANT. Technical Specifications, Proposed Change 33. (Consumers Power Co., Jackson, Mich.). 20 Jul 1972. 6p. Dep. NTIS.

Revised technical specifications are presented concerning test

requirements for the explosive valves of the liquid poison system. (D.C.C.)

47395 (DOCKET-50155-120) BIG ROCK POINT NUCLEAR PLANT. Technical Specifications, Proposed Change 34. (Consumers Power Co., Jackson, Mich.). 24 Jul 1972. 22p. Dep. NTIS.

Revised technical specifications are presented concerning the design characteristics of NFS-DA fuel bundles. (D.C.C.)

47396 (DOCKET-50213-150) HADDAM NECK PLANT, UNIT 1. Environmental Report, Appendix C, Addendum 2. (Connecticut Yankee Atomic Power Co., Hartford). 27 Jun 1972. 32p. Dep. NTIS.

The results of the Connecticut Yankee Power Reactor site environmental radiation survey program are presented for the period from July through December 1971. (H.D.R.)

47397 (DOCKET-50213-152) HADDAM NECK PLANT, UNIT 1. Environmental Report, Appendix E; Aquatic Benefit-cost. (Connecticut Yankee Atomic Power Co., Hartford). 27 Jun 1972. 77p. Dep. NTIS.

47398 (DOCKET-50219-200) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Final Safety Analysis Report, Amendment 70. (Jersey Central Power and Light Co., Morristown, N. J.). 19 Jul 1972. 31p. Dep. NTIS.

The final safety analysis report for the Oyster Creek Power Reactor Unit 1 is supplemented by the submission of requested additional data concerning maximum hurricane flood analysis. (D.C.C.)

47399 (DOCKET-50219-201) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Outside Radwaste Tank Activity. (Jersey Central Power and Light Co., Morristown, N. J.). 11 Jul 1972. 2p. Dep. NTIS.

47400 (DOCKET-50219-202) OYSTER CREEK NUCLEAR POWER PLANT, UNIT 1. Technical Specifications, Revised Pages Incorporating All Changes Through No. 12. (Jersey Central Power and Light Co., Morristown, N. J.). 1 Aug 1972. 38p. Dep. NTIS.

47401 (DOCKET-50220-98) NINE MILE POINT NUCLEAR STATION, UNIT 1. Environmental Report, Operating License Stage, Conversion to Full-Time Operating License. (Niagara Mohawk Power Corp., Syracuse, N. Y.). 5 Jul 1972. 440p. Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47402 (DOCKET-50237-244) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. QUAD-CITIES STATION, UNITS 1 AND 2. Results of Inspection of Torus Section Headers. (Commonwealth Edison Co., Chicago, Ill.). 5 Jul 1972. 8p. (DOCKET-50249-141; DOCKET-50254-104; DOCKET-50265-91). Dep. NTIS.

47403 (DOCKET-50237-246) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Steps Taken to Correct Items of Apparent Noncompliance Called to Attention in Letter of June 19, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 19 Jun 1972. 6p. (DOCKET-50249-144). Dep. NTIS.

Additional information is presented concerning the use of the railroad interlock doors of the secondary containment system. (D.C.C.)

47404 (DOCKET-50237-249) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Environmental Report; Benefit-Cost Analysis Report. (Commonwealth Edison Co., Chicago, Ill.). 31 Jul 1972. 45p. (DOCKET-50249-147). Dep. NTIS.

The environmental report for the Dresden Power Reactor Units 2 and 3 is supplemented by the submission of cost-benefit analysis for alternate designs. (D.C.C.)

47405 (DOCKET-50237-255) DRESDEN NUCLEAR POWER STATION, UNITS 2 AND 3. Technical Specifications, Proposed Change 18 (Unit 2); Technical Specifications, Change 9 (Unit 3). (Commonwealth Edison Co., Chicago, Ill.). 1 Aug 1972. 5p. (DOCKET-50249-150). Dep. NTIS.

Revised technical specifications are presented concerning the environmental radiation monitoring program. (D.C.C.)

47406 (DOCKET-50244-107) ROBERT EMMETT GINNA NUCLEAR POWER PLANT, UNIT 1. Monthly Report of Activity

Measurements. (Rochester Gas and Electric Corp., N. Y.). 3 Aug 1972. 2p. Dep. NTIS.

47407 (DOCKET-50245-139) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Abnormal Occurrence AO-72-17. (Millstone Point Co., Hartford, Conn.). 14 Jul 1972. 1p. Dep. NTIS.

Information is presented concerning incorrect trip points for pressure vessel high pressure scram switches. (D.C.C.)

47408 (DOCKET-50245-140) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Provisional Operating License DPR-21, Proposed Change 5. (Millstone Point Co., Hartford, Conn.). 12 Jul 1972. 9p. Dep. NTIS.

Additional information is presented concerning the 4.16 kV auxiliary bus system connection to the reserve station service transformer. (D.C.C.)

47409 (DOCKET-50245-141) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Technical Specifications, Proposed Change 6. (Millstone Point Co., Hartford, Conn.). 12 Jul 1972. 24p. Dep. NTIS.

Revised technical specifications are presented concerning administrative controls and operating procedures. (D.C.C.)

47410 (DOCKET-50245-144) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Technical Specifications, Proposed Change 8. (Millstone Point Co., Hartford, Conn.). 21 Jul 1972. 34p. Dep. NTIS.

Revised technical specifications are presented concerning the primary containment leak rate test requirements. (D.C.C.)

47411 (DOCKET-50245-145) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 1. Request Temporary Waiver of Technical Specification 3.5.F.6. (Millstone Point Co., Hartford, Conn.). 10 Aug 1972. 3p. Dep. NTIS.

Revised technical specifications are presented concerning maintenance procedures for the control rod drives. (D.C.C.)

47412 (DOCKET-50247-76) INDIAN POINT NUCLEAR GENERATING UNIT 2. Safety and Relief Valve Installation and Re-Analysis for ASME Class 1 and Class 2 Systems. (Consolidated Edison Co. of New York, Inc., New York). 13 Jul 1972. 22p. Dep. NTIS.

47413 (DOCKET-50250-84) TURKEY POINT PLANT, UNITS 3 AND 4. Final Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 368p. (DOCKET-50251-70). Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactors is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47414 (DOCKET-50250-85) TURKEY POINT PLANT, UNITS 3 AND 4. Facility Operating License DPR-31 With Appended Technical Specifications. (Directorate of Licensing (AEC), Washington, D. C.). 19 Jul 1972. 214p. (DOCKET-50251-71). Dep. NTIS.

47415 (DOCKET-50250-86) TURKEY POINT PLANT, UNITS 3 AND 4. Abnormal Occurrences 3-72-1 and 3-72-2. (Florida Power and Light Co.). 28 Jul 1972. 2p. (DOCKET-50251-73). Dep. NTIS.

During testing of the Turkey Point Power Reactor, Unit 3 under simulated safety injection conditions and loss of off-site power conditions the 4 kV busses failed to operate properly. Evaluations of the causes and consequences of these occurrences are presented. (H.D.R.)

47416 (DOCKET-50250-88) TURKEY POINT PLANT, UNITS 3 AND 4. Technical Specifications, Proposed Changes. (Florida Power and Light Co.). 11 Aug 1972. 4p. (DOCKET-50251-75). Dep. NTIS.

Revised technical specifications are presented concerning evaluation of power tilt limits. (D.C.C.)

47417 (DOCKET-50251-72) TURKEY POINT PLANT, UNIT 4. Structural Integrity Test Report and Containment Surveillance During the Structural Integrity Test. (Florida Power and Light Co.). 27 Jul 1972. 142p. Dep. NTIS.

47418 (DOCKET-50254-107) QUAD-CITIES STATION, UNITS 1 AND 2. Safety Analysis Report, Proposed Modification 72-1. (Commonwealth Edison Co., Chicago, Ill.). 20 Jul 1972. 12p. (DOCKET-50265-92). Dep. NTIS.

The safety analysis report for the Quad Cities Power Reactor Units 1 and 2 is supplemented by the submission of requested additional safety information. (D.C.C.)

47419 (DOCKET-50254-109) QUAD-CITIES STATION, UNITS 1 AND 2. Appendix A and Appendix B to DPR-29 and DPR-30, Proposed Change. (Commonwealth Edison Co., Chicago, Ill.). 25 Jul 1972. 110p. (DOCKET-50265-94). Dep. NTIS.

Revised and additional data are presented concerning technical specifications. (D.C.C.)

47420 (DOCKET-50254-110) QUAD-CITIES STATION, UNITS 1 AND 2. Final Report on Condenser Pump Room Modifications. (Commonwealth Edison Co., Chicago, Ill.). 31 Jul 1972. 8p. (DOCKET-50265-97). Dep. NTIS.

47421 (DOCKET-50254-112) QUAD-CITIES STATION, UNITS 1 AND 2. Operation of the Quad-Cities Diffuser Pipes. (Commonwealth Edison Co., Chicago, Ill.). 3 Aug 1972. 1p. (DOCKET-50265-99). Dep. NTIS.

47422 (DOCKET-50255-94) PALISADES PLANT. Application for Construction Permit and Operating License, Amendment 27. (Consumers Power Co., Jackson, Mich.). 14 Jul 1972. 2p. Dep. NTIS.

Additional information is presented concerning codes regulating the modifications for the radioactive waste disposal system. (D.C.C.)

47423 (DOCKET-50255-95) PALISADES PLANT. Technical Specifications, Request for Change 1. (Consumers Power Co., Jackson, Mich.). 14 Jul 1972. 13p. Dep. NTIS.

Revised technical specifications are presented concerning operating limits for the safety injection tanks. (D.C.C.)

47424 (DOCKET-50255-96) PALISADES PLANT. Environmental Impact of Plant Operation up to July 1, 1972. (Consumers Power Co., Jackson, Mich.). 19 Jul 1972. 360p. Dep. NTIS.

Information concerning radioactive and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological data for the reactor site are included. (D.C.C.)

47425 (DOCKET-50259-57) BROWNS FERRY NUCLEAR POWER STATION, UNITS 1, 2, AND 3. Final Safety Analysis Report, Amendment 35. (Tennessee Valley Authority, Chattanooga). 14 Jul 1972. 22p. (DOCKET-50260-53; DOCKET-50296-50). Dep. NTIS.

The preliminary safety analysis report for the Browns Ferry Power Reactor Units 1, 2, and 3 is supplemented by the submission of requested information concerning reactor kinetics. (D.C.C.)

47426 (DOCKET-50259-58) BROWNS FERRY NUCLEAR POWER STATION, UNITS 1, 2, AND 3. Final Safety Analysis Report, Amendment 36. (Tennessee Valley Authority, Chattanooga). 24 Jul 1972. 25p. (DOCKET-50260-54; DOCKET-50296-51). Dep. NTIS.

Revised and additional data are presented concerning the emergency electrical power systems for the reactors. (D.C.C.)

47427 (DOCKET-50259-59) BROWNS FERRY NUCLEAR POWER STATION, UNITS 1, 2, AND 3. Final Safety Analysis Report, Amendment 37. (Tennessee Valley Authority, Chattanooga). 28 Jul 1972. 108p. (DOCKET-50260-55; DOCKET-50296-52). Dep. NTIS.

The final safety analysis report for the Browns Ferry Power Reactor, Units 1, 2, and 3 is amended by the submission of revised and additional information. (H.D.R.)

47428 (DOCKET-50259-60) BROWNS FERRY NUCLEAR POWER STATION, UNITS 1, 2, AND 3. Final Safety Analysis Report, Amendment 38. (Tennessee Valley Authority, Chattanooga). 2 Aug 1972. 15p. (DOCKET-50260-56; DOCKET-50296-53). Dep. NTIS.

The final safety analysis report for the Browns Ferry Power Reactor, Units 1, 2, and 3 is amended by the submission of plant design changes. These changes incorporate charcoal filtering and pressurizing equipment in the main control room ventilation system and add physical barriers around the plant structures and service water pumps to provide protection from flooding. (H.D.R.)

47429 (DOCKET-50263-138) MONTICELLO NUCLEAR GENERATING PLANT, UNIT 1. Technical Specifications, Change Request 4. (Northern States Power Co., Minneapolis, Minn.). 24 Jul 1972. 5p. Dep. NTIS.

Revised technical specifications are presented concerning

the residual heat removal service water pump head requirements. (D.C.C.)

47430 (DOCKET-50266-74) POINT BEACH NUCLEAR PLANT, UNITS 1 AND 2. Discussion of Items of Apparent Non-compliance. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 3 Aug 1972. 10p. (DOCKET-50301-66). Dep. NTIS.

Additional information is presented concerning activities of the Off-Site Review Committee and radiation monitoring requirements. (D.C.C.)

47431 (DOCKET-50269-56) OCONEE NUCLEAR STATION, UNIT 1. Order Extending Provisional Construction Permit Completion Date. (Directorate of Licensing (AEC), Washington, D. C.). 27 Jun 1972. 2p. Dep. NTIS.

47432 (DOCKET-50269-60) OCONEE NUCLEAR STATION, UNITS 1, 2, AND 3. Final Safety Analysis Report, Amendment 33. (Duke Power Co., Charlotte, N. C.). 26 Jul 1972. 224p. (DOCKET-50270-48; DOCKET-50287-43). Dep. NTIS.

The final safety analysis report for the Oconee Power Reactor Units 1, 2, and 3 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47433 (DOCKET-50269-61) OCONEE NUCLEAR STATION, UNITS 1, 2, AND 3. Steps Taken With Respect to Item Brought to Attention in Letter Dated April 21, 1972. (Directorate of Regulatory Operations (AEC), Atlanta, Ga. (USA), Region II). 3 Aug 1972. 3p. (DOCKET-50270-50; DOCKET-50287-45). Dep. NTIS.

The provisions for use of an audit task force within the Steam Production Department of the Oconee Power Reactor, Units 1, 2, and 3 are described. (H.D.R.)

47434 (DOCKET-50270-49) OCONEE NUCLEAR STATION, UNITS 2 AND 3. Notice of Consideration of Issuance of Facility Operating Licenses and Notice of Opportunity for Hearing. (Directorate of Licensing (AEC), Washington, D. C.). 3 Aug 1972. 5p. (DOCKET-50287-44). Dep. NTIS.

47435 (DOCKET-50271-88) VERMONT YANKEE NUCLEAR POWER STATION. Final Environmental Statement. (Vermont Yankee Nuclear Power Corp., Rutland). Jul 1972. 357p. Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for facility operations is presented. (D.C.C.)

47436 (DOCKET-50271-91) VERMONT YANKEE NUCLEAR POWER STATION. Technical Specifications, Proposed Change. (Vermont Yankee Nuclear Power Corp., Rutland). 6 Jul 1972. 7p. Dep. NTIS.

Revised technical specifications are presented concerning planned tests to be performed under the existing 1 percent low power license. (D.C.C.)

47437 (DOCKET-50271-92) VERMONT YANKEE NUCLEAR POWER STATION. Operating License DPR-28, Appendix B. (Vermont Yankee Nuclear Power Corp., Rutland). 7 Jul 1972. 26p. Dep. NTIS.

Technical specifications concerning chemical and thermal discharges from the reactor are presented. (D.C.C.)

47438 (DOCKET-50271-93) VERMONT YANKEE NUCLEAR POWER STATION. Changes in Steam Tunnel Ventilation System. (Vermont Yankee Nuclear Power Corp., Rutland). 25 Jul 1972. 4p. Dep. NTIS.

47439 (DOCKET-50271-94) VERMONT YANKEE NUCLEAR POWER STATION. Technical Specifications, Change 1. (Directorate of Licensing (AEC), Washington, D. C.). 24 Jul 1972. 3p. Dep. NTIS.

Revised technical specifications are presented concerning bypass conditions for protection system scrams. (D.C.C.)

47440 (DOCKET-50271-95) VERMONT YANKEE NUCLEAR POWER STATION. Technical Specifications, Proposed Change 1, Supplement 1. (Vermont Yankee Nuclear Power Corp., Rutland). 25 Jul 1972. 18p. Dep. NTIS.

Revised technical specifications are presented concerning the off gas radioactive waste processing system. (D.C.C.)

47441 (DOCKET-50272-29) SALEM NUCLEAR GENERATING STATION, UNITS 1 AND 2. Final Safety Analysis Re-

port, Amendment 12. (Public Service Electric and Gas Co., Newark, N. J.). 31 Jul 1972. 173p. (DOCKET-50311-29). Dep. NTIS.

The final safety analysis report for the Salem Power Reactor, Units 1 and 2 is amended by the submission of revised and requested additional information. (H.D.R.)

47442 (DOCKET-50272-30) SALEM NUCLEAR GENERATING STATION, UNITS 1 AND 2. Final Safety Analysis Report, Amendment 13. (Public Service Electric and Gas Co., Newark, N. J.). 2 Jun 1972. 154p. (DOCKET-50311-30). Dep. NTIS.

The final safety analysis report for the Salem Power Reactor, Units 1 and 2 is amended by the submission of requested additional information. (H.D.R.)

47443 (DOCKET-50272-31) SALEM NUCLEAR GENERATING STATION, UNITS 1 AND 2. Final Safety Analysis Report, Volume 6. (Public Service Electric and Gas Co., Newark, N. J.). 7 Aug 1972. 109p. (DOCKET-50311-31). Dep. NTIS.

The final safety analysis report for the Salem Power Reactor, Units 1 and 2 is supplemented by the responses to specific questions. (H.D.R.)

47444 (DOCKET-50272-32) SALEM NUCLEAR GENERATING STATION, UNITS 1 AND 2. Final Safety Analysis Report, Volume 7. (Public Service Electric and Gas Co., Newark, N. J.). 7 Aug 1972. 213p. (DOCKET-50311-32). Dep. NTIS.

The final safety analysis report for the Salem Power Reactor, Units 1 and 2 is supplemented by the responses to specific questions. (H.D.R.)

47445 (DOCKET-50272-33) SALEM NUCLEAR GENERATING STATION, UNITS 1 AND 2. Environmental Report, Amendment 2. (Public Service Electric and Gas Co., Newark, N. J.). 9 Aug 1972. 206p. (DOCKET-50311-33). Dep. NTIS.

47446 (DOCKET-50275-22) DIABLO CANYON NUCLEAR POWER PLANT, UNITS 1 AND 2. Environmental Report, Supplement 2, Volume 1. (Pacific Gas and Electric Co., San Francisco, Calif.). 28 Jul 1972. 484p. (DOCKET-50323-20). Dep. NTIS.

The environmental report for the Diablo Canyon Power Reactor Units 1 and 2 is supplemented by the submission of revised and additional information for the facility. (D.C.C.)

47447 (DOCKET-50275-23) DIABLO CANYON NUCLEAR POWER PLANT, UNITS 1 AND 2. Environmental Report, Supplement 2, Volume 2. (Pacific Gas and Electric Co., San Francisco, Calif.). Jul 1972. 716p. (DOCKET-50323-21). Dep. NTIS.

47448 (DOCKET-50277-52) PEACH BOTTOM ATOMIC POWER STATION, UNITS 2 AND 3. Environmental Report, Operating License Stage, Supplement 5. (Philadelphia Electric Co., Pa.). 28 Jul 1972. 11p. (DOCKET-50278-52). Dep. NTIS.

Requested additional information is presented concerning transmission lines, chemical effluents, and thermal effluents. (D.C.C.)

47449 (DOCKET-50280-77) SURRY POWER STATION, UNITS 1 AND 2. Topical Report on General Electric Vertical Induction Motors, Inside Containment Recirculation Spray Pump Motors, Addendum. (Virginia Electric and Power Co., Richmond). 14 Jul 1972. 48p. (DOCKET-50281-75). Dep. NTIS.

47450 (DOCKET-50282-59) PRAIRIE ISLAND NUCLEAR GENERATING PLANT, UNIT 1. Application for AEC Class 104b Operating License, Amendment 13. (Northern States Power Co., Minneapolis, Minn.). 30 Dec 1971. 3p. Dep. NTIS.

A request is presented for extension of the date for completion of the Prairie Island Power Reactor, Unit 1 to February 1, 1973. (H.D.R.)

47451 (DOCKET-50282-67) PRAIRIE ISLAND NUCLEAR GENERATING PLANT, UNITS 1 AND 2. Final Safety Analysis Report, Amendment 19. (Northern States Power Co., Minneapolis, Minn.). 14 Jul 1972. 713p. (DOCKET-50306-64). Dep. NTIS.

The final safety analysis report for the Prairie Island Power Reactor Units 1 and 2 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47452 (DOCKET-50282-68) PRAIRIE ISLAND NUCLEAR GENERATING PLANT, UNITS 1 AND 2. Report of Corrosion Problem in Refueling Water Storage Tanks. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 20 Jul 1972. 4p. (DOCKET-50306-65). Dep. NTIS.

47453 (DOCKET-50282-69) PRAIRIE ISLAND NUCLEAR GENERATING PLANT, UNITS 1 AND 2. Final Safety Analysis

Report, Amendment 20. (Northern States Power Co., Minneapolis, Minn.). 4 Aug 1972. 252p. (DOCKET-50306-66). Dep. NTIS.

The final safety analysis report for the Prairie Island Power Reactor Units 1 and 2 is amended by the submission of revised and requested safety information. (D.C.C.)

47454 (DOCKET-50285-68) FORT CALHOUN STATION, UNIT 1. Safety Evaluation. (Directorate of Licensing (AEC), Washington, D. C.). 14 Aug 1972. 99p. Dep. NTIS.

47455 (DOCKET-50286-46) INDIAN POINT NUCLEAR GENERATING UNIT 3. Final Facility Description and Safety Analysis Report, Amendment 18, Supplement 4. (Consolidated Edison Co. of New York, Inc., New York). 30 Jun 1972. 36p. Dep. NTIS.

The final safety analysis report for the Indian Point Power Reactor, Unit 3 is amended by the submission of revised and requested additional information. (H.D.R.)

47456 (DOCKET-50286-47) INDIAN POINT NUCLEAR GENERATING UNIT 3. Final Safety Analysis Report, Supplement 5. (Consolidated Edison Co. of New York, Inc., New York). 20 Jul 1972. 51p. Dep. NTIS.

The final safety analysis report for the Indian Point Power Reactor Unit 3 is supplemented by the submission of requested additional information. (D.C.C.)

47457 (DOCKET-50286-48) INDIAN POINT NUCLEAR GENERATING UNIT 3. Final Facility Description and Safety Analysis Report, Supplement 6. (Consolidated Edison Co. of New York, Inc., New York). 1 Aug 1972. 104p. Dep. NTIS.

The final safety analysis report for the Indian Point Power Reactor, Unit 3 is supplemented by the submission of revised and requested additional information. (H.D.R.)

47458 (DOCKET-50286-49) INDIAN POINT NUCLEAR GENERATING UNIT 3. Final Facility Description and Safety Analysis Report, Supplement 7. (Consolidated Edison Co. of New York, Inc., New York). 1 Aug 1972. 400p. Dep. NTIS.

The final safety analysis report for the Indian Point Power Reactor Unit 3 is supplemented by the submission of requested additional safety information. (D.C.C.)

47459 (DOCKET-50289-61) THREE MILE ISLAND NUCLEAR STATION, UNIT 1. Action Taken to Correct Item of Nonconformance Found on May 4, 1972 Inspection. (Division of Compliance (AEC), Newark, N. J. Region I). 28 Jun 1972. 4p. Dep. NTIS.

Additional information is presented concerning the construction of the containment structure ring girder. (D.C.C.)

47460 (DOCKET-50289-62) THREE MILE ISLAND NUCLEAR STATION, UNIT 1. Report of Significant Deficiency/Thin Walled Valves. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 11 Jul 1972. 4p. Dep. NTIS.

47461 (DOCKET-50289-63) THREE MILE ISLAND NUCLEAR STATION, UNIT 1. Final Safety Analysis Report, Amendment 27. (Metropolitan Edison Co., Reading, Pa.). 14 Aug 1972. 136p. Dep. NTIS.

The final safety analysis report for the Three Mile Island Power Reactor Unit 1 is amended by the submission of revised and additional safety information. (D.C.C.)

47462 (DOCKET-50295-57) ZION STATION NUCLEAR POWER PLANT, UNITS 1 AND 2. Steps Taken to Correct Items of Apparent Noncompliance in Letter of June 12, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 25 Jul 1972. 10p. (DOCKET-50304-56). Dep. NTIS.

Additional information is presented concerning welding procedures, the quality assurance requirements for the main steam isolation valve, and qualifications for nondestructive testing personnel. (D.C.C.)

47463 (DOCKET-50295-58) ZION STATION NUCLEAR POWER PLANT, UNITS 1 AND 2. Accident Report, April 24, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 25 Jul 1972. 8p. (DOCKET-50304-57). Dep. NTIS.

Information is presented concerning reactor components involved in a warehouse fire at the construction site. (D.C.C.)

47464 (DOCKET-50295-59) ZION STATION NUCLEAR POWER PLANT, UNITS 1 AND 2. Steps Taken to Correct Items of Apparent Noncompliance Brought to Attention in Letter Dated June 23, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 9 Aug 1972. 7p. (DOCKET-50304-58). Dep. NTIS.

Additional information is presented concerning the installation and inspection of instrumentation and electric cables. (D.C.C.)

47465 (DOCKET-50298-43) COOPER NUCLEAR STATION. Environmental Report, Supplement 2. (Nebraska Public Power District, Columbus). 26 Jun 1972. 46p. Dep. NTIS.

The environmental report for the Cooper Power Reactor is supplemented by the submission of requested additional data. (D.C.C.)

47466 (DOCKET-50298-46) COOPER NUCLEAR STATION. Final Safety Analysis Report, Amendment 13. (Nebraska Public Power District, Columbus). 18 Jul 1972. 256p. Dep. NTIS.

The final safety analysis report for the Cooper Power Reactor is supplemented by the submission of requested additional safety information. (D.C.C.)

47467 (DOCKET-50298-47) COOPER NUCLEAR STATION. Final Safety Analysis Report, Amendment 11. (Nebraska Public Power District, Columbus). 24 Jul 1972. 8p. Dep. NTIS.

Facility modifications are described for the spent fuel storage pool, the reactor building closed cooling water system, fire protection system failure, and battery ventilation system. (D.C.C.)

47468 (DOCKET-50298-48) COOPER NUCLEAR STATION. Environmental Report, Supplement 3. (Nebraska Public Power District, Columbus). 4 Aug 1972. 41p. Dep. NTIS.

47469 (DOCKET-50301-64) POINT BEACH NUCLEAR PLANT, UNIT 2. Facility Operating License No. DPR-27, Amendment 2. (Directorate of Licensing (AEC), Washington, D. C.). 28 Jul 1972. 11p. Dep. NTIS.

47470 (DOCKET-50305-44) KEWAUNEE NUCLEAR POWER PLANT. Final Safety Analysis Report, Amendment 19. (Wisconsin Public Service Corp., Green Bay). 30 Jun 1972. 197p. Dep. NTIS.

The final safety analysis report for the Kewaunee Power Reactor is supplemented by the submission of revised and additional safety information. (D.C.C.)

47471 (DOCKET-50305-45) KEWAUNEE NUCLEAR POWER PLANT. Draft Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 249p. Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operation is presented. (D.C.C.)

47472 (DOCKET-50305-47) KEWAUNEE NUCLEAR POWER PLANT. Steps Taken to Correct Item of Apparent Noncompliance Noted in Letter Dated July 17, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 8 Aug 1972. 6p. Dep. NTIS.

Additional information is presented concerning the installation of Class 1 instrument controllers which did not meet quality control requirements. (D.C.C.)

47473 (DOCKET-50309-61) MAINE YANKEE ATOMIC POWER STATION. Final Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 231p. Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47474 (DOCKET-50309-62) MAINE YANKEE ATOMIC POWER STATION. Final Safety Analysis Report, Amendment 35. (Maine Yankee Atomic Power Co., Westboro, Mass.). 27 Jul 1972. 96p. Dep. NTIS.

The final safety analysis report for the Maine Yankee Power Reactor is supplemented by the submission of revised and additional safety information. (D.C.C.)

47475 (DOCKET-50309-63) MAINE YANKEE ATOMIC POWER STATION. Environmental Qualification of Vital Instrumentation and Electrical Cables, and Penetration Connectors. (Maine Yankee Atomic Power Co., Westboro, Mass.). 27 Jul 1972. 66p. Dep. NTIS.

47476 (DOCKET-50309-64) MAINE YANKEE ATOMIC POWER STATION. Seismic Design Review. (Maine Yankee

Atomic Power Co., Westboro, Mass.). 27 Jul 1972. 113p. Dep. NTIS.

47477 (DOCKET-50309-65) MAINE YANKEE ATOMIC POWER STATION. Seismic Qualification of Vital Instrument and Electrical Equipment. (Maine Yankee Atomic Power Co., Westboro, Mass.). 27 Jul 1972. 391p. Dep. NTIS.

47478 (DOCKET-50309-66) MAINE YANKEE ATOMIC POWER STATION. Final Safety Analysis Report, Amendment 36. (Maine Yankee Atomic Power Co., Westboro, Mass.). 10 Aug 1972. 33p. Dep. NTIS.

Revised information is presented concerning LOCA analysis for the Maine Yankee Power Reactor. (D.C.C.)

47479 (DOCKET-50312-41) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Environmental Report, Amendment 2. (Sacramento Municipal Utility District, Calif.). 30 Jun 1972. 196p. Dep. NTIS.

The environmental report for the Rancho Seco Power Reactor is supplemented by the submission of revised and additional information for the facility. (D.C.C.)

47480 (DOCKET-50312-42) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Final Safety Analysis Report, Amendment 12. (Sacramento Municipal Utility District, Calif.). 26 Jun 1972. 497p. Dep. NTIS.

The final safety analysis report for the Rancho Seco Power Reactor, Unit 1 is amended by the submission of revised and requested additional information. (H.D.R.)

47481 (DOCKET-50312-43) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Report on Minor Imperfections Found in Pipe Welds. (Sacramento Municipal Utility District, Calif.). 30 May 1972. 39p. Dep. NTIS.

47482 (DOCKET-50312-45) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Final Safety Analysis Report, Amendment 13. (Sacramento Municipal Utility District, Calif.). 19 Jul 1972. 43p. Dep. NTIS.

The final safety analysis report for the Rancho Seco Power Reactor is supplemented by the submission of requested additional safety information. (D.C.C.)

47483 (DOCKET-50312-46) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Weld Defects Found in Steam Generator Lateral Support. (Sacramento Municipal Utility District, Calif.). 24 Jul 1972. 17p. Dep. NTIS.

47484 (DOCKET-50312-47) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Report on Minor Imperfections Found in Pipe Welds. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 20 Jul 1972. 39p. Dep. NTIS.

47485 (DOCKET-50312-48) RANCHO SECO NUCLEAR GENERATING STATION, UNIT 1. Weld Defects Found in Steam Generator Lateral Support. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 4 Aug 1972. 19p. Dep. NTIS.

47486 (DOCKET-50313-44) ARKANSAS NUCLEAR ONE, UNIT 1. Environmental Report, Supplement 2. (Arkansas Power and Light Co., Little Rock). 28 Jun 1972. 44p. Dep. NTIS.

The environmental report for the Arkansas Nuclear One Power Reactor is supplemented by the submission of cost-benefit data for alternate designs for the facility. (D.C.C.)

47487 (DOCKET-50313-45) ARKANSAS NUCLEAR ONE, UNIT 1. Environmental Report, Supplement 3. (Arkansas Power and Light Co., Little Rock). 25 Jul 1972. 91p. Dep. NTIS.

The environmental report for the Arkansas Nuclear One Power Reactor is supplemented by the submission of revised and additional information for the facility. (D.C.C.)

47488 (DOCKET-50313-46) ARKANSAS NUCLEAR ONE, UNIT 1. Final Safety Analysis Report, Amendment 27. (Arkansas Power and Light Co., Little Rock). 4 Aug 1972. 167p. Dep. NTIS.

The final safety analysis report for the Arkansas Nuclear One Power Reactor is supplemented by the submission of revised and requested safety information. (D.C.C.)

47489 (DOCKET-50315-65) DONALD C. COOK NUCLEAR PLANT, UNITS 1 AND 2. Environmental Report, Supplement 3. (Indiana and Michigan Power Co., New York). 27 Jun 1972. 30p. (DOCKET-50316-65). Dep. NTIS.

The environmental report for the Donald C. Cook Power

Reactor Units 1 and 2 is supplemented by the submission of additional information on cost-benefit analysis for alternate designs. (D.C.C.)

47490 (DOCKET-50315-68) DONALD C. COOK NUCLEAR PLANT, UNITS 1 AND 2. Final Safety Analysis Report, Amendment 25. (Indiana and Michigan Power Co., New York). 21 Jul 1972. 176p. (DOCKET-50316-68). Dep. NTIS.

The final safety analysis report for the Donald C. Cook Power Reactor, Units 1 and 2 is amended by the submission of revised and requested additional information. (H.D.R.)

47491 (DOCKET-50315-69) DONALD C. COOK NUCLEAR PLANT, UNITS 1 AND 2. Defective Welding on Ice Condenser System Positioning Frames. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 31 Jul 1972. 5p. (DOCKET-50316-69). Dep. NTIS.

47492 (DOCKET-50315-70) DONALD C. COOK NUCLEAR PLANT, UNITS 1 AND 2. Deficiency in Radiography Techniques. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 14 Jul 1972. 7p. (DOCKET-50316-70). Dep. NTIS.

47493 (DOCKET-50315-71) DONALD C. COOK NUCLEAR PLANT, UNITS 1 AND 2. Final Safety Analysis Report, Amendment 26. (Indiana and Michigan Power Co., New York). 7 Aug 1972. 13p. (DOCKET-50316-71). Dep. NTIS.

Information is presented concerning the long term pressure transient in the containment structure following a LOCA. (D.C.C.)

47494 (DOCKET-50321-46) EDWIN I. HATCH NUCLEAR PLANT, UNIT 1. Final Safety Analysis Report, Amendment 17. (Georgia Power Co., Atlanta). 26 Jun 1972. 119p. Dep. NTIS.

Requested additional information is presented concerning containment systems and auxiliary equipment. Stress analysis for the pressure vessel is included. (D.C.C.)

47495 (DOCKET-50321-48) EDWIN I. HATCH NUCLEAR PLANT, UNIT 1. Steps Taken to Correct Items of Apparent Non-compliance Which Were Brought to Attention in Letter of July 5, 1972. (Directorate of Regulatory Operations (AEC), Atlanta, Ga. (USA), Region II). 3 Aug 1972. 8p. Dep. NTIS.

Requested information is presented concerning quality assurance requirements for construction, welding procedures, and storage requirements for Class 1 equipment. (D.C.C.)

47496 (DOCKET-50324-39) BRUNSWICK STEAM ELECTRIC PLANT, UNITS 1 AND 2. Expected Doses Which Might Result From Purging Containment. (Carolina Power and Light Co., Raleigh, N. C.). 31 Jul 1972. 3p. (DOCKET-50325-40). Dep. NTIS.

47497 (DOCKET-50324-40) BRUNSWICK STEAM ELECTRIC PLANT, UNITS 1 AND 2. Environmental Report, Amendment 2. (Carolina Power and Light Co., Raleigh, N. C.). 14 Aug 1972. 111p. (DOCKET-50325-41). Dep. NTIS.

The environmental report for the Brunswick Power Reactor Units 1 and 2 is amended by the submission of revised cost-benefit information. (D.C.C.)

47498 (DOCKET-50331-48) DUANE ARNOLD ENERGY CENTER. Environmental Report Revised, Amendment 1. (Iowa Electric Light and Power Co., Cedar Rapids). 6 Jul 1972. 93p. Dep. NTIS.

The environmental report for the Duane Arnold Power Reactor is supplemented by the submission of cost-benefit data for alternate design for the facility. (D.C.C.)

47499 (DOCKET-50331-49) DUANE ARNOLD ENERGY CENTER. Final Safety Analysis Report, Amendment 2. (Iowa Electric Light and Power Co., Cedar Rapids). 10 Jul 1972. 518p. Dep. NTIS.

The final safety analysis report for the Duane Arnold Power Reactor is amended by the submission of revised and requested additional information. (H.D.R.)

47500 (DOCKET-50331-51) DUANE ARNOLD ENERGY CENTER. Final Safety Analysis Report, Amendment 3. (Iowa Electric Light and Power Co., Cedar Rapids). Jul 1972. 172p. Dep. NTIS.

The final safety analysis report for the Duane Arnold Power Reactor is amended by the submission of revised and requested additional information. (H.D.R.)

47501 (DOCKET-50333-50) JAMES A. FITZPATRICK NUCLEAR POWER PLANT. Final Facility Description and Safety Analysis Report, Supplement 9. (Power Authority of the State of New York). 7 Aug 1972. 172p. Dep. NTIS.

The final safety analysis report for the James A. Fitzpatrick Power Reactor is supplemented by the submission of requested additional safety information. (D.C.C.)

47502 (DOCKET-50333-51) JAMES A. FITZPATRICK NUCLEAR POWER PLANT. Environmental Report, Supplement 3. (Power Authority of the State of New York). 11 Aug 1972. 22p. Dep. NTIS.

The environmental report for the James A. Fitzpatrick Power Reactor is supplemented by the submission of requested additional information for the facility. (D.C.C.)

47503 (DOCKET-50334-31) BEAVER VALLEY POWER STATION, UNIT 1. Action Taken to Correct Items of Nonconformance and Noncompliance Found on Inspection of April 18-20, 1972. (Directorate of Regulatory Operations (AEC), Newark, N. J. Region I). 10 Jul 1972. 8p. Dep. NTIS.

Additional information is presented concerning the quality assurance requirements for sampling and testing concrete, electrical wiring installation procedures, and storage requirements for electrical equipment. (D.C.C.)

47504 (DOCKET-50335-29) HUTCHINSON ISLAND PLANT, UNIT 1. Environmental Report, Supplement 5. (Florida Power and Light Co.). 28 Jun 1972. 34p. Dep. NTIS.

The environmental report for the Hutchinson Island Power Reactor, Unit 1 is supplemented by the submission of cost-benefit information for the facility. (H.D.R.)

47505 (DOCKET-50335-30) HUTCHINSON ISLAND PLANT, UNIT 1. Environmental Report, Supplement 6. (Florida Power and Light Co.). 6 Jul 1972. 17p. Dep. NTIS.

The alternatives are analyzed for the proposed transmission line crossings of the Indian River near the Hutchinson Island Power Reactor, Unit 1. (H.D.R.)

47506 (DOCKET-50335-32) HUTCHINSON ISLAND PLANT, UNIT 1. Environmental Report, Supplement 7. (Florida Power and Light Co.). 14 Jul 1972. 11p. Dep. NTIS.

Requested additional information is presented concerning the construction of the circulating water system intake and discharge pipelines. (D.C.C.)

47507 (DOCKET-50336-35) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 2. Operating License Application. (Millstone Point Co., Hartford, Conn.). 15 Aug 1972. 55p. Dep. NTIS.

Financial data on the owners of the Millstone Point Power Reactor Unit 2 are presented. (D.C.C.)

47508 (DOCKET-50336-36) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 2. Final Safety Analysis Report, Volume 1. (Millstone Point Co., Hartford, Conn.). 10 Aug 1972. 802p. Dep. NTIS.

Information on the Millstone Point Power Reactor Unit 2 is presented under the following broad topics: AEC general design criteria; quality assurance program; site characteristics; reactor kinetics; stress analysis for pressure vessel internals; cooling system; and structures. (D.C.C.)

47509 (DOCKET-50336-37) MILLSTONE POINT NUCLEAR POWER STATION, UNIT 2. Final Safety Analysis Report, Volume 2. (Millstone Point Co., Hartford, Conn.). 1972. 720p. Dep. NTIS.

Information on the Millstone Point Power Reactor Unit 2 is presented under the following broad topics: engineered safeguards; instrumentation; control systems; and electrical systems. (D.C.C.)

47510 (DOCKET-50338-47) NORTH ANNA POWER STATION, UNITS 1, 2, 3, AND 4. Environmental Report Supplement, Revision 2. (Virginia Electric and Power Co., Richmond). 23 Jun 1972. 42p. (DOCKET-50339-44; DOCKET-50404-17; DOCKET-50405-17). Dep. NTIS.

The environmental report for the North Anna Power Reactor Units 1, 2, 3, and 4 is supplemented by the submission of cost-benefit analysis for design alternatives for the facility. (D.C.C.)

47511 (DOCKET-50338-48) NORTH ANNA POWER STATION, UNITS 1 AND 2. Environmental Considerations. (Virginia Electric and Power Co., Richmond). 29 Sep 1970. 187p. (DOCKET-50339-45). Dep. NTIS.

The effects of impoundment of the North Anna River by the dam and the effects of Units 1 and 2 of the North Anna Power Reactor on the impoundment are discussed. (H.D.R.)

47512 (DOCKET-50338-49) NORTH ANNA POWER STATION, UNITS 1 AND 2. Request for Permission to Resume Work on Transmission Line. (Virginia Electric and Power

Co., Richmond). 1 Aug 1972. 3p. (DOCKET-50339-46). Dep. NTIS.

47513 (DOCKET-50341-42) ENRICO FERMI ATOMIC POWER PLANT, UNIT 2. Cracks in the Basement Floor Slab of the Reactor Building. (Detroit Edison Co., Mich.). 2 Jun 1972. 3p. Dep. NTIS.

47514 (DOCKET-50341-43) ENRICO FERMI ATOMIC POWER PLANT, UNIT 2. Deformation of Containment Drywell. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 17 Jul 1972. 12p. Dep. NTIS.

47515 (DOCKET-50344-29) TROJAN NUCLEAR PLANT. Initial Decision on Environmental Review. (Portland General Electric Co., Oreg.). 23 May 1972. 27p. Dep. NTIS.

The factors considered in the Atomic Safety and Licensing Board decision to allow Trojan Power Reactor construction activities to continue pending completion of the environmental review are presented. (H.D.R.)

47516 (DOCKET-50344-30) TROJAN NUCLEAR PLANT. Supplemental Environmental Report, Amendment 3. (Portland General Electric Co., Oreg.). 24 Jul 1972. 50p. Dep. NTIS.

The environmental report for the Trojan Power Reactor Unit 1 is supplemented by the submission of revised and additional information for the facility. (D.C.C.)

47517 (DOCKET-50346-27) DAVIS-BESSE NUCLEAR POWER STATION. Steps Taken to Correct Items of Apparent Nonconformance Noted in Letter Dated May 22, 1972. (Directorate of Regulatory Operations (AEC), Glen Ellyn, Ill. Region III). 22 Jun 1972. 5p. Dep. NTIS.

Additional information is presented concerning corrected inspection and certification procedures for component cooling and service water piping. (D.C.C.)

47518 (DOCKET-50346-29) DAVIS-BESSE NUCLEAR POWER STATION. Environmental Report, Supplement, Amendment 1. (Toledo Edison Co., Ohio). 14 Jul 1972. 74p. Dep. NTIS.

The environmental report for the Davis-Besse Power Reactor is supplemented by the submission of revised and additional information for the facility. (D.C.C.)

47519 (DOCKET-50348-42) JOSEPH M. FARLEY NUCLEAR PLANT, UNITS 1 AND 2. Steps Taken to Correct Items of Apparent Noncompliance Brought to Attention in Letter Dated June 21, 1972. (Directorate of Regulatory Operations (AEC), Atlanta, Ga. (USA). Region II). 3 Aug 1972. 13p. (DOCKET-50364-36). Dep. NTIS.

Actions taken to correct activities in vendor shop inspection of welds and in inspection of concrete placement for the Joseph M. Farley Power Reactor, Units 1 and 2 are described. (H.D.R.)

47520 (DOCKET-50352-27) LIMERICK GENERATING STATION, UNITS 1 AND 2. Environmental Report, Construction Permit Stage (Revised), Supplement 1. (Philadelphia Electric Co., Pa.). 11 Aug 1972. 56p. (DOCKET-50353-27). Dep. NTIS.

The environmental report for Limerick Power Reactor, Units 1 and 2 is supplemented by the submission of responses to requests for additional information. (H.D.R.)

47521 (DOCKET-50354-25) NEWBOLD ISLAND NUCLEAR GENERATING STATION, Unit 1. Inspection of Reactor Pressure Vessel Fabrication at Hitachi, Ltd. (Directorate of Regulatory Operations (AEC), Washington, D. C.). 10 Jul 1972. 11p. Dep. NTIS.

47522 (DOCKET-50361-25) SAN ONOFRE NUCLEAR GENERATING STATION, UNITS 2 AND 3. Application for a Construction Permit and a Class 104(b) License, Amendment 15. (Southern California Edison Co., Rosemead, Calif.). 30 Jun 1972. 179p. (DOCKET-50362-24). Dep. NTIS.

Financial data on the owners of the San Onofre Power Reactor Units 2 and 3 are presented. The 1971 Annual Reports for the Southern California Edison Company and the San Diego Gas and Electric Company are included. (D.C.C.)

47523 (DOCKET-50361-26) SAN ONOFRE NUCLEAR GENERATING STATION, UNITS 2 AND 3. Preliminary Safety Analysis Report, Amendment 14. (Southern California Edison Co., Rosemead, Calif.). 29 Jun 1972. 285p. (DOCKET-50362-25). Dep. NTIS.

The preliminary safety analysis report for the San Onofre Power Reactor, Units 2 and 3 is amended by the submission of revised and requested additional information. (H.D.R.)

47524 (DOCKET-50363-30) FORKED RIVER NUCLEAR GENERATING STATION, UNIT 1. Preliminary Safety Analysis Report, Amendment 19. (Jersey Central Power and Light Co., Morristown, N. J.). 15 Jul 1972. 74p. Dep. NTIS.

The preliminary safety analysis report for the Forked River Power Reactor Unit 1 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47525 (DOCKET-50363-31) FORKED RIVER NUCLEAR GENERATING STATION, UNIT 1. Preliminary Safety Analysis Report, Amendment 20. (Jersey Central Power and Light Co., Morristown, N. J.). 31 Jul 1972. 50p. Dep. NTIS.

The preliminary safety analysis report for the Forked River Power Reactor, Unit 1 is amended by the submission of revised data. (H.D.R.)

47526 (DOCKET-50363-32) FORKED RIVER NUCLEAR GENERATING STATION, UNIT 1. Environmental Report, Correction of Cost-Benefit Analysis. (Jersey Central Power and Light Co., Morristown, N. J.). 2 Aug 1972. 1p. Dep. NTIS.

47527 (DOCKET-50363-34) FORKED RIVER NUCLEAR GENERATING STATION, UNIT 1. Environmental Report, Commercial Fishing in Barnegat Bay. (Jersey Central Power and Light Co., Morristown, N. J.). 8 Aug 1972. 3p. Dep. NTIS.

47528 (DOCKET-50366-29) EDWIN I. HATCH NUCLEAR PLANT, UNIT 2. Preliminary Safety Analysis Report, Amendment 16. (Georgia Power Co., Atlanta). 23 Jun 1972. 12p. Dep. NTIS.

Additional data are presented concerning design parameters of the pressure vessel pedestal. (D.C.C.)

47529 (DOCKET-50366-31) EDWIN I. HATCH NUCLEAR PLANT, UNIT 2. Safety Evaluation. (Directorate of Licensing (AEC), Washington, D. C.). 21 Jul 1972. 182p. Dep. NTIS.

47530 (DOCKET-50366-32) EDWIN I. HATCH NUCLEAR PLANT, UNIT 2. Department of Justice Opinion With Respect to Monopoly. (Department of Justice, Washington, D. C. Antitrust Div.). 2 Aug 1972. 12p. Dep. NTIS.

47531 (DOCKET-50366-33) EDWIN I. HATCH NUCLEAR PLANT, UNIT 2. Steps Taken to Correct Items of Apparent Noncompliance Brought to Attention in Letter Dated June 28, 1972. (Directorate of Regulatory Operations (AEC), Atlanta, Ga. (USA). Region II). 3 Aug 1972. 6p. Dep. NTIS.

Requested information is presented concerning design change notices and quality assurance requirements. (D.C.C.)

47532 (DOCKET-50368-31) ARKANSAS NUCLEAR ONE, UNIT 2. Draft Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 210p. Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactor is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47533 (DOCKET-50369-26) McGUIRE NUCLEAR STATION, UNITS 1 AND 2. Application for Licenses, Amendment 12. (Duke Power Co., Charlotte, N. C.). 18 Oct 1971. 7p. (DOCKET-50370-26). Dep. NTIS.

Cost data and a forecast of construction expenditures for the McGuire Power Reactor, Units 1 and 2 are presented. (H.D.R.)

47534 (DOCKET-50369-28) McGUIRE NUCLEAR STATION, UNITS 1 AND 2. Draft Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 302p. (DOCKET-50370-28). Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactors is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47535 (DOCKET-50373-24) LA SALLE COUNTY STATION, UNITS 1 AND 2. Draft Environmental Statement. (Directorate of Licensing (AEC), Washington, D. C.). Jul 1972. 133p. (DOCKET-50374-23). Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactors is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47536 (DOCKET-50373-25) LA SALLE COUNTY STATION, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 10. (Commonwealth Edison Co., Chicago, Ill.). 26 Jul 1972. 87p. (DOCKET-50374-24). Dep. NTIS.

The preliminary safety analysis report for the La Salle County Power Reactor Units 1 and 2 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47537 (DOCKET-50376-34) AGUIRRE NUCLEAR PLANT, UNIT 1. Preliminary Facility Analysis Report, Amendment 18. (Puerto Rico Water Resources Authority, San Juan). 30 Jun 1972. 125p. Dep. NTIS.

The preliminary safety analysis report for the Aguirre Power Reactor, Unit 1 is amended by the submission of requested additional information. (H.D.R.)

47538 (DOCKET-50382-25) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Preliminary Safety Analysis Report, Amendment 17. (Louisiana Power and Light Co., New Orleans). 17 Jul 1972. 173p. Dep. NTIS.

The preliminary safety analysis report for the Waterford Power Reactor Unit 3 is supplemented by the submission of requested additional safety information. (D.C.C.)

47539 (DOCKET-50382-27) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Environmental Report, Supplement 1. (Louisiana Power and Light Co., New Orleans). 24 Jul 1972. 87p. Dep. NTIS.

The environmental report for the Waterford Power Reactor Unit 3 is supplemented by the submission of requested information for the facility. (D.C.C.)

47540 (DOCKET-50382-28) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Preliminary Safety Analysis Report, Amendment 19. (Louisiana Power and Light Co., New Orleans). 31 Jul 1972. 61p. Dep. NTIS.

The preliminary safety analysis report for the Waterford Power Reactor Unit 3 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47541 (DOCKET-50382-29) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Application for Construction Exemption. (Louisiana Power and Light Co., New Orleans). 11 May 1972. 27p. Dep. NTIS.

47542 (DOCKET-50382-30) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Basic Data for a Source Term Calculation. (Louisiana Power and Light Co., New Orleans). 17 May 1972. 21p. Dep. NTIS.

47543 (DOCKET-50382-31) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Environmental Report, Supplement 2. (Louisiana Power and Light Co., New Orleans). 15 Aug 1972. 75p. Dep. NTIS.

The environmental report for the Waterford Power Reactor Unit 3 is supplemented by the submission of revised and requested information for the facility. (D.C.C.)

47544 (DOCKET-50382-32) WATERFORD STEAM ELECTRIC STATION, UNIT 3. Preliminary Safety Analysis Report, Amendment 20. (Louisiana Power and Light Co., New Orleans). 15 Aug 1972. 85p. Dep. NTIS.

The preliminary safety analysis report for the Waterford Power Reactor Unit 3 is amended by the submission of revised and requested safety information. (D.C.C.)

47545 (DOCKET-50387-19) SUSQUEHANNA STEAM ELECTRIC STATION, UNITS 1 AND 2. Environmental Report, Revised July 1972, Vol. 1. (Pennsylvania Power and Light Co., Allentown). 7 Jul 1972. 339p. (DOCKET-50388-19). Dep. NTIS.

Information concerning chemical, radioactive, and thermal discharges from the reactors is presented. The effects of reactor operations on the environment are analyzed. Ecological, geological, hydrological, and meteorological data for the reactor site are included. Cost-benefit analysis for reactor operations is presented. (D.C.C.)

47546 (DOCKET-50387-20) SUSQUEHANNA STEAM ELECTRIC STATION, UNITS 1 AND 2. Environmental Report, Revised July 1972, Vol. 2. (Pennsylvania Power and Light Co., Allentown). 7 Jul 1972. 365p. (DOCKET-50388-20). Dep. NTIS.

Ecological, geological, hydrological, and meteorological data for the facility site are presented. (D.C.C.)

47547 (DOCKET-50387-21) SUSQUEHANNA STEAM ELECTRIC STATION, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 5, Revision. (Pennsylvania Power and Light Co., Allentown). Jul 1972. 2p. (DOCKET-50388-21). Dep. NTIS.

47548 (DOCKET-50387-22) SUSQUEHANNA STEAM ELECTRIC STATION, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 8. (Pennsylvania Power and Light Co., Allentown). 4 May 1972. 29p. (DOCKET-50388-22). Dep. NTIS.

The preliminary safety analysis report for the Susquehanna Power Reactor, Units 1 and 2 is amended by the submission of requested additional information. (H.D.R.)

47549 (DOCKET-50387-23) SUSQUEHANNA STEAM ELECTRIC STATION, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 9. (Pennsylvania Power and Light Co., Allentown). 28 Jul 1972. 28p. (DOCKET-50388-23). Dep. NTIS.

The preliminary safety analysis report of the Susquehanna Power Reactor, Units 1 and 2 is amended by the submission of revised information. (H.D.R.)

47550 (DOCKET-50390-17) WATTS BAR NUCLEAR PLANT, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 9. (Tennessee Valley Authority, Chattanooga). 19 Jun 1972. 82p. (DOCKET-50391-17). Dep. NTIS.

Maximum possible flood conditions for the reactor site are analyzed. (D.C.C.)

47551 (DOCKET-50390-18) WATTS BAR NUCLEAR PLANT, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 11. (Tennessee Valley Authority, Chattanooga). 19 Jul 1972. 319p. (DOCKET-50391-18). Dep. NTIS.

The preliminary safety analysis report for the Watts Bar Power Reactor Units 1 and 2 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47552 (DOCKET-50390-19) WATTS BAR NUCLEAR PLANT, UNITS 1 AND 2. Application for License, Amendment 12. (Tennessee Valley Authority, Chattanooga). 11 Aug 1972. 51p. (DOCKET-50391-19). Dep. NTIS.

The application for construction permits for the Watts Bar Power Reactor, Units 1 and 2 is amended by the submission of revised organizational and financial information. (H.D.R.)

47553 (DOCKET-50395-22) VIRGIL C. SUMMER NUCLEAR STATION, UNIT 1. Preliminary Safety Analysis Report, Amendment 10. (South Carolina Electric and Gas Co., Columbia). 10 Jul 1972. 494p. Dep. NTIS.

The preliminary safety analysis report of the Virgil C. Summer Power Reactor, Unit 1 is amended by the submission of revised and requested additional information. (H.D.R.)

47554 (DOCKET-50395-24) VIRGIL C. SUMMER NUCLEAR STATION, UNIT 1. Preliminary Safety Analysis Report, Amendment 12. (South Carolina Electric and Gas Co., Columbia). 1 Aug 1972. 97p. Dep. NTIS.

The preliminary safety analysis report for the Virgil C. Summer Power Reactor, Unit 1 is amended by the submission of additional information. (H.D.R.)

47555 (DOCKET-50395-25) VIRGIL C. SUMMER NUCLEAR STATION, UNIT 1. Preliminary Safety Analysis Report, Amendment 13. (South Carolina Electric and Gas Co., Columbia). 4 Aug 1972. 45p. Dep. NTIS.

The preliminary safety analysis report for the Virgil C. Summer Power Reactor Unit 1 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47556 (DOCKET-50397-19) HANFORD NO. 2 NUCLEAR POWER PLANT. Preliminary Safety Analysis Report, Amendment 7. (Washington Public Power Supply System, Richland). 7 Jul 1972. 465p. Dep. NTIS.

Revised and additional data are presented concerning the reactor site. Requested information is also presented concerning design parameters and operations aspects of the reactor. (D.C.C.)

47557 (DOCKET-50397-20) HANFORD NO. 2 NUCLEAR POWER PLANT. Preliminary Safety Analysis Report, Amendment 8. (Washington Public Power Supply System, Richland). 31 Jul 1972. 46p. Dep. NTIS.

The preliminary safety analysis report for the Hanford Power Reactor Unit 2 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47558 (DOCKET-50397-21) HANFORD NO. 2 NUCLEAR POWER PLANT. Preliminary Safety Analysis Report, Amendment 9. (Washington Public Power Supply System, Richland). 7 Aug 1972. 138p. Dep. NTIS.

The preliminary safety analysis report for the Hanford Power Reactor Unit 2 is amended by the submission of revised and requested safety information. (D.C.C.)

47559 (DOCKET-50398-11) MENDOCINO POWER PLANT, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 7. (Pacific Gas and Electric Co., San Francisco, Calif.). 5 Jul 1972. 257p. (DOCKET-50399-11). Dep. NTIS.

The preliminary safety analysis report for the Mendocino Power Reactor, Units 1 and 2 is amended by the submission of revised and requested additional information. (H.D.R.)

47560 (DOCKET-50398-15) MENDOCINO POWER PLANT, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 9. (Pacific Gas and Electric Co., San Francisco, Calif.). 21 Jul 1972. 22p. (DOCKET-50399-15). Dep. NTIS.

The preliminary safety analysis report for the Mendocino Power Reactor Units 1 and 2 is supplemented by the submission of requested information on site geology and seismology. (D.C.C.)

47561 (DOCKET-50398-16) MENDOCINO POWER PLANT, UNITS 1 AND 2. Preliminary Safety Analysis Report, Amendment 10. (Pacific Gas and Electric Co., San Francisco, Calif.). 2 Aug 1972. 313p. (DOCKET-50399-16). Dep. NTIS.

The preliminary safety analysis report for the Mendocino Power Reactor Units 1 and 2 is supplemented by the submission of revised and additional safety information. (D.C.C.)

47562 (DOCKET-50398-17) MENDOCINO POWER PLANT, UNITS 1 AND 2. Opinion of Department of Justice with Respect to Monopolistic Activity. (Department of Justice, Washington, D. C. Antitrust Div.). 2 Aug 1972. 16p. (DOCKET-50399-17). Dep. NTIS.

47563 (DOCKET-50400-19) SHEARON HARRIS NUCLEAR POWER PLANT, UNITS 1, 2, 3, AND 4. Preliminary Safety Analysis Report, Amendment 10. (Carolina Power and Light Co., Raleigh, N. C.). 17 Jul 1972. 640p. (DOCKET-50401-19; DOCKET-50402-19; DOCKET-50403-19). Dep. NTIS.

Revised and additional data are presented concerning design parameters and operations aspects of the Shearon Harris Power Reactor Units 1, 2, 3, and 4. (D.C.C.)

47564 (DOCKET-50400-20) SHEARON HARRIS NUCLEAR POWER PLANT, UNITS 1, 2, 3, AND 4. Preliminary Safety Analysis Report, Amendment 11. Environmental Report, Volume 2. (Carolina Power and Light Co., Raleigh, N. C.). 24 Jul 1972. 105p. (DOCKET-50401-20; DOCKET-50402-20; DOCKET-50403-20). Dep. NTIS.

The environmental report for the Shearon Harris Power Reactor Units 1, 2, 3, and 4 is supplemented by the submission of requested additional information for the facility. (D.C.C.)

47565 (DOCKET-50400-21) SHEARON HARRIS NUCLEAR POWER PLANT, UNITS 1, 2, 3, AND 4. Preliminary Safety Analysis Report, Amendment 12. (Carolina Power and Light Co., Raleigh, N. C.). 7 Jul 1972. 350p. (DOCKET-50401-21; DOCKET-50402-21; DOCKET-50403-21). Dep. NTIS.

The preliminary safety analysis report for the Shearon Harris Power Reactor Units 1, 2, 3, and 4 is supplemented by the submission of requested additional safety information. (D.C.C.)

47566 (DOCKET-50404-18) NORTH ANNA POWER STATION, UNITS 3 AND 4. Preliminary Safety Analysis Report, Amendment 5. (Virginia Electric and Power Co., Richmond). 23 Jun 1972. 125p. (DOCKET-50405-18). Dep. NTIS.

The preliminary safety analysis report of the North Anna Power Reactor, Units 3 and 4 is amended by the submission of revised and additional data. (H.D.R.)

47567 (DOCKET-50404-19) NORTH ANNA POWER STATION, UNITS 3 AND 4. Preliminary Safety Analysis Report, Amendment 6. (Virginia Electric and Power Co., Richmond). 3 Jul 1972. 19p. (DOCKET-50405-19). Dep. NTIS.

The preliminary safety analysis report for the North Anna Power Reactor, Units 3 and 4 is amended by the submission of requested additional information. (H.D.R.)

47568 (DOCKET-50404-20) NORTH ANNA POWER STATION, UNITS 3 AND 4. Department of Justice Opinion With Respect to Monopolistic Action. (Department of Justice, Washington, D. C. Antitrust Div.). 1 May 1972. 9p. (DOCKET-50405-20). Dep. NTIS.

47569 (DOCKET-50409-4) LA CROSSE BOILING WATER REACTOR. Technical Specifications, Change in Plant Organization. (Dairyland Power Cooperative, La Crosse, Wis.). 12 Jul 1972. 3p. Dep. NTIS.

47570 (DOCKET-50409-5) LA CROSSE BOILING WATER REACTOR. Application of Conversion of the Provisional Operating Authorization to a Provisional Operating License, Amend-

ment 1. (Dairyland Power Cooperative, La Crosse, Wis.). 21 Jul 1972. 3p. Dep. NTIS.

Additional information is presented concerning financial data, radiation sources, and fuel inventory. (D.C.C.)

47571 ARE LICENSING DELAYS REALLY NECESSARY? Males, Rene H. (Commonwealth Edison Co., Chicago). Power; 116: No. 6, 80-1 (Jun 1972).

A simplified licensing procedure and an improved method for coordinating activities between the numerous agencies involved in licensing power plant construction are recommended. (auth)

47572 SIMULATORS. Turner, James (Electronic Associates, Inc., Princeton, N. J.). Power Eng.; 76: No. 6, 26-32 (Jun 1972).

The development of simulators for nuclear power plant operator training is examined, and features of existing facilities are described. (H.D.R.)

47573 EXPERIENCE IN THE LICENSING OF NUCLEAR POWER PLANTS. Bliss, Henry E.; Stiede, Wayne L. (Commonwealth Edison Co., Chicago). Nucl. News; 15: No. 6, 59-62; 65 (Jun 1972).

Bringing a nuclear power plant into a utility system involves the acquisition of a number of permits and licenses. The permit chronology of nuclear units is described and some comments and ideas on three aspects of the licensing procedure which contribute to delay are presented. (auth)

47574 CRITERIA FOR SELECTING THE LOCATION OF NUCLEAR PLANTS. Amaldi, Ugo Jr.; Campos Venuti, Gloria; Frullani, Salvatore; Maiani, Luciano; Tabet, Eugenio. Ann. Ist. Super. Sanita; 7: 626-46 (1971). (In Italian).

Site selection criteria regarding nuclear power plants are summarized for a number of countries: USA, Japan, Germany, Canada, Great Britain, and Italy. The Italian criteria are amplified for purposes of discussion; examination of a proposed site for a nuclear plant raises questions of ecology, legislative disposition (protection of nearby human zones), natural dangers such as earthquakes, the quality of components of the nuclear installation, and the capability of the site relative to emergency handling. 30 references. (K.S.W.)

Steam and Power Conversion System

47575 (DOCKET-50220-102) NINE MILE POINT NUCLEAR STATION, UNIT 1. Results of Test on Main Steam Line Isolation Valves. (Niagara Mohawk Power Corp., Syracuse, N. Y.). 1 Aug 1972. 2p. Dep. NTIS.

47576 (DOCKET-50244-108) ROBERT EMMETT GINNA NUCLEAR POWER PLANT, UNIT 1. Pressurizer Safety Valves. (Rochester Gas and Electric Corp., N. Y.). 17 Jul 1972. 207p. Dep. NTIS.

A design study is presented for modifications to the Robert E. Ginna Power Reactor, Unit 1 main steam safety valve supports. Also included is a dynamic analysis of the pressurizer safety valve discharge piping. (H.D.R.)

Propulsion and Auxiliary Power Reactors

47577 NERVA FLIGHT ENGINE CONTROL SYSTEM DESIGN. Norman, H. H. (Westinghouse Electric Corp., Sacramento, Calif.); Parziale, E. A.; Saluja, J. K.; Schenz, R. F. Jr. Nucl. Technol.; 15: No. 3, 447-54 (Sep 1972).

Modern control system theory was applied to the design of the control system for the NERVA nuclear rocket engine. Multiloop classical controls design approach was used previously in the engine test program. The configuration and operation of the engine system with the resulting high degree of coupling and the multivariable nature of the system establishes a need for modern control techniques with considerable advantages over classical methods. The design procedure consists of system model linearization, system simplification, and the quadratic optimal control design. Transient performance results were obtained from digital implementation of the control system. 11 references. (auth)

47578 ANALOGOVOE MODELIROVANIE SUDOVYKH YADERNYKH PAROPROIZVODYASHCHIKH USTANOVOK. (Analog Models of Marine Nuclear Steam Generating Plants). Sergeev, V. L. Leningrad; Izdatel'stvo Sudostroenie (1971). 352p.

Methods of model simulation of non-steady-state processes in

marine nuclear steam-producing installations are examined on electronic analog computers. Model simulation of neutron processes, non-steady-state heat transfer in reactors, heat exchange, steam generators, and heat lag in turboducts are described. Basic units of steam generating plants are treated as systems of distributed parameters in an analysis of their performances. 81 references. (K.S.W.)

Process Heat and Desalination Reactors

47579 PRELIMINARY REMARKS ABOUT LARGE-SCALE DESALINATION AND DUAL PURPOSE PLANTS. Ahmad, Mohammad (Pakistan Atomic Energy Commission, Karachi). *Nucleus* (Karachi); 8: No. 4, 43-54(1971).

The growing need of saline water conversion vis-a-vis the global inventory of water substance is briefly discussed. Water shortage also exists in Pakistan and areas where desalination technology can be called upon to provide either the only source of potable water or to supplement the existing means of fresh water supply are described. The status of desalting methods and preliminary factors in the choice of large-scale desalination, especially by means of dual-purpose plants, are presented. (auth)

Research and Test Reactors

Refer also to abstract: 45673.

47580 (CISE-R-259) ANALOG COMPUTER MODELS FOR THE SAFETY EVALUATION OF THE CART FUEL TEST FACILITY. Leoni, G.; van Erp, J. B. (Centro Informazioni Studi Esperienze, Milan (Italy)). Dec 1967. 29p. Dep. NTIS (U. S. Sales Only).

Analog computer models used for the evaluation of some of the safety aspects of the CART fuel test facility are presented. Information is included on the consequence on the CART test-section of loss-of-flow accidents affecting directly the CART facility and the consequence on the CART test-section of a reactivity accident at power in the ESSOR reactor. (auth)

47581 (CEA-N-1483) MASURCA: STUDY OF PLUTONIUM-GRAPHITE AND URANIUM-GRAPHITE ASSEMBLIES. Tretiakoff, O.; Barberger, M.; Boyer, J.; (and others) (Commissariat a l'Energie Atomique, Saclay (France). Centre d'Etudes Nucleaires). Sep 1971. 132p. (In French). Dep. NTIS (U. S. Sales Only).

During the December 1966 to March 1969 period, assemblies with either 30% enriched uranium, or a uranium-plutonium-iron alloy and graphite mixtures were studied in a fast neutron critical mock-up (MASURCA) at Cadarache. The operation of this mock-up enabled problems associated with the reactor operation to be examined. Cores were chosen because of their simplicity and their spectrum, which approximates that of fast power reactors. Measurement techniques were developed for these cores. Future experimental programs concerning the assemblies are described. Critical size, buckling by fission rate mappings, and reactivity coefficient determination using the oscillation technique are described. The results concerning different measured parameters are presented and various values for the reactivity coefficients are given. 8 references. (France)

47582 (DOCKET-50231-120) SOUTHWEST EXPERIMENTAL FAST OXIDE REACTOR. Modification Proposed for Decommissioning Plan. (General Electric Co., Sunnyvale, Calif. Breeder Reactor Dept.). 28 Jun 1972. 5p. Dep. NTIS.

47583 (DOCKET-50231-121) SOUTHWEST EXPERIMENTAL FAST OXIDE REACTOR. Request for Issuance of Dismantling Order. (General Electric Co., Sunnyvale, Calif. Breeder Reactor Dept.). 18 Jul 1972. 2p. Dep. NTIS.

47584 (DOCKET-50231-122) SOUTHWEST EXPERIMENTAL FAST OXIDE REACTOR. Technical Specifications, Change 13. (Directorate of Licensing (AEC), Washington, D. C.). 21 Jul 1972. 1p. Dep. NTIS.

Revised technical specifications are presented concerning personnel requirements for site security. (D.C.C.)

47585 (EURFNR-1008) INFLUENCE OF FISSION NEUTRON SPECTRA ON INTEGRAL NUCLEAR QUANTITIES OF FAST REACTORS. Kieffhaber, E.; Thiem, D. (Kernforschungszentrum, Karlsruhe (West Germany). Institut fuer Neutronenphysik und Reaktortechnik). Mar 1972. 34p. (KFK-1561). Dep. NTIS (U. S. Sales Only).

Work performed under United States-Euratom Fast Reactor Exchange Program.

The influence of fission neutron spectra on important integral quantities of fast critical assemblies is discussed. The forms of the fission spectra are taken from the KEDAK and ENDF/B library respectively. The different forms are compared with each other and with the "standard" fission spectrum generally used in the calculations which belongs to $\nu = 2.8$ of the Russian ABN-set of group constants. The influence of the different forms on calculated integral quantities for fast critical assemblies and on important characteristics of large fast power reactors is investigated. Some implications for the programs used to calculate flux distributions are outlined. 19 references. (auth)

47586 (HEDL-TME-72-42) THREE DIMENSIONAL NEUTRONICS CALCULATIONS FOR THE FAST TEST REACTOR (FTR) AND THE FTR ENGINEERING MOCK UP CRITICAL ASSEMBLY (EMC). Fleischman, R. M.; Nelson, J. V. (Hanford Engineering Development Lab., Richland, Wash.). Apr 1972. Contract AT(45-1)-2170. 68p. Dep. NTIS.

The three-dimensional diffusion theory code, 3DB, was used to make full-core, detailed, thirty energy group, neutronics calculations of the FTR and EMC. Eigenvalue and reaction rate distribution calculations for the EMC are presented and the latter are compared with experiment. The FTR criticality search using target eigenvalues derived from EMC calculations and calculation of the FTR Doppler coefficient in the established critical configuration are discussed. Reaction rate and power distributions and neutron energy spectra from FTR calculations are presented. Appended to the report are complete cross section and three-dimensional modeling details, and a discussion of 3DB convergence behavior. 13 references. (auth)

47587 EVALUATION OF ABSORBER MATERIALS PERFORMANCE IN HFIR CONTROL CYLINDERS. Knight, R. W.; Richt, A. E. (Oak Ridge National Lab., Tenn.). *Nucl. Technol.*; 15: No. 3, 384-90(Sep 1972).

The irradiation behavior of the High Flux Isotope Reactor (HFIR) control cylinders was assessed. After 48,000 Mwd of irradiation, the nuclear worth of the control cylinder was not changed significantly. The maximum surface oxide buildup was 0.003 to 0.005 in. thick and occurred over the Ta-Al dispersion. The hardness of the 6061 Al cladding was found to have increased from 45 DPH at the low burnup end of the cylinder to 110 DPH over the Ta-Al region which is located approximately 5 in. below the area of maximum fast-neutron fluence. Extensive metallographic examination revealed no degradation or reaction in either the Ta-Al or Eu_2O_3 -Al portions of the cylinder. (auth)

47588 FIRST INDIAN REACTOR. Zamb, N. K. (Bhabha Atomic Research Centre, Bombay). *Vaigyanik*; 2: No. 1, 3-4; 30(2 Apr 1970). (In Hindi).

Apsara reactor that went critical on August 4, 1956, is the first Indian reactor designed and fabricated wholly by Indian scientists and engineers. It is a swimming pool type reactor and uses water as coolant. The reactor is fuelled with enriched uranium and has a maximum power rating of 1,000 kilowatts thermal but is normally operated at 400 kW. The reactor pool, core assembly, coolant system and control system are described. Apsara reactor is used for research experiments in physics, chemistry and biology, and for the production of radioisotopes. (INIS)

47589 MINERVE REACTOR. Bouchard, J.; Carre, J. C. (CEN, Fontenay-aux-Roses, France). *Bull. Inform. Sci. Tech.* (Paris); No. 168, 17-28(Mar 1972). (In French).

Minerve is low power experimental reactor designed, essentially, for neutron studies (in particular using the oscillation method). After a description of the equipment used in conjunction with this reactor, the principal experiments undertaken since 1959 are reviewed: cross section measurements, resonance integral measurements, study of the behavior of Pu in heavy water lattices, and kinetic studies. The central cavity of Minerve has great adaptability; in 1966 a simple modification enabled Minerve to be converted into a fast-thermal coupled assembly, Ermine. The results obtained with this assembly in the development of fast neutron reactor assemblies are exposed. (France)

47590 HARMONIE REACTOR. Lyon, F.; Oeraies, R.; Despret, A. (CEN de Cadarache, Saint-Paul-les-Durance, France). *Bull. Inform. Sci. Tech.* (Paris); No. 168, 45-56(Mar 1972). (In French).

The Harmonie reactor (a low power fast neutron source) is described. It can be used to provide neutrons having a thermal or hard spectrum, or as a plane source. It is well suited to exponential experiments with fast media, e.g., U or Pu. Harmonie was also used for studying shielding problems encountered with fast neutron reactors: basic research and research using mockups. (France)

47591 CALCULATIONS AND BEHAVIOR OF EXPANSION BELLOWS. Roche, R.; Vrillon, B. (CEN de Saclay, Gif-sur-Yvette, France). Bull. Inform. Sci. Tech. (Paris); No. 167, 47-51 (Feb 1972). (In French).

Because of their empirical nature, calculations concerning the behavior of expansion bellows for the channels of nuclear reactors have not always been entirely satisfactory. In 1952 the SEMT became interested in this problem. A simplified formula was developed. Recent developments in calculations using the finite element method have led to considerable progress in the treatment of this problem. The method is particularly useful for calculating the real geometry of the wave profiles. (France)

47592 EXPERIMENTAL REACTOR MASURCA. Boyer, J.; Bruna, J. (CEN de Cadarache, Saint-Paul-les-Durance, France). Bull. Inform. Sci. Tech. (Paris); No. 168, 57-69 (Mar 1972). (In French).

Masurca is a fast neutron experiment installed at Cadarache. It is used to simulate the cores of experimental and power reactors. This reactor is very adaptable and presents a very high operational safety. It has been used since 1969 to study media having compositions similar to those of the two zones of the Phenix reactor. (France)

Reactor Components, Fuels, and Accessories

Refer also to abstracts: 45424, 45933, 45939, 45957, 45959, 45960, 45974, 45996, 46067, and 46077.

47593 (RDT-E-14-6-T(6-72)) GASKETS-CONTAINMENT VESSEL AIRLOCK. (Division of Reactor Development and Technology (AEC), Washington, D. C.). Jun 1972. 10p. RSO.

The requirements for design, manufacture, installation, and test requirements for gasket installations for sealing doors in nuclear power system containment vessel airlocks, are presented. (auth)

Reactor Physics and Core Engineering

Physics

47594 (BMBW-FBK-72-6) COMPUTER-CODE FOR THE SOLUTION OF THE THREE-DIMENSIONAL-MULTIGROUP-EQUATIONS, BASED ON THE IMPROVED SYNTHESIS-METHOD.

Schaeffler, H.; Schnur, P.; Ruehle, R. (Technische Universitaet, Brunswick (West Germany)). May 1972. 81p. (In German). Dep. NTIS (U. S. Sales Only).

A FORTRAN IV-code based on the improved flux synthesis method was developed. This method allows the use of separate sets of trial functions in different axial regions of a reactor. However, the continuity of the neutron flux distribution at the region boundaries is guaranteed. The theory of the improved method is presented. The results of numerical tests described demonstrate the advantage of this technique. Computing time has been considerably reduced in comparison with conventional difference calculations of equal accuracy. Fields of application for the code are discussed and a computer-code-abstract is presented. 13 references. (auth)

47595 DETERMINATION OF RATIOS OF EFFECTIVE CROSS SECTIONS FROM MEASURED BURNUP DATA FOR YANKEE ROWE. Matsen, R. P. (Battelle-Northwest Labs., Richland, Wash.). Nucl. Technol.; 15: No. 3, 343-53 (Sep 1972).

A technique for obtaining cross-section information from experimental isotopic concentration data was applied to the data from Yankee Core I. The technique involves making a least-squares fit of appropriate transmutation equations to the data. The analysis covers two transmutation chains. One is the plutonium chain which begins with ^{238}U and ends with americium and curium. The other is a rarer isotope chain which begins with ^{235}U or ^{238}U and ends with neptunium and plutonium. Values were obtained for twelve ratios determined by the least squares analysis (e.g., $\alpha^{25} = 0.2569 \pm 0.0023$ and $\alpha^{49} = 0.622 \pm 0.082$). In order to obtain information about the less abundant transuranium isotopes, fifteen Yankee Core I samples were analyzed for their isotopic content. The ratios of effective cross sections were also deduced from the data. 9 references. (auth)

Core Thermal Hydraulics

47596 CROSS FLOW RESISTANCE IN ROD BUNDLE CORES. Weisman, Joel (Univ. of Cincinnati). Nucl. Technol.; 15: No. 3, 465-9 (Sep 1972).

Experimental data have shown a high resistance to cross flow in the presence of large axial flows. This phenomenon may be explained by use of a full lateral momentum balance. Application of the momentum balance principle shows that inter-channel cross flow at any given level depends on the cross flow at the lower level in addition to the local pressure gradient. This revised procedure for calculation of cross flow should eventually lead to improved estimation of core wide flow redistribution in open lattice cores. (auth)

GENERAL

Civilian Defense

47597 (EUR-4806) DEFINITIONS OF PROPER NAME DESCRIPTORS USED IN THE EURATOM-INIS NUCLEAR DOCUMENTATION SYSTEM. Vernimb, C. (European Communities, Brussels (Belgium). Commission. European Atomic Energy Community, Luxembourg. Center for Information and Documentation). Mar 1972. 62p. Dep. NTIS (U. S. Sales Only). EUR B. Fr. 85.-.

440 definitions are given of so-called Proper Name Terms, which are used as descriptors in the EURATOM-INIS Nuclear Documentation System. These definitions are intended to give an aid in indexing and retrieval. (auth)

Law

Refer also to abstracts: 45803 and 45804.

47598 (UCID-30047) CATALOG OF LLL COMPUTER DOCUMENTATION REPORTS. Norem, W. L. (California Univ., Livermore, Lawrence Livermore Lab.). 10 May 1972. Contract W-7405-Eng-48. 301p. Dep. NTIS.

A catalog of all unclassified reports issued by the Lawrence Livermore Laboratory and pertaining to computer documentation is given. The reports primarily deal with computers and programs of the Livermore Time-Sharing System. (auth)

Miscellaneous

47599 (ANL-TKK/RJL-2) "FORGY:" A COMPANION COMPUTER PROGRAM OF "TRIM" TO CALCULATE FORCES AND ENERGY IN ELECTROMAGNETS. Khoe, T. K.; Lari, R. J. (Argonne National Lab., Ill.). 4 Jan 1972. 26p. Dep. NTIS.

47600 (BDX-613-253(Rev.)) SYNTACTIC FOAM, FY70-3. Quarterly Report, Johnston, J. R. (Bendix Corp., Kansas City, Mo.). Feb 1972. Contract AT(29-1)-613. 54p. Dep. NTIS.

47601 (BDX-613-288(Rev.)) HIGH-TEMPERATURE THERMAL INSULATION. Quarterly Report, FY 70-3. Fotopoulos, C. U. (Bendix Corp., Kansas City, Mo.). Sep 1971. Contract AT(29-1)-613. 32p. Dep. NTIS.

47602 (BDX-613-392(Rev.)) LIQUID PROCESSING OF MOLDED DESICCANT. Hefner, Bruce A. (Bendix Corp., Kansas City, Mo.). Jan 1972. Contract AT(29-1)-613. 34p. Dep. NTIS.

47603 (BDX-613-440(Rev.)) DEVELOPMENT OF ELASTOMER CUSHIONS, FY69-4. Quarterly Report, Green, R. M. (Bendix Corp., Kansas City, Mo.). Oct 1971. Contract AT(29-1)-613. 32p. Dep. NTIS.

47604 (BDX-613-456) COMPUTER SIMULATION MODELING OF AN AUTOMATED WAREHOUSE SYSTEM. Report on Phase II of a Three-Phase Automated Warehouse Study. Smith, Jerry C. (Bendix Corp., Kansas City, Mo.). Nov 1971. Contract AT(29-1)-613. 79p. Dep. NTIS.

47605 (BDX-613-515(Rev.)) EFFECT OF SODIUM HYDROXIDE ON ALUMINUM FOIL THICKNESS. Bengtson, Glenn W. (Bendix Corp., Kansas City, Mo.). Oct 1971. Contract AT(29-1)-613. 20p. Dep. NTIS.

47606 (BDX-613-534(Rev.)) HIGH-DENSITY BRAID CABLE SHIELD, FY72-1. Quarterly Report, Galbraith, J. F. (Bendix Corp., Kansas City, Mo.). Feb 1972. Contract AT(29-1)-613. 11p. Dep. NTIS.

47607 (BDX-613-579(Rev.)) AUTOCLAVE FABRICATION TECHNIQUES FOR FLAT CABLES. Final Report, Sheeley, J. D. (Bendix Corp., Kansas City, Mo.). Feb 1972. Contract AT(29-1)-613. 24p. Dep. NTIS.

47608 (BDX-613-604(Rev.)) DETERMINATION OF SEMICONDUCTOR JUNCTION PROFILES BY ANGLE LAPPING OF SEMICONDUCTOR WAFERS. Rauch, E. L. (Bendix Corp., Kansas City, Mo.). Feb 1972. Contract AT(29-1)-613. 17p. Dep. NTIS.

47609 (BDX-613-642(Rev.)) THERMAL SHOCK DEBURRING. Final Report. Kuehnle, J. H. (Bendix Corp., Kansas City, Mo.). Jun 1972. Contract AT(29-1)-613. 45p. Dep. NTIS.

47610 (COO-2118-34) PART 1: ANNUAL TECHNICAL PROGRESS REPORT, OCTOBER 1, 1971-JUNE 30, 1972. McCor-

mick, Bruce H. (Illinois Univ., Urbana. Dept. of Computer Science). 30 Jun 1972. Contract AT(11-1)-2118. 94p. Dep. NTIS.

47611 (COO-3078-5) EARLY STEPS IN PROTEIN SYNTHESIS AND THEIR REGULATION: A BACKGROUND STUDY RELATED TO THE BIOLOGICAL EFFECTS OF RADIATION. Progress Report, September 1, 1971-August 31, 1972. (Massachusetts General Hospital, Boston. John Collins Warren Labs.). 1972. Contract AT(11-1)-3078. 26p. Dep. NTIS.

47612 (COO-3078-6) AN INVESTIGATION ON THE MECHANISM OF PROTEIN AND NUCLEIC ACID SYNTHESIS: A BACKGROUND STUDY RELATED TO THE BIOLOGICAL EFFECTS OF RADIATION. Dodecennial Progress Report, September 1, 1960-August 31, 1972. (Massachusetts General Hospital, Boston. John Collins Warren Labs.). 1972. Contract AT(11-1)-3078. 11p. Dep. NTIS.

47613 (COO-3292-1) STUDY ON THE CHEMICAL BEHAVIOR OF ZINC IN CHESAPEAKE BAY WATER USING ANODIC STRIPPING VOLTAMMETRY. Technical Report 76. Bradford, Wesley Lamont (Johns Hopkins Univ., Baltimore, Md. Chesapeake Bay Inst.). Jun 1972. Contract AT(11-1)-3292. 113p. Dep. NTIS.

47614 (COO-3498-2) FUNDAMENTAL STUDIES IN ISOTOPE CHEMISTRY. Progress Report, July 1, 1971-June 30, 1972. Bigeleisen, Jacob (Rochester Univ., N. Y.). 1972. Contract AT(11-1)-3498. 7p. Dep. NTIS.

47615 (CU-3194-22) ANELASTIC RELAXATION CAUSED BY POINT DEFECTS IN MnF₂. Srinivasan, Nemelli Krishnaswami (Columbia Univ., New York. Henry Krumb School of Mines). Apr 1972. Contract AT(11-1)-3194. 92p. Dep. NTIS.

47616 (EUR-4832) POSTGRADUATE TRAINING OF SCIENTISTS IN THE EUROPEAN COMMUNITY. Rouche, N. (Fondation Industrie-Universite, Brussels (Belgium)). 1972. 172p. (In French). Dep. NTIS (U. S. Sales Only). EUR FB 225.-.

47617 (IEA-19) SELECTED TOPICS ON DYNAMICS OF CRYSTAL LATTICES. Borgonovi, G. M. (Instituto de Energia Atomica, Sao Paulo (Brazil)). Nov 1971. 50p. Dep. NTIS (U. S. Sales Only).

47618 (JINR-P14-6407) NEW METHOD OF DETERMINATION OF PHYSICAL-CHEMICAL PROPERTIES OF MATTER. Zinov, V. G.; Konin, A. D.; Mukhin, A. I. (Joint Inst. for Nuclear Research, Dubna (USSR). Lab. of Nuclear Problems). 1972. 5p. (In Russian). Dep. NTIS (U. S. Sales Only).

47619 (KY-642) CONDITIONS FOR THE REMOVAL OF CHROMATE FROM COOLING WATER. Lemonds, Alice J. (Paducah Gaseous Diffusion Plant, Ky.). 10 Jul 1972. Contract W-7405-eng-26. 10p. Dep. NTIS.

47620 (MIT-144) MATHEMATICAL MODEL FOR THE PREDICTION OF UNSTEADY SALINITY INTRUSION IN ESTUARIES. Thatcher, M. Llewellyn; Harleman, Donald R. F. (Massachusetts Inst. of Tech., Cambridge (USA). Ralph M. Parsons Lab. for Water Resources and Hydrodynamics). Feb 1972. 232p. (R-72-10; DSR-72602; DSR-73479).

47621 (ORNL-TM-3810) EXECUTION TIME TRANSLATE SUBROUTINES. Joseph, Allen F. (Oak Ridge National Lab., Tenn.). Jul 1972. Contract W-7405-eng-26. 38p. Dep. NTIS.

47622 (ORO-4245-4) STUDIES ON NUTRITION OF THE PREPLACENTAL EMBRYOS OF THE MAMMAL. Sixth Year Progress Report. (Tennessee Univ., Knoxville). 1972. Contract AT(40-1)-4245. 8p. Dep. NTIS.

47623 (RLO-2229-T-1-18) TRACE-METAL ASSOCIATIONS IN SUB-ARCTIC AND ARCTIC MARINE ENVIRONMENTS. Progress Report, June 1971-May 1972. Burrell, David C. (Alaska Univ., College. Inst. of Marine Science). May 1972. Contract AT(45-1)-2229. 136p. (R-72-8). Dep. NTIS.

Continuing research on trace-metal associations in the marine environment of the arctic and subarctic is described. Geological studies were made of the fiords of S.E. Alaska and coordinated with geochemical studies related to sedimentation dynamics, suspended sediment size fractionation, chemical composition of bottom sediments, distribution of Zn within the environment, and sedimental interstitial water chemistry. Information is included on gas-liquid chromatographic analysis of trace metals in seawater, solvent extraction techniques, and the use of atomic ab-

sorption spectrometry for measuring trace inorganic pollutants in the marine environment. (L.C.L.)

47624 (SC-CR-723142) HIGH-PEAK-POWER, HIGH-EFFICIENCY MICROCIRCUIT AVALANCHE DIODE OSCILLATORS. Liu, S. G. (RCA Labs., Princeton, N. J.). Jul 1972. Contract 11-2475. 84p. Dep. NTIS.

For Sandia Labs., Albuquerque, N. Mex.

47625 (SC-DC-722267) ANODE TEMPERATURES AND ION ENERGY DISTRIBUTIONS DURING A VACUUM ARC DISCHARGE. Grissom, J. T.; Newton, J. C. (Sandia Labs., Albuquerque, N. Mex.). 1971. 6p. (CONF-720807-1). Dep. NTIS.

From fifth international symposium on discharges and electrical insulation in vacuum; Poznan, Poland (31 Aug 1972).

47626 (SC-T-722520) LONGITUDINAL IMPACT ON THE END FACE OF PRISMATIC RODS, CYLINDRICAL, AND CONICAL SHELLS. Egorshin, A. A. Translated by Patricia Newman (Sandia Labs., Albuquerque, N. Mex.), from Samoletost. Tekh. Vozdush. Flota; 13: 99-104(1968). 10p. Dep. NTIS.

47627 (SCL-RR-720270) AN INTRODUCTION TO COBORDISM THEORY. Rinne, R. L. (Sandia Labs., Livermore, Calif.). May 1972. Contract AT(29-1)-789. 23p. Dep. NTIS.

47628 (TID-3767) BIBLIOGRAPHIES OF ATOMIC ENERGY LITERATURE. Issued or in Progress During the Period April-June 1972. (Technical Information Center (AEC), Oak Ridge, Tenn.). 1972. 15p. TIC Free.

47629 (UCLA-12-866) SINGLE CELL TRANSPLANTS: GROSS VIRUS-INDUCED MOUSE LYMPHOMAS. Vredevoe, Donna L.; Hays, Esther F. (California Univ., Los Angeles. Lab. of Nuclear Medicine and Radiation Biology). 1971. Contract AT(04-1)-Gen-12. 15p. Dep. NTIS.

47630 (UCLA-34-P-177-11) CONTRAST EFFECTS AT GRAIN BOUNDARY LEDGES IN 304 AUSTENITIC STAINLESS STEEL. McDonald, R. C.; Ardell, A. J. (California Univ., Los Angeles. School of Engineering and Applied Science). Apr 1972. Contract AT(04-3)-34. 4p. Dep. NTIS.

47631 (UCRL-51210) SHAKER: A COMPUTER PROGRAM FOR NUMERICAL ANALYSIS OF WEIGHTED TABULAR DATA. Edwards, Arthur L. (California Univ., Livermore. Lawrence Livermore Lab.). Jan 1972. Contract W-7405-eng-48. 262p. Dep. NTIS.

The SHAKER program was developed to meet a need for a single program capable of carrying out a wide variety of numerical processing procedures on weighted tabular data, with a minimum effort on the part of the user, and with maximum flexibility in the choice and sequence of the processing steps. Available procedures include least-squares fitting of weighted tabular data with a variety of functions, regression analysis, integration and differentiation, purging, smoothing, discontinuity measurement, linear mapping of coordinates, functional transformation, reordering, finding the ranges of table variables, weight factor adjustment, reduction of multi-valued tables to single-valued tables, merging of two or more tables, reduction of two parametric tables to a single table, alignment of two tables by translation and/or scaling, and many others. Procedures are provided for input and generation of tables, labeling of tables and table variables, manipulation and replication of tables in memory, and output of tables, including printed, punched, and graphical output. During a computer run, each procedure is specified and controlled by a card in the input file. The control cards may be arranged in any desired order to obtain the desired sequences of processing. FORTRAN listings of all of the routines in the SHAKER program are given. (auth)

47632 (UCRL-51227) DETERMINANT METHOD FOR CALCULATING SMALL-CRYSTAL ENTROPY. Hindmarsh, A. C.; Hoover, W. G. (California Univ., Livermore. Lawrence Livermore Lab.). 18 May 1972. Contract W-7405-eng-48. 40p. Dep. NTIS.

47633 (UCRL-51239) AN EFFICIENT HIGH-VOLUME MINIATURE AIR MOVER. Taylor, R. D. (California Univ., Livermore. Lawrence Livermore Lab.). 30 May 1972. Contract W-7405-eng-48. 9p. Dep. NTIS.

47634 (UCRL-73587) SOLID STATE RMS RECORDING AMMETER. Parker, Robert (California Univ., Livermore. Lawrence Livermore Lab.). 16 Dec 1971. 19p. (CONF-720706-1). Dep. NTIS.

From Power Engineering Society Summer meeting; San Francisco, CA. (9 Jul 1972).

47635 (UCRL-73805) SIMPLE METHOD FOR SEPARATING SPINDLE ERROR FROM TEST BALL ROUNDNESS ERROR.

Donaldson, R. R. (California Univ., Livermore. Lawrence Livermore Lab.). 13 Mar 1972. 11p. (CONF-720809-1). Dep. NTIS.

From twenty-third meeting of the International Society of Electrochemistry; Stockholm, Sweden (27 Aug 1972).

47636 (UCRL-73925) SURFACE SCIENCE AND SURFACE DAMAGE. Khan, Jhan M. (California Univ., Livermore. Lawrence Livermore Lab.). 26 Jun 1972. 19p. (CONF-720625-1). Dep. NTIS.

From fourth symposium on damage in laser materials; Boulder, CO. (14 Jun 1972).

47637 (UR-49-1508) BIOLOGICAL PROPERTIES OF CERTAIN COMPOUNDS CONTAINING THE C-F BOND. Smith, Frank A. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1971. 36p. (CONF-710913-3). Dep. NTIS.

From 162nd national meeting of the American Chemical Society; Washington, D. C. (12 Sep 1971).

47638 (UR-3055-1) PASSIVE CATION FLUX CHANGES IN THE HUMAN ERYTHROCYTE SUSPENDED IN VARIOUS ANIONIC MEDIA. Callahan, Thomas Joseph (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 157p. Dep. NTIS.

Thesis.

47639 (UR-3055-2) BACTERIOPHAGE T4 INHIBITS COLICIN E2-INDUCED DEGRADATION OF ESCHERICHIA COLI DE-OXYRIBONUCLEIC ACID. Swift, Robert L. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 98p. Dep. NTIS.

Thesis.

47640 (UR-3490-18) PROTEIN SYNTHESIS AND THE CELL CYCLE OF L5178Y CELLS. Roti Roti, Joseph Lee (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1971. 105p. Dep. NTIS.

Thesis.

47641 (UR-3490-21) PROCEDURE FOR CHRONIC INTRA-VENOUS CATHETERIZATION IN THE RABBIT. Hall, Larry L.; De Lopez, Ofelia H.; Roberts, Anne; Smith, Frank A. (Rochester Univ., N. Y. School of Medicine and Dentistry). 1971. Contract AT(11-1)-3490. 13p. Dep. NTIS.

47642 (UR-3490-47) SPECTRAL STUDIES OF COLLAGEN. Shadbolt, Margaret Anne (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1971. 144p. Dep. NTIS.

Thesis.

47643 (UR-3490-80) EFFECT OF A POLYTHIOL SYNTHETIC RESIN ON THE ABSORPTION AND EXCRETION OF METHYLMERCURY IN EXPERIMENTAL ANIMALS. Clarkson, T. W.; Small, H.; Norseth, Tor (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 13p. Dep. NTIS.

47644 (UR-3490-85) CHRONIC LOW-LEVEL NO₂ EXPOSURE AND THE MUCOCILIARY CLEARANCE MECHANISM IN THE RAT TRACHEA IN VIVO. Giordano, Anthony Michael Jr. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 38p. Dep. NTIS.

Thesis.

47645 (UR-3490-107) FLUORINE METABOLITE CONCENTRATIONS AND RENAL CLEARANCES FOLLOWING METHOXY-FLURANE ANESTHESIA. Fry, Bill W.; Taves, Donald R.; Merin, Robert G. (Rochester Univ., N. Y. School of Medicine and Dentistry). 1972. 27p. Dep. NTIS.

47646 (UR-3490-131) EFFECT OF VARYING DOSES OF CORTISOL ON THE INCORPORATION OF ³H-THYMIDINE INTO RAT LIVER DNA AND ON THE SYNTHESIS OF FIVE PLASMA PROTEINS IN THE ISOLATED PERFUSED REGENERATING RAT LIVER. Lebda, Nancy Jean Abbott (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 59p. Dep. NTIS.

Thesis.

47647 (UR-3490-132) EFFECTS OF CHLORAMPHENICOL AND CYCLOHEXIMIDE ON THE CELL GROWTH CYCLE OF MOUSE LEUKEMIC CELLS (L5178Y). Liberman, Daniel Franklin (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 88p. Dep. NTIS.

Thesis.

47648 (UR-3490-137) PATHOPHYSIOLOGIC EFFECTS OF MICROWAVE RADIATION WITH SPECIAL REFERENCE TO THE THYROID-PITUITARY AXIS. Milroy, William C. (Rochester

Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 114p. Dep. NTIS.

Thesis.

47649 (UR-3490-147) DISTRIBUTION OF AN ^{125}I -LABELED TRANSPLANTATION ANTIBODY IN SYNGENEIC AND ALLOGENEIC RATS. Izzo, Mary Jane; Buchsbaum, Donald J.; Bale, William F. (Rochester Univ., N. Y. Dept. of Radiation Biology and Biophysics). 1972. 14p. Dep. NTIS.

47650 (UR-3490-153) EFFECTS OF ACTINOMYCIN D ON G_1 PROTEIN SYNTHESIS IN L5178Y CELLS. Roti Roti, J. L.; Okada, S.; Eberle, H. (Rochester Univ., N. Y. School of Medi-

cine and Dentistry. Tokyo Univ. (Japan). Dept. of Radiation Biophysics). 1972. 12p. Dep. NTIS.

47651 (UR-3490-154) PROTEIN SYNTHESIS DURING THE CELL CYCLE OF L5178Y CELLS. Roti Roti, J. L.; Okada, S.; Eberle, H. (Rochester Univ., N. Y. School of Medicine and Dentistry. Tokyo Univ. (Japan). Dept. of Radiation Biophysics). 1972. Contract AT(30-1)-1286. 26p. Dep. NTIS.

47652 CANADIAN NATIONAL SCIENTIFIC AND TECHNICAL INFORMATION (STI) SYSTEM: A PROGRESS REPORT. Report No. 4, May 1972. Brown, Jack E. Philadelphia; National Federation of Science Abstracting and Indexing Services (1972). 28p. (NP-19360).

CORPORATE AUTHOR INDEX

In the Corporate Author Index, technical report literature is indexed to the organization or institution responsible for issuing the report. Headings are given for all report literature and for published literature for which a corporate author approach is especially desirable, e.g., symposium and conference proceedings.

Each entry under a corporate author heading gives the publication title and the NSA volume and abstract numbers. Report numbers, where applicable, are given in parentheses at the end of the entry.

Corporate Author Headings Used by the U. S. Atomic Energy Commission in Cataloging Reports, TID-5059 (10th Rev.), is available from the National Technical Information Service (NTIS), U. S. Department of Commerce, Springfield, Virginia 22151, for \$3.00 for full size or reproduction copy and \$0.95 for microfiche copy.

A

AEROJET NUCLEAR CO., IDAHO FALLS, IDAHO.
Total neutron cross section of ^{235}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)

AKADEMIYA NAUK SSSR, MOSCOW. INSTITUT FIZIKI.
Photoproduction of neutral pions on helium nuclei, 26:46432 (SLAC-Trans-142)

AKADEMIYA NAUK UKRAINSKOI SSR, KIEV. INSTITUT TEORETICHESKOI FIZIKI.
Choice of a form factor in nonlocal quantum electrodynamics, 26:46698 (ITF-71-128-P)
Impossibility of introducing "hidden" parameters into quantum mechanics, 26:47285 (ITF-71-101-P)
On the distribution of resonance spins in the dual resonance model, 26:46609 (ITF-72-13-P)
Relativistic transformations of the one-particle wave functions, 26:46667 (ITF-71-117-P)
Threshold behavior of DAMA, 26:46774 (ITF-72-44-E)

AKTIEBOLAGET ATOMENERGI, STUDDSVIK (SWEDEN).
Depth distribution studies of carbon in steel surfaces by means of charged particle activation analysis with an account of heat and diffusion effects in the sample, 26:45466 (AE-451)
Fast neutron elastic scattering experiments, 26:46916 (AE-452)
Low-grade heat from thermal electricity production, 26:45573 (AE-448)
Nuclear chemistry. Progress report, 1971, 26:45309 (AE-453)

ALASKA UNIV., COLLEGE. INST. OF MARINE SCIENCE.
Trace-metal association in sub-Arctic and Arctic marine environments. Progress report, June 1971--May 1972, 26:47623 (RLO-2229-I-1-18)

ALLIED CHEMICAL CORP., IDAHO FALLS, IDAHO.
DTA--x-ray diffraction examination of chemical reprocessing wastes containing sodium and potassium nitrates, 26:45383 (ICP-1008)

ALLIED CHEMICAL CORP., IDAHO FALLS, IDAHO. IDAHO CHEMICAL PROGRAMS OPERATIONS OFFICE.
Evaluation of instrumentation for nuclear fuels reprocessing plant input weight measurements, 26:46093 (ICP-1014)
Idaho Chemical Programs annual technical report, fiscal year 1971, 26:45405 (ICP-1006)

ALLIED-GULF NUCLEAR SERVICES, BARNWELL, S.C.
Uranium Hexafluoride Facility. Environmental report, 26:45308 (DOCKET-701327-1)

AMES LAB., IOWA.
Testing triplet models, 26:46516 (DESY-72/21)

ARGONNE NATIONAL LAB., ILL.
200-GeV proton--proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)
"FORGY": a companion computer program of "TRIM" to calculate forces and energy in electromagnets, 26:47599 (ANL-TKK/RJL-2)
Observation of the $M(940)$ and $\delta(963)$, 26:46475 (IS-2886)
Theoretical study of the swelling of fast-neutron-irradiated materials, 26:46063 (CONF-710601-, pp 798-824)

ARKANSAS POWER AND LIGHT CO., LITTLE ROCK.
Arkansas Nuclear One, Unit 1. Environmental report, Supplement 2, 26:47486 (DOCKET-50313-44)
Arkansas Nuclear One, Unit 1. Environmental report, Supplement 3, 26:47487 (DOCKET-50313-45)
Arkansas Nuclear One, Unit 1. Final safety analysis report, Amendment 27, 26:47488 (DOCKET-50313-46)

ATLANTIC RICHFIELD HANFORD CO., RICHLAND, WASH.
Phosphate solidification of caustic process solution waste, 26:45381 (ARH-SA-120)
Plutonium--uranium partitioning by a reflux solvent extraction flowsheet, 26:45382 (ARH-SA-130)
Relative efficiency as a function of time for Atlantic Richfield Hanford Company coaxial Ge(Li) detectors, 26:45586 (ARH-2432)
Remote welding program for Atlantic Richfield Hanford Company's waste encapsulation and storage facility, 26:46099 (ARH-SA-117)
Spectrophotometric determination of phosphate in nuclear materials, 26:45219 (ARH-SA-110)

ATOMIC ENERGY COMMISSION, WASHINGTON, D. C.
Leasing of AEC controlled uranium bearing lands, Colorado, Utah, New Mexico. Environmental statement, 26:45572 (WASH-1523)
Radiation-induced voids in metals. Proceedings of the 1971 international conference held at Albany, New York, June 9--11, 1971, 26:46045 (CONF-710601)
Reactor technology. Volume 15, Number 2, 26:47301

ATOMIC ENERGY RESEARCH ESTABLISHMENT, HARWELL (ENGLAND).
Monitor for estimating tritium in urine, 26:45585 (AERE-HP/M-74)
Observation and theory of the void lattice in molybdenum, 26:46055 (CONF-710601-, pp 522-32)
Theory of void formation and growth in irradiated materials, 26:46062 (CONF-710601-, pp 769-97)
Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)

ATOMIC WEAPONS RESEARCH ESTABLISHMENT, ALDERMASTON (ENGLAND).
Some measurements of the electrical conductivity induced in polymers by pulsed x-ray irradiation, 26:47225 (AWRE-O-31/72)

ATOMICS INTERNATIONAL, CANOGA PARK, CALIF.
Studies of void formation in proton-irradiated type 316 and titanium-modified 316 stainless steels, 26:46044 (AI-AEC-13033)
Swelling as a function of displacement damage in proton-irradiated 316 stainless steel, 26:46054 (CONF-710601-, pp 511-21)

AUBURN UNIV., ALA. DEPT. OF CHEMICAL ENGINEERING.
Synthesis, identification, and purification of C₄ and C₇ fluorocarbons from radiolysis. Final report, 26:45316 (ORD-3909-3)

AUSTRALIAN ATOMIC ENERGY COMMISSION RESEARCH ESTABLISHMENT, LUCAS HEIGHTS.
Environmental hazards of fossil and nuclear power production, 26:45510 (AAEC/TM-627)
Management of wastes from irradiated nuclear fuels, 26:46098 (AAEC/TM-628)

B

BATTELLE COLUMBUS LABS., OHIO.
Stress-corrosion cracking of sensitized stainless steel in oxygenated high-temperature water, 26:45966 (BMI-1927)

BATTELLE-NORTHWEST, RICHLAND, WASH. PACIFIC NORTHWEST LAB.
 ^{252}Cf neutron radiography at Battelle-Northwest, 26:45467 (BNWL-SA-4320)

BATTELLE PACIFIC NORTHWEST LABS., RICHLAND, WASH.
High-temperature electrical conductivity of CeD/sub 2-x/, 26:47247 (BNWL-1511)
Oceanic distributions and relationships of ^7Be and fission products, 26:45530 (BNWL-SA-4205(Rev.))
Photon intensities and their importance in counting transuranic materials, 26:47011 (BNWL-1648)
Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)

BELLCOMM, INC., WASHINGTON, D. C.

Interaction of the solar wind with a planetary ionosphere, 26:46324 (N-72-18796)

BENDIX CORP., KANSAS CITY, MO.

Autoclave fabrication techniques for flat cables, 26:47607 (BDX-613-579(Rev.))

Computer simulation modeling of an automated warehouse system, 26:47604 (BDX-613-456)

Determination of semiconductor junction profiles by angle lapping of semiconductor wafers, 26:47608 (BDX-613-604(Rev.))

Development of elastomer cushions, FY69-4. Quarterly report, 26:47603 (BDX-613-440(Rev.))

Effect of sodium hydroxide on aluminum foil thickness, 26:47605 (BDX-613-515(Rev.))

High-density braid cable shield, FY72-1. Quarterly report, 26:47606 (BDX-613-543(Rev.))

High-temperature thermal insulation. Quarterly report, FY 70-3, 26:47601 (BDX-613-288(Rev.))

Liquid processing of molded desiccant, 26:47602 (BDX-613-392(Rev.))

Synthetic foam, FY70-3, 26:47600 (BDX-613-253(Rev.))

Thermal shock deburring, 26:47609 (BDX-613-642(Rev.))

BHABHA ATOMIC RESEARCH CENTRE, BOMBAY (INDIA).

Absorption of cathodic hydrogen by iron and steel, 26:45986 (BARC-585)

Analog states in ^{32}Cl by isospin-forbidden resonances in reaction $^{32}\text{S}(p,\gamma)^{32}\text{Cl}$, 26:46902 (BARC-598)

Chemical diffusion in the beta phase of zirconium--titanium system, 26:45985 (BARC-577)

Excitation energy dependence of shell effects in nuclear fission, 26:47043 (INDC-7-U, pp 3-19)

Prog. rpt. on chemistry, annual, 1970/71, 26:45311 (BARC-599)

Prompt radiations emitted in low energy fission, 26:47019 (INDC-7-U, pp 89-112)

Simple transistorised control unit for neutron crystal spectrometers, 26:45636 (BARC-591)

Spectrochemical analysis of high purity cadmium, 26:45220 (BARC-593)

Theory of fragment mass and charge distribution in fission, 26:47044 (INDC-7-U, pp 65-79)

Water chemistry studies VII: general review of water conditions and water treatment in nuclear reactors, 26:45310 (BARC-595)

BLUME (JOHN A.) AND ASSOCIATES RESEARCH DIV., SAN FRANCISCO, CALIF.

Observations of chimney damage to residential buildings in Glendale, California: San Fernando earthquake, February 1971, 26:45474 (JAB-99-88)

BONN UNIV. (WEST GERMANY). PHYSIKALISCHES INSTITUT.

Separation of sigma/sub S/ and sigma/sub T/ in the region of the Λ (1236) resonance and determination of the magnetic dipole transition form factor, 26:46431 (NP-19256)

BORDEAUX UNIV. (FRANCE).

De-excitation of compound nuclei formed in $^{181}\text{Ta} + ^{11}\text{B}$ reactions, 26:46974 (FRNC-TH-207)

BOSTON EDISON CO., MASS.

Pilgrim Nuclear Power Station. Abnormal occurrence AO-72-3, 26:47376 (DOCKET-50293-80)

Pilgrim Nuclear Power Station. Abnormal occurrence AO-72-4, 26:47377 (DOCKET-50293-82)

Pilgrim Nuclear Power Station. Abnormal occurrence AO-72-5, 26:47378 (DOCKET-50293-83)

BRANDEIS UNIV., WALTHAM, MASS.

Molecular basis for the mutagenic and lethal effects of ultraviolet irradiation. Progress report, June 16, 1971--May 31, 1972, 26:45722 (COO-3232-1)

BROOKHAVEN NATIONAL LAB., UPTON, N. Y.

Continuous sampling plan for the taking of physical inventories, 26:46092 (BNL-16930)

Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)

Evidence for existence of a new meson in the 955 MeV mass region, 26:46471 (BNL-16954)

Fourier transform of inelastic neutron scattered intensity for solid ^4He , 26:46845 (BNL-16935)

Operating characteristics of neutron well coincidence counters, 26:45587 (BNL-50332)

Photoproduction experiments at ISABELLE, 26:46426 (BNL-16968)

Scaling in $p\pi^- \rightarrow p\pi^-$ and an estimate of the triple-Pomeranchuk coupling, 26:46616 (BNL-16928)

BUNDESFORSCHUNGSANSTALT FUER LEBENSMITTELFRISSHALTUNG, KARLSRUHE (WEST GERMANY).

Development of methods for application of neutron activation analysis to determination of trace elements in foods. Progress report, 10--12/71, 26:45189 (BFL-1/1972)

Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)

C

CAIRO UNIV. (EGYPT).

Fast neutron elastic scattering experiments, 26:46916 (AE-452)

CALCUTTA UNIV. (INDIA).

Fission barrier heights, energy release, and fragment energy partition processes from the viewpoints of nuclear structural models, 26:47046 (INDC-7-U, pp 145-60)

CALIFORNIA INST. OF TECH., PASADENA.

Perturbation treatment of high energy electron diffraction from imperfect crystals, 26:46510 (CALT-767-P-3-22)

CALIFORNIA UNIV., BERKELEY. LAWRENCE BERKELEY LAB.

Annealing of loops and tetrahedra, 26:45989 (CONF-710601-, pp 647-62)

Final state hadron distributions and generalized scaling laws in deep-inelastic electroproduction processes, 26:46521 (LBL-773)

High-temperature spectroscopy and thermodynamics of titanium oxide and vanadium oxide, 26:45932 (LBL-450)

Range-momentum correlation in carbon and iron for muons of 500 to 1500 MeV/c, 26:46448 (LBL-791)

Some properties of protactinium metal and its compounds, 26:45992 (LBL-659)

Study of dipion production in the reaction $\pi^+p \rightarrow \pi^0\pi^+n$ between 1.6 and 2.4 GeV/c, 26:46477 (LBL-378)

CALIFORNIA UNIV., LIVERMORE. LAWRENCE LIVERMORE LAB.

Accumulation and loss of cobalt and cesium by the marine clam, Mya Arenaria, under laboratory and field conditions, 26:45545 (UCRL-73744)

An efficient high-volume miniature air mover, 26:47633 (UCRL-51239)

Catalog of LLL computer documentation reports, 26:47598 (UCID-30047)

Computerized quantitative analysis by gamma-ray spectrometry. Volume I. Description of the GAMANAL program, 26:45265 (UCRL-51061(Vol.1))

Contamination control hardware for remote hot cell transfers, 26:46108 (UCRL-73757)

Corrosion and scaling by steam in nuclear geothermal power plants, 26:45487 (UCRL-73939)

Determinant method for calculating small-crystal entropy, 26:47632 (UCRL-51227)

Environmental levels of radioactivity in the vicinity of the Lawrence Livermore Laboratory January through December 1971, 26:45513 (UCRL-51242)

Hazards control. Progress report No. 41, September--December 1971, 26:45673 (UCRL-50007-71-3)

Method for estimating heights of water waves generated by underwater cratering explosions, 26:45486 (UCRL-51235)

Pseudo cloud-chamber tracks and their uses, 26:45647 (UCRL-51214)

Review of problems associated with acceleration measurements, 26:45674 (UCRL-73665)

SHAKER: a computer program for numerical analysis of weighted tabular data, 26:47631 (UCRL-51210)

Shielded caves for dual-mode operation, 26:46107 (UCRL-73756)

Simple method for separating spindle error from test ball roundness error, 26:47635 (UCRL-73805)

Solid state RMS recording ammeter, 26:47634 (UCRL-73587)

Surface science and surface damage, 26:47636 (UCRL-73925)

Theoretical radiative transition rates for K- and L-shell x rays, 26:46341 (UCRL-51231)

X-ray-induced photoconductivity in dielectric films, 26:45668 (UCRL-73974)

CALIFORNIA UNIV., LOS ANGELES. LAB. OF NUCLEAR MEDICINE AND RADIATION BIOLOGY.

Irradiation and viral lymphomagenesis in thymic grafts, 26:45818 (UCLA-12-867)

Radioecology and ecophysiology of desert plants at the Nevada Test Site, 26:45755 (IID-25954)

Single cell transplants: Gross virus-induced mouse lymphomas, 26:47629 (UCLA-12-866)

CALIFORNIA UNIV., LOS ANGELES. SCHOOL OF ENGINEERING AND APPLIED SCIENCE.

Contrast effects at grain boundary ledges in 304 austenitic stainless steel, 26:47630 (UCLA-34-P-177-11)

Reactor steels studies. Quarterly progress report, October--December 1971, 26:45994 (UCLA-34-P-177-10)

CALIFORNIA UNIV., RIVERSIDE.

Radioecology and ecophysiology of desert plants at the Nevada Test Site, 26:45755 (IID-25954)

CAMBRIDGE UNIV. (ENGLAND). DEPT. OF APPLIED MATHEMATICS AND THEORETICAL PHYSICS.

Deep inelastic electroproduction and related processes, 26:46515 (DAMTP-71/24)

CAROLINA POWER AND LIGHT CO., RALEIGH, N.C.

Brunswick Steam Electric Plant, Units 1 and 2. Environmental report, Amendment 2, 26:47497 (DOCKET-50324-40)

Brunswick Steam Electric Plant, Units 1 and 2. Expected doses which might result from purging containment, 26:47496 (DOCKET-50324-39)

H. B. Robinson Unit 2. Boron injection tank dilution, 26:47366 (DOCKET-50261-102)

H. B. Robinson, Unit 2. Malfunction of 'B' steam line isolation, 26:47365 (DOCKET-50261-101)

Shearon Harris Nuclear Power Plant, Units 1, 2, 3, and 4. Preliminary safety analysis report, Amendment 10, 26:47563 (DOCKET-50400-19)

Shearon Harris Nuclear Power Plant, Units 1, 2, 3, and 4. Preliminary safety analysis report, Amendment 11. Environmental report, Volume 2, 26:47564 (DOCKET-

Shearon Harris Nuclear Power Plant, Units 1, 2, 3, and 4.
Preliminary safety analysis report, Amendment 12,
26:47565 (DOCKET-50400-21)

CENTRAL ELECTRICITY GENERATING BOARD, BERKELEY (ENGLAND).
BERKELEY NUCLEAR LABS.

Multi-compartment model for studying the dose-intake
relationship for radioactive isotopes, 26:45905
(RD/B/N-2291)

CENTRAL ELECTRICITY GENERATING BOARD, BRISTOL (ENGLAND).
BERKELEY NUCLEAR LABS.

Application of high-voltage electron microscopy to void
studies, 26:46056 (CONF-710601-, pp 533-49)

CENTRO INFORMAZIONI STUDI ESPERIENZE, MILAN (ITALY).

Actual value method evaluation of the effects of fuel
costs on the cost of power produced by a nuclear
reactor for the case of batch refueling and for the
case of continuous refueling, 26:47315 (CISE-R-178)

Analog computer models for safety evaluation of CART fuel
test facility, 26:47580 (CISE-R-259)

PDE: digital program for calculation of thermal-
hydrodynamic condition in circuit of two phase mixture
water and steam, 26:45458 (CISE-R-245)

Programs for thermohydraulic behavior of primary circuit
of reactor CIRENE, 26:47309 (CISE-R-257)

Reactor chemistry: analytical work, 26:45181 (CISE-R-
172)

Threshold of parallel channel flow instability, 26:45459
(CISE-R-275)

ČESKOSLOVENSKÁ AKADEMIE VED, PRAGUE. FYZIKÁLNÍ ÚSTAV.

Before applying the dispersion theory to the
photoproduction reactions, 26:46511 (CONF-710816-9)

ČESKOSLOVENSKÁ AKADEMIE VED, REZ. ÚSTAV JADERNEHO VÝZKUMU.

Deformation of fuel element rods during irradiation,
26:46070 (UJV-2798-M)

Mechanical characteristics of uranium alloys of type U--
Mo(Fe, Al, Si) and U--Fe--Al--(Si, Cr), 26:45995 (UJV-
2802-M)

Role of statistical fluctuations in texture during the
radiation growth in uranium, 26:46069 (UJV-2795-M)

Strontium sorption on sorbents of BaSO₄--CaSO₄ system,
26:45273 (UJV-2820-Ch)

CHALMERS TEKNISKA HOGSKOLA, GÖTEBORG (SWEDEN).

INSTITUTIONEN FÖR TEORETISK FYSIK.

Breakdown of simple cancellation in the operator multiple
scattering series, 26:46773 (CTH-ITP-71-17)

Internal quantum numbers and homogeneous spaces,
26:46665 (CTH-ITP-71-14)

Nucleon form factors in field theory on homogeneous space
of Poincaré group, 26:46514 (CTH-ITP-71-30)

CLARKSON COLL. OF TECH., POTSDAM, N.Y. DEPT. OF CHEMICAL
ENGINEERING.

Nucleation of voids in materials during high-energy
irradiation, 26:46064 (CONF-710601-, pp 825-40)

COLORADO UNIV., BOULDER (USA). DUANE PHYSICAL LAB.

Efficiency studies of Pilot Y, Pilot F, and NE 110
scintillator, 26:45588 (COO-2114-2)

COLUMBIA UNIV., NEW YORK.

Fission isomeric state in ²³⁶U, 26:47018 (INDC-7-U, pp
61-4)

Mass distributions in fission and some related phenomena,
26:47017 (INDC-7-U, pp 31-59)

Research in computational plasma physics. Progress
report, March 1, 1971--March 1, 1972, 26:47145 (COO-
3086-15)

COLUMBIA UNIV., NEW YORK. HENRY KRUMB SCHOOL OF MINES.

Anelastic relaxation caused by point defects in MnF₂,
26:47615 (CU-3194-22)

COMISION NACIONAL DE ENERGIA ATOMICA, BUENOS AIRES

(ARGENTINA).

Importance of the quadrupole pairing field in the J/sup
w/ = 0+ vibrations of shape deformed nuclei, 26:47042
(COO-3001-12)

COMISION NACIONAL DE ENERGIA ATOMICA, BUENOS AIRES

(ARGENTINA). DIVISION ESTUDIOS AMBIENTALES.

Monitoring ⁹⁰Sr and ¹³⁷Cs due to fallout in the Argentine
Republic, 26:45520 (NP-19357)

COMITATO NAZIONALE PER L'ENERGIA NUCLEARE, ROME (ITALY).

DECAT: Computer code for analysis of long term pressure
and temperature transient after loss of coolant
accident in G.E. type pressure suppression containment,
26:47318 (RT/PROT-72-4)

Measurement of ¹³⁷Cs and ⁹⁰Sr in soil, 26:45263
(RT/PROT-72-11)

COMMISSARIAT A L'ENERGIE ATOMIQUE, FONTENAY-AUX-ROSES

(FRANCE). CENTRE D'ETUDES NUCLEAIRES.

Report of committee 4 on principles of environmental
monitoring related to the handling of radioactive
materials, 26:45556

COMMISSARIAT A L'ENERGIE ATOMIQUE, GRENOBLE (FRANCE).

CENTRE D'ETUDES NUCLEAIRES.

Design considerations of heat exchangers for vaporizing
fluids, 26:45436 (CEA-CONF-2010)

Heat pipes, 26:45435 (CEA-CONF-2008)

Library of data for fission products (third edition),
26:47012 (CEA-N-1526)

COMMISSARIAT A L'ENERGIE ATOMIQUE, SACLAY (FRANCE). CENTRE

D'ETUDES NUCLEAIRES.

Computer simulation of atomic displacement cascades in
metals, 26:46061 (CONF-710601-, pp 739-56)

Geometrical properties of nuclear reactions, 26:47041
(CEA-N-1529)

MASURCA: study of plutonium--graphite and uranium--

graphite assemblies, 26:47581 (CEA-N-1483)

Prog. rpt. on nuclear physics, annual, 10/1/70--10/15/71,
26:46873 (CEA-N-1522)

Report on the CEA program of investigations of radiation-
induced cavities in metals: presentation of some
results, 26:46046 (CONF-710601-, pp 31-83)

COMMONWEALTH EDISON CO., CHICAGO, ILL.

Dresden Nuclear Power Station, Unit 1. Analysis of
minimum hydro test requirements and fluence
calculations, 26:47338 (DOCKET-50010-114)

Dresden Nuclear Power Station, Unit 1. Proposed change
17 to operating license DPR-2, 26:47389 (DOCKET-50010-
113)

Dresden Nuclear Power Station, Unit 2. Diesel generator
failed to operate, 26:47357 (DOCKET-50237-253)

Dresden Nuclear Power Station, Unit 2. Proposed change
10, supplement, 26:47355 (DOCKET-50237-251)

Dresden Nuclear Power Station, Unit 2. Sample pump
tripped, 26:47356 (DOCKET-50237-252)

Dresden Nuclear Power Station, Unit 3. During monthly
surveillance testing the inboard isolation valve failed
to close, 26:47362 (DOCKET-50249-151)

Dresden Nuclear Power Station, Unit 3. Reactor pressure
scram switch drifted, 26:47332 (DOCKET-50249-148)

Dresden Nuclear Power Station, Units 2 and 3.
Environmental report; benefit--cost analysis report,
26:47404 (DOCKET-50237-249)

Dresden Nuclear Power Station, Units 2 and 3. Oil low
pressure scram switches found to have drifted,
26:47354 (DOCKET-50237-248)

Dresden Nuclear Power Station, Units 2 and 3. Quad-
Cities Station, Units 1 and 2. Results of inspection
of torus section headers, 26:47402 (DOCKET-50237-244)

Dresden Nuclear Power Station, Units 2 and 3. Quad-
Cities Station, Units 1 and 2. Operating history of
the rod worth minimizers, 26:47358 (DOCKET-50237-254)

Dresden Nuclear Power Station, Units 2 and 3. Technical
specifications, Proposed Change 18 (Unit 2); technical
specifications, Change 9 (Unit 3), 26:47405 (DOCKET-
50237-255)

La Salle County Station, Units 1 and 2. Preliminary
safety analysis report, Amendment 10, 26:47536
(DOCKET-50373-25)

Quad-Cities Station, Unit 2. Timer found to be
inoperable, 26:47325 (DOCKET-50265-101)

Quad-Cities Station, Units 1 and 2. Appendix A and
Appendix B to DPR-29 and DPR-30, proposed change,
26:47419 (DOCKET-50254-109)

Quad-Cities Station, Units 1 and 2. Final report on
condenser pump room modifications, 26:47420 (DOCKET-
50254-110)

Quad-Cities Station, Units 1 and 2. Interim report on
condenser pump room modifications, change, 26:47364
(DOCKET-50254-108)

Quad-Cities Station, Units 1 and 2. Operation of the
Quad-Cities diffuser pipes, 26:47421 (DOCKET-50254-
112)

Quad-Cities Station, Units 1 and 2. Safety analysis
report, proposed modification 72-1, 26:47418 (DOCKET-
50254-107)

COMMONWEALTH EDISON CO., CORDOVA, ILL.

Quad-Cities Station, Unit 1. Failure of RCIC motor
operated valve MO-1-1301-49 to open on signal,
26:47363 (DOCKET-50254-105)

Quad-Cities Station, Unit 2. Failure of inboard primary
containment isolation valve in the reactor core
isolation cooling system, 26:47335 (DOCKET-50265-100)

Quad-Cities Station, Unit 2. Fire in two electrical
cable trays, 26:47370 (DOCKET-50265-95)

Quad-Cities Station, Unit 2. Reactor core isolation
cooling system inboard steam supply valve failed in
open position, 26:47371 (DOCKET-50265-96)

COMMONWEALTH EDISON CO., MORRIS, ILL. (USA).

Dresden Nuclear Power Station, Unit 2. Failure of No. 2
diesel generator, 26:47331 (DOCKET-50237-250)

Dresden Nuclear Power Station, Units 2 and 3. Drift in
calibration of some of EHC oil pressure scram switches,
26:47352 (DOCKET-50237-245)

Dresden Nuclear Power Station, Units 2 and 3. Failure
to monitor plant chimney effluents July 26, 1972,
26:47353 (DOCKET-50237-247)

CONNECTICUT UNIV., FARMINGTON. DEPT. OF RADIOLOGY.

Radiobiology and the public health, 26:45911 (COO-3013-
25)

CONNECTICUT YANKEE ATOMIC POWER CO., HADDAM.

Haddam Neck Plant, Unit 1. Operation report No. 72-6
for the month of June 1972, 26:47343 (DOCKET-50213-
154)

CONNECTICUT YANKEE ATOMIC POWER CO., HARTFORD.

Haddam Neck Plant, Unit 1. Environmental report,
Appendix C, Addendum 2, 26:47396 (DOCKET-50213-150)

Haddam Neck Plant, Unit 1. Environmental report,
Appendix E; aquatic benefit--cost, 26:47397 (DOCKET-
50213-152)

CONSOLIDATED EDISON CO. OF NEW YORK, INC., NEW YORK.

Indian Point Nuclear Generating Unit 2. Safety and
relief valve installation and re-analysis for ASME
class 1 and class 2 systems, 26:47412 (DOCKET-50247-
76)

Indian Point Nuclear Generating Unit 2. Summary report

of control rod drive line malfunction, 26:47322 (DOCKET-50247-77)

Indian Point Nuclear Generating Unit 3. Final facility description and safety analysis report, Amendment 18, Supplement 4, 26:47455 (DOCKET-50286-46)

Indian Point Nuclear Generating Unit 3. Final facility description and safety analysis report, Supplement 6, 26:47457 (DOCKET-50286-48)

Indian Point Nuclear Generating Unit 3. Final facility description and safety analysis report, Supplement 7, 26:47458 (DOCKET-50286-49)

Indian Point Nuclear Generating Unit 3. Final safety analysis report, Supplement 5, 26:47456 (DOCKET-50286-47)

Indian Point Station, Unit 1. Measures taken to repair control rod and safety implications of the incident, 26:47320 (DOCKET-50003-118)

Indian Point Station, Unit 1. Secondary coolant activity in excess of $1.5 \times 10^{-2} \mu \text{ ci/ml}$, 26:47337 (DOCKET-50003-119)

Indian Point Station, Unit 1. Topical report on General Electric induction motors: inside containment low pressure core cooling, 26:47388 (DOCKET-50003-120)

CONSUMERS POWER CO., JACKSON, MICH.

Big Rock Point Nuclear Plant. Technical specification Change Request 31, errata, 26:47393 (DOCKET-50155-117)

Big Rock Point Nuclear Plant. Technical specifications, proposed change 33, 26:47394 (DOCKET-50155-119)

Big Rock Point Nuclear Plant. Technical specifications, proposed change 34, 26:47395 (DOCKET-50155-120)

Palisades Plant. Application for construction permit and operating license, Amendment 27, 26:47422 (DOCKET-50255-94)

Palisades Plant. Environmental impact of plant operation up to July 1, 1972, 26:47424 (DOCKET-50255-96)

Palisades Plant. Technical specifications, request for change 1, 26:47423 (DOCKET-50255-95)

CORNELL UNIV., ITHACA, N. Y. DEPT. OF MATERIALS SCIENCE AND ENGINEERING.

Void formation in quenched or irradiated metals, 26:46058 (CONF-710601-, pp 563-604)

CORNELL UNIV., ITHACA, N. Y. LAB. OF NUCLEAR STUDIES.

Detection of large cosmic-ray showers by atmospheric fluorescence. Final report, 26:46150 (NYO-3038-13)

D

DAIRYLAND POWER COOPERATIVE, LA CROSSE, WIS.

La Crosse Boiling Water Reactor. Application of conversion of the provisional operating authorization to a provisional operating license, Amendment 1, 26:47570 (DOCKET-50409-5)

La Crosse Boiling Water Reactor. Forced circulation system behavior, 26:47330 (DOCKET-50409-7)

La Crosse Boiling Water Reactor. Incident report, hydrogen detonation in main condenser off-gas system, 26:47383 (DOCKET-50409-8)

La Crosse Boiling Water Reactor. Incident reports, 26:47382 (DOCKET-50409-6)

La Crosse Boiling Water Reactor. Monthly Operating Report, June 1972, 26:47384 (DOCKET-50409-9)

La Crosse Boiling Water Reactor. Technical specifications, change in plant organization, 26:47569 (DOCKET-50409-4)

DANISH ATOMIC ENERGY COMMISSION, RISØ. RESEARCH ESTABLISHMENT.

Positron lifetimes in water and ice, and in frozen aqueous solutions, 26:46362 (RISØ-254)

DEPARTMENT OF JUSTICE, WASHINGTON, D.C. ANTITRUST DIV.

Edwin I. Hatch Nuclear Plant, Unit 2. Department of Justice Opinion with respect to monopoly, 26:47530 (DOCKET-50366-32)

Mendocino Power Plant, Units 1 and 2. Opinion of Department of Justice with respect to monopolistic activity, 26:47562 (DOCKET-50398-17)

North Anna Power Station, Units 3 and 4. Department of Justice Opinion with respect to monopolistic action, 26:47568 (DOCKET-50404-20)

DETROIT EDISON CO., MICH.

Enrico Fermi Atomic Power Plant, Unit 2. Cracks in the basement floor slab of the reactor building, 26:47513 (DOCKET-50341-42)

DEUTSCHES ELEKTRONEN-SYNCHROTRON, HAMBURG (WEST GERMANY).

Currents, stress tensor, and generalized unitarity in conformal invariant quantum field theory, 26:46697 (DESY-72/20)

Diffraction--dissociation model for high-energy electroproduction, 26:46517 (DESY-72/23)

π^0 electroproduction at first resonance, 26:46512 (CONF-710816-16)

Testing triplet models, 26:46516 (DESY-72/21)

DIRECTORATE OF LICENSING (AEC), WASHINGTON, D.C.

Arkansas Nuclear One, Unit 2. Draft environmental statement, 26:47532 (DOCKET-50368-31)

Dresden Nuclear Power Station, Unit 3. Reactor protection system limit switch on No. 2 turbine control valve found to be inoperable, 26:47473 (DOCKET-50309-61)

Edwin I. Hatch Nuclear Plant, Unit 2. Safety evaluation, 26:47529 (DOCKET-50366-31)

Fort Calhoun Station, Unit 1. Safety evaluation, 26:47454 (DOCKET-50285-68)

Kewaunee Nuclear Power Plant. Draft Environmental Statement, 26:47471 (DOCKET-50305-45)

La Salle County Station, Units 1 and 2. Draft environmental statement, 26:47535 (DOCKET-50373-24)

McGuire Nuclear Station, Units 1 and 2. Draft environmental statement, 26:47534 (DOCKET-50369-28)

Midwest Fuel Recovery Plant. Notice of Consideration of Issuance of a Facility Operating License and Notice of Opportunity for Hearing, 26:45402 (DOCKET-50268-31)

Oconee Nuclear Station, Units 2 and 3. Notice of consideration of issuance of facility operating licenses and notice of opportunity for hearing, 26:47434 (DOCKET-50270-49)

Oconee Nuclear Station, Unit 1. Order extending provisional construction permit completion date, 26:47431 (DOCKET-50269-56)

Point Beach Nuclear Plant, Unit 2. Facility Operating License No. DPR-27, Amendment 2, 26:47469 (DOCKET-50301-64)

Southwest Experimental Fast Oxide Reactor. Technical specifications, Change 13, 26:47584 (DOCKET-50231-122)

Turkey Point Plant, Units 3 and 4. Facility operating license DPR-31 with appended technical specifications, 26:47414 (DOCKET-50250-85)

Turkey Point Plant, Units 3 and 4. Final Environmental Statement, 26:47413 (DOCKET-50250-84)

Vermont Yankee Nuclear Power Station. Technical specifications, Change 1, 26:47439 (DOCKET-50271-94)

West Valley Reprocessing Plant. Comments concerning nuclear criticality, 26:45401 (DOCKET-50201-128)

DIRECTORATE OF REGULATORY OPERATIONS (AEC), ATLANTA, GA. (USA). REGION II.

Edwin I. Hatch Nuclear Plant, Unit 1. Steps taken to correct items of apparent noncompliance which were brought to attention in letter of July 5, 1972, 26:47495 (DOCKET-50321-48)

Edwin I. Hatch Nuclear Plant, Unit 2. Steps taken to correct items of apparent noncompliance brought to attention in letter dated June 28, 1972, 26:47531 (DOCKET-50366-33)

Joseph M. Farley Nuclear Plant, Units 1 and 2. Steps taken to correct items of apparent noncompliance brought to attention in letter dated June 21, 1972, 26:47519 (DOCKET-50348-42)

Oconee Nuclear Station, Units 1, 2, and 3. Steps taken with respect to item brought to attention letter dated April 21, 1972, 26:47433 (DOCKET-50269-61)

DIRECTORATE OF REGULATORY OPERATIONS (AEC), GLEN ELLYN, ILL. REGION III.

Davis--Besse Nuclear Power Station. Steps taken to correct items of apparent nonconformance noted in letter dated May 22, 1972, 26:47517 (DOCKET-50346-27)

Dresden Nuclear Power Station, Units 2 and 3. Steps taken to correct items of apparent noncompliance called to attention in letter of June 19, 1972, 26:47403 (DOCKET-50237-246)

Kewaunee Nuclear Power Plant. Steps taken to correct item of apparent noncompliance noted in letter dated July 17, 1972, 26:47472 (DOCKET-50305-47)

Point Beach Nuclear Plant, Units 1 and 2. Discussion of items of apparent noncompliance, 26:47430 (DOCKET-50266-74)

Zion Station Nuclear Power Plant, Units 1 and 2. Accident report, April 24, 1972, 26:47463 (DOCKET-50295-58)

Zion Station Nuclear Power Plant, Units 1 and 2. Steps taken to correct items of apparent noncompliance in letter of June 12, 1972, 26:47462 (DOCKET-50295-57)

Zion Station Nuclear Power Plant, Units 1 and 2. Steps taken to correct items of apparent noncompliance brought to attention in letter dated June 23, 1972, 26:47464 (DOCKET-50295-59)

DIRECTORATE OF REGULATORY OPERATIONS (AEC), NEWARK, N.J. REGION I.

Beaver Valley Power Station, Unit 1. Action taken to correct items of nonconformance and noncompliance found on inspection of April 18--20, 1972, 26:47503 (DOCKET-50334-31)

DIRECTORATE OF REGULATORY OPERATIONS (AEC), WASHINGTON, D.C.

Calvert Cliffs Nuclear Power Plant, Units 1 and 2. Failure of polar crane upper limit switch, 26:47381 (DOCKET-50317-53)

Davis--Besse Nuclear Power Station. Shipping damage to containment vessel equipment hatch, 26:47326 (DOCKET-50346-30)

Donald C. Cook Nuclear Plant, Units 1 and 2. Defective welding on ice condenser system positioning frames, 26:47491 (DOCKET-50315-69)

Donald C. Cook Nuclear Plant, Units 1 and 2. Deficiency in radiography techniques, 26:47492 (DOCKET-50315-70)

Enrico Fermi Atomic Power Plant, Unit 2. Deformation of containment drywell, 26:47514 (DOCKET-50341-43)

Enrico Fermi Atomic Power Plant, Unit 2. Reactor pressure vessel ultrasonic indications, 26:47329 (DOCKET-50341-44)

Newbold Island Nuclear Generating Station, Unit 1. Inspection of reactor pressure vessel fabrication at Hitachi, Ltd, 26:47521 (DOCKET-50354-25)

Prairie Island Nuclear Generating Plant, Units 1 and 2. Report of corrosion problem in refueling water storage tanks, 26:47452 (DOCKET-50282-68)

Rancho Seco Nuclear Generating Station, Unit 1. Report on minor imperfections found in pipe welds, 26:47484 (DOCKET-50312-47)

Rancho Seco Nuclear Generating Station, Unit 1. Weld defects found in steam generator lateral support, 26:47485 (DOCKET-50312-48)

Robert Emmett Ginna Nuclear Power Plant, Unit 1. Report of exposure of an individual to radiation, 26:47359 (DOCKET-50244-106)

Three Mile Island Nuclear Station, Unit 1. Report of significant deficiency/thin walled valves, 26:47460 (DOCKET-50289-62)

Zion Station Nuclear Power Plant, Unit 2. Leaks in canopies of two control rod drive assemblies, 26:47379 (DOCKET-50304-59)

DIVISION OF COMPLIANCE (AEC), NEWARK, N.J. REGION I.

Three Mile Island Nuclear Station, Unit 1. Action taken to correct item of nonconformance found on May 4, 1972 inspection, 26:47459 (DOCKET-50289-61)

DIVISION OF NUCLEAR MATERIALS SECURITY (AEC), WASHINGTON, D.C.

Attribute sampling inspection procedure based on the hypergeometric distribution, 26:46094 (WASH-1210)

DIVISION OF REACTOR DEVELOPMENT AND TECHNOLOGY (AEC), WASHINGTON, D. C.

Gaskets--containment vessel airlock, 26:47593 (RDT-E-14-6-T(6-72))

DU PONT DE NEMOURS (E.I.) AND CO., AIKEN, S. C. SAVANNAH RIVER LAB.

Savannah River Laboratory ⁹⁰Co power and heat sources. Quarterly progress report, January--March 1972, 26:45701 (DP-1300)

DUKE POWER CO., CHARLOTTE, N.C.

McGuire Nuclear Station, Units 1 and 2. Application for licenses, Amendment 12, 26:47533 (DOCKET-50369-26)

Oconee Nuclear Station, Units 1, 2, and 3. Final safety analysis report, Amendment 33, 26:47432 (DOCKET-50269-60)

DUKE UNIV., DURHAM, N.C. DEPT. OF PHYSICS.

Progress report [on high-energy physics], July 1, 1971--June 30, 1972, 26:46479 (ORO-3065-24)

E

EG AND G, INC., GOLETA, CALIF.

Ecological and environmental effects from local fallout from Schooner. 2. The beta and gamma radiation effects from close-in fallout, 26:45542 (PNE-529)

ENVIRONMENTAL RESEARCH CORP., LAS VEGAS, NEV.

Analysis of seismic waves recorded in Las Vegas, Nevada from underground nuclear detonations, 26:45481 (NVO-1163-TM-33)

Comparison of response spectra from Yucca Flat and Pahute Mesa underground nuclear detonations, 26:45480 (NVO-1163-TM-32)

EUROPEAN ATOMIC ENERGY COMMUNITY, ISPRA (ITALY). JOINT NUCLEAR RESEARCH CENTER.

Site characteristics, 26:45570 (EUR-4841)

Trace level determination of oxygen in organic coolants and ultra-micro determination of oxygen in C-H--O organic compounds, 26:45221 (EUR-4822)

EUROPEAN ATOMIC ENERGY COMMUNITY, LUXEMBOURG. CENTER FOR INFORMATION AND DOCUMENTATION.

Definitions of Proper Name Descriptors used in the Euratom-INIS Nuclear Documentation System, 26:47597 (EUR-4806)

EUROPEAN COMMUNITIES, BRUSSELS (BELGIUM). COMMISSION.

Definitions of Proper Name Descriptors used in the Euratom-INIS Nuclear Documentation System, 26:47597 (EUR-4806)

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH, GENEVA (SWITZERLAND).

Methods of nonlinear analysis, 26:47283 (CERN-72-8)

F

FLORIDA POWER AND LIGHT CO.

Hutchinson Island Plant, Unit 1. Environmental report, Supplement 5, 26:47504 (DOCKET-50335-29)

Hutchinson Island Plant, Unit 1. Environmental report, Supplement 6, 26:47505 (DOCKET-50335-30)

Hutchinson Island Plant, Unit 1. Environmental report, supplement 7, 26:47506 (DOCKET-50335-32)

Turkey Point Plant, Unit 4. Structural integrity test report and containment surveillance during the structural integrity test, 26:47417 (DOCKET-50251-72)

Turkey Point Plant, Units 3 and 4. Abnormal occurrences 3-72-1 and 3-72-2, 26:47415 (DOCKET-50250-86)

Turkey Point Plant, Units 3 and 4. Technical specifications, proposed changes, 26:47416 (DOCKET-50250-88)

FLORIDA STATE UNIV., TALLAHASSEE. COLL. OF ARTS AND SCIENCES.

Effects of gamma radiation on the spectral sensitivity of the noctuid moth, *Heliothis Zea*, 26:45873 (TID-26078)

FLORIDA STATE UNIV., TALLAHASSEE. INST. OF MOLECULAR BIOPHYSICS.

Research proposal. Annual report No. 12, Part B:

5. CORPORATES

HANFORD ENGINEERING DEVELOPMENT LAB.,

September 1, 1971--August 31, 1972, 26:45891 (ORO-2690-12(Pt.B))

FONDATION INDUSTRIE-UNIVERSITE, BRUSSELS (BELGIUM).

Postgraduate training of scientists in the European Community, 26:47616 (EUR-4832)

FORSVARETS FORSKNINGSANSTALT, STOCKHOLM (SWEDEN).

Recent studies on prompt neutron and gamma rays from fission, 26:47020 (INDC-7-U, pp 113-29)

G

GENERAL ELECTRIC CO., CINCINNATI, OHIO.

Conceptual design study of a nuclear Brayton turboalternator--compressor, 26:45424 (N-72-15607)

GENERAL ELECTRIC CO., PLEASANTON, CALIF. VALLECITOS NUCLEAR CENTER.

Fluence and temperature dependence of void formation in highly irradiated stainless steels, 26:46048 (CONF-710601-, pp 156-73)

GENERAL ELECTRIC CO., SAN JOSE, CALIF. BREEDER REACTOR DEPT.

Deuteron-generated void formation in austenitic stainless steels, 26:46053 (CONF-710601-, pp 479-98)

GENERAL ELECTRIC CO., SAN JOSE, CALIF. NUCLEAR ENERGY DIV.

Midwest Fuel Recovery Plant. Final safety analysis report, Amendment 27, 26:45403 (DOCKET-50268-32)

Midwest Fuel Recovery Plant. Final safety analysis report, Amendment 28, 26:45404 (DOCKET-50268-33)

GENERAL ELECTRIC CO., SCHENECTADY, N. Y. RESEARCH AND DEVELOPMENT CENTER.

Solute segregation and stresses generated around growing voids in metals, 26:45988 (CONF-710601-, pp 630-46)

GENERAL ELECTRIC CO., SUNNYVALE, CALIF. BREEDER REACTOR DEPT.

Fluence and temperature dependence of void formation in highly irradiated stainless steels, 26:46048 (CONF-710601-, pp 156-73)

Southwest Experimental Fast Oxide Reactor. Modification proposed for Decommissioning Plan, 26:47582 (DOCKET-50231-120)

Southwest Experimental Fast Oxide Reactor. Request for issuance of dismantling order, 26:47583 (DOCKET-50231-121)

GEOLOGICAL SURVEY, DENVER, COLO.

Geologic and hydrologic effects of the Cannikin underground nuclear explosion, Amchitka Island, Aleutian Islands, Alaska. Amchitka-33, 26:45488 (USGS-474-148)

GEORGIA POWER CO., ATLANTA.

Edwin I. Hatch Nuclear Plant, Unit 1. Final safety analysis report, Amendment 17, 26:47494 (DOCKET-50321-46)

Edwin I. Hatch Nuclear Plant, Unit 2. Preliminary safety analysis report, Amendment 16, 26:47528 (DOCKET-50366-29)

GOSUDARSTVENNYI KOMITET PO ISPOL'ZOVANIYU ATOMNOI ENERGII SSSR, MOSCOW. INSTITUT TEORETICHESKOI I EKSPERIMENTAL'NOI FIZIKI.

Concerning the two-photon mechanism of production during collision of charged particles, 26:46518 (ITEF-848)

GOSUDARSTVENNYI KOMITET PO ISPOL'ZOVANIYU ATOMNOI ENERGII SSSR, OBNINSK. FIZIKO-ENERGETICHESKII INSTITUT.

Study of the characteristics of emergency track detectors, 26:45617 (FEI-296)

GOSUDARSTVENNYI KOMITET PO ISPOL'ZOVANIYU ATOMNOI ENERGII SSSR, SERPUKHOV. INSTITUT FIZIKI VYSOKIKH ENERGII.

Analytical properties of amplitudes and differential cross-sections for multiparticle production processes over two angular variables, 26:46497 (IFVE-STF-71-36)

Estimation of intermediate energy contribution to the dispersion sum rules and rotation of polarization in pion-nucleon scattering at high energies, 26:46617 (IFVE-STF-71-15)

Intranuclear cascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)

Total cross-sections for reaction $\pi^-p \rightarrow \pi^0n$ at high energies, 26:46618 (IFVE-STF-71-71)

GRENOBLE UNIV. (FRANCE). FACULTE DES SCIENCES.

Chemical separation and high resolution nuclear spectroscopy in $^{228}\text{Ra} + ^{228}\text{Ac} + ^{228}\text{Th}$ radioactive chain, 26:47014 (FRNC-TH-184)

GULF GENERAL ATOMIC CO., SAN DIEGO, CALIF.

Computer simulation of pulsed-neutron experiments for the Fort St. Vrain HTGR, 26:47319 (GULF-GA-B-10959)

Irradiation-induced dimensional change in HTGR control materials, 26:46077 (GULF-GA-A-12035)

H

HAMBURG UNIV. (WEST GERMANY). INSTITUT FUER THEORETISCHE PHYSIK (II).

Testing triplet models, 26:46516 (DESY-72/21)

HANFORD ENGINEERING DEVELOPMENT LAB., RICHLAND, WASH.

Effects of pore structure on the melting heat rating of oxide fuels, 26:45930 (HEDL-TME-72-81)

Recovery and recrystallization of prototypic FTR fuel cladding: 20 percent CW 316 stainless steel tubing (RM-11), 26:46067 (HEDL-TME-71-126)

- Suppression of void formation in stainless steel, 26:46049 (CONF-710601-, pp 174-86)
- Three dimensional neutronics calculations for the Fast Test Reactor (FTR) and the FTR Engineering Mock Up Critical assembly (EMC), 26:47586 (HEDL-TME-72-42)
- HEIDELBERG UNIV. (WEST GERMANY). INSTITUT FUER THEORETISCHE PHYSIK.
- Simple theory of proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46650 (KFK-1547)
- ILLINOIS UNIV., URBANA.
- Missing-mass spectrum near 960 MeV from $\pi^-p \rightarrow n + (\pi\pi)$, 26:46430 (COO-1195-241)
- Study of A_2 in $\pi^-p \rightarrow \pi^+\pi^-p$, 26:46473 (COO-1195-239)
- ILLINOIS UNIV., URBANA. DEPT. OF COMPUTER SCIENCE.
- Part 1: Annual technical progress report, October 1, 1971-June 30, 1972, 26:47610 (COO-2118-34)
- INDIANA AND MICHIGAN POWER CO., NEW YORK.
- Donald C. Cook Nuclear Plant, Units 1 and 2.
- Environmental report, Supplement 3, 26:47489 (DOCKET-50315-65)
- Donald C. Cook Nuclear Plant, Units 1 and 2. Final safety analysis report, Amendment 25, 26:47490 (DOCKET-50315-68)
- Donald C. Cook Nuclear Plant, Units 1 and 2. Final safety analysis report, Amendment 26, 26:47493 (DOCKET-50315-71)
- INDIANA-PURDUE UNIV., INDIANAPOLIS. DEPT. OF PHYSICS.
- Final states with three charged particles and a visible Λ from K^-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)
- INSTITUTO DE ENERGIA ATOMICA, SAO PAULO (BRAZIL).
- Optimization study of the precipitation of thorium oxalate, 26:45365 (IEA-183)
- Selected topics on dynamics of crystal lattices, 26:47617 (IEA-19)
- INSTITUTT FOR ATOMENERGI, KJELLER (NORWAY).
- Radioisotopes, radioactive pharmaceuticals, radioactive chemicals, radioactive sources, services, 26:45709 (INIS-mf-256)
- INTERNATIONAL ATOMIC ENERGY AGENCY, VIENNA (AUSTRIA).
- Conference on neutron induced fission, Bombay, 7/14/71, 26:47015 (INDC-7-U)
- Release of radioactive wastes into the sea, 26:46101 (IAEA-136, pp 10-17)
- INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS, TRIESTE (ITALY).
- Asymptotic numbers: one generalization of the notion of number, 26:47284 (IC-72/45)
- Conjugacy and stability theorems in connection with contraction of Lie algebras, 26:46666 (IC-72/49)
- Virtual neutrino effects, 26:46570 (IC-72/54)
- INTERNATIONAL COMMISSION ON RADIATION UNITS AND MEASUREMENTS, WASHINGTON, D.C.
- Measurement of low-level radioactivity, 26:45261 (ICRU-22)
- Radiation dosimetry: electrons with initial energies between 1 and 50 MeV, 26:45618 (ICRU-21)
- INTERNATIONAL NUCLEAR DATA COMMITTEE.
- Conference on neutron induced fission, Bombay, 7/14/71, 26:47015 (INDC-7-U)
- IOWA ELECTRIC LIGHT AND POWER CO., CEDAR RAPIDS.
- Duane Arnold Energy Center. Environmental report revised, Amendment 1, 26:47498 (DOCKET-50331-48)
- Duane Arnold Energy Center. Final safety analysis report, Amendment 2, 26:47499 (DOCKET-50331-49)
- Duane Arnold Energy Center. Final safety analysis report, Amendment 3, 26:47500 (DOCKET-50331-51)
- IOWA STATE UNIV. OF SCIENCE AND TECHNOLOGY, AMES.
- Observation of the $M(940)$ and $\delta(963)$, 26:46475 (IS-2886)
- JAGELLONIAN UNIV., KRAKOW (POLAND). DEPT. OF THEORETICAL PHYSICS.
- Comparison of the angular decay distributions in three quark models of photoproduction, 26:46526 (TPJU-14/71)
- JAPAN ATOMIC ENERGY RESEARCH INST., TOKYO.
- Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)
- JERSEY CENTRAL POWER AND LIGHT CO., MORRISTOWN, N. J.
- Forked River Nuclear Generating Station, Unit 1.
- Environmental report, correction of cost-benefit analysis, 26:47526 (DOCKET-50363-32)
- Forked River Nuclear Generating Station, Unit 1.
- Environmental report, commercial fishing in Barnegat Bay, 26:47527 (DOCKET-50363-34)
- Forked River Nuclear Generating Station, Unit 1.
- Preliminary safety analysis report, Amendment 19, 26:47524 (DOCKET-50363-30)
- Forked River Nuclear Generating Station, Unit 1.
- Preliminary safety analysis report, Amendment 20, 26:47525 (DOCKET-50363-31)
- Oyster Creek Nuclear Power Plant, Unit 1. Containment spray pump failure, 26:47348 (DOCKET-50219-204)
- Oyster Creek Nuclear Power Plant, Unit 1. Experience with the rod worth minimizer, 26:47346 (DOCKET-50219-199)
- Oyster Creek Nuclear Power Plant, Unit 1. Final safety analysis report, Amendment 70, 26:47398 (DOCKET-50219-200)
- Oyster Creek Nuclear Power Plant, Unit 1. Outside radwaste tank activity, 26:47399 (DOCKET-50219-201)
- Oyster Creek Nuclear Power Plant, Unit 1. Relay failure, 26:47344 (DOCKET-50219-196)
- Oyster Creek Nuclear Power Plant, Unit 1. Surveillance failure, 26:47347 (DOCKET-50219-203)
- Oyster Creek Nuclear Power Plant, Unit 1. Surveillance test failures, 26:47345 (DOCKET-50219-197)
- Oyster Creek Nuclear Power Plant, Unit 1. Technical specifications, revised pages incorporating all changes through No. 12, 26:47400 (DOCKET-50219-202)
- JOHNS HOPKINS UNIV., BALTIMORE, MD. CHESAPEAKE BAY INST.
- Study on the chemical behavior of zinc in Chesapeake Bay water using Anodic Stripping Voltammetry. Technical report 76, 26:47613 (COO-3292-1)
- JOHNS HOPKINS UNIV., BALTIMORE, MD. DEPT. OF BIOCHEMISTRY.
- Chemical study of the effects of radiation on nucleic acids and related compounds. Three year progress report, June 1969-May 1972, 26:45315 (COO-3286-3)
- Radiation chemistry of nucleic acids. Isolation and characterization of thymine glycols, 26:45314 (COO-3286-1)
- JOHNS HOPKINS UNIV., BALTIMORE, MD. DEPT. OF PHYSICS.
- Inelastic e-p scattering and the polarization contribution to H hyperfine structure, 26:46523 (NYO-4076-17)
- Lightlike quark model and current densities, 26:46524 (NYO-4076-18)
- Wave packet realization of lightlike states, 26:46681 (NYO-4076-19)
- JOHNS HOPKINS UNIV., BALTIMORE, MD. DEPT. OF RADIOLOGICAL SCIENCES.
- Radiation hazards in environmental medicine, 26:45904 (COO-3013-1)
- JOHNS HOPKINS UNIV., BALTIMORE, MD. DEPT. OF RADIOLOGY.
- Environmental radioactivity: radiation effects at the cellular level, 26:45890 (NYO-3970-29)
- JOINT INST. FOR NUCLEAR RESEARCH, DUBNA (USSR).
- Change in the mechanism of inelastic interactions between particles and nuclei in the energy region T approximately equal to 5 GeV, 26:47038 (BNL-tr-487)
- Cross section of pion-nucleon interactions, 26:46478 (ORNL-tr-2544)
- Influence of the electromagnetic ring acceleration on the focusing by image forces, 26:46136 (JINR-P9-6421)
- Quasi-threshold behavior of $\pi + N \rightarrow \pi + N + (\gamma\text{-virtual})$ process amplitude, 26:46620 (JINR-P4-6475)
- Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)
- JOINT INST. FOR NUCLEAR RESEARCH, DUBNA (USSR). LAB. OF HIGH ENERGY.
- Investigation of generation of particles and resonance states in interactions of 2.34 GeV/c, 5 GeV/c, and 9 GeV/c π^- -mesons with xenon nuclei, 26:46476 (JINR-P1-6408)
- Spectrum symmetry of pion longitudinal momenta in π^-p interactions at $p_c = 4$ GeV, 26:46619 (JINR-P1-6461)
- JOINT INST. FOR NUCLEAR RESEARCH, DUBNA (USSR). LAB. OF NUCLEAR PROBLEMS.
- Electromagnetic effects in nucleon polarization at meson-nucleon scattering, 26:46520 (JINR-P2-6438)
- New method of determination of physical-chemical properties of matter, 26:47618 (JINR-P14-6407)
- Signal filtration in the amplitude spectrometry systems, 26:45637 (JINR-P13-6428)
- Study of ^{179}Re decay, 26:46976 (JINR-P6-6492)
- Study of the decay $^{180}\text{Tb}/\text{sub } 1/2 = 66 \text{ min}$ and $^{180}\text{Tb}/\text{sub } 1/2 = 3.48 \pm 0.16 \text{ h}$, 26:46942 (JINR-P6-6512)
- JOINT INST. FOR NUCLEAR RESEARCH, DUBNA (USSR). LAB. OF NUCLEAR REACTIONS.
- Angular distribution of fragments of spontaneously fissioning isomers, 26:47022 (JINR-P7-6466)
- Study of transuranium element hydrolysis. I, 26:45366 (JINR-P6-6402)
- JOINT INST. FOR NUCLEAR RESEARCH, DUBNA (USSR). LAB. OF THEORETICAL PHYSICS.
- Account of nuclear boundary diffusivity in the intranuclear cascade model, 26:47047 (JINR-P2-6503)
- Analysis of inelastic collisions of particles with nuclei at very high energies, 26:46649 (JINR-P2-6395)
- Asymptotic estimates of quasipotential parameters, 26:46703 (JINR-P2-6482)
- Basic state of many-fermion system, coupled by gravitational lines, 26:46748 (JINR-P2-6270)
- Corrections to eikonal representation of amplitude in scalar field theory, 26:46699 (JINR-P2-6394)
- Feynman-Wilson-gas model as consequence of jet approach to inelastic hadron processes, 26:46586 (JINR-P2-6501)
- Investigation of large angle scattering in quasipotential approach, 26:46776 (JINR-P2-6442)
- Lower unboundedness of the Hamiltonian with scalar electrodynamics, 26:46702 (JINR-P2-6474)
- Operator method of solving the quasipotential equations and a conception of straight-line paths at high energies, 26:46700 (JINR-P2-6437)
- Parametrization of $B(E2)$ -transition in deformed even-

even nuclei within SU(3) scheme, 26:46975 (JINR-P4-6462)

Possible existence of SU(3) multiplets in spectra of deformed even-even nuclei, 26:47048 (JINR-P4-6452)
Quantum field theory: "dressing" contra divergencies, 26:46701 (JINR-P2-6454)

Reaction $\gamma + \gamma \rightarrow \pi + \pi$ and electromagnetic difference for pion masses, 26:46519 (JINR-P2-6333)

S-matrix unitarity in theories with nonpolynomial Lagrangians, 26:46775 (JINR-P2-6393)

SU(3) physical basis realization and probabilities of B(E2)-transitions in SU(3) scheme. II, 26:47049 (JINR-P4-6458)

JUNTA DE ENERGIA NUCLEAR, MADRID (SPAIN).

Green function in the secondary model of thermalization, 26:46846 (JEN-221)

New approach to beta-gamma coincidence counting.

Advance report on SAMAR electronic system, 26:45589 (JEN-247)

JUNTA DE ENERGIA NUCLEAR, MADRID (SPAIN). DIVISION DE METALLURGIA.

Table of cross section (n,p), (n, α), and (n,2n) reactions in steel components and other nuclear materials, 26:46918 (JEN-230-DME/I-25)

JUNTA DE ENERGIA NUCLEAR, MADRID (SPAIN). DIVISION DE QUIMICA.

Application of spectrographic analysis to the radioisotope production control. II, 26:45223 (JEN-232-DQ/I-81)

Volatilization mechanism of certain elements in a method of semiquantitative spectrographic analysis, 26:45222 (JEN-229-DQ/I-79)

K

KANSAS UNIV., LAWRENCE. NUCLEAR STRUCTURE LAB.

Final report, 26:46903 (COO-1120-149)

KARLSRUHE UNIV. (WEST GERMANY).

Proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46457 (KFK-1509)

KERNFORSCHUNGSANLAGE, JUELICH (WEST GERMANY). INSTITUT FUER REAKTORENTWICKLUNG.

Absolute measurement of the $^9\text{Be}(n,2n)$ cross section in the energy region of 2.37 to 3.24 MeV, 26:46874 (JUL-834-RG)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY).

Environmental burden of tritium, 26:45509 (KFK-1560)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY). INSTITUT FUER ANGEWANDTE KERNPHYSIK.

Measure of ratio of capture-to-fission cross section for ^{235}U and ^{239}Pu between 8 and 60 keV neutron energy, 26:47024 (KFK-1563)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY). INSTITUT FUER EXPERIMENTELLE KERNPHYSIK.

Proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46457 (KFK-1509)

Simple theory of proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46650 (KFK-1547)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY). INSTITUT FUER MATERIAL UND FESTKORPERFORSCHUNG.

Concept of an irradiation facility for investigation of high-temperature thermoelements at temperatures to 2000°C, 26:45438 (KFK-1556)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY). INSTITUT FUER NEUTRONENPHYSIK UND REAKTORTECHNIK.

Influence of fission neutron spectra on integral nuclear quantities of fast reactors, 26:47585 (EURFNR-1008)

Recuperative heat exchanger for a fast helium-cooled breeder reactor with helium turbine, 26:45439 (KFK-1566)

KERNFORSCHUNGSZENTRUM, KARLSRUHE (WEST GERMANY). INSTITUT FUER RADIOCHEMIE.

Mean excitation energies of the daughter nuclei on β -decay, 26:46875 (KFK-EXT-10/72-1)

Preparation and investigation of neptunium isotopes ^{231}Np , ^{232}Np , and ^{233}Np , 26:47023 (KFK-1537)

KNOLLS ATOMIC POWER LAB., SCHENECTADY, N. Y.

Fabrication of aluminum oxide films, 26:45931 (KAPL-3164)

L

LABORATOIRE CENTRAL DE L'ARMEMENT, ARCUEIL (FRANCE).

Biological effects of the intensity of irradiation, 26:45816 (FRNC-R-7)

LEHIGH UNIV., BETHLEHEM, PA. DEPT. OF METALLURGY AND MATERIALS SCIENCE.

Forming of composite for superconducting rod. Technical progress report, 26:45972 (COO-3311-1)

LOS ALAMOS SCIENTIFIC LAB., N. MEX.

Cells for nondestructive examinations of irradiated fuel pins, 26:46104 (LA-DC-72-556)

Feedback stabilization on an $l = 1$ theta-pinch plasma column, 26:47157 (LA-DC-72-707)

Importance of the quadrupole pairing field in the J/ψ $n/\pi = 0^+$ vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)

Origins and history of the Los Alamos Meson Physics Facility, 26:46119 (LA-5000)

Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)

Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)

Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)

Remotized apparatus for determining oxygen in irradiated reactor fuels and cladding materials, 26:45224 (LA-DC-72-524)

Space Electric Power R and D Program. Quarterly report for the period ending April 30, 1972, 26:45702 (LA-4978)

LOUISIANA POWER AND LIGHT CO., NEW ORLEANS.

Waterford Steam Electric Station, Unit 3. Application for construction exemption, 26:47541 (DOCKET-50382-29)

Waterford Steam Electric Station, Unit 3. Basic data for a source term calculation, 26:47542 (DOCKET-50382-30)

Waterford Steam Electric Station, Unit 3. Environmental report, supplement 1, 26:47539 (DOCKET-50382-27)

Waterford Steam Electric Station, Unit 3. Environmental report, Supplement 2, 26:47543 (DOCKET-50382-31)

Waterford Steam Electric Station, Unit 3. Preliminary safety analysis report, Amendment 17, 26:47538 (DOCKET-50382-25)

Waterford Steam Electric Station, Unit 3. Preliminary safety analysis report, Amendment 19, 26:47540 (DOCKET-50382-28)

Waterford Steam Electric Station, Unit 3. Preliminary safety analysis report, Amendment 20, 26:47544 (DOCKET-50382-32)

LOUISIANA STATE UNIV., BATON ROUGE. DEPT. OF ZOOLOGY AND PHYSIOLOGY.

Mutagenic effect of transmutation from radionuclides incorporated into DNA of *Drosophila melanogaster*. Progress report, September 1970-May 1972, 26:45912 (ORO-3728-5)

LYON UNIV. (FRANCE). INSTITUT DE PHYSIQUE NUCLEAIRE.

Final state interaction between proton and neutron from $^1\text{H}(d, pn)^1\text{H}$, $^2\text{H}(d, pn)^2\text{H}$, $^3\text{He}(d, pn)^3\text{He}$, and $^4\text{He}(d, pn)^4\text{He}$ reactions, 26:46860 (LYCEN-7186)

Generating function method and developments in quasi-boson, 26:47286 (LYCEN-7202)

M

MAGYAR TUDOMANYOS AKADEMIA KOZPONTI FIZIKAI KUTATO INTEZET, BUDAPEST.

New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)

Spontaneous decay in a time dependent external field, 26:46571 (KFKI-72-15)

MAINE YANKEE ATOMIC POWER CO., WESTBORO, MASS.

Maine Yankee Atomic Power Station. Environmental qualification of vital instrumentation and electrical cables, and penetration connectors, 26:47475 (DOCKET-50309-63)

Maine Yankee Atomic Power Station. Final safety analysis report, Amendment 35, 26:47474 (DOCKET-50309-62)

Maine Yankee Atomic Power Station. Final safety analysis report, Amendment 36, 26:47478 (DOCKET-50309-66)

Maine Yankee Atomic Power Station. Seismic qualification of vital instrument and electrical equipment, 26:47477 (DOCKET-50309-65)

MAINE YANKEE ATOMIC POWER CO., WISCASSET.

Maine Yankee Atomic Power Station. Seismic design review, 26:47476 (DOCKET-50309-64)

MASSACHUSETTS GENERAL HOSPITAL, BOSTON. JOHN COLLINS WARREN LABS.

An investigation on the mechanism of protein and nucleic acid synthesis: a background study related to the biological effects of radiation. Decennial progress report, September 1, 1960-August 31, 1972, 26:47612 (COO-3078-6)

Early steps in protein synthesis and their regulation: a background study related to the biological effects of radiation. Progress report, September 1, 1971-August 31, 1972, 26:47611 (COO-3078-5)

MASSACHUSETTS INST. OF TECH., CAMBRIDGE. CENTER FOR SPACE RESEARCH.

OGO-1 and OGO-3 MIT plasma experiments S4903. Final project report, 26:46265 (N-72-18715)

MASSACHUSETTS INST. OF TECH., CAMBRIDGE. DEPT. OF METEOROLOGY.

Correlation of D-region absorption and the 10 mb winds and temperature during the IQSY, 26:46264 (COO-2195-5)

Planetary transport processes in the upper atmosphere. Annual progress report, 26:45503 (COO-2195-6)

MASSACHUSETTS INST. OF TECH., CAMBRIDGE (USA). RALPH M. PARSONS LAB. FOR WATER RESOURCES AND HYDRODYNAMICS.

Mathematical model for the prediction of unsteady salinity intrusion in estuaries, 26:47620 (MIT-144)

MAX-PLANCK-INSTITUT FUER METALLFORSCHUNG, STUTTGART (WEST GERMANY).

Application of high-voltage electron microscopy to the study of voids and other point-defect agglomerates, 26:46057 (CONF-710601-, pp 550-62)

MAX-PLANCK-INSTITUT FUER PLASMAPHYSIK, GARCHING (WEST GERMANY).

Power production costs and possible applications of short-period combustion MHD generators, 26:47170 (IPP-IV-32)

Simultaneous spectroscopic determination of electron density and temperature as a proof of relaxation effects in a streaming argon--potassium--plasma, 26:47116 (IPP-IV-12)

METROPOLITAN EDISON CO., READING, PA.

Three Mile Island Nuclear Station, Unit 1. Final safety analysis report, Amendment 27, 26:47461 (DOCKET-50289-63)

MICHIGAN TECHNOLOGICAL UNIV., HOUGHTON. DEPT. OF METALLURGICAL ENGINEERING.

Structure and properties of solid solutions. Technical progress report, September 1, 1971--June 15, 1972, 26:45991 (COO-916-25)

MICHIGAN UNIV., ANN ARBOR.

200-GeV proton--proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)

MICHIGAN UNIV., ANN ARBOR. HARRISON M. RANDALL LAB. OF PHYSICS.

Time variation of the fine structure constant and spectral lines from quasi-stellar objects, 26:46254 (COO-1112-51)

MILLSTONE POINT CO., HARTFORD, CONN.

Millstone Point Nuclear Power Station, Unit 1. Abnormal occurrence AO-72-17, 26:47407 (DOCKET-50245-139)

Millstone Point Nuclear Power Station, Unit 1. Abnormal occurrence AO-72-18, 26:47360 (DOCKET-50245-143)

Millstone Point Nuclear Power Station, Unit 1. Provisional operating license DPR-21, proposed change 5, 26:47408 (DOCKET-50245-140)

Millstone Point Nuclear Power Station, Unit 1. Request temporary waiver of technical specification 3.5.F.6, 26:47411 (DOCKET-50245-145)

Millstone Point Nuclear Power Station, Unit 1. Semiannual report, January 1, 1972--June 30, 1972, 26:47361 (DOCKET-50245-146)

Millstone Point Nuclear Power Station, Unit 1. Technical specifications, proposed change 6, 26:47409 (DOCKET-50245-141)

Millstone Point Nuclear Power Station, Unit 1. Technical specifications, Proposed Change 8, 26:47410 (DOCKET-50245-144)

Millstone Point Nuclear Power Station, Unit 2. Final safety analysis report, Volume 1, 26:47508 (DOCKET-50336-36)

Millstone Point Nuclear Power Station, Unit 2. Final safety analysis report, Volume 2, 26:47509 (DOCKET-50336-37)

Millstone Point Nuclear Power Station, Unit 2. Operating license application, 26:47507 (DOCKET-50336-35)

MOUND LAB., MIAMISBURG, OHIO.

Carbon-13: a selected abstract bibliography, 26:45286 (MLM-1913)

Documentation of DOT-SP-5915 for 55/30 and 83/55 drum combination packaging, 26:46114 (MLM-1911)

MSA RESEARCH CORP., EVANS CITY, PA.

Entrainment moisture separators for fine (1-10 μ) water--air--steam service: their performance, development, and status. Final report, 26:45440 (MSAR-71-45)

N

NAPLES UNIV. (ITALY). ISTITUTO DI FISICA TEORICA.

Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, CLEVELAND, OHIO. LEWIS RESEARCH CENTER.

Cyclotron production of ^{123}I : an evaluation of the nuclear reactions which produce this isotope, 26:45392 (N-72-15068)

Development of dispersion-strengthened Ni--Cr--ThO₂ alloys for the space-shuttle thermal protection system, 26:45934 (N-72-18585)

Effect of two types of helium circulators on the performance of a subsonic nuclear-powered airplane, 26:45425 (N-72-16724)

Particle--hole states in ^{120}Sn , 26:46943 (N-72-18705)

Shock and vibration tests of uranium mononitride fuel pellets for a space-power nuclear reactor, 26:45933 (N-72-15609)

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, GREENBELT, MD. GODDARD SPACE FLIGHT CENTER.

Charged-particle motion in the vicinity of a neutral plane, 26:46522 (N-72-17763)

Light-ion trough, 26:46301 (N-72-18362)

Lorentz-invariant formulation of Cherenkov radiation by tachyons, 26:46704 (N-72-19766)

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, LANGLEY STATION, VA. LANGLEY RESEARCH CENTER.

Effects of 22-MeV protons on single-junction and silicon-controlled rectifiers, 26:45661 (N-72-17158)

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, MOFFETT FIELD, CALIF. AMES RESEARCH CENTER.

Radiation dosimetry for the Gemini program, 26:47203 (N-

72-18797)

NATIONAL INST. FOR METALLURGY, JOHANNESBURG (SOUTH AFRICA).

Pressed fused disc: a technique for preparation of ore samples to be analyzed for uranium by X-ray fluorescence, 26:45262 (NIM-1427)

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, ROCKVILLE, MD. ENVIRONMENTAL RESEARCH LABS.

MILROW/CANNKIN seismic effects, 26:45479 (NVO-746-122)

NAVAL RESEARCH LAB., WASHINGTON, D. C.

Production of vacancy condensates by laser bombardment, 26:46059 (CONF-710601-, pp 613-29)

NEBRASKA PUBLIC POWER DISTRICT, COLUMBUS.

Cooper Nuclear Station. Environmental report,

Supplement 2, 26:47465 (DOCKET-50298-43)

Cooper Nuclear Station. Environmental report,

Supplement 3, 26:47468 (DOCKET-50298-48)

Cooper Nuclear Station. Final safety analysis report,

Amendment 13, 26:47466 (DOCKET-50298-46)

Cooper Nuclear Station. Final safety analysis report,

Amendment 11, 26:47467 (DOCKET-50298-47)

NEVADA OPERATIONS OFFICE (AEC), LAS VEGAS.

Project WAGON WHEEL: preliminary site evaluation report, 26:45571 (NVO-55)

Reports available in Plowshare Open File, 26:45477 (NVO-86(Rev.1))

NEVADA UNIV., RENO (USA). MAX C. FLEISCHMANN COLL. OF AGRICULTURE.

Ecology of Hot Creek Valley: vegetation and soil response to underground detonation. Annual progress report FY 1971, 26:45478 (NVO-409-1)

NEW ENGLAND DEACONESS HOSPITAL, BOSTON, MASS. CANCER RESEARCH INST.

Radiation carcinogenesis. Progress report VI, May 1, 1971--April 30, 1972, 26:45814 (COO-3017-43)

NEW YORK OPERATIONS OFFICE (AEC), N. Y. HEALTH AND SAFETY LAB.

Fallout Program. Quarterly summary report, March 1,

1972--June 1, 1972, 26:45511 (HASL-257)

Fallout program. Quarterly summary report, March 1--

June 1, 1972. Appendix, 26:45512 (HASL-257(App.))

NEW YORK UNIV., N. Y.

Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)

NIAGARA MOHAWK POWER CORP., LYCOMING, N. Y.

Nine Mile Point Nuclear Station, Unit 1. Malfunctions which occurred during the periodic testing of the core spray systems, 26:47349 (DOCKET-50220-100)

NIAGARA MOHAWK POWER CORP., SYRACUSE, N. Y.

Nine Mile Point Nuclear Station, Unit 1. Delay in

delivery of nitrogen gas, 26:47350 (DOCKET-50220-101)

Nine Mile Point Nuclear Station, Unit 1. Environmental

report, operating license stage, conversion to full-

time operating license, 26:47401 (DOCKET-50220-98)

Nine Mile Point Nuclear Station, Unit 1. Results of

test on main steam line isolation valves, 26:47575

(DOCKET-50220-102)

Nine Mile Point Nuclear Station, Unit 1. Technical

specifications, proposed changes, 26:47351 (DOCKET-

50220-103)

NORTH AMERICAN ROCKWELL CORP., THOUSAND OAKS, CALIF. SCIENCE CENTER.

Nucleation of voids in materials during high-energy irradiation, 26:46064 (CONF-710601-, pp 825-40)

NORTHERN STATES POWER CO., MINNEAPOLIS, MINN.

Monticello Nuclear Generating Plant, Unit 1. Fast start failure of the high pressure coolant injection system,

26:47334 (DOCKET-50263-143)

Monticello Nuclear Generating Plant, Unit 1. High

Pressure Coolant Injection System failure, 26:47368

(DOCKET-50263-139)

Monticello Nuclear Generating Plant, Unit 1. HPCI

turbine check valve leakage, 26:47333 (DOCKET-50263-

142)

Monticello Nuclear Generating Plant, Unit 1.

Malfunction of the "D" steamline safety/relief valve

and inoperability of bellows leakage monitoring

systems, 26:47369 (DOCKET-50263-140)

Monticello Nuclear Generating Plant, Unit 1.

Supplemental letter to the May 25, 1972 report of low

APRM gain setting, 26:47323 (DOCKET-50263-141)

Monticello Nuclear Generating Plant, Unit 1. Technical

specifications, Change Request 4, 26:47429 (DOCKET-

50263-138)

Monticello Nuclear Generating Plant, Unit 1. Turbine

trip at 100 percent power and subsequent events,

26:47367 (DOCKET-50263-137)

Prairie Island Nuclear Generating Plant, Unit 1.

Application for AEC Class 104b Operating License,

Amendment 13, 26:47450 (DOCKET-50282-59)

Prairie Island Nuclear Generating Plant, Units 1 and 2.

Final safety analysis report, Amendment 19, 26:47451

(DOCKET-50282-67)

Prairie Island Nuclear Generating Plant, Units 1 and 2.

Final safety analysis report, Amendment 20, 26:47453

(DOCKET-50282-69)

NUCLEAR FUEL SERVICES, INC., ROCKVILLE, MD.

West Valley Reprocessing Plant. Low level waste

disposal, 26:45400 (DOCKET-50201-127)

West Valley Reprocessing Plant. Technical

specifications, request for approval of revisions,

26:45399 (DOCKET-50201-126)

NUCLEAR FUEL SERVICES, INC., WEST VALLEY, N. Y.

West Valley Reprocessing Plant. Quarterly report for



OAK RIDGE NATIONAL LAB., TENN.

150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)
Analytical applications of the J integral, 26:45990 (CONF-720806-2)
Burial of radioactive waste in salt, 26:46100 (CONF-720903-4)
Chemistry of HTGR, 26:45288 (TID-26106)
Compilation of crystal growers and crystal growth projects, 26:45973 (ORNL-RMIC-12)
Computer simulation of atomic displacement cascades in metals, 26:46061 (CONF-710601-, pp 739-56)
Continuous or semicontinuous leacher for leaching soluble core material from sheared spent nuclear fuel tubes, 26:45397 (CONF-720903-9)
Dependence of radiation effects on the primary recoil energy, 26:46051 (CONF-710601-, pp 397-429)
Design of high-temperature liquid metal systems, 26:45437 (CONF-720904-1)
Design studies of a molten-salt reactor demonstration plant, 26:47305 (ORNL-TM-3832)
Effect of irradiation temperature on size of defect aggregates in copper, 26:46060 (CONF-710601-, pp 672-83)
Execution time translate subroutines, 26:47621 (ORNL-TM-3810)
Fabrication of ^{252}Cf neutron sources at Oak Ridge National Laboratory, 26:45714 (CONF-720902-2)
Heat transfer performance of short twisted tapes, 26:47312 (CONF-720802-1)
Increased effective half-life of intratracheal administered $^{51}\text{Cr}_2\text{O}_3$ by respiratory infection, 26:45910 (CONF-720505-2)
KWIC Index for numerical algebra, 26:47287 (ORNL-4778)
LMFBR fuel cycle studies progress report for April 1970, No. 14, 26:45409 (ORNL-TM-2996)
LMFBR fuel cycle studies progress report for August 1970, No. 18, 26:45413 (ORNL-TM-3127)
LMFBR fuel cycle studies progress report for December 1970, No. 22, 26:45417 (ORNL-TM-3281)
LMFBR fuel cycle studies progress report for February 1970, No. 12, 26:45407 (ORNL-TM-2918)
LMFBR fuel cycle studies progress report for January 1970, No. 11, 26:45406 (ORNL-TM-2871)
LMFBR fuel cycle studies progress report for July 1970, No. 17, 26:45412 (ORNL-TM-3095)
LMFBR fuel cycle studies progress report for June 1970, No. 16, 26:45411 (ORNL-TM-3072)
LMFBR fuel cycle studies progress report for March 1970, No. 13, 26:45408 (ORNL-TM-2949)
LMFBR fuel cycle studies progress report for May 1970, No. 15, 26:45410 (ORNL-TM-3018)
LMFBR fuel cycle studies progress report for November 1970, No. 21, 26:45416 (ORNL-TM-3250)
LMFBR fuel cycle studies progress report for October 1970, No. 20, 26:45415 (ORNL-TM-3217)
LMFBR fuel cycle studies progress report for September 1970, No. 19, 26:45414 (ORNL-TM-3180)
Low-energy electron attachment to molecules at high densities, 26:46364 (ORNL-TM-3839)
ORNL isotopic power fuels quarterly report for period ending March 31, 1972, 26:45703 (ORNL-4796)
Pathophysiology and diagnosis of internal contamination, 26:45888 (CONF-720520-1)
Photoelectron chemical shifts in inorganic compounds, 26:46340 (ORNL-TM-3820)
Postirradiation mechanical properties of Types 304 and 304 + 0.15 percent Ti stainless steel, 26:46068 (ORNL-TM-3733)
Review of the cross sections of the isotopes ^6Li and ^7Li as tabulated on the ENDF/B Version III files, 26:46876 (ORNL-TM-3859)
SDT1. Iron broomstick experiment, 26:47220 (ORNL-TM-3867)
SDT2. Oxygen broomstick experiment, 26:47221 (ORNL-TM-3868)
SDT3. Nitrogen broomstick experiment, 26:47222 (ORNL-TM-3869)
SDT4. Sodium broomstick experiment, 26:47223 (ORNL-TM-3870)
SDT5. Stainless-steel broomstick experiment, 26:47224 (ORNL-TM-3871)
OAK RIDGE OPERATIONS OFFICE (AEC), TENN.
Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)
OAKLAND UNIV., ROCHESTER, MICH. DEPT. OF BIOLOGICAL SCIENCES.
Cellular proliferation and regeneration following tissue damage, 26:45813 (COO-1427-41)
OFFICE OF INFORMATION SERVICES (AEC), WASHINGTON, D.C.
Selected measurement methods for plutonium and uranium in the nuclear fuel cycle. Second edition, 26:45183 (TID-7029(2nd Ed.))
OREGON STATE UNIV., CORVALLIS. DEPT. OF GENERAL SCIENCE.
Radionuclide cycling in natural populations of amphibians. Annual progress report, June 16, 1971--June 15, 1972, 26:45913 (RLO-2227-T-8-3)

9 CORPORATES

POWER REACTOR DEVELOPMENT CO., DETROIT,

OREGON UNIV., EUGENE.

Tests of high-energy models by two-particle analyses, 26:46587 (RLO-2230-T-4-1)



PACIFIC GAS AND ELECTRIC CO., SAN FRANCISCO, CALIF.

Diablo Canyon Nuclear Power Plant, Units 1 and 2. Environmental report, Supplement 2, Volume 2, 26:47447 (DOCKET-50275-23)
Diablo Canyon Nuclear Power Plant, Units 1 and 2. Environmental report, Supplement 2, Volume 1, 26:47446 (DOCKET-50275-22)
Humboldt Bay Power Plant, Unit 3. Technical specifications, proposed change 40, 26:47392 (DOCKET-50133-89)
Humboldt Bay Power Plant, Unit 3. Transmission tripped out of service by forest fire, 26:47342 (DOCKET-50133-91)
Mendocino Power Plant, Units 1 and 2. Preliminary safety analysis report, Amendment 7, 26:47559 (DOCKET-50398-11)
Mendocino Power Plant, Units 1 and 2. Preliminary safety analysis report, Amendment 9, 26:47560 (DOCKET-50398-15)
Mendocino Power Plant, Units 1 and 2. Preliminary safety analysis report, Amendment 10, 26:47561 (DOCKET-50398-16)
PADUCAH GASEOUS DIFFUSION PLANT, KY.
Conditions for the removal of chromate from cooling water, 26:47619 (KY-642)
PARIS UNIV., ORSAY (FRANCE).
Capture of low-energy rare gas ions by metallic targets, 26:46335 (FRNC-TH-198)
PARIS UNIV., ORSAY (FRANCE). LABORATOIRE DE L'ACCELERATEUR LINEAIRE.
Hadronic production by e^+e^- collisions at 990 MeV with Orsay storage ring, 26:46429 (CONF-710816-15)
Measurement of radiative neutral decay modes of ω^0 and ϕ^0 mesons with Orsay storage ring, 26:46472 (CONF-710816-12)
 $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
 $\pi^+\pi^-$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
PARIS UNIV., ORSAY (FRANCE). LABORATOIRE DE PHYSIQUE THEORIQUE ET HAUTES ENERGIES.
Conference on high energy phenomenology, Meribel-les-Allues, France, 3/71, 26:46590
PENNSYLVANIA POWER AND LIGHT CO., ALLENTOWN.
Susquehanna Steam Electric Station, Units 1 and 2. Environmental report, revised July 1972, Vol. 2, 26:47546 (DOCKET-50387-20)
Susquehanna Steam Electric Station, Units 1 and 2. Environmental report, revised July 1972, Vol. 1, 26:47545 (DOCKET-50387-19)
Susquehanna Steam Electric Station, Units 1 and 2. Preliminary safety analysis report, amendment 5, revision, 26:47547 (DOCKET-50387-21)
Susquehanna Steam Electric Station, Units 1 and 2. Preliminary safety analysis report, Amendment 8, 26:47548 (DOCKET-50387-22)
Susquehanna Steam Electric Station, Units 1 and 2. Preliminary safety analysis report, Amendment 9, 26:47549 (DOCKET-50387-23)
PENNSYLVANIA STATE UNIV., UNIVERSITY PARK. DEPT. OF MATERIALS SCIENCE.
Research on graphite. Final report for work completed during the period 1954 to 1972, 26:46078 (NYO-1710-106)
PHILADELPHIA ELECTRIC CO., PA.
Limerick Generating Station, Units 1 and 2. Environmental report, construction permit stage (revised), Supplement 1, 26:47520 (DOCKET-50352-27)
Peach Bottom Atomic Power Station, Units 2 and 3. Amendment 17, substitution of a new Exhibit "U", 26:47375 (DOCKET-50277-51)
Peach Bottom Atomic Power Station, Units 2 and 3. Environmental report, Operating License stage, Supplement 5, 26:47448 (DOCKET-50277-52)
PITTSBURGH UNIV., PA. DEPT. OF PHYSICS.
Study of radiation induced defects in metals. Final report, January 15, 1966--January 14, 1972, 26:46066 (COO-3431-1)
PORTLAND GENERAL ELECTRIC CO., OREG.
Trojan Nuclear Plant. Initial decision on environmental review, 26:47515 (DOCKET-50344-29)
Trojan Nuclear Plant. Supplemental environmental report, Amendment 3, 26:47516 (DOCKET-50344-30)
POWER AUTHORITY OF THE STATE OF NEW YORK.
James A. Fitzpatrick Nuclear Power Plant. Environmental report, Supplement 3, 26:47502 (DOCKET-50333-51)
James A. Fitzpatrick Nuclear Power Plant. Final facility description and safety analysis report, Supplement 9, 26:47501 (DOCKET-50333-50)
POWER REACTOR DEVELOPMENT CO., DETROIT, MICH.
Enrico Fermi Atomic Power Plant, Unit 1. Answer to show cause order and amendment to request for extension of expiration date of provisional operating license,

26:47390 (DOCKET-50016-163)
 Enrico Fermi Atomic Power Plant, Unit 1. Execution of periodic test activity delayed by piping installation, 26:47327 (DOCKET-50016-160)
 Enrico Fermi Atomic Power Plant, Unit 1. Failure to measure radioactivity of sodium at specified times, 26:47340 (DOCKET-50016-165)
 Enrico Fermi Atomic Power Plant, Unit 1. Report for April 1972, 26:47339 (DOCKET-50016-164)
 PRINCETON UNIV., N. J.
 Annual progress report, June 1, 1971—May 31, 1972, 26:45313 (COO-3029-1)
 PUBLIC SERVICE ELECTRIC AND GAS CO., NEWARK, N. J.
 Salem Nuclear Generating Station, Units 1 and 2. Environmental report, Amendment 2, 26:47445 (DOCKET-50272-33)
 Salem Nuclear Generating Station, Units 1 and 2. Final safety analysis report, Amendment 13, 26:47442 (DOCKET-50272-30)
 Salem Nuclear Generating Station, Units 1 and 2. Final safety analysis report, Volume 7, 26:47444 (DOCKET-50272-32)
 Salem Nuclear Generating Station, Units 1 and 2. Final safety analysis report, Volume 6, 26:47443 (DOCKET-50272-31)
 Salem Nuclear Generating Station, Units 1 and 2. Final safety analysis report, Amendment 12, 26:47441 (DOCKET-50272-29)
 PUERTO RICO WATER RESOURCES AUTHORITY, SAN JUAN.
 Aguirre Nuclear Plant, Unit 1. Preliminary facility analysis report, Amendment 18, 26:47537 (DOCKET-50376-34)
 PURDUE UNIV., LAFAYETTE, IND.
 Observation of the $M^0(940)$ and δ^{963} , 26:46475 (IS-2886)
 PURDUE UNIV., LAFAYETTE, IND. DEPT. OF CHEMISTRY.
 Technical progress report on fundamental studies of separation processes, 26:45302 (COO-1222-55)
 PURDUE UNIV., LAFAYETTE, IND. DEPT. OF PHYSICS.
 Final states with three charged particles and a visible Λ from K-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)

Q

QUEENS COLL., NEW YORK.
 Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)

R

RCA LABS., PRINCETON, N. J.
 High-peak-power, high-efficiency microcircuit avalanche diode oscillators, 26:47624 (SC-CR-723142)
 RENSSELAER POLYTECHNIC INST., TROY, N. Y. DEPT. OF PHYSICS.
 MeV total neutron cross sections of ^{238}U , 26:47013 (COO-3058-23)
 RHODE ISLAND UNIV., KINGSTON (USA). INST. OF ENVIRONMENTAL BIOLOGY.
 Effects of ionizing radiation on salamander orientation. Progress report, 26:45815 (COO-3486-1)
 ROCHESTER GAS AND ELECTRIC CORP., N. Y.
 Robert Emmett Ginna Nuclear Power Plant, Unit 1. Loss of rod control logic system power supply, 26:47321 (DOCKET-50244-109)
 Robert Emmett Ginna Nuclear Power Plant, Unit 1. Monthly report of activity measurements, 26:47406 (DOCKET-50244-107)
 Robert Emmett Ginna Nuclear Power Plant, Unit 1. Pressurizer safety valves, 26:47576 (DOCKET-50244-108)
 ROCHESTER UNIV., N. Y.
 Effect of rat tumor allografts on the distribution of passively transferred alloantibodies studied by hemagglutination and isotopic tracer techniques, 26:45820 (UR-3490-121)
 Fundamental studies in isotope chemistry. Progress report, July 1, 1971—June 30, 1972, 26:47614 (COO-3498-2)
 ROCHESTER UNIV., N. Y. DEPT. OF RADIATION BIOLOGY AND BIOPHYSICS.
 Bacteriophage T4 inhibits colicin E2-induced degradation of *Escherichia coli* deoxyribonucleic acid, 26:47639 (UR-3055-2)
 Biological properties of certain compounds containing the C—F bond, 26:47637 (UR-49-1508)
 Chronic low-level NO_2 exposure and the mucociliary clearance mechanism in the rat trachea in vivo, 26:47644 (UR-3490-85)
 Distribution of an ^{125}I -labeled transplantation antibody in syngeneic and allogeneic rats, 26:47649 (UR-3490-147)
 Effect of a polythiol synthetic resin on the absorption and excretion of methylmercury in experimental animals, 26:47643 (UR-3490-80)
 Effect of varying doses of cortisol on the incorporation of ^3H -thymidine into rat liver DNA and on the synthesis of five plasma proteins in the isolated

perfused regenerating rat liver, 26:47646 (UR-3490-131)
 Effects of chloramphenicol and cycloheximide on the cell growth cycle of mouse leukemic cells (L5178Y), 26:47647 (UR-3490-132)
 Experimental inhalation of metallic silver, 26:45915 (UR-3490-115)
 Ionizing radiation-induced thyroid carcinogenesis in the dog, 26:45819 (UR-3490-63)
 Passive cation flux changes in the human erythrocyte suspended in various anionic media, 26:47638 (UR-3055-1)
 Pathogenesis of radionuclide-induced tumors, 26:45895 (UR-3490-102)
 Pathophysiologic effects of microwave radiation with special reference to the thyroid—pituitary axis, 26:47648 (UR-3490-137)
 Protein synthesis and the cell cycle of L5178Y cells, 26:47640 (UR-3490-18)
 Retention of ^{210}Pb in beagle dogs, 26:45916 (UR-3490-159)
 Spectral studies of collagen, 26:47642 (UR-3490-47)
 ROCHESTER UNIV., N. Y. SCHOOL OF MEDICINE AND DENTISTRY.
 Effects of actinomycin D on G₁ protein synthesis in L5178Y cells, 26:47650 (UR-3490-153)
 Fluorine metabolite concentrations and renal clearances following methoxyflurane anesthesia, 26:47645 (UR-3490-107)
 Procedure for chronic intravenous catheterization in the rabbit, 26:47641 (UR-3490-21)
 Protein synthesis during the cell cycle of L5178Y cells, 26:47651 (UR-3490-154)
 RUTHERFORD HIGH ENERGY LAB., CHILTON (ENGLAND).
 Plastic threshold Cherenkov detectors with wavelength shifter and quencher, 26:45646 (RPP/H-89)
 Regge cuts from inclusive finite mass sum rules, 26:46651 (RPP/T-18)

S

SACRAMENTO MUNICIPAL UTILITY DISTRICT, CALIF.
 Rancho Seco Nuclear Generating Station, Unit 1. Environmental report, Amendment 2, 26:47479 (DOCKET-50312-41)
 Rancho Seco Nuclear Generating Station, Unit 1. Final safety analysis report, Amendment 12, 26:47480 (DOCKET-50312-42)
 Rancho Seco Nuclear Generating Station, Unit 1. Final safety analysis report, Amendment 13, 26:47482 (DOCKET-50312-45)
 Rancho Seco Nuclear Generating Station, Unit 1. Report on minor imperfections found in pipe welds, 26:47481 (DOCKET-50312-43)
 Rancho Seco Nuclear Generating Station, Unit 1. Weld defects found in steam generator lateral support, 26:47483 (DOCKET-50312-46)
 SAHA INST. OF NUCLEAR PHYSICS, CALCUTTA (INDIA).
 Fission barrier heights, energy release, and fragment energy partition processes from the viewpoints of nuclear structural models, 26:47046 (INDC-7-U, pp 145-60)
 SANDIA LABS., ALBUQUERQUE, N. MEX.
 Anode temperatures and ion energy distributions during a vacuum arc discharge, 26:47625 (SC-DC-722267)
 Cannikins: effects on archaeological sites, 26:45485 (SC-RR-720359)
 Carrier trapping effects in irradiated pn junctions, 26:45664 (SC-DC-721237)
 Device degradation from the effects of nuclear radiation on passivation materials, 26:45662 (SC-DC-721232)
 Fused salts. A bibliography. Volume I. Citations listing. Volume II. Subject/title index. Volume III. Authors index, 26:45287 (SC-R-722769)
 High-peak-power, high-efficiency microcircuit avalanche diode oscillators, 26:47624 (SC-CR-723142)
 Large scale characterization tests of gelled slurry blasting agents, 26:45484 (SC-DC-722296)
 Oxidation and etching of 80 percent Si—20 percent Ge, 26:45935 (SC-RR-710884)
 Radiation effects in microwave bipolar transistors, 26:45663 (SC-DC-721233)
 Rapid annealing of frequency change in crystal resonators following pulsed x-irradiation, 26:45665 (SC-DC-721240)
 Semiconductor device degradation by high amplitude current pulses, 26:45666 (SC-RR-720515)
 SANDIA LABS., LIVERMORE, CALIF.
 An introduction to cobordism theory, 26:47627 (SCL-RR-720270)
 Helium migration in metals, 26:46050 (CONF-710601-, pp 230-47)
 SCIENTIFIC RESEARCH ATOMIC REACTOR INST., MELEKESS (USSR).
 Determination of the concentration of gamma-emitting isotopes in extended sources without sampling, 26:45264 (SRARI-P-123)
 Realization of synchronous rotation of some rotors suspended in a magnetic field, 26:45441 (SRARI-P-120)
 SOUTH CAROLINA ELECTRIC AND GAS CO., COLUMBIA.
 Virgil C. Summer Nuclear Station, Unit 1. Preliminary safety analysis report, Amendment 10, 26:47553 (DOCKET-50395-22)
 Virgil C. Summer Nuclear Station, Unit 1. Preliminary

safety analysis report, Amendment 12, 26:47554 (DOCKET-50395-24)

Virgil C. Summer Nuclear Station, Unit 1. Preliminary safety analysis report, Amendment 13, 26:47555 (DOCKET-50395-25)

SOUTHERN CALIFORNIA EDISON CO., ROSEMead, CALIF.
San Onofre Nuclear Generating Station, Units 2 and 3.
Application for a construction permit and a class 104(b) license, Amendment 15, 26:47522 (DOCKET-50361-25)

San Onofre Nuclear Generating Station, Units 2 and 3.
Preliminary safety analysis report, Amendment 14, 26:47523 (DOCKET-50361-26)

STAATLICHE ZENTRALE FUER STRAHLENSCHUTZ, BERLIN (EAST GERMANY).
Some problems of the discharge of radioactive gases and aerosols from nuclear facilities into the atmosphere, 26:46102 (ORNL-tr-2369)

STANFORD UNIV., CALIF. DEPT. OF CIVIL ENGINEERING.
Design of waterway connecting Orinoco and Rio Negro Rivers in Federal Territory of Amazonas, Venezuela, 26:45475 (NP-19361)

STATE UNIV. OF NEW YORK, ALBANY.
Radiation-induced voids in metals. Proceedings of the 1971 international conference held at Albany, New York, June 9--11, 1971, 26:46045 (CONF-710601)

STATE UNIV. OF NEW YORK, STONY BROOK. DEPT. OF PHYSICS.
Importance of the quadrupole pairing field in the J/ψ π/π^0 vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)

STATE UNIV. OF NEW YORK, STONY BROOK. INST. FOR THEORETICAL PHYSICS.
Importance of the quadrupole pairing field in the J/ψ π/π^0 vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)

STICHTING VOOR FUNDAMENTEEL ONDERZOEK DER MATERIE, JUTPHAAS (NETHERLANDS). INSTITUUT VOOR PLASMA-FYSICA.
Production and measurement of intense ion beams and beam-produced plasma in a cusped magnetic field, 26:47153 (NP-19364)

SUSSEX UNIV., BRIGHTON (ENGLAND).
Is the pion form factor determined by electroproduction experiments, 26:46513 (CONF-710816-17)

T

TATA INST. OF FUNDAMENTAL RESEARCH, BOMBAY (INDIA).
Did superheavy nuclei exist in our solar system, 26:47016 (INDC-7-U, pp 23-5)
Distribution of the total fission width, 26:47045 (INDC-7-U, pp 81-6)

TECHNICAL INFORMATION CENTER (AEC), OAK RIDGE, TENN.
Bibliographies of atomic energy literature. Issued or in progress during the period April--June 1972, 26:47628 (TID-3767)

TECHNISCHE HOCHSCHULE, AACHEN (WEST GERMANY).
Absolute measurement of the $^9\text{Be}(n,2n)$ cross section in the energy region of 2.37 to 3.24 MeV, 26:46874 (JUL-834-RG)

TECHNISCHE UNIVERSITAET, BRUNSWICK (WEST GERMANY).
Computer-code for solution of three-dimensional-multigroup-equations, based on improved synthesis-method, 26:47594 (BNBW-FBK-72-6)

TECHNISCHE UNIVERSITAET, HANOVER (WEST GERMANY).
Comparison of the background spectra in eight german whole-body counters and a model structure with layered shield, 26:45590 (ORNL-tr-2532)

TEKNISKE HOEJSKOLE, LYNGBY (DENMARK).
Positron lifetimes in water and ice, and in frozen aqueous solutions, 26:46362 (RISO-254)

TELEDYNE ISOTOPES, LAS VEGAS, NEV.
Distribution of radioactivity in and near the Rainier rubble chimney, 26:45483 (NVO-1229-180)
Long term release of radioactivity from Rainier melt-glass, 26:45482 (NVO-1229-177)

TENNESSEE UNIV., KNOXVILLE.
Low-energy electron attachment to molecules at high densities, 26:46364 (ORNL-TM-3839)
Measurement of liquid droplet emissions from cooling towers and process stacks, 26:45574 (CONF-720619-1)
Photoelectron chemical shifts in inorganic compounds, 26:46340 (ORNL-TM-3820)
Studies on nutrition of the preplacental embryos of the mammal. Sixth year progress report, 26:47622 (ORO-4245-4)

TENNESSEE VALLEY AUTHORITY, CHATTANOOGA.
Browns Ferry Nuclear Power Station, Units 1, 2, and 3.
Final safety analysis report, Amendment 35, 26:47425 (DOCKET-50259-57)
Browns Ferry Nuclear Power Station, Units 1, 2, and 3.
Final safety analysis report, Amendment 36, 26:47426 (DOCKET-50259-58)
Browns Ferry Nuclear Power Station, Units 1, 2, and 3.
Final safety analysis report, Amendment 37, 26:47427 (DOCKET-50259-59)
Browns Ferry Nuclear Power Station, Units 1, 2, and 3.
Final safety analysis report, Amendment 38, 26:47428 (DOCKET-50259-60)
Watts Bar Nuclear Plant, Units 1 and 2. Application for license, Amendment 12, 26:47552 (DOCKET-50390-19)
Watts Bar Nuclear Plant, Units 1 and 2. Preliminary safety analysis report, Amendment 9, 26:47550 (DOCKET-

50390-17)
Watts Bar Nuclear Plant, Units 1 and 2. Preliminary safety analysis report, Amendment 11, 26:47551 (DOCKET-50390-18)

TEXAS INSTRUMENTS INC., DALLAS. SCIENCE SERVICES DIV.
Measurement of radioactive fallout by field total absorption gamma ray spectroscopy. Report No. 3, 26:45527 (TID-26103)

TEXAS UNIV., AUSTIN. CENTER FOR PARTICLE THEORY.
Leptonic decay of vector mesons in an algebraic model for hadrons, 26:46525 (ORO-3992-80)
On a possible repulsive interaction in universal gravitation, 26:46749 (ORO-3992-94)

TOKYO UNIV. (JAPAN). DEPT. OF RADIATION BIOPHYSICS.
Effects of actinomycin D on G₁ protein synthesis in L5178Y cells, 26:47650 (UR-3490-153)
Protein synthesis during the cell cycle of L5178Y cells, 26:47651 (UR-3490-154)

TOLEDO EDISON CO., OHIO.
Davis-Besse Nuclear Power Station. Environmental report, Supplement, Amendment 1, 26:47518 (DOCKET-50346-29)

TOULOUSE UNIV. (FRANCE). CENTRE DE PHYSIQUE ATOMIQUE ET NUCLEAIRE.
Noise in Au--nSi diodes operating in the avalanche region in the presence of charges injected by radiation, 26:46560 (FRNC-CONF-29)

U

UNITED KINGDOM ATOMIC ENERGY AUTHORITY, DOUNREAY (SCOTLAND). REACTOR GROUP.
Void formation in cladding and structural materials irradiated in DFR, 26:46047 (CONF-710601-, pp 125-41)

UNITED POWER ASSOCIATION, ELK RIVER, MINN. (USA).
Elk River Reactor. Operating Authorization DPRA-3, Amendment 1, 26:47336 (DOCKET-1151-57)

UNITED STATES-EURATOM FAST REACTOR EXCHANGE PROGRAM.
Influence of fission neutron spectra on integral nuclear quantities of fast reactors, 26:47585 (EURFNR-1008)

UTAH UNIV., SALT LAKE CITY (USA). DEPT. OF SURGERY.
Biomedical engineering support. Progress report, 26:45700 (COO-2155-4)

UTAH UNIV., SALT LAKE CITY. DIV. OF MATERIALS SCIENCE AND ENGINEERING.
Preparation and characterization of biocompatible polymers by radiation grafting and radioisotope technique. Technical progress report, June 15, 1971--May 15, 1972, 26:45312 (COO-2147-2)

V

VAN POOLLEN (H.K.) AND ASSOCIATES, LITTLETON, COLO. (USA).
Projected reservoir performance. Rio Blanco Project, 26:45476 (NVO-38-33)

VERMONT YANKEE NUCLEAR POWER CORP., RUTLAND.
Vermont Yankee Nuclear Power Station. Changes in steam tunnel ventilation system, 26:47438 (DOCKET-50271-93)
Vermont Yankee Nuclear Power Station. Final environmental statement, 26:47435 (DOCKET-50271-88)
Vermont Yankee Nuclear Power Station. License application, Amendment 31, 26:47374 (DOCKET-50271-90)
Vermont Yankee Nuclear Power Station. Operating license DPR-28, Appendix B, 26:47437 (DOCKET-50271-92)
Vermont Yankee Nuclear Power Station. Technical specifications, proposed change, 26:47436 (DOCKET-50271-91)
Vermont Yankee Nuclear Power Station. Technical specifications, Proposed Change 1, Supplement 1, 26:47440 (DOCKET-50271-95)

VIRGINIA ELECTRIC AND POWER CO., RICHMOND.
North Anna Power Station, Units 1 and 2. Environmental considerations, 26:47511 (DOCKET-50338-48)
North Anna Power Station, Units 1 and 2. Request for permission to resume work on transmission line, 26:47512 (DOCKET-50338-49)
North Anna Power Station, Units 1, 2, 3, and 4. Environmental report supplement, Revision 2, 26:47510 (DOCKET-50338-47)
North Anna Power Station, Units 3 and 4. Preliminary safety analysis report, Amendment 5, 26:47566 (DOCKET-50404-18)
North Anna Power Station, Units 3 and 4. Preliminary safety analysis report, Amendment 6, 26:47567 (DOCKET-50404-19)
Surry Power Station, Unit 1. Authorization request to drain reactor coolant system water and lower water level in reactor vessel, 26:47328 (DOCKET-50280-78)
Surry Power Station, Units 1 and 2. Topical report on General Electric Vertical Induction Motors, inside containment recirculation spray pump motors, addendum, 26:47449 (DOCKET-50280-77)

VIRGINIA UNIV., CHARLOTTESVILLE. RESEARCH LABS. FOR THE ENGINEERING SCIENCES.
Interaction of heavy keV ions with matter. Progress report, 26:46365 (ORO-2911-14)

W**WASHINGTON PUBLIC POWER SUPPLY SYSTEM, RICHLAND.**

Hanford No. 2 Nuclear Power Plant. Preliminary safety analysis report, Amendment 7, 26:47556 (DOCKET-50397-19)

Hanford No. 2 Nuclear Power Plant. Preliminary safety analysis report, Amendment 8, 26:47557 (DOCKET-50397-20)

Hanford No. 2 Nuclear Power Plant. Preliminary safety analysis report, Amendment 9, 26:47558 (DOCKET-50397-21)

WASHINGTON UNIV., SEATTLE. LAB. OF RADIATION ECOLOGY.

Decline of ^{65}Zn in marine mussels following the shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)

Natural ^{210}Pb and ^{210}Po in a marine environment, 26:45544 (RLO-2225-T-14-4)

WEIZMANN INST. OF SCIENCE, REHOVOTH (ISRAEL).

Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)

WESTINGHOUSE ELECTRIC CORP., MADISON, PA. ADVANCED REACTORS DIV.

Void formation in quenched stainless steel, 26:45987 (CONF-710601-, pp 605-12)

West Valley Reprocessing Plant. Letters of May 23, June 26, 27, 30, and July 5, 1972, with respect to modifications in the plant, 26:46065 (CONF-710601-, pp 841-52)

WESTINGHOUSE ELECTRIC CORP., PITTSBURGH, PA.

Discussion on fracture mechanics concepts, 26:45996 (WCAP-7841)

WISCONSIN ELECTRIC POWER CO., MILWAUKEE.

Point Beach Nuclear Plant, Unit 1. Steam generator cladding repair, 26:47373 (DOCKET-50266-75)

Point Beach Nuclear Plant, Units 1 and 2. Boric acid concentration in "A" boric acid storage tank below technical specifications limit, 26:47372 (DOCKET-50266-73)

WISCONSIN PUBLIC SERVICE CORP., GREEN BAY.

Kewaunee Nuclear Power Plant. Application for construction permit and operating license, Amendment 20, 26:47380 (DOCKET-50305-46)

Kewaunee Nuclear Power Plant. Final safety analysis report, Amendment 19, 26:47470 (DOCKET-50305-44)

WOODS HOLE OCEANOGRAPHIC INSTITUTION, MASS.

Concentrations and distributions of long-lived fallout radionuclides in open ocean sediments, 26:45531 (COO-3563-3)

Plutonium in North Atlantic Ocean organisms; ecological relationships, 26:45541 (NYO-2174-130)

Y**YANKEE ATOMIC ELECTRIC CO., WESTBORO, MASS.**

Yankee Nuclear Power Station. Operation report 138 for month of June 1972, 26:47341 (DOCKET-50029-189)

Yankee Nuclear Power Station. Withdrawal of proposed change 101, 26:47391 (DOCKET-50029-190)

Z**ZENTRALSTELLE FUER ATOMKERNENERGIE-DOKUMENTATION,**

LEOPOLDSHAFEN (WEST GERMANY).

Surface and vacuum physics index, Vol. 7, No. 5, 1972, 26:46336

PERSONAL AUTHOR INDEX

Authors' names are indexed in the form appearing in the abstracted publication. Each entry under an author's name gives the publication title and the NSA volume and abstract numbers. Report numbers, where applicable, are given in parentheses at the end of the entry.

Because of problems involved in the mechanical preparation of this index, all accent marks are omitted. The following changes in spelling have been introduced to reduce some of the confusion which might result from omitting the accents:

Germanic ä	appears as ae
Germanic ö	oe
Germanic ü	ue
Scandinavian ø	oe

A

- AARON, H. B., Free volume as a guide to grain boundary phenomena, 26:46027
- ABABEI, L., Glucose-6-phosphate phosphohydrolase activity in erythrocytes after x-irradiation, 26:45848
- ABBE, J. Ch., Charge states of ^{208}Pb and ^{208}Tl formed by α decay of ^{212}Po and ^{212}Bi , 26:45359
Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in gas phase, 26:45361
Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in liquid phase, 26:45362
- ABDEL-HALIM, A. F., Mean lives of some high energy levels in ^{160}Dy , 26:47000
- ABDINOV, O. B., Intranuclear cascades in light nuclei, 26:46664
- ABDULLAH, T., Nonrelativistic harmonic-oscillator quark model and the ratio of longitudinal to transverse cross-sections in resonance electroproduction, 26:46544
- ABDURAKHMANOV, S. R., Analysis of experimental energies of excited states of even-even deformed nuclei using rotational formulas, 26:47080
- ABELLANAS, Lorenzo, Mathematical characterization of the breaking of hadronic internal symmetry, 26:46682
- ABERNATHEY, Robert M., Rapid mass-spectrometric determination of boron isotopic distribution in boron carbide, 26:45235
- ABIDOVA, M. F., Al-Fe-Mo oxide catalysts. I, 26:45963
- ABLAKELOV, Kh. A., Spin correlation during bremsstrahlung radiation of longitudinally polarized electrons, 26:47052
- ABOU-LEILA, Hussein, Mean lives of some high energy levels in ^{160}Dy , 26:47000
- ABRAHAM, B. M., Absence of a quadratic term in the ^4He excitation spectrum, 26:46799
- ABRAHAM, Guy E., Simultaneous radioimmunoassay of plasma testosterone and dihydrotestosterone, 26:45928
- ABRAMO, M. C., Self-consistent pair correlations and dielectric response in a plasma, 26:47150
- ABRAMOV, R. E., Characteristics of nucleotide of DNA in radiation injury, 26:45856
- ABRAMYAN, E. A., Creep of niobium under conditions of in-leakage of oxygen, 26:46000
- ABT, Helmut A., Rotational velocities of Ap stars, 26:46182
- ABURAI, Tamaru, Production of carrier-free ^{54}Mn , 26:45394
- ABU-SHUMAYS, I. K., Thermal plume data acquisition, documentation, and initial analysis, 26:45579
- ABU-WAKID, Shukri Edward, Positron-hydrogen collisions and positronium formation, 26:46372
- ACHASOV, N. N., $\rho^0 \rightarrow \omega$ interference in reactions $\gamma N \rightarrow \pi^+ \pi^- \Delta$, 26:46527
- ACHE, Hans J., Phase and temperature dependence of positron annihilation in liquid crystals, 26:46363
- ACHIMAN, Y., Possible CPT violation and the $K/\text{sub } L / \rightarrow \mu^+ \mu^-$ decay, 26:46680
- ACKERHALT, Robert E., Annealing reactions in K chromate. III, 26:45364
- ADAMCZYK, B., Partial ionization cross sections of C dioxide by electrons (25 to 600 eV), 26:46384
- ADAMS, Donald W., Shock and vibration tests of uranium mononitride fuel pellets for a space-power nuclear reactor, 26:45933 (N-72-15609)
- ADAMS, F., Instrumental neutron activation analysis of germanium, 26:45218
- ADDA, Y., Report on the CEA program of investigations of radiation-induced cavities in metals: presentation of some results, 26:46046 (CONF-710601-, pp 31-83)
- ADEISHVILI, D. I., Decrease of the off-duty factor in the operation of a linear accelerator, 26:46145
- ADELBERG, Michael L., Numerical aspects of momentum-space scattering equations for peaked potentials, 26:46787
- ADLER, Laszlo, Ultrasonic flaw determination by spectral analysis, 26:45469
- AFANAS'EV, G. N., SU(3) physical basis realization and probabilities of $B(E2)$ -transitions in SU(3) scheme. II, 26:47049 (JINR-P4-6458)
- AFANAS'EVA, T. M., Dynamics of control system incorporating 440 MW(e) reactor, 26:47324
- AGGSON, T. L., Satellite and ground-based observations of a red arc, 26:46275
- AGRAWAL, P. C., Observation of a hard x-ray flare in Cyg X-1, 26:46211
- AHLFELD, C. E., Total neutron cross section of ^{243}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- AHLGREN, Karin, Level structure of ^{187}Os from decay of ^{187}Ir , 26:46984
- AHMAD, Mohammad, Preliminary remarks about large-scale desalination and dual purpose plants, 26:47579
- AHMAD, Z., Inclusive reaction $\mu + \text{nucleon} \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV/c, 26:46452
- AHMED, E. M., Respiratory quotients of irradiated fruits, 26:45729
- AHMED, M. A., Deep inelastic scattering, scaling, and Regge behavior in a Feynman diagram model, 26:46565
- AINGER, F. W., Ferroelectrics in $\text{K}_2\text{O}-\text{SrO}-\text{Nb}_2\text{O}_5$ system, 26:45962
- AIREY, J. R., Quenching of infrared chemiluminescence: rates of energy transfer from $\text{HF}(\nu \text{ less than or equal to } 3)$ to CO_2 and HF , 26:45298
- AIVAZYAN, R. B., Compensation of the drift of charged-particle tracks in wide-gap spark chambers, 26:45652
- AJZENBERG-SELOVE, F., Energy levels of light nuclei $A = 18$ to 20, 26:46882
- AKASOFU, S.-I., Observations of the auroral oval by the Alaskan meridian chain of stations, 26:46268
- AKHIEZER, A. I., Impossibility of introducing "hidden" parameters into quantum mechanics, 26:47285 (ITF-71-101-P)
- AKIHA, Fumimasa, Production of carrier-free ^{54}Mn , 26:45394
- AKIMOV, Yu. K., Time measurements with a germanium detector, 26:45616
- AKIMOV, Yu. S., Investigation of the injection characteristics of symmetrical pin-diodes prepared by the method of ion implantation, 26:45678
- ALABISO, C., Irreducible representations of para-Bose operators, 26:46507
- ALADJEM, Abraham, Metal oxide film structure, 26:45958
- ALAIS, M., Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)
- ALAKAEVA, L. A., Microdeterminations of terbium by a luminescent method with the reagent pyrogallolsulfonic acid, 26:45231
- ALANAKYAN, K. V., Control system for beam formation after single deflection at exit of 300 MeV electron linear accelerator, 26:46131
- ALBERT, G., Recoil effect in quasi-elastic hadron-deuteron scattering, 26:46653
- ALBERT, Ph., General conclusions on the methods of activation analysis, 26:45212
- ALBERT, Philippe, Activation analysis with thermal neutrons, 26:45208
- ALBERTINI, Richard Joseph, Spontaneous and x-ray induced mutation in cultured, diploid human cells, 26:45740
- ALEINIKOV, V. G., Radiometric investigation of the absorption of potential-determining ions from aqueous solutions of non-ionic surface-active substances, 26:45279
- ALEKSANDROV, B. M., X ray fluorescence analysis of heavy element by an α -X coincidence method, 26:45270
- ALEKSANDROV, S. V., Characteristics of nucleotide of DNA in radiation injury, 26:45856

- ALEKSANDROV, Yu. A., Electron density in tungsten crystal according to neutron-electron interaction data, 26:47254
- ALEKSEEV, P. D., Accumulation of F centers in NaCl crystals as function of proton-beam intensity, 26:47246
- ALEDNARD, Roland, Anisotropy in the paramagnetic susceptibility of some rare earth elements, 26:47253
- ALESHIN, K. P., Moessbauer diffractometer having high angular resolution, 26:45683
- ALESHIN, V. P., β, γ transitions and structure of anti-analog states, 26:47050
- ALEXANDER, Edward Michael, Ultrasonic attenuation in pure niobium at low temperatures, 26:46818
- ALEXANDER, Everette E., Method of fabricating a heat pipe, 26:45449
- ALEXANDER, L. G., Design studies of a molten-salt reactor demonstration plant, 26:47305 (ORNL-TM-3832)
- ALEXANDER, T. K., Lifetime of 975-keV, $3/2^+$ state in ^{25}Mg , 26:46908
- ALEXANDROV, N. N., Chromosome aberrations in human bone-marrow cells after a single local gamma-irradiation. I, 26:45794
- ALFANO, F., Doppler-broadened annihilation radiation in metals with a Ge(Li) detector, 26:47268
- ALIKHANOV, S. G., Solitons in a two stream plasma, 26:47202
- ALLARDYCE, B. W., Test of time reversal invariance through detailed balance of the reaction $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$, 26:46881
- ALLAWAY, William H., Vanadium determination in biological materials at nanogram levels by a catalytic method, 26:45203
- ALLEN, E., Hematopoietic and immunological alterations in mice produced by ^{90}Sr -induced marrow hypoplasia, 26:45896
- ALLEN, Ronald J., Radio map of spiral galaxy Maffei 2 at 1415 MHz, 26:46237
- ALLES, W., Tests for possible class of CP invariant solutions to $K/\text{sub } L + \mu$ anti μ problem, 26:46577
- ALLES-BORELLI, V., e^+e^- annihilation into two hadrons in the energy interval 1400 to 2400 MeV, 26:46552
- ALLKOFER, Otto Claus, Radiation hazards to man in space flights, 26:45812
- ALLOIN, D., Continuum of nuclei of galaxies. II, 26:46239
- ALMAZOV, A. V., Dielectric strength of accelerator tubes, 26:46125
- ALSTON-GARNJOST, Margaret, Experimental spectra for 12 GeV/c $K^+p + pK^+\phi$ and $pK^+\omega$, 26:46484
- ALTARELLI, G., Dreili-Hearn sum rule and the lepton magnetic moment in the Weinberg model of weak and electromagnetic interactions, 26:46553
- AMALDI, Ugo Jr., Criteria for selecting the location of nuclear plants, 26:47574
- AMAN, E. G., Homogeneous anisotropic solutions of Einstein equations with cosmological term, 26:46759
- AMAT, G., Exposure measurements and spectrometry in the medium energy x-ray region, 26:47218
- AMBRO, P., New two-channel scanner. II, 26:45607
- AMBROSE, Robert T., Atomic absorption spectroscopy, 26:45251
- AMBROSI, M., Exposure measurements and spectrometry in the medium energy x-ray region, 26:47218
- AMIN, S. R., Estimation of sulphur content in human hair and nail for use in fast neutron dosimetry, 26:45788
- AMIRANASHVILI, A. G., Estimation of the turbulent diffusion coefficient and vertical component of wind velocity according to natural radioactivity distribution, 26:45522
- AMMANN, A. C., Final states with three charged particles and a visible A from K^-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)
- AMOV, B., ^{132}Ba levels excited during decay of ^{132}La and ^{132}Ce , 26:46948
- ANASHKIN, O. P., Removable current lead for power supply in superconducting magnetic systems, 26:45679
- ANDERSEN, B. V., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- ANDERSEN, N., Relative level populations in foil excited He and Li, 26:46337
- ANDERSON, Bryon Don, Study of the $^{16}\text{O}(d, n)$ reaction from 3 to 4 MeV, 26:46899
- ANDERSON, David Nels, Daily variation of the ionospheric F2 equatorial anomaly in the American and Asian sectors, 26:46293
- ANDERSSON, B., Dispersion relations for the vertex function from local commutativity. II, 26:46735
- Dispersion relations for the vertex function from local commutativity. I, 26:46736
- ANDERT, K., Time measurements with a germanium detector, 26:45616
- ANDRADE, J. D., Preparation and characterization of biocompatible polymers by radiation grafting and radioisotope technique. Technical progress report, June 15, 1971-May 15, 1972, 26:45312 (COO-2147-2)
- ANDRASSY, L., Developments in geophysical logging instruments and methods, 26:45560
- ANDREEV, G. G., Elastic scattering of polarized protons by ^{28}Si and ^{40}Ca , 26:46924
- ANDRIEUX, Claude, Semiconductor well logging probe, 26:45658
- ANDRILLAT, Y., Continuum of nuclei of galaxies. II, 26:46239
- ANGERMAN, C. L., Savannah River Laboratory ^{60}Co power and heat sources. Quarterly progress report, January-March 1972, 26:45701 (DP-1300)
- ANNO, Tosinobu, Pair correlation energies as derived from the analysis of semi-empirical values of correlation energies of atoms, 26:46412
- ANON, M. C., Thorium-acetate and thorium-propionate complexes, 26:45373
- AN'SHAKOV, A. S., Erosion of a tungsten cathode in nitrogen, 26:45968
- ANSPAUGH, L. R., Environmental levels of radioactivity in the vicinity of the Lawrence Livermore Laboratory January through December 1971, 26:45513 (UCRL-51242)
- ANTHONY, T. R., Solute segregation and stresses generated around growing voids in metals, 26:45988 (CONF-710601-, pp 630-46)
- ANTIPPA, A. F., One-dimensional causal theory of tachyons, 26:46499
- ANTOINE, J.-P., Rigged Hilbert spaces in quantum field theory, 26:46737
- ANTOKU, Shigetoshi, Death of mice irradiated with pulsed x-rays, 26:45866
- Dosimetry of pulsed x-rays and electrons from a linear accelerator, 26:45627
- ANTONIOU, Antoine, Comparative effects of ion bombardment and partial pressures on sheet polyethylene, 26:46081
- ANTROPOV, A. E., Elastic and inelastic scattering of 6-MeV protons by even isotopes of Cr, 26:46922
- APPEL, H., First forbidden beta decay of ^{209}Pb , 26:47007
- APPEL, J. A., Charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\gamma$, 26:46482
- Measurement of the charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\omega$, 26:46483
- APPLEBY, W. K., Fluence and temperature dependence of void formation in highly irradiated stainless steels, 26:46048 (CONF-710601-, pp 156-73)
- ARAI, S., Asymmetry for π^+ photoproduction from polarized protons between 300 and 900 MeV at 90°, 26:46444
- ARATA, Yoji, Nuclear magnetic relaxation study of formic acid. Isotope effect and ^{13}C relaxation, 26:45297
- ARDEBERG, A., Extended van Wijk sequence in Large Magellanic Cloud, 26:46238
- ARDELL, A. J., Contrast effects at grain boundary ledges in 304 austenitic stainless steel, 26:47630 (UCLA-34-P-177-11)
- Reactor steels studies. Quarterly progress report, October-December 1971, 26:45994 (UCLA-34-P-177-10)
- ARIFOV, P. U., Use of quasi-classical approximation for solution of system of Hartree-Fock equation, 26:47288
- ARIFOV, U. A., Properties of ruby laser radiation under action of ^{60}Co gamma radiation, 26:47227
- Secondary emission and structural properties of solid bodies, 26:47245
- ARIMA, Akira, Subtraction and positron scanning with the dual headed whole body scanner, 26:45593
- ARISAKA, Nakateru, SU(3) unitary representations, 26:46675
- ARISTOV, E. A., Determination of total coefficient of internal gamma-radiation conversion of ^{139}La , 26:46947
- ARIZUMI, Tetsuya, Recombination domain instability in high-energy electron-irradiated germanium, 26:47230
- ARKHIPOV, A. A., Estimate of the quasipotential phase shift, 26:46505
- ARLT, R., Isobars from the spallation reaction for nuclear spectroscopy, 26:46985
- Study of ^{17}Re decay, 26:46976 (JINR-P6-6492)
- ARLT, R. D., X-ray spectrometer having cooled Si(Li)-detector, 26:45644
- ARMJJO, J. S., Deuteron-generated void formation in austenitic stainless steels, 26:46053 (CONF-710601-, pp 479-98)
- ARMSTRONG, T. P., Comparison of measurements of the charge spectrum of solar cosmic rays from nuclear emulsions and the Explorer 35 solid-state detector, 26:46330
- ARNAL, Thierry, Equipment for concentrating radioactive solutions, 26:46111
- ARNAUD, A., Computerized control of an accelerator and the associated experiments, 26:46123
- ARNDT, Renate, Special techniques for the metallographic examination of hard metals, 26:45955
- ARNETT, W. David, Hydrostatic oxygen burning in stars. II. Oxygen burning at balanced power, 26:46178
- ARSEN'EV, D. A., Isomeric transition $8^- \rightarrow 6^+$ in ^{130}Ba nucleus, 26:46949
- ARTEM'EV, A. N., Moessbauer diffractometer having high angular resolution, 26:45683
- ARTEM'EV, O. I., Decay of isomers $^{126}\text{I}^{26}\text{I}^{130}\text{Sb}$ and quasiparticle levels $^{126}\text{I}^{26}\text{I}^{130}\text{Te}$ with large spin, 26:46944
- ARTSYBASHEV, V. A., Determination of density and effective atomic number of rocks and ores by gamma-gamma method in the presence of an intermediate zone, 26:45566
- ARUMI, F. N., Modified ordinary wave, 26:47192
- ARUTYUNYAN, F. R., Detector for high energy particles based on the principle of using particle radiation in a laminated medium, 26:45648
- ASARS, Juris A., Television type nuclear radiation camera system, 26:45595
- ASBRIDGE, J. R., Velocity and flux dependence of the solar-wind helium abundance, 26:46328

ASCOLI, Giulio, Study of A_3 in $\pi^+p \rightarrow \pi^+\pi^-p$, 26:46473 (COO-1195-239)

ASCUITTO, R. J., Spectroscopy with two-nucleon transfer reactions in deformed nuclei. II, 26:47062
Spectroscopy with two-nucleon transfer reactions in deformed nuclei. I, 26:47061

ASHER, Colin J., Improved method for determination of ^{35}S in plant material using oxygen flask combustion and liquid scintillation counting, 26:45266

ASHLEY, Frantz W., Age-sex specific Achilles tendon reflex time in Japanese: a study of atomic bomb survivors, Hiroshima, 26:45795

ASHMORE, J. F., Method of gauge-invariant regularization, 26:46719

ASHRAF, Mohammad, Histopathology of irradiated yellow mealworm adult and larval midguts, 26:45875

ASKEW, W. C., Synthesis, identification, and purification of C_6 and C_7 fluorocarbons from radiolysis. Final report, 26:45316 (ORO-3909-3)

ASMUSSEN, Keith E., Space-energy-dependent physics problems of plutonium fuels in HTGRs, 26:47313

ASTIER, A., K_2^0 decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)

ASTOLFI, A., Actual value method evaluation of the effects of fuel costs on the cost of power produced by a nuclear reactor for the case of batch refueling and for the case of continuous refueling, 26:47315 (CISE-R-178)

ATEN, A. H. W. Jr., Titanium and calcium content of meteorites, 26:45215

ATKINS, Patrick R., Stable isotopes of carbon as indicators of air pollution levels, 26:45245

ATOMURA, Masakatu, Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)

AU, Chi-kwan E., Retarded van der Waals potential between pairs of spinless atoms, 26:46357

AUBEAU, Raymond, Determination of trace quantities of volatile fluoride in uranium hexafluoride using an infrared spectrophotometer, 26:45238

AUBERT, C., Changes in the localization of acetylcholinesterase in neuromuscular junction after ^{60}Co irradiation, 26:45828

AUBIER, M., Exciter of Type III bursts and coronal temperature, 26:46221

AUDRETSCH, J., Neutrinos in spherically symmetric gravitational fields, 26:46772

AUGUSTSON, R. H., Nondestructive assay of power reactor fuel assemblies, 26:45269

AUSTEN, C. E., Pressed fused disc: a technique for preparation of ore samples to be analyzed for uranium by X-ray fluorescence, 26:45262 (NIM-1427)

AVAKYAN, K. M., Compensation of the drift of charged-particle tracks in wide-gap spark chambers, 26:45652

AVITZUR, Betzalel, Forming of composite for superconducting rod. Technical progress report, 26:45972 (COO-3311-1)

AVRAMOV, S. R., $\text{SU}(3)$ physical basis realization and probabilities of $\text{B}(E2)$ -transitions in $\text{SU}(3)$ scheme. II, 26:47049 (JINR-P4-6458)

AVVAKUMOV, E. G., Substance purification by zone melting in a rotating container, 26:45276

AXFORD, R. A., Burnup and power shift effects in large fast breeder reactors, 26:47306

AYPAR, A., Electron spin resonance in single crystal yttria stabilized zirconia, 26:47252

AZNAURYAN, I. G., Estimation of intermediate energy contribution to the dispersion sum rules and rotation of polarization in pion-nucleon scattering at high energies, 26:46617 (IFVE-STF-71-15)

AZUMA, Toshio, Fast-slow coincidence circuit systems for $\text{Ge}(\text{Li})$ detector, 26:45639

B

BAACKE, J., Partial wave relations from fixed-u dispersion relations, 26:46610

BABABEV, Z. R., Total cross-sections for reaction $\pi^+p \rightarrow \pi^0n$ at high energies, 26:46618 (IFVE-STF-71-71)

BABAYAN, Kh. P., Angular distribution and energy spectrum of cosmic-ray single neutrons, 26:46153

BABBITT, Donald G., Probabilistic interpretation of the quantum scattering cross section, 26:46777

BABIC, E., Superconductivity and localized spin fluctuations in concentrated aluminum-3d-transition-metal alloys, 26:46838

BACH, H., Angular dependence of erosion rate and of specific energy losses of 5.6 keV noble gas ions on glasses, 26:47238

BACIU, G., Betatron bremsstrahlung for physical research, 26:46148

BACK, B. B., Subbarrier fission resonances in Th isotopes, 26:47033

BACKE, H., Test of quantum-electrodynamical corrections in muonic atoms, 26:46708

X-rays and nuclear γ -rays in muonic thallium, 26:47031

BACON, M. E., Time-resolved diagnostic method for hydrogen plasmas, 26:47127

BADAEVA, T. A., Surfaces of solidus and liquidus in system $\text{Th}-\text{Zr}-\text{U}$, 26:46003

BADALYAN, A. M., Bound states of few-nucleon systems, 26:47088

BAER, Rudolf L., Acneform changes in areas of cobalt

irradiation, 26:45793

BAETZNER, K., Separation of $\sigma/\text{sub S/}$ and $\sigma/\text{sub T/}$ in the region of the $\Delta(1236)$ resonance and determination of the magnetic dipole transition form factor, 26:46431 (NP-19256)

BAGLEY, K. Q., Void formation in cladding and structural materials irradiated in DFR, 26:46047 (CONF-710601-, pp 125-41)

BAGLIANO, G., Measurement of ^{137}Cs and ^{90}Sr in soil, 26:45263 (RT/PROT-72-11)

BAGLIN, C., Comparison of anti $\pi^+p \rightarrow \pi^+p(K^+K^-)$ annihilations and $\pi^+p(K^+p)$ backward elastic scattering with crossing symmetry predictions, 26:46463

BAGROV, V. G., Motion of Dirac particles possessing anomalous magnetic moment in central symmetric electric field, 26:46547

BAJEK, Walter A., Radioactive radiation-type fluid interface detection system, 26:45659

BAKER, James F., Effects of gamma radiation on the spectral sensitivity of the noctuid moth, *Heliothis Zea*, 26:45873 (TID-26078)

BAKSHI, P., Linear Stark effect due to resonant interactions of static and dynamic fields, 26:46361

BALACHANDRA, J., Passivity of Type 304 stainless steel: effect of plastic deformation, 26:45967

BALAKIN, V. E., ϕ -meson mass determination, 26:46443

BALANESCU, M., Reinforced concrete vessels for nuclear power reactors: present status and future trends, 26:47311

BALDUCCI, G., Thermodynamics of rare-earth-carbon systems. IV. The lutetium-carbon system, 26:45961

BALE, William F., Distribution of an ^{125}I -labeled transplantation antibody in syngeneic and allogeneic rats, 26:47649 (UR-3490-147)

BALES, Barney L., Role of fluoride ion in irradiated aqueous systems, 26:45345

BALICK, Bruce, Fine structure in H II regions, 26:46198

BALL, J. S., Self-consistent diffraction in the multiperipheral model, 26:46602

BALLA, D., Introduction to low temperature physics, 26:46841

BALLESTRERO, A., General properties of correlation functions for inclusive processes, 26:46502

BALLET, Paul, Method of extracting fuel from nuclear fuel elements, 26:45422

BALLIN, D., Electromagnetic divergences in weak processes, 26:46582

BALLUFFI, R. W., Direct observation of grain boundary dislocation climb in ion-irradiated gold bicrystals, 26:47241

Void formation in quenched or irradiated metals, 26:46058 (CONF-710601-, pp 563-604)

BALON, Z. P., Personal photodosimetry in radiodiagnosis and radiation therapy rooms, 26:45629

BANDELLONI, G., Equal-time and equal light-cone commutator between the dilatation charge and the field in massive $\lambda\phi^4$ -theory, 26:46743

BANDL, Rolf-Eckart, Measure of ratio of capture-to-fission cross section for ^{235}U and ^{239}Pu between 8 and 60 keV neutron energy, 26:47024 (KFK-1563)

BANDO, Hiroharu, Alpha-cluster structure and weak coupling feature in light sd-shell nuclei, 26:47083

BANDYOPADHYAY, P., Parton model and the concept of duality in strong interactions, 26:46599

BANERJEE, S. N., ρ - ρ couplings and t-channel helicity conservation, 26:46638

BANIFATOV, A. E., X-ray spectrometer having cooled $\text{Si}(\text{Li})$ -detector, 26:45644

BANNER, M., Experimental study of the decays $K/\text{sub S L/} \rightarrow \gamma\gamma$, 26:46434

BANYAI, Eva, Radio-frequency alcoholic chlorobenzene dosimeter, 26:45632

BARAKAT, M. F., Chemical effects associated with the thermal neutron irradiation of solid acetanilide (antifibrin), 26:45358

BARANOV, A. A., Cosmological term, screening of gravitation, and hypothesis of negative masses, 26:46761

BARASHENKOV, V. S., Account of nuclear boundary diffusivity in the intranuclear cascade model, 26:47047 (JINR-P2-6503)

Analysis of inelastic collisions of particles with nuclei at very high energies, 26:46649 (JINR-P2-6395)

Change in the mechanism of inelastic interactions between particles and nuclei in the energy region I approximately equal to 5 GeV, 26:47038 (BNL-tr-487)

Cross section of pion-nucleon interactions, 26:46478 (ORNL-tr-2544)

Intranuclear cascades in light nuclei, 26:46664

BARAT, I., Developments in geophysical logging instruments and methods, 26:45560

BARBANEL, Yu. A., Identification of an AmCl_3 -octahedral complex from $\text{Am}(\text{III})$ absorption spectrum in ethanol solutions of HCl , 26:45376

BARBARO-GALTIERI, Angela, Experimental spectra for 12 GeV/c $K^+p \rightarrow pK^+\phi$ and $pK^+\omega$, 26:46484

BARBASHOV, B. M., Corrections to eikonal representation of amplitude in scalar field theory, 26:46699 (JINR-P2-6394)

- BARBER, P., Regulation of water--steam circuit of nuclear power station, 26:47308
- BARBERGER, M., MASURCA: study of plutonium--graphite and uranium--graphite assemblies, 26:47581 (CEA-N-1483)
- BARBIERI, G., PDE: digital program for calculation of thermal-hydrodynamic condition in circuit of two phase mixture water and steam, 26:45458 (CISE-R-245)
- BARGER, V., Nature of absorption, 26:46623
Phenomenology of total cross sections and forward scattering at high energy, 26:46633
- BARLETT, R. H., X-ray-induced photoconductivity in dielectric films, 26:45668 (UCRL-73974)
- BARLOUTAUD, R., Experimental survey of the reactions anti $\text{KN} + \gamma$ at medium and high energy, 26:46489
- BARNA, J., Hygienic investigation on consumptibility of irradiated complete rat feed, 26:45899
- BARNEDUD, D., Rotational levels with high spins in ^{171}Lu , 26:46980
Two branchings in rotational band $1/2^-$ [541] in ^{171}Lu , 26:46987
- BARNES, P. D., Measurement of the magnetic moment of the antiproton, 26:46440
Nuclear γ rays associated with stopped kaons, 26:46940
- BARON, M. J., Chatanika, Alaska, auroral-zone incoherent-scatter facility, 26:46283
- BARRY, J. F., Some measurements of the electrical conductivity induced in polymers by pulsed x-ray irradiation, 26:47225 (AWRE-D-31/72)
- BARTHELEMY, M. C., Relativistic scattering of particles with spin, 26:46793
- BARTINE, David Elliott, Low-energy electron transport by the method of discrete ordinates, 26:47210
- BARTOLI, C., Calculation and measurement of the characteristic flow parameters of a stationary supersonic plasma flow in a magnetic de Laval nozzle, 26:47123
- BARUA, A. K., Inversion from the energy dependence of total cross sections, 26:46387
- BASILE, M., Decay mode $\phi \rightarrow \text{e}^+ \gamma$ and measurement of branching ratio, 26:46435
- BASILE, R., Heavy ion nuclear reactions, 26:47056
- BASS, H. E., Acoustic measurements of rotational energy transfer, 26:46376
- BASU, S., Large scale periodic disturbances in the ionosphere, 26:46290
- BATES, H. F., Chatanika, Alaska, auroral-zone incoherent-scatter facility, 26:46283
- BATES, J. Lambert, High-temperature electrical conductivity of CeO_2 sub 2-x/y, 26:47247 (BNWL-1511)
- BATISHCHEV, A. I., Characteristics of nucleotide of DNA in radiation injury, 26:45856
- BAUD, Gilbert, Hexavalent W bronzes containing Th in interstitials: $\text{Th}/\text{sub } x/4/\text{sup IV}/(\text{B}/\text{sub } x/\text{sup V}/\text{W}/\text{sub } 1-x/\text{sup VI})\text{O}_3$, phases (B = Nb, Ta), 26:45953
- BAUDRY, A., Rotational levels with high spins in ^{171}Lu , 26:46980
Two branchings in rotational band $1/2^-$ [541] in ^{171}Lu , 26:46987
- BAUER, Arthur A., Nitride fuels: properties and potentials, 26:45960
- BAUER, H.-J., Acoustic measurements of rotational energy transfer, 26:46376
- BAUER, Josef, Dysprosium borocarbides, 26:45952
- BAUER, Walter, Helium migration in metals, 26:46050 (CONF-710601-, pp 230-47)
- BAY, Z., Measurement of an optical frequency and the speed of light, 26:46733
- BAYAR, B., Study of ^{179}Re decay, 26:46976 (JINR-P6-6492)
- BAYUKOV, Yu. D., Reaction (π^-, π^0) on ^6Li and ^{12}C , 26:46869
- BAZ, A. I., K-harmonics method to calculate nuclear properties, 26:47091
- BAZHENOV, V. A., Determination of total coefficient of internal gamma-radiation conversion of ^{139}La , 26:46947
- BAZIN, Roger, Method for the manufacture of fissile oxides and the apparatus developed, 26:45940
- BEACH, L. A., Production of carrier-free ^{197}m , ^{197}Hg with a cyclotron, 26:45395
- BEAMAN, D. R., Accuracy of quantitative electron probe microanalysis with energy dispersive spectrometers, 26:45237
- BEAN, Laurence R., Radiation control and monitoring system, 26:45603
- BEATTIE, A. G., Acoustic velocity measurements in the metal hydrides, ScH_3 , YH_3 , and ErH_3 , 26:45950
- BEATTY, Ronald L., Pulsed method for impregnation of graphite, 26:46088
- BEATTY III, Marvin E., Effects of 22-MeV protons on single-junction and silicon-controlled rectifiers, 26:45661 (N-72-17158)
- BECHER, H. J., Crystallization of boron-rich beryllium borides and aluminum beryllium borides from aluminum melts, 26:45936 (UCRL-Trans-10632)
- BECK, U., Separation of $\sigma/\text{sub } S/$ and $\sigma/\text{sub } T/$ in the region of the Δ (1236) resonance and determination of the magnetic dipole transition form factor, 26:46431 (NP-19256)
- BECKEL, Charles L., Effects of screening and polarization on vibration--rotation properties of H_2 , 26:46350
- BECKER, Donald A., Trace analysis for platinum in glasses by neutron activation, 26:45217
- BECKER, Edwin D., Some recent developments in high resolution nuclear magnetic resonance, 26:47278
- BECKER, George L., Factors governing radiographic crack detectability in steel weld specimens, 26:45471
- BECKER, W., Crossing and unitarity in context of Bethe--Salpeter equation, 26:46727
- BECKLIN, E. E., Observations of planetary nebulae at 1.65 to 3.4 microns, 26:46175
- BECKS, K. H., Separation of $\sigma/\text{sub } S/$ and $\sigma/\text{sub } T/$ in the region of the Δ (1236) resonance and determination of the magnetic dipole transition form factor, 26:46431 (NP-19256)
- BEDILOV, M. R., Properties of ruby laser radiation under action of ^{60}Co gamma radiation, 26:47227
- BEER, Lawrence P., Natural and unnatural water temperatures in Zion--Waukegan, Illinois area of southwest Lake Michigan, 26:45580
- BEER, T., Conjugate point phenomena, 26:46323
- BEERS, B., Algebraic approach to the solution of quantum-mechanical problems involving the interaction of a charged particle with electromagnetic fields, 26:46678
- BEESON, James Jackson, Investigation of the radiation polymerization of methyl methacrylate--kaolin clay composites, 26:45327
- BEGEMANN, F., $^{37}\text{Ar}/^{39}\text{Ar}$ activity ratios in meteorites and the spatial constancy of the cosmic radiation, 26:46155
- BEGZHANOV, R. B., ^{135}Ba levels, 26:46945
Collective--single-particle states of deformed odd nuclei, 26:47081
Excitation of ^{99}Tc nucleus, 26:46951
Magnetic properties of ^{153}Eu , ^{155}Eu , and ^{169}Dy , 26:46981
- BEHRENS, H., First forbidden beta decay of ^{209}Pb , 26:47007
- BELEN'KII, V. M., Analysis of experimental energies of excited states of even--even deformed nuclei using rotational formulas, 26:47080
- BELINSKAYA, L. A., Structure, phase composition, and properties of cast complex alloys of Ni--Nb, 26:46035
- BELLE, Jack, Nuclear fuel comprising uranium dioxide in a porous ceramic oxide matrix, 26:45957
- BELON, A. E., HB/N_2^+ 4709-A intensity ratio and vibrational enhancement of N_2^+ first negative bands in proton-excited auroras, 26:46276
- BELOVA, V. M., Temperature dependence of the work function of Ba--W_{110} system, 26:47257
- BELSKY, Joseph L., Age-sex specific Achilles tendon reflex time in Japanese: a study of atomic bomb survivors, Hiroshima, 26:45795
- BELYAEV, S. T., Collective low-lying nuclear states, 26:47092
- BELYAKOV, E. S., Performance logic of spark chamber information read-out by television automation and data link to a computer, 26:45650
- BELYAKOV, V. A., Compensated current-driven magnetic channel in a 240-cm isochronous cyclotron, 26:46128
Magnetic shield for a 240-cm isochronous cyclotron, 26:46129
- BEMENT, A. L. Jr., Irradiation effects on structural materials. I, 26:46071
- BENAK, Steve B., Hard and soft tissue necroses following radiation therapy for oral cancer, 26:45801
- BENAKSAS, D., Hadronic production by e^+e^- collisions at 990 MeV with Orsay storage ring, 26:46429 (CONF-710816-15)
Measurement of radiative neutral decay modes of ω^0 and ϕ^{10} mesons with Orsay storage ring, 26:46472 (CONF-710816-12)
 $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
 $\pi^+\pi^0$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
- BENAYOUN, J. J., Method for solving Faddeev equations, 26:47065
- BENEDICT, Manson, Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)
- BENESTAD, H. Breien, Regulation of granulocyte and macrophage formation in diffusion chamber cultures of mouse haematopoietic cells, 26:45741
- BENGTSOEN, Glenn W., Effect of sodium hydroxide on aluminum foil thickness, 26:47605 (BDX-613-515(Rev.))
- BENIEVA, T. Ya., Determination of the energy of dislocation of an impurity atom by an internal friction method, 26:46023
Influence of impurities and temperature on the deformation of molybdenum during dynamic loading, 26:46043
Orientation dependence of damping of elastic vibrations in tungsten single crystals, 26:46022
- BENJAMIN, R. W., Total neutron cross section of ^{235}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- BENKOE, Gyulane, Studies of the origin of the exogenous oxide inclusions in killed and unkilld cast steels through subsequently activated tracer elements, 26:46008
- BENNETT, B. G., ^{90}Sr in human bone: 1971 results for New York City and San Francisco, 26:45538 (HASL-257, pp 1,51-70)
- BENNETT, K., Galactic gamma radiation between 200 MeV and 10 GeV, 26:46154
- BENNETT, L. G. I., Activation analysis of environmental pollutants, 26:45198
- BENZ, M. G., Mechanism of sintering in cobalt--rare-earth permanent-magnet alloys, 26:45976

BERECZ, Gy., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)

BERESTNEV, V. A., Radiation damage to polymer fiber investigated by small-angle x-ray scattering, 26:45338

BEREZHOI, Yu. A., Diffraction disintegration of deuterons, 26:46923

BEREZIN, Yu. A., Effect of thermal conductivity on the structure and critical parameters of shock waves in a plasma, 26:47200

BERGER, R., French program on production and uses of ^{238}Pu , 26:45706

Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)

BERGMANN, Winfried H., Method and means for charging or discharging superconducting windings, 26:46842

BERGOUNOU, M. A., Processing of solids on fluidized beds, 26:45418

BERK, Robert N., Cholera enteropathy as a cause of diarrhea and death in radiation enteritis and its prevention with cholestyramine, 26:45821

BERMAN, Abraham S., Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)

BERNABEU, J., Quasi-elastic neutrino reaction in deuterium, 26:46581

BERNARD, M., Method of determining mean energy of secondary electrons in electron equilibrium regions using theoretical x-ray spectra (10 to 20 MV) and ionization measurements in different media, 26:47213

BERNARD, V. B., Formation of intermetallic phases in system Ir-Ta-Ni, 26:46010

BERNARDINI, M., e^+e^- annihilation into two hadrons in the energy interval 1400 to 2400 MeV, 26:46552

BERNSHTEIN, L. I., Performance logic of spark chamber information read-out by television automation and data link to a computer, 26:45650

BERNSTEIN, George J., Mixer-settler apparatus, 26:45451

BERONIUS, Per, Nanogram determination of iodide by means of isotope exchange, 26:45257

BERRY, W. T., Proceedings of the second international clean air congress held at Washington, D. C., December 6--11, 1970, 26:45205

BERRY, Warren E., Stress-corrosion cracking of sensitized stainless steel in oxygenated high-temperature water, 26:45966 (BMI-1927)

BERSI, M., Effect of taurine as a protector of human marrow cells against radiation, 26:45734

BERTON, Yves, Method of decontaminating cesium, 26:45384

BERTOZZI, W., Deformation parameters of ^{152}Sm by electron scattering, 26:46993

BERTRAND, M., Isotope effects upon kinetic energy release in metastable ion fragmentations, 26:45301

BES, D. R., Importance of the quadrupole pairing field in the $J/\text{sup } w/ = 0^+$ vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)

BESNIER, Gilbert, Approximations of the space charge field for different beam shapes in the central region of a synchrocyclotron, 26:46140

BESSEL, M. S., Primary spectrum of the eclipsing binary LR Centauri, 26:46191

BESWICK, J. A., Ferroelectrics in $\text{K}_2\text{O--SrO--Nb}_2\text{O}_5$ system, 26:45962

BETTIS, E. S., Design studies of a molten-salt reactor demonstration plant, 26:47305 (ORNL-TM-3832)

BEUERMAN, D. R., Spectrophotometric molybdenum determination with thiolactic acid, 26:45239

BEUMER III, John, Hard and soft tissue necroses following radiation therapy for oral cancer, 26:45801

BEYNON, J. H., Isotope effects upon kinetic energy release in metastable ion fragmentations, 26:45301

BEZRUK, E. T., Borocarbides of Tm, Lu, Tb, Yb, and some of their properties, 26:45964

BHANDARI, Narendra, Did superheavy nuclei exist in our solar system, 26:47016 (INDC-7-U, pp 23-5)

BHATT, B. C., Decay times, intensities, and their temperature variation of luminescence of aqueous and deuterated solutions of europic ion, 26:45295

BHATTACHARYA, D. L., Room temperature internal friction of neutron-irradiated aluminum, 26:46073

BHATTACHARYA, Chitta R., Combined ion exchange-solvent extraction (CIESE) studies of metal ions on ion exchange papers. Synergistic effects and chromatographic separations, 26:45282

BHATTACHARYA, R., Gamma-gamma angular correlation of 171- to 427-keV cascade and the g-factor of the 462-keV, $5/2^+$ level of ^{125}Te , 26:46957

Gyromagnetic ratio of 2083-keV, 4^+ state of ^{140}Ce , 26:46958

BHATTACHARYA, S. N., Radiolysis of aqueous solutions of nickel(II) ethylenediamine tetraacetate, 26:45350

BIANCHERIA, A., West Valley Reprocessing Plant. Letters of May 23, June 26, 27, 30, and July 5, 1972, with respect to modifications in the plant, 26:46065 (CONF-710601-, pp 841-52)

BICEHOUSE, Jack L., Entrainment moisture separators for fine (1-10 μ) water-air-steam service: their performance, development, and status. Final report, 26:45440 (MSAR-71-45)

BIENERT, Walter B., Thermal control and power flattening for radioisotopic thermodynamic power system, 26:45427

BIGEISEN, Jacob, Fundamental studies in isotope chemistry. Progress report, July 1, 1971--June 30, 1972, 26:47614 (COO-3498-2)

Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)

BIGELOW, J. E., Fabrication of ^{252}Cf neutron sources at Oak Ridge National Laboratory, 26:45714 (CONF-720902-2)

BIGGS, William T., Radiation-induced polymerization. I. A pyrotechnic binder, 26:45325

BIJLSMA, S., Absolute and convective instabilities, 26:47161

BIKII, I., $2^+ \rightarrow 2^+$ transition of 879 keV in ^{160}Dy investigated by electron-gamma directional correlation, 26:46991

Effective g/sub s/ factor for odd proton ^{153}Sm nucleus, 26:46994

BILCEA, M., AET on immunologic processes in irradiated rabbits, 26:45824

BILEN*KAYA, S. I., Scale relation for proton form factors, 26:46433

BILEN*KII, S. M., Scale relation for proton form factors, 26:46433

BILLING, A., New two-channel scanner. II, 26:45607

BILLINGSLEY II, Frank P., Coupled multiconfigurational self-consistent-field method for atomic dipole polarizabilities. I. Theory and application to carbon, 26:46419

BILLOIR, Pierre, Interaction of an intense quasi-monochromatic electromagnetic beam with free electrons, 26:46532

BILPUCH, E. G., New results on the fine structure of analog resonances, 26:46914

BIMBOT, R., Transfer reactions induced by ^{86}Kr on ^{232}Th , 26:47035

BINNIE, David Earl, Ultrasonic attenuation in superconducting indium and indium-tin alloys, 26:46816

BINSACK, Joseph H., Explorer 33 and 35 plasma observations of magnetosheath flow, 26:46303

BIRKS, J., Hematopoietic and immunological alterations in mice produced by ^{90}Sr -induced marrow hypoplasia, 26:45896

BITTEL, R., Method of evaluating chemical pollution levels in aquatic media and food chains, 26:45547

BITTER, M., Calculation and measurement of the characteristic flow parameters of a stationary supersonic plasma flow in a magnetic de Laval nozzle, 26:47123

BIYAJIMA, Minoru, Helicity conservation in $n\text{N} + \text{A}_1\text{N}$ scattering, 26:46646

BLACHOT, Jean, Library of data for fission products (third edition), 26:47012 (CEA-N-1526)

BLAKE, J. B., Geomagnetically trapped alpha particles. I. Off-equator particles in the outer zone, 26:46309

BLAMONT, J. E., Geomagnetic effect on the neutral temperature of the F region during the magnetic storm of September 1969, 26:46270

BLANCHET, J., Effect of heat treatments on corrosion of 18/10 stainless steels with low C contents in nitric medium in presence of ferric iron, 26:45970

BLAND, Frederick, Spectrometry and computers used for mapping deposits, 26:45507

BLANDENET, Gerard, Determination of trace quantities of volatile fluoride in uranium hexafluoride using an infrared spectrophotometer, 26:45238

BLANK, J. L., Interaction of the solar wind with a planetary ionosphere, 26:46324 (N-72-18796)

BLANKENSHIP, Charles P., Development of dispersion-strengthened Ni-Cr--ThO₂ alloys for the space-shuttle thermal protection system, 26:45934 (N-72-18585)

BLANN, M., Limits on angular momentum in heavy-ion--compound-nucleus reactions, 26:47072

BLASI, A. A., Casualty group in finite space--time, 26:46724

BLASKOVIC, L., K_2^0 decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)

BLATTMANN, Henri, Composite walls for magnetohydrodynamic apparatus, 26:45426

BLAZEK, M., Before applying the dispersion theory to the photoproduction reactions, 26:46511 (CONF-710816-9)

BLECHERMAN, Sol S., Apparatus for measuring height of a molten metal pool, 26:45655

BLIN, Marie-Francoise, Method for polymerization and/or cross-linking of acrylic polymers or copolymers, 26:45339

BLISS, Henry E., Experience in the licensing of nuclear power plants, 26:47573

BLIXT, Stig, Induced mutations in peas. XXVI, 26:45775

Mutation genetics in Pisum, 26:45776

BLOCK, Robert C., Neutron scattering and capture cross-section measurements, 26:46968

BLOEM, P. J. C., Fission recoil from surface of nuclear fuels by irradiation, 26:45939

BLOMQUIST, J., g-factor of the 12^+ isomeric state in ^{206}Pb , 26:47002

BLOOM, E. E., Postirradiation mechanical properties of Types 304 and 304 + 0.15 percent Ti stainless steel, 26:46068 (ORNL-TM-3733)

BLOSER, Manfred, Absolute measurement of the $^9\text{Be}(n,2n)$ cross section in the energy region of 2.37 to 3.24 MeV, 26:46874 (JUL-834-RG)

BLUE, James W., Cyclotron production of ^{123}I : an evaluation of the nuclear reactions which produce this

- isotope, 26:45392 (N-72-15068)
- BLUET, J. C., Computerized control of an accelerator and the associated experiments, 26:46123
- BLUM, Paul, Packaging for radioactive materials, 26:46115
- BOBROWSKI, Czeslaw, Hydrogen generators for neutron tubes, 26:45942
- BOCHEK, G. L., Angular and energy distributions of protons in (yp) and (ep) reactions on ^{12}C nucleus for 130 MeV excitation energy, 26:46888
- Control system for beam formation after single deflection at exit of 300 MeV electron linear accelerator, 26:46131
- BOCHEV, B., New isotopes of iridium, 26:46977
- New isotopes of tungsten, 26:47001
- BOCHKAREV, V. F., Magnetic compensation in iron--gadolinium and iron--terbium films, 26:47273
- BOCKRIS, J. D'M., Ionic solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
- BOCOLA, W., ESTER program: high-level radioactive waste solidification, 26:45385
- BOCQUET, J. P., On-line measurements of rare-gas fission yields in 14 MeV neutron fission, 26:47030
- BOEHM, A., Leptonic decay of vector mesons in an algebraic model for hadrons, 26:46525 (ORO-3992-80)
- BOELRIJK, N. A. I. M., Comparison of precision in single and double isotope dilution analysis, 26:45225
- BOERBOOM, A. J. H., Partial ionization cross sections of C dioxide by electrons (25 to 600 eV), 26:46384
- BOETTCHER, Alfred, Coated-particle fuel and fertile element for nuclear reactors and method of operating same, 26:45959
- BOGDANOV, G. A., Synthesis and study of lanthanide peroxides, 26:46085
- BOGDANOV, Yu. V., Age of cuprous Ukoka series (Dlekmo--Vitim mountain system) by uranium--lead method, 26:45569
- BOGEMA, Bernard LeRoy, Equilibrium and stability theory of nonneutral plasmas, 26:47167
- BOGOLYUBOV, N. N., Lectures quantum statistics. Volume 2. Quasi-averages, 26:47299
- BOISCHOT, A., Exciter of Type III bursts and coronal temperature, 26:46221
- BOK, Bart J., Birth of stars, 26:46166
- BOLLING, G. F., Free volume as a guide to grain boundary phenomena, 26:46027
- BOLLINGER, Lowell M., Gamma rays from neutron capture in resonances, 26:46969
- BOLLINI, D., e^+e^- annihilation into two hadrons in the energy interval 1400 to 2400 MeV, 26:46552
- BOLTAX, A., West Valley Reprocessing Plant. Letters of May 23, June 26, 27, 30, and July 5, 1972, with respect to modifications in the plant, 26:46065 (CONF-710601-, pp 841-52)
- BONAMY, P., Polarization in charge exchange reactions, 26:46491
- BONCHE, P., Generator-coordinate form factors for transfer reactions between heavy ions, 26:47069
- BONDAREV, V. N., Recording of alpha particles by solid track detectors based on cellulose, 26:45615
- BONHAM, R. A., Second Born zero-angle differential cross sections in high-energy limit for electron-impact excitation of He : the 2^3S -- 1^1S and 2^3P -- 1^1S transitions, 26:46373
- BONNOTE, P. A., Removal of radioactive iodine from ventilation exhaust streams at the AEC Research Establishment, 26:45502
- BONIVER, J., Variations in the bone marrow lymphoid population in the sublethally irradiated germ free mouse, 26:45843
- BONO, R., Experiments on a superposed fields theta pinch, 26:47171
- BOOKER, D. V., Monitor for estimating tritium in urine, 26:45585 (AERE-HP/M-74)
- BOOSTROM, D. R., Vacancy defects in quenched vanadium, 26:46025
- BORDES, G., t-dependence of form factors of nucleons and compositeness of rho-meson, 26:46557
- BOREJKO, Juliusz, Remarks on local metastases to lymph nodes in tumors of the larynx, tonsils, and nasopharynx after x-ray therapy, 26:45782 (IT-54002/4, pp 442-50)
- BORELOWSKI, Z., Special method in S-matrix theory with p^0 momentum representation, 26:46567
- BORGEAUD, P., Polarization in charge exchange reactions, 26:46491
- BORGHINI, M., Polarized proton and deuteron targets and high-energy physics experiments, 26:46436
- BORGONOV, G. M., Selected topics on dynamics of crystal lattices, 26:47617 (IEA-19)
- BORING, John W., Interaction of heavy keV ions with matter. Progress report, 26:46365 (ORO-2911-14)
- BORKOV, F. P., Peculiarities of formation of an increased uranium concentration in oxidizing conditions, 26:45504
- BORMANN, P., Identification of teleseismic events in records of Moxa station, 26:45496
- Study of travel-time residuals with respect to location of teleseismic events from body-wave records at Moxa station, 26:45495
- BORUKHOVICH, A. S., Magnetic susceptibility of cubic niobium and tantalum carbides in the low temperature range, 26:47266
- BOSE, S. K., Room temperature internal friction of neutron-irradiated aluminum, 26:46073
- BOUCHARD, J., Minerve reactor, 26:47589
- BOUCHER, R., French program on production and uses of ^{239}Pu , 26:45706
- Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)
- BOUDET, Roger, Dirac--Hestenes equation and Maxwell equations, 26:46568
- BOULANGER, Gilbert, Ultrasound quality control installation, 26:45472
- BOULDIN, D. P., Validity of DWBA for deuteron stripping. I, 26:47054
- BOULENGER, R., Human faecal ^{226}Ra excretion shortly after a RaSO_4 dust contamination, with alginate treatment, 26:45907
- BOUTAINE, M., Results of enquiry into source requirements of radioisotope generators in ENEA member countries for 1971 to 1975, 26:45705
- BOWEN, V. T., Concentrations and distributions of long-lived fallout radionuclides in open ocean sediments, 26:45531 (COO-3563-3)
- BOWEN, Vaughan T., Plutonium in North Atlantic Ocean organisms; ecological relationships, 26:45541 (NYO-2174-130)
- Shallow-water ^{90}Sr anomaly about the Antilles Arc, 1970, 26:45533
- BOWER, J. R., Idaho Chemical Programs annual technical report, fiscal year 1971, 26:45405 (ICP-1006)
- BOWER, John H., Histopathology of irradiated yellow mealworm adult and larval midguts, 26:45875
- BOWLEY, R. M., Microscopic theory of impurities in liquid helium. II, 26:46803
- BOWMAN, M. R., Photodetachment of electrons from major negative ions in lower D region, 26:46299
- BOX, Harold C., Radical ion saturation in some sulfur compounds x-irradiated at 4.2°K, 26:45348
- BOYADZHYAN, N. G., Angular distribution and energy spectrum of cosmic-ray single neutrons, 26:46153
- BOYD, David M. Jr., Radioactive radiation-type fluid interface detection system, 26:45659
- BOYD, Walter K., Stress-corrosion cracking of sensitized stainless steel in oxygenated high-temperature water, 26:45966 (BMI-1927)
- BOYER, J., Experimental reactor Masurca, 26:47592
- MASURCA: study of plutonium--graphite and uranium--graphite assemblies, 26:47581 (CEA-N-1483)
- BOZRIKOV, P. V., Motion of Dirac particles possessing anomalous magnetic moment in central symmetric electric field, 26:46547
- BOZSOKY, S., Tissue mast cells in human blood, 26:45810
- BRACCINI, P. L., Topics in photoproduction, 26:46438
- BRADFORD, Wesley Lamont, Study on the chemical behavior of zinc in Chesapeake Bay water using Anodic Stripping Voltammetry. Technical report 76, 26:47613 (COO-3292-1)
- BRADLEY, R. H., Explosive interaction of molten metals injected into water, 26:45461
- BRADU, E. B., Statistical treatment of relationship between solar flare-producing regions and SC + SI events, 26:46332
- BRAENDAS, Erkki, Infinite order perturbation theory with applications to atomic and molecular systems, 26:46410
- BRAMMAN, J. I., Void formation in cladding and structural materials irradiated in DFR, 26:46047 (CONF-710601-, pp 125-41)
- BRAMON, A., Nonet structure of $J/\text{sup PC} = 3^-$ mesons, 26:46611
- BRAND, G. F., Cyclotron harmonic waves in a reflex discharge, 26:47184
- BRANDES, H., Determination of the performance of steam separators with the aid of the laws of similitude, 26:45442
- BRAR, S. S., ^{137}Cs in various Chicago foods. (Collection months, January and April 1972), 26:45540 (HASL-257, pp III.45-52)
- BRASSE, F. W., π^0 electroproduction at first resonance, 26:46512 (CONF-710816-16)
- BRAUN, M. P., Structure, phase composition, and properties of cast complex alloys of Ni--Nb, 26:46035
- BRAUN, P. A., Variational bounds for imaginary frequency polarizability and dispersion interaction constant, 26:46415
- BRAUN, S. M., Reaction of Nb--Mo alloys with B, 26:46002
- BREIVIK, H., Regulation of granulocyte and macrophage formation in diffusion chamber cultures of mouse haematopoietic cells, 26:45741
- BRERETON, N. Robin, Reappraisal of $^{40}\text{Ar}/^{39}\text{Ar}$ stepwise degassing technique, 26:45557
- BRESLER, S. E., Lethal and mutagenic effect of ionizing radiations with different LET on extracellular bacteriophages, 26:45746
- BRESSANI, T., New results on ($\pi^+, 2p$) reactions in light nuclei, 26:46870
- BRETTSCHEIDER, Marcel, Numerical simulation of relativistic electrons confined in an axisymmetric mirror field, 26:47147
- BRIDGES, B. A., DNA synthesis in gamma-irradiated recombination deficient strains of *E. coli*, 26:45753
- BRINCA, A. L., Stability of obliquely propagating whistlers, 26:46314

- Whistler side-band growth due to nonlinear wave--particle interaction, 26:46315
- BRINK, L., Nucleon form factors in field theory on homogeneous space of the Poincare groups, 26:46554
- BRINK, Lars, Nucleon form factors in field theory on homogeneous space of Poincare group, 26:46514 (CTH-ITP-71-30)
- BRISCOE, William L., All-metal discharge tube, 26:47178
- BRISSET, R., On-line measurements of rare-gas fission yields in 14 MeV neutron fission, 26:47030
- BRITT, H. C., Subbarrier fission resonances in Th isotopes, 26:47033
- BRITVICH, G. I., Intracascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)
- BROADHEAD, K. G., Fire assay-emission spectrographic determination of iridium, ruthenium, and osmium, 26:45229
- BROCHU, Robert, U_3X_8 type sulfides and selenides of U and Ca, Sr, Ba, Pb, and Eu ($X = S, Se$), 26:45378
- BROGLIA, R. A., Importance of the quadrupole pairing field in the $J/\sup w = 0^+$ vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)
- BROGNIART, Guy, Determination of trace quantities of volatile fluoride in uranium hexafluoride using an infrared spectrophotometer, 26:45238
- BROOKS, Robert Richard, Gamma radiation of some plants and soils from a uraniumiferous area in New Zealand, 26:45529
- BROWN, Clifford C., Methods of nonlinear analysis, 26:47283 (CERN-72-8)
- BROWN, J. W., High-energy electron spikes at high latitudes, 26:46307
- BROWN, Jack E., Canadian National Scientific and Technical Information (STI) system: a progress report. Report No. 4, May 1972, 26:47652
- BROWN, W. D., Semiconductor device degradation by high amplitude current pulses, 26:45666 (SC-RR-720515)
- BRUCATO, R. J., Optical search for Centaurus X-3, 26:46213
- BRUECKNER, Guenter E., Space observations of the solar corona, 26:46233
- BRUGE, G., $^{58}Ni(\alpha, \alpha')$ ^{58}Ni and the nature of high-lying states in ^{58}Ni , 26:46927
- BRUININX, E., Cyclotron laboratory at Technical University of Eindhoven, 26:46146
- BRUN, Therese, Radiorecovery effect of yeast extract on carnation cuttings irradiated with ^{60}Co γ rays, 26:45777
- BRUNA, J., Experimental reactor Masurca, 26:47592
- BRUNE, D., Depth distribution studies of carbon in steel surfaces by means of charged particle activation analysis with an account of heat and diffusion effects in the sample, 26:45466 (AE-451)
- BRUNER, Allen D., Ecology of Hot Creek Valley: vegetation and soil response to underground detonation. Annual progress report FY 1971, 26:45478 (NVO-409-1)
- BRUNIAK, A., Content of Cs in muscles of broilers, hens, and partridges, 26:45548
- BRUNS, L. E., Plutonium--uranium partitioning by a reflux solvent extraction flowsheet, 26:45382 (ARL-SA-130)
- BRYANT, P. E., Changes in sensitivity of cells during exposure to radiation at low dose rate, 26:45754
- BRYMAN, Douglas Andrew, Search for the reaction $\mu^- + Cu \rightarrow e^- + Co$, 26:46941
- BRZEZOWSKI, S., Special method in S-matrix theory with p-momentum representation, 26:46567
- BRZOSKO, J. S., Role of the exit states of a compound nucleus in the emission of high-energy γ -rays, 26:47082
- BUCHALET, C., Discussion on fracture mechanics concepts, 26:45996 (WCAP-7841)
- BUCHSBAUM, Donald J., Distribution of an ^{125}I -labeled transplantation antibody in syngeneic and allogeneic rats, 26:47649 (UR-3490-147)
- BUCHSBAUM, Donald Jay, Effect of rat tumor allografts on the distribution of passively transferred alloantibodies studied by hemagglutination and isotopic tracer techniques, 26:45820 (UR-3490-121)
- BUCKINGHAM, J. Steve, Phosphate solidification of caustic process solution waste, 26:45381 (ARH-SA-120)
- BUCURESCU, D., (α, p) reaction on even nickel isotopes, 26:46934
- BUDINI, P., Possible origin of finite quantum electrodynamics from weak interactions, 26:46722
- Virtual neutrino effects, 26:46570 (IC-72/54)
- BUDNY, R., Partons may have integral charge, 26:46574
- BUDZINSKI, Edwin E., Radical ion saturation in some sulfur compounds x-irradiated at 4.20K, 26:45348
- BUENDE, R., Power production costs and possible applications of short-period combustion MHD generators, 26:47170 (IPP-IV-32)
- BUERGER, G., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- BUGAJSKA, K., Axioms of quantum mechanics, 26:47295
- BUGAJSKI, S., Axioms of quantum mechanics, 26:47295
- BUGRII, A. I., Threshold behavior of DAMA, 26:46774 (ITF-72-44-E)
- BUI AI, Comparative effects of ion bombardment and partial pressures on sheet polyethylene, 26:46081
- BUI-DUY, Q., Is regularization necessary in vacuum polarization calculations, 26:46742
- BUKIN, A. D., ϕ -meson mass determination, 26:46443
- BULLOUGH, R., Observation and theory of the void lattice in molybdenum, 26:46055 (CONF-710601-, pp 522-32)
- Theory of void formation and growth in irradiated materials, 26:46062 (CONF-710601-, pp 769-97)
- BULYAKOVA, N. V., Effect of radiation on regeneration of minced tissues of rats, 26:45827
- BURNISH, R. F., Modern analytical techniques for metals and alloys. Part 2, 26:45187
- BUNUS, Florin T., Determination of inorganic impurities in vinyl chloride by activation analysis, 26:45204
- BURAVIKHIN, V. A., Magnetic compensation in iron--gadolinium and iron--terbium films, 26:47273
- BURBIDGE, G. R., Physical associations between quasi-stellar objects and galaxies, 26:46258
- BURCH, D., Cross sections for ($\nu, \nu n$) reactions on ^{19}F and ^{21}p in the 3 to 3 resonance region, 26:46486
- BURDET, G., Many-body realization of Schroedinger algebra, 26:46673
- BURHOP, E. H. S., Interaction of K^- mesons with nuclei, 26:46487
- BURKE, John C., Plutonium in North Atlantic Ocean organisms; ecological relationships, 26:45541 (NYO-2174-130)
- Shallow-water ^{90}Sr anomaly about the Antilles Arc, 1970, 26:45533
- BURKE, Thomas J., Nuclear fuel comprising uranium dioxide in a porous ceramic oxide matrix, 26:45957
- BURKHARDT, H., Removal of ambiguities from inelastic phase-shift analysis, 26:46640
- BURKHARDT, L. C., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)
- BURNASHOVA, V. V., Formation of intermetallic phases in system Ir--Ta--Ni, 26:46010
- BURNETT, S. C., Feedback stabilization on an $l = 1$ theta-pinch plasma column, 26:47157 (LA-DC-72-707)
- BURNS, Erskine John Thomas, Nuclear energy levels in even isotopes of lead from (α, xn) reactions on separated mercury isotopes, 26:47009
- BURRELL, David C., Trace-metal association in sub-Arctic and Arctic marine environments. Progress report, June 1971--May 1972, 26:47623 (RLO-2229-T-1-18)
- BUSCHBECK, B., K^- -p interactions at 10 GeV/c, 26:46480
- BUSSOLETTI, E., Diurnal effect of cosmic rays during the period 15 October 1965--30 June 1966. I, 26:46159
- Diurnal effect of cosmic rays during the period 15 October 1965--30 June 1966. II, 26:46158
- BUSTAMANTE, N., Transmission of low-energy x-ray through various locally available materials, 26:47205
- BUSWELL, J. T., Application of high-voltage electron microscopy to void studies, 26:46056 (CONF-710601-, pp 533-49)
- BUTLER, G. C., Assessment and control of environmental contamination, 26:45554
- BYKHOVSKII, A. V., Dose to the epithelium of the upper respiratory tract from radon and its daughter products, 26:45861
- BYKOVSKII, Yu. A., Formation of the energy spectrum of the ions of a laser plasma, 26:47154
- BYLOV, Ts., ^{132}Ba levels excited during decay of ^{132}La and ^{132}La , 26:46948
- BYRSKI, E., Thermoluminescent dosimetry system for determining quality, 26:45622
- BYSTROV, V. A., Isobars from the spallation reaction for nuclear spectroscopy, 26:46985
- C**
- CABIBBO, N., Drell--Hearn sum rule and the lepton magnetic moment in the Weinberg model of weak and electromagnetic interactions, 26:46553
- CAHILL, Daniel F., Radiological surveys of Pearl Harbor, Hawaii, and environs, 1966--1968, 26:45534
- CAIRNS, Elton J., Electrochemical power-producing cell, 26:45434
- CALETKA, R., Sorption of polyvalent elements on silica gel. III, 26:45390
- Sorption of polyvalent elements on silica gel. II, 26:45389
- Sorption of polyvalent elements on silica gel. IV, 26:45391
- CALL, Nancy A., Neuropathological evaluation of monkeys exposed to body-alone x-radiation, 26:45850
- CALLAHAN, Thomas Joseph, Passive cation flux changes in the human erythrocyte suspended in various anionic media, 26:47638 (UR-3055-1)
- CALUCCI, G., Hulthen and Schwarzschild potentials in the Klein--Gordon equation, 26:46690
- Some formal developments on the eikonal treatment of production processes, 26:46778
- CALVA-TELLEZ, E., Acoplanarity and radiative corrections in Bhabha scattering, 26:46543
- CALVET, Paul, Hydrogen--deuterium exchange kinetics of cytochrome C between 20 and 38°C, 26:45305
- CAMPARGUE, R., Process and device for the separation of molecules of different masses, 26:45692
- CAMPBELL, A. M., Critical currents in Nb_3Al , 26:46829
- CAMPBELL, E. M., Workloads and use factors in diagnostic x-ray installations, 26:45811
- CAMPBELL, Milton H., Phosphate solidification of caustic process solution waste, 26:45381 (ARH-SA-120)

- CAMPOS VENUTI, Gloria, Criteria for selecting the location of nuclear plants, 26:47574
- CANOSA, Jose, Parallel shooting method for boundary-value problems: application to the neutron transport equation, 26:46859
- CAP, Ferdinand F., Possible production mechanisms of lunar magnetic fields, 26:46262
- CAPDEVILA, C., Application of spectrographic analysis to the radioisotope production control. II, 26:45223 (JEN-232-DQ/I-81)
- CAPELLA, A., K^+ photoproduction, 26:46534
- CAPPS, R. W., Measurement of polarized 10-micron radiation from cool stars with circumstellar shells, 26:46185
- CARDINAL, C. U., /sup 53m/Co: a proton-unstable isomer, 26:46925
- CARENZA, L., Flute modes in a plasma in the presence of nonuniform electric fields, 26:47162
- CARIGNAN, G. R., Angular distribution of electrons elastically scattered from N_2 , 26:46392
- CARMENA, A. O., Effects of radiation and hypoxia on the metabolic fate of erythropoietin, 26:45854
- CARMONY, D. D., Final states with three charged particles and a visible Λ from K^-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)
- CARPENTER, A. B., Analysis of five line hyperfine split Moessbauer absorber spectra, 26:47275
- CARRE, J. C., Minerve reactor, 26:47589
- CARRIVEAU, G. W., Relative level populations in foil excited He and Li, 26:46337
- CARSON, James E., Atmospheric effects of thermal discharges into a large lake, 26:45584
- CARSTEN, Arland L., Remote effects, in time and place, of x irradiation of the visual cortex in the monkey, 26:45858
- CARTAN, F. D., Evaluation of instrumentation for nuclear fuels reprocessing plant input weight measurements, 26:46093 (ICP-1014)
- CARTER, L. A., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- CASARETT, George W., Pathogenesis of radionuclide-induced tumors, 26:45895 (UR-3490-102)
- CASE, Marvin T., Whole-body gamma irradiation of newborn pigs: hematologic changes, 26:45852
- Whole-body gamma irradiation of newborn pigs: pathologic changes, 26:45853
- CASEY, Harold W., Neuropathological evaluation of monkeys exposed to body-alone x-radiation, 26:45850
- CASSANHOL, B., Noise in Au-nSi diodes operating in the avalanche region in the presence of charges injected by radiation, 26:45660 (FRNC-CONF-29)
- CASSIMATIS, Jean, Pneumatic machine for the remote controlled sealing of flasks in a shielded cell, 26:46112
- CAUSSE, R., Modifications induced by gamma irradiations in the ant, 26:45877
- CAVENESS, William, Remote effects, in time and place, of x irradiation of the visual cortex in the monkey, 26:45858
- CAWTHORNE, C., Void formation in cladding and structural materials irradiated in DFR, 26:46047 (CONF-710601-, pp 125-41)
- CECH, Bohuslav, Sintered Mg composites, cladding material of U fuel elements cooled by carbon dioxide. II, 26:46012
- CEJNAR, Frantisek, Modified macroporous styrene-divinylbenzene copolymers as the sorbents with self-fixation ability for the removal of vapors of radiiodine and its compounds. I, 26:45388
- CELASCHI, Emilio, Manufacturing details and material used for manufacture of throttle, control, interceptor, and stop valves for high rating steam turbines of conventional and nuclear type, 26:45455
- CELINSKI, Z. N., Electrical parameters in the Hall-type MHD generator, 26:45431
- CERNY, J., /sup 53m/Co: a proton-unstable isomer, 26:46925
- Beta-delayed proton decay of ^{130} , 26:46879
- Confirmed proton radioactivity of /sup 53m/Co, 26:46926
- CERNY, Joseph, Further results on proton radioactivity of /sup 53m/Co, 26:46919
- CERRAI, E., Reactor chemistry: analytical work, 26:45181 (CISE-R-172)
- CERUTTI, Peter A., Ultraviolet inactivation and miscoding of irradiated R17-RNA in vitro, 26:45728
- CEZAIIRLIYAN, A., Recent developments in thermophysical properties research. Winter Annual Meeting of the American Society of Mechanical Engineers, Washington, D. C., December 1, 1971, 26:45463
- CHADWICK, D., X-ray photoelectron spectra of uranium oxides, 26:45367
- CHAFFEE, Frederic H., Rotational velocities of Ap stars, 26:46182
- CHAFFIN, R. J., Radiation effects in microwave bipolar transistors, 26:45663 (SC-DC-721233)
- CHAIKIN, Paul Michael, Probing many-body effects with superconductivity, 26:46822
- CHAN, Hong-mo, Regge phenomenology of inclusive spectra in the central region, 26:46659
- CHANDRAN, C. R., Effective atomic numbers for photoelectric process in various shells in alloys, 26:47206
- CHANG, C. C., Evidence for an anomaly in the excitation function of the reaction $^2H(p,\gamma)^3He$, 26:46868
- CHANG, James Shih-tsih, Experimental study of wave-wave coupling processes in a beam-plasma system, 26:47193
- CHANG, Shih-Ger, High-temperature spectroscopy and thermodynamics of titanium oxide and vanadium oxide, 26:45932 (LBL-450)
- CHANG, T. H., Partons may have integral charge, 26:46574
- CHANG-LIU, C.-M., Repair of a DNA complex from x-irradiated Chinese hamster cells, 26:45745
- CHANOWITZ, M. S., Canonical anomalies and broken scale invariance, 26:46556
- CHAPMAN, Gary A., Photospheric faculae and the solar oblateness, 26:46224
- CHAPMAN, I. V., Effects of x-irradiation in vitro on the transport of potassium ions in thymocytes, 26:45744
- CHAPUIS, A. M., New reading system for an ionographic detector, 26:45598
- CHARLTON, D. E., Fluorescence photon and Auger electron spectra, 26:47209
- CHARLTON, G. R., Remarks on average pion multiplicities at high energies, 26:46662
- CHARPAK, G., New results on $(\pi^+, 2p)$ reactions in light nuclei, 26:46870
- CHASE, A. B., Crystallization of BeO , Be_2SiO_4 , and SiO_2 from Li_2MoO_4 - MoO_3 , 26:45943
- CHASSANY, J., Radioprotection practice at the Marcoule center, 26:45804
- CHASTAGNER, P., Analysis of curium and californium by single-exposure spark source mass spectrometry, 26:45260
- CHATTERJEE, A., Fission barrier heights, energy release, and fragment energy partition processes from the viewpoints of nuclear structural models, 26:47046 (INDC-7-U, pp 145-60)
- Prompt gamma decay of fission fragments, 26:47025
- CHATTOPADHYAY, A., Activation analysis of environmental pollutants, 26:45198
- CHAUDHURI, K. D., Effect of strain on ultrasonic attenuation in superconductors, 26:46824
- CHAUDRON, Georges, Recrystallization of metals of high purity, 26:45982
- CHAUMEAUX, A., $^{56}Ni(\alpha, \alpha')$ and the nature of high-lying states in ^{56}Ni , 26:46927
- CHAUVEY, Paul, Method of decontaminating cesium, 26:45384
- CHAZELAS, E., π^0 electroproduction at first resonance, 26:46512 (CONF-710816-16)
- CHECHETKIN, Yu. V., Determination of the concentration of gamma-emitting isotopes in extended sources without sampling, 26:45264 (SRARI-P-123)
- CHEISHVILI, O. D., Theory of electron tunneling in NIS systems with paramagnetic impurities, 26:46837
- CHEN, C. W., Vacancy defects in quenched vanadium, 26:46025
- CHEN, Y. H., High-speed photography of a plasma focus, 26:47121
- CHENG, Chung-chieh, Theoretical studies of flux and energy spectrum of gamma radiation from the Sun, 26:46231
- CHERN, De-ching, Exact stationary solutions for the classical relativistic three-particle problem, 26:46790
- CHERNOV, L. A., Use of solid track detectors in reactor experiments, 26:45610
- CHESHIRE, D. L., Observation of the $M^0(940)$ and $\Delta^0(963)$, 26:46475 (IS-2886)
- CHESNOKOVA, V. D., Sub-barrier (p,n) reaction in region of single-particle resonance, 26:46952
- CHEUNG, Lung, Drift-kinetic equation and electrostatic instability theory in Tokamak plasmas, 26:47166
- CHIANG, C. C., On a possible repulsive interaction in universal gravitation, 26:46749 (ORO-3992-94)
- CHIANG, I-hung, Experimental study of K^+ decay, 26:46455
- CHIBISOV, N. P., Chemical shifts in $L_{\text{sub } \alpha 1}$ x-ray line of some U compounds, 26:45377
- CHILENSKAS, Albert A., Electrochemical power-producing cell, 26:45434
- CHIN-FATT, Cecil A., Enhanced far-infrared radiation from a theta-pinch plasma, 26:47143
- CHIPLONKAR, V. T., Plasma confinement by external rf signal, 26:47114
- CHISLEAG, C., Glucose-6-phosphate phosphohydrolase activity in erythrocytes after x-irradiation, 26:45848
- CHITRE, D. M., Nature of gravitational synchrotron radiation, 26:46757
- CHIU, C. B., Model for the mixing among the 70 baryon states based on exchange degeneracy, 26:46685
- CHIZHOV, G. A., Induced emission from electron in magnetic field at level $n = 1$, 26:46551
- CHLADEK, D., Metabolism of monomeric and polymeric Pu in the rabbit. Comparison with the mouse, 26:45917
- CHLENOV, M. A., 1,6-glucodialdose as one of radiolysis products of D-glucose-6-phosphate, 26:45337
- CHO, Mann, Fast neutron time-of-flight spectrometer with high resolution, 26:46848
- CHOSA, M., γ radiolysis of gaseous carbon dioxide at high densities, 26:45346
- CHOUDHARY, D. K., Cystein protection against ethylenimine damage in barley seeds, 26:45772
- CHOUDHURY, D. K., Partons may have integral charge, 26:46574
- CHOW, S. P., High-speed photography of a plasma focus,

- CHOWDHURY, A. Roy, Charge distributions of partons and scaling in deep inelastic scattering, 26:46566
- CHOWDHURY, R. K. Roy, Charge distributions of partons and scaling in deep inelastic scattering, 26:46566
- CHRISTENSEN, Joergen, Low-grade heat from thermal electricity production, 26:45573 (AE-448)
- CHRISTIAN, J. W., Some comments on Burgers and Bogers: Burgers transformation mechanisms and their relation to mathematical theories of martensite crystallography, 26:46037
- CHRISTOFIDIS, Nicholas C., Proton E-layer Astron for producing controlled fusion reactions, 26:47175
- CHRISTOPHOROU, L. G., Low-energy electron attachment to molecules at high densities, 26:46364 (ORNL-TM-3839)
- CHRISTOV, Chr., Asymptotic numbers: one generalization of the notion of number, 26:47284 (IC-72/45)
- CHU, C. K., Research in computational plasma physics. Progress report, March 1, 1971--March 1, 1972, 26:47145 (COO-3086-15)
- CHU, Richard R. C., Aerosols produced by x-rays, 26:46082
- CHUBAROV, L. B., Effect of γ radiation on oscillation frequency of AT-cut piezoelectric plates, 26:45667 (SC-T-722519)
- CHUCHALIN, I. P., Electron acceleration in an undulator synchrotron, 26:46120
- CHUDOTVOROV, A. A., Protection from neutron emission of ^{239}Pu dioxide, 26:46852
- CHUISTOV, K. V., Separation processes in superconducting alloys of niobium--aluminum, 26:46020
- CHUMBAROV, Yu. K., Study of the characteristics of emergency track detectors, 26:45617 (FEI-296)
- CHUMIN, V. G., α -decay of neutron-deficient isotopes of Ra, 26:47004
- CHUNG, S., Excitation of the electronic states of the nitrogen molecule by electron impact, 26:46391
- CHURIN, I. N., Time measurements with a germanium detector, 26:45616
- CHUTKIN, O. A., Distribution of a radioactive aerosol in a model chamber, 26:45444
- CHYSKY, Jaromir, Radioactivity of waste waters in Nuclear Research Inst., Rez, and environment, 26:45565
- CICUTA, G. M., Analytic renormalization via continuous space dimension, 26:46729
- CIGNA, Arrigo A., Conventional and radioactive pollution in the environment, 26:45555
- CLAFLIN III, Edward Scott, Identification of solar-neutron events by observation of radiation-belt proton fluxes, 26:46320
- CLARK, A. M., New method for quantitative study of induced autosomal non-disjunction in *Drosophila melanogaster*, 26:45880
- CLARK, Leonard, Fecundity and longevity of F_1 females of *Habrobracon* from sperm x-rayed with 3000 r, 26:45878
- CLARK, Paul V., Fused salts. A bibliography. Volume I. Citations listing. Volume II. Subject/title index. Volume III. Authors index, 26:45287 (SC-R-722769)
- CLARK, Reginald Wayne, Plasma heating near the ion-cyclotron frequency and its harmonics, 26:47109
- CLARKE, E. C. W., Deuterium separation process, 26:45690
- CLARKE, R., Ferroelectrics in $\text{K}_2\text{O--SrO--Nb}_2\text{O}_5$ system, 26:45962
- CLARKE, R. H., Multi-compartment model for studying the dose-intake relationship for radioactive isotopes, 26:45905 (RD/B/N-2291)
- CLARKSON, T. W., Effect of a polythiol synthetic resin on the absorption and excretion of methylmercury in experimental animals, 26:47643 (UR-3490-80)
- CLAUDEL, Bernard, Determination of the structure of monohydrated uranyl diformate using x-ray diffraction, 26:45368
- CLAUSS, W., WAK: specifications and finishing control for equipment, 26:46105 (ORNL-tr-2371)
- CLAYTON, Donag D., Hydrostatic oxygen burning in stars. II. Oxygen burning at balanced power, 26:46178
- CLAYTON, E. D., Subcritical multiplication of ^{252}Cf neutrons and its applications, 26:46855
- CLAYTON, John C., Nuclear fuel comprising uranium dioxide in a porous ceramic oxide matrix, 26:45957
- CLEAVER, J. E., Ocular squamous cell carcinoma (cancer eye) in hereford cattle: radiation repair processes and a comparison of cultured cells with xeroderma pigmentosum in man, 26:45851
- CLEMENT, D., Consideration on direct reaction mechanisms in the unified microscopic nuclear theory, 26:47077
- CLEMENT, J. M., MeV total neutron cross sections of ^{238}U , 26:47013 (COO-3058-23)
- COCHAVI, S., Magnetic moment of the 2761 keV 8^+ state in ^{92}Mo , 26:46959
- COHEN, D. D., Cyclotron harmonic waves in a reflex discharge, 26:47184
- COHEN, P., Nuclear explosives and hydrocarbons, 26:45493
- COHN, Arthur, Linear Stark effect due to resonant interactions of static and dynamic fields, 26:46361
- COKER, W. R., Coupled-channel calculations of the energy dependence of the (p,n) charge-exchange reaction, 26:47073
- COLARD, J., Human faecal ^{226}Ra excretion shortly after a RaSO_4 dust contamination with alginate treatment, 26:45907
- COLEMAN, George E., Apparatus for irradiation in a controlled atmosphere, 26:45448
- COLLINS, Carol H., Annealing reactions in K chromate. III, 26:45364
- COLLINS, Kenneth E., Annealing reactions in K chromate. III, 26:45364
- COLLINS, P. D. B., Duality properties of Regge cuts, 26:46600
- Kinematics of coupling of Regge poles to photons, 26:46560
- COLOMBO, A., Trace level determination of oxygen in organic coolants and ultra-micro determination of oxygen in C--H--O organic compounds, 26:45221 (EUR-4822)
- COMAR, D., Simple and rapid procedure for strontium determination in rat plasma using neutron activation, 26:45202
- COMPTON, R. N., Slow electrons from electron-impact ionization of He, Ne, and Ar, 26:46390
- COMSTOCK, Irving J., Method of fabricating a heat pipe, 26:45449
- CONARD, Robert A., Effects of age and radiation exposure on chromosomes in a Marshall Island population, 26:45792
- CONDE, H., Recent studies on prompt neutron and gamma rays from fission, 26:47020 (INDC-7-U, pp 113-29)
- CONNOLLY, T. F., Compilation of crystal growers and crystal growth projects, 26:45973 (ORNL-RMIC-12)
- CONRAD, Georges, Preliminary data on tritium content of aquifer sheets in northwest and central (Algeria) Sahara; hydrogeological consequences, 26:45558
- CONWAY, R. G., Polarization of radio sources at $\lambda 49$ CM and $\lambda 73$ CM, 26:46204
- COOK, M. W., Approximate solutions of the relativistic gravitational field equations to describe clusters of galaxies, 26:46253
- COOKS, R. G., Isotope effects upon kinetic energy release in metastable ion fragmentations, 26:45301
- COOPER, A. R., Diffusion in the system $\text{K}_2\text{O--SrO--SiO}_2$. III. Interdiffusion coefficients, 26:45946
- COOPER, Bernard R., dc anisotropic magnetization at 20.4 K and the ideal crystal-field-only behavior of TmSb , 26:47258
- COOPER, David M., Spectral intensity measurements from high-pressure nitrogen plasmas, 26:47117
- COOPER, T., Deformation parameters of ^{152}Sm by electron scattering, 26:46993
- COOPS, M. S., Contamination control hardware for remote hot cell transfers, 26:46108 (UCRL-73757)
- Shielded caves for dual-mode operation, 26:46107 (UCRL-73756)
- COOTE, G. E., Gamma radiation of some plants and soils from a uraniumiferous area in New Zealand, 26:45529
- COPPAGE, F. N., Device degradation from the effects of nuclear radiation on passivation materials, 26:45662 (SC-DC-721232)
- CORBETT, James W., Radiation-induced voids in metals. Proceedings of the 1971 international conference held at Albany, New York, June 9--11, 1971, 26:46045 (CONF-710601)
- CORIOU, H., Effect of heat treatments on corrosion of 18/10 stainless steels with low C contents in nitric medium in presence of ferric iron, 26:45970
- CORLEY, J. P., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- CORONIN, J. W., Experimental study of the decays $\text{K}/\text{sub } S \text{ L} / \rightarrow \gamma\gamma$, 26:46434
- CORONITI, F. V., Changes in magnetospheric configuration during the substorm growth phase, 26:46305
- CORRUCCINI, L. R., Spin-wave phenomena in liquid ^3He systems, 26:46802
- CORY, John Slane, Mass, momentum, and energy flow from an MPD accelerator, 26:47137
- COSME, G., Hadronic production by e^+e^- collisions at 990 MeV with Orsay storage ring, 26:46429 (CONF-710816-15)
- Measurement of radiative neutral decay modes of ω^0 and ϕ^0 mesons with Orsay storage ring, 26:46472 (CONF-710816-12)
- $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
- $\pi^+\pi^-$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
- COSTA, E. C., Optimization study of the precipitation of thorium oxalate, 26:45365 (IEA-183)
- COSTA, S., Experiments on a superposed fields theta pinch, 26:47171
- COULON, R., Monitoring of the radioactive contamination of wheat flour, 26:45903
- COUNSELMAN III, C. C., Density of the solar corona from occultations of NP 0532, 26:46227
- COUPAT, B., Inelastic interaction induced by high-energy muons (6 GeV, 12 GeV) at low momentum-transfer in nuclear emulsion, 26:46451
- COURTES, G., Two new large-diameter galactic regions of faint H α emission, 26:46236
- COYTUPA, Juan, Simultaneous radioimmunoassay of plasma testosterone and dihydrotestosterone, 26:45928
- COZAR, O., Influence of ethanol on hydrated complexes of Cu ions, 26:45296
- CRAIGEN, W. J. S., Zirconium tetrafluoride process, 26:45274
- CRANCON, J., On-line measurements of rare-gas fission yields in 14 MeV neutron fission, 26:47030

- CRAWFORD, D. M., Conversion of bremsstrahlung photo neutron mean energies to monochromatic photo neutron mean energies, 26:46953
- CRAWFORD, H. J., Composition and energy spectra of heavy nuclei with $0.5 < E < 40$ MeV per nucleon in the January 24, and September 1, 1971 solar flares, 26:46232
- CRAWFORD, W. R., Thermal plume study at Douglas Point, Lake Huron, 26:45581
- CREASIA, D. A., Increased effective half-life of intratracheal administered $^{51}\text{Cr}_2\text{O}_3$ by respiratory infection, 26:45910 (CONF-720505-2)
- CREEK, Ronald B., Sodium heated steam generator, 26:45450
- CROOKS, G. B., Observation of a broad resonance in the 2^{35}S excitation of helium by electron impact, 26:46405
- CROSS, William Gunn, Method and apparatus for counting radiation damage tracks in sheet material using a capacitive discharge device, 26:46083
- CROUCH, S. R., Some comments on the signal-to-noise characteristics of real photomultiplier and photodiode detection systems, 26:45682
- CROME, A., Ionization of argon by electron impact, 26:46374
- CSABA, G., Tissue mast cells in human blood, 26:45810
- CSANADY, G. T., Thermal plume study at Douglas Point, Lake Huron, 26:45581
- CSUEROES, M., Determination of isodose curves of sealed γ -sources by calculation and film dosimetry, 26:45715
- CSUZI, S., Biochemical foundations of spore resistance, 26:45751
- CULLIFORD, E. R., Some measurements of the electrical conductivity induced in polymers by pulsed x-ray irradiation, 26:47225 (AWRE-O-31/72)
- CURRAN, M. J., Preparation of transparent Y_2O_3 -doped ThO_2 , 26:45948
- CVJETICANIN, D., $2^{+} \rightarrow 2^{+}$ transition of 879 keV in ^{160}Dy investigated by electron--gamma directional correlation, 26:46991
- CZAPLEWSKI, Stanislaw, Trial of correlating the blood picture with absorbed doses in persons exposed professionally to ionizing radiation, 26:45787 (TT-54002/5, pp 602-4)
- CZOSNOWSKA, W., Long-term study of urinary ^{90}Sr and calcium in children, 26:45908
- D**
- DA FONSECA, A. J. R., Chemical effects of isomeric transition in bromoethane mixtures, 26:45360
- DAHMEN, H. D., Method of characteristics for functional-derivative equations, 26:46728
- DAINTON, Frederick, Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. III, 26:45329
- Radiation-induced formation of excited states of aromatic hydrocarbons in cyclohexane. IV, 26:45330
- Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. II, 26:45328
- DAION, M. I., Photoproduction of neutral pions on helium nuclei, 26:46432 (SLAC-Trans-142)
- DAL'KAROV, D. D., Quantum numbers and total and partial widths of the resonance N anti $N(1970)$, 26:46458
- DALLY, E. B., Asymmetry measurements for π^0 and π^+ photoproduction by linearly polarized γ -rays in the energy range (40 to 900) MeV, 26:46442
- DALMAZZONE, Jean, Device for controlling slow drift in ionization chambers, 26:45609
- DALPIAZ, P., Decay mode ϕ + η and measurement of branching ratio, 26:46435
- DALY, P. J., Level structure of ^{187}Os from decay of ^{187}Ir , 26:46984
- DAMJANOVICH, S., Possibilities in studying the mechanism of radiosensitivity of and radiation damage to albumines and nucleic acids, 26:45726
- DAMS, Richard, Multi-element analysis of air pollution particulates by nondestructive neutron activation, 26:45207
- DANDARON, G.-N. B., Erosion of a tungsten cathode in nitrogen, 26:45968
- DANIEL, M., Analytic renormalization of the exponential interaction: the three-point time-ordered product with minimum light cone singularity, 26:46712
- DANNO, Akibumi, Radiolyses of mixtures of ammonia and hydrocarbons. II, 26:45332
- DANON, J., ESR studies of electron irradiated $\text{K}_3\text{Ir/sup III/(CN)}_6$ in KCl single crystals. I. Ligand hyperfine structure in Ir/sup II/(CN)_6 species, 26:45317
- DAR, A., Dominance of electromagnetic effects in high energy hadron and photon induced reactions, 26:46533
- DARWISH, Saher, Mean lives of some high energy levels in ^{160}Dy , 26:47000
- DAS, H. A., Determination of chromium in milk powder, 26:45196
- DAS, H. L., Effect of strain on ultrasonic attenuation in superconductors, 26:46824
- DAS, Tapan Kumar, Theoretical investigation of collective excitations in medium and heavy nuclei, 26:47102
- DAS GUPTA, M. K., Solar microwave bursts of different types in relation to optical flares, 26:46333
- DASGUPTA, S., Inducing dyeability in polypropylene by radiation induced grafting, 26:45336
- DAVIER, M., Pomeron and f exchange amplitudes in elastic $\pi^+\pi^+$ scattering, 26:46639
- DAVIES, B., Improvements on a new method for the numerical solution of the Schroedinger equation, 26:47290
- DAVIES, R. D., Variability of OH source W3, 26:46205
- DAVIS, Harold Alan, Investigation of low-density plasma shock waves in an expanding discharge, 26:47194
- DAVIS, J. Barry, Ecology of Hot Creek Valley: vegetation and soil response to underground detonation. Annual progress report FY 1971, 26:45478 (NVO-409-1)
- DAVIS, James E., Infectivity and development of x-irradiated metacercariae of Leucochloridiomorpha constantiae (Trematoda) in the chick and on the chorioallantois, 26:45879
- DAVIS, Karl C., Measurement of the isotropy of the microwave background temperature at 6-cm wavelength, 26:46255
- DAVIS, Lawrence L., Analysis of seismic waves recorded in Las Vegas, Nevada from underground nuclear detonations, 26:45481 (NVO-1163-TM-33)
- DAVIS, Monte V., Vapor transport limits of liquid metal heat pipes, 26:45460
- DAVIS, Paul Richard, Electron stimulated desorption of carbon monoxide adsorbed on niobium (111), 26:47242
- DAVIS, Philip J., Experimental spectra for $12\text{ GeV/c K}^+\pi^+\pi^0$ and $\text{K}^+\pi^0$, 26:46484
- DAVIS, Robert, Effects of gamma radiation on the respiration of Acaerus siro, 26:45884
- DAVYDOV, V. V., Elastic scattering of ^6Li ions, 26:46883
- DAWES, E. A., Effects of dietary antioxidants on lipid peroxide formation in animal tissues after whole-body irradiation, 26:45872
- DE, Anil K., Combined ion exchange-solvent extraction (CIESE) studies of metal ions on ion exchange papers. Synergistic effects and chromatographic separations, 26:45282
- DE, P. K., Passivity of Type 304 stainless steel: effect of plastic deformation, 26:45967
- DE, S. S., Parton model and the concept of duality in strong interactions, 26:46599
- DEAN, A. G., Technique for recording Langmuir probe characteristics in afterglow plasmas, 26:47119
- DEARNLEY, Geoffrey, Ion channeling, 26:47228
- DE BORTOLI, M., Site characteristics, 26:45570 (EUR-4841)
- DEBUIGNE, Jean, Dysprosium borocarbides, 26:45952
- DE CARLOS, J. E., New approach to beta--gamma coincidence counting. Advance report on SAMAR electronic system, 26:45589 (JEN-247)
- DE CARVALHO, H. G., Some remarks on the transparency of complex nuclei to photoproduced charged pions at intermediate energies, 26:46889
- DECOURREGES, M. M., K_2^0 decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)
- DEDKOV, Yu. M., Photometric determination of zirconium in natural materials with the reagent picramineepsilon, 26:45233
- DE FOUCHIER, Alain, Surface preparation of uranium parts, 26:45980
- DEGEN, V., $\text{H}\beta/\text{N}_2^+$ 4709-A intensity ratio and vibrational enhancement of N_2^+ first negative bands in proton-excited auroras, 26:46276
- DEGTARENKO, N. N., Formation of the energy spectrum of the ions of a laser plasma, 26:47154
- DEGTAREV, Yu. G., ($n,n'\gamma$) reaction in ^{27}Al , ^{51}V , ^{52}Cr , ^{55}Mn , ^{56}Fe , ^{93}Nb , ^{208}Pb , and ^{209}Bi , 26:46909
- DEGUCHI, Ichiro, Disappearance of neutron diffusion length of beryllium and graphite according to temperature, 26:46857
- DEGUCHI, Toshio, Paper chromatography of rare earth elements with developer containing of ethylene glycol monoethers, hydrophobic solvents, and oxine, 26:45275
- DE JONG, I. G., Nuclear recoil reactions in organomanganese compounds. VI, 26:45357
- DELAGRANGE, Hugues, De-excitation of compound nuclei formed in $^{181}\text{Ta} + ^{11}\text{B}$ reactions, 26:46974 (FRNC-TH-207)
- DELAHAYE, R. P., Radiobiological problems caused by supersonic transport. (With a survey of the first results established by tests performed on board the Concorde prototype), 26:45796
- DE LATIL, Pierre, Gum nebula, 26:46165
- DELBOROUGH, R., Gravitational correction to PCAC, 26:46501
- DELL'ANTONIO, G. F., Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)
- DELMAS, Pierre, Automatic analysis by absorption spectrometry, 26:45243
- DELONG, Max Myron, Prediction of heated effluent dispersion in a stream using similitude techniques, 26:45575
- DE LOPEZ, Ofelia H., Procedure for chronic intravenous catheterization in the rabbit, 26:47641 (UR-3490-21)
- DELPECH, Andre, Method for the manufacture of fissile oxides and the apparatus developed, 26:45940
- DE MARIA, G., Thermodynamics of rare-earth--carbon systems. IV. The lutetium--carbon system, 26:45961
- DEMCHENKO, V. V., Nonlinear focusing of TM waves in isotropic plasma, 26:47185
- DEMENT'EVA, T. A., Content of sulphydryl groups in mitochondria after irradiation, 26:45829
- DEMIANSKI, M., Homogeneous rotating universe with flat space, 26:46261

DEMICHELE, Don W., Vapor transport limits of liquid metal heat pipes, 26:45460

DEMMIE, Paul Norman, Characteristic development of trapped surfaces in Einstein-scalar spacetimes, 26:46764

DEMOISE, Charles F., Effects of age and radiation exposure on chromosomes in a Marshall Island population, 26:45792

DEM'YANOVA, A. S., Mechanism of $^9\text{Be}(p,p_1)^9\text{Be}$ 1.68 MeV reaction, 26:46885

DENNISON, R. A., Respiratory quotients of irradiated fruits, 26:45729

DE NORDWALL, H. J., Chemistry of HTGR, 26:45288 (TID-26106)

DENTON, R., Thermodynamic properties of small superconducting particles, 26:46813

DERADO, I., Forward production of $\Delta(1236)$ in pion-proton scattering, 26:46488

DE RAFAEL, E., Recent developments in the comparison between theory and experiments in quantum electrodynamics, 26:46710

DERENGOWSKI, M. A., Fission isomeric state in ^{236}U , 26:47018 (INDC-7-U, pp 61-4)

DERFLER, H., Cyclotron harmonic waves in an inhomogeneous plasma. I, 26:47188

DERIAN, J. C., French program on production and uses of ^{238}Pu , 26:45706

DE RIDDER, Leo I., Phenol red uptake by irradiated and nonirradiated chick mesonephros in vitro, 26:45832

DERLIKOWSKI, J., Changes of Dextran irradiated with γ radiation, 26:45730

DE SAINT-SIMON, M., Systematic study of the decomposition of very small quantities of rare earth oxides by mass spectrometry, 26:45259

DE SAUSSURE, G., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)

DES LIGNERIS, Jean, Method of decontaminating cesium, 26:45384

DESPRETS, A., Harmonie reactor, 26:47590

DE TOURREIL, Roland, Library of data for fission products (third edition), 26:47012 (CEA-N-1526)

DEUTSCH, R. V., Introduction to low temperature physics, 26:46841

DE VAUCELLES, Emmanuel, Surface preparation of uranium parts, 26:45980

DE VAULCHIER DU DESCHAUX, Louis, Surface preparation of uranium parts, 26:45980

DEVRIES, R., $^{58}\text{Ni}(\alpha,\alpha')^{58}\text{Ni}$ and the nature of high-lying states in ^{58}Ni , 26:46927

DEWEY, W. C., Confirmation of lesions having the potential for forming chromosomal aberrations, 26:45743

DEW-HUGHES, D., Effect of dislocation tangles on superconducting properties, 26:46836

DIAZ-GUERRA, J. P., Volatilization mechanism of certain elements in a method of semiquantitative spectrographic analysis, 26:45222 (JEN-229-DQ/I-79)

DI CASA, M., Radioanalytical investigation on the extraction behavior of rhodium in thiocyanate-TBP system, 26:45194

DICKE, R. H., Faculae and the solar oblateness, 26:46225

DICKERSON, Edwards S., DTA--x-ray diffraction examination of chemical reprocessing wastes containing sodium and potassium nitrates, 26:45383 (ICP-1008)

DICKEY, J. M., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)

DI CRESCENZO, Enrico, Phase sensitive method for detecting superconductive transitions in the microwave region, 26:46840

DIEHL, J. F., Disinfecting mite infested ham by ionizing radiation, 26:45886

DIENER, E. M., Evidence for an anomaly in the excitation function of the reaction $^2\text{H}(p,\gamma)^3\text{He}$, 26:46868

DI MARCO, J. N., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)

DI NAPOLI, V., Some remarks on the transparency of complex nuclei to photoproduced charged pions at intermediate energies, 26:46889

DMITRIEV, A. G., Decay of ^{76}Br , 26:46938

DOBROVOL'SKII, O. V., Size of the dust particles of a comet atmosphere and the heliocentric distance, 26:46220

DOD, Raymond Lloyd, Some properties of protactinium metal and its compounds, 26:45992 (LBL-659)

DODD, C. V., Nondestructive test for measuring the state of heat treatment in closure welds, 26:45470

DODEL, G., Simultaneous spectroscopic determination of electron density and temperature as a proof of relaxation effects in a streaming argon-potassium-plasma, 26:47116 (IPP-IV-12)

DOERING, George I., System and method for facilitating the operating setup of a radiation gauge, 26:45654

DOHERTY, L. H., Luminosity-spectral index relationship for radio galaxies, 26:46248

DOHERTY, R., Radiation menopause, 26:45807

DOKSHINA, G. A., Content of sulfhydryl groups in mitochondria after irradiation, 26:45829

DOLGOV, A. D., Interpretation of experiments on searches for $K/\text{sub } S/\text{sub } 2p$ decay, 26:46572

DOMANY, E., Algebraic derivation of generalized Talmi expansion, 26:46503

DOMBEY, Norman, Is the pion form factor determined by electroproduction experiments, 26:46513 (CONF-710816-17)

DOMOKOS, G., Lightlike quark model and current densities, 26:46524 (NYO-4076-18)

Wave packet realization of lightlike states, 26:46589

Wave packet realization of lightlike states, 26:46681 (NYO-4076-19)

DONAHUE, T. M., Analysis of Ogo 6 observations of the O I 5577-A tropical nightglow, 26:46271

DONALDSON, Gordon W., Superconductive magnets, 26:46843

DONALDSON, R. R., Simple method for separating spindle error from test ball roundness error, 26:47635 (UCRL-73805)

DONATO, A., ESTER program: high-level radioactive waste solidification, 26:45385

DONNELLY, R. G., ORNL isotopic power fuels quarterly report for period ending March 31, 1972, 26:45703 (ORNL-4796)

DONOHUE, J. T., Baryon spin determinations using polarized targets, 26:46628

Tests of the spin assignment and equal-phase assumption for high-spin baryons, 26:46621

DONOVAN, F. M. Jr., Biomedical engineering support. Progress report, 26:45700 (COO-2155-4)

DONS, R. F., Effect of various cell-free organ extracts on hematopoietic stem cells, 26:45847

DOROSZOWSKI, Jan, Statistical data on iodine uptake of the thyroid gland. II, 26:45906 (TT-54002/4, pp 462-70)

DOUSSET, N., Effect of water on the radiosensitivity of plasmatic fatty acids, 26:45725

DOUSTE-BLAZY, L., Effect of water on the radiosensitivity of plasmatic fatty acids, 26:45725

DOVER, C. B., Nucleon-nucleus potential in Hartree-Fock approximation with Skyrme's interaction, 26:47064

DOWKER, J. S., Are quarks possible, 26:46683

DOZMOROV, I. M., Solution of Einstein equations with null bundle. II, 26:46763

Solution of Einstein equations with null bundle. III, 26:46762

DRAKE, N. A., Protection against gastrointestinal effects of whole-body x-irradiation by a bile acid sequestrant in rats, 26:45844

DRICKEY, D. J., Photoproduction experiments at ISABELLE, 26:46426 (BNL-16968)

DRONKERS, J., Phenomenological analysis of $n\text{N}$ and KN charge-exchange reactions, 26:46630

DRUMHELLER, Kirk, Process of forming an isotopic heat source, 26:45707

DRYAPACHENKO, I. P., Angular distribution of neutrons from (p,n) reactions on ^{18}O , ^{37}Cl , ^{41}K , ^{53}Cr , and ^{59}Co nuclei, 26:46929

DRYNNIN, V. I., Gold assay on geological samples by neutron activation with preliminary crucible fusion, 26:45200

DUBARD, James L., Low-lying states and configurations in ^{138}La , 26:46964

DUBOIS, R. D., Observation of a broad resonance in the ^2S excitation of helium by electron impact, 26:46405

DUBROVIN, A. S., Calorimetric study of Nb-Fe-Si-Al system, 26:46007

DUBROVIN, S. A., Dose to the epithelium of the upper respiratory tract from radon and its daughter products, 26:45861

DUBROVSKAYA, L. B., Magnetic susceptibility of cubic niobium and tantalum carbides in the low temperature range, 26:47266

DUBSKY, F., Selective ion exchangers with isonitrosoacetamide functional groups. II, 26:45283

DUDNIKOVA, G. I., Effect of thermal conductivity on the structure and critical parameters of shock waves in a plasma, 26:47200

DUGAN, C. H., Plasma-sampling diagnostics using ion kinetic energy distributions, 26:47128

DUIMIO, F., Irreducible representations of para-Bose operators, 26:46507

DUKE, K. M., Environmental benefit-cost analysis for nuclear power generation, 26:47386

DUMESNIL, Patrice, Semiconductor well logging probe, 26:45658

DUMITRESCU, O., Resonance scattering theory and the structure of compound states, 26:47068

DUNAIEVSKII, S. Ya., Dynamics of control system incorporating 440 MW(e) reactor, 26:47324

DUNCAN, G. Comer, Application of Green's functions to radiative cooperative effects in multi-atom systems, 26:46418

DUNCAN, Joseph R., Measurement of liquid droplet emissions from cooling towers and process stacks, 26:45574 (CONF-720619-1)

DUPREE, S. A., Electromagnetic pulse and time-dependent escape of neutrons and gamma rays from a nuclear explosion, 26:45497

DURAND, Emile, $\Sigma/\text{sup } p q/\text{ tensor associated with the polarization quadrivector } \sigma_{\mu\nu}/\text{sub } r/\text{ and the Maxwell } \Pi/\text{sup } p q/\text{ tensor associated with the energy-momentum quadrivector } w_{\mu\nu}/\text{sub } r/\text{ in complex fields, 26:46693}$

DURELL, J. L., Missing low-lying states in ^{39}K , 26:46928

DURISEN, Richard Henry, Viscous effects in rapidly rotating stars with application to white-dwarf models, 26:46163

DURNEY, A. C., Entry of high-energy solar protons into the distant geomagnetic tail, 26:46304

DURRANI, S. A., Prolonged etching factor in solid state track detection and its applications, 26:45604

DUSHUTIN, N. K., Feynman-Wilson-gas model as consequence

- of jet approach to inelastic hadron processes, 26:46586 (JINR-P2-6501)
- DUVAL, Christian, Model for particles with spin in the gravitational and electromagnetic field, 26:46767
- DVORAK, S., Remark on neutron slowing down, 26:46856
- DWORSCHAK, F., Comparison of lattice parameter and resistivity change in electron-irradiated copper, 26:47231
- DYCK, H. M., Measurement of polarized 10-micron radiation from cool stars with circumstellar shells, 26:46185
- DYKLA, John J., On a possible repulsive interaction in universal gravitation, 26:46749 (ORD-3992-94)
- DZHELEPOV, B. S., Decay of ^{78}Br , 26:46938
- Isobaric nuclei with mass number $A = 72$, 26:46939
- DZHGARKAVA, N. N., Gamma quanta distribution at large distances from the sources, 26:47215
- DZHOLOS, R. V., Isomeric transition $8^- \rightarrow 6^+$ in ^{130}Ba nucleus, 26:46949
- DZUBAY, T. G., Energy levels of ^{36}K , 26:46904

E

- EAGLE, R., Natural ^{210}Pb and ^{210}Po in a marine environment, 26:45544 (RLD-2225-T-14-4)
- EBEL, A., Electron content of ionosphere, solar, and geomagnetic activity, 26:46287
- EBERHARD, K. A., Cross sections for $(\pi^-, \pi^+ n)$ reactions on ^{19}F and ^{21}p in the 3 to 3 resonance region, 26:46486
- EBERLE, H., Effects of actinomycin D on G₁ protein synthesis in L5178Y cells, 26:47650 (UR-3490-153)
- Protein synthesis during the cell cycle of L5178Y cells, 26:47651 (UR-3490-154)
- EBERT, D., Reggeon perturbation theory in Zachariasen-like model, 26:46747
- EBISAWA, Hiromichi, Magnetic properties of a superconductor with hexagonal symmetry, 26:46807
- ECKMANN, Jean-Pierre, Representations of CCR in $(\phi^*)_3$ model: Independence of space cutoff, 26:46715
- EDA, Bunzo, ESR study of the vinyl-type radical trapped in a single crystal of maleic acid irradiated at 77°K, 26:45347
- EDWARDS, Arthur L., SHAKER: a computer program for numerical analysis of weighted tabular data, 26:47631 (UCRL-51210)
- EDWARDS, David F., Time-resolved diagnostic method for hydrogen plasmas, 26:47127
- EFIMOV, G. F., Choice of a form factor in nonlocal quantum electrodynamics, 26:46698 (ITF-71-128-P)
- EFIMOV, G. V., Choice of form factors in non-local quantum electrodynamics, 26:46711
- EFIMOV, V. P., Peculiarities in interaction of high-energy electrons and positrons with crystals, 26:46550
- EFINGER, H. J., Example of a covariant field theory of elementary particles, 26:46723
- EGAMOV, U., Properties of ruby laser radiation under action of ^{60}Co gamma radiation, 26:47227
- EGGEN, O. J., Luminosities and motions of the F-Type stars. II. Metal-deficient stars, 26:46183
- EGIAZAROV, A. S., Separation of sulfur isotopes by a chemical exchange method, 26:45688 (MLM-1936(TR))
- EGIYAN, K. Sh., Angular and energy distributions of protons in (γp) and (γp) reactions on ^{12}C nucleus for 130 MeV excitation energy, 26:46888
- Control system for beam formation after single deflection at exit of 300 MeV electron linear accelerator, 26:46131
- EGORENKOV, V. D., Stability of a finite-pressure plasma with an anisotropic electron-velocity distribution, 26:47159 (UCRL-Trans-10630)
- EGOROV, A. A., Separate control of the injector heating and of the current amplitude in transformer accelerators, 26:46135
- EGOROV, A. V., Photoproduction of vector and axial vector resonances in broken nonlinear chiral $\text{SU}_3 \times \text{SU}_3$ dynamics, 26:46563
- EGOROV, N. I., Peculiarities of formation of an increased uranium concentration in oxidizing conditions, 26:45504
- EGOROV, V. A., Magnetic compensation in iron--gadolinium and iron--terbium films, 26:47273
- EGORSHIN, A. A., Longitudinal impact on end face of prismatic rods, cylindrical, and conical shells, 26:47626 (SC-T-722520)
- EICHLER, J., Four-particle correlations in nuclei, 26:45971 (HMI-B-121, pp VIII.1-37)
- EICHTEIN, T., Particle production in proton interactions in nuclei at 24 GeV/c, 26:46459
- EINHORN, N., Lymphocyte stimulation by thyroglobulin after local irradiation of the thyroid gland in man, 26:45889
- EISENBERG, Y., Simple diffraction considerations and relations between $\sigma/\text{sub } T$ and slope of vector meson electroproduction, 26:46564
- EISENSTEIN, R. A., Measurement of the magnetic moment of the antiproton, 26:46440
- Nuclear γ rays associated with stopped kaons, 26:46940
- EISMONT, V. P., X ray fluorescence analysis of heavy element by an α -X coincidence method, 26:45270
- EISNER, Robert L., Evidence for existence of a new meson in the 955 MeV mass region, 26:46471 (BNL-16954)
- EKIMYAN, G. A., Detector for high energy particles based on the principle of using particle radiation in a laminated medium, 26:45648
- EKIN, John W., Low-temperature electron--phonon umklapp scattering in simple metals, 26:47264
- EKSTROEM, P., Energy of levels of ^{45}Sc up to 3 MeV from $^{44}\text{Ca}(p, \gamma)^{45}\text{Sc}$ reaction, 26:46920
- ELAYAPERUMAL, K., Passivity of Type 304 stainless steel: effect of plastic deformation, 26:45967
- ELDERKIN, C. E., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- ELDRUP, Morten, Positron lifetimes in water and ice, and in frozen aqueous solutions, 26:46362 (RISO-254)
- ELENSKII, Ya. S., Source of high-intensity nanosecond light pulses, 26:45640
- EL GAVKHARI, A., Spin correlation during bremsstrahlung radiation of longitudinally polarized electrons, 26:47052
- ELISEEV, S. M., Analysis of inelastic collisions of particles with nuclei at very high energies, 26:46649 (JINR-P2-6395)
- ELKIND, M. M., Repair of a DNA complex from x-irradiated Chinese hamster cells, 26:45745
- ELLDER, J., Variability of OH source W3, 26:46205
- ELLIS, J., Canonical anomalies and broken scale invariance, 26:46556
- ELLIS, W. R., Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
- EMMONS, George Herbert, Thermodynamic study of order--disorder transitions in vanadium carbide, 26:45937
- EMOD, Gyula, Grain refining of magnesium alloys, 26:45984
- ENACHE, Ana, Effect of a particles on operating regime of a CO_2 laser, 26:45669
- ENDRZHEVSKAYA, S. N., Determination of alloying additives in artificial fluorite, 26:45232
- ENGDAHL, E. R., MILROW/CANNIKIN seismic effects, 26:45479 (NVO-746-122)
- ENGELMAN, F., Flute modes in a plasma in the presence of nonuniform electric fields, 26:47162
- ENGELMANN, Ch., Activation analysis with gamma rays and charged particles, 26:45210
- ENGEL, I. A., On-line reactivity feedback analysis of EBR-II by rod drop, 26:47307
- ENGFER, R., X-rays and nuclear γ -rays in muonic thallium, 26:47031
- ENGLUND, H. M., Proceedings of the second international clean air congress held at Washington, D. C., December 6--11, 1970, 26:45205
- ENIKOVA, M., ^{132}Ba levels excited during decay of ^{132}La and ^{132}La , 26:46948
- ENKOVSKII, L. L., Threshold behavior of DAMA, 26:46774 (ITF-72-44-E)
- ENRIETTO, J. F., Discussion on fracture mechanics concepts, 26:45996 (WCAP-7841)
- ENSSLIN, N., Deformation parameters of ^{152}Sm by electron scattering, 26:46993
- EPEL, L. G., Continuous sampling plan for the taking of physical inventories, 26:46092 (BNL-16930)
- EQUER, B., K_2^0 decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)
- ERB, K. A., ($^{16}\text{O}, ^{12}\text{C}$) studies in the $2s-1d$ shell, 26:46913
- ERGAKOV, V. A., Electrical insulation in electrostatic analyzers, 26:45687
- ERLANDSSON, B., Energy of levels of ^{45}Sc up to 3 MeV from $^{44}\text{Ca}(p, \gamma)^{45}\text{Sc}$ reaction, 26:46920
- ERLICH, Y., Determination of chromium in milk powder, 26:45196
- ERMAKOV, A. N., Photometric determination of zirconium in natural materials with the reagent picramineepsilon, 26:45233
- ERMOLAEV, E. A., Interaction of a ruby with ionizing radiation, 26:47244
- ERNST, Richard D., Pseudo cloud-chamber tracks and their uses, 26:45647 (UCRL-51214)
- ERSHOVA, L. S., Effect of Mo of aging on Fe--Co--W alloys, 26:46006
- ESAKIYA, K. E., Separation of sulfur isotopes by a chemical exchange method, 26:45688 (MLM-1936(TR))
- ESSENBERG, A., Workloads and use factors in diagnostic x-ray installations, 26:45811
- ESSINGTON, E. H., Distribution of radioactivity in and near the Rainier rubble chimney, 26:45483 (NVO-1229-180)
- Long term release of radioactivity from Rainier melt-glass, 26:45482 (NVO-1229-177)
- ESTERL, J. E., Beta-delayed proton decay of ^{130}I , 26:46879
- Confirmed proton radioactivity of $^{53}\text{M}/\text{Co}$, 26:46926
- ESTERL, John E., Further results on proton radioactivity of $^{53}\text{M}/\text{Co}$, 26:46919
- ESWARAN, M. A., Analog states in ^{33}Cl by isospin-forbidden resonances in reaction $^{32}\text{S}(p, \gamma)^{33}\text{Cl}$, 26:46902 (BARC-598)
- ETKIN, Asher, Study of the reactions $\text{anti } pp \rightarrow \text{anti } pp$ and $\text{anti } pp \rightarrow \pi^+ \pi^-$ using a polarized proton target, 26:46468
- ETSELL, T. H., Transport properties of thorium and thorium-based solid solutions, 26:45938
- EVANS, G. R., Measurement of slope parameter in decay $K/\text{sub } L^0 \rightarrow \pi^+ \pi^-$, 26:46494

EVANS, H. C., /sup 53m/Co: a proton-unstable isomer, 26:46925
 EVANS, J. H., Observation and theory of the void lattice in molybdenum, 26:46055 (CONF-710601-, pp 522-32)
 EVETTS, J. E., Critical currents in Nb₃Al, 26:46829
 EVLANOV, M. V., Diffraction disintegration of deuterons, 26:46923
 EZAWA, Z. F., Fixed-pole and virtual Compton amplitude, 26:46537
 EZHELA, V. V., Analytical properties of amplitudes and differential cross-sections for multiparticle production processes over two angular variables, 26:46497 (IFVE-STF-71-36)

F

FABREGA, S., Design considerations of heat exchangers for vaporizing fluids, 26:45436 (CEA-CONF-2010)
 FABRIKANT, Jacob I., Environmental radioactivity: radiation effects at the cellular level, 26:45890 (NYO-3970-29)
 Lymphoid-cell renewal under low-level irradiation. XI. The cell cycle and growth kinetics during radiation leukemogenesis in C57BL mice, 26:45822
 Radiation hazards in environmental medicine, 26:45904 (COO-3013-1)
 Radiobiology and the public health, 26:45911 (COO-3013-25)
 FACQUET, Louis, Surface preparation of uranium parts, 26:45980
 FADDEEV, L. D., Integral equations of scattering theory and few-nucleon systems, 26:47087
 FADNER, Willard Lee, Nuclear structure and reaction mechanism studies using the (³He,t) reaction, 26:46930
 FAES, M. H., Changes in the localization of acetylcholinesterase in neuromuscular junction after ⁶⁰Co irradiation, 26:45828
 FAESSLER, Amand, Orbital rearrangement in light nuclei, 26:46886
 FALK, Fred W., Absorbed ionizing radiation measuring device, 26:45635
 FARDEAU, P., Computerized control of an accelerator and the associated experiments, 26:46123
 FARHOOMAND, I., Observations of chimney damage to residential buildings in Glendale, California: San Fernando earthquake, February 1971, 26:45474 (JAB-99-88)
 FARKAS, J., Effect of combined methods on Clostridium sporogenes inoculated in sausage, 26:45747
 FARKAS, K., Tissue mast cells in human blood, 26:45810
 FARLEY, Thomas A., Inner-zone energetic-electron repopulation by radial diffusion, 26:46310
 FARRAR, Ruben T., Nuclear charged self-sustaining laser, 26:45677
 FARRELL, H. H., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)
 FASTLE, D. L., Large scale characterization tests of gelled slurry blasting agents, 26:45484 (SC-DC-722296)
 FAUGERAS, Pierre, Method of extracting fuel from nuclear fuel elements, 26:45422
 FAUSTOV, R. N., Relativistic transformations of the one-particle wave functions, 26:46667 (IFP-71-117-P)
 FAVIER, J., New results on (*,2p) reactions in light nuclei, 26:46870
 FAZIO, G. G., Detection of high-energy gamma rays from the Crab Nebula, 26:46214
 FEDERBUSH, Paul, Equivalence of dressing transformations, 26:46713
 FEDOROV, B. A., Effect of radioprotective preparations on the respiratory exchange of rats, 26:45823
 FEDOROV, N. A., Characteristics of nucleotide of DNA in radiation injury, 26:45856
 FEHRENBACH, W., ⁹⁰ electroproduction at first resonance, 26:46512 (CONF-710816-16)
 FELTEN, P., Study of the mechanical and thermal behavior of a high temperature reactor core, 26:47304
 FELVINCI, J., Mass distributions in fission and some related phenomena, 26:47017 (INDC-7-U, pp 31-59)
 FELVINCI, J. P., Fission isomeric state in ²³⁸U, 26:47018 (INDC-7-U, pp 61-4)
 FENEYROL, J. Y., Hexavalent W bronzes containing Th in interstitials: Th/sub x/sup IV/(B/sub x/sup V/W/sub 1-x/sup VI)/O₃ phases (B = Nb,Ta), 26:45953
 FENLON, M. W., Influences of thermal effluents upon aquatic production in Lake Ontario, 26:45578
 FERGUSON, Robert Nicholas, Isotope effects in the Barton reaction, 26:45300
 FERNANDEZ, Joseph Ignatius, Charge composition and energy spectrum of very heavy cosmic rays, 26:46151
 FERTL, W. H., Application of well logging methods in hydrocarbon prospecting, drilling, and production to the study of geostatic pressure problems, 26:45564
 FESHACH, H., Description of the interaction of high energy particles with nuclei, 26:47100
 FETTEROLF, Carlos M. Jr., Radioactivity in Lake Michigan water and biota, vicinity of Big Rock Point Nuclear Reactor, 1960-69, 26:45535
 FIALKOWSKI, K., Two-particle correlations in 60 GeV/c π -p and 67 GeV/c pp inclusive reactions, 26:46460
 FICHTEL, C. E., Comparison of measurements of the charge spectrum of solar cosmic rays from nuclear emulsions

and the Explorer 35 solid-state detector, 26:46330
 FICKES, Michael G., Scattering of CsF by electrons, 26:46406
 FIDELIS, I., Existence of regularities, tetrad, or double-double effect in changes in unit cell volumes of compounds of actinides in different oxidation states, 26:45375
 FIEL, R. J., Protein synthesis in normal, x-irradiated, and friend virus infected mouse spleen nuclei, 26:45849
 FIELD, Lamar, Organic disulfides and related substances. 33. Sodium 4-(2-acetamidoethylthio)butanesulfinate and related compounds as antiradiation drugs, 26:45290
 FIELD, R. Clive, Range-momentum correlation in carbon and iron for muons of 500 to 1500 MeV/c, 26:46448 (LBL-791)
 FINK, R. W., Gamma branching and total conversion coefficients of intraband-transitions in groundstate rotational band of ²⁴⁵Cm, 26:47034
 FINKELSTEIN, J., Some implications of a violation of the Pomeranchuk theorem, 26:46634
 FINNEY, C. D., Third-derivative method for determining electron-impact onset potentials, 26:46385
 FIRK, F. W. K., Total neutron cross section measurements, 26:46967
 FISCHER, E., Development of methods for application of neutron activation analysis to determination of trace elements in foods. Progress report, 10-12/71, 26:45189 (BFL-1/1972)
 FISCHER, W., Optical determination of the quadrupole moment of ¹³⁸La, 26:46950
 FISENKO, A. V., Miniature low-background counters for measurement of ³⁷Ar and ³⁹Ar, 26:45614
 FISH, Birney R., Measurement of liquid droplet emissions from cooling towers and process stacks, 26:45574 (CONF-720619-1)
 FISHBONE, Leslie G., Relativistic Roche problem, 26:46189
 FISHER, S. B., Application of high-voltage electron microscopy to void studies, 26:46056 (CONF-710601-, pp 533-49)
 FITZER, E., Research and development on graphite in the federal republic of germany, 26:46086
 FITZGIBBON, F. J., Cells for nondestructive examinations of irradiated fuel pins, 26:46104 (LA-DC-72-556)
 FITZWILSON, R. L., Angular distribution of metastable hydrogen formed by electron capture in a helium target, 26:46397
 FITZWILSON, Roger Louis, Angular distribution of H(2s) formed by electron capture and dissociation collisions, 26:46388
 FLAMINI, Mario, Phase sensitive method for detecting superconductive transitions in the microwave region, 26:46840
 FLEISCHER, J., Pion form factor in Bethe-Salpeter model, 26:46615
 FLEISCHMAN, R. M., Three dimensional neutronics calculations for the Fast Test Reactor (FTR) and the FTR Engineering Mock Up Critical assembly (EMC), 26:47586 (HEDL-TME-72-42)
 FLEMING, Gordon N., Spin spectrum of an unstable particle, 26:46498
 FLICHE, Henri-Hugues, Model for particles with spin in the gravitational and electromagnetic field, 26:46767
 FLIEDNER, T. M., Role of transplanted slowly proliferating bone marrow cells for regeneration of lethally x-irradiated rat bone marrow, 26:45837
 FLOBERG, Allan, Temperature dependence of the isotope effect for electromigration of rubidium ions in molten rubidium nitrate, 26:45294
 FLORES, F., Magnetohydrodynamic entrance flow in a channel, 26:46423
 FLUEKIGER, R., Identification of the superconducting phases in the molybdenum-platinum system, 26:46828
 FOELDES, J., New two-channel scanner. II, 26:45607
 FOGEL, Ya. M., Ion-ion emission: new tool for mass-spectrometric investigations, 26:47265
 FOGLIA, C., Compound atom state in neon, 26:46383
 FOIN, C., Rotational levels with high spins in ¹⁷¹Lu, 26:46980
 Two branchings in rotational band 1/2- [541] in ¹⁷¹Lu, 26:46987
 FOLDIAK, Gabor, Radio-frequency alcoholic chlorobenzene dosimeter, 26:45632
 FOLSOM, T. R., Plutonium and polonium inside giant brown algae, 26:45924
 FOMENKO, G. P., Electron acceleration in an undulator synchrotron, 26:46120
 FOMINYKH, M. I., Study of the decay ¹⁴⁰Tb(T/sub 1/2/ = 66 min) and ¹⁵⁰Tb(T/sub 1/2/ = 3,48 +- 0,16 h), 26:46942 (JINR-P6-6512)
 FONER, S., dc anisotropic magnetization at 20.4 K and the ideal crystal-field-only behavior of TmSb, 26:47258
 FOOTE, James H., Nonadiabatic energy limit versus mirror ratio in magnetic-well geometry, 26:47106
 FOOTE, R. S., Measurement of radioactive fallout by field total absorption gamma ray spectroscopy. Report No. 3, 26:45527 (TID-26103)
 FORAPONTOV, N. V., Hard gamma radiation from ²³⁸Pu, 26:47032
 FORD, John Charles, Formation of excited hydrogen atoms by charge-transfer neutralization of high-energy protons,

- 26:46378
- FORD, P. J., Superconductivity and localized spin fluctuations in concentrated aluminum--3d-transition-metal alloys, 26:46838
- FORMAN, P. R., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)
- FORSLOW, E. J., Distribution of radioactivity in and near the Rainier rubble chimney, 26:45483 (NVO-1229-180)
- FORTNEY, L. R., Progress report [on high-energy physics], July 1, 1971--June 30, 1972, 26:46479 (ORO-3065-24)
- FOSSAN, D. B., Magnetic moment of the 2761 keV 8^+ state in ^{92}Mo , 26:46959
- FOSTER, Bessie R., Lymphoid-cell renewal under low-level irradiation. XI. The cell cycle and growth kinetics during radiation leukemogenesis in C57BL mice, 26:45822
- FOSTER, J. C. Jr., Tachyon dust metric in general relativity, 26:46754
- FOTOPOULOS, C. U., High-temperature thermal insulation. Quarterly report, FY 70-3, 26:47601 (BDX-613-288(Rev.))
- FOULETIER, Bernadette, Radiorecovery effect of yeast extract on carnation cuttings irradiated with ^{60}Co γ rays, 26:45777
- FOURNELLE, Raymond Albert, Genesis of cellular precipitation in copper rich copper--indium alloys, 26:46017
- FOX, J. D., Measurement of the magnetic moment of the antiproton, 26:46440
- FRAAS, A. P., Design of high-temperature liquid metal systems, 26:45437 (CONF-720904-1)
- FRABETTI, P. L., Decay mode $\phi_1 + \eta$ and measurement of branching ratio, 26:46435
- FRADKIN, M. I., Removable current lead for power supply in superconducting magnetic systems, 26:45679
- FRAGA, S., Electric dipole polarizabilities of lanthanides and actinides, 26:46349
- FRANCE, D. M., Improved point-matching techniques applied to multi-region heat transfer problems, 26:45465
- FRANGYAN, A. A., Detector for high energy particles based on the principle of using particle radiation in a laminated medium, 26:45648
- FRANK, L. A., ELF noise bands associated with auroral-electron precipitation, 26:46308
- FRANKEVICH, E. L., Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. I, 26:45331
- FRANK-KAMENETSKII, D. A., Plasma: the fourth state of matter, 26:47104
- FRANTSEVICH, I. N., Energy spectrum and crystalline structure of wurtzite-like modification of boron nitride, 26:47255
- FRASER-SMITH, A. C., Relationship between Pc 1 micropulsations and ionospheric spread F, 26:46274
- FRATARCANGELI, S., Measurement of ^{137}Cs and ^{90}Sr in soil, 26:45263 (RT/PROT-72-11)
- FREDERICK, E. J., Burial of radioactive waste in salt, 26:46100 (CONF-720903-4)
- FREDRICKS, R. W., Detection of solar-wind electron plasma frequency fluctuations in an oblique nonlinear magnetohydrodynamic wave, 26:46329
- FREEMAN, Gordon R., Effect of pressure on the rates of reaction of solvated electrons in liquid ethanol, 26:45351
- FREMION, Lucien, Double electromagnetic valve for simultaneously controlling dangerous fluids and vacuums, 26:46109
- FRENTROP, Arthur H., Pulse neutron logging tool control and transmission system, 26:45563
- FREUNDLICH, W., Phases of scheelite structure in the systems of plutonium molybdate--lithium or sodium molybdate, 26:45954
- FRIED, Bernard, Infectivity and development of x-irradiated metacercariae of *Leucochloridiomorpha constantiae* (Trematoda) in the chick and on the chorioallantois, 26:45879
- FRIED, W., Effect of various cell-free organ extracts on hematopoietic stem cells, 26:45847
- FRIEDMAN, K. A., Photonlike particles, Compton scattering, and sum rules in the dual-resonance model, 26:46601
- FRIEL, Joseph V., Entrainment moisture separators for fine (1-10 μ) water--air--steam service: their performance, development, and status. Final report, 26:45440 (MSAR-71-45)
- FRIES, D., Diffractive fragmentation in high energy photon reactions, 26:46437
- FRIIS-CHRISTENSEN, E., Critical component of the interplanetary magnetic field responsible for large geomagnetic effects in the polar cap, 26:46266
- FRINDT, R. F., Discontinuous flux flow in superconducting NbSe_2 , 26:46831
- FRISHMAN, Yitzhak, Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)
- FRITZ, J. S., Spectrophotometric molybdenum determination with thiolactic acid, 26:45239
- FROELICH, A., K--p interactions at 10 GeV/c, 26:46480
- FROLOVA, A. V., Importance of tissue-equivalent detectors used in radiometry, 26:45619
- FROMMHOLD, Lothar, Differential- and total-cross-section calculations for models of e--H₂ collisions, 26:46399
- FRULLANI, Salvatore, Criteria for selecting the location of nuclear plants, 26:47574
- FRY, Bill W., Fluorine metabolite concentrations and renal clearances following methoxyflurane anesthesia, 26:47645 (UR-3490-107)
- FRYER, Michael Owen, Consequences of microscopic symmetries and linear response theory for the hydrodynamics of superfluid helium, 26:46796
- FUJII, Kyoichi, Acute death of mice during massive gamma irradiation. III, 26:45846
- FUJII, Shigetada, Radiation-induced copolymerization of vinylphosphonic dichloride with methyl methacrylate and styrene, 26:45318
- FUJISHIRO, Masatoshi, Intensity ratio of 2158 keV and 2505 keV gamma-rays in decay of ^{60}Co , 26:46933
- FUJITA, Yoshiaki, Slow neutron spectra in the liquid and solid methane, 26:46858
- FUJIWARA, Masato, Active species of γ -irradiated vinyl acetate and their changes on irradiation of visible light, 26:45334
- FUJIWARA, Shizuo, Nuclear magnetic relaxation study of formic acid. Isotope effect and ^{13}C relaxation, 26:45297
- FUKASAWA, Tsutomu, Determination of small amounts of molybdenum based on its catalytic effect in Methylene Blue--hydrazine sulfate reaction, 26:45227
- FUKUGITA, M., Analysis of kaon--nucleon scattering amplitudes in the intermediate energy region, 26:46624
- FUKUI, Kenichi, Long-range interactions between two hydrogen atoms and two helium atoms, 26:46351
- FUKUI, S., Asymmetry for π^+ photoproduction from polarized protons between 300 and 900 MeV at 90°, 26:46444
- FUKUOKA, Yutaka, Recombination domain instability in high-energy electron-irradiated germanium, 26:47230
- FULDE, Peter, Some crystalline field effects in metals, 26:47269
- FULLER, C. T., Oxidation and etching of 80 percent Si--20 percent Ge, 26:45935 (SC-RR-710884)
- FULTON, E. J., Void formation in cladding and structural materials irradiated in DFR, 26:46047 (CONF-710601-, pp 125-41)
- FURLAN, P., Possible origin of finite quantum electrodynamics from weak interactions, 26:46722
- FURNICA, M., Sensitization of living organisms by chemicals under effect of radiations and cytostatics. III, 26:45732
- FURUICHI, Susumu, Pion form factor and possible heavy vector meson, 26:46561
- FURUKAWA, Seijiro, Mass dependence of critical amorphizing dose in ion implantation, 26:47236
- FURUTA, Junichiro, Dose-rate measurement by several Radocon probes. II, 26:45620
- Intensity ratio of 2158 keV and 2505 keV gamma-rays in decay of ^{60}Co , 26:46933
- FUSHCHICH, V. I., Three types of relativistic equations for particles with nonzero mass, 26:46730
- FUTTERMENGER, W., Comparison of the background spectra in eight german whole-body counters and a model structure with layered shield, 26:45590 (ORNL-tr-2532)

G

- GABE, D. R., Structure of intermediate layers in diffusion coatings, 26:46041
- GABRASHVILI, A. I., Gamma quanta distribution at large distances from the sources, 26:47215
- GABRIELSSON, Ann-Britt, Nanogram determination of iodide by means of isotope exchange, 26:45257
- GADZHOKOV, V., ^{175}Hf states excited during decay of ^{175}Ta , 26:46979
- GAFFAROV, D. G., Magnetic properties of ^{153}Eu , ^{154}Eu , and ^{159}Gd , 26:46981
- GAGE, D. W., Duality-breaking effects in a model for meson--baryon scattering, 26:46642
- GAGLIONE, P., Site characteristics, 26:45570 (EUR-4841)
- GALAKHMATOVA, B. S., Mechanism of $^9\text{Be}(p,p_1)^9\text{Be}$ 1.68 MeV reaction, 26:46885
- GALAN, P., Structure of ground states of odd--odd nuclei ^{160}Tb , ^{162}Tb , ^{162}Re , and of isomer /sup 182m/Re, 26:46983
- GALATOLA, Anthony, Preliminary analysis of the atmospheric eclipse of 32 Cygni, 26:46184
- GALBRAITH, J. F., High-density braid cable shield, FY72-1. Quarterly report, 26:47606 (BDX-613-543(Rev.))
- GALIN, J., Heavy ion nuclear reactions, 26:47056
- GALLONE, F., Casualty group in finite space--time, 26:46724
- GAMALII, A. F., γ -emission in radiative capture of intermediate neutrons, 26:46917 (BNL-tr-477)
- GAMBINO, R. J., Nuclear magnetic resonance in the LaX_3 intermetallic compounds, 26:47281
- GAMMON, R. H., Ten-micron spectroscopy of circumstellar shells, 26:46174
- GANAS, P. S., Scattering by an r/sup -2n/ potential in the Born approximation and application to dipole--quadrupole scattering, 26:46380
- GANDINO, C., Site characteristics, 26:45570 (EUR-4841)
- GANEVA, A., Sensitization of living organisms by chemicals under effect of radiations and cytostatics. III, 26:45732

GANGRSKII, Yu. P., Angular distribution of fragments of spontaneously fissioning isomers, 26:47022 (JINR-P7-6466)

GANIVET, Michel, Equipment for concentrating radioactive solutions, 26:46111

GANOVICI, I., Bubble density determination in bubble chambers, 26:45649

GANZHULA, N. N., Separation processes in superconducting alloys of niobium--aluminum, 26:46020

GARAULT, Yves, Orthogonality relations in periodic waveguides having anisotropic surface impedance, 26:46141

GARCIA, W. J. Jr., Method for estimating heights of water waves generated by underwater cratering explosions, 26:45486 (UCRL-51235)

GARCIA-BELLIDO, A., Clonal parameters of tergite development in *Drosophila*, 26:45881

GARDIER, S., Early data from ultraviolet sky-scan telescope in TD1 satellite, 26:46186

GARDNER, Karl A., Sodium heated steam generator, 26:45450

GARFINKEL, A. F., Final states with three charged particles and a visible Λ from K-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)

Observation of the $M(940)$ and $\delta(963)$, 26:46475 (IS-2886)

GARI, M., Circular polarization of γ rays in ^{18}F , 26:46877

GARKUSHENKO, T. L., Synthesis and study of lanthanide peroxides, 26:46085

GARMIRE, E., Optical waveguiding in proton-implanted GaAs, 26:47232

GARNER, F. A., West Valley Reprocessing Plant. Letters of May 23, June 26, 27, 30, and July 5, 1972, with respect to modifications in the plant, 26:46065 (CONF-710601-, pp 841-52)

GARRETT, J. D., Subbarrier fission resonances in Th isotopes, 26:47033

GARRETT, Thomas B., Theoretical studies of diamagnetic properties of the hydrogen molecule ion. II. Effect of varying internuclear separation, 26:46354

Theoretical studies of diamagnetic properties of the hydrogen molecule ion. I. Approximate variation--perturbation calculation, 26:46353

GARRETT, W. R., Slow electrons from electron-impact ionization of He, Ne, and Ar, 26:46390

GARUSOV, E. A., Green's function of neutron transport equation for moving medium, 26:46650

GARVIE, Ronald C., Phase analysis in zirconia systems, 26:45945

GATAULLIN, R., Use of quasi-classical approximation for solution of system of Hartree--Fock equation, 26:47288

GATES, Olive, Radiation carcinogenesis. Progress report VI, May 1, 1971--April 30, 1972, 26:45814 (COO-3017-43)

GATROUSIS, C., Environmental levels of radioactivity in the vicinity of the Lawrence Livermore Laboratory January through December 1971, 26:45513 (UCRL-51242)

GAULT, F. D., Kinematics of coupling of Regge poles to photons, 26:46560

GAUSSENS, Gilbert, Method for polymerization and/or cross-linking of acrylic polymers or copolymers, 26:45339

GAUSTAD, J. E., Ten-micron spectroscopy of circumstellar shells, 26:46174

GAUVIN, H., Transfer reactions induced by ^{86}Kr on ^{232}Th , 26:47035

GAVRILYUK, V. I., Decay of ^{196}Au , 26:47003

Electron conversion spectrum of ^{182}Re , 26:46978

GAZSO, L., Determination of microbial contamination of disposable medical devices and the estimation of radiosensitivity of isolated microflora, 26:45748

GEER, K., Nature of absorption, 26:46623

GEHRZ, R. D., Core of eta Carinae, 26:46160

GEHSE, M., Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)

GEIER, H., Special machines and equipment of WAK, 26:46106 (ORNL-tr-2372)

GEIGER, J. S., Deorientation of nuclei of recoil atoms, 26:46935

GEL'D, P. V., Magnetic susceptibility of cubic niobium and tantalum carbides in the low temperature range, 26:47266

GELLER, Elaine N., Nature of two-particle correlations in atoms, 26:46414

GELLER, Murray, Nature of two-particle correlations in atoms, 26:46414

GEMMELL, D. S., Channeling of protons in thin BaTiO_3 crystals at temperatures above and below the ferroelectric curie point, 26:47234

GERARD, N., New reading system for an ionographic detector, 26:45598

GEROCH, Robert, Structure of the gravitational field at spatial infinity, 26:46752

GERSCHEL, C., Measurement of the quadrupole moment of a shape isomer, 26:46960

GERSTENBERGER, R. V., Magnetic moments of ^3He and ^3H , 26:46867

GERVE, A., Important wear measuring methods using radionuclides, 26:45711

Use of radionuclides for measurements in mechanical engineering, 26:45710

GETMANOV, V. B., Intranuclear cascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)

GHIRARDI, G. C., Algebraic structure of a class of solvable quantum problems, 26:46671

27 AUTHORS

GOMBOS

Hulthen and Schwarzschild potentials in the Klein--Gordon equation, 26:46690

GHYSELS, J. P., Determination of Pu in the environment by extraction and liquid scintillation counting, 26:45268

GIBBY, R. L., Effects of pore structure on the melting heat rating of oxide fuels, 26:45930 (HEDL-TME-72-81)

GIBERT, R. J., Study of pressure fluctuations generated by aerodynamic singularities in a circuit, 26:47310

GIBSON, R., Irradiation in the epidemiology of leukemia among adults, 26:45798

GIDALY, M., AET on immunologic processes in irradiated rabbits, 26:45824

GIGNOUX, C., Method for solving Faddeev equations, 26:47065

GILBERT, Ernest, Fourier transform of inelastic neutron scattered intensity for solid ^4He , 26:46845 (BNL-16935)

GILBERT, J. A., Polarization of radio sources at $\lambda 49\text{ CM}$ and $\lambda 73\text{ CM}$, 26:46204

GINESTET, J., Further results from multi-Regge models applied to $pp + pp \rightarrow \pi^+ + \pi^-$ reaction at 13.1 GeV/c, 26:46656

GINGO, Peter J., Neutron radiation characteristics of plutonium dioxide fuel, 26:45704

GINSBERG, T., Improved point-matching techniques applied to multi-region heat transfer problems, 26:45465

GINTNER, Henry, Study of the continuum states of the ^7Be system below 10 MeV, 26:46896

GIORDANO, Anthony Michael Jr., Chronic low-level NO_2 exposure and the mucociliary clearance mechanism in the rat trachea in vivo, 26:47644 (UR-3490-85)

GIPSH, A. A., Distribution and genesis of radioactivity in coals of the Pechora River basin, 26:45505

GIRARD, B., Inclusive reaction $\mu + \text{nucleon} \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV/c, 26:46452

GIRAUD, B., Generator-coordinate form factors for transfer reactions between heavy ions, 26:47069

GIZON, A., Two branchings in rotational band $1/2^- [541]$ in ^{171}Lu , 26:46987

GLABERSON, W. I., Quantized vortex ring dynamics and the vortex core radius in He II, 26:46800

GLAD'SHEV, D. A., ^{135}Ba levels, 26:46945

GLASSER, Herman, Holder for radioactive material, 26:46113

GLAZMAN, Yu. M., Radiometric investigation of the absorption of potential-determining ions from aqueous solutions of non-ionogenic surface-active substances, 26:45279

GLEW, D. N., Deuterium separation process, 26:45690

GLINSKA, A. F., Relative level populations in foil excited He and Li, 26:46337

GLINSKA, Halina, Distribution of doses in patients with Hodgkin's disease during conventional radiotherapy, 26:45786 (TT-54002/5, pp 590-3)

GLOWIAK, Bohdan J., Dependence between radioactivity and concentration of atmospheric aerosols, 26:45521

GLUSHKOVSKII, M. E., Calculation of the dead time of high-speed amplitude converters, 26:45684

GODAWSKI, Jerzy, Prognostic elements in inoperable carcinoma of the breast, 26:45783 (TT-54002/4, pp 451-5)

Selected prognostic elements in carcinoma of the larynx, 26:45781 (TT-54002/4, pp 437-41)

GODDARD III, W. A., Origin of energy barriers in excited states of He_2 , 26:46346

GOEKE, K., Orbital rearrangement in light nuclei, 26:46886

GOERLLER-WALRAND, C., Coupling scheme in uranyl complexes, 26:45380

GOFMAN, John W., Existence of nuclear weapons: a prime environmental threat, 26:45492

GOFORTH, Robert Ronda, Structure of the plasma-wave echo, 26:47195

GOKHALE, K. V. G. K., Kinetics and burst phenomenon in ZrO_2 transformation, 26:45947

GOLDBACH, C., Standard black body at 10,000°K (high pressure plasma arc), 26:47217

GOLDBACH, Claudine, Preliminary results obtained from the study of an argon plasma continuum in the near ultraviolet and visible ranges at high pressures, 26:47132

GOLDEN, D. E., Observation of a broad resonance in the 2^3S excitation of helium by electron impact, 26:46405

GOLOSKOKOV, S. V., Asymptotic estimates of quasipotential parameters, 26:46703 (JINR-P2-6482)

Investigation of large angle scattering in quasipotential approach, 26:46776 (JINR-P2-6442)

GOLOVKOV, N. A., α -decay of neutron-deficient isotopes of Ra, 26:47004

GOLUB, T. V., Influence of impurities and temperature on the deformation of molybdenum during dynamic loading, 26:46043

Orientation dependence of damping of elastic vibrations in tungsten single crystals, 26:46022

GOLUBCHINA, M. N., Use of admission system of MI-1309 instrument for precision isotopic analysis, 26:45681

GOLUBTSOV, I. V., Study of zirconium evaporation from open surface, 26:46011

GOLYAND, L. N., Dynamics of control system incorporating 440 MW(e) reactor, 26:47324

GOMBOS, P., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian

- Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- GOMES, A. A., Gadolinium g-shifts and band structure, 26:47270
- Remarks on Gd g-shifts, 26:47271
- GONCHAROV, N. F., Separate control of the injector heating and of the current amplitude in transformer accelerators, 26:46135
- GONDHALEKAR, P. M., Photodetachment of electrons from major negative ions in lower D region, 26:46299
- GOODINGS, J. M., Plasma-sampling diagnostics using ion kinetic energy distributions, 26:47128
- GORBATSEVICH, S. V., Review and analysis of the latest data on measuring fundamental physical constants, 26:46439
- GORCZYCA, B., Comparison of the angular decay distributions in three quark models of photoproduction, 26:46569
- Quark model predictions for angular decay distributions of photoproduced vector mesons in inclusive processes, 26:46536
- GORCZYCA, Barbara, Comparison of the angular decay distributions in three quark models of photoproduction, 26:46526 (TPJU-14/71)
- GORELKIN, D. S., Calorimetric study of Nb-Fe-Si-Al system, 26:46007
- GORGAN, V., AET on immunologic processes in irradiated rabbits, 26:45824
- GORINI, V., Casuality group in finite space--time, 26:46724
- GORINI, Vittorio, Structure of relativity groups, 26:46669
- GORSKI, L., Determination of ash content of coal by neutron activation analysis, 26:45195
- GORSKI, A. G., Identification of an AmCl_2^{3+} octahedral complex from Am(III) absorption spectrum in ethanol solutions of HCl, 26:45376
- GORYACHEV, B. I., New experimental data on the fine structure of dipole resonance, 26:46890
- GORYAEV, V. M., Radiation damage to polymer fiber investigated by small-angle x-ray scattering, 26:45338
- GOSEK, B., Effect of radioprotective preparations on the respiratory exchange of rats, 26:45823
- GOSHEN, S., Algebraic derivation of generalized Talmi expansion, 26:46503
- GOUGH, R. A., Confirmed proton radioactivity of ^{53}mCo , 26:46926
- Further results on proton radioactivity of ^{53}mCo , 26:46919
- GOULDING, C. A., MeV total neutron cross sections of ^{238}U , 26:47013 (COO-3058-23)
- GOVE, N. B., Log-f tables for beta decay, 26:46970
- GRABOWSKI, D., Long-term study of urinary ^{90}Sr and calcium in children, 26:45908
- GRABOWSKI, J., Parity nonconserving effects in nuclei ^{175}Lu and ^{181}Ta , 26:46982
- GRABSKA, A., Effect of x irradiation on cells, 26:45738
- GRAEME-BARBER, C., Tubular Nb/Cu conductors for ac superconductive power transmission, 26:46844
- GRAHAM, Charles D. Jr., Superconductive magnets, 26:46843
- GRAHAM, E. D. Jr., Device degradation from the effects of nuclear radiation on passivation materials, 26:45662 (SC-DC-721232)
- Radiation effects in microwave bipolar transistors, 26:45663 (SC-DC-721233)
- GRAHAM, J., X-ray photoelectron spectra of uranium oxides, 26:45367
- GRAHAM, R. L., Deorientation of nuclei of recoil atoms, 26:46935
- GRAHAM, S., Irradiation in the epidemiology of leukemia among adults, 26:45798
- GRALL, L., Effect of heat treatments on corrosion of 18/10 stainless steels with low C contents in nitric medium in presence of ferric iron, 26:45970
- GRANADOS, C. E., New approach to beta-gamma coincidence counting. Advance report on SAMAR electronic system, 26:45589 (JEN-247)
- GRAUL, E. H., Microdosimetry measurements of fast neutrons with a tissue-equivalent proportional counter of 1 cm diameter, 26:45626
- GRAY, W. J. H., DNA synthesis in gamma-irradiated recombination deficient strains of *E. coli*, 26:45753
- GREBEN, J. M., S-matrix pole trajectories in the N/D approach, 26:46789
- GREBENKINA, V. G., Carbide resistors, 26:47250
- GRECO, M., Nonet structure of $J/\text{sup PC} = 3^-$ mesons, 26:46611
- GREEN, G. F., Respiratory quotients of irradiated fruits, 26:45729
- GREEN, M. H. L., DNA synthesis in gamma-irradiated recombination deficient strains of *E. coli*, 26:45753
- GREEN, R. M., Development of elastomer cushions, FY69-4. Quarterly report, 26:47603 (BDX-613-440(Rev.))
- GREEN, William R., Irradiation apparatus for providing a high initial irradiation of the product, 26:45457
- GREENHUT, G. K., Meson photoproduction in the backward direction, 26:46529
- GREENSTADT, E. W., Analysis of Pioneer 9 low-frequency wave observations near interplanetary discontinuities, 26:46302
- GREENWAG, Gilbert S., Effects of x-irradiation on ovarian function in the pregnant hamster, 26:45825
- GREGERS-HANSEN, P. E., Sinusoidal current-phase relation for superconducting thin-film microbridges, 26:46806
- GREGORIO, M. A., Recoil effect in quasi-elastic hadron--deuteron scattering, 26:46653
- GREGORY, Derek P., New concept in energy transmission, 26:45429
- GREISEN, Kenneth, Detection of large cosmic-ray showers by atmospheric fluorescence. Final report, 26:46150 (NYO-3038-13)
- GRENON, Michel, Atomic energy in India, 26:47300
- Evaluate European siting for next enrichment plant, 26:45693
- GRESKOVICH, C., Preparation of transparent Y_2O_3 -doped ThO_2 , 26:45948
- GRIBBLE, R. F., Feedback stabilization on an $l = 1$ theta-pinch plasma column, 26:47157 (LA-DC-72-707)
- GRIBKOV, V. A., Observation of second compression in the final stage of a discharge of the plasma focus type, 26:47139
- GRIFFIN, H. C., Hindered $7/2^+ \rightarrow 3/2^-$ β -decay of ^{143}Pr , 26:46955
- GRIFFIN, J. F., Documentation of DOT-SP-5915 for 55/30 and 83/55 drum combination packaging, 26:46114 (MLM-1911)
- GRIFFIOEN, Roger D., Low-lying states and configurations in ^{138}La , 26:46964
- GRIGOR'EV, E. A., Reaction $(\pi^-, \pi^+ p)$ on ^6Li and ^{12}C , 26:46869
- GRIGOR'EV, Yu. N., Effect of the collective interaction of particles on the longitudinal dimensions of an electron bundle in an accumulator, 26:46546
- GRIGOROV, N. L., Collective interactions of nucleons of high-energy cosmic heavy nuclei, 26:46461
- GRIMM, R. C., Improvements on a new method for the numerical solution of the Schrodinger equation, 26:47290
- Viscosity and thermal conductivity effects on the containment of a toroidally confined resistive plasma with flow, 26:47105
- GRINEVA, L. P., 1,6-glucodialdose as one of radiolysis products of D-glucose-6-phosphate, 26:45337
- GRISHAEV, I. A., Angular and energy distributions of protons in (γp) and (πp) reactions on ^{12}C nucleus for 130 MeV excitation energy, 26:46888
- Decrease of the off-duty factor in the operation of a linear accelerator, 26:46145
- Effect of the collective interaction of particles on the longitudinal dimensions of an electron bundle in an accumulator, 26:46546
- Peculiarities in interaction of high-energy electrons and positrons with crystals, 26:46550
- Photoproduction of charged pions by nuclei at $E_0 = 1160$ MeV, 26:46441
- GRISHCHUK, L. P., Homogeneous rotating universe with flat space, 26:46261
- GRISHIN, V. G., Spectrum symmetry of pion longitudinal momenta in $\pi^+ p$ interactions at $p_{\text{c}} = 4$ GeV, 26:46619 (JINR-P1-6461)
- GRISHIN, V. K., Linear stability of a charged beam in an induction system, 26:46138
- GRISOM, J. T., Anode temperatures and ion energy distributions during a vacuum arc discharge, 26:47625 (SC-DC-722267)
- Slow electrons from electron-impact ionization of He, Ne, and Ar, 26:46390
- GRIWATZ, Gunther H., Entrainment moisture separators for fine (1-10 μ) water--air--steam service: their performance, development, and status. Final report, 26:45440 (MSAR-71-45)
- GROMOV, K. Ya., Study of the decay $^{148}\text{Tb}(T_{1/2} = 66 \text{ min})$ and $^{150}\text{Tb}(T_{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- GROMOVA, I. I., Study of the decay $^{148}\text{Tb}(T_{1/2} = 66 \text{ min})$ and $^{150}\text{Tb}(T_{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- GROS, G., Study of the mechanical and thermal behavior of a high temperature reactor core, 26:47304
- GROSS, S. H., Collisional losses in a very-low frequency duct associated with the lower-hybrid-resonance frequency, 26:46272
- GROSSER, D., Crossing and unitarity in context of Bethe--Salpeter equation, 26:46727
- GROSSMAN, Lawrence, Molecular basis for the mutagenic and lethal effects of ultraviolet irradiation. Progress report, June 16, 1971--May 31, 1972, 26:45722 (COO-3232-1)
- GROTDAL, T., Energy levels in ^{231}Th and ^{233}Th , 26:47029
- GROTH, F. M., Evaluation of instrumentation for nuclear fuels reprocessing plant input weight measurements, 26:46093 (ICP-1014)
- GROVE, E. L., Analytical emission spectroscopy. Part I, 26:45252
- Analytical emission spectroscopy. Part II, 26:45253
- GRUBER, Jack, Immunogenicity of purified Venezuelan equine encephalitis virus inactivated by ionizing radiation, 26:45752
- GRUENBERG, E., Gamma gamma directional correlation measurements in ^{131}Cs , 26:46963
- GRUENEWALD, Th., Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)
- GRUNDON, Noel J., Improved method for determination of ^{35}S in plant material using oxygen flask combustion and liquid scintillation counting, 26:45266
- GRUPEN, C., Stopping particles underground, 26:46157
- GRYZLOV, A. V., Particle acceleration in a H_{111} cylindrical cavity in an axial magnetic field, 26:46139

- GUASP, Jose, Green function in the secondary model of thermalization, 26:46846 (JEN-221)
- GUBERMAN, S. L., Origin of energy barriers in excited states of He_2 , 26:46346
- GUDIKSEN, P. H., Environmental levels of radioactivity in the vicinity of the Lawrence Livermore Laboratory January through December 1971, 26:45513 (UCRL-51242)
- GUDIMA, K. K., Account of nuclear boundary diffusivity in the intranuclear cascade model, 26:47047 (JINR-P2-6503)
- Change in the mechanism of inelastic interactions between particles and nuclei in the energy region T approximately equal to 5 GeV, 26:47038 (BNL-tr-487)
- GUERREAU, D., Heavy ion nuclear reactions, 26:47056
- GUGNIN, A. A., Electrical and thermal conductivities of Re, Os, and Ir and some of their alloys with Th and La in wide temperature range, 26:47248
- GUGUSHVILI, E. I., Free Riemann--Green's function of inverse problem of scattering theory, 26:46785
- Inverse problem of scattering theory and Riemann's method, 26:46788
- GUHA, Evelyn, Inversion from the energy dependence of total cross sections, 26:46387
- GUHATHAKURTA, B. K., Large scale periodic disturbances in the ionosphere, 26:46290
- GUIDO, M., Thermodynamics of rare-earth--carbon systems. IV. The lutetium--carbon system, 26:45961
- GUIHO, J. P., National standards for photon dosimetry, 26:47216
- GUILLE, B., Method of determining mean energy of secondary electrons in electron equilibrium regions using theoretical x-ray spectra (10 to 20 MeV) and ionization measurements in different media, 26:47213
- GUIMARAES, A. P., Gadolinium g-shifts and band structure, 26:47270
- Remarks on Gd g-shifts, 26:47271
- GUISAN, O., Polarization in charge exchange reactions, 26:46491
- GUL'KO, V. A., Temperature dependence of cross section of cold- and thermal-neutron scattering by O_2 and D_2 , 26:46887
- GULYAKIN, I. V., Accumulation of ^{90}Sr in wheat on sod-podzolic soils, 26:45550
- GUNNINK, R., Computerized quantitative analysis by gamma-ray spectrometry. Volume I. Description of the GAMANAL program, 26:45265 (UCRL-51061(Vol.1))
- GUPTA, Anurag B., Small-scale fluctuations in F-region critical frequency, 26:46289
- GUPTA, B. L., Estimation of sulphur content in human hair and nail for use in fast neutron dosimetry, 26:45788
- GURCHUMELIYA, A. D., Statistical consideration of retardation in relativistic Hartree--Fock equations, 26:46408
- GUREVICH, M. E., Change in linear dimensions of metals under the influence of γ radiation, 26:46075
- GURNETT, D. A., ELF noise bands associated with auroral-electron precipitation, 26:46308
- GURNEY, C. W., Hematopoietic and immunological alterations in mice produced by ^{89}Sr -induced marrow hypoplasia, 26:45896
- GURSKY, Herbert, Association of x-ray sources with bright stars, 26:46212
- GUSEV, V. M., Investigation of the injection characteristics of symmetrical pin-diodes prepared by the method of ion implantation, 26:45678
- GUSTAFSSON, B., Variations in the bone marrow lymphoid population in the sublethally irradiated germ free mouse, 26:45843
- GUTAY, L. J., Final states with three charged particles and a visible A from K-d interactions at 4.5 GeV/c, 26:46474 (COO-1428-306)
- GUTBROD, F., Pion form factor in Bethe--Salpeter model, 26:46615
- GUTOWSKI, Ryszard, Hydrogen generators for neutron tubes, 26:45942
- GUYON, E., Persistent current states in rotating superfluid helium, 26:46801
- GWAL, A. K., Reflection of microwave through laboratory plasma, 26:47118
- GWYN, C. W., Carrier trapping effects in irradiated pn junctions, 26:45664 (SC-DC-721237)
- Radiation effects in microwave bipolar transistors, 26:45663 (SC-DC-721233)
- H**
- HAAS, R. J., Role of transplanted slowly proliferating bone marrow cells for regeneration of lethally x-irradiated rat bone marrow, 26:45837
- HABENICHT, W., Isobars from the spallation reaction for nuclear spectroscopy, 26:46985
- HABERLER, P. L. F., Does the simple quark light-cone current algebra of Fritsch and Geil-Mann imply that quarks do exist, 26:46604
- HABERSTICH, A., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)
- HABERSTROH, R. A., Direct measurement of the nuclear magnetic dipole moment of ^{163}Ho , 26:46990
- HACKSTEIN, K., Process for recovering by-products from uranium dioxide production, 26:45371
- HAEUSSER, O., Lifetime of 975-keV, $3/2^+$ state in ^{25}Mg , 26:46908
- HAGEDOORN, H. L., Cyclotron laboratory at Technical University of Eindhoven, 26:46146
- HAGE-HASSAN, M., Generating function method and developments in quasi-boson, 26:47286 (LYCEN-7202)
- HAHN, Bo-Sup, Radiation chemistry of nucleic acids. Isolation and characterization of thymine glycols, 26:45314 (COO-3286-1)
- HAHN, Ernest A., Crosslinked polyvinyl chloride or polyvinylidene fluoride film, 26:45352
- HAHN, W. C. Jr., Forming of composite for superconducting rod. Technical progress report, 26:45972 (COO-3311-1)
- HAIDT, D., Particle production in proton interactions in nuclei at 24 GeV/c, 26:46459
- HAJJ, Jack P., Multipole radiation solutions for a class of massive particle wave equations, 26:46709
- HALAMA, H. J., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)
- HALDORSEN, J., Enhancement at about 2700 MeV/c² in the $\text{pp} \rightarrow \text{X}^0$ system in $\text{pp} \rightarrow \text{pp} + \text{X}^0$ at 19 GeV/c, 26:46467
- HALE, W. H. Jr., Conversion of Cf α -hydroxyisobutyrate to oxide with carboxylic acid resin, 26:45393
- HALL, Larry L., Procedure for chronic intravenous catheterization in the rabbit, 26:47641 (UR-3490-21)
- HALL, Roger P., Adhering resins to substrates, especially metal, by radiation, 26:46089
- HALPERN, I., Very inelastic scattering of alpha particles from heavy nuclei, 26:47008
- HALVERSON, W., Bremsstrahlung photon emission rate from Maxwellian plasmas, 26:47152
- HALZEN, F., Duality and backward peaks, 26:46627
- Nature of absorption, 26:46623
- HAMANOU, Kumao, Radiolyses of mixtures of ammonia and hydrocarbons. II, 26:45332
- HAMBRO, Lars, Second-order corrections to the fine structure of helium. II. Contributions from ^1P and ^3D intermediate states, 26:46360
- HAMILTON, J. H., Band mixing in ^{156}Gd and ^{158}Gd , 26:46996
- HAMMER, C. F., Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
- HAMMONS, Anna S., Increased effective half-life of intratracheal administered $^{51}\text{Cr}_2\text{O}_3$ by respiratory infection, 26:45910 (CONF-720505-2)
- HANCE, A. B., Fabrication of aluminum oxide films, 26:45931 (KAPL-3164)
- HANCOX, R., Fusion reactors, 26:47179
- HANRAHAN, L. R., Fabrication of aluminum oxide films, 26:45931 (KAPL-3164)
- HANSEN, C. Frederick, Laser power stations in orbit, 26:45428
- HANSEN, G. E., Electromagnetic pulse and time-dependent escape of neutrons and gamma rays from a nuclear explosion, 26:45497
- HANSEN, L. E., Subcritical multiplication of ^{252}Cf neutrons and its applications, 26:46855
- HANSEN, Ole, Subbarrier fission resonances in Th isotopes, 26:47033
- HANSEN, Robert L., Radioisotopes and turbine flow measurements, 26:45721
- HANSON, C. L., Contamination control hardware for remote hot cell transfers, 26:46108 (UCRL-73757)
- Shielded caves for dual-mode operation, 26:46107 (UCRL-73756)
- HANSON, W. B., Satellite and ground-based observations of a red arc, 26:46275
- Synthesis of Jicamarca data during the great storm of March 8, 1970, 26:46282
- HAOT, J., Regeneration of lymphoid bone marrow in the thymectomized adult mouse subjected to a sublethal dose of x rays, 26:45830
- Variations in the bone marrow lymphoid population in the sublethally irradiated germ free mouse, 26:45843
- HARA, Yasuo, Inclusive reactions in the Regge pole model and exotic resonances, 26:46605
- HARBOTTLE, J. E., Application of high-voltage electron microscopy to void studies, 26:46056 (CONF-710601-, pp 533-49)
- HARBOUR, R. M., Conversion of Cf α -hydroxyisobutyrate to oxide with carboxylic acid resin, 26:45393
- Mass yields from thermal-neutron induced fission of ^{235}Cm , 26:47027
- HARDER, C. R., Feedback stabilization on an $l = 1$ theta-pinch plasma column, 26:47157 (LA-DC-72-707)
- HARDIE, F. W., Synthesis, identification, and purification of C₆ and C₇ fluorocarbons from radiolysis. Final report, 26:45316 (ORO-3909-3)
- HARDING, Clifford V., Cellular proliferation and regeneration following tissue damage, 26:45813 (COO-1427-41)
- HARDY, C. J., Management of wastes from irradiated nuclear fuels, 26:46098 (AAEC/TM-628)
- HARDY, E. P., ^{90}Sr fallout in Utah, 26:45525 (HASL-257, pp I.71-94)
- Plutonium fallout in Utah, 26:45526 (HASL-257, pp I.95-118)
- HARDY, Edward P. Jr., Fallout Program. Quarterly summary report, March 1, 1972--June 1, 1972, 26:45511 (HASL-

- 257)
- HARDY, J. C., Beta-delayed proton decay of ^{130}I , 26:46879
- HARE, M. G., Mass radius of the nucleon, 26:46626
- HARKNESS, S. D., Theoretical study of the swelling of fast-neutron-irradiated materials, 26:46063 (CONF-710601-1, pp 798-824)
- HARLEMAN, Donald R. F., Mathematical model for the prediction of unsteady salinity intrusion in estuaries, 26:47620 (MIT-144)
- HARPER, John R., Effects of x irradiation upon response of mononuclear phagocytes to an induced sterile intraperitoneal inflammation, 26:45833
- HARRIS, H. W., Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)
- HARRISON, A. G., Third-derivative method for determining electron-impact onset potentials, 26:46385
- HARRISON, F. L., Accumulation and loss of cobalt and cesium by the marine clam, *Mya arenaria*, under laboratory and field conditions, 26:45545 (UCRL-73744)
- HARSANYI, Gy., Determination of isodose curves of sealed γ -sources by calculation and film dosimetry, 26:45715
- HART, Howard R. Jr., Superconductive magnets, 26:46843
- HARTEL-ULKOWSKA, Marcyza, Evaluation of the causes of failures in roentgenotherapy of carcinoma of the larynx, 26:45780 (TT-54002/4, pp 430-6)
- HARTEN, Ronald Harold, Study of neutral hydrogen gas motion within spiral arms and in the local region, 26:46246
- HARVEY, H. D. Jr., Radiological surveys of Pearl Harbor, Hawaii, and environs, 1966-1968, 26:45534
- HARVEY, J. A., Experimental neutron resonance spectroscopy, 26:46966
- Total neutron cross section of ^{233}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- HASE, Hiroto, Trapped electrons produced in ethanol glass at 4°K, 26:45319
- HASKELL, N. A., Elastodynamic near-field of a finite, propagating tensile fault, 26:45491
- HATTORI, Chuichiro, Relativistically covariant formulation in nonlinear Lagrangian theories and factor-ordering problems, 26:46744
- HAUS, A. G., Film technique for determining the absorbed dose at exit from the patient during radiation exposure, 26:45631
- HAVENS, W. W. Jr., Mass distributions in fission and some related phenomena, 26:47017 (INDC-7-U, pp 31-59)
- HAWKING, S. W., Black holes in Brans-Dicke theory of gravitation, 26:46192
- Black holes in general relativity, 26:46193
- HAYASHI, Kanji, vlf emissions during post breakup phase of polar substorm, 26:46285
- HAYES, S. H., MeV total neutron cross sections of ^{238}U , 26:47013 (COO-3058-23)
- HAYES, Stuart F., Effects of screening and polarization on vibration-rotation properties of H_2 , 26:46350
- HAYMAKER, Webb, Neuropathological evaluation of monkeys exposed to body-alone x-radiation, 26:45850
- HAYON, E., Electron and hydrogen atom attachment to aromatic carbonyl compounds in aqueous solution. Absorption spectra and dissociation constants of ketyl radicals, 26:45354
- HAYS, Esther F., Irradiation and viral lymphomagenesis in thymic grafts, 26:45818 (UCLA-12-867)
- Single cell transplants: Gross virus-induced mouse lymphomas, 26:47629 (UCLA-12-866)
- HEADY, H. H., Fire assay-emission spectrographic determination of iridium, ruthenium, and osmium, 26:45229
- HEFNER, Bruce A., Liquid processing of molded desiccant, 26:47602 (BDX-613-392(Rev.))
- HEGERFELDT, G. C., Lorentz transformations: derivation of linearity and scale factor, 26:46732
- HEIBERGER, J. J., Containment of fission product iodine in the reprocessing of LMFBR fuels by pyrochemical reactions, 26:45421
- HEIDENTHAL, Gertrude, Fecundity and longevity of F_1 females of *Habrobracon* from sperm x-rayed with 3000 r, 26:45878
- HEIJNEKER, H. L., In vitro repair of uv-inactivated transforming DNA from *Bacillus subtilis*, 26:45727
- HEIKKILA, W. J., Fermi acceleration of auroral particles, 26:46267
- HEINRICH, Wolfgang, Radiation hazards to man in space flights, 26:45812
- HEINZELMANN, M., Scattered neutrons and their effects on calibration of Bonner sphere counters, 26:45599
- HELINIAK, Andrzej, Selected prognostic elements in carcinoma of the larynx, 26:45781 (TT-54002/4, pp 437-41)
- HELLER, Eric J., Interpolation of variationally obtained scattering cross sections, 26:46402
- HELM, S., Power production costs and possible applications of short-period combustion MHD generators, 26:47170 (IPP-IV-32)
- HELMKEN, H. F., Detection of high-energy gamma rays from the Crab Nebula, 26:46214
- HEMENWAY, Robert Clark Jr., Effects of gamma radiation on the respiration of *Acarus siro*, 26:45884
- HENDRICKS, J. S., Nondestructive assay of power reactor fuel assemblies, 26:45269
- HENDRICKS, Melvin Elton, Magnetic properties of some actinide halide compounds, 26:47272
- HENDRICKSON, A. A., Structure and properties of solid solutions. Technical progress report, September 1, 1971-June 15, 1972, 26:45991 (COO-916-25)
- HENLEY, E. M., Charge-symmetry breaking and N-N scattering lengths, 26:47055
- HENRIKSEN, Thorodd, Electron spin resonance studies and INDO MO-calculations of electron irradiated peptides. A single crystal of N-acetylglycine, 26:45349
- HENRY, Marian S., Iron(II) and iron(III) isotope exchange in water-dimethyl sulfoxide mixtures, 26:45304
- HERBST, J. F., Single-electron energies, many-electron effects, and the renormalized-atom scheme as applied to rare-earth metals, 26:47260
- HERKSTROETER, William G., Absorption spectroscopy of transient species, 26:45340
- HERMANN, Bruce Raymond, Polarization aperture synthesis of five galactic nonthermal radio sources, 26:46207
- HERMENT, Micheline, Chemical separation and high resolution nuclear spectroscopy in ^{228}Ra , ^{228}Ac , ^{228}Th radioactive chain, 26:47014 (FRNC-TH-184)
- HERNANDEZ, G., Determination of the quenching of $\text{O}(^1\text{D})$ by molecular nitrogen using the ionospheric modification experiment, 26:46278
- HERSHKOWITZ, N., Analysis of five line hyperfine split Moessbauer absorber spectra, 26:47275
- HERSKIND, B., g-factor of the 12^+ isomeric state in ^{206}Pb , 26:47002
- HESPE, E. D., Management of wastes from irradiated nuclear fuels, 26:46098 (AAEC/TM-628)
- HESS, Kenneth H., Remote effects, in time and place, of x irradiation of the visual cortex in the monkey, 26:45858
- HESSER, J. E., Optical observations of supernova in NGC 5253, 26:46187
- HETTINGER, G., Commercial thimble chambers for absorbed dose measurements at high energy electron radiation, 26:45623
- HEUKELS, W. F., Quantum-mechanical scattering of asymmetric molecules. II. Comparison between the systems $\text{He}-\text{D}_2$ and $\text{He}-\text{HT}$, 26:46379
- HEUMANN, K. G., Dependence of the formation of uranium and thorium ions and their oxide ions on various conditions of thermal ionization, 26:45258
- HIGASHIMURA, Takenobu, Trapped electrons produced in ethanol glass at 4°K, 26:45319
- HIIDA, K., Many-body forces and the effect of the matter distribution in the Universe to the gravitational constant, 26:46765
- HIIDA, Kichiro, Gauge transformation and gravitational potentials, 26:46766
- HILL, I. E., Observations of supernova remnant IC 443 at 1.42 GHz, 26:46203
- HILL, J. R., Method for rapid computation of vlf, steep-incidence reflections from the ionospheric D region, 26:46263 (AD-734392)
- HILL, Lemuel L., Spectra of neutrons emitted after muon capture in ^{16}O , ^{28}Si , ^{32}S and ^{40}Ca , 26:46880
- HILLNHAGEN, Emil, Special techniques for the metallographic examination of hard metals, 26:45955
- HILLS, P. R., Radiation polymerization of impregnated porous material, 26:45718
- HINDMARSH, A. C., Determinant method for calculating small-crystal entropy, 26:47632 (UCRL-51227)
- HIRAI, Hidefumi, Active species of γ -irradiated vinyl acetate and their changes on irradiation of visible light, 26:45334
- HIRAKI, Keizo, Fluorometric determination of Al and Ga with Superchome Garnet Y, 26:45236
- HIRAMATSU, Hiroshi, Quantitative evaluation of radioisotope distribution in vivo by isosensitive scanner plus 4096-word multi-channel analyzer coupling, 26:45601
- HIRIKAWA, N., Asymmetry for π^+ photoproduction from polarized protons between 300 and 900 MeV at 90°, 26:46444
- HIRSCH, P. B., Dynamical theory for the contrast of perfect and imperfect crystals in the scanning electron microscope using backscattered electrons, 26:47267
- HIRSCH, Robert L., Electrostatic containment in fusion reactors, 26:47113
- HIRSHBERG, J., Velocity and flux dependence of the solar-wind helium abundance, 26:46328
- HIRST, Eric, Livestock shelters as horizontal cooling towers, 26:45576
- HISADA, Kin-ichi, Quantitative evaluation of radioisotope distribution in vivo by isosensitive scanner plus 4096-word multi-channel analyzer coupling, 26:45601
- HIXSON, J. D., Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)
- HJORTH, S. A., Validity of rotor model in odd-Z nuclide ^{171}Lu , 26:46995
- HLASNIK, I., Study of losses in an assembly of superconducting filaments transposed in a conducting matrix and subjected to a pulsed magnetic field, 26:46810
- HLINIAK, Andrzej, Prognostic elements in inoperable carcinoma of the breast, 26:45783 (TT-54002/4, pp 451-5)
- HOANG, C. T., Meridional transport of ^{90}Sr in the troposphere, 26:45516 (HNSL-257, pp I.2-17)
- HOBBS, L. M., Circumstellar molecules in the Pleiades, 26:46190
- HOCH, R. J., Satellite and ground-based observations of a

HODGE, Paul W., Some current studies of galaxies, 26:46234
HODGE, V. F., Plutonium and polonium inside giant brown algae, 26:45924

HODGSON, R. J. W., Solution of the Bethe--Goldstone equation in finite nuclei from N--N phase shift data, 26:47074

HODINKA, L., Tissue mast cells in human blood, 26:45810
HOEDE, D., Determination of chromium in milk powder, 26:45196

HOEHLER, G., Review of near-forward π N scattering amplitudes at high energies, 26:46636

HOFER, K. F., Biological toxicity of intracellular radioisotope decay by beta emission and electron capture, 26:45892 (ORO-2690-12(Pt.B), Paper 4)

HOFER, W., X-ray-induced photoconductivity in dielectric films, 26:45668 (UCRL-73974)

HOFFMAN, C. M., Experimental study of the decays $K/\text{sub } S \text{ L/} + \gamma$, 26:46434

HOFFMAN, E. A., Effect of ^{137}Cs radiation on growth, development, and mortality of larvae of the tussock moth, *Hemerocampa leucostigma*, 26:45885

HOFFMAN, E. W., Observation of the $M^0(940)$ and $\Delta^0(963)$, 26:46475 (IS-2886)

HOFFMANN, A., Study of vibrations of the centering parts of fuel element cartridges, 26:47314

HOFFMANN, G. W., Coupled-channel calculations of the energy dependence of the (p,n) charge-exchange reaction, 26:47073

HOFFMANN, W., Special machines and equipment of WAK, 26:46106 (ORNL-tr-2372)

HOLLANDER, J. J., Decay of ^{210}At to levels in ^{210}Po , 26:47006

HOLLAS, C. L., $^{90}\text{Zr}(d,n)^{91}\text{Nb}$ and $^{96}\text{Zr}(d,n)^{97}\text{Nb}$ reactions, 26:46956

HOLLOWAY, A. F., Workloads and use factors in diagnostic x-ray installations, 26:45811

HOLLOWAY, L., Missing-mass spectrum near 960 MeV from $\pi^-p \rightarrow n + (\text{mm})$, 26:46430 (COO-1195-241)

HOLLSTEIN, M., De-excitation of fast He, Ne, and Ar metastable atoms in various gases, 26:46394

HOLLWEG, Joseph V., Heat conduction in a turbulent magnetic field with application to solar-wind electrons, 26:46326

HOLMES, J. J., Recovery and recrystallization of prototypic FTR fuel cladding: 20 percent CW 316 stainless steel tubing (RM-11), 26:46067 (HEDL-TME-71-126)

HOMAYOONFAR, Mehdi, Application of track structure theory to dosimetry and radiobiology, 26:45611

HONIG, Richard E., Mass spectrometric techniques, 26:45248
HOOVER, W. G., Determinant method for calculating small-crystal entropy, 26:47632 (UCRL-51227)

HOPEGARTEN, N., Cathode heating in an asymmetrically burning Penning discharge, 26:47136

HOPKINS, H. W. K., Measurement of slope parameter in decay $K/\text{sub } L^0 \rightarrow \pi^+\pi^-$, 26:46494

HORNSEY, S., Sensitization of hypoxic cells in vivo with tracetoneamine N-oxyl (TAN), 26:45869

HORTON, J. L., $^{90}\text{Zr}(d,n)^{91}\text{Nb}$ and $^{96}\text{Zr}(d,n)^{97}\text{Nb}$ reactions, 26:46956

HORVATH, B., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)

HORVATH, Zsuzsanna, Radio-frequency alcoholic chlorobenzene dosimeter, 26:45632

HOSAKA, Koji, Effects of cepharanthine on fluctuation in white blood cell count by radiotherapy for gynecologic cancer, 26:45805

HOSHI, Michio, Precipitation of some actinide element complex ions by using hexamine Co(III) cation. IV, 26:45370

HOSTE, J., Instrumental neutron activation analysis of germanium, 26:45218

HOUSEHOLDER, A. S., KWIC Index for numerical algebra, 26:47287 (ORNL-4778)

HOWARD, Boyd D., Process of forming an isotopic heat source, 26:45707

HOWARD, J. S., Chemi-ionization in collisions between helium metastable atoms and argon, 26:46404

HOWE, Herbert C. Jr., Explorer 33 and 35 plasma observations of magnetosheath flow, 26:46303

HOYER, P., Regge phenomenology of inclusive spectra in the central region, 26:46659

HOYLE, F., Mass difference of the muon and the electron, 26:46259

HRASKO, P., Spontaneous decay in a time dependent external field, 26:46571 (KFKI-72-15)

HRUSKA, A., Anisotropy of high-latitude electron fluxes during substorms and structure of the magnetotail, 26:46306

HSU, Chia-jung, Effect of lateral rod displacement on laminar-flow transfer, 26:45462

HU, C.-R., Dynamic structure of vortices in superconductors. III. Graphs for B approximately = $H/\text{sub } c2/$, 26:46814

HU, Chia-ren, Numerical constants for isolated vortices in superconductors, 26:46812

HUB, R., Consideration on direct reaction mechanisms in the unified microscopic nuclear theory, 26:47077

HUBER, J. F. K., Rapid separation of metal chelates by column liquid-liquid chromatography using ultraviolet detection, 26:45281

HUBER, Kent Alvin, Investigation of two nucleon transfer

reactions in ^{19}F using 45 MeV protons, 26:46895

HUCKABEE, John W., Neutron activation analysis of trace elements in hair, 26:45214

HUDSON, J. A., Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)

HUEBOTTER, Paul R., Effects of metal swelling and creep on fast reactor design and performance, 26:47303

HUEHNERMANN, H., Optical determination of the quadrupole moment of ^{138}La , 26:46950

HUG, O., Radiation hazards and radiation protection, 26:45778 (NP-tr-1941)

HUGHES, D., Radiation accidents in university laboratories, 26:45791

HUIZENGA, J. R., Level densities from spectra of different reactions populating the same nucleus, 26:46932

HUIZENGA, W., Production and measurement of intense ion beams and beam-produced plasma in a cusped magnetic field, 26:47153 (NP-19364)

HULD, B., Missing-mass spectrum near 960 MeV from $\pi^-p \rightarrow n + (\text{mm})$, 26:46430 (COO-1195-241)

HUMBLE, S., Triple Regge couplings and the central region of inclusive distributions, 26:46661

HUMPHREYS, C. J., Dynamical theory for the contrast of perfect and imperfect crystals in the scanning electron microscope using backscattered electrons, 26:47267

HUNSPERGER, R. G., Optical waveguiding in proton-implanted GaAs, 26:47232

HUNSTEAD, R. W., Accurate positions of radio sources at 408 MHz, 26:46202

HUNTER, James H., Influence of local conditions in the interstellar medium upon star formation, 26:46169

HUNTLEY, D. J., Discontinuous flux flow in superconducting NbSe_2 , 26:46831

HUPPERT, K. L., Special machines and equipment of WAK, 26:46106 (ORNL-tr-2372)

HUQ, S. S., Systematic methods for the analysis of a sample of nuclear interactions in hydrogen bubble chamber, 26:45651

HURSH, John B., Retention of ^{210}Pb in beagle dogs, 26:45916 (UR-3490-159)

HUSEBYE, Eystein S., Seismic arrays and seismology, 26:45498

HUSLER, John, Determination of niobium in rocks, ores, and alloys by atomic-absorption spectrophotometry, 26:45234

HUSSEIN, A. M., Nonlinear focusing of TM waves in isotropic plasma, 26:47185

HUTT, P. K., Duality properties of Regge cuts, 26:46600

HUYGHE, J., Design considerations of heat exchangers for vaporizing fluids, 26:45436 (CEA-CONF-2010)

HWA, Rudolph C., Tests of high-energy models by two-particle analyses, 26:46587 (RL0-2230-T-4-1)

HWANG, A. E., On a possible repulsive interaction in universal gravitation, 26:46749 (ORO-3592-94)

HWANG, Jae Y., Atomic absorption and flame emission spectroscopy in the analysis of trace metals in atmospheric particulates, 26:46244

HYDE, James S., Powder ENDOR line shapes: nuclear relaxation induced by motion of nearby electron spins, 26:47279

Proton ENDOR of γ -irradiated Y-type zeolites, 26:45324

IANNAIRELLA, L., Gadolinium g-shifts and band structure, 26:47270

Remarks on Gd g-shifts, 26:47271

IANNIELLO, Louis C., Radiation-induced voids in metals. Proceedings of the 1971 international conference held at Albany, New York, June 9--11, 1971, 26:46045 (CONF-710601)

IBADINOV, Kh., Size of the dust particles of a comet atmosphere and the heliocentric distance, 26:46220

IBATA, T., Electron and hydrogen atom attachment to aromatic carbonyl compounds in aqueous solution. Absorption spectra and dissociation constants of ketyl radicals, 26:45354

ICHIKAWA, Masaru, Identification of reaction intermediates by microwave spectroscopy. The catalytic reactions between propylene and deuterium over zinc oxide, 26:45303

IGALI, S., Determination of microbial contamination of disposable medical devices and the estimation of radiosensitivity of isolated microflora, 26:45748
Microbe-genetical problems of radiosterilization, 26:45750

IGARASI, Sin-iti, (N--Z) dependence of (n, γ) cross sections, 26:47085

IHLE, H., Optical determination of the quadrupole moment of ^{138}La , 26:46950

IKEDA, Takayoshi, Statistical analysis of radioinduced abnormal development, 26:45867

IKUTA, A., Optimization study of the precipitation of thorium oxalate, 26:45365 (IEA-183)

ILLE, Bernard, Final state interaction between proton and neutron from $^1\text{H}(d,pn)^1\text{H}$, $^2\text{H}(d,pn)^2\text{H}$, $^3\text{He}(d,pn)^3\text{He}$, and $^4\text{He}(d,pn)^4\text{He}$ reactions, 26:46860 (LYCEN-7186)

IMBERT, Pierre, Automatic analysis by absorption spectrometry, 26:45243

- IMSHENNIK, V. C., Two-dimensional nonsteady-state solution of magnetohydrodynamic equations, 26:46425
- INADA, Koji, Electron spin resonance study of the far-ultraviolet photolysis of some inorganic and organic compounds, 26:45323
- INAMI, T., Analysis of kaon--nucleon scattering amplitudes in the intermediate energy region, 26:46624
- INCZE, K., Effect of combined methods on Clostridium sporogenes inoculated in sausage, 26:45747
- INDOVINA, Pietro L., Phase sensitive method for detecting superconductive transitions in the microwave region, 26:46840
- INGERSOLL, Andrew P., Photospheric faculae and the solar oblateness, 26:46224
- INGHAM, David Ransford, Small angle inelastic scattering of 40 MeV protons by ^{12}C , ^{40}Ca , ^{56}Fe , and ^{60}Ni , 26:46894
- INGLE, J. D. Jr., Some comments on the signal-to-noise characteristics of real photomultiplier and photodiode detection systems, 26:45682
- INGLE, R. W., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)
- INHABER, H., Magnetic field distributions in Type II superconductors, 26:46833
- INOAGMOV, Sh. V., Spectrum symmetry of pion longitudinal momenta in $\pi\text{-p}$ interactions at $p_{\text{c}} = 4 \text{ GeV}$, 26:46619 (JINR-P1-6461)
- INOUE, Chuya, Localized magnetic states in superconductor, 26:46809
- INOUE, Kazuhiko, Slow neutron spectra in the liquid and solid methane, 26:46858
- IOANNU, P. D., Elastic and inelastic scattering of 6-MeV protons by even isotopes of Cr, 26:46922
- IONKINA, E. M., Structure, phase composition, and properties of cast complex alloys of Ni--Nb, 26:46035
- JORDANISHVILI, E. A., Gamma quanta distribution at large distances from the sources, 26:47215
- IRIE, Koichi, Automatic frequency control against temperature change in linac's accelerating structure, 26:46127
- Q-value effects of accelerating structure in traveling-wave linacs, 26:46122
- ISAEV, P. S., Reaction $\gamma + \gamma \rightarrow \pi + \pi$ and electromagnetic difference for pion masses, 26:46519 (JINR-P2-6333)
- ISAICHEV, V. I., Textures of rolling and annealing in tungsten foils of various compositions, 26:46042
- ISAILA, M. V., Acceleration of argon ions to 1.17×10^{10} electron volts, 26:46124
- ISHIDA, Koji, Thin-layer chromatography of metal ions on microcrystalline cellulose impregnated with potassium chloride, 26:45278
- ISHII, Chikara, Thermodynamical properties of Josephson junction with normal-metal barrier, 26:46808
- ISHIKAWA, Koki, Phonon dispersion and contribution to the heat capacity in liquid ^4He , 26:46795
- ISHIMARU, Shingo, Long-range interactions between two hydrogen atoms and two helium atoms, 26:46351
- ISHIMORI, Tomitaro, Ranges of fission fragments in uranium--aluminum alloys and uranium oxides, 26:46074
- ISHIWARA, Hiroshi, Mass dependence of critical amorphizing dose in ion implantation, 26:47236
- ISHIZAKI, Y., $^2\text{H}(n,p)^2\text{n}$ reaction at 50 MeV and neutron--neutron scattering length, 26:46861
- ISHKhanov, B. S., New experimental data on the fine structure of dipole resonance, 26:46890
- ISKANDEROVA, A. D., Age of cuprous Ukoka series (Olekma--Vitim mountain system) by uranium--lead method, 26:45569
- ISKHAKOV, G. I., Study of the decay $^{140}\text{Tb}(T_{1/2} = 66 \text{ min})$ and $^{150}\text{Tb}(T_{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- ISLER, R. C., Optical excitation and polarization in low-energy collisions of $\text{He}^+ + \text{H}_2$, 26:46396
- ISMAIL, M., Analog states in ^{33}Cl by isospin-forbidden resonances in reaction $^{32}\text{S}(p,\gamma)^{33}\text{Cl}$, 26:46902 (BARC-598)
- ISOMOTO, Yukuo, (N--Z) dependence of (n, γ) cross sections, 26:47085
- ISOZUMI, Y., Possible evidence for a contribution from direct collisions in K-shell internal ionization accompanying β decay, 26:46962
- ISRAELSSON, Sven, Meteorological influences on atmospheric radioactivity and electricity at ground level, 26:45524
- ISSEL, W., WAK: specifications and finishing control for equipment, 26:46105 (ORNL-tr-2371)
- ITO, Haruhide, Effects of cepharanthine on fluctuation in white blood cell count by radiotherapy for gynecologic cancer, 26:45805
- ITO, Yukio, Mixing parameter epsilon in neutral K decay, 26:46585
- IUCCI, N., Diurnal effect of cosmic rays during the period 15 October 1965--30 June 1966. II, 26:46158
- Diurnal effect of cosmic rays during the period 15 October 1965--30 June 1966. I, 26:46159
- IVANCHENKO, V. G., Alloys of chromium with terbium, 26:46033
- Solubility of Ce, Pr, and Nd in Cr containing O, 26:46034
- IVANENKO, D. D., Toward the theory of nuclear shells, 26:47040 (BNL-tr-489)
- IVANENKO, M. A., Allowance for mass-spectrometer background and the reproducibility in the determination of isotope compositions for nitrogen and oxygen, 26:45241
- IVANOV, E. A., Radiometric unit for measuring wear of materials and machine parts, 26:45657
- IVANOV, I. N., Influence of the electromagnetic ring acceleration on the focusing by image forces, 26:46136 (JINR-P9-6421)
- Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)
- IVANOV, L. I., Creep of niobium under conditions of in-leakage of oxygen, 26:46000
- IVANOV, R. B., α -decay of neutron-deficient isotopes of Ra, 26:47004
- IVANOV, S. N., Effect of soil conditions and K on uptake of ^{86}Rb in corn plants, 26:45925
- IVANOV, V. B., γ -emission in radiative capture of intermediate neutrons, 26:46917 (BNL-tr-477)
- IVANTSOV, I. I., Electron transfer during zone melting of yttrium, 26:45975
- IVANYUKOVICH, G. A., Determination of density and effective atomic number of rocks and ores by gamma--gamma method in the presence of an intermediate zone, 26:45566
- IVASCU, M., (α,p) reaction on even nickel isotopes, 26:46934
- IWAI, Yoshiaki, Culture of murine leukemic cells induced by x-irradiation, 26:45855
- IWASAKI, M., Fluorination of UO_2 and U_3O_8 by gaseous Br trifluoride, 26:45379
- Infrared absorption spectra of potassium uranium fluorides, 26:45230
- IWASAKI, Machio, ESR study of the vinyl-type radical trapped in a single crystal of maleic acid irradiated at 77°K , 26:45347
- IWASAKI, Zensuke, Radiation effects on bone, 26:45870
- IZZO, Mary Jane, Distribution of an ^{125}I -labeled transplantation antibody in syngeneic and allogeneic rats, 26:47649 (UR-3490-147)
- JACKSON, K. P., ^{53}mCo : a proton-unstable isomer, 26:46925
- JACOBEL, R. W., Observation of the $\text{M}^0(940)$ and $\text{delta}^0(963)$, 26:46475 (IS-2886)
- JACOBI, Siegfried, New electrical procedure for the compaction of technical powders, 26:45951
- JACOBI, W., Diffusion deposition of small-sized particles in the bronchial tree, 26:45921
- JACOBSEN, T., Enhancement at about 2700 MeV/c ^{2} in the pp--X^0 system in $\text{pp} + \text{pp}^*\text{--X}^0$ at 19 GeV/c, 26:46467
- JACROT, Bernard, Neutron scattering, 26:47251
- JAFFE, A. A., Energy levels of ^{36}K , 26:46904
- JAHNKE, U., X-rays and nuclear γ -rays in muonic thallium, 26:47031
- JAHOBA, F. C., Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)
- Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
- JAIN, A. K., Charge form factor and quadrupole moment of ^6Li , 26:46878
- JAIN, P. L., Inclusive reaction $\mu + \text{nucleon} \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV/c, 26:46452
- JAKES, Dusan, Exploitation of solid electrolytes in research and technology of nuclear power, 26:46013
- JAMAR, C., Early data from ultraviolet sky-scan telescope in TD1 satellite, 26:46186
- JANCEL, Raymond, Kinetic analysis of collision effects on nonlinearities related to electromagnetic wave propagation in a weak dissipative-ionized medium, 26:47183
- JARDINE, L. J., Decay of ^{210}At to levels in ^{210}Po , 26:47006
- JAVELLAUD, Jean, Composite walls for magnetohydrodynamic apparatus, 26:45426
- Semiconductor well logging probe, 26:45658
- JEAN-MARIE, B., Hadronic production by e^+e^- collisions at 990 MeV with Orsay storage ring, 26:46429 (CONF-710816-15)
- Measurement of radiative neutral decay modes of ω^0 and ϕ^{10} mesons with Orsay storage ring, 26:46472 (CONF-710816-12)
- $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
- $\pi^+\pi^-$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
- JECKO, Bernard, Orthogonality relations in periodic waveguides having anisotropic surface impedance, 26:46141
- JELLEY, N. A., ^{53}mCo : a proton-unstable isomer, 26:46925
- JENSEN, D., Inelastic e-p scattering and the polarization contribution to H hyperfine structure, 26:46523 (NYO-476-17)
- JENSEN, K., Relative level populations in foil excited He and Li, 26:46337
- JERVIS, R. E., Activation analysis of environmental pollutants, 26:45198
- JHA, Kamal N., Effect of pressure on the rates of reaction of solvated electrons in liquid ethanol, 26:45351
- JOE, E. G., Zirconium tetrafluoride process, 26:45274

JOELSSON, I., Stenosis of the small bowel as a complication in radiation therapy of carcinoma of the uterine cervix, 26:45802

JOHANSON, Karl J., Deoxyribonucleic acid metabolism in small intestine in vivo and in vitro, 26:45842

JOHN, James E. A., Thermal pollution: a potential threat to our aquatic environment, 26:45583

JOHNSON, J. L., Viscosity and thermal conductivity effects on the containment of a toroidally confined resistive plasma with flow, 26:47105

JOHNSON, Noah R., Band mixing in ^{156}Gd and ^{158}Gd , 26:46996

JOHNSTON, J. R., Syntactic foam, FY70-3, 26:47600 (BDX-613-253(Rev.))

JOHNSTON, K. J., Microwave celestial water-vapor sources, 26:46206

JOINER, W. C. H., Voltages induced in superconductors in the absence of transport currents, 26:46835

JOKELA, T., Natural ^{210}Pb and ^{210}Po in a marine environment, 26:45544 (RLO-2225-T-14-4)

JOKIPII, J. R., Heat conduction in a turbulent magnetic field with application to solar-wind electrons, 26:46326

JOLIVET, Andre, Relativistic scattering of particles with spin, 26:46793

JONA-LASINIO, G., Method of characteristics for functional-derivative equations, 26:46728

JONARD, Robert, Radiorecovery effect of yeast extract on carnation cuttings irradiated with ^{60}Co γ rays, 26:45777

JONES, Frank C., Lorentz-invariant formulation of Cherenkov radiation by tachyons, 26:46704 (N-72-19766)

JONES, R. W., Determination of the electron-phonon enhancement factor from specific-heat data, 26:47262

JOSEPH, Allen F., Execution time translate subroutines, 26:47621 (ORNL-TM-3810)

JOSHI, S. K., Excitation of hydrogen in collisions with alkali-metal atoms, 26:46403

JULLIAN, S., e^+e^- production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)

e^+e^- production by e^+e^- annihilation at $\sqrt{s}$ mass, 26:46427 (CONF-710816-13)

JUNJEN, J. L., Effects of thyroidian hormones and γ irradiation on adenylyclase of plasma membranes of rat liver, 26:45831

K

KABANOV, Vitaly Yakovlevich, Method of modifying polymeric materials, wood, and articles made therefrom by radiation-initiated graft copolymerization thereof, 26:45355

KABYSH, G. M., Radiometric investigation of the absorption of potential-determining ions from aqueous solutions of non-ionogenic surface-active substances, 26:45279

KADA, T., Semi-in vitro repair of radiation-induced damage in transforming DNA of *Bacillus subtilis*, 26:45731

KADOYA, Yoshiki, Development of 50Hz 44 inch turbine blade, 26:45454

KAFATOS, Minas C., Time-dependent ionization equilibrium and line radiation under flarelike conditions, 26:46226

KAIMAL, V. N. P., Spectrochemical analysis of high purity cadmium, 26:45220 (BARC-593)

KAINER, R. A., Ocular squamous cell carcinoma (cancer eye) in hereford cattle: radiation repair processes and a comparison of cultured cells with xeroderma pigmentosum in man, 26:45851

KAISOV, D. K., Lifetime of 1.013-MeV state in ^{27}Al , 26:46905

KAISER, R., Radiobiological problems caused by supersonic transport. (With a survey of the first results established by tests performed on board the Concorde prototype), 26:45796

KAKEHI, Hirotake, Subtraction and positron scanning with the dual headed whole body scanner, 26:45593

KAKIHANA, Hidetake, Anion exchange resin separation of Gd, Eu, Sm, Nd, and Pr, 26:45280

KALEBIN, S. M., Realization of synchronous rotation of some rotors suspended in a magnetic field, 26:45441 (SRARI-P-120)

KALIBJIAN, R., Line-focus electron beam lens, 26:46147

KALINAUSKAS, R. K., Two-particle orbital matrix of density for one-quanta oscillator-excited SU_3 -configurations for p-nuclei, 26:47067

KALININ, A. I., Time measurements with a germanium detector, 26:45616

X-ray spectrometer having cooled $\text{Si}(\text{Li})$ -detector, 26:45644

KALMAN, G., Linear Stark effect due to resonant interactions of static and dynamic fields, 26:46361

KALMYKOV, L. Z., Influence of a container's material for indications rendered by chemical methyl-orange dosimeter, 26:45633

KAMALI, A. R., Effect of gamma irradiation on Fuerte avocado fruits, 26:45724

KAMINSKAYA, A. M., Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)

KAMMERER, O. F., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)

KANADA, Hiroyuki, Pion form factor and possible heavy

vector meson, 26:46561

KANAZAWA, M., Electrical properties of n-type Si layers doped with proton bombardment induced shallow donors, 26:47237

KANAZAWA, Tamotsu, Dose-rate measurement by several Radocon probes. II, 26:45620

KANE, G. L., Simple description of high energy elastic scattering, 26:46660

KANEKO, T., $K/\text{sub } 13/-$ decay parameters and possible scalar coupling in weak interactions of urbaryons, 26:46583

KANTZ, Asher D., Ecological and environmental effects from local fallout from Schooner. 2. The beta and gamma radiation effects from close-in fallout, 26:45542 (PNE-529)

KAPATURIN, G. G., Distribution and genesis of radioactivity in coals of the Pechora River basin, 26:45505

KAPOOR, S. S., Excitation energy dependence of shell effects in nuclear fission, 26:47043 (INDC-7-U, pp 3-19)

KARACHENTSEV, I., Predominant motion in spherical systems of galaxies, 26:46235

KARASUND, T., s-channel helicity amplitudes without pomeron for elastic $K^+ \rightarrow p$ and $w \rightarrow p$ charge exchange scattering, 26:46637

KARLSSON, B. R., Breakdown of simple cancellation in the operator multiple scattering series, 26:46773 (CTH-ITP-71-17)

KARMANOV, V. A., Many-neutron resonances, 26:46644

KAROL, I. L., Assessment and control of environmental contamination, 26:45554

KARR, H. J., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)

KASHEVSKAYA, O. N., Solubility of Ce, Pr, and Nd in Cr containing O, 26:46034

KASHIHIRA, Nobuyuki, Ionizing collisions of fast alkali atoms with Cl_2 , Br_2 , and O_2 , 26:46366

KASUGA, Toshio, Hepatic injuries following irradiation, 26:45809

KASULOV, V. I., Peculiarities in interaction of high-energy electrons and positrons with crystals, 26:46550

KASYMOV, A. Kh., Distribution of alkali ions in nickel by ionic bombardment, 26:47235

KATAYAMA, Yoshishige, Spontaneous emission of plasma waves in the presence of a finite-amplitude wave, 26:47180

KATHREN, R. L., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)

KATKEVICH, Yu. Yu., Gamma induced changes in spruce wood, 26:45774

KATO, Motohiro, Remote effects, in time and place, of x irradiation of the visual cortex in the monkey, 26:45858

KATO, S., Strong-interaction form factor of proton in high-energy proton-proton elastic scattering, 26:46654

KATRIEL, J., Algebraic derivation of generalized Talmi expansion, 26:46503

KATZ, Joseph L., Nucleation of voids in materials during high-energy irradiation, 26:46064 (CONF-710601-, pp 825-40)

KAUL, Ashok Kumar, Processes and measurements in an abnormal negative glow plasma, 26:47138

KAUL, B. L., Cystein protection against ethylenimine damage in barley seeds, 26:45772

KAVALENKO, P. N., Extractive photometric determination of U(VI) in mine waters, 26:45242

KAVILADZE, M. Sh., Separation of sulfur isotopes by a chemical exchange method, 26:45688 (MLM-1936(TR))

KAWABATA, Kiyosh, Continuum near the H and K lines of Ca^+ in the solar spectrum, 26:46218

KAWASHIMA, Nobuki, Neutral-point discharge experiment, 26:47142

KAWASHIMA, Teruko, Flux flow in nonideal Type-II superconductors, 26:46819

KAZAMANOVA, Iraida Vasilievna, Method of modifying polymeric materials, wood, and articles made therefrom by radiation-initiated graft copolymerization thereof, 26:45355

KAZARINOV, Yu. M., Scale relation for proton form factors, 26:46433

KAZILIN, E. E., Creep of niobium under conditions of in-leakage of oxygen, 26:46000

KEARNEY, P. D., Evidence for solar particle production above approximately 75 GeV, 26:46219

KEEPER, D. W., Studies of void formation in proton-irradiated type 316 and titanium-modified 316 stainless steels, 26:46044 (AI-AEC-13033)

Swelling as a function of displacement damage in proton-irradiated 316 stainless steel, 26:46054 (CONF-710601-, pp 511-21)

KEIL, E., CERN proton storage ring, 26:46143

KEILIN, V. E., Cryogenic problems of forced cooled superconducting systems, 26:46839

KEIRIM-MARKUS, I. B., Comparison of neutron and gamma-ray personnel dosimeters at critical assemblies, 26:45630

KELEMEN, V. I., Photoproduction of vector and axial vector resonances in broken nonlinear chiral $\text{SU}_3 \times \text{SU}_3$ dynamics, 26:46563

KELIHER, T. E., Charge-symmetry breaking and N-N scattering lengths, 26:47055

- KELLER, K. A., Mean excitation energies of the daughter nuclei on β^- -decay, 26:46875 (KFK-EXT-10/72-1)
- KENG, Edward Y. H., Aerosols produced by x-rays, 26:46082
- KENNEL, C. F., Changes in magnetospheric configuration during the substorm growth phase, 26:46305
- KENNEL, Charles F., Pitch-angle diffusion of radiation-belt electrons within the plasmasphere, 26:46311
- KERBIKOV, B. O., Quantum numbers and total and partial widths of the resonance N anti N(1970), 26:46458
- KERIMOV, B. K., Spin correlation during bremsstrahlung radiation of longitudinally polarized electrons, 26:47052
- KERN, Robert Allyn, Plasma current conduction to a cold electrode, 26:47130
- KERNEL, G., Test of time reversal invariance through detailed balance of the reaction $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$, 26:46881
- KERNER, M. N., Separation of sulfur isotopes by a chemical exchange method, 26:45688 (MLM-1936(TR))
- KERRIDGE, John F., Accretionary processes in the early solar system: an experimental approach, 26:46229
- KESLEV, D., Effect of Al phosphate gel on whole-body retention of simultaneously administered ^{22}Ra , ^{85}Sr , and ^{45}Ca in mice, 26:45918
- KESSLER, P., Acoplanarity and radiative corrections in Bhabha scattering, 26:46543
- KETTERSON, J. B., Absence of a quadratic term in the ^4He excitation spectrum, 26:46799
- KEVAN, Larry, Role of fluoride ion in irradiated aqueous systems, 26:45345
- KHABAKHPASHEV, A. G., ϕ -meson mass determination, 26:46443
- KHACHIROV, Dzh. G., Dose to the epithelium of the upper respiratory tract from radon and its daughter products, 26:45861
- KHAIDAROV, K., Properties of ruby laser radiation under action of ^{60}Co gamma radiation, 26:47227
- KHAKIMOV, U. B., Al-Fe-Mo oxide catalysts. I, 26:45963
- KHALILOV, V. R., Induced emission from electron in magnetic field at level $n = 1$, 26:46551
- KHAM'YANOV, L. P., γ -emission in radiative capture of intermediate neutrons, 26:46917 (BNL-tr-477)
- KHAN, H. A., Prolonged etching factor in solid state track detection and its applications, 26:45604
- KHAN, Jhan M., Surface science and surface damage, 26:47636 (UCRL-73925)
- KHARE, A. V., Low-energy theorem for pion-nucleon scattering from pion source dynamics, 26:46641
- KHARTONOV, Yu. I., Regularity in multiplet properties of odd-odd nuclei, 26:47051
- KHIL'CHEVSKAYA, T. V., Structure, phase composition, and properties of cast complex alloys of Ni-Nb, 26:46035
- KHIM, Yong H., Organic disulfides and related substances. 33. Sodium 4-(2-acetamidoethylthio)butanesulfinate and related compounds as antiradiation drugs, 26:45290
- KHLESKOV, V. I., Reaction $\gamma + \gamma \rightarrow \pi + \pi$ and electromagnetic difference for pion masses, 26:46519 (JINR-P2-6333)
- KHOE, T. K., "FORGY": a companion computer program of "TRIM" to calculate forces and energy in electromagnets, 26:47599 (ANL-TKK/RJL-2)
- KHOLIN, V. M., Review and analysis of the latest data on measuring fundamental physical constants, 26:46439
- KHOLODOV, A. G., Solubility of Ce, Pr, and Nd in Cr containing O, 26:46034
- KHOOT, T. L., Mixing of two-proton bands with two-neutron bands in ^{176}Hf , 26:46992
- KHOSLA, M. M. L., Selective extraction of Ga with N-benzylaniline in chloroform from hydrochloric acid media, 26:45284
- KHRUSTALEV, O. A., N. N. Bogolyubov's transformation in strong coupling theory. II, 26:46691
- KHUNDZHU, T. G., Estimation of the turbulent diffusion coefficient and vertical component of wind velocity according to natural radioactivity distribution, 26:45522
- KHURSHUDYAN, L. S., Preamplifier for Cherenkov spectrometer, 26:45608
- KIBA, Nobutoshi, Polarographic determination of Sn in Cu-, Al-, and Zr-base alloys, 26:45256
- KICHIGIN, G. N., Solitons in a two stream plasma, 26:47202
- KIEFER, H., Radiation protection measurement, 26:45808
- KIEFHABER, E., Influence of fission neutron spectra on integral nuclear quantities of fast reactors, 26:47585 (EURFNR-1008)
- KIKUCHI, Kozo, Transformation properties of operators in local field theory, 26:46745
- KILCHER, P., Decay of ^{155}Dy , 26:46986
- KIM, Hyunyoung, Microwave spectrum, structure, and dipole moment of ethylene sulfone, 26:45291
- KIMURA, Iwane, Cherenkov instability and vlf emissions generated outside the plasmopause, 26:46318
- KING, Ivan R., Structure of the Coma Cluster of galaxies, 26:46242
- KING, J. C., Rapid annealing of frequency change in crystal resonators following pulsed x-irradiation, 26:45665 (SC-DC-721240)
- KING, Thomas Richard, Study of ^2H and ^3He elastic scattering and the reaction $^2\text{H}(^3\text{He}, ^4\text{He})^1\text{H}$ between 17.5 and 44.1 MeV, 26:46871
- KINGSLEY, R. L., $\Delta I = 1/2$ rule in the three-triplet and paraquark models, 26:46580
- KINTNER, Eric C., Structure of the Coma Cluster of galaxies, 26:46242
- KIPLINGER, Dale V., Pulsed method for impregnation of graphite, 26:46088
- KIRALYHIDI, L., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- KIRGINTSEV, A. N., Substance purification by zone melting in a rotating container, 26:45276
- KIRK, John H., Body weight decreases in some proton exposed primates, 26:45841
- Neuropathological evaluation of monkeys exposed to body-alone x-radiation, 26:45850
- KIRKLAND, Leland Rose Jr., Electron irradiation effects in aluminum single crystals, 26:47229
- KIRKLIN, C. W., Cumulative dose from diagnostic irradiation, 26:45799
- KIRSANOVA, T. S., Temperature dependence of the work function of Ba- V_{110} system, 26:47257
- KIRSCHNER, H., K conversion coefficient near threshold: /sup 191mOs, 26:47005
- KISELEV, V. G., Decay of isomers ^{126}I , ^{128}Sb and quasiparticle levels ^{126}I , ^{128}Sb with large spin, 26:46944
- KISELEVA, L. S., Electrolytic deposition of W-Co alloys, 26:45977
- KISELEVA, Z. P., Elastic scattering of protons by deuterons at energies of 14, 16, and 17.6 MeV, 26:46863
- KISO, Yoshiyuki, Direct measurement of γ -ray energies and half-life of ^{134}Cs , 26:46971
- KISS, I., Investigations on the radiosensitivity of yeasts, 26:45749
- KITAMURA, Teruo, Determination of small amounts of molybdenum based on its catalytic effect in Methylene Blue-hydrazine sulfate reaction, 26:45227
- KLASSEN, L., Hematopoietic and immunological alterations in mice produced by ^{90}Sr -induced marrow hypoplasia, 26:45896
- KLEPPER, O. H., Heat transfer performance of short twisted tapes, 26:47312 (CONF-720802-1)
- KLEVALIN, V. A., Performance logic of spark chamber information read-out by television automation and data link to a computer, 26:45650
- KLIMENKO, E. Yu., Cryogenic problems of forced cooled superconducting systems, 26:46839
- KLINE, D. E., Research on graphite. Final report for work completed during the period 1954 to 1972, 26:46078 (NYO-1710-106)
- KLOPPER, E., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- KLOTZ, A. H., Revised nonsymmetric unified field theory, 26:46694
- KLUCHAROV, A. P., Elastic scattering of polarized protons by ^{20}Si and ^{40}Ca , 26:46924
- KLUK, A. S., Band mixing in ^{156}Gd and ^{158}Gd , 26:46996
- KLYUCH, V. E., Dose to the epithelium of the upper respiratory tract from radon and its daughter products, 26:45861
- KNAELMANN, M., Dependence of iodide sorption on various soil components, 26:45528
- KNAPP, B. C., Experimental study of the decays $\text{K}/\text{sub S L}/ + \gamma\gamma$, 26:46434
- KNAPP, G. S., Determination of the electron-phonon enhancement factor from specific-heat data, 26:47262
- KNAPP, Gillian Revill, High-frequency resolution H I line observations of interstellar dust clouds, 26:46247
- KNAUER, J. B., Fabrication of ^{252}Cf neutron sources at Oak Ridge National Laboratory, 26:45714 (CONF-720902-2)
- KNEIPP, Kenneth George, Kinetic isotope effects in free radical chemistry, 26:45342
- KNIGHT, James A. Jr., Aerosols produced by x-rays, 26:46082
- KNIGHT, R. W., Evaluation of absorber materials performance in HFIR control cylinders, 26:47587
- KNIGHTON, James B., Mixer-settler apparatus, 26:45451
- KNOBLOCH, V., Sorption of polyvalent elements on silica gel. IV, 26:45391
- KNOLL, Kenneth C., Method of disposing of radioactive organic waste solutions, 26:45386
- KNOSPE, W. H., Effect of various cell-free organ extracts on hematopoietic stem cells, 26:45847
- KNOWLES, S. H., Microwave celestial water-vapor sources, 26:46206
- KOBAYASHI, Akizo, Urbaryon rearrangement and inelastic two-body processes, 26:46647
- KOBELT, M., First forbidden beta decay of ^{209}Pb , 26:47007
- KOBILOV, O. Sh., ^{135}Ba levels, 26:46945
- KOBYLINSKII, N. A., Threshold behavior of DAMA, 26:46774 (ITF-72-44-E)
- KOBZENKO, G. F., Alloys of chromium with terbium, 26:46033
- Solubility of Ce, Pr, and Nd in Cr containing O, 26:46034
- KOCHETKOV, N. K., 1,6-glucodialdose as one of radiolysis products of D-glucose-6-phosphate, 26:45337
- KOCHI, R., Conversion of bremsstrahlung photo neutron mean energies to monochromatic photo neutron mean energies, 26:46953
- KOCHKIN, Dmitry Alexandrovich, Method of modifying polymeric materials, wood, and articles made therefrom by radiation-initiated graft copolymerization thereof, 26:45355
- KOENIG, L. A., Environmental burden of tritium, 26:45509

- KOETZ, U., Missing-mass spectrum near 960 MeV from $\pi^-p \rightarrow n + (\text{mm})$, 26:46430 (COO-1195-241)
- KOJIMA, H., Persistent current states in rotating superfluid helium, 26:46801
- KOJIMA, Kazuhiko, Quantitative evaluation of radioisotope distribution in vivo by isosensitive scanner plus 4096-word multi-channel analyzer coupling, 26:45601
- KOK, L. P., S-matrix pole trajectories in the N/D approach, 26:46789
- Vertex parameters and coupling constants for the virtual decay of light nuclei, 26:47063
- KOKENGE, Bernard R., Radioisotopic fuel capsule, 26:45708
- KOKUBUN, Susumu, Association between quasi-periodic vlf emission and micropulsation, 26:46319
- vlf emissions during post breakup phase of polar substorm, 26:46285
- KOLACHKOVSKI, A., α -decay of neutron-deficient isotopes of Ra, 26:47004
- KOLESIKOV, E. M., Miniature low-background counters for measurement of ^{37}Ar and ^{39}Ar , 26:45614
- KOLFF, W. J., Biomedical engineering support. Progress report, 26:45700 (COO-2155-4)
- KOLISHNYK, Z., Bruce auxiliary steam supply, 26:45695
- KOLOMIETS, V. M., β, γ transitions and structure of anti-analog states, 26:47050
- Excitation of input states during thermal neutron capture by ^{66}Zn , ^{70}Ge , and ^{80}Se , 26:46937
- KOLOTOILOVA, V. G., Importance of tissue-equivalent detectors used in radiometry, 26:45619
- KOLYBASOV, V. M., Conference on problems of modern nuclear physics, Novosibirsk, USSR, 6/70, 26:47086
- Mechanism of direct processes at high energies, 26:47101
- KOMEL'KOV, V. S., Generation of shock waves by exploding current sheath, 26:47186
- KOMEM, Ya., Direct observation of grain boundary dislocation climb in ion-irradiated gold bicrystals, 26:47241
- KOMPANEETS, A., Self-consistent field method as applied to the nucleus, 26:47039 (BNL-tr-488)
- KONDO, Toshikiko, Identification of reaction intermediates by microwave spectroscopy. The catalytic reactions between propylene and deuterium over zinc oxide, 26:45303
- KONDO, Yoshio, Solvent extraction of Eu by Versatic Acid 911, 26:45277
- KONDRAT'EV, A. A., Electron energy distribution and striations in the Langmuir-paradox region, 26:47133
- KONDRATYUK, L. A., Many-neutron resonances, 26:46644
- KONIN, A. D., New method of determination of physical-chemical properties of matter, 26:47618 (JINR-P14-6407)
- KONKASBAEV, I. K., Solitons in a two stream plasma, 26:47202
- KONOBEEVSKII, E. S., Excitation cross section of first 2+ levels of Ni and Zn nuclei, 26:46936
- KONRADY, John Andrew Jr., Retardation effects in low-energy atom-atom scattering, 26:46377
- KONYAEV, V. F., Conversion transitions in ^{226}Ra , 26:47028
- KOONS, H. C., Observation of very-low-frequency whistler-mode waves in the region of the radiation-belt slot, 26:46312
- KOPP, J., Discontinuous flux flow in superconducting NbSe_2 , 26:46831
- KORNIENKO, T. G., Determination of alloying additives in artificial fluorite, 26:45232
- KOROLEVA, V. P., Use of solid track detectors in reactor experiments, 26:45610
- KOROTKIN, Yu. S., Study of transuranium element hydrolysis. I, 26:45366 (JINR-P6-6402)
- KORSAKOVA, N. V., Photometric determination of zirconium in natural materials with the reagent picramineepsilon, 26:45233
- KORYAKIN, Yu. I., Methodology employed in economic calculations relating to nuclear power stations, 26:47387
- KORYTOWSKI, Adam, Hydrogen generators for neutron tubes, 26:45942
- KORZHAVIN, V. M., Observation of second compression in the final stage of a discharge of the plasma focus type, 26:47139
- KORZIN, E. G., Recording of alpha particles by solid track detectors based on cellulose, 26:45615
- KOSHKAREV, D. G., Instability of axial oscillations in an electron-ring accelerator, 26:46142
- KOSIARA, Andrzej, Search for parity and isobaric spin violating decay of the 3.56 MeV level in ^6Li , 26:46900
- KOSIN, L. F., Thermodynamic properties of solutions in system $\text{In}-\text{Sn}-\text{Hg}$, 26:46004
- KOSS, D. A., Structure and properties of solid solutions. Technical progress report, September 1, 1971--June 15, 1972, 26:45991 (COO-916-25)
- KOSTKA, P., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- KOSYAK, Yu. G., Lifetime of 1.013-MeV state in ^{27}Al , 26:46905
- KOTAKE, Yashige, Electron spin resonance study of the far-ultraviolet photolysis of some inorganic and organic compounds, 26:45323
- KOTLEWSKI, A., Charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\gamma$, 26:46482
- Measurement of the charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, 26:46483
- KOTLIN, V. P., Identification of an AmCl_2^{3-} octahedral complex from Am(III) absorption spectrum in ethanol solutions of HCl , 26:45376
- KOTOV, G. V., Personal photodisimetry in radiodiagnosis and radiation therapy rooms, 26:45629
- KOURI, Donald J., Differential- and total-cross-section calculations for models of e^-H_2 collisions, 26:46399
- KOVAL, A. D., Effect of micro-additives of rare earth metals on the grain boundary energy of nickel, 26:45999
- KOVALENKO, V. G., Characteristics of charged-particle detection sections having channel electron multipliers, 26:45645
- Multichannel modular spectrometer for low-energy electrons and protons, 26:45642
- KOVALEV, A. I., Introduction of correction for nonlinearity of direct-current Q-meter, 26:45680
- KOVALEV, N. N., Electrical conductivity, thermo-emf, and thermionic emission of BaO and SrO crystals, 26:47249
- KOVALEVSKII, A. V., Thermodynamic properties and phase composition of $\text{La}-\text{Zn}$ alloys, 26:46001
- KOVESI-DOMOKOS, S., Inelastic e^-p scattering and the polarization contribution to H hyperfine structure, 26:46523 (NYO-4076-17)
- KOZIK, M., Effect of ionizing radiation on enzyme activity in the neuroglia of the developing rabbit brain, 26:45836
- KOZLOV, A. P., Improvement of electron beam shaper in B5M-25 model betatron, 26:46132
- KOZLOV, B. D., Collective interactions of nucleons of high-energy cosmic heavy nuclei, 26:46461
- KOZLOV, Stanislav Fedorovich, Nuclear radiation detection device utilizing diamond detector with injecting and blocking contacts, 26:45597
- KRAAK, J. C., Rapid separation of metal chelates by column liquid-liquid chromatography using ultraviolet detection, 26:45281
- KRAITOR, S. N., Comparison of neutron and gamma-ray personnel dosimeters at critical assemblies, 26:45630
- KRAJEWSKI, B. T., Updating stratospheric inventories to July 1971, 26:45518 (HASL-257, pp I.33-50)
- KRAMER, D., Studies of void formation in proton-irradiated type 316 and titanium-modified 316 stainless steels, 26:46044 (AI-AEC-13033)
- Swelling as a function of displacement damage in proton-irradiated 316 stainless steel, 26:46054 (CONF-710601-pp 511-21)
- KRAMER, G., Diffraction-dissociation model for high-energy electroproduction, 26:46517 (DESY-72/23)
- KRASOVITSKII, V. B., Excitation of a regular plasma wave by a modulated beam with high energy density, 26:47201
- KRASZNAI, I., New two-channel scanner. II, 26:45607
- KRAUSS, Morris, Coupled multiconfigurational self-consistent-field method for atomic dipole polarizabilities. I. Theory and application to carbon, 26:46419
- KRENZ, W., Measurement of slope parameter in decay $K/\text{sub } L^0 \rightarrow \pi^+\pi^-$, 26:46494
- KREPELKA, Jiri, Radioactivity of waste waters in Nuclear Research Inst., Rez, and environment, 26:45565
- KREY, P. W., ^{90}Sr fallout in Utah, 26:45525 (HASL-257, pp I.71-94)
- Plutonium fallout in Utah, 26:45526 (HASL-257, pp I.95-118)
- Updating stratospheric inventories to July 1971, 26:45518 (HASL-257, pp I.33-50)
- KRIEGER, J. B., Analytic calculation of the coherent atomic scattering factor for the ground state of the helium isoelectronic sequence, 26:46359
- Variational principles for single-particle expectation values in the Hartree approximation, 26:46417
- KRIGAS, Thomas Michael, Electron spin resonance study of radicals formed by high-energy irradiation of K_2PtCl_6 and K_2PtCl_4 , 26:45341
- KRIKORIAN, Oscar H., Corrosion and scaling by steam in nuclear geothermal power plants, 26:45487 (UCRL-73939)
- KRIMIGIS, S. M., Comparison of measurements of the charge spectrum of solar cosmic rays from nuclear emulsions and the Explorer 35 solid-state detector, 26:46330
- KRISCH, A. D., 200-GeV proton-proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)
- KRISHNAMURTHY, M. V., EPR study of VO^{2+} in $\text{UO}_2(\text{NO}_3)_2 \cdot \text{X} \cdot 6\text{H}_2\text{O}$ single crystal, 26:45372
- KRISTIAN, J., Optical search for Centaurus X-3, 26:46213
- KRITSKII, G. A., Characteristics of nucleotide of DNA in radiation injury, 26:45856
- KRIVAN, V., Systematics of excitation functions for fast neutron induced (n,p)- and (n, α)-reactions, 26:47066
- KRIVISKII, A. S., Lethal and mutagenic effect of ionizing radiations with different LET on extracellular bacteriophages, 26:45746
- KROKHIN, O. N., Observation of second compression in the final stage of a discharge of the plasma focus type, 26:47139
- KRONBERG, P. P., Polarization of radio sources at $\lambda 49 \text{ CM}$ and $\lambda 243 \text{ CM}$, 26:46204
- KRONGAUZ, A. N., Importance of tissue-equivalent detectors used in radiometry, 26:45619
- KROPAC, B., Remark on neutron slowing down, 26:46856
- KRPIC, D., Effective $g/\text{sub } s/\text{factor}$ for odd proton ^{153}Sm

- nucleus, 26:46994
- KRUBASIK, E., Amplitude analysis of πN Coulomb interference data, 26:46625
- KRUEGER, H., Microdosimetry measurements of fast neutrons with a tissue-equivalent proportional counter of 1 cm diameter, 26:45626
- KRUGER, Paul, Design of waterway connecting Orinoco and Rio Negro Rivers in Federal Territory of Amazonas, Venezuela, 26:45475 (NP-19361)
- KRUMPELT, M., Containment of fission product iodine in the reprocessing of LMFBR fuels by pyrochemical reactions, 26:45421
- KRUPA, E., Multipole order of transitions and characteristics of excited state of ^{147}Eu , 26:46954
- KRYUCHKOV, I. I., Electrical insulation in electrostatic analyzers, 26:45687
- KRZYWICKI, A., Large angle scattering of hadrons, 26:46591
- KUBO, Sohei, Effects of cepharanthine on fluctuation in white blood cell count by radiotherapy for gynecologic cancer, 26:45805
- KUCERA, J., Comments on diffusion of cobalt in plutonium by M. J. Daphy, M. P. Dariel, and A. Rosen in *Scripta Met.*; 5: 191-4(1971), 26:46024
- KUCHAR, Karel, Bubble-time canonical formalism for geometrodynamics, 26:46751
- KUCHNIR, M., Absence of a quadratic term in the ^4He excitation spectrum, 26:46799
- KUDO, Hiroshi, Effects of pre-irradiation treatments on chemical behavior of ^{64}Cu recoil atoms in neutron-irradiated Cu phthalocyanine, 26:45356
- KUDRYASHOV, L. I., 1,6-glucodialdose as one of radiolysis products of D-glucose-6-phosphate, 26:45337
- KUDRYAVTSEV, I. V., Effect of mechanical surface hardening on the low-cycle fatigue strength of titanium alloys, 26:46009
- KUEHNLE, J. H., Thermal shock deburring, 26:47609 (BDX-613-642(Rev.))
- KUEHMEL, H., Circular polarization of γ rays in ^{18}F , 26:46877
- Resonance scattering theory and the structure of compound states, 26:47068
- KUENZLE, H. P., Canonical dynamics of spinning particles in gravitational and electromagnetic fields, 26:46753
- KUHLMANN, E., Excited states of ^{21}Ne , 26:46907
- KUIKEN, R., Fission of aligned ^{233}U nuclei by neutrons from 0.4 to 2000 eV, 26:47026
- KULESHOV, S. P., Operator method of solving the quasipotential equations and a conception of straight-line paths at high energies, 26:46700 (JINR-P2-6437)
- KULIBABA, V. I., Angular and energy distributions of protons in (γp) and (γp) reactions on ^{12}C nucleus for 130 MeV excitation energy, 26:46888
- KULISH, Yu. V., Electroproduction of π mesons near threshold and current algebra, 26:46549
- Tensor meson photoproduction and $\text{SU}(6)$ symmetry, 26:46684
- KUL'KIN, A. G., Induced radiation from neutrons moving through a medium in a magnetic field, 26:46541
- Spin effects in theory of synchrotron radiation due to motion of an electron in a medium, 26:46548
- KUNDRA, A. A., Spectral and scintillation properties of the pyrimidine derivatives, 26:45606
- KUNDU, K. P., Radiolysis of aqueous solutions of nickel(II) ethylenediamine tetraacetate, 26:45350
- KUNKEL, W. E., Optical observations of supernova in NGC 5253, 26:46187
- KUPRINA, V. V., Formation of intermetallic phases in system Ir--Ta--Ni, 26:46010
- KUPRYASHKIN, V. T., Decay of ^{198}Au , 26:47003
- Electron conversion spectrum of ^{182}Re , 26:46978
- KURAMOTO, Yasuo, Development of 50Hz 44 inch turbine blade, 26:45454
- KURDYUMOV, A. V., Energy spectrum and crystalline structure of wurtzite-like modification of boron nitride, 26:47255
- KURINNYI, V. I., Investigation of the injection characteristics of symmetrical pin-diodes prepared by the method of ion implantation, 26:45678
- KURLYANDSKII, A. S., Isotopic structure of the giant dipole resonance and the isospin frequency in the ground state of the nucleus, 26:47053
- KURNOSOVA, L. V., Removable current lead for power supply in superconducting magnetic systems, 26:45679
- KURODA, M., Polarization in inclusive reactions, 26:46595
- KURODA, Masaaki, Parity tests in elastic $e-p$ scattering at large angles, 26:46562
- KUROKAWA, Kazuo, Anion exchange resin separation of Gd, Eu, Sm, Nd, and Pr, 26:45280
- KURYSHKIN, V. V., Possible generalization of nonrelativistic quantum mechanics, 26:47297
- KUSAYANAGI, Eichi, Flux flow in nonideal Type-II superconductors, 26:46819
- KUTSCHERA, W., Neutron deficient even-even isotopes of Kr, Se, Ge, and pseudo SU_3 coupling, 26:46931
- KUTTER, G. S., Studies of hydrodynamic events in stellar evolution. II. Dynamic instabilities in stellar envelopes, 26:46176
- KUWATA, Keiji, Electron spin resonance study of the far-ultraviolet photolysis of some inorganic and organic compounds, 26:45323
- KUZ'MIN, V. V., Comparison of neutron and gamma-ray personnel dosimeters at critical assemblies, 26:45630
- KUZNETSOV, V. A., Use of solid track detectors in reactor experiments, 26:45610
- KUZNETSOV, V. V., Study of the decay $^{140}\text{Tb}(T_{\text{sub}}^{1/2} = 66 \text{ min})$ and $^{150}\text{Tb}(T_{\text{sub}}^{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- KUZNETSOVA, M. Ya., Study of the decay $^{140}\text{Tb}(T_{\text{sub}}^{1/2} = 66 \text{ min})$ and $^{150}\text{Tb}(T_{\text{sub}}^{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- KUZNETSOVA, R. I., Surfaces of solidus and liquidus in system Th--Zr--U, 26:46003
- LACINA, J., Longitudinal waves in plasmas and physical mechanism of Landau damping, 26:47191
- LACOSTE, Robert, Comparative effects of ion bombardment and partial pressures on sheet polyethylene, 26:46081
- LACOURLY, G., Method of evaluating chemical pollution levels in aquatic media and food chains, 26:45547
- LADES, Heinrich, Beta decay asymmetry and nuclear magnetic resonance of ^{28}Al , ^{38}Cl , and ^{116}In produced by capture of polarized neutrons, 26:46911
- LAFONTAINE, I., Plutonium fuel production plant and breeder projects, 26:45974
- LAGARDE, G., Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in gas phase, 26:45361
- Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in liquid phase, 26:45362
- LAGARRIGUE, M., Particle physics, 26:46509
- LAI, Kwan-wu, Pion exchange in the reaction $\pi p + pN$ at 6 GeV/c, 26:46481
- LAIDLER, J. J., Suppression of void formation in stainless steel, 26:46049 (CONF-710601-, pp 174-86)
- LAL, K. B., Cylindrical wave solutions of Bonnor's and Schroedinger's unified field equations, 26:46695
- LALLI, M. G., Measurement of ^{137}Cs and ^{90}Sr in soil, 26:45263 (RT/PROT-72-11)
- LAM, Siu Kee, Two-dimensional transport problems in a quarter space, 26:46854
- LAM, W. C., Nuclear γ rays associated with stopped kaons, 26:46940
- LAMB, Eugene, ORNL isotopic power fuels quarterly report for period ending March 31, 1972, 26:45703 (ORNL-4796)
- LAMB, R. C., Observation of the $M(940)$ and $\delta(963)$, 26:46475 (IS-2886)
- LAMBERT, Irma, Isotopic enrichment of hydrogen for heavy water production, 26:45696
- Method of preparing secondary alkaline amine derivatives, 26:45289
- LAMBERT, M., Generating function method and developments in quasi-boson, 26:47286 (LYCEN-7202)
- LANDER, G. H., Magnetic phase diagram of the UAs--US system, 26:47261
- LANDSHOFF, P. V., Deep inelastic electroproduction and related processes, 26:46515 (DAMP-71/24)
- LANE, A. M., Theory of anomalies observed in (d,p) excitation curves, 26:47058
- LAPIDUS, L. I., Electromagnetic effects in nucleon polarization at meson-nucleon scattering, 26:46520 (JINR-P2-6438)
- LAPIN, N. I., Photoproduction of charged pions by nuclei at $E_0 = 1160 \text{ MeV}$, 26:46441
- LAPLANCHE, F., $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
- $\pi^+\pi^-$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
- LARI, R. J., "FORGY": a companion computer program of "TRIM" to calculate forces and energy in electromagnets, 26:47599 (ANL-TKK/RJL-2)
- LARIKOV, L. N., Disintegration of supersaturated solid solutions according to a heterogeneous mechanism, 26:46032
- LARSON, B. C., Effect of irradiation temperature on size of defect aggregates in copper, 26:46060 (CONF-710601-, pp 672-83)
- LARSON, H. A., On-line reactivity feedback analysis of EBR-II by rod drop, 26:47307
- LARSSON, Sven, Correlation effects in atomic systems, 26:46409
- LASHMORE-DAVIES, C. N., Collisionless heating of a plasma by means of oscillating external currents, 26:47107
- LASIJO, Romedon, Low-lying states and configurations in ^{138}La , 26:46964
- LASSEN, K., Critical component of the interplanetary magnetic field responsible for large geomagnetic effects in the polar cap, 26:46266
- LASTUSHKIN, N. I., Substance purification by zone melting in a rotating container, 26:45276
- LASZLO, Andrassy, Calibration of radioactive logging apparatus and neutron modeling, 26:45653 (CSIRO-Trans-10705)
- LATHROP, K. D., Discrete-ordinates methods for the numerical solution of the transport equation, 26:47211
- LATSIS, R. V., Role of thiols in radiosensitivity of sea urchin eggs, 26:45882
- LATYPOV, Z. Z., Oscillations of charge-exchange cross section in system Li^+-He , 26:46368
- LATYSHEV, G. D., Electron conversion spectrum of ^{182}Re , 26:46978

LATYSHEV, I. N., Type A stars in the Ursa Major stream, 26:46164

LATYSHEVA, V. I., Phase diagram of the niobium--gallium system, 26:46018

Variations in the critical temperature of superconducting compounds Nb₃Al and Nb₃Al/sub x/Ge/sub 1-x/, 26:46021

LAURENS, P., Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)

LAURITZEN, T., Deuteron-generated void formation in austenitic stainless steels, 26:46053 (CONF-710601-, pp 479-98)

LAURSEN, H. G., Large scale characterization tests of gelled slurry blasting agents, 26:45484 (SC-DC-722296)

LAUTRUP, B. E., Recent developments in the comparison between theory and experiments in quantum electrodynamics, 26:46710

LAVERLOCHERE, J., Activation analysis with fast neutrons, 26:45209

Use of computers in activation analysis, 26:45211

LAVLOV, V. S., Electron transfer during zone melting of yttrium, 26:45975

LAWRENCE, L. A., Effects of pore structure on the melting heat rating of oxide fuels, 26:45930 (HEDL-TME-72-81)

LAYSSAC, J., e⁺e⁻ annihilation into hadrons, 26:46540

LAYER, J. G., Measurement of the charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, 26:46483

LEADABRAND, R. L., Chatanika, Alaska, auroral-zone incoherent-scatter facility, 26:46283

LEBDA, Nancy Jean Abbott, Effect of varying doses of cortisol on the incorporation of ³H--thymidine into rat liver DNA and on the synthesis of five plasma proteins in the isolated perfused regenerating rat liver, 26:47646 (UR-3490-131)

LEBEDEV, B. A., Performance logic of spark chamber information read-out by television automation and data link to a computer, 26:45650

LEBEDEV, K. P., Structure, phase composition, and properties of cast complex alloys of Ni--Nb, 26:46035

LEBEDEV, V. A., Thermodynamic properties and phase composition of La--Zn alloys, 26:46001

LEBEDEV, V. N., Intranuclear cascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)

LE BEYEC, Y., Transfer reactions induced by ⁸⁴Kr on ²³²Th, 26:47035

LEBRUN, C., Uranium isotope separation studies in France, 26:45691

LEBRUN, J. P., Factorization in a multiperipheral configuration, 26:46726

LECOUTURIER, D., Decay of ¹⁵⁵Dy, 26:46986

LEE, B. W., Process $\nu/\text{sub } \mu/ + p + \nu/\text{sub } \mu/ + p + \pi^0$ in Weinberg's model of weak interactions, 26:46614

LEE, Byoung Whie, Activated sintering mechanism of UO₂ powder, 26:45956

LEE, D. M., Spin-wave phenomena in liquid ³He systems, 26:46802

LEE, F., Conversion of bremsstrahlung photo neutron mean energies to monochromatic photo neutron mean energies, 26:46953

LEE, George, Laser power stations in orbit, 26:45428

LEE, H., Transmission of low-energy x-ray through various locally available materials, 26:47205

LEE, Paul Lung Sang, Isotope shift studies in the rare earths and lead, 26:46973

LEE, R. S., X-ray-induced photoconductivity in dielectric films, 26:45668 (UCRL-73974)

LEE, S., High-speed photography of a plasma focus, 26:47121

LEE, W., Experimental limit on the neutral current in the semileptonic processes, 26:46453

LEE, William R., Mutagenic effect of transmutation from radionuclides incorporated into DNA of Drosophila melanogaster. Progress report, September 1970--May 1972, 26:45912 (DRO-3728-5)

LEER, Egil, Problems in solar-system plasma physics, 26:46331

LEFORT, Guy, Method of extracting fuel from nuclear fuel elements, 26:45422

LEFORT, M., Heavy ion nuclear reactions, 26:47056

LEFRANCOIS, J., $\pi^+\pi^-$ production by e⁺e⁻ annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)

$\pi^+\pi^-$ production by e⁺e⁻ annihilation at ω mass, 26:46427 (CONF-710816-13)

LE GALLIC, Y., Metrology of ionizing radiations, 26:47219

LEGEAY, Biological effects of the intensity of irradiation, 26:45816 (FRNC-R-7)

LE GUYADER, J., Further results from multi-Regge models applied to $pp \rightarrow p\bar{p}\pi^+\pi^-$ reaction at 13.1 GeV/c, 26:46656

LEHANE, J. A., Parametric amplification of Alfvén waves, 26:47187

LEIN, E. L., Low-energy-electron spectrometer having a multiplate analyzer, 26:45643

LEIPUNSKAYA, D. I., Gold assay on geological samples by neutron activation with preliminary crucible fusion, 26:45200

LEKSIN, G. A., Investigation of many-nucleon resonances, 26:46658

LEMONDS, Alice J., Conditions for the removal of chromate from cooling water, 26:47619 (KY-642)

LE NADAN, Andre, Absolute measurement of lifetime of ⁴D level of He atom using electron--photon coincidence method, 26:46345

LENGER, V., Measurement of ²²⁶Ra in mice, 26:45920

LENIART, Daniel S., Powder ENDOR line shapes: nuclear relaxation induced by motion of nearby electron spins, 26:47279

Proton ENDOR of γ -irradiated Y-type zeolites, 26:45324

LEONARD, Regis F., Particle--hole states in ¹²⁰Sn, 26:46943 (N-72-18705)

LEONI, G., Analog computer models for safety evaluation of CART fuel test facility, 26:47580 (CISE-R-259)

LEPLAE, L., Time-dependent phenomena in weakly-coupled superconductors, 26:46827

LEPRINCE, P., Parametric coupling between electronic and ionic wave modes in an auto-resonant plasma, 26:47190

LERMAN, Abraham, ⁹⁰Sr in the Great Lakes: concentration-time model, 26:45532

LETELLIER, P. R., Recombination products from the radiolysis of tricaproin, 26:45321

LEUTERER, F., Cyclotron harmonic waves in an inhomogeneous plasma. I, 26:47188

Cyclotron harmonic waves in an inhomogeneous plasma. II, 26:47189

LEVEDAHL, William J., Thermal control and power flattening for radioisotopic thermodynamic power system, 26:45427

LEVENES, Guy, Improvements to equipments for measuring thickness by backscattering, 26:45656

LEVIN, F. S., Validity of DWBA for deuteron stripping. I, 26:47054

LEVINSEN, M. T., Sinusoidal current-phase relation for superconducting thin-film microbridges, 26:46806

LEVKOVSKII, V. N., Decay of isomers ¹²⁶ ¹²⁸ ¹³⁰Sb and quasiparticle levels ¹²⁶ ¹²⁸ ¹³⁰Te with large spin, 26:46944

LEVSKII, L. K., Age of cuprous Ukokan series (Dlekmo--Vitim mountain system) by uranium--lead method, 26:45569

LEWIS, Alan L., Perturbation treatment of high energy electron diffraction from imperfect crystals, 26:46510 (CALT-767-P-3-22)

LEWIS, B. M., 21 cm observations of NGC 45, 26:46252

LI, Che-Yu, Theoretical study of the swelling of fast-neutron-irradiated materials, 26:46063 (CONF-710601-, pp 798-824)

LIBBY, L. M., Dynamic interpretation of baryon and meson mass formulae, 26:46594

LIBERMAN, A. D., $\pi^+\pi^-$ production by e⁺e⁻ annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)

$\pi^+\pi^-$ production by e⁺e⁻ annihilation at ω mass, 26:46427 (CONF-710816-13)

LIBERMAN, Daniel Franklin, Effects of chloramphenicol and cycloheximide on the cell growth cycle of mouse leukemic cells (L5178Y), 26:47647 (UR-3490-132)

LICHTIN, N. N., Electron and hydrogen atom attachment to aromatic carbonyl compounds in aqueous solution. Absorption spectra and dissociation constants of ketyl radicals, 26:45354

LIE, K. H., Contrast patterns produced by self-interstitial atoms in field ion microscope images of a b.c.c. metal, 26:46029

LIGON, Jerry Lloyd, Plasma injection and collision phenomena in an octupole magnetic field, 26:47110

LILIENFELD, A., Irradiation in the epidemiology of leukemia among adults, 26:45798

LIMPAECHEER, Rudolf, Parallel magnetic multipole confinement of a magnetic-field-free plasma, 26:47111

LIMSTRAND, J., Energy levels in ²³¹Th and ²³³Th, 26:47029

LIN, Chun C., Excitation of the electronic states of the nitrogen molecule by electron impact, 26:46391

LINCHEVSKAYA, G. A., Measurement of absorbed doses with fluid ionization chambers, 26:45634

Phantom-dosimeter with liquid-filled ionization chamber, 26:45621

LINCOLN, T. A., Pathophysiology and diagnosis of internal contamination, 26:45888 (CONF-720520-1)

LIND, N. C., Consistent theory for spherical shells in equilibrium, 26:45453

LINDEKEN, C. L., Environmental levels of radioactivity in the vicinity of the Lawrence Livermore Laboratory January through December 1971, 26:45513 (UCRL-51242)

LINDELL, B. G., Assessment and control of environmental contamination, 26:45554

LINDENBAUM, A., Metabolism of monomeric and polymeric Pu in the rabbit. Comparison with the mouse, 26:45917

LIPPMANN, G., Local magnetic field distributions in superconducting Nb at 4.2 K, 26:46826

LISITSA, V. S., Hydrogen atom in a rotating electric field, 26:46352

LISKA, Donald J., Multipactoring electron gun for high duty linacs, 26:46134

LISZT, Ferenc, Developments in geophysical logging instruments and methods, 26:45560

LITUNOVSKII, R. N., Compensated current-driven magnetic channel in a 240-cm isochronous cyclotron, 26:46128

Magnetic shield for a 240-cm isochronous cyclotron, 26:46129

LITVAK, M. M., Time-resolved diagnostic method for hydrogen plasmas, 26:47127

LIU, S. G., High-peak-power, high-efficiency microcircuit avalanche diode oscillators, 26:47624 (SC-CR-723142)

LIVINGSTON, M. Stanley, Origins and history of the Los Alamos Meson Physics Facility, 26:46119 (LA-5000)

- LIVOLANT, M., Study of vibrations of the centering parts of fuel element cartridges, 26:47314
- LJUNG, Donovan Allen, Experimental search for some rare K⁺ decay modes, 26:46454
- LLORÉ, A., K₂⁰ decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)
- LOBODYUK, V. A., Electron microscopic investigation of niobium--oxygen alloys with additions of zirconium and hafnium, 26:46039
- LOBODZIEC, Włodzimierz, Trial of correlating the blood picture with absorbed doses in persons exposed professionally to ionizing radiation, 26:45787 (TT-54002/5, pp 602-4)
- LOBOV, G. A., Limitations on currents of second class in process $v/\text{sub } \mu/ + n + p + \mu^-$, 26:46573
- LOCKWOOD, G. W., Near-infrared photometry of Mira variables, 26:46194
- Observations of the He II $\lambda 10124$ line in O and Of stars, 26:46180
- LODHI, M. A. K., Nuclear level sequence of single-particle energies in a momentum-dependent potential, 26:47071
- LOESCHER, D. H., Oxidation and etching of 80 percent Si--20 percent Ge, 26:45935 (SC-RR-710884)
- LOGAN, B. A., Optimisation of possible tests of the validity of time-reversal, 26:46677
- LOGINOV, V. F., Nature of a semi-secular cycle in the troposphere, 26:46296
- LOKTIONOV, V. A., Electron microscopic investigation of niobium--oxygen alloys with additions of zirconium and hafnium, 26:46039
- LOMBARDO, U., Superluminal frames and group of generalized Lorentz transformations in four dimensions, 26:46734
- LONDERGAN, J. T., Charge-dependent effects in the photodisintegration of ⁴He, 26:46866
- LONGAIR, M. S., Observational cosmology, 26:46256
- LONGUEUE, J. P., Low-lying positive-parity states of ³⁵Cl and ³⁶Cl from (p,d) reactions and shell-model interpretations, 26:46906
- LONSKIL, E. D., Possibilities of using radioisotope instruments in fabrication of high power plants of supercritical steam parameters, 26:45473
- LOOS, Eckhard, Apparatus for continuous measurement of active uptake of radioactive substances, 26:45592
- LOPEZ JIMENEZ, Jose, Table of cross section (n,p), (n, α), and (n,2n) reactions in steel components and other nuclear materials, 26:46918 (JEN-230-DMe/I-25)
- LORENTE, Ambrosio, Numerical simulation of evolution of an excited monochromatic wave in one-dimensional Maxwellian plasma, 26:47182
- LORENTE-ARCAS, A., Model for the hollow cathode discharge, 26:47140
- LORENTS, D. C., De-excitation of fast He, Ne, and Ar metastable atoms in various gases, 26:46394
- LORENZEN, J., Depth distribution studies of carbon in steel surfaces by means of charged particle activation analysis with an account of heat and diffusion effects in the sample, 26:45466 (AE-451)
- LOSKA, L., Determination of ash content of coal by neutron activation analysis, 26:45195
- LOWELL, J., Resistivity of the intermediate state of Type-I superconductors, 26:46823
- LOWY, D. N., Single-electron energies, many-electron effects, and the renormalized-atom scheme as applied to rare-earth metals, 26:47260
- LU, C. C., Level densities from spectra of different reactions populating the same nucleus, 26:46932
- Nondestructive test for measuring the state of heat treatment in closure welds, 26:45470
- LU, Shin-tsu, Ionizing radiation-induced thyroid carcinogenesis in the dog, 26:45819 (UR-3490-63)
- LUCA, I., Influence of neutron irradiation on seed germination and plant growth of tomatoes, 26:45773
- LUDEMAN, C. A., Energy shifts and relative intensities of K x rays produced by swift heavy ions, 26:46339
- LUDWIG, E. J., Energy levels of ³⁶K, 26:46904
- LUGIATO, Luigi A., Structure of relativity groups, 26:46669
- LUKASIEWICZ, M., Partial ionization cross sections of C dioxide by electrons (25 to 600 eV), 26:46384
- LUK'YANOV, L. A., Measurement of absorbed doses with fluid ionization chambers, 26:45634
- Phantom-dosimeter with liquid-filled ionization chamber, 26:45621
- LUMPKIN, Alexander Henderson, Development of a two-section Ge(Li) Compton polarimeter with applications in the 2s--1d shell, 26:46893
- LUNDEN, Arnold, Temperature dependence of the isotope effect for electromigration of rubidium ions in molten rubidium nitrate, 26:45294
- LUNDQUIST, S., Model studies of density fluctuations of small and large ions in the atmosphere, 26:46288
- LUTHER, G. G., Measurement of an optical frequency and the speed of light, 26:46733
- LUTON, J. M., Geomagnetic effect on the neutral temperature of the F region during the magnetic storm of September 1969, 26:46270
- LUTZ, Julie Haynes, Interstellar dust and distances to planetary nebulae, 26:46171
- L'VOV, S. N., Electrical and thermal conductivities of Re, Os, and Ir and some of their alloys with Th and La in wide temperature range, 26:47248
- LYALINA, L. I., Personal photodosimetry in radiodiagnosis and radiation therapy rooms, 26:45629
- LYNCH, John H., Correlation of irradiation data using activation fluences and irradiation temperature, 26:46072
- LYNCH, Robert D., Comparison of response spectra from Yucca Flat and Pahute Mesa underground nuclear detonations, 26:45480 (NVO-1163-TM-32)
- LYNDS, Roger, Ability of the luminosity-volume test to reveal the statistical evolution of the luminosity of quasi-stellar sources, 26:46208
- LYON, F., Harmonic reactor, 26:47590
- LYONS, Lawrence R., Pitch-angle diffusion of radiation-belt electrons within the plasmasphere, 26:46311
- M**
- MACADAM, K. B., Molecular-beam magnetic-resonance measurements of the anisotropies of the electric polarizabilities of H₂ and D₂, 26:46358
- MACCALLUM, M. A. H., Variational principles and spatially-homogeneous universes, including rotation, 26:46758
- MACCRAGH, Adolfo, Extraction of fission product noble metals from spent nuclear fuels, 26:45423
- MACDONALD, A. B., Lifetime of 975-keV, ³/₂⁺ state in ²⁵Mg, 26:46908
- MACDOUGALL, C. S., Remotized apparatus for determining oxygen in irradiated reactor fuels and cladding materials, 26:45224 (LA-DC-72-524)
- MACFADYEN, N. W., Nondegenerate complementary series of representations of group SU₂, 26:46672
- MACHIDA, Kazushige, Localized magnetic states in superconductor, 26:46809
- MACK, G., Currents, stress tensor, and generalized unitarity in conformal invariant quantum field theory, 26:46697 (DESY-72/20)
- MACLEAN, Lee M., Review of problems associated with acceleration measurements, 26:45674 (UCRL-73665)
- MACLEOD, J. M., Luminosity-spectral index relationship for radio galaxies, 26:46248
- MACMURDO, K. W., Mass yields from thermal-neutron induced fission of ²³⁵Cm, 26:47027
- MACPHERSON, H. G., Development of materials and systems for the molten-salt reactor concept, 26:47302
- MADAN, Rabinder N., Fermi pressure shift and electron--rare-gas-atom scattering lengths, 26:46401
- MADDING, R. D. Jr., Documentation of DOT-SP-5915 for 55/30 and 83/55 drum combination packaging, 26:46114 (MLM-1911)
- MADDOCK, B. J., Tubular Nb/Cu conductors for ac superconductive power transmission, 26:46844
- MADER, D. L., Fine structure of the helium negative ion, 26:46343
- MADUSKAR, K. D., Synthesis, identification, and purification of C₆ and C₇ fluorocarbons from radiolysis. Final report, 26:45316 (ORO-3909-3)
- MAECK, William J., System for total iodine retention, 26:45387
- MAEDA, Katsuki, Radiation-induced copolymerization of vinylphosphonic dichloride with methyl methacrylate and styrene, 26:45318
- MAENHAUT, W., Instrumental neutron activation analysis of germanium, 26:45218
- MAERKER, R. E., SDT1. Iron broomstick experiment, 26:47220 (ORNL-TM-3867)
- SDT2. Oxygen broomstick experiment, 26:47221 (ORNL-TM-3868)
- SDT3. Nitrogen broomstick experiment, 26:47222 (ORNL-TM-3869)
- SDT4. Sodium broomstick experiment, 26:47223 (ORNL-TM-3870)
- SDT5. Stainless-steel broomstick experiment, 26:47224 (ORNL-TM-3871)
- MAGARIK, K. N., Radiometric unit for measuring wear of materials and machine parts, 26:45657
- MAGER, T. R., Discussion on fracture mechanics concepts, 26:45996 (WCAP-7841)
- MAH, S. Q., Electron temperature determination from the rate of ionization, 26:47122
- MAH, Seck Qui, Turbulent heating of the ions in a toroidal discharge, 26:47112
- MAHANTA, M. N., Approximate solution for the static, spherically symmetric metric due to a point charged mass in Brans--Dicke theory, 26:46750
- MAHDAVI, Mohamad Ali, Structure of a large-amplitude stationary hydromagnetic disturbance in a collisionless plasma, 26:47196
- MAHER, J. V., (¹⁶O,¹²C) studies in the 2s--1d shell, 26:46913
- MAHESHWARI, R. C., Isotope effect on Franck--Condon factors and r-centroids for the new (¹E/_{sub u}/⁺--¹E/_{sub g}/⁺) system of isotopic C₂ molecule, 26:45293
- MAIANI, L., Dreil--Hearn sum rule and the lepton magnetic moment in the Weinberg model of weak and electromagnetic interactions, 26:46553
- MATANI, Luciano, Criteria for selecting the location of nuclear plants, 26:47574
- MAILLARD, M. J., Simple and rapid procedure for strontium determination in rat plasma using neutron activation, 26:45202
- MAISTRENKO, A. N., Temperature dependence of cross section

of cold- and thermal-neutron scattering by O_2 and D_2 , 26:46887

MATII, H. S., Kinetics and burst phenomenon in ZrO_2 transformation, 26:45947

MAJUMDAR, Chanchal K., Solution of Faddeev equations for a one-dimensional system, 26:46782

MAKAI, E., Remarks on the three body problem in quantum mechanics, 26:47296

MAKARENKO, A. I., Hard gamma radiation from ^{238}Pu , 26:47032

MAKAROV, L. L., Chemical shifts in $L_{\alpha 1}$ x-ray line of some U compounds, 26:45377

MAKAROVICH, K. I., Accumulation of ^{90}Sr in wheat on sod-podzolic soils, 26:45550

MAKI, Kazumi, Inelastic neutron scattering from vortex lattice in Type-II superconductors. II. The London model, 26:46820

MAKI, Ziro, Quartet scheme of hadrons in chiral $U(4) \times U(4)$, 26:46687

MALES, R. H., Environmental benefit-cost analysis for nuclear power generation, 26:47386

MALES, Rene H., Are licensing delays really necessary, 26:47571

MALHERBE, J., Study of the mechanical and thermal behavior of a high temperature reactor core, 26:47304

MAL'KO, P. I., Electrical and thermal conductivities of Re, Os, and Ir and some of their alloys with Th and La in wide temperature range, 26:47248

MAL'KOV, V. V., Intranuclear cascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)

MALMIN, R. E., Evidence for shape or quasimolecular resonances in the $^{16}O + ^{16}O$ interaction, 26:46884

MAL'TSEV, V. M., Feynman-Wilson-gas model as consequence of jet approach to inelastic hadron processes, 26:46586 (JINR-P2-6501)

MALUCCI, R. D., Inclusive reaction $\mu + \text{nucleon} \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV/c, 26:46452

MAMIDZHANYAN, E. A., Angular distribution and energy spectrum of cosmic-ray single neutrons, 26:46153

MANCINI, F., Time-dependent phenomena in weakly-coupled superconductors, 26:46827

MANDI, Erika, Iron metabolism in mice exposed to mixed radiations from a reactor, 26:45840

MANDREK, K., Optical determination of the quadrupole moment of ^{138}La , 26:46950

MANN, A., Derivation of the critical temperature for the superfluid phase of hadronic matter, 26:46596

MANN, L. A., Fluid irradiating apparatus using gamma rays and helical passageways, 26:45719

MANN, Leland A., Irradiator apparatus with a fluid flowpath determined by communicating core and casting baffles, 26:45456

MANDWITZ, Bernard, Isotope ratio tracer method: applications in atmospheric sulfur pollution studies, 26:45568

MANSON, Steven Trent, Inelastic collisions of fast charged particles with atoms: ionization of the aluminum L shell, 26:46393

MANSOURI, Freydoon, Space-time description of hadrons, 26:46603

MARC, J.-C., Effects of thyroidian hormones and γ irradiation on adenylyclase of plasma membranes of rat liver, 26:45831

MARCHAL, P., Utilization of alumina radiothermoluminescence in zone dosimetry, 26:45628

MARCHESIN, Dan, Time-dependent and time-independent potential scattering for asymptotically Coulomb potentials, 26:46780

MARCHIONNI, R., Effect of taurine as a protector of human marrow cells against radiation, 26:45734

MARCILHACY, G., Are vector potentials measurable quantities in electromagnetic theory, 26:46542

MARCINKOWSKI, A., Energy of levels of ^{55}Sc up to 3 MeV from $^{44}Ca(p,\gamma)^{55}Sc$ reaction, 26:46920

MARCUSE, W., Continuous sampling plan for the taking of physical inventories, 26:46092 (BNL-16930)

MARDYKIN, I. P., Thermal properties of liquid cerium, 26:45998

MAREEL, Marc M., Phenol red uptake by irradiated and nonirradiated chick mesonephros in vitro, 26:45832

MARGOLIS, B., Reggeon contributions in a two-fire-ball model, 26:46663

MARINKOV, L., $2^{+} \rightarrow 2^{+}$ transition of 879 keV in ^{160}Dy investigated by electron-gamma directional correlation, 26:46991

MARIS, Georgeta, Active longitudes of sunspots and solar flares in the 20th cycle, 26:46222

Radio and x-ray emissions associated with chromospheric flares of November 15 and 16, 1970, 26:46223

MARIWALLA, K. H., Conformal invariance of zero mass Klein-Gordon-type equation, 26:46731

MARKER, T. F., Large scale characterization tests of gelled slurry blasting agents, 26:45484 (SC-DC-722296)

MARKOVSKII, L. Ya., Borocarbides of Tm, Lu, Tb, Yb, and some of their properties, 26:45964

MARKS, J. E., Film technique for determining the absorbed dose at exit from the patient during radiation exposure, 26:45631

MARKYTAN, M., K $^{-}$ -p interactions at 10 GeV/c, 26:46480

MARLOW, K. W., Production of carrier-free ^{197m}Hg with a cyclotron, 26:45395

MARNELIUS, R., Nucleon form factors in field theory on homogeneous space of the Poincare groups, 26:46554

39 AUTHORS

MATVEEV

MARNELIUS, Robert, Nucleon form factors in field theory on homogeneous space of Poincare group, 26:46514 (CTH-ITP-71-30)

MARONI, V. A., Containment of fission product iodine in the reprocessing of LMFBR fuels by pyrochemical reactions, 26:45421

MARRI, Aluisio, Installation of radioactive sources in the No. 4 blast furnace at Cia. Ferro Brasileiro for studying lining wear, 26:45720

MARTIN, D., Diffusion deposition of small-sized particles in the bronchial tree, 26:45921

MARTIN, D. L., Mechanism of sintering in cobalt-rare-earth permanent-magnet alloys, 26:45976

MARTIN, M. J., Log-f tables for beta decay, 26:46970

MARTIN, P. J., Critical currents in Nb_3Al , 26:46829

MARTIN, P. N., Method of determining mean energy of secondary electrons in electron equilibrium regions using theoretical x-ray spectra (10 to 20 MV) and ionization measurements in different media, 26:47213

MARTINIC, Nicolas J., Solar modulation of galactic particles, 26:46152

MARTINS, J. B., Some remarks on the transparency of complex nuclei to photoproduced charged pions at intermediate energies, 26:46889

MARTYNCHUK, E. L., Solubility of Ce, Pr, and Nd in Cr containing O, 26:46034

MASHKAROVA, V. T., Electron transfer during zone melting of yttrium, 26:45975

MASINI, G., Threshold of parallel channel flow instability, 26:45459 (CISE-R-275)

MASKAWA, Toshihide, Quartet scheme of hadrons in chiral $U(4) \times U(4)$, 26:46687

MASNOU, J. L., Investigation of low- q^2 inelastic neutrino events, 26:46450

MASON, W. N., Test of time reversal invariance through detailed balance of the reaction $^{12}C(\alpha,\gamma)^{16}O$, 26:46881

MASTERSON, Thomas G., Differential cross sections for small angle n-p scattering at 24 MeV, 26:46872

MATEI, V., Influence of neutron irradiation on seed germination and plant growth of tomatoes, 26:45773

MATEICIUC, V. A., Neutron spectrometry of Sb nucleus, 26:46946

MATEVOSYAN, E. M., Compensation of the drift of charged-particle tracks in wide-gap spark chambers, 26:45652

MATHEWS, P. M., Gauge problem in quantum field theory, 26:46720

Poincare and TCP invariance in the determination of wave equation for particles of arbitrary spin, 26:46714

MATHUR, A., Ring current effect on magnetospheric electron density profiles derived from plasmopause whistlers, 26:46284

MATHUR, K. C., Excitation of hydrogen in collisions with alkali-metal atoms, 26:46403

MATHUR, P. K., Water chemistry studies VII: general review of water conditions and water treatment in nuclear reactors, 26:45310 (BARC-595)

MATSEN, R. P., Determination of ratios of effective cross sections from measured burnup data for Yankee Rowe, 26:47595

MATSIBORKO, N. G., Non-linear theory of instability of a mono-energetic electron beam in plasma, 26:47151

MATSUDA, Tatsuo, Synthesis of O-containing halogenated compounds. II, 26:45335

MATSUDAIRA, Masamichi, Quantitative evaluation of radioisotope distribution in vivo by isosensitive scanner plus 4096-word multi-channel analyzer coupling, 26:45601

MATSUMOTO, H., Attempt at generalizing dual-resonance models, 26:46593

MATSUOKA, Kazuo, (N $\rightarrow$ Z) dependence of (n, γ) cross sections, 26:47085

MATSUOKA, Takeo, Urbaryon rearrangement and inclusive phenomenology, 26:46606

MATSUSHIMA, Satoshi, Continuum near the H and K lines of Ca $^{+}$ in the solar spectrum, 26:46218

MATSUSHITA, Rokuji, Direct measurement of γ -ray energies and half-life of ^{134}Cs , 26:46971

MATSUZAKI, Kei, Radiation-induced graft copolymerization of mixtures of styrene and n-butyl acrylate onto cellulose and cellulose triacetate, 26:45320

MATTHAEY, H., Proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46457 (KFK-1509)

Simple theory of proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46650 (KFK-1547)

MATTHIAE, G., Elastic scattering and isobar production in proton-proton collisions, 26:46464

MATTSOON, Ronny, Temperature dependence of the isotope effect for electromigration of rubidium ions in molten rubidium nitrate, 26:45294

MATUMOTO, Ken-iti, Size of hadronic constituents and applicability of scaling law, 26:46608

MATUURA, Nobuo, Theoretical models of ionospheric storms, 26:46300

MATVEENKO, I. I., Magnetic susceptibility of cubic niobium and tantalum carbides in the low temperature range, 26:47266

MATVEEV, V. A., Asymptotic estimates of quasipotential parameters, 26:46703 (JINR-P2-6482)

Operator method of solving the quasipotential equations and a conception of straight-line paths at high

- energies, 26:46700 (JINR-P2-6437)
- MATVIISHINA, N. N., Temperature dependence of cross section of cold- and thermal-neutron scattering by O_2 and D_2 , 26:46887
- MATYUSHENKO, N. N., Structure of binary silicides, 26:45965
- MAUSHART, R., Radiation protection measurement, 26:45808
- MAXIE, E. C., Effect of gamma irradiation on Fuerte avocado fruits, 26:45724
- MAY, G. G., Removal of radioactive iodine from ventilation exhaust streams at the AEC Research Establishment, 26:45502
- MAY, R., Radiation-induced formation of excited states of aromatic hydrocarbons in cyclohexane. IV, 26:45330
- MAYOUX, Christian, Comparative effects of ion bombardment and partial pressures on sheet polyethylene, 26:46081
- MAZARAKIS, Michael Geras, Experimental investigation of the $^{12}C+^{12}C$ reaction at low energies, 26:46898
- MAZEY, D. J., Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)
- MAZIERE, B., Simple and rapid procedure for strontium determination in rat plasma using neutron activation, 26:45202
- MCALPINE, I., Comparison of the radiolysis of liquid bromotrifluoromethane with the radiolysis of liquid trifluoroiodomethane, 26:45322
- MCCALLEY, Roderick Canfield, I. Radiation damage in malonic acid: an acyl radical. II. Deuteron distant ENDOR studies, 26:45343
- MCCONKEY, J. W., Ionization of argon by electron impact, 26:46374
- MCCORKLE, D. L., Low-energy electron attachment to molecules at high densities, 26:46364 (ORNL-TM-3839)
- MCCORMICK, Bruce H., Part 1: Annual technical progress report, October 1, 1971--June 30, 1972, 26:47610 (COO-2118-34)
- MCCRAY, Richard, Formation of clouds in a cooling interstellar medium, 26:46245
- MCCREA, W. H., Relativistic beaming, 26:46201
- MCCURRY, D. C., Radiological surveys of Pearl Harbor, Hawaii, and environs, 1966--1968, 26:45534
- MCDIARMID, I. B., Anisotropy of high-latitude electron fluxes during substorms and structure of the magnetotail, 26:46306
- MCDONALD, J. M., Magnetic moment of the 2761 keV 8^+ state in ^{92}Mo , 26:46959
- MCDONALD, R. C., Contrast effects at grain boundary ledges in 304 austenitic stainless steel, 26:47630 (UCLA-34-P-177-11)
- MCSELFRESH, P. M., Comparison of theoretical and experimental limits of detection in atomic absorption spectrometry using air-acetylene and nitrous oxide-acetylene flames, 26:45228
- MCGAHAN, D. J., Radiation polymerization of impregnated porous material, 26:45718
- MCGOWAN, F. K., Coulomb excitation of $^{117} 119Sn$, 26:46961
- MCGUIRE, Gary Elder, Photoelectron chemical shifts in inorganic compounds, 26:46340 (ORNL-TM-3820)
- MCKNIGHT, Allan, Atomic safeguards: a study in international verification, 26:46096
- MCNAUGHT, D. C., Influences of thermal effluents upon aquatic production in Lake Ontario, 26:45578
- MCPHERSON, D. A., Observation of very-low-frequency whistler-mode waves in the region of the radiation-belt slot, 26:46312
- MCRAE, Thomas G., Observations of precursor ionization in electromagnetic T tubes, 26:47131
- MEDIONI, R., Utilization of alumina radiothermoluminescence in zone dosimetry, 26:45628
- MEDITSCH, J. O., Determination of silica in silicates containing P, Ti, and Zr, 26:45255
- MEDVEDEV, A. I., Isobaric nuclei with mass number $A = 72$, 26:46939
- MEDVEY, Wojciech, Distribution of doses in the lesser pelvis in the treatment of carcinoma of the cervix uteri, 26:45784 (IT-54002/4, pp 456-61)
- MEESSEN, A., Finite difference Schroedinger equation, 26:47293
- MEGAW, W. J., Monitor for estimating tritium in urine, 26:45585 (AERE-HP/M-74)
- MEGGS, W. J., Reggeon contributions in a two-fire-ball model, 26:46663
- MEGLEN, Robert Rudolf, Studies of continuous dielectric models for solvated electrons, 26:45344
- MEHTANI, Usha, Partial wave analysis of $\pi^+p \rightarrow \pi^0\Delta^+(1236)$ in the center of mass energy region 1820 to 2090 MeV, 26:46495
- MEININGHAUS, H., WAK: specifications and finishing control for equipment, 26:46105 (ORNL-tr-2371)
- MELIK-BARKHUDAROV, T. K., Kinetic equation for superconducting alloys located in weak static fields, 26:46825
- MELKONIAN, E., Fission isomeric state in ^{236}U , 26:47018 (INDC-7-U, pp 61-4)
- Mass distributions in fission and some related phenomena, 26:47017 (INDC-7-U, pp 31-59)
- Total neutron cross section measurements, 26:46967
- MELDE, Thor B., Electron spin resonance studies and INDO MO-calculations of electron irradiated peptides. A single crystal of N-acetylglycine, 26:45349
- MELSKENS, J., Relative level populations in foil excited He and Li, 26:46337
- MELVANIN, F. W., Zirconium tetrafluoride process, 26:45274
- MENKE, Michael Matthew, Study of non-strange boson resonances produced by quasi-monochromatic photon beams of energies between 4.3 and 7.5 GeV in a hydrogen bubble chamber, 26:46446
- MENTKOVSKII, Yu. L., Free Riemann--Green's function of inverse problem of scattering theory, 26:46785
- Inverse problem of scattering theory and Riemann's method, 26:46788
- MENTZEN, Bernard, Determination of the structure of monohydrated uranyl diformate using x-ray diffraction, 26:45368
- MERCEREAU, J. E., Time-dependent superconductivity and quantum dissipation, 26:46821
- MERCIER, C., Generalized nonlinear expression for plasma displacement in Harmonica 2 and Tokamak, 26:47115
- MERIN, Robert G., Fluorine metabolite concentrations and renal clearances following methoxyflurane anesthesia, 26:47645 (UR-3490-107)
- MERKLE, J. G., Analytical applications of the J integral, 26:45990 (CONF-720806-2)
- MERONEY, Robert N., Gaseous plume diffusion about isolated structures of simple geometry, 26:45514
- MERRIAM, J. R., Clonal parameters of tergite development in Drosophila, 26:45881
- MERRITT, M. L., Cannikin: effects on archaeological sites, 26:45485 (SC-RR-720359)
- MERZYN, G., Hyperfine structure of ground state of ^{165}Ho , 26:46989
- MESSNER, Robert, Efficiency studies of Pilot Y, Pilot F, and NE 110 scintillator, 26:45588 (COO-2114-2)
- MESTELAN-NONNENMACHER, Genevieve, Hydrogen--deuterium exchange kinetics of cytochrome C between 20 and 380°C, 26:45305
- METAG, V., Missing low-lying states in ^{39}K , 26:46928
- METCALF, W. G., Shallow-water ^{90}Sr anomaly about the Antilles Arc, 1970, 26:45533
- METZ, S. A., Production of vacancy condensates by laser bombardment, 26:46059 (CONF-710601-, pp 613-29)
- MEYER, J., Charge states of ^{208}Pb and ^{208}Tl formed by α decay of ^{212}Po and ^{212}Bi , 26:45359
- MEYER, Joachim, 12-month wave in geomagnetic activity, 26:46316
- MEYER, V. D., Measurements of the absolute radiative efficiency of cathodoluminescent phosphors, 26:45591
- MEYER, Walter Thomas, PI-zero photoproduction from complex nuclei, 26:46447
- MEYER-HAMME, K., Role of transplanted slowly proliferating bone marrow cells for regeneration of lethally x-irradiated rat bone marrow, 26:45837
- MEYERHOF, W. E., New results on $(\pi^+, 2p)$ reactions in light nuclei, 26:46870
- Recent experiments on the four and five nucleon systems, 26:47089
- MEZIN, M., Uranium isotope separation studies in France, 26:45691
- MICHAEL, C., Line reversed reactions, 26:46629
- MICHAELIDES, Paraskevas, Acneiform changes in areas of cobalt irradiation, 26:45793
- MICHAELSON, S. M., Ionizing radiation-induced thyroid carcinogenesis in the dog, 26:45819 (UR-3490-63)
- MICHALSKA-KASZCZYNSKA, Maria, Effects of ionizing radiation on the structure and metabolism of mitochondria in rat liver cells, 26:45838
- Lysosomal system of liver cells in rats following a radiation dose of 1000 r to the whole body, 26:45839
- MICHAUD, Georges, Stellar reaction rates for the $^{12}C + ^{12}C$, $^{12}C + ^{16}O$, and $^{16}O + ^{16}O$ reactions, 26:46179
- MICHEL, B., Inelastic interaction induced by high-energy muons (6 GeV, 12 GeV) at low momentum-transfer in nuclear emulsion, 26:46451
- MICHEL, R., Effects of thyroidal hormones and γ irradiation on adenylylase of plasma membranes of rat liver, 26:45831
- MICU, M., Bounds for K /sub 13/ decay parameters, 26:46575
- MIDZUNO, Yukio, Nonlocal drift waves in a cylindrical plasma, 26:47181
- MIERZEWSKA, H., Passage of ^{90}Sr from bovine bones to received food products, 26:45549
- MIETTINEN, H. I., Regge phenomenology of inclusive spectra in the central region, 26:46659
- MIFUNE, Tsuyoshi, Use of irradiation in plastic wastes disposal, 26:46090
- MIGDAL, A. B., Comparison of various approaches to nuclear theory, 26:47093
- MIGNANI, R., Superluminal frames and group of generalized Lorentz transformations in four dimensions, 26:46734
- MIHALAS, Dimitri, Observations of the He II $\lambda 10124$ line in O and Of stars, 26:46180
- MIKAMO, S., Possible bounds on muon pair and W^+ production in super-high-energy hadron collisions, 26:46598
- MIKHAILOVSKII, A. B., Micro-instabilities of high pressure plasma associated with dependence of the ion viscosity on the derivatives of the longitudinal thermal flux, 26:47165
- MIKKELSON, R. C., Channeling of protons in thin $BaTiO_3$ crystals at temperatures above and below the ferroelectric curie point, 26:47234
- MIKUL'SKAYA, G. Yu., Study of zirconium evaporation from open surface, 26:46011
- MILANOVIC, J., Effective g /sub a/ factor for odd proton

MILENIN, V. M., Electron energy distribution and striations in the Langmuir-paradox region, 26:47133

MILLER, D. H., Final states with three charged particles and a visible Λ from K-d interactions at 4.5 GeV/c, 26:46474 (CDD-1428-306)

MILLER, E. C., Various levels of DNA repair synthesis in xeroderma pigmentosum cells exposed to carcinogens N-hydroxy and N-acetoxy-2-acetyl-aminofluorene, 26:45742

MILLER, J. A., Various levels of DNA repair synthesis in xeroderma pigmentosum cells exposed to carcinogens N-hydroxy and N-acetoxy-2-acetyl-aminofluorene, 26:45742

MILLER, R. W., (¹⁶O, ¹²C) studies in the 2s--1d shell, 26:46913

MILLER, William E., Mixer-settler apparatus, 26:45451

MILLION, B., Comments on diffusion of cobalt in plutonium by M. J. Daph, M. P. Dariel, and A. Rosen in Scripta Met., 5: 191-4(1971), 26:46024

MILLS, Alfred Leonard, Processing of irradiated nuclear fuels, 26:45420

MILLS, James Ignatius Jr., Electrical shock-tube precursor, 26:47197

MIL'MAN, Y. V., Peculiarities in the structural sensitivity and mechanical characteristics of niobium, 26:46019

MILNE, D. K., Polarized radio emission from five supernova remnants, 26:46216

MILNER, W. T., Coulomb excitation of ¹¹⁷ ¹¹⁹Sn, 26:46961

MILOSZ, Jacek, Hydrogen generators for neutron tubes, 26:45942

MILROY, William C., Pathophysiologic effects of microwave radiation with special reference to the thyroid--pituitary axis, 26:47648 (UR-3490-137)

MINAKATA, H., Inclusive reactions and the asymptotic behavior of proton--proton collisions, 26:46657

MINAMI, S., s-channel helicity amplitudes without pomeron for elastic K⁺--p and π --p charge exchange scattering, 26:46637

MINARDI, E., Statistical thermodynamics of collisionless plasma. II, 26:47164

Statistical thermodynamics of collisionless plasma. I, 26:47163

MINELLI, T. A., Solution of the standing-wave integral equation, 26:46786

MINKOWSKI, P., Dangerous scaling limit for virtual Compton scattering off the forward direction, 26:46530

MINKOWSKI, R., Bright nebulae near concentrations of high-velocity gas, 26:46250

MINYAEV, D. A., Compensated current-driven magnetic channel in a 240-cm isochronous cyclotron, 26:46128

Magnetic shield for a 240-cm isochronous cyclotron, 26:46129

MIRANSKII, V. A., Duality, scaling, and fixed poles in virtual Compton scattering, 26:46559

MIRZABABAEV, R. M., Moessbauer diffractometer having high angular resolution, 26:45683

MISGUICH, J. H., Generalized nonlinear expression for plasma displacement in Harmonica 2 and Tokamak, 26:47115

MISHRA, R., Room temperature internal friction of neutron-irradiated aluminum, 26:46073

MISRA, K. D., Reflection of microwave through laboratory plasma, 26:47118

MISRA, S., Thermodynamic properties of the phase In₅Bi₃, 26:46026

MITRA, A. N., Model for the mixing among the 70 baryon states based on exchange degeneracy, 26:46685

MITTER, P. K., Analytic renormalization of the exponential interaction: the three-point time-ordered product with minimum light cone singularity, 26:46712

MITTLEMAN, M. H., Optimized pseudostates, intermediate-energy scattering, and collective modes in atoms, 26:46389

MIYABARA, Shinichi, Statistical analysis of radioinduced abnormal development, 26:45867

MIZERAS, A. V., Some observations on radiation safety in Virginia dental offices, 26:45500

MOCHESNIKOV, N. I., Control system for beam formation after single deflection at exit of 300 MeV electron linear accelerator, 26:46131

Decrease of the off-duty factor in the operation of a linear accelerator, 26:46145

MODZOLEVSKII, V. I., Generation of shock waves by exploding current sheath, 26:47186

MOGILEVSKII, O. A., Choice of a form factor in nonlocal quantum electrodynamics, 26:46698 (ITF-71-128-P)

Choice of form factors in non-local quantum electrodynamics, 26:46711

MOGIL'NER, A. I., Use of solid track detectors in reactor experiments, 26:45610

MOHAN, N., Application of perturbation theory for the calculation of molecular constants due to small mass changes and on the determination of force constants from very heavy isotope substitution, 26:45299

MOISENKO, B. A., Protection from neutron emission of ²³⁸Pu dioxide, 26:46852

MOISIU, M., Glucose-6-phosphate phosphohydrolase activity in erythrocytes after x-irradiation, 26:45848

MOLDovanu, A., Statistical treatment of relationship between solar flare-producing regions and SC + SI events, 26:46332

MOLNAR, I., ¹⁷⁵Hf states excited during decay of ¹⁷⁵Ta, 26:46979

MONAGHAN, William, Method of effective potentials:

scattering of positrons and electrons by light atoms, 26:46400

MONDINO, P. L., Experiments on a superposed fields theta pinch, 26:47171

MONNIER, Bernard, Effect of nuclear radiations on rubber joints, 26:46087

MONTALDI, E., Analytic renormalization via continuous space dimension, 26:46729

MONTANET, L., Properties of meson resonances as observed in anti π --p annihilations at rest, 26:46465

MONTGOMERY, C. D., Cells for nondestructive examinations of irradiated fuel pins, 26:46104 (LA-DC-72-556)

MONTRET, J. C., Inelastic interaction induced by high-energy muons (6 GeV, 12 GeV) at low momentum-transfer in nuclear emulsion, 26:46451

MOORE, C. Bradley, Isotopic separation by photopredissociation, 26:45307

MOORE, M. S., Measurements on fissile nuclides, 26:47036

MOORE, Ronald Lee, Structure and heating of the chromosphere--corona transition region, 26:46228

MORAN, T. I., Direct measurement of the nuclear magnetic dipole moment of ¹⁶⁵Ho, 26:46990

MORCOS, Ikram, Surface tension of stress-annealed pyrolytic graphite, 26:46091

MORETTI, E., Metabolism of monomeric and polymeric Pu in the rabbit. Comparison with the mouse, 26:45917

MORFILL, G. E., Entry of high-energy solar protons into the distant geomagnetic tail, 26:46304

MORGAN, Carlisle Lee Jr., Investigation of the pressure shift of the ground state hyperfine lines of H, D, and T, 26:46347

MORI, Ryoichi, Vascular alteration in irradiated bone marrow of rabbit mandible, 26:45835

MORICI, A., DECAT: Computer code for analysis of long term pressure and temperature transient after loss of coolant accident in G.E. type pressure suppression containment, 26:47318 (RT/PROT-72-4)

MORINAGA, H., Neutron deficient even-even isotopes of Kr, Se, Ge, and pseudo SU₃ coupling, 26:46931

MORIoka, Akira, Daytime irregular geomagnetic pulsation, 26:46317

MORITA, K., Scaling and finite electromagnetic self-mass of hadrons, 26:46558

MORIUCHI, Yasuyuki, Present status of personnel dosimeters, 26:45624

MORRIS, T., Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. II, 26:45328

MORRISON, Clyde A., Nuclear charged self-sustaining laser, 26:45677

MORRISON, D. L., Environmental benefit-cost analysis for nuclear power generation, 26:47386

MORRISON, G. C., ⁵⁸Ni(α, α')⁵⁸Ni and the nature of high-lying states in ⁵⁸Ni, 26:46927

MORROW, Paul E., Experimental inhalation of metallic silver, 26:45915 (UR-3490-115)

MORROW, Roy W., Determining boron in glass by flame emission spectroscopy, 26:45254

MORROW, T., Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. III, 26:45329

Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. I, 26:45331

Radiation-induced formation of excited states of aromatic hydrocarbons in cyclohexane. IV, 26:45330

MOSEL, U., Single-particle effects on fission probabilities for the lighter elements, 26:47070

MOSELEY, John T., De-excitation of fast He, Ne, and Ar metastable atoms in various gases, 26:46394

MOSS, D. L., Angular momentum loss from contact binary systems, 26:46167

MOSSOTTI, Victor G., Emission spectroscopy including dc arc, spark, and other methods, 26:45247

MOTZ, H., Note on plasma acceleration caused by plasma turbulence, 26:47134

MOUSSA, A., On-line measurements of rare-gas fission yields in 14 MeV neutron fission, 26:47030

MUBARAK, M. A., Radiation safety around hot facilities at PINSTECH, 26:45499

MUEHLSCHLEGEL, B., Thermodynamic properties of small superconducting particles, 26:46813

MUELER, H. J. W., Regge poles for large coupling constants in the Wick--Cutkosky model, 26:46725

MUELLER, A., Application of perturbation theory for the calculation of molecular constants due to small mass changes and on the determination of force constants from very heavy isotope substitution, 26:45299

MUELLER, Erwin W., Imaging process in field ion microscopy, 26:46038

MUELLER, M. H., Magnetic phase diagram of the UAs--US system, 26:47261

MUENZEL, H., Mean excitation energies of the daughter nuclei on β --decay, 26:46875 (KFK-EXT-10/72-1)

Systematics of excitation functions for fast neutron induced (n,p)- and (n, α)-reactions, 26:47066

MUETHER, H., Orbital rearrangement in light nuclei, 26:46886

MUIJE, Pieter, Kinetics of solid vanadium--nitrogen reaction at high temperatures, 26:45969

MUIR, J., Measurement of slope parameter in decay K/_{sub} L/⁰

- *w-w0, 26:46494
- MUKHIN, A. I., New method of determination of physical-chemical properties of matter, 26:47618 (JINR-P14-6407)
- MUKOYAMA, T., Possible evidence for a contribution from direct collisions in K-shell internal ionization accompanying β decay, 26:46962
- MUKOYAMA, Tomoyuki, Polarographic determination of Sn in Cu-, Al-, and Zr-base alloys, 26:45256
- MULLER, H., Process for recovering by-products from uranium dioxide production, 26:45371
- MULLER, J., Identification of the superconducting phases in the molybdenum-platinum system, 26:46828
- MULLIGAN, David Dwight, Radiomodifying chemicals for bacteria and mice, 26:45859
- MUMINOV, A. I., Magnetic properties of ^{153}Eu , ^{155}Eu , and ^{157}Dy , 26:46981
- MUNIZ, R. P. A., ESR studies of electron irradiated $\text{K}_2\text{Ir/sup III/}(\text{CN})_6$ in KCl single crystals. I. Ligand hyperfine structure in Ir/sup II/(\text{CN})₆ species, 26:45317
- MUNSON, B. R., Protein synthesis in normal, x-irradiated, and friend virus infected mouse spleen nuclei, 26:45849
- MUNTENBRUCH, H., Power production costs and possible applications of short-period combustion MHD generators, 26:47170 (IPP-IV-32)
- MURADYAN, E. G., Preamplifier for Cherenkov spectrometer, 26:45608
- MURAI, Masafumi, Recessive genes in sugi, *Cryptomeria japonica*, 26:45771
- MURAKAMI, Yukio, Production of carrier-free ^{54}Mn , 26:45394
- MURDIN, Paul, Photoelectric measurements of the 6284 Å diffuse interstellar line, 26:46168
- MURGALESCU, Sanda, Determination of inorganic impurities in vinyl chloride by activation analysis, 26:45204
- MURINO, Pasquale, Spectrographic measurement of the temperature of a plasma analyzed by statistical methods, 26:47129
- MURPHY, J., Paramagnetic hole centers in CeO_2 , 26:47233
- MURTY, P. Sreerama, Spectrochemical analysis of high purity cadmium, 26:45220 (BARC-593)
- MURTY, R., Model studies of density fluctuations of small and large ions in the atmosphere, 26:46288
- MURZIN, A. V., Excitation of input states during thermal neutron capture by ^{60}Zn , ^{70}Ge , and ^{80}Se , 26:46937
- MUSAEV, R. M., Excitation cross section of first 2+ levels of Ni and Zn nuclei, 26:46936
- MUSCETTOLA, M., PDE: digital program for calculation of thermal-hydrodynamic condition in circuit of two phase mixture water and steam, 26:45458 (CISE-R-245)
- Programs for thermohydraulic behavior of primary circuit of reactor CIRENE, 26:47309 (CISE-R-257)
- MYTISOV, F. F., Dielectric strength of accelerator tubes, 26:46125
- N**
- NABEDRYK-VIALA, Eliane, Hydrogen-deuterium exchange kinetics of cytochrome C between 20 and 38°C, 26:45305
- NADKARNI, D. M., Prompt radiations emitted in low energy fission, 26:47019 (INDC-7-U, pp 89-112)
- NADZHAKOV, E., New isotopes of iridium, 26:46977
- New isotopes of tungsten, 26:47001
- NAGAI, Hideo, Paper chromatography of rare earth elements with developer containing ethylene glycol monoethers, hydrophobic solvents, and oxine, 26:45275
- NAGAI, Osamu, Effects of impurities on gamma-irradiated silicon crystal examined by photovoltaic effect of p-n junction diode, 26:45671
- NAGY, A. F., Satellite and ground-based observations of a red arc, 26:46275
- NAITO, Shuichi, Identification of reaction intermediates by microwave spectroscopy. The catalytic reactions between propylene and deuterium over zinc oxide, 26:45303
- NAKAGIRI, Masaru, Effects of x-ray irradiation on MAS diodes, 26:45670
- NAKAGOMI, Yoshihito, Thin-layer chromatography of metal ions on microcrystalline cellulose impregnated with potassium chloride, 26:45278
- NAKAI, K., g-factor of the 12^+ isomeric state in ^{206}Pb , 26:47002
- NAKAI, Kiyoto, Paper chromatography of rare earth elements with developer containing ethylene glycol monoethers, hydrophobic solvents, and oxine, 26:45275
- NAKAJIMA, N., Effect of heat treatments on corrosion of 18/10 stainless steels with low C contents in nitric medium in presence of ferric iron, 26:45970
- NAKAMURA, Shigeo, Radiation-induced graft copolymerization of mixtures of styrene and n-butyl acrylate onto cellulose and cellulose triacetate, 26:45320
- NAKAMURA, W., Effects of partial body irradiation of mice on quantity of 5-hydroxyindol acetic acid in urine, the quantity of water consumed, and the volume of urine excreted, 26:45826
- NAKANISHI, Fumiko, Hepatic injuries following irradiation, 26:45809
- NAKKAGAWA, Hisao, Explanation of strong core in hadron reactions, 26:46607
- NAMIKI, Akira, Trapped electrons produced in ethanol glass at 40K, 26:45319
- NAMYSLOWSKI, J. M., Breakdown of simple cancellation in the operator multiple scattering series, 26:46773 (CTH-ITP-71-17)
- NARASIMHA RAJU, M. L., Effective atomic numbers for photoelectric process in various shells in alloys, 26:47206
- NARAY-SZABO, G., Decoupled Hartree-Fock methods. I. Calculations for atoms with orthogonalized orbitals, 26:46416
- NARLIKAR, J. V., Mass difference of the muon and the electron, 26:46259
- NASRALLAH, N. F., Collinear dispersion relations and rho- π and A_1 -rho- π couplings, 26:46612
- NASTOYASHCHII, A. F., Thermonuclear combustion waves in a bounded plasma, 26:47172
- NATHAN, R. D., Optical excitation and polarization in low-energy collisions of $\text{He}^+ + \text{H}_2$, 26:46396
- NATHAN, Richard David, Optical excitation from low-energy $\text{He}^+ + \text{H}_2$ and $\text{H}_2^+ + \text{He}$ collisions, 26:46371
- NAUENBERG, Uriel, Efficiency studies of Pilot Y, Pilot F, and NE 110 scintillator, 26:45588 (COO-2114-2)
- NAUGOL'NYI, N. N., Effect of the collective interaction of particles on the longitudinal dimensions of an electron bundle in an accumulator, 26:46546
- NAVARRO, Alain, Determination of the structure of monohydrated uranyl diformate using x-ray diffraction, 26:45368
- NAWAR, W. W., Recombination products from the radiolysis of tricaproin, 26:45321
- NAYLOR, Mark Dennis Thomson, Equilibrium structure of close binary stars, 26:46173
- NAZARETH, R. A. M. S., Some remarks on the transparency of complex nuclei to photoproduced charged pions at intermediate energies, 26:46889
- NAZARYAN, A. A., Detector for high energy particles based on the principle of using particle radiation in a laminated medium, 26:45648
- NEE, Soe-mie Foeng, Synchrotron radiation from a magnetically confined hot-electron plasma, 26:47144
- NEGELE, J. W., Progress in the theory of nuclear matter and its application to finite nuclei, 26:47096
- NELSON, C. A., Algebraic solution for the Kaellen-Pauli state vectors in the Vtheta sector by a congruence transformation, 26:46705
- NELSON, Donald M., Radionuclide concentrations near the Big Rock Point Nuclear Power Station, 26:45536
- NELSON, J. V., Three dimensional neutronics calculations for the Fast Test Reactor (FTR) and the FTR Engineering Mock Up Critical assembly (EMC), 26:47586 (HEDL-TME-72-42)
- NELSON, Neal S., Bone and soft-tissue neoplasms in cats exposed to radiostrontium, 26:45901
- NELSON, R. S., Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)
- NELSON, Victor A., Decline of ^{65}Zn in marine mussels following the shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)
- NELSON, Wayne, Fecundity and longevity of F_1 females of *Habrobracon* from sperm x-rayed with 3000 r, 26:45878
- NEMCHENKO, V. F., Electrical and thermal conductivities of Re, Os, and Ir and some of their alloys with Th and La in wide temperature range, 26:47248
- NEMETS, O. F., Angular distribution of neutrons from (p,n) reactions on ^{140}La , ^{137}Cl , ^{41}K , ^{53}Cr , and ^{59}Co nuclei, 26:46929
- NEMOTO, Fumiki, Alpha-cluster structure and weak coupling feature in light sd-shell nuclei, 26:47083
- NENOV, N., ^{175}Hf states excited during decay of ^{175}Ta , 26:46979
- NESTERENKO, V. V., Corrections to eikonal representation of amplitude in scalar field theory, 26:46699 (JINR-P2-6394)
- NETSEROV, B. V., γ -emission in radiative capture of intermediate neutrons, 26:46917 (BNL-tr-477)
- NETTESHEIM, P., Increased effective half-life of intratracheal administered $^{51}\text{Cr}_2\text{O}_3$ by respiratory infection, 26:45910 (CONF-720505-2)
- NETZEL, Daniel A., Pulse fourier transform nuclear magnetic resonance spectroscopy, 26:47277
- NEUGEBAUER, M., Detection of solar-wind electron plasma frequency fluctuations in an oblique nonlinear magnetohydrodynamic wave, 26:46329
- NEWBY, B. J., DTA-x-ray diffraction examination of chemical reprocessing wastes containing sodium and potassium nitrates, 26:45383 (ICP-1008)
- NEWELL, Reginald E., Correlation of D-region absorption and the 10 mb winds and temperature during the IQSY, 26:46264 (COO-2195-5)
- Planetary transport processes in the upper atmosphere. Annual progress report, 26:45503 (COO-2195-6)
- NEWMAN, Charles Edward Jr., Charged-particle transport in turbulent astrophysical plasmas, 26:46294
- NEWMAN, Leonard, Isotope ratio tracer method: applications in atmospheric sulfur pollution studies, 26:45568
- NEWTON, J. C., Anode temperatures and ion energy distributions during a vacuum arc discharge, 26:47625 (SC-DC-722267)
- NEY, E. P., Core of eta Carinae, 26:46160
- NGUYEN CONG KHAN, Angular distribution of fragments of spontaneously fissioning isomers, 26:47022 (JINR-P7-

- NGUYEN LONG-DEN, Systematic study of the decomposition of very small quantities of rare earth oxides by mass spectrometry, 26:45259
- NGUYEN VAN GIAI, Nucleon-nucleus potential in Hartree-Fock approximation with Skyrme's interaction, 26:47064
- NGUYEN VAN HIEU, Essentially nonlinear scalar fields and the space-time geometry, 26:46692
- NICHITIN, Fl., Bubble density determination in bubble chambers, 26:45649
- NICHKOV, I. F., Thermodynamic properties and phase composition of La-Zn alloys, 26:46001
- NICHOLAS, J. Blair, Phase and temperature dependence of positron annihilation in liquid crystals, 26:46363
- NICHOLSON, Patrick S., Phase analysis in zirconia systems, 26:45945
- NICKLE, H. H., Algebraic approach to the solution of quantum-mechanical problems involving the interaction of a charged particle with electromagnetic fields, 26:46678
- NIDAY, J. B., Computerized quantitative analysis by gamma-ray spectrometry. Volume I. Description of the GAMANAL program, 26:45265 (UCRL-51061(Vol.1))
- NIEWIADOMSKI, Michal, Statistical data on iodine uptake of the thyroid gland. II, 26:45906 (TT-54002/4, pp 462-70)
- NIFONG, Gordon D., Multi-element analysis of air pollution particulates by nondestructive neutron activation, 26:45207
- NIGAM, A. K., Gamma-gamma angular correlation of 171- to 427-keV cascade and the g-factor of the 462-keV, 5/2⁺ level of ¹²⁵Te, 26:46957
- Gyromagnetic ratio of 2083-keV, 4⁺ state of ¹⁴⁰Ce, 26:46958
- NIGMETOVA, R. Sh., Thermodynamic properties of solutions in system In-Sn-Hg, 26:46004
- NIKA, G. Gerald, Study of exchange reactions employing a complementary shock tube facility, 26:45306
- NIKIFOROV, V. I., Photoproduction of charged pions by nuclei at E₀ = 1160 MeV, 26:46441
- NIKITENKO, N. F., Effect of soil conditions and K on uptake of ⁸⁶Rb in corn plants, 26:45925
- NIKITIN, V. I., Distribution of a radioactive aerosol in a model chamber, 26:45444
- NIKOLAISHVILI, Sh. S., Gamma quanta distribution at large distances from the sources, 26:47215
- NIKOL'SKII, S. N., Realization of synchronous rotation of some rotors suspended in a magnetic field, 26:45441 (SRARI-P-120)
- NILSSON, B., Importance of the quadrupole pairing field in the J/sup w/ = 0⁺ vibrations of shape deformed nuclei, 26:47042 (COO-3001-12)
- NISHIKAWA, M., γ radiolysis of gaseous carbon dioxide at high densities, 26:45346
- NISHIMOTO, Y., Effects of partial body irradiation of mice on quantity of 5-hydroxyindol acetic acid in urine, the quantity of water consumed, and the volume of urine excreted, 26:45826
- NISHIMURA, Fumio, Energy and angular distributions of secondary electrons resulting from ionizing collisions of electrons with helium and krypton, 26:46367
- NISHIMURA, Kazuaki, (N--Z) dependence of (n, γ) cross sections, 26:47085
- NISHIMURA, Sanji, Solvent extraction of Eu by Versatic Acid 911, 26:45277
- NISHIO, Kazuo, Effects of ⁹⁰Sr and ¹³⁷Cs administered continuously upon mice. VIII, 26:45900
- NODA, Masato, Trapped electrons produced in ethanol glass at 40K, 26:45319
- NOGAMI, Y., Magnetic moments of ³He and ³H, 26:46867
- NOGUTI, T., Semi-in vitro repair of radiation-induced damage in transforming DNA of Bacillus subtilis, 26:45731
- NOLLEZ, G., Standard black body at 10,000°K (high pressure plasma arc), 26:47217
- NOLLEZ, Gerard, Preliminary results obtained from the study of an argon plasma continuum in the near ultraviolet and visible ranges at high pressures, 26:47132
- NOLTE, E., Neutron deficient even-even isotopes of Kr, Se, Ge, and pseudo SU₃ coupling, 26:46931
- NOREM, W. L., Catalog of LLL computer documentation reports, 26:47598 (UCID-30047)
- NORMAN, H. H., NERVA flight engine control system design, 26:47577
- NORRIS, D. I. R., Application of high-voltage electron microscopy to void studies, 26:46056 (CONF-710601-, pp 533-49)
- NORSEEV, Yu. V., α -decay of neutron-deficient isotopes of Ra, 26:47004
- NORSETH, Tor, Effect of a polythiol synthetic resin on the absorption and excretion of methylmercury in experimental animals, 26:47643 (UR-3490-80)
- NORTHAM, D., Momentum transfer and plasma formation above a surface with high-power CO₂ laser, 26:47155
- NORTHROP, John, Long-range detection and location of some shallow-depth explosions in the Northwest Pacific, 26:45489
- NORTON, R. B., Thermospheric molecular oxygen from solar extreme-ultraviolet occultation measurements, 26:46269
- NORWOOD, W. D., Cumulative dose from diagnostic irradiation, 26:45799
- NOSAL, V. N., Review and analysis of the latest data on measuring fundamental physical constants, 26:46439
- NOSHKIN, V. E., Concentrations and distributions of long-lived fallout radionuclides in open ocean sediments, 26:45531 (COO-3563-3)
- NOSHKIN, Victor E., Plutonium in North Atlantic Ocean organisms; ecological relationships, 26:45541 (NYO-2174-130)
- NOTEA, A., Investigation of freon fire-extinguishing systems with a nucleonic gage, 26:45712
- NOVEL, J. P., Bleomycin labeled with ⁵⁷Co, 26:45929
- NOVAK, J., Deformation of fuel element rods during irradiation, 26:46070 (UJV-2798-M)
- Role of statistical fluctuations in texture during the radiation growth in uranium, 26:46069 (UJV-2795-M)
- NOVAK, L., Effect of radioprotective preparations on the respiratory exchange of rats, 26:45823
- NOVATSKII, B. G., Elastic scattering of ⁶Li ions, 26:46883
- NOVICK, R., Fine structure of the helium negative ion, 26:46343
- NOZAKI, Fumio, Study on catalysis of uranium oxide and its mixed catalysts. II, 26:45374
- NOZAKI, Tadashi, Production of carrier-free ⁵⁴Mn, 26:45394
- NULMAN, R., General properties of correlation functions for inclusive processes, 26:46502
- NYBO, K., Energy levels in ²³¹Th and ²³³Th, 26:47029
- OAKES, M. E., Modified ordinary wave, 26:47192
- OBATUROV, G. M., Study of the characteristics of emergency track detectors, 26:45617 (FEI-296)
- OBERT, W., Multiple ionization of rare gas clusters by electron impact, 26:46386
- O'CARROLL, Michael L., Time-dependent and time-independent potential scattering for asymptotically Coulomb potentials, 26:46780
- OCERAIES, R., Harmonie reactor, 26:47590
- OCHS, Wilhelm, Covering law in quantal proposition systems, 26:47294
- O'CLAIR, C. R., Preparation of transparent Y₂O₃-doped ThO₂, 26:45948
- O'CONNOR, G. P., Field-ion microscope study of ion-implantation in Ir. II, 26:47240
- Field-ion microscope study on ion-implantation in Ir. I, 26:47239
- ODA, Nobuo, Energy and angular distributions of secondary electrons resulting from ionizing collisions of electrons with helium and krypton, 26:46367
- O'DELL, S. L., Physical associations between quasi-stellar objects and galaxies, 26:46258
- ODINTSOV, P. N., Gamma induced changes in spruce wood, 26:45774
- ODOM, C. H., Continuous or semicontinuous leacher for leaching soluble core material from sheared spent nuclear fuel tubes, 26:45397 (CONF-720903-9)
- O'DONOVAN, P. J., Duality-breaking effects in a model for meson-baryon scattering, 26:46642
- ODORICO, R., Experimental low-energy behavior of dips and Veneziano duality, 26:46635
- OGLOBLIN, A. A., Elastic scattering of ⁶Li ions, 26:46883
- OHATA, Takeo, Hepatic injuries following irradiation, 26:45809
- OHBA, Kihachiro, Recessive genes in sugi, Cryptomeria japonica, 26:45771
- OHLENSCHLAEGER, L., Radiation protection considered from the medical viewpoint, 26:45803
- OHLWEILER, O. A., Determination of silica in silicates containing P, Ti, and Zr, 26:45255
- OHMER, Melvin C., Voltages induced in superconductors in the absence of transport currents, 26:46835
- OHMURA, Y., Electrical properties of n-type Si layers doped with proton bombardment induced shallow donors, 26:47237
- OHMURA, Yamichi, Determination of the spatial distribution of deep centers from capacitance measurements of pn junctions, 26:47243
- OHNISHI, Tokuhiko, Dose-rate measurement by several Radocon probes. II, 26:45620
- OHTANI, Teruya, Variational principle for nonlinear Lagrangian in quantum mechanics, 26:47298
- OHWADA, K., Infrared absorption spectra of potassium uranium fluorides, 26:45230
- OHYABU, Nobuyoshi, Neutral-point discharge experiment, 26:47142
- OIKAWA, Masaru, Pulmonary radiation injury, 26:45863
- OKABAYASHI, Takao, Breakdown of symmetry in the quantum theory of the nonlinear realization of a group on its subgroup, 26:46676
- OKABE, Shigeru, Effect of temperature on various types of the ionization chamber, 26:45594
- OKADA, Hiroshi, Study on catalysis of uranium oxide and its mixed catalysts. II, 26:45374
- OKADA, S., Effects of actinomycin D on G₁ protein synthesis in L5178Y cells, 26:47650 (UR-3490-153)
- Protein synthesis during the cell cycle of L5178Y cells, 26:47651 (UR-3490-154)
- OKAGAWA, Seigo, Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)
- OKAMOTO, Jiro, Radiolyses of mixtures of ammonia and hydrocarbons. II, 26:45332

- OKAMURA, Hiroshi, Gauge transformation and gravitational potentials, 26:46766
- OKHLOPKOVA, V. A., Cross section of pion--nucleon interactions, 26:46478 (ORNL-tr-2544)
- OKROASHVILI, G. G., Estimation of the turbulent diffusion coefficient and vertical component of wind velocity according to natural radioactivity distribution, 26:45522
- OKUBO, Atsuo, Development of 50Hz 44 inch turbine blade, 26:45454
- OKUDA, Takayoshi, Impedance of a partially ionized cesium plasma, 26:47135
- OKUMOTO, Masaaki, Culture of murine leukemic cells induced by x-irradiation, 26:45855
- OKUN, L. B., Interpretation of experiments on searches for $K/\text{sub } S/ + 2\mu$ decay, 26:46572
- OKUNEWICK, James P., Suppression and recovery of colony forming units in the rat following sub-lethal x irradiation, 26:45834
- OL, A. I., Disturbance of the Earth's magnetic field and its variations in the eleven-year cycle, 26:46322
- OL'DEKOP, L. G., Low-energy-electron spectrometer having a multiplate analyzer, 26:45643
- OLDS, F. C., Fast breeder: schedule lengthens, cost escalates, 26:47316
- OLINESCU, R., Effect of x-ray irradiation on glutathione peroxidase, 26:45733
- OLIVE, Philippe, Preliminary data on tritium content of aquifer sheets in northwest and central (Algeria) Sahara; hydrogeological consequences, 26:45558
- OLIVER, G. D. Jr., Aspects of radiation control for clinical implants of ^{252}Cf , 26:45790
- OLKOWSKI, Z., Ribonucleic acid synthesis and ionizing radiation, 26:45868
- OL'SHANETSKII, V. E., Effect of micro-additives of rare earth metals on the grain boundary energy of nickel, 26:45999
- OLSON, Ronald E., Semiempirical calculations of the $\text{He}^+(2^3\text{S} \text{ and } 2^1\text{S}) + \text{Ar}$ ionization total cross sections, 26:46395
- O'MONGAIN, E., Detection of high-energy gamma rays from the Crab Nebula, 26:46214
- OMORI, Yoshiaki, Age-sex specific Achilles tendon reflex time in Japanese: a study of atomic bomb survivors, Hiroshima, 26:45795
- O'NEIL, R. A., Mixing of two-proton bands with two-neutron bands in ^{17}HF , 26:46992
- ONISHCHENKO, I. N., Non-linear theory of instability of a mono-energetic electron beam in plasma, 26:47151
- ONISHI, Hiroshi, Separation and spectrophotometric determination of Ce and Mn by TTA extraction, 26:45226
- ONO, K., Reactor steels studies. Quarterly progress report, October--December 1971, 26:45994 (UCLA-34-P-177-10)
- ONO, Mitsuo, Electron spin resonance study of the far-ultraviolet photolysis of some inorganic and organic compounds, 26:45323
- ONORI, Sandro, Phase sensitive method for detecting superconductive transitions in the microwave region, 26:46840
- OOSTERBANN, R. A., In vitro repair of uv-inactivated transforming DNA from *Bacillus subtilis*, 26:45727
- ORLOV, B. N., Elastic and inelastic scattering of 6-MeV protons by even isotopes of Cr, 26:46922
- ORLOVA, K. B., Allowance for mass-spectrometer background and the reproducibility in the determination of isotope compositions for nitrogen and oxygen, 26:45241
- ORR, Clyde Jr., Aerosols produced by x-rays, 26:46082
- ORWIG, Larry Eugene, Observation of pulsed gamma rays with energies greater than 250 keV from the Crab Nebula pulsar NP 0532, 26:46199
- OSHEROFF, D. D., Spin-wave phenomena in liquid ^3He systems, 26:46802
- OSMAN, A., Repulsive core effects in (t,p) stripping reactions, 26:47078
- OSMER, Judith A., Crystallization of BeO , Be_2SiO_4 , and SiO_2 from Li_2MoO_4 -- MoO_3 , 26:45943
- OSMER, P. S., Optical observations of supernova in NGC 5253, 26:46187
- OSTASHENKO, V. F., Dynamics of control system incorporating 440 MW(e) reactor, 26:47324
- OSTASHEVSKII, I. Ya., Use of a model of two classes of tryptophan residues for interpreting the results of ultraviolet fluorescence of cells exposed to x-rays, 26:45736
- OSTERMEIER, R. M., Surface-bound electrons on liquid helium: a negative result, 26:46798
- OSTRETISOV, L. A., Hard gamma radiation from ^{238}Pu , 26:47032
- OTOMO, Norio, Slow neutron spectra in the liquid and solid methane, 26:46858
- OVERMAN, Irving Dean, Coherent photoproduction of rho mesons from deuterium, 26:46445
- OWSIANKOWSKI, M., Effect of ionizing radiation on enzyme activity in the neuroglia of the developing rabbit brain, 26:45836
- OYA, Hiroshi, Turbulence of electrostatic electron cyclotron harmonic waves observed by Ugo 5, 26:46313
- OYANAGI, Y., Polarization in inclusive reactions, 26:46595
- OZAWA, Hiroshi, Nuclear magnetic relaxation study of formic acid. Isotope effect and ^{13}C relaxation, 26:45297
- OZEROV, R. P., Electron density in tungsten crystal according to neutron--electron interaction data, 26:47254
- OZIASHVILI, E. L., Separation of sulfur isotopes by a chemical exchange method, 26:45688 (MLM-1936(TR))

P

- PAC, Pong Youl, Time-dependent neutron transport equation with delayed neutrons, 26:46847
- PACHADZHANOV, D. N., Phenylthiazolylazochromotropic acid as a photocolometric reagent for thorium, 26:45240
- PACHNER, Jaroslav Jr., Optimization of magnetic probe measurements in transient plasmas, 26:47124
- PACK, R. T., Relations between some exponential approximations in rotationally inelastic molecular collisions, 26:46369
- PACKALEN, T., Lymphocyte stimulation by thyroglobulin after local irradiation of the thyroid gland in man, 26:45889
- PACKER, Louis L., Apparatus for measuring height of a molten metal pool, 26:45655
- PADE, B., Thermal plume study at Douglas Point, Lake Huron, 26:45581
- PADIOU, Jean, U_3X_5 type sulfides and selenides of U and Ce, Sr, Ba, Pb, and Eu ($\text{X} = \text{S, Se}$), 26:45378
- PAEDA, R. I., Spectral and scintillation properties of the pyrimidine derivatives, 26:45606
- PAGE, Thornton L., Structure of the Coma Cluster of galaxies, 26:46242
- PAGES, M., Phases of scheelite structure in the systems of plutonium molybdate--lithium or sodium molybdate, 26:45954
- PAIGE, Frank E., Scaling in $\text{p}\bar{\text{v}} + \text{pX}$ and an estimate of the triple-Pomeranchuk coupling, 26:46616 (BNL-16928)
- PAK, Namik K., Final state hadron distributions and generalized scaling laws in deep-inelastic electroproduction processes, 26:46521 (LBL-773)
- PAKHUSOVA, E. V., phi-meson mass determination, 26:46443
- PALCZYNSKI, Ryszard J., Dependence between radioactivity and concentration of atmospheric aerosols, 26:45521
- PALEVSKY, Harry, Direct reactions induced by high energy protons, 26:46891
- PALUSAMY, S., Consistent theory for spherical shells in equilibrium, 26:45453
- PAM ZUI HIEN, Angular distribution of fragments of spontaneously fissioning isomers, 26:47022 (JINR-P7-6466)
- PAN, V. M., Phase diagram of the niobium--gallium system, 26:46018
- Separation processes in superconducting alloys of niobium--aluminum, 26:46020
- System Nb--Al--Ge, its phase diagram, and superconducting properties, 26:46031
- Variations in the critical temperature of superconducting compounds Nb_3Al and $\text{Nb}_3\text{Al}/\text{sub } x/\text{Ge}/\text{sub } 1-x/$, 26:46021
- PANASYUK, V. S., Particle acceleration in a H_{111} cylindrical cavity in an axial magnetic field, 26:46139
- PANNEKOEK, H., In vitro repair of uv-inactivated transforming DNA from *Bacillus subtilis*, 26:45727
- PANT, D. D., Decay times, intensities, and their temperature variation of luminescence of aqueous and deuterated solutions of europic ion, 26:45295
- PAOLI, A., Identification of the superconducting phases in the molybdenum--platinum system, 26:46628
- PAOLONI, F. J., Parametric amplification of Alfvén waves, 26:47187
- PAPAPETROU, A., Elastic excitations excited by gravitational waves, 26:46768
- PAPINI, G., Mass radius of the nucleon, 26:46626
- PARAK, F., Moessbauer effect studies of ^{57}Co and ^{57}Fe in β -rhombohedral B, 26:47280
- PARD, A. G., Studies of void formation in proton-irradiated type 316 and titanium-modified 316 stainless steels, 26:46044 (AI-AEC-13033)
- Swelling as a function of displacement damage in proton-irradiated 316 stainless steel, 26:46054 (CONF-710601, pp 511-21)
- PARFANOVICH, D. M., Recording of alpha particles by solid track detectors based on cellulose, 26:45615
- PARIS, P., Decay of ^{155}Dy , 26:46986
- PARISI, J., Acoplanarity and radiative corrections in Bhabha scattering, 26:46543
- PARK, Richard Dee, X-irradiation of the skin of the beagle, 26:45860
- PARKER, B. R., Synthesis, identification, and purification of C_6 and C_7 fluorocarbons from radiolysis. Final report, 26:45316 (ORO-3909-3)
- PARKER, Donald Lester, Antiproton--proton interactions from 1.51 to 2.90 GeV/c, 26:46469
- PARKER, Robert, Solid state RMS recording ammeter, 26:47634 (UCRL-73587)
- PARKINSON, S., Resistivity of the intermediate state of Type-I superconductors, 26:46823
- PARKINSON, T. M., Protection against gastrointestinal effects of whole-body x-irradiation by a bile acid sequestrant in rats, 26:45844
- PARKINSON, William W., Radiation-resistant plastic insulators, 26:45672

PARLOW, Albert F., Simultaneous radioimmunoassay of plasma testosterone and dihydrotestosterone, 26:45928

PARRISH, C. F., Radiation-induced polymerization. I. A pyrotechnic binder, 26:45325

PARROUR, G., $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)

$\pi^+\pi^-$ production by e^+e^- annihilation at a mass, 26:46427 (CONF-710816-13)

PARSON, M. L., Comparison of theoretical and experimental limits of detection in atomic absorption spectrometry using air-acetylene and nitrous oxide-acetylene flames, 26:45228

PARTENSKY, A., Eigenvalues of the invariant operators of the unitary unimodular group SU(n), 26:46668

PARTHASARADHI, K., Effective atomic numbers for differential processes of gamma-ray interactions, 26:47207

Effective atomic numbers for photoelectric process in various shells in alloys, 26:47206

PARZIALE, E. A., NERVA flight engine control system design, 26:47577

PASCUAL, P., Quasi-elastic neutrino reaction in deuterium, 26:46581

PASYUK, A. S., Production and acceleration of multicharged ions of phosphorus and zinc, 26:46133

PASZTOR, E., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)

PATERA, J., Incompatibility of gauge invariance and nonrelativistic locality in path integral formulation, 26:47292

PATI, J. C., Tests for possible class of CP invariant solutions to K/sub L/ + μ anti μ problem, 26:46577

PATRAULEA, N. N., Random method for Boltzmann's equation, 26:47208

PATTENDEN, N. J., Fission of aligned ^{235}U nuclei by neutrons from 0.4 to 2000 eV, 26:47026

PATTISON, J. B. M., Particle production in proton interactions in nuclei at 24 GeV/c, 26:46459

PATYSHEVA, V. I., Separation processes in superconducting alloys of niobium--aluminum, 26:46020

System Nb--Al--Ge, its phase diagram, and superconducting properties, 26:46031

PAUL, John, Experimental and numerical investigations of rectangular jet, 26:45582

PAULIKAS, G. A., Geomagnetically trapped alpha particles. I. Off-equator particles in the outer zone, 26:46309

PAULUS, J. M., Charge states of ^{208}Pb and ^{208}Tl formed by α decay of ^{212}Po and ^{212}Bi , 26:45359

Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in liquid phase, 26:45362

Chemical consequences of $^{127}\text{I}(n,\gamma)^{128}\text{I}$ in mixtures of I and n-butane in gas phase, 26:45361

PAVLENKO, Yu. G., Induced radiation from neutrons moving through a medium in a magnetic field, 26:46541

Spin effects in theory of synchrotron radiation due to motion of an electron in a medium, 26:46548

PAVLICSEK, I., Determination of isodose curves of sealed γ -sources by calculation and film dosimetry, 26:45715

PAVLOVA, L. M., Study of zirconium evaporation from open surface, 26:46011

PAWASKAR, P. B., Activation analysis with a Pu--Be neutron source, 26:45199

PAXTON, M. M., Recovery and recrystallization of prototypic FTR fuel cladding: 20 percent CW 316 stainless steel tubing (RM-11), 26:46067 (HEDL-TMC-71-126)

PAZ-CASTILLO, Fernando A., Design of waterway connecting Orinoco and Rio Negro Rivers in Federal Territory of Amazonas, Venezuela, 26:45475 (NP-19361)

PEACH, K. J., Measurement of slope parameter in decay K/sub L/0 + $\pi^+\pi^-$, 26:46494

PEDERSEN, G. Fog, Sinusoidal current-phase relation for superconducting thin-film microbridges, 26:46806

PEGEL, V. A., Content of sulphydryl groups in mitochondria after irradiation, 26:45829

PEKER, L. K., Collective states of even--even nuclei of Mo and Ru including unpaired nucleons, 26:46965

Decay of isomers ^{126}I , ^{128}I , ^{130}Sb and quasiparticle levels ^{126}I , ^{128}I , ^{130}Te with large spin, 26:46944

Regularity in multiplet properties of odd--odd nuclei, 26:47051

PELICK, E., Deformation of fuel element rods during irradiation, 26:46070 (UJV-2798-M)

PELEVIN, G. A., Realization of synchronous rotation of some rotors suspended in a magnetic field, 26:45441 (SRARI-P-120)

PELRAS, M., Effect of heat treatments on corrosion of 18/10 stainless steels with low C contents in nitric medium in presence of ferric iron, 26:45970

PELZL, J., Optimum absorber thickness for single line Moessbauer effect in the presence of strong non-resonant absorption, 26:47274

PENAFIEL, H. R., Parallel shooting method for boundary-value problems: application to the neutron transport equation, 26:46859

PENDERE, M. K., Gamma induced changes in spruce wood, 26:45774

PENENGO, P., Galactic gamma radiation between 200 MeV and 10 GeV, 26:46154

PENNER, H., Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)

PENSELIN, S., Direct measurement of the nuclear magnetic

dipole moment of ^{165}Ho , 26:46990

Hyperfine structure of ground state of ^{165}Ho , 26:46989

PENZEL, R., Single-particle spin-orbit potential following from the nucleon--nucleon spin-orbit interaction, 26:47060

PEREAU-LEROY, Pierre, Radiorecovery effect of yeast extract on carnation cuttings irradiated with ^{60}Co γ rays, 26:45777

PEREL'SHEIN, E. A., Instability of axial oscillations in an electron-ring accelerator, 26:46142

Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)

PERES, A., Relativistic wave functions for composite particles, 26:46506

PERESSE, Joseph, Absolute measurement of lifetime of ^4D level of He atom using electron--photon coincidence method, 26:46345

PEREZ, R. B., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)

PERKINS, R. W., Subcritical multiplication of ^{252}Cf neutrons and its applications, 26:46855

PERRET, William R., Seismic-source energies of underground nuclear explosions, 26:45490

PERRIN, D. D., Masking and demasking of chemical reactions. Theoretical aspects and practical applications, 26:45188

PERRIN, M., Many-body realization of Schroedinger algebra, 26:46673

PERRIN, N., Measurement of the quadrupole moment of a shape isomer, 26:46960

PERRIN, R. C., Theory of void formation and growth in irradiated materials, 26:46062 (CONF-710601-, pp 769-97)

PERROT, Michel, New method of nuclear fuel reprocessing, 26:45419

PERSSON, Sven Eric, Interpretation of the neutral helium triplet spectrum in planetary nebulae, 26:46162

PESCHEL, Ingo, Some crystalline field effects in metals, 26:47269

PESSARA, W., Gamma- and x-rays in decay of ^{166}Ho , 26:46998

PETEL, R., Transmission of low-energy x-ray through various locally available materials, 26:47205

PETERMAN, A., Recent developments in the comparison between theory and experiments in quantum electrodynamics, 26:46710

PETERSEN, M. R., Oceanic distributions and relationships of ^7Be and fission products, 26:45530 (BNWL-SA-4205(Rev.))

PETERSON, F. C., Observation of the $M^0(940)$ and $\delta^{00}(963)$, 26:46475 (IS-2886)

PETERSON, J. R., De-excitation of fast He, Ne, and Ar metastable atoms in various gases, 26:46394

PETR, Ivo, Directional properties of cylindrical scintillators screened by a sleeve-type screening, 26:45605

PETRAKOV, A. V., Performance logic of spark chamber information read-out by television automation and data link to a computer, 26:45650

PETRASSO, Richard David, Electric field experiments on the metastable states of helium, 26:46342

PETRICEKS, J., Chatanika, Alaska, auroral-zone incoherent-scatter facility, 26:46283

PETROFF, P., Direct observation of grain boundary dislocation climb in ion-irradiated gold bicrystals, 26:47241

PETROSIAN, Vahe, Ability of the luminosity-volume test to reveal the statistical evolution of the luminosity of quasi-stellar sources, 26:46208

PETROV, M. M., X-ray spectrometer having cooled Si(Li)-detector, 26:45644

PETROV, Yu. V., Green's function of neutron transport equation for moving medium, 26:46850

PETROVA, S. A., Chromosome aberrations in human bone-marrow cells after a single local gamma-irradiation. I, 26:45794

PETRYNA, Alicja, Prognostic elements in inoperable carcinoma of the breast, 26:45783 (IT-54002/4, pp 451-5)

PETERSSON, C., Commercial thimble chambers for absorbed dose measurements at high energy electron radiation, 26:45623

PEYRE-LAVIGNE, A., Noise in Au--nSi diodes operating in the avalanche region in the presence of charges injected by radiation, 26:45660 (FRNC-CONF-29)

PEYRE-LAVIGNE, M., Noise in Au--nSi diodes operating in the avalanche region in the presence of charges injected by radiation, 26:45660 (FRNC-CONF-29)

PEYTURAUX, R., Standard black body at 10,000°K (high pressure plasma arc), 26:47217

PFEIFER, Charles Dean, Energy transfer from a pulsed thermal source to He II below 0.3 K, 26:46797

PFEIFFER, Wayne, Computer simulation of pulsed-neutron experiments for the Fort St. Vrain HTGR, 26:47319 (GULF-GA-B-10959)

PFISTER, A., Radiobiological problems caused by supersonic transport. (With a survey of the first results established by tests performed on board the Concorde prototype), 26:45796

PHALEN, Robert F., Experimental inhalation of metallic silver, 26:45915 (UR-3490-115)

- PHAM DINH LIEN, Deuteron-deuteron scattering, 26:46865
- PHILBIN, J. S., Burnup and power shift effects in large fast breeder reactors, 26:47306
- PHILIBERT, J., Electron probe microanalysis, 26:45246
- Secondary ion emission analysis, 26:45249
- PHILLIPS, A. C., Non-resonant attraction between neutrons in the $\pi^- + {}^3\text{He} \rightarrow \pi^+ + 3n$ reaction, 26:46862
- PHILLIPS, D. L., Thermal plume data acquisition, documentation, and initial analysis, 26:45579
- PHILLIPS, Eleanor L., Suppression and recovery of colony forming units in the rat following sub-lethal x irradiation, 26:45834
- PHILLIPS, J. A., Recent results from the shock-heated toroidal z-pinch experiment ZT-1, 26:47158 (LA-DC-72-726)
- PHILLIPS, James A., All-metal discharge tube, 26:47178
- PHILLIPS, P. E., Influence of reflected ions on the magnetic structure of a collisionless shock front, 26:47199
- PHILLIPS, Perry Edward, Study of the structure of collisionless shock waves propagating perpendicular to an initial field, 26:47198
- PHIPPS, Claude Raymond Jr., Study of optical-scattering methods in laboratory plasma diagnostics, 26:47125
- PICCIOTTO, C., Soft-photon cross sections for $\pi^- \rightarrow n\gamma$, 26:46531
- PIEL, D., Concept of an irradiation facility for investigation of high-temperature thermochemicals at temperatures to 2000°C, 26:45438 (KFK-1556)
- PIEPENBRING, R., Coriolis coupling of bands $1/2^-$ [541] and $3/2^-$ [532] in ${}^{171}\text{Lu}$, 26:46988
- PIETRZAK-FLIS, Z., Long-term study of urinary ${}^{90}\text{Sr}$ and calcium in children, 26:45908
- PIIPARINEN, M., Gamma decay of excited states in ${}^{28}\text{Al}$, 26:46910
- PILIPCHENKO, V. A., Angular distribution of neutrons from (p,n) reactions on ${}^{18}\text{O}$, ${}^{37}\text{Cl}$, ${}^{41}\text{K}$, ${}^{52}\text{Cr}$, and ${}^{59}\text{Co}$ nuclei, 26:46929
- PILYAVETS, V. I., Investigation of positron spectra obtained on linear 8 to 30 MeV electronic accelerator, 26:46149
- PINK, E., Brittle-ductile transition of body-centered cubic metals with special consideration of tungsten and molybdenum. I, 26:45993 (ORNL-tr-2377)
- PINKWART, Harry, Trial of correlating the blood picture with absorbed doses in persons exposed professionally to ionizing radiation, 26:45787 (TT-54002/5, pp 602-4)
- PIOTROWSKI, J., Role of the exit states of a compound nucleus in the emission of high-energy γ -rays, 26:47082
- PIPER, B. C., Fire assay-emission spectrographic determination of iridium, ruthenium, and osmium, 26:45229
- PIRK, H., Process for recovering by-products from uranium dioxide production, 26:45371
- PIRRI, A. N., Momentum transfer and plasma formation above a surface with high-power CO_2 laser, 26:47155
- PISKUNOV, L. I., Radioisotope accumulation in hydrobionts, 26:45926
- PITTENDRIGH, L. W. D., Monitor for estimating tritium in urine, 26:45585 (AERE-HP/M-74)
- PITTMAN, Roland Nathan, Forward differential cross section in anti pp + anti nn, 26:46470
- PIWNICA, A., Radioisotopic pacemaker developments in France, 26:45699 (CEA-CONF-1998)
- PLANCK, R., Equation of state for liquid ammonia, 26:46079 (UCRL-Trans-10622)
- PLASIL, F., Limits on angular momentum in heavy-ion--compound-nucleus reactions, 26:47072
- PLAVNIK, G. M., Radiation damage to polymer fiber investigated by small-angle x-ray scattering, 26:45338
- PLENDL, H. S., Cross sections for (π^- , π^-n) reactions on ${}^{19}\text{F}$ and ${}^{21}\text{p}$ in the 3 to 3 resonance region, 26:46486
- PLESCH, Rolf, Radioisotopic thickness measurements in the manufacture of wire, 26:45717
- PLUMB, I. C., Technique for recording Langmuir probe characteristics in afterglow plasmas, 26:47119
- POCHAT, Amedee, Absolute measurement of lifetime of 4d level of He atom using electron--photon coincidence method, 26:46345
- POHLIT, W., Sensibilisation in radiobiology and radiotherapy, 26:45845
- POHLMAYER, K., Equation curl $\mathbf{W}/\text{sub } \mu/(\mathbf{X}) = 0$ in quantum field theory, 26:46717
- Higher order perturbation theory for exponential Lagrangians, 26:46706
- POLAK, B., Thermoluminescent dosimetry system for determining quality, 26:45622
- POLAK, Bogumil, Distribution of doses in the lesser pelvis in the treatment of carcinoma of the cervix uteri, 26:45784 (TT-54002/4, pp 456-61)
- POLENOV, B. V., Characteristics of charged-particle detection sections having channel electron multipliers, 26:45645
- Low-energy-electron spectrometer having a multiplate analyzer, 26:45643
- POLKINGHORNE, J. C., Deep inelastic electroproduction and related processes, 26:46515 (DAMP-71/24)
- POLKOW, Bogdan, Distribution of doses in the lesser pelvis in the treatment of carcinoma of the cervix uteri, 26:45784 (TT-54002/4, pp 456-61)
- POLOTSKII, I. G., Determination of the energy of dislocation of an impurity atom by an internal friction method, 26:46023
- POLOVIN, R. V., Impossibility of introducing "hidden" parameters into quantum mechanics, 26:47285 (ITF-71-101-P)
- POLSON, H. J., Removal of radioactive iodine from ventilation exhaust streams at the AAEC Research Establishment, 26:45502
- POLUEKTOV, N. S., Microdeterminations of terbium by a luminescent method with the reagent pyrogallolsulfonic acid, 26:45231
- POLYAKOV, V. I., Determination of the concentration of gamma-emitting isotopes in extended sources without sampling, 26:45264 (SRARI-P-123)
- POLYAKOV, V. N., Computer systems for physics research centers, 26:45686
- PONS, Roland, Integration of Newman--Penrose equations in Minkowski space, 26:46769
- PONTECORVO, B., Possible existence of hadronic isomers, 26:46588
- POPESCU, G., New method for potassium determination by neutron activation in mineral and rock samples, 26:45197
- POPESCU, Ion M., Effect of a particles on operating regime of a CO_2 laser, 26:45669
- POPLEY, R. A., Tubular Nb/Cu conductors for ac superconductive power transmission, 26:46844
- POPOV, V. I., Excitation cross section of first $2+$ levels of Ni and Zn nuclei, 26:46936
- POPPEMA, O. J., Cyclotron laboratory at Technical University of Eindhoven, 26:46146
- PORTAL, G., Utilization of alumina radiothermoluminescence in zone dosimetry, 26:45628
- PORTER, S. G., Ferroelectrics in $\text{K}_2\text{O--SrO--Nb}_2\text{O}_5$ system, 26:45962
- POSSA, G., Threshold of parallel channel flow instability, 26:45459 (CISE-R-275)
- POST, Richard F., Conversion of high-temperature plasma energy into electrical energy, 26:45433
- POST, Roy G., Techniques for source determination in the Tucson basin, 26:45206
- POSTMA, H., Fission of aligned ${}^{233}\text{U}$ nuclei by neutrons from 0.4 to 2000 eV, 26:47026
- POTEMPA, A. V., Study of the decay ${}^{186}\text{Tb}$ ($T_{1/2} = 66$ min) and ${}^{180}\text{Tb}$ ($T_{1/2} = 3.48 \pm 0.16$ h), 26:46942 (JINR-P6-6512)
- POTOCZAK, M. J., Inclusive reaction $\mu + \text{nucleon} \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV/c, 26:46452
- POTTER, William R., Radical ion saturation in some sulfur compounds x-irradiated at 4.2°K , 26:45348
- POUBEAU, Pierre, Ultracentrifuge rotor, 26:45697
- POUILLANGE, J.-P., Study of losses in an assembly of superconducting filaments transposed in a conducting matrix and subjected to a pulsed magnetic field, 26:46810
- POULET, M., Polarization in $\pi^+ \rightarrow \pi^-$ backward elastic scattering at high energy, 26:46490
- POULSEN, Lise Lotte, Optical model for vibrational relaxation in reactive systems, 26:46375
- POWELL, Jack E., Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)
- POWERS, John Richard, Nuclear structure of ${}^{23}\text{Na}$: charged particle reaction studies, 26:46915
- PRADHAN, T., Low-energy theorem for pion--nucleon scattering from pion source dynamics, 26:46641
- PRAHL, Joseph, Experimental and numerical investigations of rectangular jet, 26:45582
- PRASAD, Kadambi Narasimha, Investigation into the application of post-cutting chip activation analysis to the study of tool wear, 26:45713
- PRASTEIN, S. M., Thermal plume data acquisition, documentation, and initial analysis, 26:45579
- PRATT, Ivor, Adhering resins to substrates, especially metal, by radiation, 26:46089
- PREDA, Alexandru M., Effect of a particles on operating regime of a CO_2 laser, 26:45669
- PREDAZZI, E., General properties of correlation functions for inclusive processes, 26:46502
- PREPEJCHAL, William, Radionuclide concentrations near the Big Rock Point Nuclear Power Station, 26:45536
- PRESTON, J. A., Ionization of argon by electron impact, 26:46374
- PRETRE, S., Quantities for description of neutron irradiations of man, 26:45789
- PRICE, P. B., Composition and energy spectra of heavy nuclei with $0.5 < E < 40$ MeV per nucleon in the January 24, and September 1, 1971 solar flares, 26:46232
- PRICE, Richard H., Nature of gravitational synchrotron radiation, 26:46757
- PRIGENT, Jacques, U_3X_5 type sulfides and selenides of U and Ca, Sr, Ba, Pb, and Eu ($X = \text{S, Se}$), 26:45378
- PRILEZHAEVA, N. A., Manifestation of structure in electron energy distribution functions and determination of the effective cross sections of inelastic processes, 26:46407
- PRILUTSKII, R. E., Use of admission system of MI-1309 instrument for precision isotopic analysis, 26:45681
- PRINCEHOUSE, D. W., High-resolution heat-capacity study of ${}^4\text{He}$ adsorbed on bare copper, 26:46804
- PROSEN, Marijan, Stars with dark companions, 26:46197
- PROTOPOESCU, V., New results in theory of pulsed thermal

neutrons, 26:46849
 PROTOPOPOV, V. N., ($n, n'\gamma$) reaction in ^{27}Al , ^{51}V , ^{52}Cr , ^{55}Mn , ^{56}Fe , ^{93}Nb , ^{208}Pb , and ^{209}Bi , 26:46909
 PRUGOVECKI, Eduard, Scattering theory in Fock space, 26:46783
 PRUSSIN, S. G., Decay of ^{210}At to levels in ^{210}Po , 26:47006
 PUAUX, J. P., Determination of the structure of monohydrated uranyl diformate using x-ray diffraction, 26:45368
 PUGACHEV, G. D., Photoproduction of charged pions by nuclei at $E_0 = 1160$ MeV, 26:46441
 PUGACHEVA, T. S., Distribution of alkali ions in nickel by ionic bombardment, 26:47235
 PULSTOV, D. D., Angular distribution of fragments of spontaneously fissioning isomers, 26:47022 (JINR-P7-6466)
 PURSIHEIMO, V., Decay of $1.4 \text{ min } ^{173}\text{Er}$, 26:46997
 PURYEAR, Dean A., Phosphate solidification of caustic process solution waste, 26:45381 (ARH-SA-120)
 PUTNAM, John H. Jr., Sequential sampler, 26:45613
 PYATKIN, E. K., Chromosome aberrations in human bone-marrow cells after a single local gamma-irradiation. I, 26:45794
 PYRAGAS, Kazimieras A., Covariant canonical equations of the test-particle motion in general relativity, 26:46770

Q

QUENBY, J. J., Entry of high-energy solar protons into the distant geomagnetic tail, 26:46304
 QUINLAN, W. J. Jr., Ionizing radiation-induced thyroid carcinogenesis in the dog, 26:45819 (UR-3490-63)
 QUINN, H. R., Diffraction-dissociation model for high-energy electroproduction, 26:46517 (DESY-72/23)
 QUINN, W. E., Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
 QUINTILIANI, Marcello, Factors affecting radiation sensitivity, 26:45739

R

RADHAKRISHNAN, T. P., Absorption of cathodic hydrogen by iron and steel, 26:45986 (BARC-585)
 RADYUK, G. A., Elastic scattering of protons by deuterons at energies of 14, 16, and 17.6 MeV, 26:46863
 RADZHAPOV, P. S., ^{135}Ba levels, 26:46945
 Magnetic properties of ^{153}Eu , ^{155}Eu , and ^{189}Os , 26:46981
 RAE, Ernest R., Neutron scattering and capture cross-section measurements, 26:46968
 Pulsed accelerator time-of-flight spectrometers, 26:45641
 RAE, H. L., Effect of gamma irradiation on Fuerte avocado fruits, 26:45724
 RAEF, L., Stenosis of the small bowel as a complication in radiation therapy of carcinoma of the uterine cervix, 26:45802
 RAFAELOFF, R., Chemistry of gallium [1--5]. VI, 26:45285
 RAGHUNATHAN, V. S., Chemical diffusion in the beta phase of zirconium-titanium system, 26:45985 (BARC-577)
 RAGOOWANSI, N. L., Analog states in ^{33}Cl by isospin-forbidden resonances in reaction $^{32}\text{S}(p, \gamma)^{33}\text{Cl}$, 26:46902 (BARC-598)
 RAGSDALE, H. L., Ecological and environmental effects from local fallout from Schooner. 2. The beta and gamma radiation effects from close-in fallout, 26:45542 (PNE-529)
 RAHMAN, M. Matlur, Utilization of research reactors for radioisotope production, 26:45396
 RAHMAN, Naseem K., Multiphoton transition in hydrogen between 1S and 2P levels, 26:46356
 RAHN, John Phillip, Magnetic surface states in zinc and cadmium, 26:47256
 RAHN, Kenneth A., Multi-element analysis of air pollution particulates by nondestructive neutron activation, 26:45207
 RAI, D. K., Cross sections for single-electron capture in collisions of protons and deuterons with atoms, 26:46398
 RAICHEV, P. P., Parametrization of $B(E2)$ -transition in deformed even-even nuclei within $SU(3)$ scheme, 26:46975 (JINR-P4-6462)
 Possible existence of $SU(3)$ multiplets in spectra of deformed even-even nuclei, 26:47048 (JINR-P4-6452)
 $SU(3)$ physical basis realization and probabilities of $B(E2)$ -transitions in $SU(3)$ scheme. II, 26:47049 (JINR-P4-6458)
 RAIMOND, Ernst, Radio map of spiral galaxy Maffei 2 at 1415 MHz, 26:46237
 RAJAGOPALAN, Mahalingam, Long range alpha particle emission in the fission of heavy elements, 26:47037
 RAJARAMAN, R., Some consequences of the vanishing of the triple-pomeronchukon-vertex, 26:46597
 RAKHMATULLAEV, N., Al-Fe-Mo oxide catalysts. I, 26:45963
 RAKOV, Yu. A., Photoproduction of neutral pions on helium nuclei, 26:46432 (SLAC-Trans-142)
 RALPH, B., Field-ion microscope study of ion-implantation in Ir. II, 26:47240
 Field-ion microscope study on ion-implantation in Ir. I, 26:47239
 RAM, Rishi, Vorticity jump across a stationary magnetogasdynamic discontinuity, 26:46421
 RAMACHANDRAN, J., Dynamic response of a plate to time-dependent boundary conditions, 26:45443
 RAMAMURTHY, V. S., Theory of fragment mass and charge distribution in fission, 26:47044 (INDC-7-U, pp 65-79)
 RAMAMURTY, S., Effective atomic numbers for photoelectric process in various shells in alloys, 26:47206
 RAMAN, S., Coulomb excitation of ^{117}In , ^{119}Sn , 26:46961
 RAMA RAO, J., Effective atomic numbers for differential processes of gamma-ray interactions, 26:47207
 RAMASWAMY, M. K., Doppler-broadened annihilation radiation in metals with a Ge(Li) detector, 26:47268
 RAMOND, P., Virasoro group in dual-resonance models, 26:46686
 RAMSEY, N. F., Molecular-beam magnetic-resonance measurements of the anisotropies of the electric polarizabilities of H_2 and D_2 , 26:46358
 RANDOW, F., Artificial radioactivity of fallout and food in the northern districts of GDR in the year 1969, 26:45546
 RANEV, N. V., Electron density in tungsten crystal according to neutron-electron interaction data, 26:47254
 RANKIN, J. M., Density of the solar corona from occultations of NP 0532, 26:46227
 RANSOHOFF, Jackson A., Pool irradiator with radiation source in a rotating container, 26:45446
 Production irradiator, 26:45447
 RAO, K. R., Simple transistorised control unit for neutron crystal spectrometers, 26:45636 (BARC-591)
 RAO, M. H., Thermodynamic properties of the phase In_3Bi_3 , 26:46026
 RAO, S. P., Selective extraction of Ga with N-benzylaniline in chloroform from hydrochloric acid media, 26:45284
 RAO, V. V. Ramana, Secondary flow of a conducting fluid in a rotating straight annular pipe, 26:46422
 RAPP, R. A., Physicochemical measurements in metals research. Part 1, 26:46014
 Physicochemical measurements in metals research. Part 2, 26:46015
 RASMUSSEN, J. J., Surface tension, density, and volume change on melting of Al_2O_3 systems, Cr_2O_3 , and Sm_2O_3 , 26:45949
 RASPOPIN, S. P., Thermodynamic properties and phase composition of La-Zn alloys, 26:46001
 RASULOV, E. N., Elastic scattering of protons by deuterons at energies of 14, 16, and 17.6 MeV, 26:46863
 RASZILLIER, I., Information on pion electromagnetic form factor inside its analyticity region provided by bounds on its modulus on cut (t greater than or equal to $4m_\pi^2$), 26:46528
 RATCLIFF, Blair Norman, Rho-meson production by pions at 15 GeV/c: a study of vector dominance and of rho-omega interference, 26:46496
 RATHI, S. K., Increased androgen production in fish following irradiation, 26:45898
 RATNER, L. G., 200-GeV proton-proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)
 RAUBENHEIMER, B. C., Rigidity dependence of cosmic-ray modulation at rigidities > 2 GV in August 1969, 26:46327
 RAUCH, E. L., Determination of semiconductor junction profiles by angle lapping of semiconductor wafers, 26:47608 (BDX-613-604(Rev.))
 RAUNEMAA, T., Decay of $1.4 \text{ min } ^{173}\text{Er}$, 26:46997
 RAVOIRE, Jean, Isotopic enrichment of hydrogen for heavy water production, 26:45696
 Method of preparing secondary alkaline amine derivatives, 26:45289
 RAY, J. R., Tachyon dust metric in general relativity, 26:46754
 RAY, Swati, Inversion from the energy dependence of total cross sections, 26:46387
 RAYNAL, Jacques, Geometrical properties of nuclear reactions, 26:47041 (CEA-N-1529)
 RAZBUDEI, V. F., Temperature dependence of cross section of cold- and thermal-neutron scattering by O_2 and D_2 , 26:46887
 RAZORENOV, L. A., Removable current lead for power supply in superconducting magnetic systems, 26:45679
 READ, B. J., Is the pion form factor determined by electroproduction experiments, 26:46513 (CONF-710816-17)
 REAMES, D. V., Comparison of measurements of the charge spectrum of solar cosmic rays from nuclear emulsions and the Explorer 35 solid-state detector, 26:46330
 REBANE, T. K., Variational bounds for imaginary frequency polarizability and dispersion interaction constant, 26:46415
 REBENKO, A. L., Equations for matrix elements of Euclidean quantum electrodynamics, 26:46739
 RECAMI, E., Superluminal frames and group of generalized Lorentz transformations in four dimensions, 26:46734
 RECUIERO, A., Magnetohydrodynamic entrance flow in a channel, 26:46423
 REDDY, D. R. K., Approximate solution for the static, spherically symmetric metric due to a point charged mass in Brans-Dicke theory, 26:46750

- REDDY, G. R., Activation analysis with a Pu--Be neutron source, 26:45199
- REDDY, J. P., Magnetic phase diagram of the UAs--US system, 26:47261
- REED-HILL, Robert E., Tensile properties of α --Zr, 26:46040
- REHLER, Nicholas E., Theory of cooperative emission from two-level atoms, 26:46411
- REIN, James E., Rapid mass-spectrometric determination of boron isotopic distribution in boron carbide, 26:45235
- REISS, Howard R., Multiphoton transition in hydrogen between 1S and 2P levels, 26:46356
- REKALO, A. P., Interaction processes of nonzero rest mass ν and anti ν with leptons, 26:46579
- REKALO, M. P., Violation of Pomeranchuk theorem and relations between the differential cross sections, 26:46645
- REMSEN, Joyce F., Ultraviolet inactivation and miscoding of irradiated R17-RNA in vitro, 26:45728
- RENARD, F. M., Connections between low energy and high energy K^+ photoproduction, 26:46535
- e^+e^- annihilation into hadrons, 26:46540
- RENAULT, H., Bleomycin labeled with ^{57}Co , 26:45929
- RENGAN, K., Hindered $7/2^+ + 3/2^-$ β -decay of ^{133}Pr , 26:46955
- REPELLIN, J. P., $\pi^+\pi^-$ production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
- $\pi^+\pi^0$ production by e^+e^- annihilation at ω mass, 26:46427 (CONF-710816-13)
- REPNOW, R., Missing low-lying states in ^{39}K , 26:46928
- REVESZ, L., Variations in the bone marrow lymphoid population in the sublethally irradiated germ free mouse, 26:45843
- REYNOLDS, T., Doppler-broadened annihilation radiation in metals with a Ge(Li) detector, 26:47268
- REYNOLDS, Warren L., Iron(II) and Iron(III) isotope exchange in water--dimethyl sulfoxide mixtures, 26:45304
- RHEE, S. K., Critical surface energies of Al_2O_3 and graphite, 26:45944
- RHOADS, William A., Ecological and environmental effects from local fallout from Schooner. 2. The beta and gamma radiation effects from close-in fallout, 26:45542 (PNE-529)
- RIBE, F. L., Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
- RIBLET, G., Anomalous critical field curves of Kondo superconductor $(\text{La}/\text{sub } 1-x/\text{Ce}/\text{sub } x)/\text{Al}_2$, 26:46832
- RIBON, Pierre, Radiations emitted in fission, 26:47021 (INDC-7-U, pp 131-44)
- RICCI, J. L., (^{10}O , ^{12}C) studies in the $2s$ -- $1d$ shell, 26:46913
- RICHARDSON, R. C., Spin-wave phenomena in liquid ^3He systems, 26:46802
- RICHMOND, Robert G., Radiation dosimetry for the Gemini program, 26:47203 (N-72-18797)
- RICHT, A. E., Evaluation of absorber materials performance in HFIR control cylinders, 26:47587
- RICQUE, R., Design considerations of heat exchangers for vaporizing fluids, 26:45436 (CEA-CONF-2010)
- RIEDL, P., New 5 MeV Van de Graaff ion accelerator at Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, 26:46118 (KFKI-71-47)
- RIEGER, T. J., Time-dependent superconductivity and quantum dissipation, 26:46821
- RIESS, F., Excited states of ^{21}Ne , 26:46907
- RILEY, J. M., Observations of $^{30}\text{Cl } 272.1$ at 2.7 and 5.0 GHz, 26:46200
- RILEY, P. J., $^{90}\text{Zr}(d,n)^{91}\text{Nb}$ and $^{90}\text{Zr}(d,n)^{91}\text{Nb}$ reactions, 26:46956
- RINAT, A. S., Vertex parameters and coupling constants for the virtual decay of light nuclei, 26:47063
- RINAUDO, G., Evidence for double diffractive dissociation in $\pi^+\text{p}$ reactions at 5 GeV/c, 26:46493
- RINGLAND, G. A., Phenomenological prescription for Regge cuts, 26:46622
- RINNE, R. L., An introduction to cobordism theory, 26:47627 (SCL-RR-720270)
- RIOLA, J. P., Chemi-ionization in collisions between helium metastable atoms and argon, 26:46404
- RISING, F. L., Cumulative dose from diagnostic irradiation, 26:45799
- RISKA, D. O., Two-pion exchange AN potential, 26:46652
- RITCEY, G. M., Zirconium tetrafluoride process, 26:45274
- RITZMAN, R. L., Environmental benefit-cost analysis for nuclear power generation, 26:47386
- RIVET, P., K_2^0 decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)
- RIZZUTO, D., Superconductivity and localized spin fluctuations in concentrated aluminum--3d-transition-metal alloys, 26:46838
- ROACH, Pat R., Absence of a quadratic term in the ^3He excitation spectrum, 26:46799
- ROBBINS, D. E., Velocity and flux dependence of the solar-wind helium abundance, 26:46328
- ROBBINS, J. A., Multi-element analysis of air pollution particulates by nondestructive neutron activation, 26:45207
- ROBERT, J., Bleomycin labeled with ^{57}Co , 26:45929
- ROBERTS, Anne, Procedure for chronic intravenous catheterization in the rabbit, 26:47641 (UR-3490-21)
- ROBERTS, J. B., 200-GeV proton--proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)
- ROBERTS, Paul A., Differences in synaptic affinity of chromosome arms of *Drosophila melanogaster* revealed by differential sensitivity to translocation heterozygosity, 26:45876
- ROBERTS, R. G., Phenomenological prescription for Regge cuts, 26:46622
- Regge cuts from inclusive finite mass sum rules, 26:46651 (RPP/T-18)
- ROBINSON, Mark T., Computer simulation of atomic displacement cascades in metals, 26:46061 (CONF-710601-, pp 739-56)
- Dependence of radiation effects on the primary recoil energy, 26:46051 (CONF-710601-, pp 397-429)
- ROBINSON, R. L., Coulomb excitation of $^{117}\text{In}^{95}\text{Sn}$, 26:46961
- ROBINSON, Ramon K., Process of forming an isotopic heat source, 26:45707
- ROBLE, R. G., Thermospheric molecular oxygen from solar extreme-ultraviolet occultation measurements, 26:46269
- ROBSON, A. E., Influence of reflected ions on the magnetic structure of a collisionless shock front, 26:47199
- ROBSON, R. J., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)
- ROCA, M., Application of spectrographic analysis to the radioisotope production control. II, 26:45223 (JEN-232-DQ/I-81)
- ROCHE, R., Calculations and behavior of expansion bellows, 26:47591
- ROCHESTER, G. K., Galactic gamma radiation between 200 MeV and 10 GeV, 26:46154
- ROCQUET, G., Effects of thyroidian hormones and γ irradiation on adenylylase of plasma membranes of rat liver, 26:45831
- RODDEN, Clement J., Selected measurement methods for plutonium and uranium in the nuclear fuel cycle. Second edition, 26:45183 (TID-7029(2nd Ed.))
- ROE, Kenneth A., Trends in capital costs of generating plants, 26:47385
- ROGANI, A., Surface excitations of Type-I superconductors in a static magnetic field, 26:46830
- ROGERS, L. B., Technical progress report on fundamental studies of separation processes, 26:45302 (COO-1222-55)
- ROGGEN, R., Identification of the superconducting phases in the molybdenum--platinum system, 26:46828
- ROHLOFF, F., Scattered neutrons and their effects on calibration of Bonner sphere counters, 26:45599
- ROLFS, C., Excited states of ^{21}Ne , 26:46907
- ROMANOV, V. S., Elastic and inelastic scattering of 6-MeV protons by even isotopes of Cr, 26:46922
- ROMANOV, Yu. I., Lepton charge and weak neutrino--lepton interaction, 26:46578
- ROMBERG, Gerald P., Radionuclide concentrations near the Big Rock Point Nuclear Power Station, 26:45536
- ROMICK, G. J., HB/N_2^+ 4709-A intensity ratio and vibrational enhancement of N_2^+ first negative bands in proton-excited auroras, 26:46276
- ROMNEY, E. M., Radioecology and ecophysiology of desert plants at the Nevada Test Site, 26:45755 (TID-25954)
- ROOD, Herbert J., Structure of the Coma Cluster of galaxies, 26:46242
- ROSATI, Alfredo, Phase sensitive method for detecting superconductive transitions in the microwave region, 26:46840
- ROSCHKE, E. J., Experimental investigation of heat transfer from partially ionized argon with an applied transverse magnetic field, 26:45464
- ROSE-INNES, A. C., Resistivity of the intermediate state of Type-I superconductors, 26:46823
- ROSEN, G., General formula for obtaining expectation values from the hypercharacteristic function, 26:47289
- ROSEN, Gerald, Exact solutions to the Yang--Mills field equations, 26:46689
- ROSEN, Lon, $(\phi/\text{sup } 2\pi/\text{)}_2$ field Hamiltonian for complex coupling constant, 26:46718
- ROSEN, V., Model studies of density fluctuations of small and large ions in the atmosphere, 26:46288
- ROSENBERG, Leonard, Method of effective potentials: scattering of positrons and electrons by light atoms, 26:46400
- ROSENTHAL, M. W., Metabolism of monomeric and polymeric Pu in the rabbit. Comparison with the mouse, 26:45917
- ROSENZWEIG, C., Photonlike particles, Compton scattering, and sum rules in the dual-resonance model, 26:46601
- ROSS, D. A., Radiative corrections to muon decay, 26:46576
- ROSS, J. S., Electron spin resonance in single crystal yttria stabilized zirconia, 26:47252
- ROSS, John, Optical model for vibrational relaxation in reactive systems, 26:46375
- ROSSELLE, N., Changes in the localization of acetylcholinesterase in neuromuscular junction after ^{60}Co irradiation, 26:45828
- ROSSITER, Bryant W., Physical methods of chemistry. Part IIIB, 26:45250
- ROSSMOORE, H. W., Effect of ^{137}Cs radiation on growth, development, and mortality of larvae of the tussock moth, *Homocampa leucostigma*, 26:45885
- ROST, M., Evidence for double diffractive dissociation in $\pi^+\text{p}$ reactions at 5 GeV/c, 26:46493

ROSTAD, Lynne C., Nuclear-cytoplasmic relations in the mitosis of sea urchin eggs. III. γ -ray-induced damage to whole eggs and nucleate and anucleate half-eggs, 26:45874

ROSTAGNI, G., Experiments on a superposed fields theta pinch, 26:47171

ROSZMAN, Larry Joe, Effects of ion motion on spectral line broadening in plasmas, 26:47148

ROTH, B. F., Special machines and equipment of WAK, 26:46106 (ORNL-tr-2372)

ROTH, Etienne, Isotopic enrichment of hydrogen for heavy water production, 26:45696

ROTHMULLER, I. J., Conjugate point phenomena, 26:46323

ROTI ROTI, J. L., Effects of actinomycin D on G₁ protein synthesis in L5178Y cells, 26:47650 (UR-3490-153)

Protein synthesis during the cell cycle of L5178Y cells, 26:47651 (UR-3490-154)

ROTI ROTI, Joseph Lee, Protein synthesis and the cell cycle of L5178Y cells, 26:47640 (UR-3490-18)

ROUCHE, N., Postgraduate training of scientists in the European Community, 26:47616 (EUR-4832)

ROUGE, A., Reactions $K^- + p \rightarrow \Lambda(\Sigma^0) +$ vector meson at 3.95 GeV/c, 26:46485

ROUSSEL, Joseph, Capture of low-energy rare gas ions by metallic targets, 26:46335 (PRNC-TH-198)

ROUTTI, Jorma T., Dose estimates outside the shielding of the CERN Intersecting Storage Rings by the Moyer method, 26:46121

ROWCLIFFE, A. F., Void formation in quenched stainless steel, 26:45987 (CONF-710601-, pp 605-12)

ROY, D. P., Complex conjugate Regge poles, 26:46631

Phenomenological prescription for Regge cuts, 26:46622

Regge cuts from inclusive finite mass sum rules, 26:46651 (RPP/T-18)

Regge phenomenology of inclusive spectra in the central region, 26:46659

ROY, T., ρ - ρ couplings and t-channel helicity conservation, 26:46638

Bound states in quantum electrodynamics, 26:46721

ROYCHOWDHURY, A., ρ - ρ couplings and t-channel helicity conservation, 26:46638

RUBIN, Sylvan, Elastic nuclear scattering techniques, 26:45271

RUDD, M. E., Observation of a broad resonance in the 2^3S excitation of helium by electron impact, 26:46405

RUDDICK, I., Persistent current states in rotating superfluid helium, 26:46801

RUEHL, Franklin R. Jr., Quadrupole moments of light nuclides, 26:46892

RUEHLE, R., Computer-code for solution of three-dimensional-multigroup-equations, based on improved synthesis-method, 26:47594 (BMBW-FBK-72-6)

RUKOLAINE, G. V., Realization of synchronous rotation of some rotors suspended in a magnetic field, 26:45441 (SRARI-P-120)

RUMI, G.-C., New method for the determination of N(h) in the D-region, 26:46298

RUMPF, G., Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)

RUNDEL, R. D., Chemi-ionization in collisions between helium metastable atoms and argon, 26:46404

RUSSEL, J. J., Metabolism of monomeric and polymeric Pu in the rabbit. Comparison with the mouse, 26:45917

RUSSELL, B. G., Pressed fused disc: a technique for preparation of ore samples to be analyzed for uranium by X-ray fluorescence, 26:45262 (NIM-1427)

RUSSELL, C. T., Detection of solar-wind electron plasma frequency fluctuations in an oblique nonlinear magnetohydrodynamic wave, 26:46329

RUSTAD, Ronald C., Nuclear-cytoplasmic relations in the mitosis of sea urchin eggs. III. γ -ray-induced damage to whole eggs and nucleate and anucleate half-eggs, 26:45874

RUSU, F., Influence of neutron irradiation on seed germination and plant growth of tomatoes, 26:45773

RUTH, R. D., Analysis of five line hyperfine split Moessbauer absorber spectra, 26:47275

RYABCHIKOVA, G. G., Radiation damage to polymer fiber investigated by small-angle x-ray scattering, 26:45338

RYABOV, V. P., Ternary system Y--Fe--Al in region 0 to 33.3 percent Y, 26:46005

RYBCHIN, V. N., Lethal and mutagenic effect of ionizing radiations with different LET on extracellular bacteriophages, 26:45746

RYBICKI, K., Two-particle correlations in 60 GeV/c π -p and 67 GeV/c pp inclusive reactions, 26:46460

RYBIN, Yu. A., Substance purification by zone melting in a rotating container, 26:45276

RYCROFT, M. J., Ring current effect on magnetospheric electron density profiles derived from plasmopause whistlers, 26:46284

RYDE, H., Validity of rotor model in odd-Z nuclide ^{171}Lu , 26:46995

RYKHAL, R. M., Ternary system Y--Fe--Al in region 0 to 33.3 percent Y, 26:46005

RYZHIKOV, I. V., Investigation of the injection characteristics of symmetrical pin-diodes prepared by the method of ion implantation, 26:45678

RYZHKOV, V. M., Particle acceleration in a H_{111} cylindrical cavity in an axial magnetic field, 26:46139

S

SABATIER, Pierre C., Complete solution of the inverse scattering problem at fixed energy, 26:46784

SABATIER, Rene, Hexavalent W bronzes containing Th in interstitials: $\text{Th}/\text{sub } x/4/\text{sup IV}/(\text{B}/\text{sub } x/\text{sup V}/\text{W}/\text{sub } 1-x/\text{sup VI})\text{O}_3$ phases (B = Nb, Ta), 26:45953

SABIROV, Kh. S., ^{135}Ba levels, 26:46945

Excitation of ^{99}Tc nucleus, 26:46951

SABO, V. I., Photoproduction of vector and axial vector resonances in broken nonlinear chiral $\text{SU}_3 \times \text{SU}_3$ dynamics, 26:46563

SABOL, Jozef, Circuit for continuous acquisition of rate ratio of two pulse trains, 26:45675

SACHARIDIS, E. J., Plastic threshold Cherenkov detectors with wavelength shifter and quencher, 26:45646 (RPP/H-89)

SADOWSKA, Barbara, Method for labeling human albumin with ^{99m}Tc , 26:45927 (TT-54002/4, pp 471-4)

SAEGUSA, Kenji, Subtraction and positron scanning with the dual headed whole body scanner, 26:45593

SAEKI, Masakatsu, Ranges of fission fragments in uranium--aluminum alloys and uranium oxides, 26:46074

SAENKO, E. M., Electron transfer during zone melting of yttrium, 26:45975

SAFAROV, A., Distribution of alkali ions in nickel by ionic bombardment, 26:47235

SAGAN, Carl, Interstellar organic chemistry, 26:46240

SAGREDO, J. L., Ring current effect on magnetospheric electron density profiles derived from plasmopause whistlers, 26:46284

SAHNI, V., Analytic calculation of the coherent atomic scattering factor for the ground state of the helium isoelectronic sequence, 26:46359

SAITO, Hiroshi, Convergent hole collimator for transmission scintiphotograph, 26:45602

SAITO, Takao, Daytime irregular geomagnetic pulsation, 26:46317

SAJI, Y., $^2\text{H}(n,p)^2\text{n}$ reaction at 50 MeV and neutron--neutron scattering length, 26:46861

SAKAI, Yoshiko, Pair correlation energies as derived from the analysis of semi-empirical values of correlation energies of atoms, 26:46412

SAKAMOTO, Yoshio, Hepatic injuries following irradiation, 26:45809

SAKURAI, T., Fluorination of UO_2 and U_3O_8 by gaseous Br trifluoride, 26:45379

SALAKHITDINOV, A. N., Collective--single-particle states of deformed odd nuclei, 26:47081

SALAM, A., Gravitational correction to PCAC, 26:46501

SALAMA, M., Fast neutron elastic scattering experiments, 26:46916 (AE-452)

SALAMONI, E., Superconductivity and localized spin fluctuations in concentrated aluminum--3d-transition-metal alloys, 26:46838

SALLACH, R. A., Initial reaction of nitrogen with zirconium at 1440°, 26:46030

SALMON, G. A., Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. II, 26:45328

Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. I, 26:45331

Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. III, 26:45329

Radiation-induced formation of excited states of aromatic hydrocarbons in cyclohexane. IV, 26:45330

SALOMAA, Pentti, Solvent deuterium isotope effects in acid-base reactions. VII, 26:45292

SALPETER, E. E., Abundances of interstellar molecules, 26:46244

SALTMARSH, M. J., Energy shifts and relative intensities of K x rays produced by swift heavy ions, 26:46339

SALUJA, J. K., NERVA flight engine control system design, 26:47577

SALUJA, P. P. S., Ionic solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

SAMIGOV, K. A., Al--Fe--Mo oxide catalysts. I, 26:45963

SAMOEL, Alain, Ultrasound quality control installation, 26:45472

SAMOVAROV, V. S., Use of solid track detectors in reactor experiments, 26:45610

SAMS, W. Neal, Differential- and total-cross-section calculations for models of e^-H_2 collisions, 26:46399

SAN, R. H. C., Various levels of DNA repair synthesis in xeroderma pigmentosum cells exposed to carcinogens N-hydroxy and N-acetoxy-2-acetyl-aminofluorene, 26:45742

SANDER, H. H., Rapid annealing of frequency change in crystal resonators following pulsed x-irradiation, 26:45665 (SC-DC-721240)

SANDERS, William A., Simple perturbation and perturbation--variation treatments of the $1s_{1/2}$ g/ and $2p_{3/2}$ sub u/ states of H_2^+ , 26:46355

SANDERSON, T. R., Galactic gamma radiation between 200 MeV and 10 GeV, 26:46154

SANDMEIER, H. A., Electromagnetic pulse and time-dependent escape of neutrons and gamma rays from a nuclear

- explosion, 26:45497
- SANDUSKY, D. W., Fluence and temperature dependence of void formation in highly irradiated stainless steels, 26:46048 (CONF-710601-, pp 156-73)
- SANKAR DAS, M., Activation analysis with a Pu-Be neutron source, 26:45199
- SANO, Mituo, (N-Z) dependence of (n, γ) cross sections, 26:47085
- SANTAROSSA, G., DECAT: Computer code for analysis of long term pressure and temperature transient after loss of coolant accident in G.E. type pressure suppression containment, 26:47318 (RT/PROT-72-4)
- SANTHA, Andras, Iron metabolism in mice exposed to mixed radiations from a reactor, 26:45840
- SANTINI, R. E., Some comments on the signal-to-noise characteristics of real photomultiplier and photodiode detection systems, 26:45682
- SANWAL, D. N., Decay times, intensities, and their temperature variation of luminescence of aqueous and deuterated solutions of europic ion, 26:45295
- SAPERSTEIN, Alvin M., Numerical aspects of momentum-space scattering equations for peaked potentials, 26:46787
- SAPRE, A. V., Plasma confinement by external rf signal, 26:47114
- SARKAR, R., Prompt gamma decay of fission fragments, 26:47025
- SARKAR, S. K., Solar microwave bursts of different types in relation to optical flares, 26:46333
- SARKOZI, Lewis P., Production irradiator, 26:45447
- SARMA, L. V. K. V., Motion of a sphere in an electrically conducting rotating fluid, 26:46424
- SARMA, N., Charge form factor and quadrupole moment of ^6Li , 26:46878
- SASS, S. L., Structure and decomposition of Zr and Ti bcc solid solutions, 26:46036
- SATO, Humitaka, Hot Universe, cosmic rays of ultrahigh energy, and absolute reference system, 26:46257
- SATO, Tomohiko, Development of 50Hz 44 inch turbine blade, 26:45454
- SATO, Yuji, Fast-slow coincidence circuit systems for Ge(Li) detector, 26:45639
- SATOW, Yukio, Statistical analysis of radioinduced abnormal development, 26:45867
- SATZ, H., Limiting behavior of partially inclusive reactions, 26:46613
- SAULIT, V. R., Magnetic field inversely proportional to r with a periodic azimuthal inhomogeneity, 26:45638
- SAUNDERS, Neal T., Development of dispersion-strengthened Ni-Cr-ThO₂ alloys for the space-shuttle thermal protection system, 26:45934 (N-72-18585)
- SAUVAGE, G., e^+e^- production by e^+e^- annihilation in rho energy range with Orsay storage ring, 26:46428 (CONF-710816-14)
- $e^+e^- \rightarrow 0$ production by e^+e^- annihilation at w mass, 26:46427 (CONF-710816-13)
- SAVATTERI, C., Recuperative heat exchanger for a fast helium-cooled breeder reactor with helium turbine, 26:45439 (KFK-1566)
- SAVITCH, D., Influence of neutron irradiation on seed germination and plant growth of tomatoes, 26:45773
- SAVOSIN, S. I., Gold assay on geological samples by neutron activation with preliminary crucible fusion, 26:45200
- SAWYER, G. A., Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)
- SAXEBOEL, Gunnar, Electron spin resonance studies and INDO MO-calculations of electron irradiated peptides. A single crystal of N-acetylglycine, 26:45349
- SAXENA, K. M. S., Electric dipole polarizabilities of lanthanides and actinides, 26:46349
- SAXL, I., Mechanical characteristics of uranium alloys of type U-Mo(Fe, Al, Si) and U-Fe-Al-(Si, Cr), 26:45995 (UJV-2802-M)
- SAYERS, Robert P., Radioactive waste storage system and method, 26:46103
- SAZONOV, B. I., Circulation in the troposphere and the stratosphere and high-energy particles, 26:46297
- SAZONOV, M. I., Erosion of a tungsten cathode in nitrogen, 26:45968
- SCALAPINO, D. J., Thermodynamic properties of small superconducting particles, 26:46813
- Time-dependent superconductivity and quantum dissipation, 26:46821
- SCARF, F. L., Analysis of Pioneer 9 low-frequency wave observations near interplanetary discontinuities, 26:46302
- Detection of solar-wind electron plasma frequency fluctuations in an oblique nonlinear magnetohydrodynamic wave, 26:46329
- SCARR, J. Michael, Pion exchange in the reaction $\pi p \rightarrow \pi N$ at 6 GeV/c, 26:46481
- SCHAEFFLER, H., Computer-code for solution of three-dimensional-multigroup-equations, based on improved synthesis-method, 26:47594 (BMBW-FBK-72-6)
- SCHAEFER, H., Transport of radioactive materials by rail, 26:46117
- SCHLEENZ, R., Development of methods for application of neutron activation analysis to determination of trace elements in foods. Progress report, 10-12/71, 26:45189 (BFL-1/1972)
- SHELL, W. R., Natural ^{210}Po and ^{210}Pb in a marine environment, 26:45544 (RLO-2225-T-14-4)
- SCHELTEN, J., Local magnetic field distributions in superconducting Nb at 4.2 K, 26:46826
- SCHENZ, R. F. Jr., NERVA flight engine control system design, 26:47577
- SCHERZER, Walter O., Sample-conveying mechanism for scintillation counting, 26:46110
- SCHIFF, H., Dually charged particles and dual invariance, 26:46539
- SCHIFFER, John P., Coulomb energies and nuclear radii, 26:47097
- SCHIMMERLING, W., Acceleration of argon ions to 1.17×10^{10} electron volts, 26:46124
- SCHINDLER, S. M., Evidence for solar particle production above approximately 75 GeV, 26:46219
- SCHLATTER, R., Vacuum induction melting, 26:45981
- SCHLIEDER, S., Some remarks on the null plane development of a relativistic quantum field theory, 26:46716
- SCHLIER, R., Momentum transfer and plasma formation above a surface with high-power CO₂ laser, 26:47155
- SCHLITT, Leland Glenn, Effects of parallel wavelength on the collisional drift instability, 26:47168
- SCHMIDT, K. H., Application of perturbation theory for the calculation of molecular constants due to small mass changes and on the determination of force constants from very heavy isotope substitution, 26:45299
- SCHMIDT, P. J., Cosmic-ray electron spectrum and its modulation near solar maximum, 26:46325
- SCHMIDT-OTT, W.-D., Gamma branching and total conversion coefficients of intraband transitions in groundstate rotational band of ^{238}Cm , 26:47034
- SCHMITZ, A., Regeneration of lymphoid bone marrow in the thymectomized adult mouse subjected to a sublethal dose of x rays, 26:45830
- SCHNEEBERGER, Robert J., Television type nuclear radiation camera system, 26:45595
- SCHNEIDER, Rolf, New electrical procedure for the compaction of technical powders, 26:45951
- SCHNEIDER, Thomas Richard, Nuclear three-body problem and generalized spherical harmonics, 26:47103
- SCHNUR, P., Computer-code for solution of three-dimensional-multigroup-equations, based on improved synthesis-method, 26:47594 (BMBW-FBK-72-6)
- SCHOFIELD, Aldred E., All-metal discharge tube, 26:47178
- SCHOLL, R. E., Observations of chimney damage to residential buildings in Glendale, California: San Fernando earthquake, February 1971, 26:45474 (JAB-99-88)
- SCHOLZ, Kenneth L., Cyclotron production of ^{123}I : an evaluation of the nuclear reactions which produce this isotope, 26:45392 (N-72-15068)
- SCHONBERG, E., Inelastic e-p scattering and the polarization contribution to H hyperfine structure, 26:46523 (NYO-4076-17)
- SCHROEDER, G. D., Influences of thermal effluents upon aquatic production in Lake Ontario, 26:45578
- SCHUEREN, H., Scattered neutrons and their effects on calibration of Bonner sphere counters, 26:45599
- SCHULZE, H.-A., Artificial radioactivity of fallout and food in the northern districts of GDR in the year 1969, 26:45546
- SCHURINK, F., Fission recoil from surface of nuclear fuels by irradiation, 26:45939
- SCHUSTER, Jack S., Radioisotopes and turbine flow measurements, 26:45721
- SCHUTZ, Bernard F. Jr., Linear pulsations and stability of differentially rotating stellar models. I. Newtonian analysis, 26:46195
- Linear pulsations and stability of differentially rotating stellar models. II. General relativistic analysis, 26:46196
- SCHUTZ, Bernard Frederick Jr., Relativistic velocity-potential hydrodynamics and stellar stability, 26:46161
- SCHWANDT, P., Evidence for shape or quasimolecular resonances in the $^{16}\text{O} + ^{16}\text{O}$ interaction, 26:46884
- SCHWARTZ, P. R., Microwave celestial water-vapor sources, 26:46206
- SCHWARZ, Joseph, Formation of clouds in a cooling interstellar medium, 26:46245
- SCHWARZ, K. W., Surface-bound electrons on liquid helium: a negative result, 26:46798
- SCHWENDIMAN, L. C., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- SCOFIELD, James H., Theoretical radiative transition rates for K- and L-shell x rays, 26:46341 (UCRL-51231)
- SCOVILLE, N. Z., Kinematics of molecular clouds near the galactic center, 26:46251
- SCUKA, V., Model studies of density fluctuations of small and large ions in the atmosphere, 26:46288
- SEARS, Richard, Internal quantum numbers and homogeneous spaces, 26:46665 (CTH-ITP-71-14)
- SEAY, David G., Choleraic enteropathy as a cause of diarrhea and death in radiation enteritis and its prevention with cholestyramine, 26:45821
- SECHRIST, C. F. Jr., Investigation of the ionospheric D region at sunrise. II. Estimation of some photodetachment rates, 26:46280
- Investigation of the ionospheric D region at sunrise. III. Time variations of negative-ion and electron

- Investigation of the ionospheric D region at sunrise.
I. Time variations of ozone, metastable molecular oxygen, and atomic oxygen, 26:46279
- SEDA, Josef, Circuit for continuous acquisition of rate ratio of two pulse trains, 26:45675
- SEDLATSCHEK, K., Brittle-ductile transition of body-centered cubic metals with special consideration of tungsten and molybdenum. I, 26:45993 (ORNL-tr-2377)
- SEEBURGER, D. James, Radioactivity in Lake Michigan water and biota, vicinity of Big Rock Point Nuclear Reactor, 1960-69, 26:45535
- SEEGER, A., Application of high-voltage electron microscopy to the study of voids and other point-defect agglomerates, 26:46057 (CONF-710601-, pp 550-62)
- SEETHARAMAN, M., Poincare and TCP invariance in the determination of wave equation for particles of arbitrary spin, 26:46714
- SEETHARAMASWAMY, R., Motion of a sphere in an electrically conducting rotating fluid, 26:46424
- SEGAL, Y., Investigation of freon fire-extinguishing systems with a nucleonic gage, 26:45712
- SEGALL, Stephen Barrett, Interpretation of signals from a cylindrical Langmuir probe in a streaming laser-produced plasma, 26:47126
- SEGUN, J. G., Plasma-sampling diagnostics using ion kinetic energy distributions, 26:47128
- SEIDEN, Joseph, Energy flux between two superfluid helium baths separated by a thin layer, 26:46794
- SEIDMAN, D. N., Contrast patterns produced by self-interstitial atoms in field ion microscope images of a b.c.c. metal, 26:46029
- Void formation in quenched or irradiated metals, 26:46058 (CONF-710601-, pp 563-604)
- SEILER, E., Some remarks on the null plane development of a relativistic quantum field theory, 26:46716
- SEILER, R., Quantum theory of particles with spin zero and one half in external fields, 26:46741
- SEITZ, Martin George, Heavy-ion radiation studies in terrestrial and extraterrestrial materials, 26:46156
- SEKI, Ryoichi, K-nucleus scattering lengths, 26:46912
- SEKIYA, Tamotsu, Disappearance of neutron diffusion length of beryllium and graphite according to temperature, 26:46857
- SELBY, J. M., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- SELIVANOVA, M. N., Synthesis and study of lanthanide peroxides, 26:46085
- SELLE, James E., Radioisotopic fuel capsule, 26:45708
- SELUCKA, H., Strontium sorption on sorbents of BaSO₄--CaSO₄ system, 26:45273 (UJV-2820-Ch)
- SELVUTIN, V. P., Average-pulse-amplitude meter, 26:45685
- SEMENESCU, G., (α, p) reaction on even nickel isotopes, 26:46934
- SEMENKIN, V. A., Determination of background of Moessbauer transmission experiments using Fe alloys with 3d-transition elements, 26:47282
- SEMENTOV, B. Yu., Signal filtration in the amplitude spectrometry systems, 26:45637 (JINR-P13-6428)
- SEMENTOV, L. F., Effect of radioprotective preparations on the respiratory exchange of rats, 26:45823
- SEMERIA, R., Heat pipes, 26:45435 (CEA-CONF-2008)
- SEN, D. K., Weyl and Lyra manifolds, 26:46688
- SENE, M., Further results from multi-Regge models applied to $pp \rightarrow p\bar{p}\pi^+\pi^-$ reaction at 13.1 GeV/c, 26:46656
- SERDYUK, L. G., Lethal and mutagenic effect of ionizing radiations with different LET on extracellular bacteriophages, 26:45746
- SERGEEV, V. L., Analog models of marine nuclear steam generating plants, 26:47578
- SERVANT, J., Meridional transport of ⁹⁰Sr in the troposphere, 26:45516 (HASL-257, pp 1-2-17)
- SETH, V. P., Comment on the backward K-n $\rightarrow$ $\Lambda\pi^-$ scattering, 26:46648
- Recoil effect in quasi-elastic hadron--deuteron scattering, 26:46653
- SETTY, C. S. G. K., Small-scale fluctuations in F-region critical frequency, 26:46289
- SEXTRO, R. G., Beta-delayed proton decay of ¹³⁰I, 26:46879
- Confirmed proton radioactivity of ^{53m}Co, 26:46926
- Further results on proton radioactivity of ^{53m}Co, 26:46919
- SEYMOUR, Allyn H., Decline of ⁶⁵Zn in marine mussels following the shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)
- SHACTER, John, Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)
- SHADBOLT, Margaret Anne, Spectral studies of collagen, 26:47642 (UR-3490-47)
- SHAGIMARDANOV, N. Sh., Importance of tissue-equivalent detectors used in radiometry, 26:45619
- SHAH, K. Tahir, Conjugacy and stability theorems in connection with contraction of Lie algebras, 26:46666 (IC-72/49)
- SHAH, V. M., Simple transistorized control unit for neutron crystal spectrometers, 26:45636 (BARC-591)
- SHAIKH, Unis, Radiation safety around hot facilities at PINSTECH, 26:45499
- SHAKIN, C. M., Charge-dependent effects in the photodisintegration of ⁴He, 26:46866
- SHALAEV, A. M., Change in linear dimensions of metals under the influence of γ radiation, 26:46075
- Formation of defects in a crystalline lattice during irradiation of metals and alloys, 26:46076
- SHAPIRO, I. S., Many-neutron resonances, 26:46644
- Overlapping levels and giant resonances, 26:47098
- Quantum numbers and total and partial widths of the resonance N anti N(1970), 26:46458
- SHAPIRO, V. D., Non-linear theory of instability of a mono-energetic electron beam in plasma, 26:47151
- SHAPORENKO, A. A., Oscillations of charge-exchange cross section in system Li⁺⁺-He, 26:46368
- SHARAF, M. A., Transfer reactions to unbound states by the diffraction model, 26:47075
- SHARBER, J. R., Fermi acceleration of auroral particles, 26:46267
- SHARIPOV, Sh., Collective--single-particle states of deformed odd nuclei, 26:47081
- SHARMA, B. D., Chemical diffusion in the beta phase of zirconium--titanium system, 26:45985 (BARC-577)
- SHARMA, R. C., Alpha intensities to rotational bands of odd-A nuclei, 26:47079
- SHARMA, Subhash Chander, Track effects in physical, chemical, and biological systems, 26:45612
- SHARMA, V. D., Vorticity jump across a stationary magnetohydrodynamic discontinuity, 26:46421
- SHARP, V. L., Environmental benefit-cost analysis for nuclear power generation, 26:47386
- SHAW, D., Chemical effects of isomeric transition in bromoethane mixtures, 26:45360
- SHAW, P. F. D., Chemical effects of isomeric transition in bromoethane mixtures, 26:45360
- SHCHEMELEVA, G. G., Extractive photometric determination of U(VI) in mine waters, 26:45242
- SHCHERBANOVSKII, A., Predominant motion in spherical systems of galaxies, 26:46235
- SHEELEY, J. D., Autoclave fabrication techniques for flat cables, 26:47607 (BDX-613-579(Rev.))
- SHEFFIELD, John, Method for measuring the direction of the magnetic field in a Tokamak plasma by laser light scattering, 26:47120
- SHELEGEDIN, V. N., Lethal and mutagenic effect of ionizing radiations with different LET on extracellular bacteriophages, 26:45746
- SHELEKHIN, Yu. L., Interaction of a ruby with ionizing radiation, 26:47244
- SHELEST, V. P., Duality, scaling, and fixed poles in virtual Compton scattering, 26:46559
- SHELINE, Raymond K., Low-lying states and configurations in ¹³⁸La, 26:46964
- SHEN, Shu-wei, Study of flux-flow noise in Type-II superconductors, 26:46815
- SHEPTII, V. I., Feynman-Wilson-gas model as consequence of jet approach to inelastic hadron processes, 26:46586 (JINR-P2-6501)
- SHER, R., Operating characteristics of neutron well coincidence counters, 26:45587 (BNL-50332)
- SHERETOV, E. P., Shape of the mass peak in a three-dimensional quadrupole mass spectrometer, 26:45676
- SHERR, Theodore S., Attribute sampling inspection procedure based on the hypergeometric distribution, 26:46094 (WASH-1210)
- SHESTAKOV, G. N., $\rho\omega$ interference in reactions $\gamma N \rightarrow \pi^+\pi^-\Delta$, 26:46527
- SHEVCHENKO, L. P., Thermonuclear combustion waves in a bounded plasma, 26:47172
- SHEVCHENKO, V. G., New experimental data on the fine structure of dipole resonance, 26:46890
- SHEVCHENKO, V. I., Non-linear theory of instability of a mono-energetic electron beam in plasma, 26:47151
- SHEVCHENKO, Yu. A., Determination of background of Moessbauer transmission experiments using Fe alloys with 3d-transition elements, 26:47282
- SHEVCHUK, A. I., Accumulation and distribution of radioactive elements in natural phytocenoses of the Carpathians, 26:45551
- SHIBATA, Fumiaki, Localized magnetic states in superconductor, 26:46809
- SHIDA, Y., Neutron deficient even-even isotopes of Kr, Se, Ge, and pseudo SU₃ coupling, 26:46931
- SHIFRINE, M., Radiation-induced transplantable myelomonocytic leukemia in beagles: hematologic parameters, 26:45897
- SHIGEMATSU, Tatsuhiko, Solvent extraction of Eu by Versatic Acid 911, 26:45277
- SHIKIN, I. S., Gravitational fields with groups of motions on two-dimensional transitivity hypersurfaces in a model with matter and a magnetic field, 26:46755
- SHIMIZU, S., Possible evidence for a contribution from direct collisions in K-shell internal ionization accompanying β decay, 26:46962
- SHIMOTAKE, Hiroshi, Electrochemical power-producing cell, 26:45434
- SHINDO, Shuichi, Radiation-induced graft copolymerization of mixtures of styrene and n-butyl acrylate onto cellulose and cellulose triacetate, 26:45320
- SHINAKA, Kyoji, Theoretical consideration of electron-scavenging process in liquid hydrocarbons. III, 26:45333
- SHIROKOV, M. I., Quantum field theory: "dressing" contra divergencies, 26:46701 (JINR-P2-6454)

- SHISHKANOV, N. G., Dose to the epithelium of the upper respiratory tract from radon and its daughter products, 26:45861
- SHISHOV, V. A., Improvement of electron beam shaper in BSM-25 model betatron, 26:46132
- SHKLYAREVSKII, V. M., Introduction of correction for nonlinearity of direct-current Q-meter, 26:45680
- SHMARIOVICH, E. M., Problem of epigenetic zonality in uranium ore deposits of carbonate rocks, 26:45506
- SHMATKO, G. A., Disintegration of supersaturated solid solutions according to a heterogeneous mechanism, 26:46032
- SHOCHET, M. J., Experimental study of the decays $K/\text{sub } S \text{ L} / + \gamma$, 26:46434
- SHOOP, C. Robert, Effects of ionizing radiation on salamander orientation. Progress report, 26:45815 (COO-3486-1)
- SHRAMENKO, B. I., Photoproduction of charged pions by nuclei at $E_0 = 1160$ MeV, 26:46441
- SHTAN, A. S., Radiation technology. VI, 26:45432
- SHUBNYI, Yu. K., Lifetime of 1.013-MeV state in ^{27}Al , 26:46905
- SHUI, Ven H., Monte Carlo trajectory calculations of the three-body recombination and dissociation of diatomic molecules, 26:46382
- SHUKLA, P. K., Reflection of microwave through laboratory plasma, 26:47118
- Total energy loss of a test charge in a collisional magnetoplasma, 26:47149
- SHULMAN, A. R., Temperature dependence of the work function of Be-W₁₁₀ system, 26:47257
- SHUNK, Francis A., Constitution of binary alloys, second supplement, 26:45997
- SHURAVIN, V. M., Time measurements with a germanium detector, 26:45616
- SHYN, T. W., Angular distribution of electrons elastically scattered from N₂, 26:46392
- SHAKOLAH, Mohammed Ali, Morphology and mechanical properties of cellulose acetate grafted with polyacrylonitrile, 26:46084
- SIAUD, J., K₂⁰ decay in a Wilson chamber, 26:46449 (SLAC-Trans-143)
- SIDDAPPA, K., Effective atomic numbers for differential processes of gamma-ray interactions, 26:47207
- SIDOROV, A. A., Photoproduction of hypernuclei and current algebra, 26:46545
- SIDOROV, V. A., phi-meson mass determination, 26:46443
- SIDYAKINA, T. M., Removable current lead for power supply in superconducting magnetic systems, 26:45679
- SIEKIERSKI, S., Existence of regularities, tetrad, or double-double effect in changes in unit cell volumes of compounds of actinides in different oxidation states, 26:45375
- SIEMON, R. E., Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)
- Plasma experiments on $l = 1, 0$ helical equilibria in the Scyllac 5-meter, theta-pinch toroidal sector, 26:47156 (LA-DC-72-706)
- SIEMSEN, R. H., Evidence for shape or quasimolecular resonances in the $^{140} + ^{140}$ interaction, 26:46884
- SILENKO, P. N., Gravitational energy emitted by isolated material source. I, 26:46760
- SILK, Joseph, Bright nebulae near concentrations of high-velocity gas, 26:46250
- SILKER, W. B., Oceanic distributions and relationships of ^7Be and fission products, 26:45530 (BNWL-SA-4205(Rev.))
- SILL, W. R., Interaction of the solar wind with a planetary ionosphere, 26:46324 (N-72-18796)
- SILUK, Richard S., Bright nebulae near concentrations of high-velocity gas, 26:46250
- SILVA, Sandra, Determination of silica in silicates containing P, Ti, and Zr, 26:45255
- SILVEIRA, Carmem L. P., Determination of silica in silicates containing P, Ti, and Zr, 26:45255
- SILVER, E. G., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)
- SILVERMAN, Sol Jr., Hard and soft tissue necroses following radiation therapy for oral cancer, 26:45801
- SIMIC, M., Electron and hydrogen atom attachment to aromatic carbonyl compounds in aqueous solution. Absorption spectra and dissociation constants of ketyl radicals, 26:45354
- SIMMONS, Michael Fred, X-ray and neutron diffraction study of the holmium-iron system, 26:46016
- SIMON, Barry, $(\text{phi}/\text{sup } 2n)/_2$ field Hamiltonian for complex coupling constant, 26:46718
- SIMON, Joseph, Whole-body gamma irradiation of newborn pigs: hematologic changes, 26:45852
- Whole-body gamma irradiation of newborn pigs: pathologic changes, 26:45853
- SIMON, Manfred, Radiation hazards to man in space flights, 26:45812
- SIMON, Sandor, Studies of the origin of the exogenous oxide inclusions in killed and unkilld cast steels through subsequently activated tracer elements, 26:46008
- SIMONOV, Yu. A., Hyperspherical harmonics method in the theory of nuclear bound states, 26:47090
- SIMPSON, F. B., Total neutron cross section of ^{243}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- SIMPSON, O. D., Total neutron cross section of ^{243}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- SINGH, H. P., Thermodynamic properties of the phase In_2Bi_3 , 26:46026
- SINGH, I. D., Isotope effect on Franck-Condon factors and r-centroids for the new $(^{15}\text{I}/\text{sub } u/^{+} - ^{15}\text{I}/\text{sub } g/^{+})$ system of isotopic C₂ molecule, 26:45293
- SINGH, K. P., Reflection of microwave through laboratory plasma, 26:47118
- SINGH, P. P., Evidence for shape or quasimolecular resonances in the $^{140} + ^{140}$ interaction, 26:46884
- SINGH, R. N., Total energy loss of a test charge in a collisional magnetoplasma, 26:47149
- SINGH, T., Cylindrical wave solutions of Bonnor's and Schroedinger's unified field equations, 26:46695
- SINGH, Virender, Simple transistorised control unit for neutron crystal spectrometers, 26:45636 (BARC-591)
- SINHA, B. C., Isospin dependence of optical model potentials, 26:47059
- SINHA, Om P., Isotope separation rate in Haefner effect, 26:45694
- SINK, D. A., Evidence for shape or quasimolecular resonances in the $^{140} + ^{140}$ interaction, 26:46884
- SIPLER, Dwight Passmore, Optical detection of radio-frequency heating of electrons in the F region, 26:46292
- SIROTA, N. N., Radiation physics of crystals and p-n junctions, 26:47214
- SIROTKIN, V. K., Sub-barrier (p,n) reaction in region of single-particle resonance, 26:46952
- SISAKYAN, A. N., Operator method of solving the quasipotential equations and a conception of straight-line paths at high energies, 26:46700 (JINR-P2-6437)
- SITENKO, M. L., Control system for beam formation after single deflection at exit of 300 MeV electron linear accelerator, 26:46131
- SIUDA, Jerzy, Distribution of doses in the lesser pelvis in the treatment of carcinoma of the cervix uteri, 26:45784 (TT-54002/4, pp 456-61)
- SIVAN, J. P., Two new large-diameter galactic regions of faint H α emission, 26:46236
- SIX, J., Investigation of low-q² inelastic neutrino events, 26:46450
- SJOEDEN, A.-M., Effect of irradiation on thyroid growth in mouse foetuses and goitrogen challenged adult mice, 26:45865
- SKAANBERG, B., Validity of rotor model in odd-Z nuclide ^{171}Lu , 26:46995
- SKAR, K., Energy levels in ^{231}Th and ^{233}Th , 26:47029
- SKARSGARD, H. M., Electron temperature determination from the rate of ionization, 26:47122
- SKOLYSZEWSKI, Jan, Distribution of doses in patients with Hodgkin's disease during conventional radiotherapy, 26:45786 (TT-54002/5, pp 590-3)
- SKOPIN, V. S., Determination of the energy of dislocation of an impurity atom by an internal friction method, 26:46023
- SKORODUMOV, B. G., Elastic scattering of protons by deuterons at energies of 14, 16, and 17.6 MeV, 26:46863
- SKUJA, Andris, Study of dipion production in the reaction $\pi^+ p \rightarrow \pi^0 \pi^0 n$ between 1.6 and 2.4 GeV/c, 26:46477 (LBL-378)
- SLABOSPITSKII, R. P., Elastic scattering of polarized protons by ^{28}Si and ^{40}Ca , 26:46924
- SLATER, Richard C., Scattering of CsF by electrons, 26:46406
- SLAUGHTER, G. G., Total neutron cross section of ^{243}Am from 0.5 to 1000 eV, 26:47010 (ANCR-1060)
- SLEEPER, Arthur, Nonlinear saturation of Type-I irregularities in the equatorial electrojet, 26:46277
- SLEPCHENKO, I. G., Electrochemical reduction of U(VI) to U(IV) in sulfuric acid solutions, 26:45978
- SLIV, L. A., Parameters of effective particle interaction in nucleus, 26:47094
- SLOMSKA, Janina, Prognostic elements in inoperable carcinoma of the breast, 26:45783 (TT-54002/4, pp 451-5)
- SLOVINSKII, B., Investigation of generation of particles and resonance states in interactions of 2.34 GeV/c, 5 GeV/c, and 9 GeV/c π -mesons with xenon nuclei, 26:46476 (JINR-PI-6408)
- SMALL, H., Effect of a polythiol synthetic resin on the absorption and excretion of methylmercury in experimental animals, 26:47643 (UR-3490-80)
- SMART, B. C., Zirconium tetrafluoride process, 26:45274
- SMEDLEY, P. N., Tubular Nb/Cu conductors for ac superconductive power transmission, 26:46844
- SMIDT, F. A. Jr., Production of vacancy condensates by laser bombardment, 26:46059 (CONF-710601-, pp 613-29)
- SMIRIN, L. N., Lifetime of 1.013-MeV state in ^{27}Al , 26:46905
- SMIRNOVA, I. B., Role of thiols in radiosensitivity of sea urchin eggs, 26:45882
- SMITH, C. H. Llewellyn, Neutrino reactions at accelerator energies, 26:46584
- SMITH, Cody, Numerical evaluation of Chapman's grazing incidence integral ch (X, chi), 26:46273
- SMITH, D., Technique for recording Langmuir probe characteristics in afterglow plasmas, 26:47119
- SMITH, David E., Pull-through operation for radiation-induced rectovaginal fistula. Report of a case, 26:45800

SMITH, Frank A., Biological properties of certain compounds containing the C—F bond, 26:47637 (UR-49-1508)
 Procedure for chronic intravenous catheterization in the rabbit, 26:47641 (UR-3490-21)

SMITH, Harding E., Radial cyanogen-band gradients near the nuclei of galaxies, 26:46243

SMITH, I. W. M., Quenching of infrared chemiluminescence: rates of energy transfer from HF(ν less than or equal to 3) to CO₂ and HF, 26:45298

SMITH, J. Marshall, Special biophysics projects, 26:45893 (ORO-2690-12(Pt.B), Paper 13)

SMITH, James C., Studies in the uses of ionizing radiations as stimuli, 26:45817 (ORO-2690-12(Pt.B), Paper 12)

SMITH, Jay Edward, Neutronics calculations for toroidal thermonuclear reactor blankets containing iron, 26:47176

SMITH, Jerry C., Computer simulation modeling of an automated warehouse system, 26:47604 (BDX-613-456)

SMITH, M. E., Remotized apparatus for determining oxygen in irradiated reactor fuels and cladding materials, 26:45224 (LA-DC-72-524)

SMITH, Maynard E., Isotope ratio tracer method: applications in atmospheric sulfur pollution studies, 26:45568

SMITH, Myron A., Metallicism in border regions of the Am domain. I. Extremely young Am stars in the Orion 1c association, 26:46181

SMITH, T. I., Dynamic structure of vortices in superconductors. III. Graphs for B approximately = H/sub c2/, 26:46814

SMITH, V. H., Effect of mass on promethium retention and removal from the rat, 26:45922

SMITH III, F. L., Numerical evaluation of Chapman's grazing incidence integral ch(X, chi), 26:46273

SMONDYREV, M. A., Operator method of solving the quasipotential equations and a conception of straight-line paths at high energies, 26:46700 (JINR-P2-6437)

SMONIEWSKI, J., Sensitometric measurements of non-screen industrial films exposed by x or gamma radiation, 26:45468

SNYDER, A. L., Observations of the auroral oval by the Alaskan meridian chain of stations, 26:46268

SOBELS, F. H., New method for quantitative study of induced autosomal non-disjunction in Drosophila melanogaster, 26:45880

SODD, Vincent J., Cyclotron production of ¹²³I: an evaluation of the nuclear reactions which produce this isotope, 26:45392 (N-72-15068)

SOEDERBERG, G., Stenosis of the small bowel as a complication in radiation therapy of carcinoma of the uterine cervix, 26:45802

SOEREN MATSSON, L. J., ²²Na in the food-chain: lichen--reindeer--man, 26:45909

SOERENSEN, Bent, Spectroscopy with two-nucleon transfer reactions in deformed nuclei. II, 26:47062
 Spectroscopy with two-nucleon transfer reactions in deformed nuclei. I, 26:47061

SOFIA, Sabatino, Extragalactic origin of the transient x-ray sources, 26:46215

SOGA, T., Infrared absorption spectra of potassium uranium fluorides, 26:45230

SOKOL, Ludvik, Sintered Mg composites, cladding material of U fuel elements cooled by carbon dioxide. II, 26:46012

SOKOLOV, A. A., Particle acceleration in a H₁₁₁ cylindrical cavity in an axial magnetic field, 26:46139
 Spin correlation during bremsstrahlung radiation of longitudinally polarized electrons, 26:47052
 Toward the theory of nuclear shells, 26:47040 (BNL-tr-489)

SOLDATOV, A. N., Manifestation of structure in electron energy distribution functions and determination of the effective cross sections of inelastic processes, 26:46407

SOLHEIM, P. Oe., Radiation injury of the spinal cord, 26:45797

SOLINGER, Alan B., Relationship between x-ray luminosity and velocity dispersion in clusters of galaxies, 26:46249

SOLODOVNIKOV, A. P., Dielectric strength of insulators in a vacuum, 26:46130

SOLODOVNIKOVA, E. P., N. N. Bogolyubov's transformation in strong coupling theory. II, 26:46691

SOLOSKY, L. F., Accuracy of quantitative electron probe microanalysis with energy dispersive spectrometers, 26:45237

SOLOV'EV, L. D., Estimation of intermediate energy contribution to the dispersion sum rules and rotation of polarization in pion--nucleon scattering at high energies, 26:46617 (IFVE-STF-71-15)

SOLOV'EV, S. M., X ray fluorescence analysis of heavy element by an α -X coincidence method, 26:45270

SOLTAN, A. Jr., Role of the exit states of a compound nucleus in the emission of high-energy γ -rays, 26:47082

SOMA, Y., γ radiolysis of gaseous carbon dioxide at high densities, 26:45346

SOMON, Roland, Equipment for concentrating radioactive solutions, 26:46111

SOOD, R. K., Galactic gamma radiation between 200 MeV and 10 GeV, 26:46154

SORTIOU, M., New method for potassium determination by neutron activation in mineral and rock samples,

26:45197

SOROKIN, O. V., Electrical conductivity, thermo-emf, and thermionic emission of BaO and SrO crystals, 26:47249

SOSNIVKER, W. A., Photoproduction of vector and axial vector resonances in broken nonlinear chiral SU₃ x SU₃ dynamics, 26:46563

SOUDAIN, G., New reading system for an ionographic detector, 26:45598

SOUFFRIN, S., Continuum of nuclei of galaxies. II, 26:46239

SOURIAU, Jean-Marie, Model for particles with spin in the gravitational and electromagnetic field, 26:46767

SPAAS, H. A. C. M., Stress distribution in nozzle-to-cylinder connections for small diameter ratios, 26:45445

SPARKS, Warren M., Studies of hydrodynamic events in stellar evolution. II. Dynamic instabilities in stellar envelopes, 26:46176

SPENCER, J. P., Dynamical theory for the contrast of perfect and imperfect crystals in the scanning electron microscope using backscattered electrons, 26:47267

SPINRAD, Hyron, Radial cyanogen-band gradients near the nuclei of galaxies, 26:46243

SPITSYN, Vikt. I., Radiation damage to polymer fiber investigated by small-angle x-ray scattering, 26:45338

SPITSYN, Viktor Ivanovich, Method of modifying polymeric materials, wood, and articles made therefrom by radiation-initiated graft copolymerization thereof, 26:45355

SPRAY, P. H. G., Bruce auxiliary steam supply, 26:45695

SPRY, David Jon, Nuclear spins as determined by the method of gamma--gamma angular correlations, 26:46901

SRINIVASAN, Nemell Krishnaswami, Anelastic relaxation caused by point defects in MnF₂, 26:47615 (CU-3194-22)

SRIVASTAVA, B. N., Pressure dependence of Moessbauer recoilless fraction, 26:47276

STABROVSKII, A. I., Electrochemical reduction of U(VI) to U(IV) in sulfuric acid solutions, 26:45978

STAMP, A. P., Calculation of the nuclear quadrupole moment, 26:47057

STANISIC, Milomir M., Weak fluctuation modes of conducting viscous compressible flow, 26:46420

STANKE, F., Moessbauer effect studies of ⁵⁷Co and ⁵⁷Fe in β -rhombohedral B, 26:47280

STANKUNAS, Alexander Robert, I. Radiation chemistry of n-butylphenone in benzene solution. II. Radiation-induced alkylation of caffeine by amines, 26:45326

STANSFIELD, O. M., Irradiation-induced dimensional change in HTR control materials, 26:46077 (GULF-GA-A-12035)

STARODUBOV, I. P., Electrical and thermal conductivities of Re, Os, and Ir and some of their alloys with Th and La in wide temperature range, 26:47248

STARODUBTSEV, S. V., Effect of γ radiation on oscillation frequency of AT-cut piezoquartz plates, 26:45667 (SC-T-722519)

STEBBINGS, R. F., Chemi-ionization in collisions between helium metastable atoms and argon, 26:46404

STEIN, Robert F., Formation of clouds in a cooling interstellar medium, 26:46245

STEINBERG, Meyer, Isotope ratio tracer method: applications in atmospheric sulfur pollution studies, 26:45568

STEINDLER, M. J., Containment of fission product iodine in the reprocessing of LMFBR fuels by pyrochemical reactions, 26:45421

STEINFELD, Jeffrey I., Optical model for vibrational relaxation in reactive systems, 26:46375

STEINGART, M., Quantized vortex ring dynamics and the vortex core radius in He II, 26:46800

STEINNES, E., Determination of Mn in high-purity Fe by neutron activation analysis after pre-irradiation separation, 26:45216

STELLA, R., Radioanalytical investigation on the extraction behavior of rhodium in thiocyanate-TBP system, 26:45194

STELSON, P. H., Coulomb excitation of ¹¹⁷ ¹¹⁹Sn, 26:46961

STEPANENKO, Yu. V., Extractive photometric determination of U(VI) in mine waters, 26:45242

STEPANOV, K. N., Stability of a finite-pressure plasma with an anisotropic electron-velocity distribution, 26:47159 (UCRL-Trans-10630)

STEPIC, R., Effective g/sub s/ factor for odd proton ¹⁵³Sm nucleus, 26:46994

STEPLEWSKI, Z., Effect of x irradiation on cells, 26:45738

STERLING, D. L., Synthesis of Jicamarca data during the great storm of March 8, 1970, 26:46282

STERN, David, Charged-particle motion in the vicinity of a neutral plane, 26:46522 (N-72-17763)

STERN, Richard C., Scattering of CsF by electrons, 26:46406

STEUNENBERG, Robert K., Electrochemical power-producing cell, 26:45434

STEVENS, D. J., Assessment and control of environmental contamination, 26:45554

STEWART, A. M., Paramagnetic susceptibilities of metallic samarium compounds, 26:47259

STEYERL, A., Interference in the ultracold neutron transmission through thin layers, 26:46853

STICH, H. F., Various levels of DNA repair synthesis in xeroderma pigmentosum cells exposed to carcinogens N-

- hydroxy and N-acetoxy-2-acetyl-aminofluorene, 26:45742
- STIEDE, Wayne L., Experience in the licensing of nuclear power plants, 26:47573
- STIEGLER, J. O., Postirradiation mechanical properties of Types 304 and 304 + 0.15 percent Ti stainless steel, 26:46068 (ORNL-TM-3733)
- STINGL, M., Vertex parameters and coupling constants for the virtual decay of light nuclei, 26:47063
- STIX, Thomas H., Controlled-fusion reactor, 26:47174
- Heating of toroidal plasmas by neutral injection, 26:47108
- STOCKER, W., Single-particle spin-orbit potential following from the nucleon-nucleon spin-orbit interaction, 26:47060
- STOCKTON, A. N., Ton 155 and 156: a double quasar, 26:46210
- STOKER, P. H., Rigidity dependence of cosmic-ray modulation at rigidities > 2 GV in August 1969, 26:46327
- STOLARSKI, N. S., Angular distribution of electrons elastically scattered from N_2 , 26:46392
- STOLER, P., MeV total neutron cross sections of ^{238}U , 26:47013 (COO-3058-23)
- STOLL, H., Optical waveguiding in proton-implanted GaAs, 26:47232
- STONE, E. C., High-energy electron spikes at high latitudes, 26:46307
- STONEHAM, A. M., Observation and theory of the void lattice in molybdenum, 26:46055 (CONF-710601-, pp 522-32)
- STORER, R. G., Improvements on a new method for the numerical solution of the Schrodinger equation, 26:47290
- STOTHERS, Richard, Red supergiants and neutrino emission. II, 26:46177
- STRACK, William C., Effect of two types of helium circulators on the performance of a subsonic nuclear-powered airplane, 26:45425 (N-72-16724)
- STRAZHEKSKO, D. N., Radiometric investigation of the absorption of potential-determining ions from aqueous solutions of non-ionic surface-active substances, 26:45279
- STREATER, R. F., Connection between spectrum condition and Lorentz invariance of $P(\phi)_2$, 26:46707
- STREB, Alan J., Thermal control and power flattening for radioisotopic thermodynamic power system, 26:45427
- STRICKER, A., $^2H(n,p)^2n$ reaction at 50 MeV and neutron-neutron scattering length, 26:46861
- STRILCHUK, A. R., Electron temperature determination from the rate of ionization, 26:47122
- STRINDEHAG, O. M., Apparatus for measuring D_2O content in a flow of water, 26:45267
- STRINGHAM, W. S., Carbon-13: a selected abstract bibliography, 26:45286 (MLN-1913)
- STRITTMATTER, P. A., Physical associations between quasi-stellar objects and galaxies, 26:46258
- STROM, R. G., Polarization of radio sources at $\lambda 49$ CM and $\lambda 73$ CM, 26:46204
- STRONGIN, M., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)
- STRUBLER, K., Film technique for determining the absorbed dose at exit from the patient during radiation exposure, 26:45631
- STRUGAL'SKII, Z. S., Investigation of generation of particles and resonance states in interactions of 2.34 GeV/c, 5 GeV/c, and 9 GeV/c π -mesons with xenon nuclei, 26:46476 (JINR-P1-6408)
- STRUMINSKII, B. V., On the distribution of resonance spins in the dual resonance model, 26:46609 (ITF-72-13-P)
- STUMPF, H., Functional definition and calculation of global observables in nonlinear spinor field quantum theory, 26:46746
- STURM, Bernard J., Radiation-resistant plastic insulators, 26:45672
- STURM, R., Gamma gamma directional correlation measurements in ^{131}Cs , 26:46963
- STURROCK, David, Neuropathological evaluation of monkeys exposed to body-alone x-radiation, 26:45850
- STURROCK, M. G., Effects of x-irradiation in vitro on the transport of potassium ions in thymocytes, 26:45744
- SUBBARAO, E. C., Kinetics and burst phenomenon in ZrO_2 transformation, 26:45947
- SUDAKOV, V. A., Thermodynamic properties of solutions in system In-Sn-Hg, 26:46004
- SUDBERY, A., Non-relativistic de Sitter space-time and harmonic oscillator, 26:46670
- SUDOVTSOV, A. I., Variations in the critical temperature of superconducting compounds Nb_3Al and $Nb_3Al/sub x/Ge/sub 1-x/$, 26:46021
- SUFFOLK, Graham, Rotational velocities of Ap stars, 26:46182
- SUGANO, Reiji, Variational principle for nonlinear Lagrangian in quantum mechanics, 26:47298
- SUGLOBOVA, I. G., Chemical shifts in $L/sub \alpha 1/$ x-ray line of some U compounds, 26:45377
- SUH, Kyung Soo, Activated sintering mechanism of UO_2 powder, 26:45956
- SULIVAN, James D., Composition and energy spectra of heavy nuclei with $0.5 < E < 40$ MeV per nucleon in the January 24, and September 1, 1971 solar flares, 26:46232
- SUMP, Kenneth R., Process of forming an isotopic heat source, 26:45707
- SUNAYASHIKI, Tadashi, Dosimetry of pulsed x-rays and electrons from a linear accelerator, 26:45627
- SURKOVA, I. V., Excitation cross section of first $2+$ levels of Ni and Zn nuclei, 26:46936
- SUROVTSOV, Yu. S., Quasi-threshold behavior of $\pi + N + N + (\gamma\text{-virtual})$ process amplitude, 26:46620 (JINR-P4-6475)
- SUSKOV, I. I., Chromosome aberrations in human bone-marrow cells after a single local gamma-irradiation. I, 26:45794
- SUTCLIFFE, H., Comparison of the radiolysis of liquid bromotrifluoromethane with the radiolysis of liquid trifluoriodomethane, 26:45322
- SUTHERLAND, Peter Gordon, White dwarfs and neutron stars, 26:46170
- SUURA, H., Testing triplet models, 26:46516 (DESY-72/21)
- SUZUKI, Takashi, Determination of small amounts of molybdenum based on its catalytic effect in Methylene Blue-hydrazine sulfate reaction, 26:45227
- SUZUKI, Tsuneo, Relativistically covariant formulation in nonlinear Lagrangian theories and factor-ordering problems, 26:46744
- SVECHNIKOV, V. N., Alloys of chromium with terbium, 26:46033
- SVENSSON, H., Commercial thimble chambers for absorbed dose measurements at high energy electron radiation, 26:45623
- SWAROOP, Janardhan, Versatile heavy water, 26:45698
- SWARTZ, Harold M., Radiation effects on plasma electron-spin-resonance (ESR) spectra of cancer patients, 26:45723
- SWENSON, G. W., VRS for VRO, 26:46209
- SWIATNICKA, Jolanta, Selected prognostic elements in carcinoma of the larynx, 26:45781 (TT-54002/4, pp 437-41)
- SWIFT, Robert L., Bacteriophage T4 inhibits colicin E2-induced degradation of Escherichia coli deoxyribonucleic acid, 26:47639 (UR-3055-2)
- SWINTH, K. L., ^{252}Cf neutron radiography at Battelle-Northwest, 26:45467 (BNWL-SA-4320)
- Photon intensities and their importance in counting transuranic materials, 26:47011 (BNWL-1648)
- SYCHEV, B. S., Intracuclear cascade in concrete shield of a 70-GeV proton synchrotron, 26:46126 (ORNL-tr-2617)
- SYKORA, V., Selective ion exchangers with isonitrosoacetamide functional groups. II, 26:45283
- SYMANKIK, K., Currents, stress tensor, and generalized unitarity in conformal invariant quantum field theory, 26:46697 (DESY-72/20)
- SZARKA, Gyula, Studies of the origin of the exogenous oxide inclusions in killed and unkilld cast steels through subsequently activated tracer elements, 26:46008
- SZEFLINSKI, Z., Role of the exit states of a compound nucleus in the emission of high-energy γ -rays, 26:47082
- SZTANYIK, Laszlo, Iron metabolism in mice exposed to mixed radiations from a reactor, 26:45840
- SZULC, M., Content of Cs in muscles of broilers, hens, and partridges, 26:45548
- Passage of ^{90}Sr from bovine bones to received food products, 26:45549
- SZYMCZYK, W., Thermoluminescent dosimetry system for determining quality, 26:45622
- SZYMCZYK, Wieslawa, Distribution of doses in the lesser pelvis in the treatment of carcinoma of the cervix uteri, 26:45784 (TT-54002/4, pp 456-61)
- SZYSZKO, E., Changes of Dextran irradiated with γ radiation, 26:45730

T

- TABATA, Tatsuo, Effect of temperature on various types of the ionization chamber, 26:45594
- TABET, E., Surface excitations of Type-I superconductors in a static magnetic field, 26:46830
- TABET, Eugenio, Criteria for selecting the location of nuclear plants, 26:47574
- TABUTEAU, A., Phases of scheelite structure in the systems of plutonium molybdate-lithium or sodium molybdate, 26:45954
- TACCONI, F. A., Threshold of parallel channel flow instability, 26:45459 (CISE-R-275)
- TAGUCHI, Yukio, Transformation properties of operators in local field theory, 26:46745
- TAHERZADEH, Mojtaba, Neutron radiation characteristics of plutonium dioxide fuel, 26:45704
- TAHIRA, Sadamu, Energy and angular distributions of secondary electrons resulting from ionizing collisions of electrons with helium and krypton, 26:46367
- TAKADA, Jitsuya, Direct measurement of γ -ray energies and half-life of ^{14}Cs , 26:46971
- TAKAHASHI, Akira, Study on RBE of various types of radiation, 26:45864
- TAKAHASHI, Hisao, Observation of the intensity ratio between (5,1) and (9,4) bands of OH emission in the night airglow, 26:46286
- TAKAMORI, Yasuhiko, Culture of murine leukemic cells induced by x-irradiation, 26:45855
- TAKANAKA, Kenji, Magnetic properties of a superconductor with hexagonal symmetry, 26:46807
- TAKEDA, Ryuichi, Permeability of lingual mucosa in radiation glossitis, 26:45862

TAKEOKA, Seiji, Dosimetry of pulsed x-rays and electrons from a linear accelerator, 26:45627

TAKESHITA, Kenji, Acute death of mice during massive gamma irradiation. III, 26:45846

Death of mice irradiated with pulsed x-rays, 26:45866
Dosimetry of pulsed x-rays and electrons from a linear accelerator, 26:45627

TALANIN, Yu. N., Elastic scattering of protons by deuterons at energies of 14, 16, and 17.6 MeV, 26:46863

TALPOV, G. Sh., Al-Fe-Mo oxide catalysts. I, 26:45963
TAMAI, Tadaharu, Direct measurement of γ -ray energies and half-life of ^{144}Cs , 26:46971

TAMARU, Kenzi, Identification of reaction intermediates by microwave spectroscopy. The catalytic reactions between propylene and deuterium over zinc oxide, 26:45303

TAMIYA, Tadashi, Convergent hole collimator for transmission scintiphoto, 26:45602

TANABE, Teruo, Solvent extraction of Eu by Versatic Acid 911, 26:45277

TANAKA, Masao, Polarographic determination of Sn in Cu-, Al-, and Zr-base alloys, 26:45256

TANAKA, Masaru, Use of irradiation in plastic wastes disposal, 26:46090

TANAKA, Syu, High energy production processes in the relativistic eikonal approximation, 26:46791

TANNER, N. W., Test of time reversal invariance through detailed balance of the reaction $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$, 26:46881

TARASENKO, N. D., Radiation mutagenesis of potato, 26:45770

TARASOV, A. V., Electromagnetic effects in nucleon polarization at meson-nucleon scattering, 26:46520 (JINR-P2-6438)

TARNAUCEANU, D., Statistical treatment of relationship between solar flare-producing regions and SC + SI events, 26:46332

TARNOVSKII, A. S., Lorentz transformations for electromagnetic field vectors and the motion of charges in constant and homogeneous electromagnetic fields, 26:46679

TARNSTROM, Guy LaVern, Solar radio spike bursts, 26:46217

TARRAGO, X., Heavy ion nuclear reactions, 26:47056

TARSKI, J., Method of characteristics for functional-derivative equations, 26:46728

TARUISHI, Koji, (44) and (60) + (06) interactions in the s-d shell, 26:47084

TARUTIN, I. G., Investigation of positron spectra obtained on linear 8 to 30 MeV electronic accelerator, 26:46149

TATI, Takao, Hot Universe, cosmic rays of ultrahigh energy, and absolute reference system, 26:46257

TAUB, A. H., Variational principles and spatially-homogeneous universes, including rotation, 26:46758

TAUSSIG, R. T., Research in computational plasma physics. Progress report, March 1, 1971--March 1, 1972, 26:47145 (COO-3086-15)

TAVARES, O. A. P., Some remarks on the transparency of complex nuclei to photoproduced charged pions at intermediate energies, 26:46889

TAVES, Donald R., Fluorine metabolite concentrations and renal clearances following methoxyflurane anesthesia, 26:47645 (UR-3490-107)

TAVKHELIDZE, A. N., N. N. Bogolyubov's transformation in strong coupling theory. II, 26:46691

TAYLOR, Brian G., Scientific results from HEOS 1, 26:46334

TAYLOR, C. G., Radiometric determination of trace amounts of anionic surfactants in ground water and potable water, 26:45272

TAYLOR, Donald J., Radial cyanogen-band gradients near the nuclei of galaxies, 26:46243

TAYLOR, Harry A. Jr., Light-ion trough, 26:46301 (N-72-18362)

TAYLOR, J. Herbert, Chromosomes of eukaryotes: regulation of DNA replication, organization and integration of DNA and effects of decay of incorporated isotopes (^3H and ^{125}I), 26:45894 (ORO-2690-12(Pt.B), Paper 14)

TAYLOR, R. D., An efficient high-volume miniature air mover, 26:47633 (UCRL-51239)

TEDESCO, P. H., Thorium--acetate and thorium--propionate complexes, 26:45373

TEEL, Dale M., Sequential sampler, 26:45613

TERENT'EV, M. V., Concerning the two-photon mechanism of production during collision of charged particles, 26:46518 (ITEF-848)

TERESHKIN, Yu. M., Particle acceleration in a H_{111} cylindrical cavity in an axial magnetic field, 26:46139

TERHUNE, Robert W., Prediction of underground nuclear explosion effects in Wagon Wheel sandstone, 26:45494

TER-MIKAEYAN, M. L., High-energy electromagnetic processes in condensed media, 26:46538

TERNOV, I. M., Induced emission from electron in magnetic field at level $n = 1$, 26:46551

Motion of Dirac particles possessing anomalous magnetic moment in central symmetric electric field, 26:46547

TERRADO, J., Transmission of low-energy x-ray through various locally available materials, 26:47205

TERWILLIGER, K. M., 200-GeV proton--proton elastic scattering at high transverse momentum, 26:46456 (COO-1112-75)

TETELMAN, A. S., Reactor steels studies. Quarterly progress report, October--December 1971, 26:45994 (UCLA-34-P-177-10)

THALER, J. J., Charge asymmetry in the decay $\eta \rightarrow \pi^+\pi^-\gamma$,

26:46482

THATCHER, M. Llewellyn, Mathematical model for the prediction of unsteady salinity intrusion in estuaries, 26:47620 (MIT-144)

THEILE, F. W., Gamma gamma directional correlation measurements in ^{131}Cs , 26:46963

THEIN AUNG, Absolute differential scattering cross sections of 0.662 MeV photons, 26:47204

THEIN HTUT, Hla Pe, Absolute differential scattering cross sections of 0.662 MeV photons, 26:47204

THIELMANN, R., Gamma gamma directional correlation measurements in ^{131}Cs , 26:46963

THIEM, D., Influence of fission neutron spectra on integral nuclear quantities of fast reactors, 26:47585 (EURFNR-1008)

THIES, H. H., Conversion of bremsstrahlung photo neutron mean energies to monochromatic photo neutron mean energies, 26:46953

THIES, W. G., First forbidden beta decay of ^{209}Pb , 26:47007

THOMA, B. W., Conversion of bremsstrahlung photo neutron mean energies to monochromatic photo neutron mean energies, 26:46953

THOMAS, E. W., Angular distribution of metastable hydrogen formed by electron capture in a helium target, 26:46397

THOMAS, Edward W., Sample-conveying mechanism for scintillation counting, 26:46110

THOMAS, G. H., Remarks on average pion multiplicities at high energies, 26:46662

THOMAS, Gordon Albert, Resistive transition of a superconductor, 26:46817

THOMAS, K. S., Plasma experiments in the Scyllac 5-meter, linear theta pinch, 26:47141 (LA-DC-72-703)

THOMAS, L., Photodetachment of electrons from major negative ions in lower D region, 26:46299

THOMAS, R. J., Analysis of Ogo 6 observations of the O I 5577-A tropical nightglow, 26:46271

THOMFORD, Neil R., Pull-through operation for radiation-induced rectovaginal fistula. Report of a case, 26:45800

THOMPSON, Daniel Thomas, $^3\text{He}(^6\text{Li}, \alpha)^4\text{He}$ reaction at 1.54 MeV, 26:46897

THOMPSON, G. F., Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. II, 26:45328

Radiation-induced formation of excited states of aromatic hydrocarbons in cyclohexane. IV, 26:45330

Radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. III, 26:45329

THOMPSON, R. S., Dynamic structure of vortices in superconductors. III. Graphs for B approximately = $H_{\text{sub c2}}/2$, 26:46814

THOMSON, K. C., Elastodynamic near-field of a finite, propagating tensile fault, 26:45491

THORNE, Richard Mansergh, Pitch-angle diffusion of radiation-belt electrons within the plasmasphere, 26:46311

THORP, J. S., Electron spin resonance in single crystal yttria stabilized zirconia, 26:47252

THORSTEINSEN, T. F., Energy levels in ^{231}Th and ^{233}Th , 26:47029

THOUROUDE, Daniel, Approximations of the space charge field for different beam shapes in the central region of a synchrocyclotron, 26:46140

THROWER, P. A., Research on graphite. Final report for work completed during the period 1954 to 1972, 26:46078 (NYO-1710-106)

TIFFANY, B., Analysis of curium and californium by single-exposure spark source mass spectrometry, 26:45260

TIFFT, W. G., Correlation of redshift with magnitude and morphology in the Coma cluster, 26:46241

TILLMAN, J., Energy of levels of ^{48}Sc up to 3 MeV from $^{44}\text{Ca}(p, \gamma)^{48}\text{Sc}$ reaction, 26:46920

TINKO, D. J., Application of well logging methods in hydrocarbon prospecting, drilling, and production to the study of geostatic pressure problems, 26:45564

TINKHAM, M., Tunneling generation, relaxation, and tunneling detection of hole-electron imbalance in superconductors, 26:46811

TISHCHENKO, M. A., Microdeterminations of terbium by a luminescent method with the reagent pyrogallolsulfonic acid, 26:45231

TIITIRICI, M., (α, p) reaction on even nickel isotopes, 26:46934

TIWARI, G. P., Chemical diffusion in the beta phase of zirconium--titanium system, 26:45985 (BARC-577)

KEBUCHAVA, F. G., Quasi-threshold behavior of $\pi + N + N + (\gamma\text{-virtual})$ process amplitude, 26:46620 (JINR-P4-6475)

TOBIAS, Melvin, Review of the cross sections of the isotopes ^6Li and ^7Li as tabulated on the ENDF/B Version III files, 26:46876 (ORNL-TM-3859)

TODD, Richard Ross, Gamma ray spectroscopic studies of members of the A = 141 decay chain, 26:46972

TODOROV, T., Asymptotic numbers: one generalization of the notion of number, 26:47284 (IC-72/45)

TOEPFFER, Christian, Quasi-classical treatment of multinucleon transfer reactions, 26:47076

TOEROEK, G., Development of methods for application of

- neutron activation analysis to determination of trace elements in foods. Progress report, 10--12/71, 26:45189 (BFL-1/1972)
- TOEROEK, I., Measurement of ^{137}Cs and ^{90}Sr in soil, 26:45263 (RT/PROT-72-11)
- TOITA, Yukio, Separation and spectrophotometric determination of Ce and Mn by TTA extraction, 26:45226
- TOKSUBAEV, A. I., Use of admission system of MI-1309 instrument for precision isotopic analysis, 26:45681
- TOKUNAGA, Okihiro, Radiolyses of mixtures of ammonia and hydrocarbons. II, 26:45332
- TOLSTOI, A. E., Decrease of the off-duty factor in the operation of a linear accelerator, 26:46145
- TOMASSIAN, Albert D., Inner-zone energetic-electron repopulation by radial diffusion, 26:46310
- TOMMASINO, Luigi, Method and apparatus for counting radiation damage tracks in sheet material using a capacitive discharge device, 26:46083
- TOMOZAWA, Yukio, Time variation of the fine structure constant and spectral lines from quasi-stellar objects, 26:46254 (COO-1112-51)
- TONE, Hirohito, Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)
- TONEEV, V. D., Account of nuclear boundary diffusivity in the intranuclear cascade model, 26:47047 (JINR-P2-6503)
- Change in the mechanism of inelastic interactions between particles and nuclei in the energy region T approximately equal to 5 GeV, 26:47038 (BNL-tr-487)
- TORGOMYAN, F. B., Detector for high energy particles based on the principle of using particle radiation in a laminated medium, 26:45648
- TORIUMI, Tatsuo, Effects of cepharanthine on fluctuation in white blood cell count by radiotherapy for gynecologic cancer, 26:45805
- TOROSSIAN, C., Modifications induced by gamma irradiations in the ant, 26:45877
- TORRENS, Ian M., Computer simulation of atomic displacement cascades in metals, 26:46061 (CONF-710601-, pp 739-56)
- TORRES, J. P., Decay of ^{155}Dy , 26:46986
- TOSI, M. P., Self-consistent pair correlations and dielectric response in a plasma, 26:47150
- TOWNER, I. S., Low-lying positive-parity states of ^{35}Cl and ^{36}Cl from (p,d) reactions and shell-model interpretations, 26:46906
- TOWNSEND, John R., Study of radiation induced defects in metals. Final report, January 15, 1966--January 14, 1972, 26:46066 (COO-3431-1)
- TOXEN, A. M., Nuclear magnetic resonance in the LaX_3 intermetallic compounds, 26:47281
- TRABALKA, John Richard, Radioecology of Ce-- ^{144}Pr in a littoral fresh-water microcosm, 26:45553
- TRADOWSKY, Klaus, New electrical procedure for the compaction of technical powders, 26:45951
- TRAN THANH VAN, J., Conference on high energy phenomenology, Meribel-les-Allues, France, 3/71, 26:46590
- Exchange degeneracy and dip mechanisms in high energy collisions, 26:46592
- Phenomenological prescription for Regge cuts, 26:46622
- TRAUTVETTER, H. P., Excited states of ^{21}Ne , 26:46907
- TRAVIS, Larry Dean, Convection in stellar atmospheres, 26:46172
- TREBUKHOVSKII, Yu. V., Electrical insulation in electrostatic analyzers, 26:45687
- TREFFERS, R. R., Ten-micron spectroscopy of circumstellar shells, 26:46174
- TREILLE, D., Elastic scattering of π^+ , π^- , and anti p on protons at 5 GeV, 26:46492
- TRETIKOFF, O., MASURCA: study of plutonium-graphite and uranium-graphite assemblies, 26:47581 (CEA-N-1483)
- TRET'YAKOV, E. F., Conversion transitions in ^{226}Ra , 26:47028
- TRET'YAKOV, Yu. P., Production and acceleration of multicharged ions of phosphorus and zinc, 26:46133
- TRET'YAKOVA, N. I., Conversion transitions in ^{226}Ra , 26:47028
- TRIPATHI, A. N., Excitation of hydrogen in collisions with alkali-metal atoms, 26:46403
- TRIPATHI, D. N., Cross sections for single-electron capture in collisions of protons and deuterons with atoms, 26:46398
- TRIULZI, C., Reactor chemistry: analytical work, 26:45181 (CISE-R-172)
- TROBAUGH, F. E. Jr., Effect of various cell-free organ extracts on hematopoietic stem cells, 26:45847
- TRUCCO, R., Reactor chemistry: analytical work, 26:45181 (CISE-R-172)
- TRUJILLO, J. R., Cells for nondestructive examinations of irradiated fuel pins, 26:46104 (LA-DC-72-556)
- TRUNNELL, Thomas N., Acneiform changes in areas of cobalt irradiation, 26:45793
- TRUSOV, V. F., Theoretical coefficients of internal conversion for ^{51}V , 26:46921
- TSAREV, V. A., Photoproduction of neutral pions on helium nuclei, 26:46432 (SLAC-Trans-142)
- TSENDIN, L. D., Electron energy distribution and striations in the Langmuir-paradox region, 26:47133
- TSUJIDE, Tohu, Effects of x-ray irradiation on MAS diodes, 26:45670
- TSUMORI, Kunihiko, Effect of temperature on various types of the ionization chamber, 26:45594
- TSYPIN, V. S., Micro-instabilities of high pressure plasma associated with dependence of the ion viscosity on the derivatives of the longitudinal thermal flux, 26:47165
- TSYTOVICH, V. N., Note on plasma acceleration caused by plasma turbulence, 26:47134
- TUCKER, Wallace H., Relationship between x-ray luminosity and velocity dispersion in clusters of galaxies, 26:46249
- Time-dependent ionization equilibrium and line radiation under flarelike conditions, 26:46226
- TUCKER, Walter D., Isotope ratio tracer method: applications in atmospheric sulfur pollution studies, 26:45568
- TUELLER, Paul T., Ecology of Hot Creek Valley: vegetation and soil response to underground detonation. Annual progress report FY 1971, 26:45478 (NVO-409-1)
- TUNG, Eleanor W., Simple perturbation and perturbation--variation treatments of the $1s\sigma$ and $2p\sigma$ states of H_2^+ , 26:46355
- TURCO, R. P., Investigation of the ionospheric D region at sunrise. I. Time variations of ozone, metastable molecular oxygen, and atomic oxygen, 26:46279
- Investigation of the ionospheric D region at sunrise. III. Time variations of negative-ion and electron densities, 26:46281
- Investigation of the ionospheric D region at sunrise. II. Estimation of some photodetachment rates, 26:46280
- TURKEVICH, John, Annual progress report, June 1, 1971--May 31, 1972, 26:45313 (COO-3029-1)
- TURNER, G. I., Temperature dependence of the friction stress for basal dislocations in beryllium in the range 300 to 500°K, 26:46028
- TURNER, James, Simulators, 26:47572
- TURSKI, L. A., Model of the Josephson effect, 26:46834
- TUURNALA, T., Decay of 1.4 min ^{173}Er , 26:46997
- TYAGI, R. N., Pressure dependence of Moessbauer recoilless fraction, 26:47276
- TYLER, A. F., Design considerations in the measurement of electron temperature in the ionosphere, 26:46295
- TYUL'KIN, V. A., Radiation-induced centers of paramagnetic and optical absorption in glasses of systems SrO--SiO_2 and BaO--SiO_2 , 26:47226
- TYVIN, L. I., X ray fluorescence analysis of heavy element by an α -X coincidence method, 26:45270
- X-ray spectrometer having cooled $\text{Si}(\text{Li})$ -detector, 26:45644
- U, Raymond, Enhancing effect of lucanthone on frequency of x-ray-induced chromosome loss in *Drosophila* male germ cells, 26:45887
- UDOVENKO, A. A., Peculiarities in the structural sensitivity and mechanical characteristics of niobium, 26:46019
- UEBERALL, H., Spectra of neutrons emitted after muon capture in ^{18}O , ^{28}Si , ^{32}S and ^{40}Ca , 26:46880
- UENO, Kaoru, Precipitation of some actinide element complex ions by using hexamine $\text{Co}(\text{III})$ cation. IV, 26:45370
- UENO, Takehisa, Statistical analysis of radioinduced abnormal development, 26:45867
- UKAI, Seiji, Solution of multi-dimensional neutron transport equation by finite element method, 26:46851
- ULION, Nicholas E., Apparatus for measuring height of a molten metal pool, 26:45655
- ULKOWSKI, Mirosław, Evaluation of the causes of failures in roentgentherapy of carcinoma of the larynx, 26:45780 (TT-54002/4, pp 430-6)
- ULLAH, Nazakat, Distribution of the total fission width, 26:47045 (INDC-7-U, pp 81-6)
- ULLMAIER, H., Local magnetic field distributions in superconducting Nb at 4.2 K, 26:46826
- UMAYAHARA, Akira, Acute death of mice during massive gamma irradiation. III, 26:45846
- UMEMURA, Isao, Quartet scheme of hadrons in chiral $\text{U}(4) \times \text{U}(4)$, 26:46687
- UMEMURA, Sunao, Development of 50Hz 44 inch turbine blade, 26:45454
- UMEZAWA, H., Time-dependent phenomena in weakly-coupled superconductors, 26:46827
- UMNOVA, M. M., Vascularization of muscular autotransplants in irradiated site, 26:45857
- UNRUH, C. M., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- URAY, Z., AET on immunologic processes in irradiated rabbits, 26:45824
- URBAN, K., Application of high-voltage electron microscopy to the study of voids and other point-defect agglomerates, 26:46057 (CONF-710601-, pp 550-62)
- URCH, D. S., Estrup--Wolfgang theory of hot reactions. II, 26:45363
- UTSUDO, Masahiko, Slow neutron spectra in the liquid and solid methane, 26:46858



- VAHALA, George Martin, Kinetic theory of a two-dimensional magnetized plasma, 26:47146
- VAINSHTAIN, V. G., Effect of mechanical surface hardening on the low-cycle fatigue strength of titanium alloys, 26:46009
- VAISBERG, O. L., Multichannel modular spectrometer for low-energy electrons and protons, 26:45642
- VAISBURD, D. I., Accumulation of F centers in NaCl crystals as function of proton-beam intensity, 26:47246
- VAKHTEL, V. M., Signal filtration in the amplitude spectrometry systems, 26:45637 (JINR-P13-6428)
- VALENTIN, J., Two branchings in rotational band $1/2^-$ [541] in ^{171}Lu , 26:46987
- VALENTIN, L., Measurement of the quadrupole moment of a shape isomer, 26:46960
- VALLEE, R. E., Documentation of DOT-SP-5915 for 55/30 and 83/55 drum combination packaging, 26:46114 (MLM-1911)
- VAMPOLA, Alfred L., Inner-zone energetic-electron repopulation by radial diffusion, 26:46310
- VANAGAS, V. V., Theory of low-lying nuclear states in scheme U/sub 3(A-1)/, 26:47095
- Two-particle orbital matrix of density for one-quanta oscillatory-excited SU_3 -configurations for p-nuclei, 26:47067
- VAN CAMPEN, D. H., Stress distribution in nozzle-to-cylinder connections for small diameter ratios, 26:45445
- VAN CLEVE, J. E. Jr., Fabrication of ^{252}Cf neutron sources at Oak Ridge National Laboratory, 26:45714 (CONF-720902-2)
- VANDEMBEMDEN, E., Plutonium fuel production plant and breeder projects, 26:45974
- VANDENBOSCH, R., Single-particle effects on fission probabilities for the lighter elements, 26:47070
- VAN DER BORGH, O., Effect of Al phosphate gel on whole-body retention of simultaneously administered ^{22}Ra , ^{85}Sr , and ^{47}Ca in mice, 26:45918
- Human faecal ^{22}Ra excretion shortly after a RaSO_4 dust contamination, with alginate treatment, 26:45907
- VAN DE REE, J., Quantum-mechanical scattering of asymmetric molecules. II. Comparison between the systems $\text{He}-\text{D}_2$ and $\text{He}-\text{HT}$, 26:46379
- VAN DER PLOEG, F. A., S-matrix pole trajectories in the N/D approach, 26:46789
- VAN DER SCHUEREN, G., Changes in the localization of acetylcholinesterase in neuromuscular junction after ^{60}Co irradiation, 26:45828
- VAN DER WALT, A. J., Rigidity dependence of cosmic-ray modulation at rigidities > 2 GV in August 1969, 26:46327
- VAN DER WOUDE, A., Energy shifts and relative intensities of K x rays produced by swift heavy ions, 26:46339
- VAN DE VOORDE, M. H., Dose estimates outside the shielding of the CERN Intersecting Storage Rings by the Moyer method, 26:46121
- VAN DE VOORDE, Marcel, Effect of nuclear radiations on rubber joints, 26:46087
- VANDLIK, T. B., New isotope ^{180}Tl , 26:46999
- VANDLIK, Ya., New isotope ^{180}Tl , 26:46999
- VAN ERP, J. B., Analog computer models for safety evaluation of CART fuel test facility, 26:47580 (CISE-R-259)
- VAN PARADIJS, J. A., Evidence for the hydrodynamic character of microturbulence, 26:46188
- VAN POOLLEN, H. K., Projected reservoir performance. Rio Blanco Project, 26:45476 (NVO-38-33)
- VAN PUYNBROECK, S., Effect of Al phosphate gel on whole-body retention of simultaneously administered ^{22}Ra , ^{85}Sr , and ^{47}Ca in mice, 26:45918
- VANQUICKENBORNE, L. G., Coupling scheme in uranyl complexes, 26:45380
- VANSTONE, J. R., Weyl and Lyra manifolds, 26:46688
- VANSTRUM, Paul R., Report of uranium isotope separation review Ad Hoc Committee, 26:45689 (ORO-694)
- VAN ZELST, L., Titanium and calcium content of meteorites, 26:45215
- VANZHA, A. P., Electromagnetic effects in nucleon polarization at meson-nucleon scattering, 26:46520 (JINR-P2-6438)
- VARMA, M. N., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)
- VARMAZIS, C., Effect of surface conditions on some superconducting properties of niobium, 26:46805 (BNL-16913)
- VARSHNEYA, A. K., Diffusion in the system $\text{K}_2\text{O}-\text{SrO}-\text{SiO}_2$. III. Interdiffusion coefficients, 26:45946
- VARTERESZ, V., Determination of microbial contamination of disposable medical devices and the estimation of radiosensitivity of isolated microflora, 26:45748
- VASILENKO, M. N., Energy spectrum and crystalline structure of wurtzite-like modification of boron nitride, 26:47255
- VASIL'EV, M. N., Formation of the energy spectrum of the ions of a laser plasma, 26:47154
- VASIL'EV, S. S., Mechanism of $^9\text{Be}(p,p_1)^*{}^8\text{Be}$ 1.68 MeV reaction, 26:46885
- VASIL'EV, V. I., Magnetic shield for a 240-cm isochronous cyclotron, 26:46129
- VAZ, L. C., Level densities from spectra of different reactions populating the same nucleus, 26:46932
- VAZEILLE, F., Inelastic interaction induced by high-energy muons (6 GeV, 12 GeV) at low momentum-transfer in nuclear emulsion, 26:46451
- VDOVENKO, V. M., Chemical shifts in L/sub α_1 / x-ray line of some U compounds, 26:45377
- VEDDER, James F., Accretionary processes in the early solar system: an experimental approach, 26:46229
- VEDRINE, Jacques C., Powder ENDOR line shapes: nuclear relaxation induced by motion of nearby electron spins, 26:47279
- Proton ENDOR of γ -irradiated Y-type zeolites, 26:45324
- VEENING, Hans, Rapid separation of metal chelates by column liquid-liquid chromatography using ultraviolet detection, 26:45281
- VEITH, W., Persistent current states in rotating superfluid helium, 26:46801
- VEJE, E., Relative level populations in foil excited He and Li, 26:46337
- VELINOV, P. I., Conditions for acceleration of particles in solar atmosphere, 26:46230
- VENANZI, Thomas J., Higher multipole coefficients of the hydrogen-hydrogen interaction as determined by dynamic multipole polarizabilities, 26:46344
- VENKATESWARLU, K. S., Water chemistry studies VII: general review of water conditions and water treatment in nuclear reactors, 26:45310 (BARC-595)
- VENKOVA, Ts., New isotopes of iridium, 26:46977
- New isotopes of tungsten, 26:47001
- VENTURA, E., Evidence for an anomaly in the excitation function of the reaction $^2\text{H}(p,\gamma)^3\text{He}$, 26:46868
- VERHAAR, B. J., Cyclotron laboratory at Technical University of Eindhoven, 26:46146
- VERNIMB, C., Definitions of Proper Name Descriptors used in the Euratom-INIS Nuclear Documentation System, 26:47597 (EUR-4806)
- VERTEBNYI, V. P., Temperature dependence of cross section of cold- and thermal-neutron scattering by O_2 and D_2 , 26:46887
- VERTMAN, A. A., Thermal properties of liquid cerium, 26:45998
- VERZARIU, Pompiliu, Magnetohydrodynamic effects at the magnetosphere boundary, 26:46321
- VIDEAU, H., Reactions $\text{K}^- + p + \Lambda(\Sigma^0) + \text{vector meson}$ at 3.95 GeV/c, 26:46485
- VIDEAU, I., Reactions $\text{K}^- + p + \Lambda(\Sigma^0) + \text{vector meson}$ at 3.95 GeV/c, 26:46485
- VIE, Robert, Pneumatic machine for the remote controlled sealing of flasks in a shielded cell, 26:46112
- VIENNET, R., Analysis of neutron-deuteron elastic scattering at low energy, 26:46864
- VIGIER, J. P., Are vector potentials measurable quantities in electromagnetic theory, 26:46542
- VIGNON, B., Low-lying positive-parity states of ^{35}Cl and ^{36}Cl from (p,d) reactions and shell-model interpretations, 26:46906
- VILLAGRANA, Robert E., Perturbation treatment of high energy electron diffraction from imperfect crystals, 26:46510 (CALT-767-P-3-22)
- VILLORESI, G., Diurnal effect of cosmic rays during the period 15 October 1965-30 June 1966. I, 26:46159
- VISHNEVSKII, I. N., Decay of ^{198}Au , 26:47003
- VISOTSKAITE, S. B., Spectral and scintillation properties of the pyrimidine derivatives, 26:45606
- VISVANATHAN, N., Observations of planetary nebulae at 1.65 to 3.4 microns, 26:46175
- VITMAN, V. D., Isobaric nuclei with mass number $A = 72$, 26:46939
- VITOL, E. N., Allowance for mass-spectrometer background and the reproducibility in the determination of isotope compositions for nitrogen and oxygen, 26:45241
- VIVCHAR, O. I., Ternary system $\text{Y}-\text{Fe}-\text{Al}$ in region 0 to 33.3 percent Y, 26:46005
- VIVERS, Robert William Jackson, Processing of irradiated nuclear fuels, 26:45420
- VLASOV, N. A., Neutron matter, 26:46466
- VOEGELE, A. L., Contamination control hardware for remote hot cell transfers, 26:46108 (UCRL-73757)
- VOGELANG, W. F., Breeding ratio, inventory, and doubling time in a D-T fusion reactor, 26:47173
- VOGT, Oscar, dc anisotropic magnetization at 20.4 K and the ideal crystal-field-only behavior of TmSb , 26:47258
- VOJTECH, O., Strontium sorption on sorbents of BaSO_4 - CaSO_4 system, 26:45273 (UJV-2820-Ch)
- VOLCHOK, H. L., ^{90}Sr fallout in Utah, 26:45525 (HASL-257, pp I.71-94)
- High fallout in the western United States. An explanation, 26:45517 (HASL-257, pp I.18-32)
- Plutonium fallout in Utah, 26:45526 (HASL-257, pp I.95-118)
- VOLF, Frantisek, Sintered Mg composites, cladding material of U fuel elements cooled by carbon dioxide. II, 26:46012
- VOLKOV, M. K., Nonpolynomial Lagrangians, 26:46738
- S-matrix unitarity in theories with nonpolynomial Lagrangians, 26:46775 (JINR-P2-6393)
- VOLKOV, O. S., Possibilities of using radioisotope instruments in fabrication of high power plants of supercritical steam parameters, 26:45473

- VOLKOV, V. A., Chemical shifts in $L_{\alpha 1}$ x-ray line of some U compounds, 26:45377
- VON BRENTANO, Peter, Spectroscopic studies with isobaric analog resonances, 26:47099
- VON DROSTE, G. Frh, Gamma- and x-rays in decay of ^{166}Ho , 26:46998
- VON WIMMERSPERG, U., Test of time reversal invariance through detailed balance of the reaction $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$, 26:46881
- VORBRDT, A., Effect of x irradiation on cells, 26:45738
- VORBRDTOWA, Janina, Selected prognostic elements in carcinoma of the larynx, 26:45781 (IT-54002/4, pp 437-41)
- VOROB'EV, A. I., Chromosome aberrations in human bone-marrow cells after a single local gamma-irradiation. I, 26:45794
- VOROB'EV, L. S., Reaction (π^-, π^+) on ^6Li and ^{12}C , 26:46869
- VOROB'EV, V. D., Regularity in multiplet properties of odd-odd nuclei, 26:47051
- VORONKOV, Nikolai Andreevich, Method of modifying polymeric materials, wood, and articles made therefrom by radiation-initiated graft copolymerization thereof, 26:45355
- VOS, O., Stem cell renewal in spleen and bone marrow of mice after repeated total-body irradiations, 26:45871
- VOSBURGH, K. G., Acceleration of argon ions to 1.17×10^{10} electron volts, 26:46124
- VOTINOV, M. P., Interaction of a ruby with ionizing radiation, 26:47244
- VREDEVOE, Donna L., Single cell transplants: Gross virus-induced mouse lymphomas, 26:47629 (UCLA-12-866)
- VRILLON, B., Calculations and behavior of expansion bellows, 26:47591
- VUGMAN, N. V., ESR studies of electron irradiated $\text{K}_2\text{Ir}/\text{sup III}/(\text{CN})_6$ in KCl single crystals. I. Ligand hyperfine structure in $\text{Ir}/\text{sup II}/(\text{CN})_6$ species, 26:45317
- VUILLEUMIER, J. H., Test of quantum-electrodynamical corrections in muonic atoms, 26:46708
- VYLOV, Ts., Study of ^{17}Re decay, 26:46976 (JINR-P6-6492)
- Study of the decay $^{180}\text{Tb}(T_{1/2} = 66 \text{ min})$ and $^{180}\text{Tb}(T_{1/2} = 3.48 \pm 0.16 \text{ h})$, 26:46942 (JINR-P6-6512)
- VYVRYSHCHUK, Ya., Multipole order of transitions and characteristics of excited state of ^{147}Eu , 26:46954
- W**
- WAAK, B. T., Nuclear level sequence of single-particle energies in a momentum-dependent potential, 26:47071
- WACKS, Morton E., Techniques for source determination in the Tucson basin, 26:45206
- WADA, Takao, Recombination domain instability in high-energy electron-irradiated germanium, 26:47230
- WADDINGTON, J. C., Mixing of two-proton bands with two-neutron bands in ^{174}Hf , 26:46992
- WAELEI, Paul, Space-energy-dependent physics problems of plutonium fuels in HTGRs, 26:47313
- WAGNER, G. R., Paramagnetic hole centers in CeO_2 , 26:47233
- WAGNER, H., Comparison of lattice parameter and resistivity change in electron-irradiated copper, 26:47231
- WAHL, John S., Pulse neutron logging tool control and transmission system, 26:45563
- WALDMAN, Helio, Coupling between the ionosphere and the protonosphere and its implications on the long-term variations of ionospheric electron content, 26:46291
- WALINDER, G., Determination of ^{131}I dose to mouse thyroid, 26:45902
- Effect of irradiation on thyroid growth in mouse foetuses and goitrogen challenged adult mice, 26:45865
- WALKER, P. L. Jr., Research on graphite. Final report for work completed during the period 1954 to 1972, 26:46078 (NYO-1710-106)
- WALKER, W. D., Progress report [on high-energy physics], July 1, 1971--June 30, 1972, 26:46479 (ORO-3065-24)
- WALKER, William G., Scintillation camera having improved peripheral response, 26:45596
- WALLACE, A., Radioecology and ecophysiology of desert plants at the Nevada Test Site, 26:45755 (TID-25954)
- WALSH, T. F., Testing triplet models, 26:46516 (DESY-72/21)
- WALTER, H. K., Test of quantum-electrodynamical corrections in muonic atoms, 26:46708
- WALTERS, G. P., Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)
- WALZ, Albert, Use of reinforced plastic in a uranium production factory, 26:46080
- WAMBACH, U. M., Circular polarization of γ rays in ^{18}F , 26:46877
- WANAGEL, John, Study of hydrogen in vanadium, 26:47263
- WANG, Ling-Lie, Scaling in $p\pi^- + p\pi^-$ and an estimate of the triple-Pomeranchukon coupling, 26:46616 (BNL-16928)
- WANG, P. S. C., Upper and lower bounds to excited states of two-electron atoms, 26:46413
- WANG, Shih Y., Chemical study of the effects of radiation on nucleic acids and related compounds. Three year progress report, June 1969--May 1972, 26:45315 (COO-3286-3)
- WANG, Shih Yi, Radiation chemistry of nucleic acids. Isolation and characterization of thymine glycols, 26:45314 (COO-3286-1)
- WARASHINA, Tetsuo, Trapped electrons produced in ethanol glass at 4°K , 26:45319
- WARCHOLD, L., WAK: specifications and finishing control for equipment, 26:46105 (ORNL-tr-2371)
- WARD, C. H., Synthesis, identification, and purification of C_6 and C_7 fluorocarbons from radiolysis. Final report, 26:45316 (ORO-3909-3)
- WARD, D., Deorientation of nuclei of recoil atoms, 26:46935
- WARD, Jerrold M., Bone and soft-tissue neoplasms in cats exposed to radiostrontium, 26:45901
- WARREN, Shields, Radiation carcinogenesis. Progress report VI, May 1, 1971--April 30, 1972, 26:45814 (COO-3017-43)
- WASHBURN, Jack, Annealing of loops and tetrahedra, 26:45989 (CONF-710601-, pp 647-62)
- WASSERMAN, J., Lymphocyte stimulation by thyroglobulin after local irradiation of the thyroid gland in man, 26:45889
- WATAGHIN, G., Remarks on some symmetry problems in cosmology and in the theory of particles and fields, 26:46674
- WATANABE, Keiji, Pion form factor and possible heavy vector meson, 26:46561
- WATANABE, Toshikazu, Hepatic injuries following irradiation, 26:45809
- WATERBURY, G. R., Remotized apparatus for determining oxygen in irradiated reactor fuels and cladding materials, 26:45224 (LA-DC-72-524)
- WATERS, J., Radiometric determination of trace amounts of anionic surfactants in ground water and potable water, 26:45272
- WATSON, E. C., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)
- WATSON, G. M., Environmental hazards of fossil and nuclear power production, 26:45510 (AAEC/TM-627)
- WATSON, R. E., Single-electron energies, many-electron effects, and the renormalized-atom scheme as applied to rare-earth metals, 26:47260
- WATSON, W. D., Abundances of interstellar molecules, 26:46244
- WATTS, H. L., Design studies of a molten-salt reactor demonstration plant, 26:47305 (ORNL-TM-3832)
- WEAVER, David L., Multipole radiation solutions for a class of massive particle wave equations, 26:46709
- WEAVER, H., 150-meter time-of-flight facility at ORELA, 26:46144 (CONF-720607-22)
- WEBSTER, A. S., Can the lumpy distribution of galaxies be detected by x-ray observations, 26:46260
- WEDBERG, G. H., ($^{16}\text{O}, ^{12}\text{C}$) studies in the $2s-1d$ shell, 26:46913
- WEEKES, T. C., Detection of high-energy gamma rays from the Crab Nebula, 26:46214
- WEHENKEL, Claude, Calculation of spectra of electron energy losses in Au from 1 to 12 eV, 26:47212
- WEHRL, Alfred, Methods of nonlinear analysis, 26:47283 (CERN-72-8)
- WEINER, R., Derivation of the critical temperature for the superfluid phase of hadronic matter, 26:46596
- WEINGART, R. C., X-ray-induced photoconductivity in dielectric films, 26:45668 (UCRL-73974)
- WEINHOLD, F., Upper and lower bounds to excited states of two-electron atoms, 26:46413
- WEINSTOCK, Jerome, Nonlinear saturation of Type-I irregularities in the equatorial electrojet, 26:46277
- WEIR, R., Removal of radioactive iodine from ventilation exhaust streams at the AAEC Research Establishment, 26:45502
- WEISBIN, C. R., Nondestructive assay of power reactor fuel assemblies, 26:45269
- WEISMAN, Joel, Cross flow resistance in rod bundle cores, 26:47596
- WEISS, U., Practical relativistic three-particle theory with arbitrary particle interactions, 26:46779
- WEISSBERGER, Arnold, Physical methods of chemistry. Part IIB, 26:45250
- WEISSKOPF, Victor F., Physics in the twentieth century: selected essays, 26:46508
- WEISS-REUTER, R., Preparation and investigation of neptunium isotopes 231, 232, and 233, 26:47023 (KFK-1537)
- WELCH, Ross M., Vanadium determination in biological materials at nanogram levels by a catalytic method, 26:45203
- WELLENSTEIN, Hermann Fridenart, Relaxation processes of the $n = 3$ states of helium, 26:46348
- WELLMAN, Henry N., Cyclotron production of ^{123}I : an evaluation of the nuclear reactions which produce this isotope, 26:45392 (N-72-15068)
- WELSH, L. B., Nuclear magnetic resonance in the LaX_3 intermetallic compounds, 26:47281
- WENDER, M., Effect of ionizing radiation on enzyme activity in the neuroglia of the developing rabbit brain, 26:45836
- WENDER, S. A., Analysis of five line hyperfine split Moessbauer absorber spectra, 26:47275
- WERBROUCK, A. E., Evidence for double diffractive dissociation in $\pi^+\pi^+$ reactions at 5 GeV/c, 26:46493

WERNER, Richard W., Method of fabricating a heat pipe, 26:45449

WESTPHAL, J. A., Optical search for Centaurus X-3, 26:46213

WHALEY, Hubert L., Ultrasonic flaw determination by spectral analysis, 26:45469

WHITE, Earl L., Stress-corrosion cracking of sensitized stainless steel in oxygenated high-temperature water, 26:45966 (BML-1927)

WHITE, J. A., Measurement of an optical frequency and the speed of light, 26:46733

WHITE, J. S., Temperature dependence of the friction stress for basal dislocations in beryllium in the range 300 to 500°K, 26:46028

WHITE, Sheldon Andrew, Effect of ionizing radiation on the stable fly, *Stomoxys calcitrans* (Linnaeus), 26:45883

WHITEHEAD, Neil E., Gamma radiation of some plants and soils from a uraniferous area in New Zealand, 26:45529

WHITWORTH, James Bradley, Development and application of a systematic approach to elemental analysis in forensic investigations, 26:45213

WICART, L., Bleomycin labeled with ^{57}Co , 26:45929

WICHMANN, P. A., Application of neutron activation to the determination of elements in boreholes, 26:45567

WIDEROE, R., Quantitative and qualitative aspects of radiobiology and their significance in radiation therapy, 26:45735

WIEDERSICH, Hartmut, Nucleation of voids in materials during high-energy irradiation, 26:46064 (CONF-710601-, pp 825-40)

WIESNER, James, Radiation effects on plasma electron-spin-resonance (ESR) spectra of cancer patients, 26:45723

WILDERMUTH, K., Consideration on direct reaction mechanisms in the unified microscopic nuclear theory, 26:47077

WILES, D. R., Nuclear recoil reactions in organomanganese compounds. VI, 26:45357

WILHELM, J., Critical component of the interplanetary magnetic field responsible for large geomagnetic effects in the polar cap, 26:46266

WILK, G., Photoproduction of rho-mesons on deuterons, 26:46555

WILKENS, M., Application of high-voltage electron microscopy to the study of voids and other point-defect agglomerates, 26:46057 (CONF-710601-, pp 550-62)

WILKNISS, P. E., Production of carrier-free $^{197}\text{m}\text{Tl}$, 197/Hg with a cyclotron, 26:45395

WILLIAMS, K. R., Application of high-voltage electron microscopy to void studies, 26:46056 (CONF-710601-, pp 533-49)

WILLIAMS, L. C., Fabrication of ^{252}Cf neutron sources at Oak Ridge National Laboratory, 26:45714 (CONF-720902-2)

WILLIAMS, Spencer L., Comparison of response spectra from Yucca Flat and Pahute Mesa underground nuclear detonations, 26:45480 (NVO-1163-IM-32)

WILLIAMS, T. M., Void formation in metals during ion bombardment, 26:46052 (CONF-710601-, pp 430-48)

WILLIS, David L., Radionuclide cycling in natural populations of amphibians. Annual progress report, June 16, 1971—June 15, 1972, 26:45913 (RLD-2227-T-8-3)

WILLNER, S. P., Observations of planetary nebulae at 1.65 to 3.4 microns, 26:46175

WILLS, E. D., Effects of dietary antioxidants on lipid peroxide formation in animal tissues after whole-body irradiation, 26:45872

WILSON, A. J., Variability of OH source W3, 26:46205

WILSON, F. D., Radiation-induced transplantable myelomonocytic leukemia in beagles: hematologic parameters, 26:45897

WILSON, R., Early data from ultraviolet sky-scan telescope in TD1 satellite, 26:46186

WILSON, W. D., Helium migration in metals, 26:46050 (CONF-710601-, pp 230-47)

WILSON, Wayne H., Pull-through operation for radiation-induced rectovaginal fistula. Report of a case, 26:45800

WINCEL, Henryk, Ion-molecule reactions in ammonia at high pressures, 26:45353

WINCHESTER, John W., Multi-element analysis of air pollution particulates by nondestructive neutron activation, 26:45207

WINTERS, William I., Spectrophotometric determination of phosphate in nuclear materials, 26:45219 (ARH-SA-110)

WINZER, K., Anomalous critical field curves of Kondo superconductor $(\text{La}/\text{sub } 1-x/\text{Ce}/\text{sub } x)/\text{Al}_2$, 26:46832

WIT, R., Two-particle correlations in 60 GeV/c $\pi^+\pi^-$ and 67 GeV/c $\pi^+\pi^-$ inclusive reactions, 26:46460

WITALIS, E., Depth distribution studies of carbon in steel surfaces by means of charged particle activation analysis with an account of heat and diffusion effects in the sample, 26:45466 (AE-451)

WITCOMB, M. J., Effect of dislocation tangles on superconducting properties, 26:46836

WITTE, L. C., Explosive interaction of molten metals injected into water, 26:45461

WOBIG, Horst, Convection of a plasma in a gravitational field, 26:47160

WOGMAN, N. A., Subcritical multiplication of ^{252}Cf neutrons and its applications, 26:46855

WOJCIESZEK, Zbigniew, Prognostic elements in inoperable carcinoma of the breast, 26:45783 (TT-54002/4, pp 451-5)

WOLBER, G., Hyperfine structure of ground state of ^{165}Ho , 26:46989

WOLF, Norman S., Effects of x irradiation upon response of mononuclear phagocytes to an induced sterile intraperitoneal inflammation, 26:45833

WOLF, W., Industrial processing of irradiated potatoes, 26:45716 (BFL-2/1972)

WOLFENDALE, A. W., Stopping particles underground, 26:46157

WOLFER, W. G., West Valley Reprocessing Plant. Letters of May 23, June 26, 27, 30, and July 5, 1972, with respect to modifications in the plant, 26:46065 (CONF-710601-, pp 841-52)

WOLFF, U. E., Fluence and temperature dependence of void formation in highly irradiated stainless steels, 26:46048 (CONF-710601-, pp 156-73)

WOMBACHER, P., Comparison of lattice parameter and resistivity change in electron-irradiated copper, 26:47231

WONG, K. M., Plutonium and polonium inside giant brown algae, 26:45924

WONG, Kai M., Plutonium in North Atlantic Ocean organisms; ecological relationships, 26:45541 (NYO-2174-130)

WOODBRIIDGE, D. D., Fluid irradiating apparatus using gamma rays and helical passageways, 26:45719

WOODBRIIDGE, David D., Irradiator apparatus with a fluid flowpath determined by communicating core and casting baffles, 26:45456

WOODMAN, Ronald F., Synthesis of Jicamarca data during the great storm of March 8, 1970, 26:46282

WOODRUFF, R. K., Technological considerations in emergency instrumentation preparedness. Phase II-A. Emergency radiological and meteorological instrumentation criteria for reactors, 26:47317 (BNWL-1635)

WOOSLEY, S. E., Hydrostatic oxygen burning in stars. II. Oxygen burning at balanced power, 26:46178

WORMELI, John C., Remote welding program for Atlantic Richfield Hanford Company's waste encapsulation and storage facility, 26:46099 (ARH-SA-117)

WORTMAN, Donald E., Nuclear charged self-sustaining laser, 26:45677

WRESZINSKI, W. F., Note on construction of Bogolyubov scattering operator in $(\phi^4)_2$ theory, 26:46740

WRIGHT, James F., Bone and soft-tissue neoplasms in cats exposed to radiostrontium, 26:45901

WU, Mao-Fou, Correlation of D-region absorption and the 10 mb winds and temperature during the IQSY, 26:46264 (COO-2195-5)

Y

YAES, R. J., Kinematically corrected correlation functions for inclusive reactions, 26:46500

YAGUNOV, A. S., Use of a model of two classes of tryptophan residues for interpreting the results of ultraviolet fluorescence of cells exposed to x-rays, 26:45736

YAMABE, Tokio, Long-range interactions between two hydrogen atoms and two helium atoms, 26:46351

YAMADA, Jun, Impedance of a partially ionized cesium plasma, 26:47135

YAMADA, Kazuo, Phonon dispersion and contribution to the heat capacity in liquid ^4He , 26:46795

YAMAFUJI, Kaoru, Flux flow in nonideal Type-II superconductors, 26:46819

YAMAGISHI, Tomejiro, Disappearance of neutron diffusion length of beryllium and graphite according to temperature, 26:46857

YAMAGUCHI, Masafumi, Effects of impurities on gamma-irradiated silicon crystal examined by photovoltaic effect of p-n junction diode, 26:45671

YAMAGUCHI, Yoshio, Parity tests in elastic $e^+\pi^-$ scattering at large angles, 26:46562

YAMAMOTO, Katumune, Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)

YAMAMOTO, Kunio, Transformation properties of operators in local field theory, 26:46745

YAMANE, Takeshi, Determination of small amounts of molybdenum based on its catalytic effect in Methylene Blue-hydrazine sulfate reaction, 26:45227

Polarographic determination of Sn in Cu-, Al-, and Zr-base alloys, 26:45256

YAMAZAKI, Hideo, Theoretical consideration of electron-scavenging process in liquid hydrocarbons. III, 26:45333

YAMAZAKI, Masakatsu, Multiple-scattering expansion and the Glauber approximation, 26:46792

YANCHO, G., Spectrum symmetry of pion longitudinal momenta in $\pi^+\pi^-$ interactions at $\sqrt{s} = 4$ GeV, 26:46619 (JINR-P1-6461)

YANG, Bing-tsuen, Gaseous plume diffusion about isolated structures of simple geometry, 26:45514

YANG, T. C., Energy-momentum conservation in inclusive processes, 26:46504

YANO, Kyo, Observation of the intensity ratio between (5,1) and (9,4) bands of OH emission in the night airglow, 26:46286

YARIV, A., Optical waveguiding in proton-implanted GaAs, 26:47232

YASINSKAS, L. L., Spectral and scintillation properties of

- the pyrimidine derivatives, 26:45606
- YATANI, Ryuichi, Induction of unscheduled DNA synthesis by half-nucleus irradiation of HeLa cells with uv-microbeam, 26:45737
- YATES, Albert C., Multiple-scattering effects in high-energy electron-molecule collisions. II. Polyatomic molecules, 26:46381
- YEN, W. L., Final states with three charged particles and a visible Λ from K-d interactions at 4.5 GeV/c, 26:46474 (COD-1428-306)
- YEROUHALMI, David, Composite walls for magnetohydrodynamic apparatus, 26:45426
- YEUNG, Edward S., Isotopic separation by photopredissociation, 26:45307
- YOKOSAWA, A., K ρ phase shift analysis and exotic resonances, 26:46632
- Polarization at large angles, 26:46462
- YOKOUCHI, Iichiro, Chemical analysis procedures for JMTR, 26:45182 (JAERI-M-4594)
- YOO, Kun Joong, Time-dependent neutron transport equation with delayed neutrons, 26:46847
- YOSHIKAWA, Shoichi, Nuclear fusion device of the air-core tokamak type, 26:47177
- YOSHIMURA, M., γ radiolysis of gaseous carbon dioxide at high densities, 26:45346
- YOUNG, Arthur H., Radioactivity well logging system having a pad mounted neutron source and a centralized radiation detector that provides compensation for borehole fluid density and borehole diameter variations, 26:45562
- Thermal neutron activation radioactivity logging method, 26:45561
- Well logging method for determining the presence of aluminum and magnesium, 26:45559
- YOUNG, Bing-Lin, Testing triplet models, 26:46516 (DESY-72/21)
- YOUNG, E. C. M., Stopping particles underground, 26:46157
- YOUNG, F. W. Jr., Effect of irradiation temperature on size of defect aggregates in copper, 26:46060 (CONF-710601-, pp 672-83)
- YOUNG, J. A., Oceanic distributions and relationships of ^7Be and fission products, 26:45530 (BNWL-SA-4205(Rev.))
- YOUNG, Richard A., Adhering resins to substrates, especially metal, by radiation, 26:46089
- YOUNG, William H., Trends in capital costs of generating plants, 26:47385
- YUANG, J. T., Lifetime of 975-keV, $3/2^+$ state in ^{25}Mg , 26:46908
- YUDINTSEVA, E. V., Accumulation of ^{90}Sr in wheat on sod-podzolic soils, 26:45550
- YUDOVICH, Ya. E., Distribution and genesis of radioactivity in coals of the Pechora River basin, 26:45505
- YUITA, Koichi, Production of carrier-free ^{54}Mn , 26:45394
- YULDASHEV, B. S., Spectrum symmetry of pion longitudinal momenta in πp interactions at $pc = 4$ GeV, 26:46619 (JINR-P1-6461)
- YUMOTO, Takaari, Synthesis of O-containing halogenated compounds. II, 26:45335
- YUR'EV, B. P., Electrolytic deposition of W-Co alloys, 26:45977
- Influence of temperature on electrodeposition of tungsten-cobalt alloy, 26:45983
- YUSHKOV, Yu. G., Electron acceleration in an undulator synchrotron, 26:46120
- YUSUPOV, M. Yu., Phenylthiazolylazochromotropic acid as a photocalorimetric reagent for thorium, 26:45240
- YUYAMA, Shuhei, Nuclear-cytoplasmic relations in the mitosis of sea urchin eggs. III. γ -ray-induced damage to whole eggs and nucleate and anucleate half-eggs, 26:45874
- YVON, K., Identification of the superconducting phases in the molybdenum-platinum system, 26:46828
- Z**
- ZABELIN, I. V., Electrolytic deposition of W-Co alloys, 26:45977
- Influence of temperature on electrodeposition of tungsten-cobalt alloy, 26:45983
- ZABIYAKIN, G. I., Computer systems for physics research centers, 26:45686
- ZACHARIASEN, F., Self-consistent diffraction in the multiperipheral model, 26:46602
- ZACHARY, W. W., Nonrelativistic time-dependent scattering theory and von Neumann algebras. I. Single channel scattering, 26:46781
- ZAGER, B. A., Production and acceleration of multicharged ions of phosphorus and zinc, 26:46133
- ZAGORSKI, Z. P., Changes of Dextran irradiated with γ radiation, 26:45730
- ZAHARAN, A. H., Chemical effects associated with the thermal neutron irradiation of solid acetanilide (antifibrin), 26:45358
- ZAITSSEV, E. V., Peculiarities of formation of an increased uranium concentration in oxidizing conditions, 26:45504
- ZAITSSEVA, N. G., New isotope ^{187}Tl , 26:46999
- ZAKHAROV, V. I., Interpretation of experiments on searches for K/sub S/ $\rightarrow 2\mu$ decay, 26:46572
- ZALMANZON, Yu. E., Distribution of a radioactive aerosol in a model chamber, 26:45444
- ZAMB, N. K., First Indian reactor, 26:47588
- ZAMPAGLIONE, V., Threshold of ionization instability in an unbounded low temperature plasma, 26:47169
- ZAPOROWSKI, B., Investigation of optimum conditions for energy conversion in MHD generator, 26:45430
- ZARDI, F., Solution of the standing-wave integral equation, 26:46786
- ZARECHNYUK, O. S., Ternary system Y-Fe-Al in region 0 to 33.3 percent Y, 26:46005
- ZARUBIN, P. P., Elastic and inelastic scattering of 6-MeV protons by even isotopes of Cr, 26:46922
- ZASIMCHUK, E. E., Textures of rolling and annealing in tungsten foils of various compositions, 26:46042
- ZASTAVENKO, L. G., Basic state of many-fermion system, coupled by gravitational lines, 26:46748 (JINR-P2-6270)
- Lower unboundedness of the Hamiltonian with scalar electrodynamics, 26:46702 (JINR-P2-6474)
- ZDARKO, R. W., Asymmetry measurements for π^0 and π^+ photoproduction by linearly polarized γ -rays in the energy range (40 to 900) MeV, 26:46442
- ZECCA, A., Casualty group in finite space-time, 26:46724
- ZEH, H. D., Simple theory of proton-proton final state interaction in reaction $d + p \rightarrow p + p + n$, 26:46650 (KFK-1547)
- ZELENAY, Krystyna, Method for labeling human albumin with ^{99m}Tc , 26:45927 (IT-54002/4, pp 471-4)
- ZELENY, V., Assessment and control of environmental contamination, 26:45554
- ZELLE, M. R., Ocular squamous cell carcinoma (cancer eye) in hereford cattle: radiation repair processes and a comparison of cultured cells with xeroderma pigmentosum in man, 26:45851
- ZEMAN, Herbert David, Optimization of low-energy electron scattering for producing a beam of spin-polarized electrons, 26:46338
- ZEMTSEV, B. V., γ -emission in radiative capture of intermediate neutrons, 26:46917 (BNL-tr-477)
- ZENKEVICH, P. R., Instability of axial oscillations in an electron-ring accelerator, 26:46142
- ZENKIN, V. A., Shape of the mass peak in a three-dimensional quadrupole mass spectrometer, 26:45676
- ZEROKA, Daniel, Theoretical studies of diamagnetic properties of the hydrogen molecule ion. II. Effect of varying internuclear separation, 26:46354
- Theoretical studies of diamagnetic properties of the hydrogen molecule ion. I. Approximate variation--perturbation calculation, 26:46353
- ZHABITSKII, V. M., Influence of the electromagnetic ring acceleration on the focusing by image forces, 26:46136 (JINR-P9-6421)
- Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)
- ZHARKOV, V. A., Protection from neutron emission of ^{238}Pu dioxide, 26:46852
- ZHELINOROVICH, V. A., Energy-momentum determination in the general theory of relativity, 26:46756
- ZHEREGII, F., Account of nuclear boundary diffusivity in the intranuclear cascade model, 26:47047 (JINR-P2-6503)
- ZHUK, V., Multipole order of transitions and characteristics of excited state of ^{147}Eu , 26:46954
- ZHUKOV, M. F., Erosion of a tungsten cathode in nitrogen, 26:45968
- ZHUKOVSKII, N. N., Decay of ^{78}Br , 26:46938
- ZHUNKOVSKII, G. L., Reaction of Nb-Mo alloys with B, 26:46002
- ZHURAKOVSKII, E. A., Energy spectrum and crystalline structure of wurtzite-like modification of boron nitride, 26:47255
- ZHURASKENE, E. A., Spectral and scintillation properties of the pyrimidine derivatives, 26:45606
- ZHURAVLEV, A. F., Induced emission from electron in magnetic field at level $n = 1$, 26:46551
- ZHURAVLEV, D. A., Collective interactions of nucleons of high-energy cosmic heavy nuclei, 26:46461
- ZHURINA, L. S., Multichannel modular spectrometer for low-energy electrons and protons, 26:45642
- ZIETEK, W. J., Unitary operator form of general Bogolyubov-Tyablikov transformation, 26:47291
- ZIMA, V. G., Photon exchange in $\pi + N \rightarrow \rho + N$ reactions, 26:46643
- ZIMANYI, J., Repulsive core effects in (t,p) stripping reactions, 26:47078
- ZIMMER, William H., Relative efficiency as a function of time for Atlantic Richfield Hanford Company coaxial Ge(Li) detectors, 26:45586 (ARH-2432)
- ZIMMERMAN, Ivan Harold, Coupled-state calculations for proton-hydrogen scattering in the Reading representation, 26:46370
- ZIMMERMANN, G., Special machines and equipment of WAK, 26:46106 (ORNL-tr-2372)
- ZINDLER, H., Some problems of the discharge of radioactive gases and aerosols from nuclear facilities into the atmosphere, 26:46102 (ORNL-tr-2369)
- ZINOV, V. G., New method of determination of physical-chemical properties of matter, 26:47618 (JINR-P14-6407)
- ZINOV'EV, G. M., Duality, scaling, and fixed poles in virtual Compton scattering, 26:46559
- ZIYAKAEV, R. G., Electron acceleration in an undulator

synchrotron, 26:46120

ZMIJEWSKA, W., Activation methods of analysis, 26:45201

ZNAMIROVSKI, V., Influence of ethanol on hydrated complexes of Cu ions, 26:45296

ZOHTA, Y., Electrical properties of n-type Si layers doped with proton bombardment induced shallow donors, 26:47237

ZOHTA, Yasuhito, Determination of the spatial distribution of deep centers from capacitance measurements of pn junctions, 26:47243

ZOLINA, I. A., Stability of the screened ring at the center shift, 26:46137 (JINR-P9-6514)

ZONDERHUIS, J., Determination of chromium in milk powder, 26:45196

ZONN, W., Predominant motion in spherical systems of galaxies, 26:46235

ZORCHEV, S. N., Possibilities of using radioisotope instruments in fabrication of high power plants of supercritical steam parameters, 26:45473

ZORINA, T. A., Problems of production, sanitation, and technical hydrobiology of the southern seas, 26:45919

ZUCKER, M. S., Continuous sampling plan for the taking of physical inventories, 26:46092 (BNL-16930)

61 AUTHORS

ZYWIETZ

ZUND, J. D., Special class of states of total pure radiation in general relativity, 26:46771

ZUPANCIC, M., $2^+ \rightarrow 2^+$ transition of 879 keV in ^{160}Dy investigated by electron--gamma directional correlation, 26:46991

ZUPANIC, C., New results on $(\pi^+, 2p)$ reactions in light nuclei, 26:46870

ZVOL'SKI, I., ^{175}Hf states excited during decay of ^{175}Ta , 26:46979

ZWANZIGER, Daniel, Thirring model in terms of currents: solution and light cone expansions, 26:46696 (CONF-710816-19)

ZYSS, Ryszard, Effects of ionizing radiation on the structure and metabolism of mitochondria in rat liver cells, 26:45838

Lysosomal system of liver cells in rats following a radiation dose of 1000 r to the whole body, 26:45839

ZYWIETZ, Friedrich, Dosimetry problems with bremsstrahlung in the GeV range, 26:45625

SUBJECT INDEX

Cross references are included in cumulative subject indexes and in *Subject Headings Used by the USAEC Technical Information Center* (USAEC report TID-5001).

The reader is also referred to *Guide to Abstracting and Indexing for Nuclear Science Abstracts* (USAEC report TID-4576).

Most headings in the subject index are the names of specific materials, things, and processes. Each heading is followed by a modifier that describes the properties of, or processes applied to, that heading.

All index entries cite *NSA* volume and abstract numbers; volume number precedes the colon and the abstract number follows it. The designation (R) following the abstract number indicates a progress report; (P) indicates a patent; and (T) indicates a translation. Abstracts for reports other than progress reports carry no letter designation. Report numbers, where applicable, are included in parentheses after the abstract number of the citation. Published literature (journal article or book) can be identified by the absence of a report number.

Ideas presented in the original literature, rather than key words, have been indexed. The indexers have selected what seemed the most probable and logical place to look for typical information. Since it is not always possible to anticipate where a user would look for information, the cumulative indexes include numerous cross references. Consideration of synonymous or closely related terms will prove valuable.

Information is indexed for the greatest use to the greatest number of users. This requires that a topic be indexed to headings that most specifically describe it and not ordinarily to the headings that describe its ultimate use or to the general class headings that include the specific headings. For example, information on boiler design for the Fermi Fast Breeder Reactor would be indexed under the headings BOILERS and FERMI FAST BREEDER REACTOR but not necessarily under the headings REACTORS, FAST or REACTORS, BREEDER.

General or broad headings are reserved for general summaries, reviews, comprehensive reports, and similar literature.

Subject Heading Styles

Generally, the direct form of subject entry is used rather than inverted headings; e.g., AMINO ACIDS rather than ACIDS, AMINO; HELIUM ISOTOPES He-4 rather than HELIUM—ISOTOPES He-4 or HELIUM—He-4 ISOTOPES. Note, however, that in some cases parenthetical modifying terms or inverted headings such as NEUTRONS, FAST, are used where the alphabetical arrangement of a direct entry would cause information to be scattered. In addition, organic compounds are usually entered in the index in the inverted form to facilitate grouping like or related compounds with the parent compounds; e.g., BENZENE, BROMO- rather than BROMOBENZENE.

Alphabetization

Subject headings are alphabetized word by word. Parenthetical qualifiers, numbers, and Greek letters within subject headings are disregarded in alphabetizing unless there are two or more headings that are otherwise identical, in which case the qualifier, numerical progression, or spelled names for the Greek letters determine the order.

Examples:

M-Capture
see ELECTRON CAPTURE
MAGNESIUM
MAGNESIUM ALLOYS AND SYSTEMS
MESONS
MESONS (π)
MESONS (ρ)
1-PENTANOL
2-PENTANOL
URANIUM OXIDES
URANIUM OXIDES UO_2

Modifying Phrases

The modifiers appearing under a subject heading are written specifically to show the relation between the item being indexed and the heading. The modifier therefore contains a maximum of useful information. Although the modifier begins with the word judged to be of greatest use to the greatest number of users, there is still need for the searcher to consider all modifiers under the heading(s) of interest.

Modifiers usually form meaningful phrases when combined with the subject heading. Some are intended to be read following the subject heading, e.g., the entry HYDROGEN—*sorption by charcoal*, should be read *hydrogen sorption by charcoal*. As a result of placing the important word first, however, many modifiers must be rearranged for smooth reading, e.g., FORMALDEHYDE—*yields in radiolysis of ethyl ether, effects of deuterium on*, should be read *effects of deuterium on formaldehyde yields in radiolysis of ethyl ether* and URANIUM—*crystal structure of, effect of cooling rate on allotropic transformations in*, should be read *effect of cooling rate on allotropic transformations in crystal structure of uranium*.

Special characters are added to modifiers in the fields of physics and reactor technology; if the modifier does not establish the experimental or theoretical character of a reference, a symbol (E), (T), or (E/T) is placed immediately after the modifying phrase and before the abstract number.

Physical States and Configurations of Materials

The specific physical state, if other than normal, or the geometric configuration of the material is included in the modifying phrase; e.g., information on liquid metals is indexed under the specific metal with "liquid" in the modifier and carbon crucibles is indexed under CARBON with "crucibles" appropriately placed in the modifier. The information on carbon crucibles is also indexed under CRUCIBLES.

Nomenclature of Inorganic Compounds

Inorganic compounds are indexed to subject headings in the direct form. Cations are listed in the order of increasing valence for polycationic salts of varying valence and in alphabetic order for polycationic salts in which all cations have the same valence; e.g., SODIUM ALUMINUM SILICATES and POTASSIUM SODIUM SULFATES. If more than one anionic species is presented in a compound, the anions are arranged alphabetically, e.g., BERYLLIUM FLUORIDE HYDROXIDES.

Nomenclature of Organic Compounds

Complex organic compounds occur in such variety as to require a systematic nomenclature. Headings for most substituted organic compounds are found in an inverted form following names of the parent compounds. Examples of two such headings are:

2,4-HEPTANEDIONE
2,4-HEPTANEDIONE, 1-PHENYL-

Headings are included for broad class names of enzymes, e.g., HYDROLASES and TRANSFERASES. Specific enzyme names are to be found in modifiers under these headings. Similarly, broad class names for hormones are included as headings, except in this case the headings are entered in the inverted form as, e.g., HORMONES, ADRENAL and HORMONES, SEX. The more specific names of hormones are to be found in the modifiers under the appropriate broad headings.

For compounds used as drugs, an effort is made to use subject headings that reflect the recommendations of the USAN (United States Adopted Names) compilation or the "New Names" columns of the *Journal of the American Medical Association*. The *Merck Index* has also been used in selecting subject headings for drugs.

Nomenclature of Life Sciences

Except for those in the phylum Tracheophyta, subject headings for plants are common names. For example, ALGAE, FUNGI, SLIME MOLDS, MOSSES, and LIVERWORTS are subject headings. The phylum name TRACHEOPHYTA and several classes and subclasses belonging to it are entered as subject headings. The specific classes entered are PSILOPSIDA, LYCOPSIDA, SPHENOPSIDA, and PTEROPSIDA. The three subclasses FILICINIDAE, GYMNOSPERMIDAE, and ANGIOSPERMIDAE which belong under Pteropsida are also subject headings. Scientific names (genus and species) and common names if known are used in the modifying phrase.

Subject headings for microorganisms are VIRUSES, RICKETTSIA, and BACTERIA. The name of the type or kind of virus or the genus and species names of the organisms should appear in the modifying phrase.

The names of animal phyla are subject headings. The arthropods are also indexed to subject headings for classes

INSECTA, CRUSTACEA, ARACHNIDA, DIPLOPODA, and CHILOPODA. The invertebrate chordates are indexed to subject headings UROCHORDATA and CEPHALOCHORDATA. The vertebrates are indexed to headings for the classes AGNATHA, CHONDRICHTHYES, OSTEICHTHYES, AMPHIBIA, REPTILIA, AVES, and MAMMALIA as well as to the general heading VERTEBRATA. Common names are used as subject headings in the class Mammalia.

Subject headings are entered for various classes of commodities, and specific products are identified in the file-word position of the modifier. In the following examples, approved subject headings are capitalized and typical modifier file words are listed in lower case.

Example 1

CEREALS
bread
flour
oats
rice
rye
wheat

Example 2

MEATS
beef
fish
mutton
pork
poultry
seafood

Diseases, pathological conditions, and in vitro studies of atypical cells and tissues are indexed under subject headings for the organism affected, under the appropriate anatomical structure of physiological phenomenon affected, and under the appropriate disease heading. Subject headings for diseases (defined as any deviation from the norm) are common names such as ANEMIA, FIBROSIS, INFECTIONS, LEUKEMIA, NEOPLASMS, and RADIATION SICKNESS. Also, the general headings DISEASES and DISEASES, PLANTS are available as subject headings.

Multicomponent Alloys and Other Systems

Multicomponent alloys and systems, except those with well-known trade names, are indexed under a subject heading for each component in the form "element systems" for nonmetals or "element alloys and systems" for metals. The modifier begins with element symbols for each component arranged alphabetically. In addition, if significance is placed on the base metal of the alloy, this is indicated by adding "-base" to the element symbol.

Example: A 1.5C, 12Cr, 0.25V, 0.8Mo, bal Fe alloy is indexed as

CARBON SYSTEMS
C--Cr--Fe--base--Mo--V, ...
CHROMIUM ALLOYS AND SYSTEMS
C--Cr--Fe--base--Mo--V, ...
IRON ALLOYS AND SYSTEMS
C--Cr--Fe--base--Mo--V, ...
MOLYBDENUM ALLOYS AND SYSTEMS
C--Cr--Fe--base--Mo--V, ...
VANADIUM ALLOYS AND SYSTEMS
C--Cr--Fe--base--Mo--V, ...

Well-known alloys, such as ZIRCALOY-2, are indexed under the trade name; cross references from each component of the alloy to the trade name appear in cumulative indexes.

Multicomponent systems in which nonmetallic elements are present in large percentage and which are commonly referred to as systems or mixtures are indexed in a manner similar to alloys.

Example: A tungsten-silicon preparation is indexed as

TUNGSTEN ALLOYS AND SYSTEMS
Si--W, ...
SILICON SYSTEMS
Si--W, ...

Multicomponent systems and mixtures of inorganic compounds are indexed under the subject heading for each component compound; the modifiers begin with an alphabetical listing of compound formulas.

Example: A lead titanate-lead zirconate mixture is indexed as

LEAD TITANATES
PbTiO₃--PbZrO₃
LEAD ZIRCONATES
PbTiO₃--PbZrO₃

Mixtures of organic compounds are indexed under subject headings for each component of the mixture; the modifiers begin with an alphabetical listing of compound names.

Example:

ACETIC ACID
acetic acid--BF₃, ...
BORON FLUORIDES
acetic acid--BF₃, ...

Example:

BENZENE, CHLORO-
bromophenol--chlorobenzene, ...
PHENOL, BROMO-
bromophenol--chlorobenzene, ...

A mixed system of compounds and elements is indexed by a combined form of the preceding conventions; e.g., a dispersion of copper and alumina would be indexed as

ALUMINUM OXIDES
Al₂O₃--Cu, ...
COPPER ALLOYS AND SYSTEMS
Al₂O₃--Cu, ...

If modifiers for alloys and systems are included with other modifiers under the same subject heading, they are filed last.

Example:

CARBON SYSTEMS
deposition on nuclear fuel
particles
molecular structure of
neutron total cross section for
sorptive properties for xenon
Young's modulus for
Al--C--Cr, superconductivity in
B--C, phase studies for
C--U--Zr, fabrication and
physical properties of

Nomenclature of High Energy Physics

Information on particle-particle events is indexed under subject headings for the specific particles involved. The modifier file word *interactions* is followed by the specific interaction written in symbolic form. Particle nomenclature and symbols follow that of the Particle Data Group published in *Review of Particle Properties* (latest revision). For consistency the incident particle is always written first. The mass (in MeV) of a

resonance state is included in parentheses after the symbol to avoid ambiguity. Hyperfragments are indicated in modifiers as in the following examples: Λ -He⁵, Σ -H³, $\Lambda\Lambda$ -Be¹⁰. The superscript on the right indicates the total number of baryons in the composite system.

Examples:

MESONS (π^-)
interactions $\pi^- + p + \Lambda + K^+ + \pi^-$ at 6 GeV/c in
hydrogen bubble chamber, AK(1700) enhancement
in, (E)
BARYONS, RESONANCE (Δ)
production of $\Delta^{++}(1236)$ and $\Delta^0(1236)$ from $K^- +$
 $p + K^- + p + \pi^+ + \pi^-$ interactions at 2.63 and
2.70 GeV/c, (E)
HYPERNUCLEI
binding energy of $\Lambda\Lambda$ -He⁶, three-body $\pi^- \Lambda^- \Lambda^-$
model for

Subject headings are also provided for specific models and theories.

Examples:

ELEMENTARY PARTICLE MODELS
quark, hadron total cross sections in,
numerical analysis of
CURRENT ALGEBRA
PCAC hypothesis for A_1 and rho dominance in $\pi^- +$
 $\pi^+ + \nu + \gamma$ decay
REGGE POLE THEORY
predictions for polarization in $K^- + p + \text{anti}$
 $K^0 + n$, $\pi^- + p + \pi^0 + n$, and $\pi^- + p + \pi^+ + n$
interactions at high energies
SYMMETRY GROUPS
SU(2) x SU(4) intermediate-coupling model for
nucleon and hyperon isobars

Nomenclature of Reactor Technology

Subject headings for reactors are of three types:

(1) Specific reactors, e.g., ARGONNE RESEARCH REACTOR, SAXTON POWER REACTOR

(2) Reactors having a common physics or engineering characteristic, e.g., REACTORS, FAST; REACTORS, ORGANIC-COOLED; REACTORS, PRESSURIZED-WATER

(3) Reactors having a common purpose, e.g., REACTORS, POWER; REACTORS, RESEARCH

Information on specific reactors, type (1), is indexed under the one specific subject heading. Information on classes or types of reactors, types (2) (characteristics) and (3) (purpose), is indexed under all applicable headings. For example, information on a heavy-water-moderated organic-cooled power reactor is indexed under REACTORS, HEAVY WATER-MODERATED; REACTORS, ORGANIC-COOLED; and REACTORS, POWER.

Subject headings exist for selected reactor components, e.g., REACTOR FUEL ELEMENTS. When applicable, headings for components are indexed even though they may have been included in a modifier under a type (2) or (3) heading. The materials in reactor components are indexed when the information is peculiar to the material.

Modifier file words under reactor headings are of two primary types: one type expresses reactor characteristics, e.g., reactivity; the other, components, e.g., control rod. If no subject heading appears for a reactor component, the heading REACTORS is used, and the component is used as the modifier file word.

ALUMINUM ISOTOPES Al-27

- energy level at 1.013 MeV from γ resonance scattering, lifetime of, (E), 26:46905
- energy-level transitions in, multipole mixing ratios for, (E), 26:46893
- neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909
- neutron reactions (n,p) with, calculation of excitation functions for fast, 26:47066
- oxygen-16 reactions at 50 to 170 MeV, calculation of fusion excitation function using equilibrium model with fission competition, (T), 26:47072
- photon reactions (γ ,n) and (γ ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
- photon reactions at 1.160 GeV at 150°, photoemission study of kinetic energy and total yield of charged pions from, (E), 26:46441
- proton reactions (p,n) at 20.3 MeV, DWBA analysis of differential asymmetry from polarized, p. 129-3, (E), 26:46873(R) (CEA-N-1522)
- transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889

ALUMINUM ISOTOPES Al-28

- beta decay of polarized, asymmetry of, (E), 26:46911
- nuclear magnetic moment from NMR measurements, (E), 26:46911
- nuclear magnetic resonance in aluminum phosphides, (E), 26:46911

ALUMINUM OXIDES

- compaction of powdered, electrodynamic method for, 26:45951
- density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
- fabrication of films of, from foil, 26:45931 (KAPL-3164)
- properties for use as thermoluminescent dosimeter, 26:45628
- radiation effects on elasticity and swelling of, fast neutron, 26:45702(R) (LA-4978)
- surface energy of, wetting studies on critical, 26:45944
- Al₂O₃--CaO--Cr₂O₃--F--MgO--MoO₃--TiO₂--ZrO₂, use as welding flux, 26:45941(P)
- Al₂O₃--Fe₂O₃, phase studies on catalysts of, 26:45963
- Al₂O₃--HgO, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
- Al₂O₃--Mo, brittle--ductile transition in dispersions of, transition temperatures for, 26:45993(T) (ORNL-tr-2377)
- Al₂O₃--MoO₃, phase studies on, 26:45963
- Al₂O₃--Sm₂O₃, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
- Al₂O₃--Y₂O₃, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949

ALUMINUM PHOSPHATES

- effects on uptake and retention of calcium-47, radium-226, and strontium-85 by mice following gastric tube administration, 26:45918

ALUMINUM PHOSPHIDES

- aluminum-28 NMR in, (E), 26:46911

AMCHITKA ISLAND AREA

- geology and hydrology of, effects of Cannikin Event on, (E), 26:45488 (USGS-474-148)
- ground motion effects on archaeological sites in, effects of Cannikin Event-generated, (E), 26:45485 (SC-RR-720359)
- seismicity during 1969 to 1972, effects of Cannikin and Milrow Events on, (E), 26:45479 (NWD-746-122)

AMERICIUM

- complexes, identification of octahedral AmCl₆³⁻, 26:45376
- spectra of trivalent, in ethanol--hydrochloric acid solutions, 26:45376
- AMERICIUM ISOTOPES Am-241
- use as radiation source for wire thickness measurement, 26:45717
- use as source in evaluation of Freon 13 fire extinguisher operation by gamma attenuation, 26:45712

AMERICIUM ISOTOPES Am-243

- neutron absorption resonance integral at 0.525 to 1000 eV, (E), 26:47010 (ANCR-1060)
- neutron reactions at 0.5 to 1000 eV, resonance parameters, total cross sections, and transmission for, (E), 26:47010 (ANCR-1060)

AMINES

- preparation of alkaline derivatives of secondary, for use in isotopic exchange reactions, 26:45235(P)

AMMETERS

- design and performance of solid state rms, using metal foil with liquid crystal coating, 26:47634 (UCRL-73587)

AMMONIA

- equation of state for liquid, 26:46479(T) (UCRL-Trans-10622)
- positron lifetime in frozen aqueous solutions of, effects of temperature and positronium trapping on, (E), 26:46362 (RISO-254)
- reactions with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
- recovery as by-product from uranium dioxide production, method for, 26:45371(P)
- butadiene--NH₃, radiolysis of mixtures of, yields of hydrogen from γ , 26:45332
- ethylene--NH₃, radiolysis of mixtures of, yields of

67 SUBJECTS

ANGIOSPERMIDAE

- hydrogen from γ , 26:45332
- NH₃--propene, radiolysis of mixtures of, yields of hydrogen from γ , 26:45332
- AMMONIUM ACID FLUORIDES
- recovery as by-product from uranium dioxide production, method for, 26:45371(P)
- AMMONIUM IONS
- reactions in mass spectrometer, test of clustering in radiolysis by, 26:45353
- stability of solvated, 26:45353
- AMPHIBIA
- Ambystoma maculatum (salamander), radiation effects on migration of, gamma, 26:45815(R) (COD-3486-1)
- Ambystoma opacum (salamander), radiation effects on migration of, gamma, 26:45815(R) (COD-3486-1)
- Rana esculenta (frog), radiation effects on localization of acetylcholinesterase in muscle and neuromuscular junction of, gamma, 26:45823
- Rana pretiosa (frog), radiation effects on developing, chronic cesium-134, 26:45913(R) (RLD-2227-I-8-3)
- Taricha granulosa (newt), radioisotope distribution and retention in, 26:45913(R) (RLD-2227-I-8-3)
- AMPLIFIERS
- circuits for fast, p. 151-2, 26:45873(R) (CEA-N-1522)
- circuits for, p. 79-80, 26:45873(R) (CEA-N-1522)
- description of differential MOSFET, p. 238-40, 26:46873(R) (CEA-N-1522)
- ANALYTICAL CHEMISTRY
- book: Masking and Demasking of Chemical Reactions, Theoretical Aspects and Practical Applications, 26:45188
- book: Modern Analytical Techniques for Metals and Alloys, Part 2, 26:45187
- methods at Japanese Materials Testing Reactor, 26:45182 (JAERI-M-4594)
- secondary ion emission methods in, review on, 26:45249
- ANESTHETICS
- effects of methoxyflurane, on fluorine metabolite concentrations and renal clearance in man, 26:47645 (UR-3490-107)
- ANGIOSPERMIDAE
- Artemisia sp. (sagebrush), revegetation at Nevada Test Site following destruction by fallout, 26:45763 (TID-25954, pp 87-101)
- Artemisia sp. (sagebrush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
- Artemisia tridentata (sagebrush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
- Astragalus sp. (desert plant), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
- Atriplex canescens (salt bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
- Atriplex sp. (salt bush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
- Dianthus caryophyllus (carnation), radiation effects on cuttings of, effects of yeast extract on recovery from γ , 26:45777
- Ephedra nevadensis (mormon tea), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
- Ephedra nevadensis (mormon tea), radiation effects on growth and reproduction of, chronic γ , 26:45756 (TID-25954, pp 5-11)
- Ephedra sp. (mormon tea), ecology at Nevada Test Site, 26:45764 (TID-25954, pp 102-9)
- Ephedra sp. (mormon tea), radiation effects on survival of, lethal dose for γ , 26:45764 (TID-25954, pp 102-9)
- Franseria dumosa (burr bush or burro bush), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
- Franseria dumosa (burr bush or burro bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
- Franseria dumosa (burr sage or burro bush), ecology at Nevada Test Site, 26:45768 (TID-25954, pp 153-61)
- Franseria dumosa (burr sage or burro bush), radiation effects on growth and survival of, acute γ , 26:45768 (TID-25954, pp 153-61)
- Grayia spinosa (spiny hop sage), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
- Larrea divaricata (creosote bush), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
- Larrea divaricata (creosote bush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
- Larrea divaricata (creosote bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)

Pelagophycus porra, radioisotope uptake and retention by, correlation between content of seawater and polonium-210 and plutonium-239, 26:45924
radioisotope content in saltwater, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)

ALGERIA

tritium content in aquifers in central and northwest, 26:45558

ALGINIC ACID

effectiveness of dietary, for removal of radium-226 from man following oral uptake as radium sulfate dust, 26:45907

ALKALI METALS

hydrogen-atom reactions with, Born-approximation calculations of excitation to 2s, 2p, 3s, 3p, and 3d states in, 26:46403
spectra in presence of rare gases, Fermi pressure shift in, (T), 26:46401

ALKANES

photolysis of solid, ESR spectral studies of far-uv, 26:45323

ALKYLAMINES

mercapto-, radioprotective effects of, on Escherichia coli and mice, 26:45859

ALLOYS

analysis for niobium by atomic absorption spectrophotometry, 26:45234
book: Constitution of Binary Alloys, Second Supplement, 26:45997
book: Modern Analytical Techniques for Metals and Alloys. Part 2, 26:45187
decomposition of supersaturated solid solutions of, effects of aging on mechanism of, 26:46032
grain boundary models for, relations with free volume, 26:46027
radioinduced crystal lattice defects in, review on fast electron, γ , and neutron, 26:46076

ALPHA DECAY

width for, 8-matrix theory for, (T), 26:47068

ALPHA PARTICLES

detection of, etching conditions for nitrocellulose detectors for, 26:45610
detection of, performance of cellulose-based track detectors for, 26:45615
distribution spectra of trapped, in outer radiation belt, June--July 1969, (E), 26:46309
dose to respiratory tract epithelial cells of animals following inhalation of radon-222 aerosols, relation of content of daughter products in air to, 26:45861
effects on CO₂ lasers, 26:45669
effects on survival of cultured mammalian cells, relation of mathematical analysis of, to radiotherapy, 26:45735
reactions ($\alpha, 2n\gamma$) with thulium-169 at 28 MeV, γ -ray angular distributions and excitation functions for, (E), 26:46995
reactions ($\alpha, 2n\gamma$) with thulium-169, γ spectra from, (E), 26:46980
reactions ($\alpha, 2n\gamma$) with thulium-169, γ - γ coincidence spectra from, (E), 26:46987
reactions (α, γ) with carbon-12 near 8 MeV, detailed balance test of time reversal invariance in, (E), 26:46881
reactions ($\alpha, He-3$) with tantalum-181, gold-197, and lead-208 up to 80 MeV, energy spectra and cross sections for, (E), 26:47008
reactions ($\alpha, n\gamma$) with oxygen-18 at 4.9 to 11.5 MeV, γ -ray angular distributions and γ - γ and n - γ angular correlations for, (E), 26:46907
reactions (α, p) with iron-56, cobalt-59, nickel-60, and nickel-62 at 17 MeV, proton spectra from, (E), 26:46932
reactions (α, p) with nickel-58, -60, -62, and -64 at 19.3 MeV, zero-range DWBA analysis of angular distributions for, (E), 26:46934
reactions with natural chromium, production of carrier-free manganese-54 from, 26:45394
scattering by iron-56, cobalt-59, nickel-60, nickel-62, copper-63, and copper-65 at 17 MeV, α spectra from inelastic, (E), 26:46932
scattering by lithium-6 at 36.6 MeV, angular distribution for elastic, (E), 26:46883
scattering by nickel-58 at 30 MeV, DWBA analysis of differential cross sections and spectrum from elastic, (E), 26:46927
scattering by tantalum-181, gold-197, and lead-208 up to 80 MeV, energy spectra and cross sections for inelastic, (E), 26:47008
scattering by tin-117 and -119 at 10, 10.5, and 11 MeV, γ -ray angular distributions and spectra for, (E), 26:46961

ALPHA SOURCES
determination of low-activity, recommendations for, 26:45261 (ICRU-22)
handling of high-activity, description of dual-mode operation hot cell for, 26:46107 (UCRL-73756)
release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)

ALUMINUM
deposition on niobium--titanium wires by extrusion, parameters for, 26:45972(R) (COO-3311-1)

determination by fluorometry using Superchrome Garnet Y, 26:45236
determination in boreholes, neutron activation logging method for, 26:45567
determination in coal and coal ash by neutron activation, 26:45195
determination in high-purity cadmium oxide, spectrographic, 26:45220 (BARC-593)
determination of, chemical procedures for, 26:45181 (CISE-R-172)
diffusion and outgassing of ion implanted-helium in, effects of temperature on, 26:46050 (CONF-710601-, pp 230-47)
dissolution in iron, 26:46007
effects on mechanical properties of molybdenum--uranium-base alloys at 20 to 640°C, 26:45995 (UJV-2802-M)
electron transport in, discrete ordinates method for 1- to 8-MeV, (E/T), 26:47210
electron--phonon umklapp interaction at low temperatures in, (E), 26:47264
positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
prospecting for deposits of, pulsed neutron well logging method for, 26:45559(P)
proton reactions at 24 GeV/c, w , K, p, and anti p spectra from, (E), 26:46459
proton reactions with, Born-approximation calculations of energy losses and L-shell ionization in, 26:46393
radiation effects on internal friction of wires of, at room temperature, neutron, 26:46073
radiation effects on length of, γ , 26:46075
radioinduced defect aggregates in, relations of temperature with fast neutron, 26:46060 (CONF-710601-, pp 672-83)
radioinduced defects in single crystals of, study of I/sub B/ stage in annealing of electron, 26:47229
recrystallization of, effects of impurities on, 26:45982
separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281
separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
solute segregation and stresses around growing voids in, during cooling from high-temperature anneals, 26:45988 (CONF-710601-, pp 630-46)
transitions of superconducting, resistive, (E/T), 26:46817

ALUMINUM ALLOYS AND SYSTEMS

analysis for tin by alternating current polarographic method, 26:45256
Al--Ce--La, critical field curves of Kondo superconducting, 26:46832
Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009
Al--Cr--Ni--ThO₂, properties of dispersion-strengthened, for use in space shuttle, 26:45934 (N-72-18585)
Al--Fe--Mo--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Al--Fe--Nb--Si, dissolution in, 26:46007
Al--Fe--Si--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Al--Fe--U-base, mechanical properties of, at 20 to 640°C, effects of chromium and silicon additions on, 26:45995 (UJV-2802-M)
Al--Fe--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Al--Fe--Y, phase studies on, 26:46005
Al--Ge--Nb, critical temperature of superconducting intermetallic Nb₃Al/sub x/Ge/sub 1-x/, variations in, 26:46021
Al--Ge--Nb, phase diagram of niobium corner of, 26:46031
Al--Ge--Nb, superconducting properties of, 26:46031
Al--Mg, grain refining of, mechanism and method for, 26:45984
Al--Mo--Si--U, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Al--Nb, critical currents in superconducting Nb₃Al, (E), 26:46829
Al--Nb, critical temperature of superconducting intermetallic Nb₃Al, variations in, 26:46021
Al--Nb, phase transformations and decomposition in superconducting, effects of heat treatments on, 26:46020
Al-base--Co, superconductivity and localized spin fluctuations in, (E/T), 26:46838
Al-base--Cr, superconductivity and localized spin fluctuations in, (E/T), 26:46838
Al-base--Fe, superconductivity and localized spin fluctuations in, (E/T), 26:46838
Al-base--Ti, superconductivity and localized spin fluctuations in, (E/T), 26:46838
Al-base--U, fission fragment ranges in, 26:46074
Al-base--V, superconductivity and localized spin fluctuations in, (E/T), 26:46838

ALUMINUM ISOTOPES

photon reactions (γ, n^0) at 3.5 to 8.3 GeV, A dependence of, 26:46447

ALUMINUM ISOTOPES Al-25

energy levels from magnesium-24 (p, γ) reactions, lifetimes of, (E), 26:46910
energy-level transitions from magnesium-24 (p, γ) reactions, branching and mixing ratios for, (E), 26:46910

ALUMINUM ISOTOPES Al-27

energy level at 1.013 MeV from γ resonance scattering, lifetime of, (E), 26:46905
 energy-level transitions in, multipole mixing ratios for, (E), 26:46893
 neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909
 neutron reactions (n,p) with, calculation of excitation functions for fast, 26:47066
 oxygen-16 reactions at 50 to 170 MeV, calculation of fusion excitation function using equilibrium model with fission competition, (T), 26:47072
 photon reactions (γ ,n) and (γ ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
 photon reactions at 1.160 GeV at 1500, photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441
 proton reactions (p,n) at 20.3 MeV, DWBA analysis of differential asymmetry from polarized, p. 120-3, (E), 26:46873(R) (CEA-N-1522)
 transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889

ALUMINUM ISOTOPES Al-28

beta decay of polarized, asymmetry of, (E), 26:46911
 nuclear magnetic moment from NMR measurements, (E), 26:46911
 nuclear magnetic resonance in aluminum phosphides, (E), 26:46911

ALUMINUM OXIDES

compaction of powdered, electrodynamic method for, 26:45951
 density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
 fabrication of films of, from foil, 26:45931 (KAPL-3164)
 properties for use as thermoluminescent dosimeter, 26:45628
 radiation effects on elasticity and swelling of, fast neutron, 26:45702(R) (LA-4978)
 surface energy of, wetting studies on critical, 26:45944
 Al₂O₃--CaO--Cr₂O₃--F--MgO--MoO--TiO₂--ZrO₂, use as welding flux, 26:45941(P)
 Al₂O₃--Fe₂O₃, phase studies on catalysts of, 26:45963
 Al₂O₃--MgO, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
 Al₂O₃--Mo, brittle--ductile transition in dispersions of, transition temperatures for, 26:45993(T) (ORNL-tr-2377)
 Al₂O₃--MoO₃, phase studies on, 26:45963
 Al₂O₃--Sm₂O₃, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
 Al₂O₃--Y₂O₃, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949

ALUMINUM PHOSPHATES

effects on uptake and retention of calcium-47, radium-226, and strontium-85 by mice following gastric tube administration, 26:45918

ALUMINUM PHOSPHIDES

aluminum-28 NMR in, (E), 26:46911

AMCHITKA ISLAND AREA

geology and hydrology of, effects of Cannikin Event on, (E), 26:45488 (USGS-474-148)
 ground motion effects on archaeological sites in, effects of Cannikin Event-generated, (E), 26:45485 (SC-RR-720359)
 seismicity during 1969 to 1972, effects of Cannikin and Milrow Events on, (E), 26:45479 (NVD-746-122)

AMERICIUM

complexes, identification of octahedral AmCl₆³⁻, 26:45376

spectra of trivalent, in ethanol--hydrochloric acid solutions, 26:45376

AMERICIUM ISOTOPES Am-241

use as radiation source for wire thickness measurement, 26:45717
 use as source in evaluation of Freon 13 fire extinguisher operation by gamma attenuation, 26:45712

AMERICIUM ISOTOPES Am-243

neutron absorption resonance integral at 0.625 to 1000 eV, (E), 26:47010 (ANCR-1060)
 neutron reactions at 0.5 to 1000 eV, resonance parameters, total cross sections, and transmission for, (E), 26:47010 (ANCR-1060)

AMINES

preparation of alkaline derivatives of secondary, for use in isotopic exchange reactions, 26:45289(P)

AMMETERS

design and performance of solid state rms, using metal foil with liquid crystal coating, 26:47634 (UCRL-73587)

AMMONIA

equation of state for liquid, 26:46079(T) (UCRL-Trans-10622)
 positron lifetime in frozen aqueous solutions of, effects of temperature and positronium trapping on, (E), 26:46362 (RISO-254)
 reactions with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
 recovery as by-product from uranium dioxide production, method for, 26:45371(P)
 butadiene--NH₃, radiolysis of mixtures of, yields of hydrogen from γ , 26:45332
 ethylene--NH₃, radiolysis of mixtures of, yields of

67 SUBJECTS

ANGIOSPERMIDAE

hydrogen from γ , 26:45332

NH₃--propene, radiolysis of mixtures of, yields of hydrogen from γ , 26:45332

AMMONIUM ACID FLUORIDES

recovery as by-product from uranium dioxide production, method for, 26:45371(P)

AMMONIUM IONS

reactions in mass spectrometer, test of clustering in radiolysis by, 26:45353
 stability of solvated, 26:45353

AMPHIBIA

Ambystoma maculatum (salamander), radiation effects on migration of, gamma, 26:45815(R) (COO-3486-1)
 Ambystoma opacum (salamander), radiation effects on migration of, gamma, 26:45815(R) (COO-3486-1)
 Rana esculenta (frog), radiation effects on localization of acetylcholinesterase in muscle and neuromuscular junction of, gamma, 26:45828
 Rana pretiosa (frog), radiation effects on developing, chronic cesium-134, 26:45913(R) (RLO-2227-T-8-3)
 Taricha granulosa (newt), radioisotope distribution and retention in, 26:45913(R) (RLO-2227-T-8-3)

AMPLIFIERS

circuits for fast, p. 151-2, 26:46873(R) (CEA-N-1522)
 circuits for, p. 79-80, 26:46873(R) (CEA-N-1522)
 description of differential MOSFET, p. 238-40, 26:46873(R) (CEA-N-1522)

ANALYTICAL CHEMISTRY

book: Masking and Demasking of Chemical Reactions. Theoretical Aspects and Practical Applications, 26:45188
 book: Modern Analytical Techniques for Metals and Alloys. Part 2, 26:45187
 methods at Japanese Materials Testing Reactor, 26:45182 (JAERI-M-4594)
 secondary ion emission methods in, review on, 26:45249

ANESTHETICS

effects of methoxyflurane, on fluorine metabolite concentrations and renal clearance in man, 26:47645 (UR-3490-107)

ANGIOSPERMIDAE

Artemisia sp. (sagebrush), revegetation at Nevada Test Site following destruction by fallout, 26:45763 (TID-25954, pp 87-101)
 Artemisia sp. (sagebrush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
 Artemisia tridentata (sagebrush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
 Astragalus sp. (desert plant), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
 Atriplex canescens (salt bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
 Atriplex sp. (salt bush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
 Dianthus caryophyllus (carnation), radiation effects on cuttings of, effects of yeast extract on recovery from γ , 26:45777
 Ephedra nevadensis (mormon tea), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
 Ephedra nevadensis (mormon tea), radiation effects on growth and reproduction of, chronic γ , 26:45756 (TID-25954, pp 5-11)
 Ephedra sp. (mormon tea), ecology at Nevada Test Site, 26:45764 (TID-25954, pp 102-9)
 Ephedra sp. (mormon tea), radiation effects on survival of, lethal dose for γ , 26:45764 (TID-25954, pp 102-9)
 Franseria dumosa (burr bush or burro bush), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
 Franseria dumosa (burr bush or burro bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)
 Franseria dumosa (burr bush or burro bush), ecology at Nevada Test Site, 26:45768 (TID-25954, pp 153-61)
 Franseria dumosa (burr sage or burro bush), radiation effects on growth and survival of, acute γ , 26:45768 (TID-25954, pp 153-61)
 Grayia spinosa (spiny hop sage), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
 Larrea divaricata (creosote bush), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)
 Larrea divaricata (creosote bush), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
 Larrea divaricata (creosote bush), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)

Larrea sp. (creosote bush), survival at Nevada Test Site, relation of radioresistance to, 26:45765 (TID-25954, pp 115-32)

Lycium andersonii (desert shrub), radiation effects on growth and reproduction of, chronic γ , 26:45756 (TID-25954, pp 5-11)

Lycium andersonii (desert shrub), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)

Lycium pallidum (desert shrub), radiation effects on seeds of, relation of dose and moisture content to germination and seedling survival, 26:45761 (TID-25954, pp 73-4)

Lycium sp. (desert shrub), radiation effects on growth and reproduction of, chronic γ , 26:45757 (TID-25954, pp 12-48)

Lycium sp. (desert shrub), survival at Nevada Test Site, relation of radiosensitivity to, 26:45766 (TID-25954, pp 133-46)

Lycopersicon sp. (tomato), radiation effects on seed of, effects on rate of germination and plant growth, effects of dose and moisture on neutron, 26:45773

Oryzopsis hymenoides (range grass), strontium-90 uptake and distribution in tissues of, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)

Pisum sp. (pea), radiation effects on survival of, effects of genotype on gamma, 26:45775

Pisum sp. (pea), radioinduced mutations in, review of, 26:45776

Salsola pestifera (desert plant), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)

Salsola sp. (Russian thistle or tumbleweed) ecology at Nevada Test Site, 26:45767 (TID-25954, pp 147-52)

Salsola sp. (Russian thistle or tumbleweed), radiation effects on seeds of, dose dependence of germination and seedling growth, γ , 26:45767 (TID-25954, pp 147-52)

Solanum tuberosum (potato), radioinduced mutations in, for protein and starch contents and yield, fast neutron and γ , 26:45770

Triticum sp. (wheat), radioisotope content of, in France during 1960 to 1969, cesium-137 and strontium-90, 26:45903

Triticum sp., strontium-90 accumulation in, from Podzolic soils, effects of calcium on, 26:45550

Yucca brevifolia (yucca), radiation effects on growth and survival of, following seed exposure, dose dependence of γ , 26:45758 (TID-25954, pp 49-53)

Yucca sp. (yucca), radiation effects on growth of, dose dependence of acute γ , 26:45769 (TID-25954, pp 188-98)

Yucca sp. (yucca), survival at Nevada Test Site, relation to radiosensitivity, 26:45769 (TID-25954, pp 188-98)

Zea mays (maize), rubidium-86 uptake by, effects of calcium oxide and potassium content of soil on, 26:45925

ANILINE
radiolysis of aqueous solutions of pure and substituted, yield of cyclohexadienyl-type radical from pulse, 26:45309(R) (AE-453)

ANIMAL FEEDS
radiation processing of, effects on rats following ingestion, 26:45899

ANIMALS
analysis for cesium by neutron activation, 26:45193 (TID-25954, pp 376-8)
analysis for potassium by potassium-40 γ spectroscopy, 26:45193 (TID-25954, pp 376-8)
radiation dose to respiratory tract epithelial cells of, following inhalation of radon-222 aerosols, relation of content of daughter products in air to α , 26:45861
radioinduced injuries in, following chronic low dose internal exposure, repair mechanisms at cellular and molecular level for, 26:45911 (COO-3013-25)
radioinduced neoplasms in, following inhalation or injection of plutonium-239, dose and time dependence of, 26:45888 (CONF-720520-1)
shelters for raising small farm, use of power plant waste heat for heating, 26:45576

ANNEALING
mechanisms of dislocation loops and tetrahedra in metals, 26:45989 (CONF-710601-1, pp 647-62)

ANTENNAS
operation of radar, for incoherent scattering experiments in auroral zone, 26:46283

ANTHRACENE
electron reactions with, cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)
radioinduced fluorescence in benzene and cyclohexane solutions, effects of concentration on, 26:45328
radioinduced fluorescence in cyclohexane, effects of scavengers on electron pulse, 26:45331
radiolysis in benzene and cyclohexane solutions, effects of concentration on yield of excited triplet states from pulse, 26:45328

2-ANTHRACENESULFONIC ACID, 9,10-DIHYDRO-9,10-DIOXO-
reactions with hydrated electrons and hydrogen atoms from radiolysis of aqueous solutions, 26:45354

ANTHRONE

reactions with hydrated electrons from pulse radiolysis of aqueous solutions, 26:45354

ANTIMONY

determination in hair by neutron activation and γ spectroscopy, 26:45214
determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
effects on recrystallization of iron, 26:45982
neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)
neutron reactions at 0.05 to 150 eV, cross reactions and resonance parameters for, (E), 26:46946
purification by zone melting in rotating container, 26:45276

ANTIMONY ALLOYS AND SYSTEMS

Sb-Tm, magnetization of TmSb at 1.5 and 20.4°K, anisotropic dc, 26:47258

ANTIMONY ISOTOPES Sb-122

energy levels from neutron resonances in antimony-121, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

ANTIMONY ISOTOPES Sb-124

energy levels from antimony-123 (n, γ) reactions, spins, parities, and widths of, p. 300-2, (E), 26:46873(R) (CEA-N-1522)

energy levels from neutron resonances in antimony-123, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

ANTIMONY ISOTOPES Sb-125

beta decay of, γ - γ angular correlation for, (E), 26:46957

content in atmosphere in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)

content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)

solubility of Rainier Event rubble-contained, measurement of long-term, 26:45482 (NVO-1229-177)

ANTIMONY ISOTOPES Sb-126

beta decay of ground-state and isomeric, γ spectra from, (E), 26:46944

ANTIMONY ISOTOPES Sb-128

beta decay of ground-state and isomeric, γ spectra from, (E), 26:46944

ANTIMONY ISOTOPES Sb-130

beta decay of ground-state and isomeric, γ spectra from, (E), 26:46944

ANTIMONY URANIUM OXIDES

structure of, relation of catalytic activity in oxidation of propylene to acrolein and, 26:45374

ARCHAEOLOGICAL SPECIMENS

damage to Amchitka Island in-situ, from Cannikin Event-generated ground motion, (E), 26:45485 (SC-RR-720359)

ARCTIC OCEAN

trace metal content and distribution in waters and sediments of, during June 1971 to June 1972, 26:47623(R) (RLO-2229-T-1-18)

ARGON

abundance in solar-flare particles, (E), 26:46232

electron reactions in eV range with, trapped-electron measurements of integral cross sections for production of secondary electrons of 0 to 1 eV by ionization in, 26:46390

electron reactions up to 300 eV with, ionization efficiency curves for Ar⁺ and Ar²⁺ in, (E), 26:46374
electron reactions with clusters of, multiple ionization in, (E), 26:46386

heat transfer from partially ionized, effects of transverse magnetic fields on, 26:45464

helium-metastable-atom reactions at 150 to 1600 eV with, absolute cross sections for de-excitation of 2¹S and 2³S states in, (E), 26:46394

helium-metastable-atom reactions at 50 to 600 eV with, cross sections for de-excitation in, (E), 26:46394

helium-metastable-atom reactions at thermal energies with, crossed-beam and gas-cell measurements of cross sections for ionization in, 26:46404

interactions with argon and neon atoms, retarded van der Waals potential for, as function of interatomic distance, 26:46357

neon-metastable-atom reactions at 50 to 600 eV with, cross sections for de-excitation in, (E), 26:46394

plasma of, continuum spectra in near-ultraviolet and visible ranges, 26:47132

plasma of, parallel magnetic multipole confinement of, (E), 26:47111

plasma of, shock wave propagation in expanding discharge, (E/T), 26:47194

proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s, 3p, and 3d hydrogen atoms by charge exchange and excitation in, (E), 26:46378

proton scattering by, reduced cross sections for, (T), 26:46365(R) (ORO-2911-14)

reactions between atoms of, potentials for calculation of average energy going to electronic processes in, 26:46365(R) (ORO-2911-14)

reactions with acetylene at 50 to 600 eV of metastable atoms of, cross sections for de-excitation in, (E),

reactions with helium 2^3S and 2^1S atoms at 10^{-3} to 10^4 eV, semi-empirical calculations of total cross sections for associative and Penning ionization in, 26:46395
 reactions with nitrogen, Monte Carlo trajectory calculations of three-body recombination and dissociation in, 26:46382

ARGON IONS

acceleration to 11.7 GeV in Princeton Synchrotron, 26:46124
 effects of 10.4-MeV/amu, on crystal structure of minerals, 26:46156
 reactions with molybdenum monocrystals at low energies, mechanism of ion capture by surface in, (E/T), 26:46335 (FRNC-TH-198)

ARGON ISOTOPES

abundance in Lost City meteorite, (E), 26:46155

ARGON ISOTOPES Ar-36

helium-3 reactions ($\text{He}-3, t$) at 22.2 MeV, triton spectra and Q-value for, (E), 26:46904
 nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892

ARGON ISOTOPES Ar-37

activity measurements of, miniature low-background G--M or proportional counter for, 26:45614

ARGON ISOTOPES Ar-39

activity measurements of, miniature low-background G--M or proportional counter for, 26:45614
 content ratio to argon-40 in samples, use for age estimation, 26:45557

ARGON ISOTOPES Ar-40

content ratio to argon-39 in samples, use for age estimation, 26:45557

ARGON SYSTEMS

Ar--K, spectra measurements of electron density and temperature in plasma of, 26:47116 (IPP-IV-12)

ARKANSAS NUCLEAR ONE POWER REACTOR

environmental report for, cost-benefit analysis for alternate designs, 26:47486 (DOCKET-50313-44)
 environmental report for, revised and additional data for, 26:47487 (DOCKET-50313-45)
 safety of, revised and requested data on, 26:47488 (DOCKET-50313-46)

ARKANSAS NUCLEAR ONE POWER REACTOR, UNIT 2

effluents from, effects on environment of chemical, radioactive, and thermal, 26:47532 (DOCKET-50368-31)
 operation of, cost-benefit analysis for, 26:47532 (DOCKET-50368-31)
 site for, ecology, geology, hydrology, and meteorology of, 26:47532 (DOCKET-50368-31)

ARSENIC

determination as impurity in vinyl chloride by neutron activation, 26:45204
 determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
 determination of trace amounts of, in environmental samples by neutron activation, 26:45198
 effects on recrystallization of iron, 26:45982

ARSENIC ISOTOPES As-72

energy levels and properties of, compilation of, (E), 26:46939

ARSENIC ISOTOPES As-75

neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)
 neutron reactions (n, α) with, calculation of excitation functions for fast, 26:47066
 neutron reactions (n, p) with, calculation of excitation functions for fast, 26:47066

ARSENIC ISOTOPES As-76

beta decay of, γ spectra from, (E), 26:46938

ASHES

determination in coal by neutron activation determination of silicon and aluminum, 26:45195

ASTATINE ISOTOPES At-210

electron-capture decay of, branching ratio and γ and conversion electron spectra from, (E), 26:47006

ASTRON DEVICE

E-layer simulation in axisymmetric mirror field, (T), 26:47147
 proton E-layer for, properties of, 26:47175(P)

ASTRONAUTS

radiation hazards to, during space flight, cosmic, 26:45812

ATLANTIC OCEAN

fallout content and distribution in silts in, 26:45531 (COO-3563-3)
 fallout content in north, ecological relations for cesium-137, plutonium-238, -239, and -240, and strontium-90, 26:45541 (NYO-2174-130)

ATMOSPHERE

beryllium-7 content in, relations of fission product deposition in oceans to, 26:45530 (BNWL-SA-4205(Rev.))
 contamination by gaseous radioactive wastes, problems associated with, 26:46102(T) (ORNL-tr-2369)
 effects of thermal effluent discharge into large lakes on, 26:45584
 electric phenomena in ground-level, effects on atmospheric natural radioactivity, 26:45524
 fallout content in, Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
 fallout monitoring in surface, at world sites, 26:45512(R) (HASL-257(App.))
 fluorescence due to large cosmic showers, detection of,

(E), 26:46150 (NYO-3038-13)

particle diffusion in, zinc sulfide tracer detector for measuring, pp.1,2, 26:45673(R) (UCRL-50007-71-3)
 radioactivity in ground-level, effects of meteorological conditions on natural, 26:45524
 radioactivity monitoring of ground-level California, during Feb. 1972, gross beta, 26:45523
 radioactivity monitoring of ground-level, in Canada, South America, and US during Feb. 1972, gross beta, 26:45523
 radioactivity monitoring of, in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
 radioactivity of aerosols in Polish, relations with aerosol dust content, (E), 26:45521
 radioisotope content in Argentina, 1960 through 1970, cesium-137 and strontium-90, 26:45520 (NP-19357)
 radioisotope content in western US, measurement of cesium-137, plutonium-239, and strontium-90, 26:45517 (HASL-257, pp I.18-32)
 radon distribution in, use for calculating turbulent diffusion coefficient and vertical component of wind velocity, 26:45522
 sulfur content in, isotope ratio tracer method for determining source of, 26:45568

ATMOSPHERES

control of gaseous, in industrial irradiation facility, design of equipment for, 26:45448(P)

ATMOSPHERIC PRECIPITATION

radioactivity monitoring in Canadian, South American, and US, during Feb. 1972, gross beta, 26:45523
 strontium-90 content in western US, 26:45517 (HASL-257, pp I.18-32)

ATOMIC ABSORPTION SPECTROMETRY

development for trace amounts of metals in atmospheric particulate, 26:45244
 review on, 26:45251
 sensitivity of air--acetylene and nitrous oxide--acetylene flames in, 26:45228

ATOMIC ENERGY

bibliographies on, April--June, 1972, 26:47628 (TID-3767)

developments in India, 26:47300

ATOMIC ENERGY, LEGAL ASPECTS

legislation on maximum permissible radiation doses to personnel in Federal Republic of Germany, 26:45803
 licensing of U.S. nuclear power plants, proposed procedures for, 26:47571

ATOMIC PHYSICS

bibliography on surface and vacuum physics, 26:46336

ATOMIC STRUCTURE

symmetry for, (T), 26:46508

ATOMS

deuteron reactions with, classical calculations of cross sections for single-electron capture into ground and excited states in, 26:46398
 electron correlation in, analytical evaluation of integrals in calculation of, 26:46414
 electron pair correlation energies in, derivation from total correlation energies, (T), 26:46412
 electron reactions with, bound states in quantum electrodynamics for, (T), 26:46721
 electron reactions with, third-derivative method for determination of ionization and appearance potentials for, (E/T), 26:46385
 electron scattering by, average inelastic channel and collective modes of excitation in, 26:46389
 electron scattering by, optimization of basis set in calculation of amplitudes for, 26:46389
 energy of, decoupled Hartree--Fock method for calculation of, 26:46416
 exchange energy in, statistical calculation of retardation in relativistic Hartree--Fock equations for, 26:46408
 excitation energy of, decoupled Hartree--Fock method for calculation of, 26:46416
 expectation values of single-particle operators for, in Hartree approximation, calculations using Delves' variational principle, 26:46417
 hyperfine structure of, correlation corrections to expectation values for, 26:46409
 interactions between, inversion method for calculation of potentials for, from scattering cross sections, 26:46387
 ionization potentials of, decoupled Hartree--Fock method for calculation of, 26:46416
 photon reactions with, bound states in quantum electrodynamics for, (T), 26:46721
 polarizability of, coupled multicoufigurational method for calculation of dipole, 26:46419
 polarizability under perturbation having imaginary frequency, variational bounds for, (T), 26:46415
 properties of, infinite-order perturbation theory for calculation of, 26:46410
 proton reactions with, classical calculations of cross sections for single-electron capture into ground and excited states in, 26:46398
 radiation emission from systems of N two-level, Green-function calculation of cooperative effects in spontaneous, 26:46418

- radiation emission from systems of two-level, time evolution and direction of cooperative, (T), 26:46411
 reactions with atoms and molecules, Monte Carlo trajectory calculations of three-body recombination and dissociation in, 26:46382
 reactions with atoms, bound states in quantum electrodynamics for, (T), 26:46721
 reactions with diatomic molecules, comparison of S matrices calculated by various exponential approximations for rotationally inelastic, 26:46369
 reactions with diatomic molecules, optical-model calculations of probabilities for vibrational transitions in, 26:46375
 scattering of, bound states in quantum electrodynamics for, (T), 26:46721
 wave functions for, electron correlation in calculation of, 26:46409
 x-ray emission from filling of K- and L-shell vacancies in $Z = 5$ to 102, calculated rates of, 26:46341 (UCRL-51231)
- AURORAE**
 excitation by protons, $H\beta/N_2 + 4709$ -A intensity ratio and vibrational enhancement of N_2^+ first negative bands in, (E), 26:46276
 particle fluxes in nighttime, Fermi acceleration mechanism for keV, (E/T), 26:46267
- AURORAL ZONE**
 electron precipitation in, association of emission of intense bands of ELF noise at low altitudes with low-energy, (E), 26:46308
 oval observations by all-sky photographs from Alaskan meridian chain of stations, dynamics of, (E), 26:46268
 pulsations in daytime, relation of polar and magnetospheric substorms with Pld irregular, (E), 26:46317
 radar incoherent scattering facility for, operation of, 26:46283
- AUSTRALIA**
 mapping of uranium deposits in, using gamma spectrometry, 26:45507
 radioactive waste management in, from fuel reprocessing, programs for, 26:46098 (AAEC/TM-628)
- AVES**
 cesium-137 content in muscles of partridges, Warsaw, Poland, 1968 to 1969, 26:45548
 Gallus sp. (chicken), radiation effects on infectivity and development of metacercariae in bursa of Fabricius and chorioallantois of, dose dependence of, 26:45879
 Gallus sp., cesium-137 content in muscles of, Warsaw, Poland, 1968 to 1969, 26:45548
- AVIATION PERSONNEL**
 radiation dose to supersonic, measurements on board the Concorde prototype, 26:45796
- B**
- BACTERIA**
 Bacillus subtilis, radioinduced single-strand breaks in transforming DNA of, effectiveness of toluene treatment for enhancing semi-in vitro repair of, 26:45731
 Chlamydomonas reinhardtii, radiation effects on survival of, following high dose rate exposure, effects of prior low dose rate exposure on sensitivity to electron, 26:45754
 Clostridium sp., radioinduced inhibition of sporulation in, in packaged sausage, combined effects of heat treatment on, 26:45747
 Escherichia coli, radiation effects on DNA synthesis in recombination-deficient strains of, 26:45753
 Micrococcus luteus, radioinduced damage to DNA of, role of enzymes in repair of, 26:45722(R) (COO-3232-1)
- BACTERIOPHAGES**
 R17, radiation effects on in vitro coding activity of RNA of, 26:45728
 radioinduced inactivation of extracellular, effects of LET on, heavy ion, and proton, 26:45746
 radioinduced mutations in extracellular, effects of LET on frequency of, heavy ion, and proton, 26:45746
- BARIIUM**
 mass flow choking limit for heat pipe working fluid of, (E), 26:45460
 polarizability of ground-state, electric dipole, (T), 26:46349
 solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
 work function on tungsten(110), temperature dependence of, 26:47257
- BARIIUM IONS**
 implantation in nickel at 1 to 10 keV, penetration and spatial distribution from, 26:47235
- BARIIUM ISOTOPES Ba-130**
 energy level of, model analysis of 8- isomeric, (T), 26:46949
- BARIIUM ISOTOPES Ba-131**
 electron-capture decay of, γ - γ angular correlations in, (E), 26:46963
- BARIIUM ISOTOPES Ba-132**
 energy-level scheme from lanthanum-132 decay, (E), 26:46948
- BARIIUM ISOTOPES Ba-135**
 energy levels of, lifetimes and spin-parity assignments for, (E), 26:46945
- BARIIUM ISOTOPES Ba-140**
 range of fission fragments of, in aluminum-uranium alloys and uranium oxides, 26:46074
 use as tracer to study oxide inclusions in steel, 26:46008
- BARIIUM OXIDES**
 electric, thermionic, and thermoelectric properties of crystals of, 26:47249
 BaO-SiO₂, radioinduced optical absorption and paramagnetic centers in glassy, spectral studies on, 26:47226
- BARIIUM SULFATES**
 BaSO₄-CaSO₄, sorptive properties for strontium, 26:45273 (UJV-2820-Ch)
- BARIIUM TITANATES**
 proton channeling at 3.8 MeV in paraelectric and ferroelectric BaTiO₃ crystals, 26:47234
- BARIIUM URANIUM SELENIDES**
 preparation and structure of BaU₂Se₈, 26:45378
- BARNWELL NUCLEAR FUEL PROCESSING PLANT**
 uranium hexafluoride facility at, environmental studies on, 26:45308 (DOCKET-701327-1)
- BARYONS**
 interactions with mesons, duality properties of Regge cuts in, 26:46600
 interactions with mesons, partial wave relations from fixed-u dispersion relations for, 26:46610
 interactions with mesons, test of equal-phase assumption for helicity amplitudes for, (T), 26:46621
 interactions with mesons, urbaryon rearrangement diagrams for differential cross sections and polarizations for, (T), 26:46647
 interactions with photons, comparison of excited quark, interacting quarks, and additive quark models for constraints on angular decay distributions of resonances produced in, (T), 26:46526 (TPJU-14/71)
 interactions with photons, comparison of excited quark, interacting quark, and additive quark models for constraints on angular decay distributions of resonances produced in, (T), 26:46569
 interactions with pseudoscalar mesons, duality constraints for baryon exchange degeneracy patterns and absence of parity doublets and backward peak behavior in octet pseudoscalar meson and decuplet baryon production in, (T), 26:46627
 interactions with spin-0 mesons using polarized targets, lecture on method for determination of spins of baryon resonances produced in quasi-two-body, (T), 26:46628
 mass formulas for, dynamic interpretation of, (T), 26:46594
 mixing in SU(6) 70, L = 1 supermultiplet, exchange degeneracy for model of, (T), 26:46685
- BARYONS, RESONANCE**
 decay angular distributions in photon interactions with spin-1/2 baryons, comparison of excited quark, interacting quarks, and additive quark models for constraints on, (T), 26:46526 (TPJU-14/71)
 decay of, comparison of excited quark, interacting quarks, and additive quark models for constraints on angular distributions for, (T), 26:46569
 existence of exotic, in inclusive interactions in Regge pole model, 26:46605
 interactions $K^+ + p \rightarrow K^+ + p$ at 0.52 to 1.89 GeV/c, partial-wave amplitudes and exotic resonance in P/sub 3/2/ partial wave from phase shift analysis of, (T), 26:46632
 production in $e + N$ interactions, predictions of nonrelativistic harmonic-oscillator quark model for cross sections for, 26:46544
 production in $K^- + p$ interactions at 10 GeV/c in bubble chamber, review of, (E), 26:46480
 production in photon interactions with baryons, comparison of excited quark, interacting quark, and additive quark models for constraints on angular decay distributions in, (T), 26:46569
 production in π reactions with xenon at 2.34 to 9 GeV/c, (E), 26:46476 (JINR-P1-6408)
 properties of exotic, phase shift analysis of $K^+ + p \rightarrow K^+ + p$ interactions at 0.52 to 1.89 GeV/c for, (T), 26:46632
 spin assignment for high-spin, tests of, (T), 26:46621
 spin determination in spin-zero meson interactions with spin-1/2 baryons using polarized targets, lecture on method of, (T), 26:46628
 spin distribution in dual resonance model, statistical mechanics calculation of, (T), 26:46609 (ITF-72-13-P)
- BARYONS, RESONANCE (Δ)**
 form factor for $\Delta(1236)$ in $e + N \rightarrow \pi^0 + N$ interactions in first resonance region, multipole amplitude analysis of transition, (T), 26:46512 (CONF-710816-16)
 form factor for γp vertex determination from longitudinal and transverse cross sections for inelastic $e + p$ interactions in $\Delta(1236)$ region at 20 to 810, magnetic dipole transition, (E), 26:46431 (NP-19256)
 interactions $\Delta + \text{anti } \Delta$, duality in parton model for, (T), 26:46599
 interactions $\Delta + N$, models for induced nonexchange and exchange, (T), 26:46644
 interactions $\Delta + N$, properties of $\pi + N \rightarrow \pi + N + N$

amplitude for amplitude of, (T), 26:46644
interactions of many-baryon, contradiction between
isonuclei and many-baryon resonance theories for energy
dependence of total and differential cross sections and
number of reaction occurrences as function of invariant
and deficient mass of system for, (T), 26:46658
production in $\gamma + N \rightarrow \rho + \Delta(1236)$ interactions, method
for dispersion theory for, (T), 26:46511 (CONF-710816-
9)
production of $\Delta(1236)$ in $\nu + N$ interactions, 26:46584
production of $\Delta(1236)$ in $\pi^- + p \rightarrow 2\pi^0 + n$ interactions at
1.60 to 2.40 GeV/c, (E), 26:46477 (LBL-378)
production of $\Delta(1950)$, $\Delta(1910)$, $\Delta(1890)$, and $\Delta(1960)$ in
 $\pi^+ + p \rightarrow \pi^0 + \Delta^+(1236)$ interactions at 1820 to 2090
MeV, partial-wave analysis of, (E), 26:46495
production of $\Delta^-(1236)$ and $\Delta^+(1236)$ in $\pi^- + p \rightarrow \pi^- +$
 $\pi^+(p^0) + n(p)$ interactions at 4 GeV/c, effective mass
and angular distributions and differential cross
sections for forward, (E), 26:46488
production of $\pi^-X^0(2700)$ enhancement in $p + p \rightarrow p + p +$
 $\pi^+ + \pi^- + X^0$ interactions at 19 GeV/c, (E), 26:46467
theory for light Δ nuclei containing $\Delta(1236)$, (T),
26:46644
theory for two-baryon Δ nuclei, isotriplet and isoquintet
energy levels in, (T), 26:46644
BARYONS, RESONANCE (N)
interactions $N^* + N$, properties of $\pi + N + N \rightarrow \pi + N + N$
amplitude for amplitudes of, (T), 26:46644
interactions of many-baryon, contradiction between
isonuclei and many-baryon resonance theories for energy
dependence of total and differential cross sections and
number of reaction occurrences as function of invariant
and deficient mass of system for, (T), 26:46658
production of $N(1470)$, $N(1520)$, $N(1688)$, and $N(2190)$ in
elastic and inelastic $p + p$ interactions in GeV range,
review of cross sections for, (E), 26:46464
BARYONS, RESONANCE (E)
decay $\Sigma(1640) \rightarrow \text{anti } K + N$ in deuterium bubble chamber,
branching ratio for, (E), 26:46474 (COO-1428-306)
decay $\Sigma(1640) \rightarrow \Sigma(A) + \pi$ in deuterium bubble chamber,
upper limit for branching ratio for, (E), 26:46474
(COO-1428-306)
mass and width of $\Sigma(1640)$ in $K^- + n$ interactions at 4.5
GeV/c in deuterium bubble chamber, (E), 26:46474 (COO-
1428-306)
production of $\Sigma(1385)$, $\Sigma^+(1640)$, and $\Sigma^-(1640)$, in $K^- + n$
interactions at 4.5 GeV/c in deuterium bubble chamber,
cross sections for, (E), 26:46474 (COO-1428-306)
BATTERIES
description of lithium/selenium, incorporating paste
electrolyte disk separating anode and cathode,
26:45434(P)
BEAM MONITORS
description for 1- to 50-MeV electron beams, 26:45618
(ICRU-21)
design using transition radiation, p. 261-2, 26:46873(R)
(CEA-N-1522)
BEAVER VALLEY POWER REACTOR, UNIT 1
electrical equipment for, additional data on storage
requirements for, 26:47503 (DOCKET-50334-31)
electrical wiring for, additional data on installation
procedures for, 26:47503 (DOCKET-50334-31)
quality assurance requirements for, additional data on
concrete sampling and testing, 26:47503 (DOCKET-50334-
31)
BELLOWS
performance of expansion, for research reactor
experimental facilities, 26:47591
BENZAMIDE
reactions with hydrated electrons from pulse radiolysis
of aqueous solutions, 26:45354
BENZANTHRACENE
electron reactions with, cross sections for electron
attachment in, (E), 26:46364 (ORNL-IM-3839)
BENZENE
production of aerosols of, by exposure of vapors to x
rays, 26:46082
BENZENE, DIVINYLBENZENE
polydivinylbenzene--polystyrene, sorptive properties for
iodomethane, 26:45388
BENZENE, (1,2-EPOXYETHYL)-
radioinduced polymerization of, effects of peroxide
sensitizers on dose rate in γ , 26:45325
BENZIL
reactions with hydrated electrons and hydrogen atoms from
radiolysis of aqueous solutions, 26:45354
BENZOIC ACID, p-(HYDROXYMERCURY)-
effects on deoxyribonuclease activity in liver of normal
and irradiated rats, 26:45732
BENZON
reactions with hydrated electrons from pulse radiolysis
of aqueous solutions, 26:45354
BENZOPHENONE
reactions with hydrated electrons and hydrogen atoms from
radiolysis of aqueous solutions, 26:45354
BENZOYL DISULFIDE
radiolysis of single crystals of, at 4.2°K, ESR spectral
study of radical ion saturation in x, 26:45348
BENZOYL PEROXIDE
effects on dose rate in radioinduced polymerization of
monomers, 26:45325
BENZYL (FREE RADICAL)
spectra of, produced by pulse radiolysis of aqueous acid

71 SUBJECTS

BIBLIOGRAPHIES

toluene solutions, 26:45309(R) (AE-453)
BERKELIUM IONS
magnetic properties of cesium sodium lutetium chloride
doped with, 26:47272
BERYLLIUM
dislocations at 300 to 500°K in, effects of temperature
on friction stress for basal, 26:46028
neutron diffusion in, temperature dependence of diffusion
length for, (T), 26:46857
separation from other metals as β -diketonates by column
liquid-liquid chromatography, 26:45281
BERYLLIUM ALUMINUM BORIDES
crystallization from aluminum melts, 26:45936(T) (UCRL-
Trans-10632)
BERYLLIUM BORIDES
crystallization from aluminum melts, 26:45936(T) (UCRL-
Trans-10632)
BERYLLIUM ISOTOPES
photon reactions (γ, p^0) at 3.5 to 8.3 GeV, A dependence
of, (E), 26:46447
proton reactions at 24 GeV/c, π , K, anti p, and p spectra
from, (E), 26:46459
BERYLLIUM ISOTOPES Be-7
content in atmosphere and vegetation in vicinity of
Lawrence Livermore Laboratory, Jan. through Dec. 1971,
26:45513 (UCRL-51242)
content in surface atmosphere at world sites, 1970
through Feb. 1972, 26:45512(R) (HASL-257(App.))
distribution in atmosphere and oceans, relations of
fission products to, 26:45530 (BNWL-SA-4205(Rev.))
energy levels of, DWBA K-matrix calculation of continuum,
(T), 26:46896
BERYLLIUM ISOTOPES Be-8
nuclear quadrupole moment of, calculation using effective
N-N interactions, 26:46892
BERYLLIUM ISOTOPES Be-9
neutron reactions ($n, 2n$) at 2.37 to 3.24 MeV, absolute
cross sections for, (E), 26:46874 (JUL-834-RG)
proton elastic and inelastic scattering at 6.36 and 6.48
MeV, angular distributions for, (E), 26:46885
BERYLLIUM OXIDES
crystallization from lithium molybdate--molybdenum oxide
flux, effects of solute composition on, 26:45943
BERYLLIUM SILICATES
crystallization from lithium molybdate--molybdenum oxide
flux, effects of solute composition on, 26:45943
BETA DECAY
half-lives for, tables of mean excitation energies of
daughters for prediction of, 26:46875 (KFK-EXT-10/72-
1)
log-f tables for, 26:46970
transitions between anti-analog states in, shell-model
analysis of, 26:47050
BETA PARTICLES
content in atmosphere in Ispra, Italy, Sept. 1971 through
March 1972, 26:45519 (HASL-257, pp III.53-8)
dose to thyroid gland of adult and fetal mice following
injection of iodine-131, mathematical model for
calculating, 26:45902
effects on survival of cultured mammalian cells, relation
of mathematical analysis of, to radiotherapy, 26:45735
effects on tongue of rabbits, permeability of lingual
mucosa during development and recovery from
pathological, 26:45862
emission from iodine-131 and tritium, toxic effects of,
on mouse cells, 26:45892 (ORO-2690-12(Pt.B), Paper 4)
monitoring in atmospheric precipitation and ground-level
atmosphere in Canada, South America, and US during Feb.
1972, 26:45523
monitoring in California sewage and water supplies during
1970, 26:45537
monitoring in Lake Michigan aquatic environment near Big
Rock Point power reactor during 1960 to 1969, 26:45535
monitoring in liquid waste effluents at Brookhaven
National Laboratory during Jan. to July 1968, 26:45515
BETA SOURCES
determination of low-activity, recommendations for,
26:45261 (ICRU-22)
monitoring in vicinity of Lawrence Livermore Laboratory,
Jan. through Dec. 1971, 26:45513 (UCRL-51242)
release at West Valley Processing Plant, April-June,
1972, 26:45398 (DOCKET-50201-125)
BETA SPECTROMETERS
description of cylindrical multiplate electrostatic, for
electrons at 1 to 10 keV, 26:45643
description of electrostatic, with channel electron
multipliers for electrons at 0.03 to 10 keV, 26:45642
BETATONS
beam shaper for B5M-25, design and performance of
improved, 26:46132
research at Bucharest, with bremsstrahlung beam,
26:46148
BEVERAGES
coffee, fallout cesium-137 and strontium-90 content in,
Mecklenburg, GDR, 1969, 26:45546
tea, fallout cesium-137 and strontium-90 content in,
Mecklenburg, GDR, 1969, 26:45546
BIBLIOGRAPHIES
on carbon-13, 26:45286 (MLM-1913)
on fused salts, author, citation, and subject--title list

- for, 26:45287 (SC-R-722769)
on LLL computer documentation reports, 26:47598 (UCID-30047)
on numerical algebra, 26:47287 (ORNL-4778)
on open-file reports for Gasbuggy, Rio Blanco, Rulison, and Wagon Wheel Projects in Project Plowshare, 26:45477 (NVO-86(Rev.1))
on organizations and projects for crystal growing, 26:45973 (ORNL-RMIC-12)
on positron lifetimes in light and heavy ice and water and frozen aqueous solutions, 26:46362 (RISO-254)
on surface and vacuum physics, 26:46336
- BIG ROCK POINT POWER REACTOR**
emergency systems for, revised test requirements for explosive valves of liquid poison, 26:47394 (DOCKET-50155-119)
fuel bundles for, design characteristics of NFS-DA, 26:47395 (DOCKET-50155-120)
fuel rod bundle for, illustration of Reload G, 26:47393 (DOCKET-50155-117)
radioactive effluents from, gross beta radioactivity in Lake Michigan aquatic environment during 1960 to 1969 from, 26:45535
radioactive effluents from, radioisotope content of Lake Michigan biota, sediments, and water during 1970 from, 26:45536
- BIOLOGICAL MATERIALS**
analysis for plutonium by extraction and liquid scintillation counting, 26:45268
analysis for vanadium, comparison of neutron activation and spectrophotometric methods for, 26:45203
- BIOLOGICAL MODELS**
development for calculating radiation dose to thyroid gland of adult and fetal mice following injection of iodine-131, 26:45902
development of compartmental, for determining relation between radiation dose and radioisotope distribution and retention in man, 26:45905 (RD/B/N-2291)
use of respiratory tract, for measurements of radioisotope deposition and retention kinetics following inhalation as aerosols, 26:45921
- BIOLOGY**
book: Masking and Demasking of Chemical Reactions. Theoretical Aspects and Practical Applications, 26:45188
- BIOPHYSICS**
research programs on molecular, at Florida State University, Tallahassee, during 1971 and 1972, 26:45891(R) (ORO-2690-12(Pt.B))
- BISMUTH**
determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
purification by zone melting in rotating container, 26:45276
- BISMUTH ALLOYS AND SYSTEMS**
Bi-In, thermodynamic properties of intermetallic In_3Bi_3 , 26:46026
- BISMUTH ISOTOPES BI-209**
neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909
proton reactions (p,n) at 20 to 45 MeV, with subsequent proton decay, coupled-channels calculations of energy dependence of, (T), 26:47073
- BISMUTH ISOTOPES BI-212**
alpha decay of, charge states of thallium-208 from, 26:45359
- BISMUTH ISOTOPES BI-214**
content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)
- BLACK BODY RADIATION**
standard source for, performance of argon arc, 26:47217
- BLACK HOLES**
collision of two, Brans-Dicke theory for absence of scalar gravitational radiation in, (T), 26:46192
energy extractable from, Kerr solution and surface area for upper limits on amounts of, (T), 26:46193
force between two, upper limits on amounts of extractable energy for spin-dependent, (T), 26:46193
gravitational theory for, gravitational mass loss in formation, equivalence principle violation, and absence of scalar gravitational radiation in collision in Brans-Dicke, (T), 26:46192
production of, Brans-Dicke theory for gravitational mass loss in, (T), 26:46192
relativistic Roche problem associated with, (T), 26:46189
- BLACK SEA**
book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
- BLOOD**
radiation effects on activity of glutathione peroxidase in, of swine, x, 26:45733
radiation effects on count of mast cells in, of man following exposure of mediastinum, x, 26:45810
- BLOOD CELLS**
radiation effects on yield of granulocytes and macrophages in peritoneal diffusion chamber cultures, effects of exposure of host mice on, 26:45741
- BLOOD PLASMA**
analysis of rat, for strontium by neutron activation and radiochemical separation, 26:45202
radiation effects on ESR spectra of human, following therapy, 26:45723
radiation effects on fatty acids of in vitro, effects of water on x, 26:45725
- BLOOD VESSELS**
radiation effects on regeneration of, in muscle transplants in rats, 26:45857
radioinduced injuries to, of rabbit mandible bones, dose dependence of gamma, 26:45835
radioinduced malformations in, of rat fetuses following sublethal whole-body exposure in utero, statistical analysis of relation between age, dose, and incidence of fast neutron, 26:45867
- BLOWERS**
design of efficient high-volume miniature, 26:47633 (UCRL-51239)
performance of helium, in nuclear aircraft engines, effects of power take-off design on, 26:45425 (N-72-16724)
- BODY CAVITIES**
radiation dose distribution in human pelvis, from radium applicator implanted in uterus, method for calculating, 26:45784(T) (TT-54002/4, pp 456-61)
- BOILERS**
deposits chemistry in HTGR, 26:45288 (TID-26106)
design of sodium-heated steam, 26:45450(P)
welded joints in large-capacity, radiometric method for inspection of, 26:45473
- BONE MARROW**
radiation effects on rat, following lethal whole-body exposure, role of transplanted slowly proliferating bone marrow cells in regeneration of x, 26:45837
radiation effects on stem cell renewal in mouse, following fractionated whole-body exposure, x, 26:45871
radioinduced chromosome aberrations in cells of human, following single local exposure, effects of dose and post-exposure time on frequency of γ , 26:45794
radioinduced injuries to, of rabbit mandible bones, dose dependence of gamma, 26:45835
strontium-89-induced hypoplasia in, of mice, effects of, on hematopoiesis and immune reactions, 26:45896
- BONES**
radiation effects on growth of tissues of, effects of dose and parotin on, 26:45870
radioinduced injuries of rabbit mandible, dose dependence of gamma, 26:45835
radioinduced neoplasms in, following uptake of radioisotopes, review of pathology of, 26:45895 (UR-3490-102)
radioinduced neoplasms in, of cats following chronic ingestion of strontium-89, dose dependence of β , 26:45901
strontium-90 content in human, in New York City and San Francisco, 1971, effects of diet on, 26:45538 (HASL-257, pp 1.51-70)
strontium-90 passage from bovine, to processed food products, 26:45549
- BORON**
determination in borosilicate glass by flame emission spectroscopy, 26:45254
determination of, chemical procedures for, 26:45181 (CISE-R-172)
effective electron mass estimate from Doppler-broadened positron annihilation radiation, 26:47268
Moessbauer spectra of chromium-57 and iron-57 in β -rhombohedral, 26:47280
neutron capture in, cellulose track detector measurement of spatial distribution of thermal, 26:45615
positron annihilation radiation in, correlation of Doppler-broadened, 26:47268
reactions with molybdenum-niobium alloys at 1100 to 1400°C, effects of composition on, 26:46002
- BORON CARBIDES**
analysis for boron-11/boron-10 ratio by thermionic ionization mass spectrometry, 26:45235
proton reactions at 24 GeV/c, w , K, p, and anti p spectra from, (E), 26:46459
 B_4C -graphite-base, radiation effects on boron-10-containing control materials of, fast neutron, 26:46077 (GULF-GA-A-12035)
- BORON ISOTOPES B-10**
determination of abundance of, relative to boron-11 in boron carbide by thermionic ionization mass spectrometry, 26:45235
energy levels of, spin determinations from γ - γ angular correlations, 26:46901
meson (K^-) elastic scattering by, scattering lengths for, (E), 26:46912
reactions with tantalum-181 at 50 to 100 MeV, analysis of compound nucleus deexcitation using statistical evaporation model, (E/T), 26:46974 (FRNC-TH-207)
- BORON ISOTOPES B-11**
determination of abundance of, relative to boron-10 in boron carbide by thermionic ionization mass spectrometry, 26:45235
lithium-6 elastic scattering at 28 MeV, optical-model analysis of angular distribution from, p. 26-30, (E), 26:46873(R) (CEA-N-1522)
meson (K^-) elastic scattering by, scattering lengths for,

reactions with tantalum-181 at 50 to 100 MeV, analysis of compound nucleus deexcitation using statistical evaporation model, (E/T), 26:46974 (FRNC-TH-207)

BORON NITRIDES

structure of hexagonal wurtzite-like, 26:47255

BOROSILICATES

analysis of glass of, for boron by flame emission spectroscopy, 26:45254

BOSONS

interactions with bosons, regularization method in S-matrix theory in p^4 momentum representation for, (T), 26:46567

quantum theory for spin-0, in external electric fields, (T), 26:46741

S-matrix for spin-0, in external electric fields, (T), 26:46741

theory for states of, generating function method for expansion of Hamiltonian operator from isomorphism between space of fermion states and space in, (T), 26:47286 (LYCEN-7202)

theory for, unitary operator form of general Bogolubov--Tyablikov transformation in, (T), 26:47291

BOSONS, INTERMEDIATE

production and decay of χ^0 , (T), 26:46577
production of W in $\nu + N$ interactions, 26:46584
production of $W^+(W^-)$ in hadron interactions at super-high energies, (T), 26:46598

BRAIN

radiation effects on enzyme activity in neuroglia of developing, of rabbits, x, 26:45836

radiolinduced injuries in visual cortex of, in monkeys, latent x, 26:45858

BREMSSTRAHLUNG

dosimetry in GeV range using $FeSO_4$ dosimeter, 26:45625
dosimetry of, design of tissue-equivalent liquid ionization chamber for, 26:45621

emission from Maxwellian plasma, (T), 26:47152

production of soft, in $\pi^- + d + 2n + \gamma$ at 150 to 300 MeV, phenomenological calculation of cross sections for, (T), 26:46531

reactions (γ, f) with thorium-232, uranium-235, and uranium-238, cross sections for, (E), 26:46890

reactions (γ, n) and ($\gamma, 3n$) with holmium-165 at 90 MeV, relative cross sections for, (E), 26:46998

reactions (γ, n) and (γ, p) with $A = 12$ to 208 nuclei up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

reactions (γ, p) with carbon-12 at 130 MeV, analysis of differential cross sections and spectra from, (E/T), 26:46888

reactions ($\gamma, p\pi^-$) with helium-4 and carbon-12 at 300 MeV, differential cross sections for, p. 182-3, (E), 26:46873(R) (CEA-N-1522)

reactions with nuclei, conversion of bremsstrahlung photon neutron mean energies to monochromatic photon neutron mean energies for, 26:46953

reactions with nuclei, research on, 26:46148

BRICKS

development of aluminum oxide, as thermoluminescent dosimeter, 26:45628

BROMIDES

solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

BROMINE

cesium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366

determination as impurity in vinyl chloride by neutron activation, 26:45204

determination in hair by neutron activation and γ spectroscopy, 26:45214

potassium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366

Br--bromoethane, bromine-80m isomeric transition in mixtures of, effects of initial form on yield of labeled compounds and chemical damage in, 26:45360

BROMINE FLUORIDES

reactions of gaseous, with uranium dioxide and U_3O_8 , gravimetric study of, 26:45379

BROMINE ISOTOPES Br-72

nuclear properties of, compilation of, (E), 26:46939

BROMINE ISOTOPES Br-76

beta decay of, β and γ spectra from, (E), 26:46938

BROMINE ISOTOPES Br-80

isomeric transition in bromine--bromoethane mixtures, effects of initial form on yield of labeled compounds and chemical damage in, 26:45360

BROMINE ISOTOPES Br-82

reactions of recoil, in neutron-irradiated bromopentacarbonylmanganese, 26:45357
use as tracer for measuring mixing efficiency in aerosol chamber, 26:45444

BROWNS FERRY POWER REACTOR, UNIT 1

control room ventilation system for, use of charcoal filtering and pressurizing equipment in, 26:47428 (DOCKET-50259-60)

emergency electrical power systems for, revised and additional data on, 26:47426 (DOCKET-50259-58)
flood protection for, use of physical barriers for,

26:47428 (DOCKET-50259-60)

kinetics for, requested additional data on, 26:47425 (DOCKET-50259-57)

safety of, revised and additional data on, 26:47427 (DOCKET-50259-59)

BROWNS FERRY POWER REACTOR, UNIT 2

control room ventilation system for, use of charcoal filtering and pressurizing equipment in, 26:47428 (DOCKET-50259-60)

emergency electrical power systems for, revised and additional data on, 26:47426 (DOCKET-50259-58)

flood protection for, use of physical barriers for, 26:47428 (DOCKET-50259-60)

kinetics for, requested additional data on, 26:47425 (DOCKET-50259-57)

safety of, revised and additional data on, 26:47427 (DOCKET-50259-59)

BROWNS FERRY POWER REACTOR, UNIT 3

control room ventilation system for, use of charcoal filtering and pressurizing equipment in, 26:47428 (DOCKET-50259-60)

emergency electrical power systems for, revised and additional data on, 26:47426 (DOCKET-50259-58)

flood protection for, use of physical barriers for, 26:47428 (DOCKET-50259-60)

kinetics for, requested additional data on, 26:47425 (DOCKET-50259-57)

safety of, revised and additional data on, 26:47427 (DOCKET-50259-59)

BRUNSWICK POWER REACTOR, UNIT 1

containment system for, radiation doses from purging of, 26:47496 (DOCKET-50324-39)

environmental report for, revised cost-benefit information for, 26:47497 (DOCKET-50324-40)

BRUNSWICK POWER REACTOR, UNIT 2

containment system for, radiation doses from purging of, 26:47496 (DOCKET-50324-39)

environmental report for, revised cost-benefit information for, 26:47497 (DOCKET-50324-40)

BUTADIENE

butadiene-- NH_3 , radiolysis of mixtures of, yields of hydrogen from γ , 26:45332

BUTANE

butane--I, neutron reactions (n, γ) with iodine-127 in gaseous mixtures of, chemical state of recoil iodine-128 from, 26:45361

butane--I, neutron reactions (n, γ) with liquid mixtures of, chemical state of recoil iodine-128 from, 26:45362

1-BUTANESULFINIC ACID, 4-[(2-ACETAMIDOETHYL)DITHIO]-sodium salt, properties as radioprotective drug, effects of disproportionation reactions on, 26:45290

BUTYROPHENONE

radiolysis in benzene solutions, yields from, 26:45326

**CABLES**

fabrication method for flat, autoclave, 26:47607 (BDX-613-579(Rev.))

shielding for, method for preparation of braided wire, 26:47606(R) (BDX-613-543(Rev.))

CADMIUM

analysis as high-purity oxide, for trace amounts of elements, spectroscopic, 26:45220 (BARC-593)

determination of trace amounts of, in environmental samples by neutron activation, 26:45198

electron-phonon unklapp interaction at low temperatures in, (E), 26:47264

neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)

positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268

surface electron band structure of single crystals of, magnetically induced, 26:47256

CADMIUM ISOTOPES Cd-109

content in surface atmosphere at world sites, 1964 through Nov. 1967, 26:45512(R) (HASL-257(App.))

CADMIUM ISOTOPES Cd-110

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CADMIUM ISOTOPES Cd-111

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CADMIUM ISOTOPES Cd-112

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CADMIUM ISOTOPES Cd-113

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CADMIUM ISOTOPES Cd-114

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CADMIUM ISOTOPES Cd-116

photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

CAFFEINE

effects on deoxyribonuclease activity in liver of normal and irradiated rats, 26:45732
radioinduced alkylation by amines, γ , 26:45326

CALCIUM

determination in calcium-45 solutions, spectrographic, 26:45223 (JEN-232-DQ/I-81)
determination in meteorites by proton activation and bremsstrahlung activation, 26:45215
determination in soils, 26:45263 (RT/PROT-72-11)
effects on strontium-90 accumulation in wheat in podzolic soils, 26:45550
excretion in urine of children in Poland, relation between dietary intake and, 26:45908
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

CALCIUM FLUORIDES

CaF_2 -- YF_3 , thermodynamics of solid electrolytes of, review on, 26:46013

CALCIUM ISOTOPES Ca-40

electron reactions ($e, e'p$) at 501 MeV, cross sections for, p. 194-6, (E), 26:46873(R) (CEA-N-1522)
energy levels from potassium-39 ($\text{He}-3, d, \gamma$) reactions, lifetimes and partial widths for, p. 41-53, (E), 26:46873(R) (CEA-N-1522)
lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
muon (μ^-) capture by, particle-hole calculation of capture rate and neutron spectra for, (T), 26:46880
nuclear quadrupole moment rearrangement due to proton removal, calculation of, (T), 26:46886
photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
proton elastic scattering at 2.36 to 2.78 MeV, polarization from polarized, (E), 26:46924
proton inelastic scattering at 40 MeV, DWBA and coupled channel analysis of, (E), 26:46894
proton separation and orbital rearrangement energies, calculations of, (T), 26:46886

CALCIUM ISOTOPES Ca-44

proton reactions (p, γ) at 1658 keV, gamma spectra from, (E), 26:46920

CALCIUM ISOTOPES Ca-45

analysis of solutions of, for calcium-45 and target element, spectrographic, 26:45223 (JEN-232-DQ/I-81)

CALCIUM ISOTOPES Ca-47

uptake and retention in mice following gastric tube administration, effects of aluminum phosphate gel on, 26:45918

CALCIUM OXIDES

Al_2O_3 -- CaO -- Cr_2O_3 -- F -- MgO -- MoO_3 -- TiO_2 -- ZrO_2 , use as welding flux, 26:45941(P)
 CaO -- La_2O_3 , thermodynamics of solid electrolytes of, review on, 26:46013
 CaO -- ZrO_2 , phase studies on partially stabilized, 26:45945
 CaO -- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

CALCIUM PHOSPHATES

cathodoluminescence of single crystals of, 26:45591

CALCIUM SULFATES

BaSO_4 -- CaSO_4 , sorptive properties for strontium, 26:45273 (UJV-2820-Ch)

CALCIUM URANIUM SELENIDES

preparation and structure of CaU_2Se_6 , 26:45378

CALIFORNIA

radiation monitoring of sewage and water supplies in, during 1970, 26:45537
radioactivity monitoring in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
radioactivity monitoring of ground-level atmosphere in, during Feb. 1972, gross beta, 26:45523
strontium-90 content in diets in San Francisco, Jan. to March 1972, 26:45539 (HASL-257, pp II.1-6)

CALIFORNIUM ISOTOPES Cf-252

analysis for impurities by spark source mass spectrometry, 26:45260
fabrication of neutron sources of, description of method for, 26:45714 (CONF-720902-2)
fission of, angular distributions of long range alpha particles from, (E), 26:47037
fission of, calculation of prompt gamma yields using improved renormalized Fermi gas model for spontaneous, (T), 26:47025
neutron fission of, gamma and neutron emission in, p. 314-8, (E), 26:46873(R) (CEA-N-1522)
production as oxide by calcination after sorption of hydroxyisobutyrate on carboxylic acid resin, 26:45393
production of, mass spectrometric analysis for quality control, 26:45260
use as neutron source for radiography, 26:45467 (BNWL-SA-4320)
use for therapy, methods for reducing neutron dose to medical personnel during interstitial and

intercavitary, 26:45790

CALIFORNIUM ISOTOPES Cf-253

energy levels of, calculation using phenomenological model, (T), 26:47081

CALIFORNIUM OXIDES

production of californium-252, by calcination after sorption of hydroxyisobutyrate on carboxylic acid resin, 26:45393

CALVERT CLIFFS POWER REACTOR, UNIT 1

crane for, failure of upper limit switch for polar, 26:47381 (DOCKET-50317-53)

CALVERT CLIFFS POWER REACTOR, UNIT 2

crane for, failure of upper limit switch for polar, 26:47381 (DOCKET-50317-53)

CANADA

nuclear power plants for, site selection criteria for, 26:47574
radiation protection programs of diagnostic radiography facilities in, 26:45811
radioactivity monitoring of atmospheric precipitation and ground-level atmosphere in, during Feb. 1972, gross beta, 26:45523
radioisotope content of milk in, during Mar. 1971 to Mar. 1972, cesium-137 and strontium-90, 26:45552

CANALS

construction of inland, between Orinoco and Rio Negro Rivers in Venezuela, feasibility study of nuclear excavation method for, 26:45475 (NP-19361)

CANDU REACTOR

thermal effluents from, temperature structure of Lake Huron underwater plumes of, (E), 26:45581

CARBAMIC ACID, DITHIO-

ammonium salt, radioprotective effects of, on Escherichia coli and mice, 26:45859

CARBON

burning in supergiant stars, effects of neutrino emission on, (T), 26:46177
deposition of pyrolytic, in graphite, pulsed method for, 26:46088(P)
determination in stainless steel and Zircaloy by deuterium and proton activation, 26:45309(R) (AE-453)
distribution in steel surfaces, measurement by charged particle and proton activation, 26:45466 (AE-451)
effects on recrystallization of iron, 26:45982
helium-ion reactions at 40 keV with foils of, relative level populations excited by, as functions of quantum number and spin, (E), 26:46337
lithium-ion reactions at 40 keV with foils of, relative level populations excited by, as functions of quantum number and spin, (E), 26:46337
mechanical properties of, research program on, 26:46078 (NYO-1710-106)
neutron transmission below 10^{-4} eV through films of, interference effects in cold, (E), 26:46853
polarizability of ^{13}C , ^{13}D , and ^{31}P states of, static dipole, (T), 26:46419
preparation of graphitized, research program on, 26:46078 (NYO-1710-106)
reactions with nitrogen and uranium dioxide, 26:45417(R) (ORNL-TM-3281)
reactions with nitrogen and uranium dioxide, 26:45408(R) (ORNL-TM-2949)

CARBON DIOXIDES

carbon-13 concentration in, by gas chromatography, 26:45302(R) (COO-1222-55)
effects on yields of excited states from pulse radiolysis of deaerated naphthalene-cyclohexane solutions, 26:45330
electron reactions at 25 to 600 eV with, cross sections for production of CO^+ , O^+ , C^+ , and O_2^+ by dissociation and ionization in, (E), 26:46384
proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s hydrogen atoms by charge exchange and excitation in, (E), 26:46378
radiolysis of gaseous, effects of density on γ , 26:45346
reactions with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
recovery as by-product from uranium dioxide production, method for, 26:45371(P)

CARBON IONS

implantation in iridium at keV energies, field-ion microscopic study of damage profiles produced in, 26:47240
reactions with cobalt, iron, tin, and zirconium at 5 and 10 MeV/amu, energy shifts and relative intensities of K x rays from, (E), 26:46339

CARBON ISOTOPES

determination of ratio of stable, as indicators of air pollution levels, 26:45245
neutrino reactions in propane bubble chamber with, test of PCAC from differential cross sections for inelastic, (E/T), 26:46450
particle reactions at 300 to above 1000 MeV, intranuclear cascade model for density effects on cascade particle properties and density and cluster-structure effects on isotope yield in, (T), 26:46664
photon reactions (γ, π^0) at 3.5 to 8.3 GeV, Λ dependence of, (E), 26:46447
photon reactions at 1.160 GeV at 150 μ , photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441

CARBON ISOTOPES C-12

alpha reactions (α, γ) near 8 MeV, detailed balance test

- of time reversal invariance in, (E), 26:46881
- carbon-12 reactions (C-12, α) and (C-12,p) at 2.4 to 5 MeV, angular distributions and excitation functions for, (E), 26:46898
- carbon-12 reactions (C-12,p) at 30 MeV, angular distributions for, (E), 26:46915
- electron reactions (e,e'p) at 501 MeV, cross sections for, p. 194-6, (E), 26:46873(R) (CEA-N-1522)
- electron reactions (e,p) at 130 MeV, analysis of differential cross sections and spectra from, (E/T), 26:46888
- lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
- meson (K-) elastic scattering by, scattering lengths for, (E), 26:46912
- meson reactions (π^+ ,2p) at 76 MeV, cross section for, 26:46870
- meson reactions (π^- , π^- p) at 1 GeV/c, π^- -meson momenta from, (E), 26:46869
- nitrogen-14 reactions (N-14, Li-6) at 45 MeV, energy spectra from, p. 22-4, (E), 26:46873(R) (CEA-N-1522)
- nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892
- photon reactions (γ ,n) and (γ ,p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
- photon reactions (γ ,p) at 130 MeV, analysis of differential cross sections and spectra from, (E/T), 26:46888
- photon reactions (γ , π^-) at 300 MeV, differential cross sections for, p. 182-3, (E), 26:46873(R) (CEA-N-1522)
- proton elastic scattering at 24.5 MeV, optical-model analysis of angular distribution and polarization from polarized, p. 134-7, (E), 26:46873(R) (CEA-N-1522)
- proton inelastic scattering at 24.5 MeV, coupled-channels and DWBA analyses of differential cross sections and asymmetries from polarized, p. 138-42, (E), 26:46873(R) (CEA-N-1522)
- proton inelastic scattering at 40 MeV, DWBA and coupled channel analysis of, (E), 26:46894
- reactions (C-12, Be-10) with oxygen-18 at 46 MeV, energy spectra from, p. 20-1, (E), 26:46873(R) (CEA-N-1522)
- reactions (C-12, α) and (C-12,p) with carbon-12 at 2.4 to 5 MeV, angular distributions and excitation functions for, (E), 26:46898
- reactions (C-12, α) with carbon-14 at 35 and 40 MeV, cross sections and spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- reactions (C-12,p) with carbon-12 at 30 MeV, angular distributions for, (E), 26:46915
- reactions with carbon-12 in stars, rates for, (T), 26:46179
- reactions with oxygen-16 in stars, rates for, (T), 26:46179
- transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889
- CARBON ISOTOPES C-13**
- bibliography on properties, analytical and production methods, and uses of, 26:45286 (MLM-1913)
- concentration in carbon dioxide by gas chromatography, 26:45302(R) (COO-1222-55)
- effects on Franck-Condon factors and r-centroids for diatomic carbon molecules, isotopic, 26:45293
- effects on spin-lattice relaxation times of formic acid, 26:45297
- helium-3 reactions (He-3, α) at 29 to 42 MeV, excitation functions for, p. 31-32, (E), 26:46873(R) (CEA-N-1522)
- lithium-6 elastic scattering at 28 MeV, optical-model analysis of angular distribution from, p. 26-30, (E), 26:46873(R) (CEA-N-1522)
- proton elastic scattering at 24.5 MeV, optical-model analysis of angular distribution and polarization from polarized, p. 134-7, (E), 26:46873(R) (CEA-N-1522)
- CARBON ISOTOPES C-14**
- carbon-12 reactions (C-12, α) at 35 and 40 MeV, cross sections and spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- design of aluminum-backed and self-supporting targets of, p. 93-4, 26:46873(R) (CEA-N-1522)
- effects on *Drosophila* following ingestion, genetic, dose dependence of β , 26:45912(R) (ORO-3728-5)
- oxygen-16 reactions (O-16, α) at 35 MeV, cross sections and spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- reactions of recoil, in neutron-irradiated solid acetanilide, 26:45358
- CARBON ISOTOPES C-18**
- energy levels of, data compilation on, 26:46882
- CARBON ISOTOPES C-19**
- energy levels of, data compilation on, 26:46882
- CARBON ISOTOPES C-20**
- energy levels of, data compilation on, 26:46882
- CARBON MONOXIDES**
- abundances in interstellar clouds, (E/T), 26:46244
- proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s hydrogen atoms by charge exchange and excitation in, (E), 26:46378
- radioinduced desorption from niobium (111), temperature dependence of electron, 26:47242
- CARBON OXIDES**
- content variations of CO₃⁻ and CO₃⁰ with time in ionospheric D region, 26:46281
- electron photodetachment from CO₃⁻ in ionospheric D region, (E/T), 26:46299
- CARBON TETRACHLORIDE**
- electron reactions with, cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- radioinduced reaction with cyclohexanone, yields from γ , 26:45335
- CARBON TETRAFLUORIDE**
- radiolysis of, research progress on production of large-molecular-weight fluorocarbons by γ , 26:45316(R) (ORO-3909-3)
- CARBONACEOUS ROCKS**
- uranium mineralization in, epigenetic zonality in, 26:45506
- CARBONYL FLUORIDE**
- determination in nitrogen by automatic absorption spectrometry, 26:45243(P)
- CARDIAC PACEMAKERS**
- development of implantable radioisotope-fueled, 26:45699 (CEA-CONF-1998)
- heat sources for, French production of plutonium-238, 26:45706
- CARIBBEAN SEA**
- strontium-90 content and distribution in Antilles Arc of, during 1970, vertical profiles of, 26:45533
- CASCADE SHOWERS**
- production of high-energy hadron intranuclear, in concrete, (E/T), 26:46126(T) (ORNL-tr-2617)
- CATS**
- radioinduced neoplasms in bones and soft tissues adjacent to bones in, following chronic ingestion of strontium-89, dose dependence of β , 26:45901
- CATTLE**
- radioinduced ocular squamous cell carcinoma in, mechanisms of uv, in comparison with human xeroderma pigmentosum, 26:45851
- CAVITY RESONATORS**
- particle acceleration in H₁₁₁ cylindrical, with axial magnetic drive field, 26:46139
- CELL MEMBRANES**
- radiation effects on permeability of rat thymocyte, to potassium ions following exposure in vitro, x, 26:45744
- CELL NUCLEI**
- radiation effects on protein synthesis in, of mouse spleen following whole-body exposure, comparison of effects of Friend virus infection with x, 26:45849
- CELL PHASE**
- relation to radiosensitivity of living cells, review of, 26:45845
- CELLS**
- growth of mouse leukemic (L5178Y), effects of chloramphenicol and cycloheximide on, 26:47647 (UR-3490-132)
- injection of leukemia, labeled with iodine-125, iodine-131, and tritium into mice, toxic effects following, 26:45892 (ORO-2690-12(Pt.B), Paper 4)
- iodine-125 effects on, following uptake, biophysical evaluation of toxic, 26:45893 (ORO-2690-12(Pt.B), Paper 13)
- radiation effects on DNA in, of higher animals and plants, relation of organizational properties of genome to internal, 26:45894 (ORO-2690-12(Pt.B), Paper 14)
- radiation effects on hematopoietic stem, in mice, effects of cell-free tissue extracts on recovery from x, 26:45847
- radiation effects on mammalian, model for recovery from, 26:45816 (FRNC-R-7)
- radiation effects on proliferation and regeneration of lens, during injury response and wound healing in rabbits, x, 26:45813(R) (COO-1427-41)
- radiation effects on renewal of hematopoietic stem, in bone marrow and spleen of mice following fractionated whole-body exposure, x, 26:45871
- radiation effects on spleen colony formation by bone marrow, of rats following transplantation into mice, x, 26:45834
- radiation injuries to, in relation to sensitization, review of, 26:45845
- radioinduced injuries of, in mammals following internal irradiation, review of, 26:45890 (NYO-3970-29)
- CELLS, CULTURED**
- properties of x-radioinduced leukemia, from mice, cytological, enzymatic, and growth, 26:45855
- radiation effects on DNA complex in Chinese hamster, dose dependence of repair processes following x, 26:45745
- radiation effects on DNA in human Xeroderma pigmentosum, in comparison to effects of chemical carcinogens and mutagens, extent of repair synthesis following uv, 26:45742
- radiation effects on DNA synthesis in HeLa, following microbeam exposure of half-nuclei, effects of hydroxyurea on uv, 26:45737
- radiation effects on human bone marrow, protective effects of taurine against, 26:45734
- radiation effects on phagocytic activity of macrophage, co-cultured with tumor cells, x, 26:45738
- radiation effects on survival of mammalian, relation of mathematical analysis of, to radiotherapy, 26:45735
- radiation effects on survival of, predictions of delta-ray theory of track effects for, 26:45611

- radiation effects on survival of, predictions of delta-ray theory of track structure for, 26:45612
- radiation effects on survival of, relation of cell cycle and environmental and intrinsic factors to, 26:45739
- radiation effects on uv fluorescence of Ehrlich ascites, x, 26:45736
- radioinduced aberrations of chromosomes in lymphocyte, from human population of Marshall Island in relation to age and sex, 26:45792
- radioinduced chromosome aberrations in Chinese hamster ovary, effects of post-exposure temperature on repair of x, 26:45743
- radioinduced mutations in human fibroblast, effects of dose on frequency of x, 26:45740
- CELLULOSE**
- radioinduced graft polymerization of butyl acrylate--styrene mixtures on, 26:45320
- CELLULOSE ACETATES**
- radioinduced graft polymerization of butyl acrylate--styrene mixtures on, 26:45320
- cellulose acetate--polyacrylonitrile, mechanical properties and structure of, produced by γ -radioinduced grafting, 26:46084
- CENTRAL AMERICA**
- radioisotope content of milk in, during Mar. 1971 to Mar. 1972, cesium-137 and strontium-90, 26:45552
- CENTRIFUGES**
- fabrication of ultra-, use of carbon composites in, 26:46086
- rotor for ultra-, design for isotope separation in fluids, 26:45697(P)
- CEPHARANTHINE**
- radioprotective effect on leukocyte count in man, 26:45805
- CEREALS**
- fallout cesium-137 and strontium-90 content in, Mecklenburg, GR, 1969, 26:45546
- flour, radioisotope content of, in France during 1960 to 1969, cesium-137 and strontium-90, 26:45903
- CERIUM**
- determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
- determination in hair by neutron activation and γ spectroscopy, 26:45214
- determination in presence of manganese by solvent extraction using TTA and spectrophotometry, 26:45226
- effects on grain boundary energy of nickel, 26:45999
- separation from other metals by combined ion exchange--solvent extraction chromatography, 26:45282
- solubility in oxygen-containing chromium at 1000 to 1800°C, microhardness studies on, 26:46034
- solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
- thermal capacity of liquid, at 1100 to 1600°K, 26:45998
- thermal conductivity of liquid, at 1100 to 1600°K, 26:45998
- CERIUM ALLOYS AND SYSTEMS**
- Al--Ce--La, critical field curves of Kondo superconducting, 26:46832
- Ge--La, critical-field and proximity-effect behavior of superconducting, many-body effects from, (E/T), 26:46822
- CERIUM ISOTOPES Ce-139**
- electron-capture decay of, γ -x-ray coincidence spectra from, (E), 26:46947
- CERIUM ISOTOPES Ce-140**
- energy level at 2083 keV, g-factor for, (E), 26:46958
- CERIUM ISOTOPES Ce-141**
- content in atmosphere in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
- content in surface atmosphere at world sites, 1965 through Jan. 1968, 26:45512(R) (HASL-257(App.))
- range of fission fragments of, in aluminum--uranium alloys and uranium oxides, 26:46074
- CERIUM ISOTOPES Ce-144**
- accumulation by organisms and plants in littoral freshwater microecosystem, 26:45553
- content in atmosphere, milk, soils, and vegetation in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
- content in fallout deposited on Dallas, Texas, soil during Jan. to July 1964, (E), 26:45527 (TID-26103)
- content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
- content in surface atmosphere at world sites, 1963 through Feb. 1972, 26:45512(R) (HASL-257(App.))
- effects on gastrointestinal tract following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
- effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
- CERIUM OXIDES**
- electric conductivity of CeO₂/sub 2-x/ at 16 to 1998°C, 26:47247 (BNWL-1511)
- radioinduced hole centers in CeO₂ crystals, ESR spectra of uv-, 26:47233
- CESIUM**
- determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
- determination in animal, plant, and soil samples by neutron activation, 26:45193 (TID-25954, pp 376-8)
- determination in plant leaves by neutron activation, 26:45192 (TID-25954, pp 373-5)
- mass flow choking limit for heat pipe working fluid of, (E), 26:45460
- plasma of, impedance of partially ionized, (E/T), 26:47135
- reactions with bromine, chlorine, and oxygen molecules at 50 to 350 eV, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366
- separation from liquid wastes using colloidal suspension, 26:45384(P)
- solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
- uptake of radioactive, by larval and hatching amphibians and reptiles, 26:45815(R) (COO-3486-1)
- CESIUM FLUORIDES**
- scattering by electron beams at 0.6 to 6.5 eV, molecular-beam recoil measurements of absolute differential and total cross sections for, 26:46406
- CESIUM IONS**
- implantation in nickel at 1 to 10 keV, penetration and spatial distribution from, 26:47235
- CESIUM ISOTOPES Cs-131**
- energy levels from barium-131 electron-capture decay, spin--parity assignments for, (E), 26:46963
- energy-level transitions from barium-131 electron-capture decay, multipolarities of, (E), 26:46963
- CESIUM ISOTOPES Cs-133**
- neutron reactions (n, α) with, calculation of excitation functions for fast, 26:47066
- CESIUM ISOTOPES Cs-134**
- content in biota, sediments, and water in Lake Michigan near Big Rock Point power reactor during 1970, 26:45536
- content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
- effects on developing frogs following chronic exposure, 26:45913(R) (RLD-2227-T-8-3)
- CESIUM ISOTOPES Cs-137**
- accumulation and loss of, by marine clams, model for food chains from kinetics of, 26:45545 (UCRL-73744)
- content and distribution in open ocean silts from Atlantic Ocean and Mediterranean Sea, 26:45531 (COO-3563-3)
- content in atmosphere and man in Argentina, 1960 through 1970, 26:45520 (NP-19357)
- content in atmosphere in Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
- content in atmosphere, milk, soils, and vegetation in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
- content in biota, sediments, and water in Lake Michigan near Big Rock Point power reactor during 1970, 26:45536
- content in cereals, coffee, fruits, milk, and tea in Mecklenburg, GDR, 1969, 26:45546
- content in drinking water in New York City, 1965 through 1971, 26:45512(R) (HASL-257(App.))
- content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)
- content in human diets in Chicago, Jan. to June 1972, 26:45540 (HASL-257, pp III.45-52)
- content in marine organisms in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)
- content in muscles of broilers, hens, and partridges in Warsaw, Poland area, 1968 to 1969, 26:45548
- content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
- content in surface atmosphere at world sites, 1963 through Feb. 1972, 26:45512(R) (HASL-257(App.))
- content in surface atmosphere in western US, 26:45517 (HASL-257, pp I.18-32)
- content in Western Hemisphere milk samples, during Mar. 1971 to Mar. 1972, 26:45552
- content of flour and wheat in France during 1960 to 1969, 26:45903
- content stratosphere to July 1971, effects of nuclear weapons testing on, 26:45518 (HASL-257, pp I.33-50)
- deposition at Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
- determination in soils, 26:45263 (RT/PROT-72-11)
- effects on gastrointestinal tract following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
- effects on mice following chronic ingestion, effects on litter size and reproduction span of, effects of concentration on, 26:45900
- encapsulation of waste, remote welding methods for, 26:46099 (ARH-SA-117)
- solubility of Rainier Event rubble-contained, measurement of long-term, 26:45482 (NVO-1229-177)
- transport through soil-plant-small mammal ecosystems at Nevada Test Site, 26:45914 (TID-25954, pp 58-64)
- CESIUM ISOTOPES Cs-141**
- beta decay of, γ spectra and half-life for, (E), 26:46971
- CESIUM SODIUM AMERICIUM CHLORIDES**
- magnetic properties of, 26:47272
- CESIUM SODIUM LUTETIUM CHLORIDES**
- magnetic properties of curium- and berkelium-doped, 26:47272

magnetic properties of, 26:47272

CESIUM SODIUM URANIUM CHLORIDES

magnetic properties of, 26:47272

CESIUM URANIUM CHLORIDES

 Cs_2UCl_6 -- UCl_4 , spectra of, effects of chemical state of uranium on, 26:45377 CsUCl_5 -- UCl_4 , spectra of, effects of chemical state of uranium on, 26:45377

CHARGED-PARTICLE BEAMS

interactions with afterglow plasma, wave coupling in, (E), 26:47193

CHARGED PARTICLES

acceleration in H_{111} cylindrical cavity with axial magnetic drive field, 26:46139

acceleration in solar atmosphere, effect of single- and multiple-ionization losses on, (T), 26:46230

detection at 1 GeV, development of proportional counter for, p. 160-3, 26:46873(R) (CEA-N-1522)

detection of high-energy, design of large-angle spectrometer for, p. 165-6, 26:46873(R) (CEA-N-1522)

interactions with electromagnetic field, algebraic Poincare group algebra for, 26:46678

motion in electromagnetic fields, regularization in vacuum polarization calculations of vector currents for, 26:46742

motion in two-dimensional neutral sheet with linear magnetic field variation, Hamiltonian methods for oscillation frequency and analytic treatment of perturbed, (T), 26:46522 (N-72-17763)

motion of spinning, in gravitational and electromagnetic field, model for, (T), 26:46767

CHEMICALS AND REAGENTS

availability and cost of radioactive, produced by Norway Institute for Atomic Energy, 26:45709 (INIS-mf-256)

CHEMISTRY

research progress on, India, 1970-71, 26:45311(R) (BARC-599)

CHERENKOV RADIATION

emission by tachyons, Lorentz-invariant formulation for, (T), 26:46704 (N-72-19766)

CHLORIDES

reactions of molten, with zinc iodides, 26:45421

solvation numbers from compressibilities and ionic

vibration potentials measurements, 26:45369

CHLORINE

cesium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366

determination of, chemical procedures for, 26:45181 (CISE-R-172)

potassium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366

CHLORINE FLUORIDES

determination in nitrogen by automatic absorption spectrometry, 26:45243(P)

CHLORINE ISOTOPES Cl-33

beta decay of, half-life for, (E), 26:46902 (BARC-598)

energy levels from proton resonances in sulfur-32,

isospin and strengths of, (E), 26:46902 (BARC-598)

energy-level transitions from proton resonances in sulfur-32, branching ratios and multipolarities for, (E), 26:46902 (BARC-598)

CHLORINE ISOTOPES Cl-34

energy levels from chlorine-35 (p,d) reactions, spin-parity assignments, spectroscopic factors, and shell-model interpretations for, (E/T), 26:46906

CHLORINE ISOTOPES Cl-35

meson (K^-) elastic scattering by, scattering lengths for, (E), 26:46912

proton reactions (p,d) at 40 MeV, DWBA analysis of angular distributions for, (E), 26:46906

CHLORINE ISOTOPES Cl-36

energy levels from chlorine-37 (p,d) reactions, spin-parity assignments, spectroscopic factors, and shell-model interpretations for, (E/T), 26:46906

CHLORINE ISOTOPES Cl-37

proton reactions (p,n) at 3.7 and 3.9 MeV, angular distributions from, (E), 26:46929

CHLORINE ISOTOPES Cl-38

beta decay of polarized, asymmetry of, (E), 26:46911

nuclear magnetic moment from NMR measurements, (E), 26:46911

nuclear magnetic resonance in sodium chlorides, (E), 26:46911

CHLOROFORM

radioinduced reaction with cyclohexanone, yields from γ , 26:45335

CHROMATES

separation from cooling water by ferrous sulfate reduction and lime precipitation, 26:47619 (KY-642)

CHROMIUM

alpha reactions with natural, production of carrier-free manganese-54 from, 26:45394

determination in milk powder by thermal neutron

activation and radiochemical separation, 26:45196

determination of trace amounts of, in germanium by

instrumental neutron activation, 26:45218

determination of, chemical procedures for, 26:45181 (CISE-R-172)

effects on mechanical properties of aluminum--iron--uranium-base alloys at 20 to 640°C, 26:45995 (UJV-2802-M)

effects on recrystallization of iron, 26:45982

helium-3 reactions with natural, production of carrier-free manganese-54 from, 26:45394

neutron elastic scattering at 1.77 to 2.76 MeV, angular distributions for, (E), 26:46916 (AE-452)

separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281

solvent properties of oxygen-containing, for cerium, neodymium, and praseodymium at 1000 to 1800°C,

microhardness studies on, 26:46034

CHROMIUM ALLOYS AND SYSTEMS

Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009

Al--Cr--Ni--ThO₂, properties of dispersion-strengthened, for use in space shuttle, 26:45934 (N-72-18585)

Al-base--Cr, superconductivity and localized spin fluctuations in, (E/T), 26:46838

Cr--Tb, hardness and metallography of, 26:46033

Cr--Tb, phase diagram at 600 to 1900°C, 26:46033

CHROMIUM ISOTOPES Cr-50

helium-3 reactions (He-3,t) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930

proton elastic and inelastic scattering at 6 MeV, proton spectra and angular distributions for, (E), 26:46922

CHROMIUM ISOTOPES Cr-51

distribution in tissues of newts, 26:45913(R) (RLO-2227-T-8-3)

reactions of dopant-atoms and recoil atoms of, in potassium chromate, thermal annealing, 26:45364

retention in lungs of mice following intratracheal injection as oxide, effects of respiratory infection on effective half-life for, 26:45910 (CONF-720505-2)

CHROMIUM ISOTOPES Cr-52

neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909photon reactions (γ,n) and (γ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

proton elastic and inelastic scattering at 6 MeV, proton spectra and angular distributions for, (E), 26:46922

CHROMIUM ISOTOPES Cr-53

proton reactions (p,n) at 2.8 to 4.0 MeV, angular distributions and total cross sections for, (E), 26:46929

CHROMIUM ISOTOPES Cr-54

proton elastic and inelastic scattering at 6 MeV, proton spectra and angular distributions for, (E), 26:46922

CHROMIUM OXIDES

density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949

 Al_2O_3 -- CaO -- Cr_2O_3 -- F -- MgO -- MoO_3 -- TiO_2 -- ZrO_2 , use as welding flux, 26:45941(P)

CHROMOSOMES

radiation effects on, effects of iodine-125 and tritium decay on, in eukaryote cells, 26:45894 (ORO-2690-12(Pt.B), Paper 14)

radioinduced aberrations of, in cultured Chinese hamster ovary cells, effects of post-exposure temperature on repair of x, 26:45743

radioinduced aberrations of, in cultured lymphocytes from Marshall Island human population, combined effects of age and sex on frequency of, 26:45792

radioinduced aberrations of, in human bone marrow cells following single local exposure, effects of dose and post-exposure time on frequency of γ , 26:45794

radioinduced breaks in, of human bone marrow cells, use for determining protective effects of taurine, 26:45734

radioinduced non-disjunction in *Drosophila*, quantitative study of x, 26:45880radioinduced translocations of, in male *Drosophila*, effects on frequency of crossing over in chromosomes of F_1 females, x, 26:45876

CHROMOSPHERE

heating of transition region with corona, (E/T), 26:46228

structure of transition region with corona, (E/T), 26:46228

CIRCUITS

design for continuously finding ratio of two pulse trains, 26:45675

CIRCULATORY SYSTEM

power requirements for human, calculation of, 26:45700(R) (COO-2155-4)

CLINTON P. ANDERSON MESON PHYSICS FACILITY

development of, origins and history of, 26:46119 (LA-5000)

COAL

analysis for ash content by neutron activation determination of silicon and aluminum, 26:45195

COAL DEPOSITS

radioactivity in Pechora River basin, distribution and genesis of, 26:45505

uranium content in Pechora River basin, distribution and genesis of, 26:45505

COATINGS

analysis of irradiated fuel element, for oxygen,
development of remote apparatus for, 26:45224 (LA-DC-72-524)

COBALT

carbon-ion reactions at 10 MeV/amu with, energy shift and relative intensity of K x rays from, (E), 26:46339
compatibility with refractory metal fuel capsules at 850 to 1000°C, 26:45701(R) (DP-1300)
corrosion by marine environment, effects of nickel plating on, 26:45701(R) (DP-1300)
determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
determination of, chemical procedures for, 26:45181 (CISE-R-172)
diffusion in delta and epsilon phases in plutonium, 26:46024
effects on recrystallization of iron, 26:45982
ion reactions at 5 MeV/amu with, dependence of energy shift and relative intensity of K x rays from, on ion stopping power, (E), 26:46339
separation from other metals as β -diketonates by column liquid-liquid chromatography, 26:45281

COBALT ALLOYS AND SYSTEMS

production of wrought superalloys of, vacuum induction melting method for, 26:45981
Al-base--Co, superconductivity and localized spin fluctuations in, (E/T), 26:46838
Co--Fe--W, crystal structure of, during aging, effects of molybdenum on, 26:46006
Co--Pt, microscopy of, imaging process in field ion, 26:46038
Co--TiC--WC, metallography of, methods for, 26:45955
Co--W, deposition of, effects of current density and temperature on electrolytic, 26:45977
Co--W, electrodeposition of, effects of temperature on, 26:45983
Co--WC, metallography of, methods for, 26:45955
Co-base--Sm, sintering of Co₂Sm permanent magnet alloys, mechanism of, 26:45976

COBALT ALUMINATES

corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)

COBALT ISOTOPES Co-53

decay of isomeric, proton spectra and half-life for, (E), 26:46925
proton decay of isomeric, partial half-life and branching ratio for, (E), 26:46919
proton decay of isomeric, proton spectra and half-life for, (E), 26:46926

COBALT ISOTOPES Co-57

Mössbauer spectra in β -rhombohedral boron, 26:47280
use of bleomycin-incorporated, for localization of neoplasms, 26:45929

COBALT ISOTOPES Co-59

alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932
alpha reactions (α, p) at 17 MeV, proton spectra from, (E), 26:46932
energy-level density from alpha and proton reactions, (E), 26:46932
neutron reactions (n, p) with, calculation of excitation functions for fast, 26:47066
proton inelastic scattering at 13 MeV, proton spectra from, (E), 26:46932
proton reactions (p, α) at 13 MeV, α spectra from, (E), 26:46932
proton reactions (p, n) at 2.4 to 4.0 MeV, angular distributions and total cross sections for, (E), 26:46929

COBALT ISOTOPES Co-60

accumulation and loss of, by marine clams, model for food chains from kinetics of, 26:45545 (UCRL-73744)
availability and cost of radiation sources of, produced by Norway Institute for Atomic Energy, 26:45709 (INIS-mf-256)
beta decay of, intensity ratio of 2158- and 2505-keV γ -rays from, (E), 26:46933
content in biota, sediments, and water in Lake Michigan near Big Rock Point power reactor during 1970, 26:45536
content in Pearl Harbor marine environment during 1966 and 1968, 26:45534
content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
heat sources of, data on AEC loan program for, 26:45701(R) (DP-1300)
use as tracer for measurement of blast furnace lining wear, 26:45720

COBALT OXIDES

corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
CoO--MgO, corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)

COINCIDENCE CIRCUITS

description of $4\pi\beta\gamma$, with reduced dead time loss, 26:45589 (JEN-247)
description of fast-slow, using NaI(Tl) and Ge(Li)

detectors, 26:45639

design of priority, p. 81, 26:46873(R) (CEA-N-1522)
design of, p. 154-6, 26:46873(R) (CEA-N-1522)

COLLAGEN

structure of, uv spectral studies of, 26:47642 (UR-3490-47)

COLLIMATORS

description for scintillation camera for improved peripheral response, 26:45596(P)
description of convergent multihole focusing, for scintillation camera, 26:45602

COLLISION PROCESSES

amplitudes of atomic, bounds on second term of Born series and regularization of Born approximation for, 26:46377
exponential approximations for rotationally inelastic, relations between distorted-wave, first-order sudden, and infinite-order sudden, 26:46369
rotational energy transfer in, measurement using acoustic absorption and dispersion, 26:46376

COLORADO

leasing of USAEC-controlled uranium-bearing lands in, environmental impact report on, 26:45572 (WASH-1523)

COLUMBIA RIVER

zinc-65 content in mussels in, following shutdown of Hanford reactors, 26:45543 (RLD-2225-T-1-9)

COMETS

dust particles of, size of, (T), 26:46220

COMPOSITE MATERIALS

development and uses of carbon, 26:46086

COMPOSITE PARTICLES

coupling of massless composite lepton-antilepton states to free leptons, (T), 26:46570 (IC-72/54)
production of massless lepton-antilepton states from exchange of virtual neutrino pairs in anti $l + l$ interactions, (T), 26:46570 (IC-72/54)
spin of massless lepton-antilepton states from exchange of virtual neutrino pairs in anti $l + l$ interactions, (T), 26:46570 (IC-72/54)
wave functions of, relativistic, 26:46506

COMPUTER PROGRAMS

catalog of LLL documentation reports on, 26:47598 (UCID-30047)

CRIP, for thermohydraulic analysis of CIRENE cooling system, FORTRAN IV for IBM 7040, 26:47309 (CISE-R-257)

DECAT, for long term pressure and temperature transients in pressure suppression containment following loss of coolant accident, MIMIC for IBM 7094, 26:47318 (RT/PROT-72-4)

FLUB, for neutron flux distribution in reactors, FORTRAN IV, 26:47594 (BMW-FBK-72-6)

for Fortran translation table setup, 26:47621 (ORNL-TM-3810)

GAMANAL, for γ spectral analysis, 26:45265 (UCRL-51061(Vol.1))

HENRE, for calculation of gamma radiation and neutron escape rates from 24-kt nuclear explosions, 26:45497

PDE, for calculation of thermal-hydrodynamic conditions in steam-water circuits, FORTRAN IV for IBM 7040, 26:45458 (CISE-R-245)

PECVAR, for thermohydraulic analysis of CIRENE cooling system, FORTRAN IV for IBM 7040, 26:47309 (CISE-R-257)

SHAKER, for numerical analysis of weighted tabular data, in FORTRAN, 26:47631 (UCRL-51210)

COMPUTERS

interfaces for CII 10020 and Varian 620i, design of, 26:46873(R) (CEA-N-1522)

programming for simulation of radioinduced atomic displacement cascades in metals, 26:46061 (CONF-710601-, pp 739-56)

uses in activation analysis, review on, 26:45211

CONCRETES

cascade shower production in, high energy intranuclear hadron, (E/T), 26:46126(T) (ORNL-tr-2617)
concretes--polymers, properties of radiation-polymerized, for use in low-porosity pipes, 26:45718

CONES

impact attenuation equation for, longitudinal, 26:47626(T) (SC-T-722520)

CONNECTICUT YANKEE POWER REACTOR

environmental report for, aquatic cost-benefit analysis for, 26:47397 (DOCKET-50213-152)
operations during June 1972, 26:47343 (DOCKET-50213-154)
site for, environmental radiation survey for July through December 1971, 26:47396 (DOCKET-50213-150)

CONNECTORS (ELECTRIC)

design of removable, for superconducting magnet power supplies, 26:45679

CONVEYERS

design and operation of irradiator, for continuous irradiation of packaged materials, 26:45447(P)

COOLING TOWERS

liquid droplets in effluents from, method for measuring size of, 26:45574 (CONF-720619-1)

COOPER POWER REACTOR

auxiliary cooling system for, modifications for building, 26:47467 (DOCKET-50298-47)
battery for, ventilation system for, 26:47467 (DOCKET-50298-47)
environmental report for, requested additional data for, 26:47465 (DOCKET-50298-43)
environmental report on, additional data for, 26:47468 (DOCKET-50298-48)

fire protection system for, modification for, 26:47467 (DOCKET-50298-47)
safety of, requested additional data on, 26:47466 (DOCKET-50298-46)
spent fuel storage pool for, proposed modifications for, 26:47467 (DOCKET-50298-47)

COPPER

complexes with phthalocyanine, neutron reactions with, radiation and quenching effects on chemical behavior of recoil copper-64 from, 26:45356
determination in high-purity cadmium oxide, spectrographic, 26:45220 (BARC-593)
determination in solutions of copper-64, spectrographic, 26:45223 (JEN-232-DQ/I-81)
determination of, chemical procedures for, 26:45181 (CISE-R-172)
diffusion and outgassing of ion implanted-helium in, effects of temperature on, 26:46050 (CONF-710601-, pp 230-47)
effects on recrystallization of iron, 26:45982
gamma scattering at 0.662 MeV by, absolute differential cross sections for, (E), 26:47204
meson (K^-) capture by, gamma spectra from, (E), 26:46940
neutron coherent scattering below 10^{-4} eV, scattering length for cold, (E), 26:46853
neutron transmission below 10^{-4} eV through films of, interference effects in cold, (E), 26:46853
photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
proton reactions at 24 GeV/c, w , K , p , and anti p spectra from, (E), 26:46459
radiation effects on lattice parameter and electric conductivity of, annealing of 60K 2.6-MeV electron, 26:47231
radiation effects on length of, γ , 26:46075
radiation effects on mechanical properties of, at low temperatures, 10-MeV proton, 26:46066(R) (COO-3431-1)
radioinduced atomic displacement cascades in, statistical analysis of fast neutron, 26:46061 (CONF-710601-, pp 739-56)
radioinduced defect aggregates in, relations of temperature with fast neutron, 26:46060 (CONF-710601-, pp 672-83)
recrystallization of, effects of impurities on, 26:45982
separation from other metals as β -diketonates by column liquid-liquid chromatography, 26:45281
separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285

COPPER ALLOYS AND SYSTEMS

analysis for tin by alternating current polarographic method, 26:45256
Cu-Mg, dislocation energy of impurity atoms in solid solutions of, internal friction method for studying, 26:46023
Cu-base--In, precipitation in, mechanisms of cellular, 26:46017

COPPER ISOTOPES

muons (μ^-) reactions (μ^-, e^+) with, upper limit on branching ratio relative to ordinary capture for, (E), 26:46941
photon reactions (γ, π^0) at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
photon reactions at 1.160 GeV at 150°, photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441

COPPER ISOTOPES Cu-61

energy levels from nickel-58 (α, p) reactions, spin assignments for, (E), 26:46934

COPPER ISOTOPES Cu-63

alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932
energy levels from nickel-60 (α, p) reactions, spin assignments for, (E), 26:46934
energy-level density from alpha and proton reactions, (E), 26:46932
proton inelastic scattering at 13 MeV, proton spectra from, (E), 26:46932
proton reactions (p, α) at 13 MeV, α spectra from, (E), 26:46932

COPPER ISOTOPES Cu-64

analysis of solutions of, for calcium-45 and target element, spectrographic, 26:45223 (JEN-232-DQ/I-81)
chemical behavior of recoil atoms of, in neutron-irradiated copper phthalocyanine, effects of preirradiation quenching on, 26:45356
radiation effects on chemical behavior of recoil atoms of, in neutron-irradiated copper phthalocyanine, 26:45356

COPPER ISOTOPES Cu-65

alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932
energy levels from nickel-62 (α, p) reactions, spin assignments for, (E), 26:46934
energy-level density from alpha and proton reactions, (E), 26:46932
proton inelastic scattering at 13 MeV, proton spectra from, (E), 26:46932
proton reactions (p, α) at 13 MeV, α spectra from, (E), 26:46932

COPPER ISOTOPES Cu-67

energy levels from nickel-64 (α, p) reactions, spin assignments for, (E), 26:46934

CORUNDUM

radiation effects on crystals of, thermal-neutron, γ , and x , 26:47244

COSMIC GAMMA SOURCES

detection of, (E), 26:46154

COSMIC PARTICLES

charge composition of very heavy primary, (E), 26:46151
detection of heavy, delta-ray theory for, 26:45612
energy spectra of very heavy primary, (E), 26:46151
interactions of ultrahigh-energy, with cosmic microwave background, (T), 26:46257
modulation of primary, integration of steady Fokker--Planck equation for solar, (T), 26:46152
neutron angular distribution and spectrum above 5×10^{11} eV, (E), 26:46153
nucleon reactions with complex heavy nuclei at high energies, pion spectra and energy transfer to π^0 for study of collective, (E), 26:46461
spectra and modulation near solar maximum, 20-MeV to 15-GeV electron, (E), 26:46325

COSMIC RADIATION

anisotropy of primary, relation of interplanetary magnetic field structure to equatorial, 26:46158
charge spectrum of solar, from nuclear emulsions and Explorer-35 solid-state detector, (E), 26:46330
dose rate to man during high-altitude supersonic flight, effects of solar flares on, 26:45806
gamma background, origin of, (E), 26:46154
hazards to astronauts during space flight, 26:45812
heliocentric gradient of galactic, (E), 26:46155
modulation at rigidities > 2 GV in Aug. 1969, rigidity dependence of, (E/T), 26:46327
observations from HEOS-1 launched Dec. 5, 1968, (E), 26:46334
variations of primary, method of analysis and statistical distribution of diurnal, 26:46159

COSMIC RADIO SOURCES

map of 3C 272.1 at 2.7 and 5.0 GHz, (E), 26:46200
nomenclature for variable, VRS rather than VRO, 26:46209
polarization aperture synthesis of five galactic nonthermal, (E), 26:46207
positions of small-diameter, at 408 MHz, (E), 26:46202
variability at 1665 MHz in W3, (E), 26:46205

COSMIC SHOWERS

detection of large, by atmospheric fluorescence, (E), 26:46150 (NYO-3038-13)

COSMIC X-RAY SOURCES

flare in Cyg X-1, observation of hard x-ray, (E), 26:46211
optical counterpart for Cen X-3, search for, (E), 26:46213
origin of 2U 1543-47, extragalactic, (E/T), 26:46215
origin of transient, extragalactic, (E/T), 26:46215
relation to bright stars, (E), 26:46212

COSMOLOGY

fine-structure constant in, time variation of, (T), 26:46254 (COO-1112-51)
mass difference of muon and electron, cosmological origin of, (T), 26:46259
model formulation for rotating Universe with flat space in, Einstein field equations for homogeneous anisotropic, 26:46261
model with positive- and negative-mass particles, for interacting galaxies and gravitational screening, 26:46761
models of, implications of recent observations for, (T), 26:46256
models with primordial magnetic field, exact solutions of Einstein equations for gravitational fields with sources of dust and magnetic field for anisotropic, (T), 26:46755
reference system in, absolute, (T), 26:46257
relativistic gravitational field equations for clusters of galaxies, (T), 26:46253
symmetry problems in, 26:46674

CRAB NEBULA

gamma emission from, detection of, (E), 26:46214
p-CRESOL, 2,6-DI-tert-BUTYL--effects of dietary, on rate of lipid peroxidation in tissues of whole-body x-irradiated mice, 26:45872

CRITICAL ASSEMBLIES

development of Ermine, history of, 26:47589
development of Harmonie, summary of, 26:47590
development of Masurca, history of, 26:47592
kinetics of ZPR-9 EMC-BOL, computer program calculation of three dimensional, 26:47586 (HEDL-TME-72-42)
physics measurements for fast, effects of fission neutron spectra on, 26:47585 (EURFNR-1008)
physics measurements in MASURCA, for plutonium--graphite and uranium--graphite mixtures, (E), 26:47581 (CEA-N-1483)

CROLOY

microstructure and tensile properties of, effects of long-term tempering on, 26:45994(R) (UCLA-34-P-177-10)

CRUSTACEA

bosmina, population in Lake Ontario, effects of thermal effluents from nuclear power plants on, (E), 26:45578

daphnia retrocurva, populations in Lake Ontario, effects of thermal effluents from nuclear power plants on, (E), 26:45578
 radioisotope content in, of Puget Sound basin, 26:45544 (RL0-2225-T-14-4)

CRYSTAL STRUCTURE
 lattice dynamics, (T), 26:47617 (IEA-19)

CRYSTALLOGRAPHY
 theory of martensite, relation of Burgers and Bogers--Burgers transformation mechanisms to, 26:46037

CRYSTALS
 book: Radiation Physics of Crystals and p-n Junctions (In Russian), 26:47214
 electron diffraction from imperfect, modified Bloch wave expansion for formulation of theory for high energy, (T), 26:46510 (CALI-767-P-3-22)
 electron interactions with niobium and silicon, at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550
 electron microscopy of perfect and imperfect, dynamic theory of backscattering in scanning, 26:47267
 entropy of small, determinant method for calculation of, 26:47632 (UCRL-51227)
 growth of, bibliography of organizations and projects for, 26:45973 (ORNL-RMIC-12)
 ion channeling in, review of, 26:47228
 positron annihilation in liquid, positronium-phase and temperature dependence of lifetimes in, (E), 26:46363
 positron interactions with niobium and silicon, at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550

CURIUM IONS
 magnetic properties of cesium sodium lutetium chloride doped with, 26:47272

CURIUM ISOTOPES Cm-243
 fission of, angular distribution of fragments from spontaneous, (E), 26:47022 (JINR-P7-6466)

CURIUM ISOTOPES Cm-244
 analysis for impurities by spark source mass spectrometry, 26:45260
 production of, mass spectrometric analysis for quality control, 26:45260
 properties for use in isotope-fueled thermoelectric converters, 26:45703(R) (ORNL-4796)

CURIUM ISOTOPES Cm-245
 energy-level transitions from californium-249 alpha decay, branching ratio and total conversion coefficients for, (E), 26:47034
 neutron fission of, mass yields from thermal, (E), 26:47027

CURIUM OXIDES
 fabrication of radioisotope fuels of, use of thorium oxide as stand-in for, 26:45703(R) (ORNL-4796)
 inspection of heat source of, following operation at high temperatures, 26:45703(R) (ORNL-4796)

CURRENT ALGEBRA
 axial-vector form factor for weak interactions, mass limitations for, (T), 26:46573
 PCAC theory in, gravitational correction to, 26:46501
 quark propagator in light-cone, implications for existence of quarks, 26:46604
 second-class currents in, upper bound for tensor component of, (T), 26:46573
 SU(1,1) classification for wave packet realization of lightlike particle states for formulation of parton model for hadrons, 26:46589
 SU(1,1) representations for wave packet realization of lightlike particle states for formulation of parton model for hadrons in, 26:46681 (NYO-4076-19)

CYANOGEN
 spectral lines in galactic nuclei due to, (E), 26:46243

1,3-CYCLOHEXADIENE
 effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331

CYCLOHEXANONE
 radioinduced reaction with chlorinated hydrocarbons, yields from γ , 26:45335

CYCLONE SEPARATORS
 design for concentration of radioactive solutions, 26:46111(P)
 performance of steam, analysis by similitude laws, 26:45442

CYCLOTRONS
 beam extraction system of Ukrainian Inst. of Nuclear Research 240-cm isochronous, design of magnetic shield for, 26:46129
 beam extraction system of Ukrainian Inst. of Nuclear Research 240-cm isochronous, design of compensated current-driven magnetic channel for, 26:46128
 description of Technical University of Eindhoven, 26:46146
 experimental facilities at Technical University of Eindhoven, survey of, 26:46146
 ion source for multicharged ions of solid materials, development of, 26:46133

CYLINDERS
 impact attenuation equation for, longitudinal, 26:47626(T) (SC-T-722520)
 joints with nozzles, method for calculation of stress

distribution in, 26:45445

CYSTAMINE
 effects on respiratory exchange in rats, 26:45823

CYSTEAMINE
 effects on respiratory exchange in rats, 26:45823

CYSTEINE
 effects on recovery of glucose-6-phosphatase activity in erythrocytes of whole-body x-irradiated rabbits and rats, 26:45848
 protective effects on barley seeds against mutagenic and toxic effects of ethylenimine, 26:45772

D

DATA ACQUISITION SYSTEMS
 description of on-line, for (γ ,xn) experiments, p. 274-6, 26:46873(R) (CEA-N-1522)
 development for studying thermal effluent dispersion in surface waters, 26:45579
 development of multicomputer, for physics research centers, review on, 26:45686

DATA PROCESSING SYSTEMS
 development of Illiac III-computer based, for interactive picture processing, 26:47610(R) (COO-2118-34)
 development of multicomputer, for physics research centers, review on, 26:45686

DATA TABULATIONS
 log-ft values for beta decay, 26:46970
 on energy levels for A = 18 to 20 nuclei, 26:46882
 on mean excitation energies of daughters in β^- decay, 26:46875 (KFK-EXT-10/72-1)
 on nuclear data from fission products, 26:47012 (CEA-N-1526)

DAVIS-BESSE POWER REACTOR
 containment building equipment hatch for, shipping damage to, 26:47326 (DOCKET-50346-30)
 environmental report for, revised and additional data for, 26:47518 (DOCKET-50346-29)
 piping for, corrected inspection and certification procedures for component cooling and service water, 26:47517 (DOCKET-50346-27)

DECANE, 1-BROMO-
 electron reactions with, cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)

DECANOIN
 tri-, radiolysis of, reaction kinetics in production of yields from, 26:45321

DECAY SCHEMES
 book: Isobaric Nuclei with Mass Number A = 72 (In Russian), 26:46939

DENSIMETERS
 design of gamma radiometric, for locating liquid-liquid interfaces, 26:45659(P)

DENSITOMETERS
 description for dielectric track detectors, 26:45598

DESALTING
 equipment for seawater, design of heat exchanger, 26:45436 (CEA-CONF-2010)
 membrane studies for, ion-control, 26:45405(R) (ICP-1006)
 method for large-scale, status of, 26:47579

DETONATION WAVES
 characteristics of gelled-slurry blasting agents, large-scale tests of, 26:45484 (SC-DC-722296)

DEUTERIUM
 deuteron reactions (d,pn) at 27.5 MeV, analysis of p-n final-state interaction in, (E), 26:46860 (LYCEN-7186)
 effects on abstraction of hydrogen from thiols by free radicals, isotope, 26:45342
 effects on dissociation of picric acid, solvent transfer and isotopic exchange isotopic, 26:45292
 effects on ESR spectra of copper complex hydrates in ethanol-water systems, solvent isotopic, 26:45296
 effects on fluorescence decay time and luminescence intensity of europium ions in aqueous solutions, solvent isotopic, 26:45295
 effects on hydrogen abstraction in photolysis of nitrites of epimeric alcohols, isotope, 26:45300
 effects on kinetic energy release in metastable ion fragmentations, isotopic, 26:45301
 effects on quenching by carbon dioxide of infrared chemiluminescence from vibrationally excited hydrogen fluoride, isotopic, 26:45298
 effects on radiolysis of single crystals of malonic acid, isotope, 26:45343
 exchange in deuterium hydrides in shock tube facility, isotope, 26:45306
 exchange reactions with hydrogen and hydrochloric acid in shock tube facility, isotope, 26:45306
 exchange reactions with hydrogen in cytochrome c at 20 to 38°C, 26:45305
 exchange with propylene over zinc oxide, mechanisms of, 26:45303
 helium-3 elastic scattering at 17.5 to 44.1 MeV, angular distributions for, (E), 26:46871
 helium-3 reactions ($\text{He-3}, \alpha$) at 17.5 to 44.1 MeV, angular distributions for, (E), 26:46871
 helium-atom scattering by molecular, coupled-channel calculations of effects of molecular symmetry on cross sections for, 26:46379
 hyperfine structure of ground-state, pressure shift in argon, (E), 26:46347

meson reactions (π^+ , 2p) at 76 MeV, cross section for, 26:46870

neutron elastic scattering at -100 to +100°C, total cross sections for cold and thermal, (E), 26:46887

neutron elastic scattering below 3.34 MeV, effective-range analysis of, (T), 26:46864

neutron reactions (n,p) at 50 MeV, proton spectra from, (E), 26:46861

plasma of, parallel magnetic multipole confinement of, (E), 26:47111

polarizability of molecular, molecular-beam magnetic-resonance measurement of anisotropy of static electric, 26:46358

proton elastic scattering at 14 to 17.6 MeV, differential cross sections for, (E), 26:46863

proton reactions (p, γ) at 10 to 17.5 MeV, excitation function for, (E), 26:46868

separation from hydrogen in formaldehyde by photopredissociation, 26:45307

separation from liquid mixtures of deuterium sulfide and hydrogen sulfide by distillation, 26:45690(P)

DEUTERIUM IONS

reactions with atoms, classical calculations of cross sections for single-electron capture into ground and excited states in, 26:46398

DEUTERIUM SULFIDES

D₂S--H₂S, distillation of liquid, for production of deuterium, 26:45690(P)

DEUTERIUM SYSTEMS

D--T, breeding ratios and inventory calculations for use in thermonuclear reactors, 26:47173

DEUTERONS

effects on survival of cultured mammalian cells, relation of mathematical analysis of, to radiotherapy, 26:45735

effects on void formation in Types 304 and 321 stainless steel, 26:46053 (CONF-710601-, pp 479-98)

interactions d + d, analysis of elastic scattering in the case of nonconservation of channel spin, (T), 26:46865

interactions d + p + 2p + n at 52.3 MeV, angular distributions and scattering lengths for p + p and p + n final-state interactions and model for p + p final-state interactions in, (E/T), 26:46457 (KFK-1509)

interactions d + p + 2p + n at 52.3 MeV, differential cross sections and scattering length for p + p final-state interactions for amplitudes of, (T), 26:46650 (KFK-1547)

interactions γ + d + ρ^0 + d at 6, 12, and 18 GeV, forward differential cross section for, (E), 26:46445

interactions γ + d + ρ^0 + d in Glauber theory, differential cross sections and density matrix elements for, (T), 26:46555

interactions ν + d + μ^- + p + p in impulse approximation, differential cross sections for, (T), 26:46581

interactions p + d in Glauber theory, differential cross sections, polarization, and asymmetry for, (T), 26:46655

interactions π^- + d + 2n + γ at 150 to 300 MeV, phenomenological calculation of soft-photon cross sections for, (T), 26:46531

interactions π^+ + d at high energies, research at Duke Univ., July 1971 to June 1972, (E), 26:46479(R) (ORO-3065-24)

interactions with hadrons, recoil effect in Glauber theory for quasi-elastic, 26:46653

reactions (d,n) and (d,p) with nuclei, DWBA formulation with Mitra three-body model, (T), 26:47054

reactions (d,n) with carbon-12 in steel, use in measurement of carbon distribution, 26:45466 (AE-451)

reactions (d,n) with oxygen-16 at 3 to 4 MeV, DWBA analysis of angular distributions for, (E), 26:46899

reactions (d,n) with zirconium-90 and -96 at 12 MeV, DWBA analysis of angular distributions for, (E), 26:46956

reactions (d,p) and (d,t) with thorium-232 at 13.1 MeV, Q-value and angular distributions for, (E), 26:47029

reactions (d,p) with A = 80 to 100 nuclei, analysis of threshold anomalies in, (T), 26:47058

reactions (d,p) with oxygen-16 at 12 MeV, diffraction-model analysis of differential cross sections for, (T), 26:47075

reactions (d,pf) with thorium-230 at 15 MeV, comparison of subbarrier resonances with predictions of Strutinsky theory, (E/T), 26:47033

reactions (d,pn) with hydrogen, deuterium, helium-3, and helium-4 at 27.5 MeV, analysis of p-n final-state interaction in, (E), 26:46860 (LYCEN-7186)

reactions (d,t) with lanthanum-139 at 16 MeV, DWBA analysis of differential cross section and Q value for, (E), 26:46964

scattering by nickel-58 at 100 MeV, effect of diffuse-edge potential and Coulomb interaction on elastic, (T), 26:46923

scattering by polarized hydrogen at 27.5 MeV, spin correlation coefficients from vector-polarized elastic, p. 142-5, (E), 26:46873(R) (CEA-N-1522)

DEXTRAN

radioinduced decomposition of, γ , 26:45730

DIABLO CANYON POWER REACTOR, UNIT 1

environmental report for, revised and additional data for, 26:47446 (DOCKET-50275-22)

environmental report on, appendices to, 26:47447 (DOCKET-50275-23)

DIABLO CANYON POWER REACTOR, UNIT 2

environmental report for, revised and additional data

81 SUBJECTS

DRESDEN POWER REACTOR, UNIT 2

for, 26:47446 (DOCKET-50275-22)

environmental report on, appendices to, 26:47447 (DOCKET-50275-23)

DIACETAMIDE, N-FLUOREN-2-YL-

effects of N-acetoxy and N-hydroxy derivatives of, on DNA in cultured human Xeroderma pigmentosum cells in comparison to effects of uv radiation, extent of repair synthesis following, 26:45742

DIELECTRIC FILMS

particle track counting in, using capacitive discharge device, 26:46083(P)

radioinduced photoconductivity in, x, (E/T), 26:45668 (UCRL-73974)

DIET

cesium-137 content in human, in Chicago, Jan. to June 1972, 26:45540 (HASL-257, pp III.45-52)

effects on strontium-90 content in human bones in New York City and San Francisco, 1971, 26:45538 (HASL-257, pp I.51-70)

strontium-90 content in New York City and San Francisco, Jan. to March 1972, 26:45539 (HASL-257, pp II.1-6)

DIFRACTOMETERS

description of Moessbauer, with high angular resolution, 26:45683

DIODES

radioinduced noise in Au--nSi, operating in avalanche region, (E/T), 26:45660 (FRNC-CONF-29)

DIOXANE

effects on sorption of protactinium on silica gel from aqueous solutions of hydrochloric and nitric acids, 26:45389

DIRECT ENERGY CONVERSION DEVICES

design for fusion reactors, 26:45433(P)

heat sources for radioisotope-fueled, in ENEA countries, demand to 1975 for, 26:45705

DISEASES

edema, radioinduced, in human laryngeal mucosa following therapy of laryngeal carcinomas, incidence of x, 26:45781(T) (TT-54002/4, pp 437-41)

DISPERSION RELATIONS

theory for one-dimensional, for vertex functions from local commutativity in scalar field theory, 26:46736

theory for two-dimensional, for vertex functions from local commutativity in field theory, 26:46735

DISSOLVERS

forward mixing in continuous, 26:45414(R) (ORNL-TM-3180)

DOGS

distribution and retention of metastable silver-110 in tissues of, following inhalation, 26:45915 (UR-3490-115)

hematologic changes toward aleukemic signs in peripheral blood of fetal, following serial transplantation of radioinduced canine myelomonocytic leukemia, 26:45897

lead-210 retention in beagle, following intravenous injection, 26:45916 (UR-3490-159)

radioinduced injuries to skin of beagle, following fractionated or single partial body exposure, effects of dose and fractionation interval on x, 26:45860

radioinduced neoplasms in thyroid gland of, combined iodine-131 β and γ and external x, 26:45819 (UR-3490-63)

DONALD C. COOK POWER REACTOR, UNIT 1

containment structure for, analysis of pressure transient during LOCA in, 26:47493 (DOCKET-50315-71)

environmental report for, cost-benefit analysis for alternate designs, 26:47489 (DOCKET-50315-65)

ice condenser system for, weld defects in positioning frames for, 26:47491 (DOCKET-50315-69)

pressure vessel for, deficiency in radiography techniques for inspection of, 26:47492 (DOCKET-50315-70)

safety of, revised and requested additional data on, 26:47490 (DOCKET-50315-68)

DONALD C. COOK POWER REACTOR, UNIT 2

containment structure for, analysis of pressure transient during LOCA in, 26:47493 (DOCKET-50315-71)

environmental report for, cost-benefit analysis for alternate designs, 26:47489 (DOCKET-50315-65)

ice condenser system for, weld defects in positioning frames for, 26:47491 (DOCKET-50315-69)

pressure vessel for, deficiency in radiography techniques for inspection of, 26:47492 (DOCKET-50315-70)

safety of, revised and requested additional data on, 26:47490 (DOCKET-50315-68)

DRESDEN POWER REACTOR, UNIT 1

coolant loss accidents in, hydrogen production in containment following, 26:47389 (DOCKET-50010-113)

pressure vessel for, minimum hydro test temperature and belline neutron flux calculations for, 26:47338 (DOCKET-50010-114)

DRESDEN POWER REACTOR, UNIT 2

containment structure drywell for, inspection of torus section headers of, 26:47402 (DOCKET-50237-244)

containment system for, use of railroad interlock doors of secondary, 26:47403 (DOCKET-50237-246)

control rod worth minimizer for, performance of, 26:47358 (DOCKET-50237-254)

diesel generator for, failure of, 26:47357 (DOCKET-50237-253)

diesel generators for, failure of, 26:47331 (DOCKET-50237-250)

environmental report for, cost-benefit analysis for alternate designs, 26:47404 (DOCKET-50237-249)
 inspection of, inservice, 26:47355 (DOCKET-50237-251)
 protection system for, drift in calibration of electrohydraulic control oil pressure scram switches of, 26:47352 (DOCKET-50237-245)
 protection system for, failure of electrohydraulic control oil low pressure scram switches for, 26:47354 (DOCKET-50237-248)
 radiation monitoring sampling system for, malfunction of stack, 26:47353 (DOCKET-50237-247)
 site for, revised environmental radiation monitoring program for, 26:47405 (DOCKET-50237-255)
 stack monitoring system for, malfunction of sample pump for, 26:47356 (DOCKET-50237-252)

DRESDEN POWER REACTOR, UNIT 3
 containment structure drywell for, inspection of torus section headers of, 26:47402 (DOCKET-50237-244)
 containment system for, use of railroad interlock doors of secondary, 26:47403 (DOCKET-50237-246)
 control rod worth minimizer for, performance of, 26:47358 (DOCKET-50237-254)
 environmental report for, cost-benefit analysis for alternate designs, 26:47404 (DOCKET-50237-249)
 isolation condenser for, failure of steam line high flow valve for, 26:47362 (DOCKET-50249-151)
 protection system for, drift in calibration of electrohydraulic control oil pressure scram switches of, 26:47352 (DOCKET-50237-245)
 protection system for, failure of electrohydraulic control oil low pressure scram switches for, 26:47354 (DOCKET-50237-248)
 radiation monitoring sampling system for, malfunction of stack, 26:47353 (DOCKET-50237-247)
 scram switch for, setpoint drift on pressure, 26:47332 (DOCKET-50249-148)
 site for, revised environmental radiation monitoring program for, 26:47405 (DOCKET-50237-255)

DRILL HOLES
 element content in walls of, neutron activation logging method for determination of, 26:45567

DRYING AGENTS
 processing through liquid preparation, molding, curing, machining, activation, and packaging, 26:47602 (BDX-613-392(Rev.))

DUANE ARNOLD POWER REACTOR
 environmental report for, cost-benefit analysis for alternate designs, 26:47498 (DOCKET-50331-48)
 safety of, revised and requested additional data on, 26:47500 (DOCKET-50331-51)
 safety of, revised and requested additional data on, 26:47499 (DOCKET-50331-49)

DUSTS
 analysis for plutonium by extraction and liquid scintillation counting, 26:45268
 content in atmospheric aerosols, in Poland, relations with natural radioactivity of aerosols, (E), 26:45521

DYSPROSIUM
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 magnetic susceptibility of single crystals of, anisotropy in para-, (E), 26:47253
 separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285

DYSPROSIUM BOROCARBIDES
 phase studies of, 26:45952

DYSPROSIUM ISOTOPES
 x-ray spectra from, isotope shifts in, (E), 26:46973

DYSPROSIUM ISOTOPES Dy-155
 energy levels from holmium-155 decay, spin-parity assignments for, (E), 26:46986
 energy-level transitions from holmium-155 decay, multipolarities and conversion coefficients for, (E), 26:46986

DYSPROSIUM ISOTOPES Dy-158
 energy-level transitions in deformed, SU(3) parametrization of E2, (T), 26:46975 (JINR-P4-6462)

DYSPROSIUM ISOTOPES Dy-160
 energy levels of, lifetimes of high-lying, (F), 26:47000
 energy-level transitions from terbium-160 beta decay, multipole mixing ratio for, (E), 26:46991

DYSPROSIUM ISOTOPES Dy-166
 energy-level transitions in deformed, SU(3) parametrization of E2, (T), 26:46975 (JINR-P4-6462)

E

EARTHQUAKES
 discrimination between underground nuclear explosions and, effects of tensile faults on, (T), 26:45491
 ground motion from Feb. 9, 1971 San Fernando, damage to Glendale, Calif. residential chimneys from, (E), 26:45474 (JAB-99-88)
 location from seismic data measured at single station, method for improving accuracy of, 26:45495
 seismic identification of, at East German Moxa Station, description of simple method for, 26:45496

ECHINODERMATA
 Arbacia punctulata (sea urchin), radiation effects on

anucleate and nucleate half-eggs and whole eggs of, role of cytoplasm in repair of gamma, 26:45874
 Arbacia punctulata (sea urchin), radiation effects on survival of eggs of, effects of cell phase and thiol content on x, 26:45882

ECOSYSTEMS
 book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
 cerium-144 accumulation by organisms and plants in freshwater, 26:45553
 community composition of desert plant, at Nevada Test Site, 26:45757 (TID-25954, pp 12-48)
 effects of underground nuclear explosion-generated ground motions on Hot Creek Valley, Nev, 26:45478(R) (NVO-409-1)
 praseodymium-144 accumulation by organisms and plants in littoral freshwater, 26:45553
 radiation effects on desert plant, at Nevada Test Site, fallout beta and gamma, 26:45759 (TID-25954, pp 54-7)
 radiation effects on plant, at Nevada Test Site following exposure to close-in fallout, revegetation following, 26:45762 (TID-25954, pp 75-86)
 radioactivity monitoring in Lake Michigan aquatic, near Big Rock Point power reactor during 1960 to 1969, gross beta, 26:45535
 radioecology and ecophysiology of desert plant, at Nevada Test Site, 26:45755 (TID-25954)
 radioisotope content of Lake Michigan aquatic, near Big Rock Point power reactor during 1970, measurements of cesium-134, cesium-137, cobalt-60, and zinc-65, 26:45536
 radioisotope transport through aquatic, relation of plant retention to, 26:45926
 radioisotope transport through soil-plant-small mammal, at Nevada Test Site, fallout, 26:45914 (TID-25954, pp 58-64)
 strontium-90 transport through soil to plants in desert, 26:45923 (TID-25954, pp 69-72)
 thermal effluent effects on aquatic, evaluation of, 26:45583
 thermal effluent effects on Lake Ontario aquatic, (E), 26:45578

EDWIN I. HATCH POWER REACTOR, UNIT 1
 auxiliary equipment for, requested additional data on, 26:47494 (DOCKET-50321-46)
 construction of, information on quality assurance requirements and welding procedures for, 26:47495 (DOCKET-50321-48)
 containment systems for, requested additional data on, 26:47494 (DOCKET-50321-46)
 electrical equipment for, information on storage requirements for Class 1, 26:47495 (DOCKET-50321-48)
 mechanical equipment for, information on storage requirements for Class 1, 26:47495 (DOCKET-50321-48)
 pressure vessel for, stress analysis for, 26:47494 (DOCKET-50321-46)

EDWIN I. HATCH POWER REACTOR, UNIT 2
 design change notices for, requested information on, 26:47531 (DOCKET-50366-33)
 owners of, antitrust considerations for, 26:47530 (DOCKET-50366-32)
 quality assurance requirements for, requested information on, 26:47531 (DOCKET-50366-33)
 safety evaluation of, Division of Reactor Licensing, 26:47529 (DOCKET-50366-31)
 structures for, design parameters of pressure vessel pedestal, 26:47528 (DOCKET-50366-29)

EGGS
 radiation effects on sea urchin, role of cytoplasm in repair of gamma, 26:45874
 radiation effects on survival of sea urchin, effects of cell phase and thiol content on x, 26:45882

ELECTRIC ARCS
 anode temperatures during vacuum, 26:47625 (SC-DC-722267)
 ion energy distributions during vacuum, 26:47625 (SC-DC-722267)
 use of argon, as standard black-body radiator, 26:47217

ELECTRIC CHARGES
 motion in constant and homogeneous electromagnetic fields, Lorentz transformations for electromagnetic field vectors for, (T), 26:46679

ELECTRIC CONDUCTIVITY
 effects of atmospheric, on natural radioactivity in ground-level atmosphere, 26:45524

ELECTRIC CONDUCTORS
 performance of tubular niobium/copper, for ac superconducting power transmission, 26:46844

ELECTRIC CURRENTS
 effects on erosion of thoriated tungsten cathode, protective effects of nitrogen against, 26:45968

ELECTRIC DISCHARGE
 cathode heating in asymmetrically burning Penning, 26:47136
 cyclotron harmonic waves in reflex, dispersion relations for, (E/T), 26:47184
 electron density determination by laser Thomson scattering on hydrogen, (E), 26:47125
 electron energy distribution and striations in Langmuir paradox region, (E/T), 26:47133
 model for hollow-cathode, 26:47140
 properties in plasma through magnetic neutral point, (E), 26:47142

shock wave propagation in expanding argon, (E/T),

26:47194

striations in helium, critical conditions for appearance and disappearance of, 26:46348

ELECTRIC FIELDS

bosons in external, S-matrix and quantum theory for spin-0, (T), 26:46741

electron motion in central-symmetric, energy level classification for relativistic, with anomalous magnetic moment, (T), 26:46547

fermions in external, S-matrix and quantum theory for spin-1/2, (T), 26:46741

neutron motion in central, (T), 26:46547

ELECTRIC INSULATORS

description of radiation resistant plastic, for pocket dosimeters, 26:45672(P)

joining to electrode for vacuum application, method for, 26:46130

radioinduced photoconductivity in films of, x, (E/T), 26:45668 (UCRL-73974)

ELECTRIC POWER

transmission, performance of tubular Nb/Cu conductors for ac superconducting, 26:46844

ELECTRIC PROBES

properties of Langmuir in afterglow plasmas, technique for recording, 26:47119

signal interpretation from cylindrical Langmuir, in laser-produced plasmas, 26:47126

ELECTRODES

joining to insulators for vacuum application, method for, 26:46130

plasma boundary sheath near cold, (T), 26:47130

ELECTROJETS

irregularities in equatorial, nonlinear theory for saturation of Type-1 E-layer instability, (T), 26:46277

ELECTROLYTES

properties of solid, for use in research in nuclear history, 26:46013

ELECTROMAGNETIC FIELDS

charged-particle interactions with, Poincare group algebra for, 26:46678

charged-particle motion in gravitational and, model for spinning, (T), 26:46767

charged-particle motion in, regularization in vacuum polarization calculations of vector currents for, 26:46742

Einstein--Maxwell equations for, null bundle transformations for solutions to, (T), 26:46762

Einstein--Maxwell equations for, null bundle transformations for solutions to, (T), 26:46763

electric charge motion in constant and homogeneous, Lorentz transformations for electromagnetic field vectors for, (T), 26:46679

field equations for, cylindrical wave solutions of Bonner and Schroedinger unified, (T), 26:46695

field equations in formulation of Einstein--Kaufman nonsymmetric unified field theory for, static, spherically symmetric solutions of, (T), 26:46694

field theory for, formulation of Einstein--Kaufman nonsymmetric unified, (T), 26:46694

Maxwell equations for singular source-free, for nonspecial recurrent space--time for states of total pure radiation, 26:46771

Maxwell equations for, null bundle transformations for Kerr solution for gravitation field of homogeneous rotating ring for, (T), 26:46762

particle motion in, canonical dynamics of spinning, 26:46753

ELECTROMAGNETIC WAVES

coupling with electron space-charge waves outside plasmopause, Cherenkov instability and vlf emissions from, (E/T), 26:46318

emission between 3.9 and 10.4 kHz in high-altitude radiation-belt slot, whistler-mode vlf, (E), 26:46312

emission from electron in magnetic field at $n = 1$ level, (T), 26:46551

emission of intense bands of elf noise in association with low-energy auroral electron precipitation at low altitudes, (E), 26:46308

focusing in isotropic plasma, nonlinear self-, (T), 26:47185

production from neutron motion through medium in magnetic field, coherent generation mechanism for, (T), 26:46541

propagation in dissipative plasma, collision effects on nonlinearities in, (T), 26:47183

ELECTROMAGNETISM

vector potentials in, measurement of, 26:46542

ELECTRON BEAMS

dosimetry at 8 to 30 MeV, calibration and performance of thimble ionization chambers for, 26:45623

focusing of, electrostatic lens for line, 26:46147

interactions with plasma, wave excitation by modulated, (E), 26:47201

interactions with plasma, wave excitation from cold, (T), 26:47151

monitoring at 1 to 50 MeV, instruments and methods for, 26:45618 (ICRU-21)

polarization of, by scattering by atomic beams or gases, (E), 26:46338

simulation of intense relativistic, in axisymmetric magnetic mirror fields, (T), 26:47147

space-charge waves from, Cherenkov instability in coupling with electromagnetic waves, (E/T), 26:46318

ELECTRON CAPTURE

log-ft tables for, 26:46970

spectra as function of atomic number from, Auger-electron and fluorescence-photon, (T), 26:47209

ELECTRON MICROSCOPES

development for localization of trace elements in biological materials, 26:45893 (ORO-2690-12(Pt.B), Paper 13)

use of high-voltage, in studies of electron-induced defect clusters at high temperatures, 26:46057 (CONF-710601-, pp 550-62)

use of high-voltage, in studies of electron-induced voids in metal films, 26:46056 (CONF-710601-, pp 533-49)

ELECTRON MICROSCOPY

backscattering from perfect and imperfect crystals in scanning, dynamic theory of, 26:47267

ELECTRON PROBE MICROANALYSIS

accuracy of quantitative, using energy dispersive spectrometers, 26:45237

equipment, methods, and errors in, review on, 26:45246

ELECTRON-RING ACCELERATORS

electron and electron--ion rings in, calculations of axial stability of, 26:46142

electron-ring focusing by image forces in cylindrical screen in, 26:46136 (JINR-P9-6421)

electron-ring stability at center shift in screening tube, 26:46137 (JINR-P9-6514)

ELECTRON SOURCES

description of secondary emission multipactoring 20-mA, for linacs, 26:46134

ELECTRONIC FILTERS

design optimization from amplitude spectrometry systems, 26:45637 (JINR-P13-6428)

ELECTRONS

cesium fluoride scattering by 0.6- to 6.5-eV, molecular-beam recoil measurements of absolute differential and total cross sections for, 26:46406

detection and discrimination at 2.8 and 3.7 GeV, description of resonance radiation detector for, 26:45648

dielectric models for solvated, continuous, 26:45344

diffraction of high energy, from imperfect crystals, modified Bloch wave expansion for formulation of theory for, (T), 26:46510 (CALT-767-P-3-22)

diffusion of plasmaspheric radiation-belt, association with plasmaspheric whistler-mode waves of pitch-angle, (E/T), 26:46311

dosimetry at 1 to 50 MeV, recommendations for dosimeters and methods for, 26:45618 (ICRU-21)

dosimetry of pulsed, at 10 to 30 MeV, performance of ionization chambers for, 26:45627

effects of pulsed, on anthracene, naphthalene, diphenyl-2,5-oxazole, and pyrene in benzene and cyclohexane solutions, 26:45328

effects of pulsed, on fluorescence of anthracene, naphthalene, and pyrene in cyclohexane, 26:45331

effects of pulsed, on ice and ice-d₂, optical absorption spectra of transient electrons from, 26:45309(R) (AE-453)

effects of pulsed, on liquid hydrocarbons, theoretical study of electron scavenging process in, 26:45333

effects of pulsed, on naphthalene and pyrene in benzene or cyclohexane solutions, effects of quenchers on yields of singlet state from, 26:45329

effects of pulsed, on naphthalene in cyclohexane solutions, effects of scavengers on yield of excited states from, 26:45330

effects on aluminum single crystals, study of I/sub B/ stage in annealing of, 26:47229

effects on copper at 2.6 MeV and 60K, annealing of lattice parameter and electric resistivity changes from, 26:47231

effects on crosslinking and polymerization of polyacrylates dissolved in vinyl monomers, 26:45339(P)

effects on crystal lattice defect formation in alloys and metals, review on, 26:46076

effects on desorption of carbon monoxide from niobium (111), 26:47242

effects on embrittlement of polyethylene and polyvinyl chloride at 1.3 MeV, 26:46090

effects on hypoxic mice following whole-body exposure, sensitizing effects of TAN on lethal, 26:45869

effects on Kapron fibers, x-ray scattering studies on, 26:45338

effects on mites infesting ham, lethal, 26:45886

effects on potassium cyanoiridate in potassium chloride lattice, ESR studies on, 26:45317

effects on secondary emission and properties of solids, Russian book on, 26:47245

effects on single crystals of acetylglycine, ESR and INDO MO studies of radicals from, 26:45349

effects on single crystals of potassium chloropalladate and potassium chloroplatinate, ESR studies on, 26:45341

effects on survival of Chlamydomonas reinhardtii, dose and dose rate dependence of, 26:45754

effects on volt--amp characteristics of n-type germanium at 8.5 MeV, 26:47230

- electromagnetic radiation emission from, in magnetic field at $n = 1$ level, (T), 26:46551
- energy of secondary, from gamma and x-ray attenuation in matter, method for determining mean, 26:47213
- fluxes in nighttime aurorae, Fermi acceleration mechanism for keV, (E/T), 26:46267
- g-factor of, review and analysis of data on ratio to muon for, (E), 26:46439
- interactions $e + e \rightarrow 2e + 2\nu$, cross sections for, (T), 26:46519 (JINR-P2-6333)
- interactions $e + e$, invariant method for differential cross sections, angular distributions, and two-photon mechanism for meson production in, (T), 26:46518 (ITEF-848)
- interactions $e + e$, test of Han-Nambu and Gell-Mann--Zweig triplet quark models for 2γ production in, (T), 26:46516 (DESY-72/21)
- interactions $e + N + e + \text{hadron} + \text{anything}$ via virtual photon $N + \text{hadron} + \text{anything}$ interactions at high energies and momentum transfer, model-independent derivation of generalized scaling laws and produced-particle multiplicity for, (T), 26:46521 (LBL-773)
- interactions $e + N + \pi + N$ in first resonance region, model with axial vector form factor for pion form factor from, (T), 26:46513 (CONF-710816-17)
- interactions $e + N + \pi + N$ near threshold, current algebra and PCAC hypothesis for amplitudes and cross section sensitivity to nucleon axial-vector form factor for, (T), 26:46549
- interactions $e + N + \pi^0 + N$ in first resonance region, multipole amplitude analysis of angular distributions and transition form factor for $\Delta(1236)$ in, (T), 26:46512 (CONF-710816-16)
- interactions $e + N + \pi^+(\pi^-) + e + \text{anything}$, PCAC hypothesis and soft pion theorems for cross sections for, (T), 26:46521 (LBL-773)
- interactions $e + N$ at 20 GeV, evidence for partons from inelastic, (E), 26:46466
- interactions $e + N$, formulation of field theory of quarks for structure functions of nucleon in inelastic, (T), 26:46524 (NYO-4076-18)
- interactions $e + N$, predictions of nonrelativistic harmonic-oscillator quark model for cross sections for resonance production in, 26:46544
- interactions $e + p \rightarrow e + p + \pi^+ + \pi^-$ in rho resonance region, extension of diffraction--dissociation model for polarization, momentum transfer, and virtual photon and dipion mass dependence of cross sections for, (T), 26:46517 (DESY-72/23)
- interactions $e + p$ at 5 + 200 GeV in colliding beams in ISABELLE, feasibility of photoproduction experiments from total cross sections for, (E), 26:46426 (BNL-16968)
- interactions $e + p$ at large angles, parity violation in elastic, (E/T), 26:46562
- interactions $e + p$ in $\Delta(1236)$ region at 20 to 810, magnetic dipole transition form factor of $\gamma p \Delta$ vertex determination from longitudinal and transverse cross sections for inelastic, (E), 26:46431 (NP-19256)
- interactions $e + p$ in parton model, scaling behavior of structure functions for deep inelastic, (T), 26:46566
- interactions $e + p$, invariant method for differential cross sections, angular distributions, and two-photon mechanism for meson production in, (T), 26:46518 (ITEF-848)
- interactions $e + p$, polarization contribution to ground-state hyperfine splitting of hydrogen from structure functions from resonance model of inelastic, (T), 26:46523 (NYO-4076-17)
- interactions $e^+ + e^- \rightarrow 2\pi^+ + 2\pi^-$ at 990 MeV in colliding beams, upper limit for cross sections for, (E), 26:46429 (CONF-710816-15)
- interactions $e^+ + e^- \rightarrow e^+ + e^-$, acoplanarity distributions and radiative corrections in Bhabha, (T), 26:46543
- interactions $e^+ + e^- \rightarrow \gamma + \text{hadrons}$, canonical anomaly for high-energy behavior of, (T), 26:46556
- interactions $e^+ + e^- \rightarrow \text{hadron pairs}$ at 1400 to 2400 MeV, energy dependence of, (E/T), 26:46552
- interactions $e^+ + e^- \rightarrow \text{hadrons} + \text{anything}$, review of dual quark-parton model for semi-inclusive, (T), 26:46515 (DAMTP-71/24)
- interactions $e^+ + e^- \rightarrow \mu^+ + \mu^- + \text{anything}$, review of dual quark-parton model for semi-inclusive, (T), 26:46515 (DAMTP-71/24)
- interactions $e^+ + e^- \rightarrow \phi + K_1^0 + K_2^0, K_1^0 + \pi^+ + \pi^-$, mass of phi from, (E), 26:46443
- interactions $e^+ + e^- \rightarrow \pi + \text{hadrons}$, behavior of structure functions and cross sections for, (T), 26:46540
- interactions $e^+ + e^- \rightarrow \pi^+ + \pi^- + \pi^0 (2\pi^0)$ at 990 MeV in colliding beams, total cross sections for, (E), 26:46429 (CONF-710816-15)
- interactions $e^+ + e^- \rightarrow \pi^+ + \pi^- + \pi^0$ in ω region in storage ring, cross sections for, (E), 26:46427 (CONF-710816-13)
- interactions $e^+ + e^- \rightarrow \pi^+ + \pi^-$ in colliding beams, $\pi^+\pi^-$ excitation curve in rho region for amplitude and phase of ω --rho interference and rho production cross sections in, (E), 26:46428 (CONF-710816-14)
- interactions $e^+ + e^-$, tests of Han-Nambu and Gell-Mann--Zweig triplet quark models for 2γ production in, (T), 26:46516 (DESY-72/21)
- interactions $e^- + p \rightarrow e^- + \mu^+ + \mu^- + p$, scaling limit for virtual Compton scattering off forward direction from, (T), 26:46530
- interactions $\gamma + e$, quantum-mechanical calculation of, (E), 26:46532
- interactions $\nu + e \rightarrow \text{anti } \nu + e^-(\mu^-)$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions and energy dependence of ν energy loss and scattering angle in, (T), 26:46579
- interactions $\nu + e \rightarrow \nu + e$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions and energy dependence of ν energy loss and scattering angle for, (T), 26:46579
- interactions $\nu + e \rightarrow \nu + \mu^-$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions and energy dependence of ν energy loss and scattering angle for, (T), 26:46579
- interactions of, test of quantum electrodynamics from, 26:46710
- interactions with niobium and silicon crystals at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550
- magnetic moment of, quantum electrodynamic contribution to anomalous, 26:46710
- magnetic moment of, review and analysis of data on relation for, (E), 26:46439
- mass of, cosmological significance of difference of muon mass and, (T), 26:46259
- mass of, review and analysis of data on ratio to proton and muon for, (E), 26:46439
- motion in medium in magnetic field, spin effects on synchrotron radiation intensity and lifetime of ground and first excited states for ultra-luminous, (T), 26:46548
- motion of bundle of relativistic, in ring accumulator, particle collective interaction effects on longitudinal dimensions and phase for, (T), 26:46546
- motion of relativistic, with anomalous magnetic moment, in central-symmetric electric field, energy level classification for, (T), 26:46547
- observations of high-energy, from HEOS-1 launched Dec. 5, 1968, (E), 26:46334
- polarization in magnetic field, external magnetic field wave effects on, (T), 26:46551
- radial-diffusion coefficients in inner radiation belts, natural and artificial, (E/T), 26:46310
- reactions ($e,e'p$) with carbon-12 and calcium-40 at 501 MeV, cross sections for, p. 194-6, (E), 26:46873(R) (CEA-N-1522)
- reactions (e,p) with carbon-12 at 130 MeV, analysis of differential cross sections and spectra from, (E/T), 26:46888
- reactions of hydrated, with aromatic carbonyl compounds, 26:45354
- reactions of solvated, in γ radiolysis of liquid ethanol, effects of pressure on, 26:45351
- reactions with argon up to 300 eV, ionization efficiency curves for Ar^+ and Ar^{2+} in, (E), 26:46374
- reactions with argon, helium, and neon in eV range, integral cross sections for production of secondary electrons of 0 to 1 eV by ionization in, (E), 26:46390
- reactions with atoms and molecules, third-derivative method for determination of ionization and appearance potentials for, (E/T), 26:46385
- reactions with atoms, bound states in quantum electrodynamics for, (T), 26:46721
- reactions with carbon dioxide at 25 to 600 eV, cross sections for production of $\text{CO}^+, \text{O}^+, \text{C}^+$, and O_2^+ by dissociation and ionization in, (E), 26:46384
- reactions with gold at 35 keV, calculation of energy loss spectra from 1 to 12 eV from, (T), 26:47212
- reactions with helium at 40 to 70 eV, resonance in differential cross sections for 2^3S excitation in, (E), 26:46405
- reactions with helium at 500 eV and 1 keV, angular and energy distributions of secondary electrons from direct ionization and autoionization in, (E), 26:46367
- reactions with helium at zero angle, second Born differential cross sections in high-energy limit for excitation of 2^3S and 2^3P states in, 26:46373
- reactions with helium, structure in electron energy distribution, (E), 26:46407
- reactions with krypton at 1 keV, angular and energy distributions of secondary electrons from direct ionization and autoionization in, (E), 26:46367
- reactions with nitrogen molecules at 0 to 2000 eV, Born-approximation calculations of cross sections for excitation in, effects of choice of wave functions on, 26:46391
- reactions with nitrous oxide, cross sections for O^- production by dissociative electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- reactions with oxygen molecules, cross sections for electron attachment and dissociative electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- reactions with polyatomic molecules, cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- reactions with rare gas clusters, multiple ionization in, (E), 26:46386
- reactions with tantalum at 12 to 30 MeV, target thickness effects on positron spectra from, (E), 26:46149

collective modes of excitation in, 26:46389
 scattering by atoms, optimization of basis set in calculation of amplitudes for, 26:46389
 scattering by helium, calculation using method of effective potentials and variational formulations, 26:46400
 scattering by hydrogen molecules, close-coupling calculations of differential and total cross sections for, using square-well potentials, 26:46399
 scattering by lead-208 at 400 and 600 MeV, phase-shift analysis of form factor for elastic, p. 192-4, (E), 26:46873(R) (CEA-N-1522)
 scattering by molecules at high energies, Glauber-theory calculations of amplitudes and differential cross sections for multiple elastic, 26:46381
 scattering by neon, bound-state calculation of resonance due to Ne⁻ ion in, 26:46383
 scattering by nitrogen molecules, crossed-beam measurements of angular distribution and total elastic-scattering and momentum-transfer cross sections for, 26:46392
 scattering by nuclei with bremsstrahlung emission, effect of nuclear multipole moments on polarized, (T), 26:47052
 scattering by rare gases, scattering lengths for, (T), 26:46401
 scattering by samarium-152 at 76 MeV, electron spectra from inelastic, (E), 26:46993
 scattering by zirconium-90 at 302 MeV, differential cross sections for elastic, p. 186-8, (E), 26:46873(R) (CEA-N-1522)
 scattering by zirconium-90, molybdenum-92, and tin-116, -120, and -124 at 209 MeV, differential cross sections for inelastic, p. 189-91, (E), 26:46873(R) (CEA-N-1522)
 scattering of, bound states in quantum electrodynamics for, (T), 26:46721
 spectra of 20-MeV to 15-GeV cosmic-ray, modulation near solar maximum, (E), 26:46325
 spectra of trapped, produced by γ radiolysis of ethanol glass, optical absorption and ESR, 26:45319
 spike detection at high latitudes in magnetosphere, phenomena of equal to or greater than 425 keV, (E), 26:46307
 theory for, review of, (T), 26:46508
 transport at 1 to 8 MeV, discrete-ordinates method for, 26:47210
 wave equations for, fusion method for Dirac--Hestenes and Maxwell, (T), 26:46568
 yields of trapped, in radiolysis of doped ice, effects of fluoride on, 26:45345

ELECTROSTATIC ACCELERATORS
 beam injection and modulation system for pulsed, control system for, 26:46135

ELECTROSTATIC ANALYZERS
 electrical insulation for plane-parallel electrode type, review on, 26:45587

ELECTROSTATIC GENERATORS
 accelerator tubes for, method for joining electrodes and insulators in fabrication of, 26:46130

ELECTROSTATIC LENSES
 description for transforming circular electron beam into line-focus beam, 26:46147

ELEMENTARY PARTICLE MODELS
 bootstrap, matrix elements of internal symmetry currents, energy--momentum density tensor, and generalized unitarity in conformal invariant quantum field theory in, 26:46697 (DESY-72/20)
 diffraction dissociation, extension for polarization, momentum transfer, and virtual photon and dipion mass dependence of cross sections for $e + p \rightarrow e + p + \pi^+ + \pi^-$ interactions at high energies, 26:46517 (DESY-72/23)
 diffractive-excitation, comparison with multiperipheral model for two-particle distribution functions for strong interactions at high energies for, 26:46587 (RLO-2230-T-4-1)
 dual quark-parton, review for $\nu + N$, $e^+ + e^-$, and strong interactions in, 26:46515 (DAMP-71/24)
 dual resonance, finite transformations of Virasoro group in, 26:46686
 dual resonance, SL(2,R) group for generalized, 26:46593
 dual resonance, statistical mechanics calculation of resonance spin distribution in, 26:46609 (ITF-72-13-P)
 dual resonance, sum rules for Compton scattering of photon-like particles in, 26:46601
 dual, factorization in multiperipheral configuration in, 26:46726
 Feynman gas, correlation functions and generalization in formulation from jet approach to deep inelastic hadron interactions, 26:46586 (JINR-P2-6501)
 fireball, Reggeon contributions to inelastic interactions in two-, 26:46663
 harmonic-oscillator quark, for resonance electroproduction, nonrelativistic, 26:46544
 multiperipheral, comparison with diffractive-excitation model for two-particle distribution functions for strong interactions at high energies for, 26:46587 (RLO-2230-T-4-1)
 multiperipheral, self-consistent solution for diffraction scattering in, 26:46602
 paraquark, $\Delta I = 1/2$ rule for nonleptonic weak Hamiltonian

in, 26:46580
 parton, duality in strong interactions, 26:46599
 parton, formulation for hadrons, SU(1,1) algebra representations for wave packet realization of lightlike particle states for, 26:46681 (NYO-4076-19)
 parton, SU(1,1) algebra classification for wave packet realization of lightlike particle states for formulation for hadrons, 26:46589
 parton, weak and electromagnetic interactions in integral-charge, 26:46574
 quark, comparison of excited, interacting, and additive, for constraints on decay angular distributions for resonances produced in photon interactions with baryons, 26:46569
 quark, comparison of excited, interacting, and additive, for constraints on resonance angular decay distributions in photon interactions with spin- $1/2$ baryons, 26:46526 (TPJU-14/71)
 quark, current densities in formulation of field theoretic lightlike, 26:46524 (NYO-4076-18)
 quark, tests of Han--Nambu and Gell-Mann--Zweig triplet, using meson radiative decays and e^+ and e^- interactions, 26:46516 (DESY-72/21)
 quartet, for hadrons in chiral U(4) x U(4), 26:46687
 Regge, for cross sections of $\gamma + N \rightarrow K^+ + N$ interactions at 5 to 15 GeV/c, 26:46534
 three-triplet, $\Delta I = 1/2$ rule for nonleptonic weak Hamiltonian in, 26:46580
 urbaryon rearrangement, differential cross sections for hadron inclusive interactions in, 26:46606

ELEMENTARY PARTICLES
 decay of unstable, spin spectrum from, 26:46498
 electromagnetic interactions at high energies in condensed media, book on, (T), 26:46538
 interactions at high energies, relativistic eikonal approximation for production, (T), 26:46791
 interactions of relativistic, phase shift in first and second orders with respect to quasipotential for, (T), 26:46505
 interactions of spinning, relativistic generalization of Faxen--Holtzmark method for amplitudes of, (T), 26:46793
 interactions of, algebraic derivation of generalized Talmi expansion for two-body, 26:46503
 interactions of, analytical properties of amplitudes and differential cross sections for multiparticle production over two angular variables in, (T), 26:46497 (IFVE-STF-71-36)
 interactions of, breakdown of simple cancellation in eikonal multiple scattering series for three-body, (T), 26:46773 (CTH-ITP-71-17)
 interactions of, energy--momentum conservation in inclusive, (T), 26:46504
 interactions of, explicit Poincare group representations labelled by additional discrete quantum numbers for, (T), 26:46665 (CTH-ITP-71-14)
 interactions of, kinematically corrected correlation functions in phase-space model for inclusive, (T), 26:46500
 interactions of, properties of correlation functions for cross sections of inclusive, (T), 26:46502
 interactions of, Regge pole model, duality, and exotic resonances in inclusive, 26:46605
 interactions of, removal of ambiguities in phase-shift analysis of, 26:46640
 interactions of, self-consistent solution in multiperipheral model for diffractive, (T), 26:46602
 interactions with fields, Bogoliubov transformation in strong coupling theory for, (T), 26:46691
 lectures on, (T), 26:46509
 operators for, irreducible representations of Sp(2n,R) algebra for para-Bose, 26:46507
 production at high energies, relativistic eikonal approximation for, (T), 26:46791
 production of, analytical properties of amplitudes and differential cross sections over two angular variables in multiple, (T), 26:46497 (IFVE-STF-71-36)
 production processes for, eikonal formalism for, 26:46778
 quantum theory of, review of, 26:46508
 reactions with carbon-type light nuclei at 300 to above 1000 MeV, intranuclear cascade model for density effects on cascade particle properties and density and cluster-structure effects on isotope yield in, (T), 26:46664
 scattering by nuclei at high energies, formalism for, 26:47100
 spin spectrum of unstable, 26:46498
 strong interactions at high energies and large angles, review of elastic, (E/T), 26:46591
 strong interactions at high energies, comparison of two-particle distribution functions in diffractive-excitation and multiperipheral models for, (T), 26:46587 (RLO-2230-T-4-1)
 strong interactions at high energies, conference on, (E/T), 26:46590
 strong interactions at high energies, exchange degeneracy and dip mechanism in differential cross sections of, (E/T), 26:46592
 strong interactions in parton model, duality in, (T),

- 26:46599
strong interactions of, review of dual quark-parton model for, (T), 26:46515 (DAMTP-71/24)
wave equations for arbitrary-spin, Poincare and CPT invariance in determination of, 26:46714
wave equations for massive, multipole radiation solutions, 26:46709
wave equations for nonzero-mass, arbitrary-spin, three types of relativistic, 26:46730
wave functions of free arbitrary-spin, Poincare transformations for properties of one-particle, (T), 26:46667 (ITF-71-117-P)
weak interactions of, theory of, 26:46584
weak interactions of, upper bound for tensor component of second-class currents and mass limitation for axial-vector form factor for, (T), 26:46573
- ELEMENTARY PARTICLES, POSTULATED**
theory for dyons, dual invariant Lagrangian for dual invariance for, (T), 26:46539
- ELEMENTS**
content in geophysical formations, neutron activation method for determining, 26:45561(P)
- ELK RIVER POWER REACTOR**
operating authorization for, transfer request for, 26:47336 (DOCKET-1151-57)
- EMBRYOS**
nutrition of mammalian, 26:47622(R) (ORO-4245-4)
radiation effects on phenol red uptake by mesonephros of chick, in vitro, temperature dependence of x, 26:45832
- ENERGY**
transmission to user sites as hydrogen, evaluation of nuclear systems for, 26:45429
- ENTRAINMENT SEPARATORS**
development, performance, and testing of commercial 1- to 10 μ air--water, survey on, 26:45440 (MSAR-71-45)
- ENVIRONMENT**
book: Radiation Protection Measurement, 26:45886
contamination by artificial radioactivity, assessment and control of, 26:45554
contamination from peaceful uses of nuclear energy, evaluation of, 26:45555
hazards from fossil- and nuclear-fueled power plants, methods for comparing, 26:45510 (AAEC/TM-627)
hazards to world, from existence of nuclear weapons, 26:45492
impact report on, leasing of AEC-controlled uranium-bearing lands in Colorado, New Mexico, and Utah, 26:45572 (WASH-1523)
impact report, amendment to Brunswick Power Reactor Units 1 and 2, 26:47497 (DOCKET-50324-40)
impact report, amendment to Salem Power Reactor, Units 1 and 2, 26:47445 (DOCKET-50272-33)
impact report, Arkansas Nuclear One Power Reactor Unit 2, 26:47532 (DOCKET-50368-31)
impact report, Barnwell Nuclear Fuel Plant uranium hexafluoride facility, 26:45308 (DOCKET-701327-1)
impact report, Kewaunee Power Reactor, 26:47471 (DOCKET-50305-45)
impact report, LaSalle County Power Reactor Units 1 and 2, 26:47535 (DOCKET-50373-24)
impact report, Maine Yankee Power Reactor, 26:47473 (DOCKET-50309-61)
impact report, McGuire Power Reactor Units 1 and 2, 26:47534 (DOCKET-50369-28)
impact report, Nine Mile Point Power Reactor Unit 1, 26:47401 (DOCKET-50220-98)
impact report, North Anna Power Reactor, Units 1 and 2, 26:47511 (DOCKET-50338-48)
impact report, Palisades Power Reactor, 26:47424 (DOCKET-50255-96)
impact report, supplement to Arkansas Nuclear One Power Reactor, 26:47487 (DOCKET-50313-45)
impact report, supplement to Arkansas Nuclear One Power Reactor, 26:47486 (DOCKET-50313-44)
impact report, supplement to Connecticut Yankee Power Reactor, 26:47397 (DOCKET-50213-152)
impact report, supplement to Cooper Power Reactor, 26:47465 (DOCKET-50298-43)
impact report, supplement to Cooper Power Reactor, 26:47468 (DOCKET-50298-48)
impact report, supplement to Davis--Besse Power Reactor, 26:47518 (DOCKET-50346-29)
impact report, supplement to Diablo Canyon Power Reactor Units 1 and 2, 26:47446 (DOCKET-50275-22)
impact report, supplement to Diablo Canyon Power Reactor, Units 1 and 2, 26:47447 (DOCKET-50275-23)
impact report, supplement to Donald C. Cook Power Reactor Units 1 and 2, 26:47489 (DOCKET-50315-65)
impact report, supplement to Dresden Power Reactor Units 2 and 3, 26:47404 (DOCKET-50237-249)
impact report, supplement to Duane Arnold Power Reactor, 26:47498 (DOCKET-50331-48)
impact report, supplement to James A. Fitzpatrick Power Reactor, 26:47502 (DOCKET-50333-51)
impact report, supplement to Linerick Power Reactor, Units 1 and 2, 26:47520 (DOCKET-50352-27)
impact report, supplement to North Anna Power Reactor Units 1, 2, 3, and 4, 26:47510 (DOCKET-50338-47)
impact report, supplement to Peach Bottom Power Reactor Units 2 and 3, 26:47448 (DOCKET-50277-52)
- impact report, supplement to Rancho Seco Power Reactor, 26:47479 (DOCKET-50312-41)
impact report, supplement to Shearon Harris Power Reactor Units 1, 2, 3, and 4, 26:47564 (DOCKET-50400-20)
impact report, supplement to Trojan Power Reactor Unit 1, 26:47516 (DOCKET-50344-30)
impact report, supplement to Waterford Power Reactor Unit 3, 26:47539 (DOCKET-50382-27)
impact report, supplement to Waterford Power Reactor Unit 3, 26:47543 (DOCKET-50382-31)
impact report, Susquehanna Power Reactor Units 1 and 2, 26:47545 (DOCKET-50387-19)
impact report, Susquehanna Power Reactor Units 1 and 2, 26:47546 (DOCKET-50387-20)
impact report, Turkey Point Power Reactor Units 3 and 4, 26:47413 (DOCKET-50250-84)
impact report, Vermont Yankee Power Reactor, 26:47437 (DOCKET-50271-92)
impact report, Vermont Yankee Power Reactor, 26:47435 (DOCKET-50271-88)
radiation monitoring near nuclear facilities, development of programs for, 26:45556
radiation monitoring of Pearl Harbor marine, during 1966 and 1968, 26:45534
radiation protection programs for, at Marcoule nuclear center, 26:45804
radioactivity in human, hazards to public health from, 26:45890 (NYO-3970-29)
radioactivity in human, hazards to public health from, 26:45904 (COO-3013-1)
radioactivity of human, relation to public health, 26:45911 (COO-3013-25)
tritium burden in, following release and migration, 26:45509 (KFK-1560)
- ENZYMES**
ion reactions with, binary encounter method for calculation of cross sections for inactivation in, 26:46365(R) (ORO-2911-14)
radiation effects on activity and regulatory functions of, 26:45726
radiation effects on activity of, in neuroglia of developing rabbit brain, x, 26:45836
radiation effects on activity of, in spinal neurons of mice, 26:45868
- EPITHELIUM**
radioinduced injuries in midgut, of adult and larval yellow mealworms, dose and time dependence of γ , 26:45875
- EPOXY RESINS**
radioinduced photoconductivity in films of, x, (E/T), 26:45668 (UCRL-73974)
- ERBIUM**
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
magnetic susceptibility of single crystals of, anisotropy in para-, (E), 26:47253
- ERBIUM HYDRIDES**
elastic properties of $\text{ErH}_{1.81}$, from acoustic velocity measurements, 26:45950
- ERBIUM ISOTOPIES**
x-ray spectra from, isotope shifts in, (E), 26:46973
- ERBIUM ISOTOPIES Er-166**
energy-level transitions in deformed, $\text{SU}(3)$ parametrization of E_2 , (T), 26:46975 (JINR-P4-6462)
photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
- ERBIUM ISOTOPIES Er-168**
energy-level transitions in deformed, $\text{SU}(3)$ parametrization of E_2 , (T), 26:46975 (JINR-P4-6462)
- ERBIUM ISOTOPIES Er-173**
beta decay of, gamma spectra, half-life, and Q value for, (E), 26:46997
- ERBIUM PEROXIDES**
preparation and properties of, 26:46085
- ERYTHROCYTES**
cation flux changes in human, suspended in anionic media, 26:47638 (UR-3055-1)
radioinduced inhibition of glucose-6-phosphatase activity in, of rabbits and rats following whole-body exposure, effects of cysteine and reduced glutathione on extent of recovery from x, 26:45848
- ERYTHROPOIETIN**
radiation effects on metabolism of, in rats following whole-body exposure, additive effects of hypoxia on, 26:45854
- ESTUARIES**
salinity intrusions in, mathematical model for predicting unsteady, 26:47620 (MIT-144)
- ETHANE**
proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s hydrogen atoms by charge exchange and excitation in, (E), 26:46378
- ETHANE, BROMO-**
Br--bromoethane, bromine-80m isomeric transition in mixtures of, effects of initial form on yield of labeled compounds and chemical damage in, 26:45360
- ETHANE, 1,1-DICHLORO-**
radioinduced reaction with cyclohexanone, yields from γ , 26:45335
- ETHANE, HEXAFLUORO-**
radiolysis of, research progress on production of large-molecular-weight fluorocarbons by γ , 26:45316(R) (ORO-

ETHANOL

- effects on sorption of protactinium on silica gel from aqueous solutions of hydrochloric and nitric acids, 26:45389
- radiolysis of glasses of, optical absorption and ESR spectral studies of trapped electrons from γ , 26:45319
- radiolysis of neutral and acidic liquid, effects of pressure on rates of reaction of solvated electrons from γ , 26:45351

ETHERS

- photolysis of solid, ESR spectral studies of far-uv, 26:45323

ETHYLENE

- ethylene-- NH_3 , radiolysis of mixtures of, yields of hydrogen from γ , 26:45332

ETHYLENE, CHLORO-

- analysis for arsenic, bromine, and phosphorus impurities by neutron activation, 26:45204

ETHYLENE, CHLORO-, POLYMERS

- radioinduced crosslinking and curing of coatings of, 26:45352(P)
- radioinduced embrittlement for mechanical waste consolidation, 1.3-MeV electron, 26:46090

ETHYLENE, 1,2-DICHLORO-

- radioinduced reaction with cyclohexanone, yields from γ , 26:45335

ETHYLENE, 1,2-DIFLUORO-

- radioinduced crosslinking and curing of coatings of, 26:45352(P)

ETHYLENE POLYMERS

- radiation effects on sheets of, comparison of partial discharges with ion, 26:46081
- radioinduced electric conductivity of, effects of dose rate of pulsed γ - and x -, 26:47225 (AWRE-0-31/72)
- radioinduced embrittlement for mechanical waste consolidation, 1.3-MeV electron, 26:46090
- radioinduced photoconductivity in films of, x , (E/T), 26:45668 (UCRL-73974)

ETHYLENE, TETRACHLORO-

- radioinduced reaction with cyclohexanone, yields from γ , 26:45335

ETHYLENE, TETRAFLUORO-

- radioinduced photoconductivity in films of, x , (E/T), 26:45668 (UCRL-73974)

ETHYLENE, TETRAFLUORO-, POLYMERS

- radioinduced electric conductivity of, effects of dose rate of pulsed γ - and x -, 26:47225 (AWRE-0-31/72)

ETHYLENE, TRICHLORO-

- radioinduced reaction with cyclohexanone, yields from γ , 26:45335

ETHYLENIMINE

- effects on barley seeds, protective effects of cysteine against mutagenic and toxic, 26:45772

EUROPE

- uranium enrichment in, plans for development of international civilian gaseous diffusion plant for, 26:45693

EUROPIUM

- determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
- separation from gadolinium, neodymium, praseodymium, and samarium by anion exchange using acetic acid--methanol--nitric acid mixtures, 26:45280
- separation from nitric acid solutions by solvent extraction using Versatic acid 911, 26:45277
- separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
- uptake of radioactive, by larval and hatchling amphibians and reptiles, 26:45815(R) (COO-3486-1)

EUROPIUM ISOTOPES Eu-147

- energy levels from gadolinium-147 decay, spin--parity assignments and transition multipolarities for, (E), 26:46954

EUROPIUM ISOTOPES Eu-152

- content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)

EUROPIUM ISOTOPES Eu-153

- energy level at 103.2 keV, g -factor for, (E), 26:46981

EUROPIUM ISOTOPES Eu-154

- content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)

EUROPIUM ISOTOPES Eu-155

- energy level at 105 keV, g -factor for, (E), 26:46981

EUROPIUM ISOTOPES Eu-156

- beta decay of, γ -- γ coincidence spectra from, (E), 26:46996

EUROPIUM ISOTOPES Eu-158

- beta decay of, γ -- γ coincidence spectra from, (E), 26:46996

EUROPIUM URANIUM SELENIDES

- preparation and structure EuU_2Se_5 , 26:45378

EUROPIUM URANIUM SULFIDES

- preparation and structure of EuU_2S_5 , 26:45378

EVAPORATORS

- criticality in intercycle, at West Valley, 26:45401 (DOCKET-50201-128)
- heat exchangers for, design criteria for, 26:45436 (CEA-CONF-2010)

EXPERIMENTAL BREEDER REACTOR-II

- reactivity feedback analysis by rod drop in, digital data reduction procedure for on-line, 26:47307

EXPLOSIONS

- detection and location of shallow-depth chemical, in northwest Pacific Ocean, performance of hydrophone method for, 26:45489

EXPLOSIVES

- description of nuclear, for production stimulation and storage of hydrocarbons underground, 26:45493
- detonation properties of gelled-slurry Types DBA-22M, DBA-65T2, and DBA-105T2, large--scale tests of, 26:45484 (SC-DC-722296)

EYES

- radiation effects on lens cell proliferation and regeneration in rabbit, during injury response and wound healing, x , 26:45813(R) (COO-1427-41)
- radioinduced injuries to crystalline lens in rabbit, effects of dose on RBE for fast neutron, γ , mixed γ -neutron, and x , 26:45864

F

FALLOUT

- effects of Schooner Event close-in, on vegetation, (E), 26:45542 (PNE-529)
- effects on desert plant survival and reproduction at Nevada Test Site, combined beta and γ , 26:45759 (TID-25954, pp 54-7)
- effects on plants at Nevada Test Site, revegetation following lethal or pathological close-in, 26:45762 (TID-25954, pp 75-86)
- monitoring in atmosphere, bones, diets, and soils in Italy and U.S., 26:45511(R) (HASL-257)
- monitoring of close-in, following Schooner Event, 26:45542 (PNE-529)
- radioisotope content of soil surface-deposited, performance of continuously operating gamma spectrometer for determining, 26:45527 (TID-26103)
- transport through soil-plant-small mammal ecosystems at Nevada Test Site, 26:45914 (TID-25954, pp 58-64)

FAST TEST REACTOR

- core components for, testing of Type 316 stainless steel for, 26:46067 (HEDL-TME-71-126)
- kinetics of, computer program calculation of three dimensional, 26:47586 (HEDL-TME-72-42)

FECES

- radium-226 excretion in human, following oral uptake as radium sulfate dust, effects of alginate ingestion on, 26:45907

FEDERAL REPUBLIC OF GERMANY

- nuclear power plants for, site selection criteria for, 26:47574
- radiation protection programs for nuclear personnel in, medical aspects of, 26:45803

FERMI FAST BREEDER REACTOR

- coolant for, failure to measure radioactivity of liquid sodium, 26:47340 (DOCKET-50016-165)
- coolant storage tank for, request for delay in drainage test of emergency, 26:47327 (DOCKET-50016-160)
- operation of, request for extension of license expiration date for, 26:47390 (DOCKET-50016-163)
- operations during April 1972, 26:47339 (DOCKET-50016-164)

FERMI POWER REACTOR, UNIT 2

- building for, cracks in concrete basement floor slab of, 26:47513 (DOCKET-50341-42)
- containment structure drywell for, deformation of, 26:47514 (DOCKET-50341-43)
- pressure vessel for, ultrasonic discontinuities in welds on, 26:47329 (DOCKET-50341-44)

FERMIONS

- energy of basic state of many-body system of, coupled by gravitational lines, (T), 26:46748 (JINR-P2-6270)
- quantum theory for spin-1/2, in external electric fields, (T), 26:46741
- S-matrix for spin-1/2, in external electric fields, (T), 26:46741
- theory for states of, generating function method for expansion of Hamiltonian operator from isomorphism between space of boson states and space in, (T), 26:47286 (LYCEN-7202)

- theory for, unitary operator form of general Bogolubov--Tyablikov transformation in, (T), 26:47291

FERROCENE

- radiation effects on, adsorbed on glass, γ , 26:45313(R) (COO-3029-1)

FETUSES

- hematologic changes toward aleukemic signs in peripheral blood of dog, following serial transplantation of radioinduced canine myelomonocytic leukemia, 26:45897
- radiation dose to thyroid gland of mouse, following injection of iodine-131, mathematical model for calculating β , 26:45902
- radioinduced cardiovascular malformations in rat, following sublethal whole-body exposure in utero, statistical analysis of relation between age, dose, and incidence of fast neutron, 26:45867

FIBERS

- radiation effects on Kapron, x-ray scattering studies on electron or γ , 26:45338

FIELD EMISSION MICROSCOPY

imaging of defects in ion-implanted solids, 26:47239
imaging process in, 26:46038

FIELD THEORY

energy--momentum quadrivector in complex fields in, Maxwell tensor for, 26:46693
field equations in formulation of Einstein--Kaufman nonsymmetric unified, static, spherically symmetric solutions of, 26:46694
field equations in, cylindrical wave solutions of Bonner and Schroedinger unified, 26:46695
field interactions with essentially nonlinear field in, space--time curvature for, 26:46692
formulation of Einstein--Kaufman nonsymmetric unified, 26:46694
geometry for unified, Lyra and Weyl, 26:46688
Klein--Gordon equation for Hulthén and Schwarzschild potentials in, 26:46690
Maxwell tensor for energy--momentum quadrivector in complex fields in, 26:46693
metrics determination in field with given Lagrangian in, space--time curvature in essentially nonlinear fields for, 26:46692
polarization quadrivector in complex fields in, tensor for, 26:46693
space--time curvature in essentially nonlinear fields in, 26:46692
tensor for polarization quadrivector in complex fields in, 26:46693
Yang--Mills classical field equations, exact solutions to, 26:46689

FILMS

preparation of refractory oxide, for electrolytic membranes and radiation windows, method for, 26:45958(P)
thickness of thin, beta backscattering gage for measuring, 26:45656(P)

FILTERS

performance of impregnated charcoal bed, for removing iodine-131 from ventilation exhaust streams, 26:45502
performance testing of high efficiency air, pp.11-13, 26:45673(R) (UCRL-50007-71-3)

FIRE EXTINGUISHING AGENTS

performance of Freon 13, radiometric evaluation by gamma attenuation, 26:45712

FISSION

conference on neutron-induced, 26:47015 (INDC-7-U)
mass and charge distributions in, theory for, 26:47044 (INDC-7-U, pp 65-79)
mass distribution in, systematics of, 26:47017 (INDC-7-U, pp 31-59)
measurements on neutron-induced, review on, 26:47036
microscopic models for, development of, (T), 26:47046 (INDC-7-U, pp 145-60)
probabilities in $A = 208$ region, single-particle effects on, (T), 26:47070
prompt γ -ray and neutron emission in, review of, 26:47020 (INDC-7-U, pp 113-29)
prompt radiation emission in, survey on, 26:47019 (INDC-7-U, pp 89-112)
radiation emission in, review on, 26:47021 (INDC-7-U, pp 131-44)
shell effects in, excitation energy dependence of, (T), 26:47043 (INDC-7-U, pp 3-19)
widths for, distribution of total, (T), 26:47045 (INDC-7-U, pp 81-6)

FISSION FRAGMENTS

angular distribution from spontaneous fission of plutonium-236 and -237 and curium-243, (E), 26:47022 (JINR-P7-6466)
detection of, etching conditions for mica detectors for, 26:45610
effects on crystal structure of thorium dioxide--uranium dioxide fuel elements, 26:45939
effects on tungsten thermionic fuel cladding, 26:45702(R) (LA-4978)
gamma emission from fission of californium-252 and uranium-235, p. 314-8, (E), 26:46873(R) (CEA-N-1522)
mass and charge distributions of, theory for, 26:47044 (INDC-7-U, pp 65-79)
mass distributions of, systematics of, 26:47017 (INDC-7-U, pp 31-59)
neutron emission from fission of californium-252 and uranium-235, p. 314-8, (E), 26:46873(R) (CEA-N-1522)
prompt γ -ray yields from, calculation using improved renormalized Fermi gas model, (T), 26:47025

FISSION PRODUCTS

behavior in molten salt fuels, review of, 26:47302
chemistry in HTGR, 26:45288 (TID-26106)
chemistry in oxide fuel dissolution, 26:45407(R) (ORNL-TM-2918)
containment in pyrochemical reprocessing of LMFBR fuels, 26:45421
data tabulation on, 26:47012 (CEA-N-1526)
deposition in oceans, relations with atmospheric beryllium-7 content, 26:45530 (BNWL-SA-4205(Rev.))
determination of isotope distribution of, in californium-252 and curium-244 by mass spectroscopy, 26:45260
diffusion from oxide fuel, 26:45412(R) (ORNL-TM-3095)
diffusion in oxide fuels, 26:45417(R) (ORNL-TM-3281)

diffusion of gaseous, from oxide fuels, 26:45411(R) (ORNL-TM-3072)
fixation in phosphate glasses, 26:45385
reactions in oxide fuels, 26:45413(R) (ORNL-TM-3127)
release in reprocessing of LMFBR fuels, 26:45406(R) (ORNL-TM-2871)
removal from molten salt fuels, development of processes for, 26:47302
sampling in reactor effluents, design of remote-controlled sequential sampler for, 26:45613(P)
separation from gas streams by sorption and reaction on zeolites, 26:45387(P)
solubility in nitric acid, 26:45408(R) (ORNL-TM-2949)
solubility in nitric acid, 26:45409(R) (ORNL-TM-2996)
volatilization from oxide fuels, 26:45409(R) (ORNL-TM-2996)
volatilization from oxide fuels, 26:45406(R) (ORNL-TM-2871)
volatilization from oxide fuels, 26:45410(R) (ORNL-TM-3018)
volatilization from oxide fuels, 26:45411(R) (ORNL-TM-3072)
volatilization from oxide fuels, 26:45412(R) (ORNL-TM-3095)
volatilization from oxide fuels, 26:45416(R) (ORNL-TM-3250)
volatilization from oxide fuels, 26:45415(R) (ORNL-TM-3217)

FISSION TRACK ANALYSIS

errors in, prolonged etching factor for correcting, 26:45604

FISSIONABLE MATERIAL

book: Atomic Safeguards: A Study in International Verification, 26:46096
shipping of, design and testing of drum-type shielded container for, 26:46114 (MLM-1911)

FLUID FLOW

calculation of cross-flow component in rod bundles, method for, 26:47596
heat transfer in laminar, of partially ionized argon, effects of transverse magnetic field on, 26:45464
heat transfer in longitudinal laminar, effects of lateral rod displacement on, 26:45462
hydrodynamics of two-phase, computer program for analysis of, 26:45458 (CISE-R-245)
magnetohydrodynamic entrance, in channel, (T), 26:46423
magnetohydrodynamic rotating, sphere motion in electrically conducting, (T), 26:46424
magnetohydrodynamic secondary, in rotating straight annular pipe, (T), 26:46422
magnetohydrodynamic, weak fluctuation modes of electrically and thermally conducting viscous compressible, (T), 26:46420
stability of forced convection, in parallel channels, (E), 26:45459 (CISE-R-275)

FLUIDIZED BEDS

solids processing on, 26:45418

FLUIDS

conference on thermophysical properties of, Washington, D. C., Dec. 1, 1971, 26:45463
mixing and separation of multi-phase, design of mixer-settler for simultaneous, 26:45451(P)

FLUORENONE

reactions with hydrated electrons and hydrogen atoms from radiolysis of aqueous solutions, 26:45354

FLUORIDE VOLATILITY PROCESSES

evaluation for use in Europe, 26:45419

FLUORIDES

determination of volatile, in uranium hexafluoride by infrared spectrophotometry, 26:45238
effects on yields of trapped electrons in radiolysis of doped ice, 26:45345
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

FLUORINE

determination in fluorites and plants by neutron activation using plutonium--beryllium source, 26:45199

FLUORINE ISOTOPES F-18

energy levels of, calculation using Bethe--Goldstone equation in shell-model basis, 26:47074
energy levels of, data compilation on, 26:46882
energy-level transition in, calculation of circular polarization of 1.08-MeV γ ray using various weak interaction theories, (T), 26:46877

FLUORINE ISOTOPES F-19

energy levels of, calculation using alpha cluster model with weak coupling, 26:47083
energy levels of, data compilation on, 26:46882
energy-level transitions in, multipole mixing ratios for, (E), 26:46893
lithium-6 reactions (Li-6,d) at 16 MeV, angular distributions for, (E), 26:46915
meson reactions (π^- , π^0) at 140 to 240 MeV, cross sections for, (E), 26:46486
proton reactions (p,He-3) and (p,t) at 45 MeV, cross sections and angular distributions for, (E), 26:46895

FLUORINE ISOTOPES F-20

energy levels of, data compilation on, 26:46882

FLUORINE SYSTEMS

Al_2O_3 -- CaO -- Cr_2O_3 -- F -- MgO -- MoO_3 -- TiO_2 -- ZrO_2 , use as welding flux, 26:45941(P)

FLUORITES

analysis for fluorine by neutron activation using

- plutonium--beryllium source, 26:45199
analysis of artificial, for rare earths, use of paper chromatography in, 26:45232
- FLUOROCARBONS**
radiolysis of, research progress on synthesis, identification, and purification of C₆ and C₇ fluorocarbons from, 26:45316(R) (ORO-3909-3)
- FOAMS**
molding process for syntactic resin, 26:47600(R) (BDX-613-253(Rev.))
- FOOD**
analysis for trace elements by neutron activation and radiochemical separation, 26:45189(R) (BFL-1/1972)
strontium-90 passage from bovine bones to processed products, 26:45549
- FOOD CHAINS**
radioisotope transport in aquatic, mathematical model for calculation of, 26:45547
sodium-22 content of lichen-reindeer-man, in Sweden during 1961 to 1969, fallout, 26:45909
- FORBUSH DECREASES**
relation with solar-flare-associated interplanetary streams in solar wind, (E/T), 26:46332
- FORENSIC SCIENCE**
activation analysis uses in, 26:45213
x-ray fluorescence analysis uses in, nondispersive, 26:45213
- FORESTS**
strontium-90 accumulation and distribution in natural plant associations of, in Carpathians, 26:45551
- FORKED RIVER POWER REACTOR, UNIT 1**
operation of, correction of cost-benefit analysis for, 26:47526 (DOCKET-50363-32)
safety of, revised and additional data on, 26:47524 (DOCKET-50363-30)
safety of, revised data on, 26:47525 (DOCKET-50363-31)
site for, commercial fishing in Barnegat Bay, 26:47527 (DOCKET-50363-34)
- FORMIC ACID**
carbon-13 NMR study of, 26:45297
reactions with plutonium dioxide sols, 26:45408(R) (ORNL-TM-2949)
spin-lattice relaxation times in, effects of carbon-13 on, 26:45297
- FORT CALHOUN POWER REACTOR, UNIT 1**
safety evaluation of, Directorate of Licensing, 26:47454 (DOCKET-50285-68)
- FORT ST. VRAIN POWER REACTOR, UNIT 1**
control rods for, computer simulation of pulsed neutron experiments for reactivity worth of, 26:47319 (GULF-GA-B-10959)
- FRANCE**
cardiac pacemaker development in, review on, 26:45699 (CEA-COINF-1998)
plutonium-238 production and uses in medicine in, review of, 26:45706
radiation protection programs for environment and personnel at Marcoule nuclear center, 26:45804
radioisotope content of flour and wheat in, during 1960 to 1969, cesium-137 and strontium-90, 26:45903
uranium enrichment in, economic studies of gaseous diffusion process for, 26:45691
- FREE RADICALS**
reactions of radiolytic, photolytic, and thermolytic, with mercaptans, isotope effects on, 26:45342
- FRUITS**
avocado, radiation effects on ethylene production, respiration, and ripening time of, effects of dose and storage time on γ , 26:45724
fallout cesium-137 and strontium-90 content in citrus and tropical, Mecklenburg, GDR, 1969, 26:45546
radiation processing of, effects on respiration rate, γ , 26:45729
- FUEL CAPSULES**
design of plutonium-238 heat source, for thermoelectric and thermionic generators, 26:45708(P)
fabrication of radioisotope heat source, method for, 26:45707(P)
materials for high-temperature, development and testing of, 26:45703(R) (ORNL-4796)
testing of cermet fuel-containing, development of equipment for impact, 26:45703(R) (ORNL-4796)
- FURNACES**
lining wear in blast, use of implanted cobalt-60 sources for measuring, 26:45720
- FUSED SALTS**
bibliography on, author, citation, and subject--title list for, 26:45287 (SC-R-722769)
- GADOLINIUM**
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
electron spin resonance in metallic systems, relation between s--d hybridization and g shifts in, (T), 26:47271
electron spin resonance in metallic systems, relation between band structure and g shifts in, (T), 26:47270
magnetic susceptibility of single crystals of, anisotropy in para-, (E), 26:47253
properties as soluble nuclear poison, 26:45405(R) (ICP-
- 1006)
separation from europium, neodymium, praseodymium, and samarium by anion exchange using acetic acid--methanol--nitric acid mixtures, 26:45280
- GADOLINIUM ALLOYS AND SYSTEMS**
Fe--Gd, magnetic compensation in films of, 26:47273
Gd--La, critical-field and proximity-effect behavior of superconducting, many-body effects from, (E/T), 26:46822
- GADOLINIUM ISOTOPES**
x-ray spectra from, isotope shifts in, (E), 26:46973
- GADOLINIUM ISOTOPES Gd-148**
energy-level scheme from terbium-148 decay, (E), 26:46942 (JINR-P6-6512)
- GADOLINIUM ISOTOPES Gd-150**
energy-level scheme from terbium-150 decay, (E), 26:46942 (JINR-P6-6512)
- GADOLINIUM ISOTOPES Gd-154**
energy-level transitions in deformed, SU(3) parametrization of E₂, (T), 26:46975 (JINR-P4-6462)
- GADOLINIUM ISOTOPES Gd-156**
energy levels from europium-156 beta decay, band mixing in, (E), 26:46996
energy-level transitions from europium-156 beta decay, branching ratios and probabilities for, (E), 26:46996
- GADOLINIUM ISOTOPES Gd-158**
energy levels from europium-158 beta decay, band mixing in, (E), 26:46996
energy-level transitions from europium-158 beta decay, branching ratios and probabilities for, (E), 26:46996
- GAGES AND METERS**
nonlinearity of dc Q-meter, corrections for, 26:45680
- GALAXIES**
associations between quasars and, physical, (E/T), 26:46258
distribution of, x-ray study of lumpy, (E/T), 26:46260
field equations for clusters of, relativistic gravitational, (T), 26:46253
hydrogen in NGC 45, mass of neutral, (E), 26:46252
infrared emission from nuclei of, (T), 26:46234
interactions in groups of, (T), 26:46234
interactions with galaxies, cosmological model with positive- and negative-mass particles for, (T), 26:46761
mass distribution in Coma Cluster of, (E), 26:46242
mass of NGC 45, (E), 26:46252
motions in clusters and groups of, (T), 26:46235
photometric standard sequence for LMC, (E), 26:46238
properties of peculiar, (T), 26:46234
radio map of M84 at 2.7 and 5.0 GHz, (E), 26:46200
radio map of Maffei 2 at 1415 MHz, (E), 26:46237
radio observations at 21 cm of NGC 45, (E), 26:46252
spectra of nuclei of NGC 1052, 2655, 2903, and 4569, (E/T), 26:46239
spectra of nuclei of, radial cyanogen-band gradients in, (E), 26:46243
spectral redshifts in Coma Cluster, relation to magnitude and morphology, (E/T), 26:46241
spectral redshifts of, (T), 26:46234
star birth and death processes in, survey of, (T), 26:46166
stellar population of nuclei of, (E/T), 26:46239
structure of Coma Cluster of, from analysis of radial velocities, (E), 26:46242
x-ray luminosity and velocity dispersion in clusters of, (T), 26:46249
- GALLIUM**
determination by fluorometry using Superchrome Garnet Y, 26:45236
effects on recrystallization of iron, 26:45982
separation from aluminum, copper, iron, germanium, and rare earths by solvent extraction using methyl isobutyl ketone, 26:45285
separation from hydrochloric acid media by solvent extraction using N-benzylaniline in chloroform, 26:45284
- GALLIUM ALLOYS AND SYSTEMS**
Ga--Nb, phase diagram of, 26:46018
- GALLIUM ARSENIDES**
proton implantation at 300 keV in n-type, production of optical waveguides by, carrier elimination in, 26:47232
- GALLIUM ISOTOPES Ga-72**
energy levels and properties of, compilation of, (E), 26:46939
- GAMETES**
radiation effects on frequency of chromosome loss in male, of Drosophila, enhancing effects of dose fractionation and lucanthone on x, 26:45887
- GAMMA-ABSORPTION ANALYSIS**
use for evaluation of Freon 13 fire extinguisher operation, 26:45712
- GAMMA CAMERAS**
readout system for, television type, 26:45595(P)
- GAMMA DECAY**
transitions between anti-analog states in, shell-model analysis of, 26:47050
- GAMMA RADIATION**
attenuation in matter, method for determining mean energy of secondary electrons from, 26:47213

- detection by monkeys and rats, visual, 26:45817 (ORO-2690-12(Pt.B), Paper 12)
- detection of, directional properties of cylindrical scintillation detectors with sleeved lead screen for, 26:45605
- distribution at large distances from source, generalized method for calculation, 26:47215
- dose rate measurements of, radiation and temperature effects on Radocon detectors for, 26:45620
- dosimetry of, design of tissue-equivalent liquid ionization chamber for, 26:45621
- dosimetry of, performance of LiF, photographic film, and photoluminescent dosimeters for, 26:45630
- effects of cobalt-60, on oscillation frequency of AT-cut quartz, 26:45667(T) (SC-T-722519)
- effects of pulsed, on electric conductivity of polymers, 26:47225 (AWRE-O-31/72)
- effects on alkylation of caffeine by amines, 26:45326
- effects on ammonia-olefin mixtures, yields of hydrogen from, 26:45332
- effects on anatomical structure and chemical composition of wood in spruce seedlings, 26:45774
- effects on ants, effects on fecundity of female and worker, dose dependence of, 26:45877
- effects on aqueous solutions of glucose phosphates, 26:45337
- effects on aqueous solutions of thymine, 26:45314 (COO-3286-1)
- effects on avocados, effects of dose and storage time on ethylene production, respiration, and ripening time, 26:45724
- effects on blood vessels and bone marrow of rabbit mandible, dose dependence of pathological, 26:45835
- effects on bone marrow of newborn swine, effects of dose on induction of anemia and leukopenia by, 26:45852
- effects on bones and uterus of parabiont and single rats following focal internal exposure, neoplastic cobalt-60, 26:45814(R) (COO-3017-43)
- effects on carbon tetrafluoride and hexafluoroethane, production of large-molecular-weight fluorocarbons by, 26:45316(R) (ORO-3909-3)
- effects on carnation cuttings, effects of yeast extracts on recovery from, 26:45777
- effects on carrier lifetime and isochronal annealing of silicon p-n diodes, 26:45671
- effects on chemical behavior of recoil copper-64 atoms in neutron-irradiated copper phthalocyanine, 26:45356
- effects on chromosomes in human bone marrow cells following single local exposure, effects of dose and post-exposure time on frequency of aberrant, 26:45794
- effects on crosslinking and polymerization of polyacrylates dissolved in vinyl monomers, 26:45339(P)
- effects on crystal lattice defect formation in alloys and metals, review on, 26:46076
- effects on crystalline and glassy vinyl acetate, effects of light on ESR spectra of free radicals from, 26:45334
- effects on dextran, effects on chemical and physical properties of, 26:45730
- effects on DNA of *Bacillus subtilis*, effectiveness of toluene treatment for enhancing semi-in vitro repair of transforming, 26:45731
- effects on DNA synthesis in recombination-deficient strains of *Escherichia coli*, inhibitory, 26:45753
- effects on eggs of sea urchins, effects on mitosis, 26:45874
- effects on enzyme activity in plasma membranes of rat liver, combined effects of thyroid hormones on, 26:45831
- effects on Ephedra (mormon tea) plants, lethal dose determination, 26:45764 (TID-25954, pp 102-9)
- effects on epithelium in midgut of adult and larval yellow mealworms, dose and time dependence of pathological, 26:45875
- effects on ethanol glass, optical absorption and ESR spectral studies of trapped electrons from, 26:45319
- effects on extracellular bacteriophages, lethal and mutagenic, 26:45746
- effects on ferrocene adsorbed on glass, 26:45313(R) (COO-3029-1)
- effects on *Franseria dumosa* (burr sage or burro bush) following acute exposure, dose dependence of growth and survival, 26:45768 (TID-25954, pp 153-61)
- effects on frog muscles, effects on localization of acetylcholinesterase in neuromuscular junction of, 26:45828
- effects on fruits, effects on rate of respiration in, 26:45729
- effects on gaseous carbon dioxide at high densities, 26:45346
- effects on glassy barium oxide-silicon oxide and silicon oxide-strontium oxide systems, spectral studies on color centers induced by, 26:47226
- effects on graft polymerization of hydroxyalkyl methacrylates on polymers, 26:45312(R) (COO-2147-2)
- effects on graft polymerization of unsaturated monomer containing hydrophilic groups in polypropylene for inducing dyeability, 26:45336(P)
- effects on growth of yucca plants following acute exposure, dose dependence of, 26:45769 (TID-25954, pp 188-98)
- effects on hematopoietic cells in bone marrow, lymph nodes and spleen of newborn swine, effects of dose on pathological, 26:45853
- effects on human liver following therapy of adjacent organs, dose and time dependence of pathological, 26:45809
- effects on human skin following therapy of internal tumors, pathological, 26:45793
- effects on immune reactions of rabbits to *Salmonella typhimurium*, protective effects of AET against suppressive, 26:45824
- effects on infective activity of purified Venezuelan equine encephalitis virus immunogenicity following inactivating, 26:45752
- effects on insects, effects on growth, development, and survival of tussock moth larvae, effects of dose on, 26:45885
- effects on insects, effects on life span, mating competitiveness, and reproductive capacity of adult stable flies following exposure of pupae, effects of dose and pupal age at exposure on, 26:45883
- effects on insects, effects on respiration rate in grain mites, effects of dose, life stage, sex, and temperature on, 26:45884
- effects on iron metabolism in mice following whole-body exposure, protective effects of AET and mexamine against pathological mixed neutron and, tracer study using iron-59, 26:45840
- effects on Japanese cedars, mutagenic for albinism, 26:45771
- effects on Kapron fibers, x-ray scattering studies on, 26:45338
- effects on large intestine and vagina of man following combined external and intracavitary therapy of uterine carcinoma, surgical repair of pathological, 26:45800
- effects on length of aluminum, copper, molybdenum, and nickel, 26:46075
- effects on lymphomagenic properties of neonatal mouse thymus gland subsequently transplanted into Gross murine leukemia virus-infected hosts, 26:45818 (UCLA-12-867)
- effects on mechanical properties and structure of cellulose acetate-polyacrylonitrile, 26:46084
- effects on mice following acute whole-body exposure, comparison of RBE with pulsed x rays for lethal, 26:45866
- effects on mice, effects on DNA metabolism in intestine of, in vitro and in vivo, RBE in comparison to proton, 26:45842
- effects on mice, effects on heart rate and survival time of, effects of dose, dose rate, and phenobarbital on, 26:45846
- effects on microwave bipolar transistors, (E/T), 26:45663 (SC-DC-721233)
- effects on migration of salamanders in Rhode Island, effects of dose rate, species, and temperature on, 26:45815(R) (COO-3486-1)
- effects on neutral and acidic liquid ethanol, effects of pressure on reactions of solvated electrons from, 26:45351
- effects on plants at Nevada Test Site, effects on growth and reproduction of, chronic, 26:45757 (TID-25954, pp 12-48)
- effects on polymerization of butyl acrylate-styrene mixture onto cellulose and cellulose acetate, 26:45320
- effects on polymerization of kaolin-methyl methacrylate composites, 26:45327
- effects on polymerization of monomers, effects of peroxide sensitizers on dose rate in, 26:45325
- effects on polymerization of trialkylacryloxystannanes in wood, 26:45355(P)
- effects on polymerization of vinylphosphonic dichloride with methyl methacrylate and styrene, 26:45318
- effects on potato plants, mutagenic, 26:45770
- effects on rabbits, effects on crystalline lens of, effects of dose on RBE for, and mixed neutron and, in comparison with x radiation, 26:45864
- effects on rats, relation to development of taste aversions, 26:45817 (ORO-2690-12(Pt.B), Paper 12)
- effects on reaction between cyclohexanone and chlorinated hydrocarbons, yields from, 26:45335
- effects on ruby and corundum crystals, 26:47244
- effects on ruby laser radiation, 26:47227
- effects on *Salsola* (Russian thistle) following seed exposure, relation of dose to germination and seedling survival, 26:45767 (TID-25954, pp 147-52)
- effects on seeds of desert plants, effects of dose and dose rate on germination and growth of, 26:45758 (TID-25954, pp 49-53)
- effects on seeds of *Lycium palladium* (desert shrub), relation of dose and moisture content to germination and seedling survival, 26:45761 (TID-25954, pp 73-4)
- effects on semiconductor devices due to induced high-amplitude current pulses in circuits, 26:45666 (SC-RR-720515)
- effects on shrubs at Nevada Test Site, effects on growth and reproduction during chronic five-year exposure, 26:45756 (TID-25954, pp 5-11)
- effects on single crystals of deuterated maleic acid, ESR spectral studies of trapped vinyl-type radical from, 26:45347
- effects on single crystals of potassium chloropalladate

- and potassium chloroplatinate, ESR studies on, 26:45341
- effects on small intestine of man following therapy of uterine carcinomas, symptoms and treatment of pathological, 26:45802
- effects on survival of 30 species of desert plants, dose dependence of, 26:45760 (TID-25954, pp 65-8)
- effects on survival of peas, effects of genotype on, 26:45775
- effects on tongue of rabbits, permeability of lingual mucosa during development and recovery from pathological, 26:45862
- effects on visual response of noctuid moths, 26:45873 (TID-26078)
- effects on γ -type zeolites, proton ENDOR studies on, 26:45324
- emission from 24-kt nuclear explosions, calculation of time-dependent, 26:45497
- emission from Crab Nebula, detection of, (E), 26:46214
- emission of pulsed, from NP 0532, (E), 26:46199
- energy spectrum from Sun, review of solar-flare-associated production mechanisms of flux and, 26:46231
- interactions with alloys, effective atomic numbers for photoelectric process in various shells in, (T), 26:47206
- interactions with alloys, effective atomic numbers for differential processes in, (T), 26:47207
- monitoring in environs of Brookhaven National Laboratory during June to July 1968, 26:45515
- monitoring in off-site environs of Nevada Test Site during July through Dec. 1971, 26:45515
- origin of galactic, (E), 26:46154
- polarization measurements on, development of two-section Ge(Li) compton, (E), 26:46893
- production of monochromatic, at CEA EL3 reactor, p. 309-11, 26:46873(R) (CEA-N-1522)
- reactions (γ, α) with oxygen-16 at 13.1 MeV, detailed balance test of time reversal invariance in, (E), 26:46881
- scattering by copper at 0.662 MeV, absolute differential cross sections for, (E), 26:47204
- scattering by lead-208 at 7.170 MeV, gamma spectrum from resonant, p. 313-4, (E), 26:46873(R) (CEA-N-1522)
- spectra from resonance neutron capture, review on, 26:46969
- GAMMA SOURCES**
- determination by γ spectroscopy, description of GAMANAL program in, 26:45265 (UCRL-51061(Vol.1))
- determination of extended, without sampling, 26:45264 (SRARI-P-123)
- determination of low-activity, recommendations for, 26:45261 (ICRU-22)
- development for photon dosimetry standards, 26:47216
- fabrication of specified dose rate distribution, method for, 26:45715
- isodose curves for sealed, calculation and measurement of, 26:45715
- preparation of standard, 26:45181 (CISE-R-172)
- GAMMA SPECTROMETERS**
- calibration of scintillation, for extended source analysis, 26:45264 (SRARI-P-123)
- circuit for Ge(Li)-NaI(Tl), fast-slow coincidence, 26:45639
- GAMMA SPECTROSCOPY**
- performance for measuring soil surface-deposited fallout, 26:45527 (TID-26103)
- spectra analysis in, description of GAMANAL program for, 26:45265 (UCRL-51061(Vol.1))
- GAS CHROMATOGRAPHY**
- diffusion-limited phenomena in, 26:45302(R) (COO-1222-55)
- peak shape analysis in, 26:45302(R) (COO-1222-55)
- GAS FLOW**
- magnetohydrodynamic, vorticity jump across stationary discontinuity in, 26:46421
- GASEOUS DIFFUSION PLANTS**
- development of international civilian, in Europe, plans for, 26:45693
- GASEOUS DIFFUSION PROCESS**
- efficiency for uranium enrichment, comparison with other isotope separation processes, 26:45689 (ORO-694)
- uranium enrichment using, French economic studies on, 26:45691
- GASES**
- reactions with graphite, research program on, 26:46078 (NYO-1710-106)
- transport in rarefied, random method for solution of Boltzmann's equation for, 26:47208
- GASTROINTESTINAL TRACT**
- radioinduced injuries to, of rats, protective effects of bile acid sequestrant against x, 26:45844
- radioinduced neoplasms in, following uptake of radioisotopes, review of pathology of, 26:45895 (UR-3490-102)
- GATING SYSTEMS**
- circuits for fast, p. 152-3, 26:46873(R) (CEA-N-1522)
- GENES**
- effects on gamma-radiosensitivity of peas, 26:45775
- GENETIC RECOMBINATION**
- radioinduced, in *Drosophila* in relation to tergite development, 26:45881
- GEOMAGNETIC FIELDS**
- activity of, 12-month wave data from 1884 to 1968, (E/T), 26:46316
- activity of, effect on 1965-1968 electron content in low and middle latitudes of ionosphere, (E/T), 26:46287
- activity on Feb. 2, 1969, 1f-wave observations near interplanetary discontinuities, (E), 26:46302
- disturbances from 1884 to 1967, variations in 11-year solar cycles, (E), 26:46322
- pulsations in daytime auroral zone, relation of polar and magnetospheric substorms with Pld irregular, (E), 26:46317
- pulsations in, association of quasi-periodic vlf emissions with micropulsations, (E), 26:46319
- pulsations in, relation of ionospheric spread F with Pc 1 micropulsations, Jan. 1, 1970 to June 30, 1971, (E/T), 26:46274
- GEOPHYSICS**
- equipment and instruments for, review of Hungarian, 26:45560
- GERMANIUM**
- analysis for trace amounts of impurities by instrumental neutron activation, 26:45218
- domain instability in 8.5-MeV electron irradiated, recombination, 26:47230
- purification by zone melting in rotating container, 26:45276
- radiation effects on volt-amp characteristics of n-type, 8.5-MeV electron, 26:47230
- separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
- GERMANIUM ALLOYS AND SYSTEMS**
- Al-Ge-Nb, critical temperature of superconducting intermetallic Nb₃Al/sub x/Ge/sub 1-x/, variations in, 26:46021
- Al-Ge-Nb, phase diagram of niobium corner of, 26:46031
- Al-Ge-Nb, superconducting properties of, 26:46031
- Ge-Si-base, etching by hydrofluoric acid-nitric acid solutions, 26:45935 (SC-RR-710884)
- Ge-Si-base, oxidation by air and water, effects of temperature on, 26:45935 (SC-RR-710884)
- GERMANIUM ISOTOPES Ge-68**
- energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo-SU₃ coupling schemes, 26:46931
- GERMANIUM ISOTOPES Ge-70**
- neutron capture by, γ spectra from thermal, (E), 26:46937
- GERMANIUM ISOTOPES Ge-71**
- energy levels from germanium-70 neutron capture, (E), 26:46937
- GERMANIUM ISOTOPES Ge-72**
- energy levels and properties of, compilation of, (E), 26:46939
- GERMANIUM ISOTOPES Ge-74**
- heavy ion scattering by, time-dependent deorientation of recoils in vacuum from, (E), 26:46935
- GLASS**
- analysis for trace amounts of platinum by neutron activation, 26:45217
- analysis of borosilicate, for boron by flame emission spectroscopy, 26:45254
- radiation effects on, sputtering, energy loss, and luminescence from 5.6-keV noble-gas-ion, 26:47238
- radioactive waste fixation in phosphate, 26:45385
- radioinduced optical absorption and paramagnetic centers in, of systems SrO-SiO₂ and BaO-SiO₂, 26:47226
- radioinduced photoconductivity in films of, x, (E/T), 26:45668 (UCRL-73974)
- GLOVE BOXES**
- description of dual-mode operation, for alpha-active materials, 26:46107 (UCRL-73756)
- GLOW DISCHARGES**
- properties of abnormal glow, 26:47138
- GLUCOSE PHOSPHATES**
- radiolysis of aqueous solutions of, yields of glucodialdose from γ , 26:45337
- GLUTATHIONE**
- effects on recovery of glucose-6-phosphatase activity in erythrocytes of whole-body x-irradiated rabbits and rats, 26:45848
- GLYCINE, N-ACETYL-**
- radiolysis of single crystals of, ESR and INDO MO studies of radicals from electron, 26:45349
- GOLD**
- determination in hair by neutron activation and γ spectroscopy, 26:45214
- determination in ores by lead crucible fusion and neutron activation, 26:45200
- determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
- effects on recrystallization of iron, 26:45982
- electron reactions at 35 keV, calculation of energy loss spectra from 1 to 12 eV from, (T), 26:47212
- electron transport in, discrete ordinates method for 1- to 8-MeV, (E/T), 26:47210
- neutron transmission below 10⁻⁴ eV through films of, interference effects in cold, (E), 26:46853
- positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
- radioinduced grain-boundary dislocation climb in

bicrystals of, 20-keV gold-ion, 26:47241

GOLD IONS
effects on grain-boundary dislocation climb in gold bicrystals at 20 keV, 26:47241

GOLD ISOTOPIES Au-194
decay of, γ and conversion electron spectra from, (E), 26:47003

GOLD ISOTOPIES Au-197
alpha inelastic scattering up to 80 MeV, α spectra and cross section for, (E), 26:47008
alpha reactions (α ,He-3) up to 80 MeV, energy spectra and cross section for, (E), 26:47008
helium-3 inelastic scattering up to 80 MeV, energy spectra and cross section for, (E), 26:47008
helium-3 reactions (He-3, α) up to 80 MeV, α spectra and cross section for, (E), 26:47008
lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
lithium-7 breakup reactions with, mechanism for, p. 24-6, (E), 26:46873(R) (CEA-N-1522)
neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)
proton reactions (p,n) with, production of carrier-free mercury-197 by, 26:45395

GOLD ISOTOPIES Au-198
availability and cost of radiation sources of, produced by Norway Institute for Atomic Energy, 26:45709 (INIS-mf-256)
effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

GRAPHITE
diffusion of fission-product xenon-133 from, to 1400°C, 26:46078 (NYO-1710-106)
impregnation with pyrolytic carbon, pulsed method for, 26:46088(P)
mechanical properties of, research program on, 26:46078 (NYO-1710-106)
muon (μ^-) range--momentum correlation in, 500- to 1500-MeV/c, (E), 26:46448 (LBL-791)
muon (μ^+) range--momentum correlation in, 500- to 1500-MeV/c, (E), 26:46448 (LBL-791)
neutron diffusion in, temperature dependence of diffusion length for, (T), 26:46857
preparation of fiber-reinforced composites of, 26:46086
radiation effects on, 26:46078 (NYO-1710-106)
radiation effects on high temperature properties of, neutron, 26:47302
reactions with gases, research program on, 26:46078 (NYO-1710-106)
research programs on, in West Germany, 26:46086
surface energy of, wetting studies on critical, 26:45944
surface tension of stress-annealed Type ZYH pyrolytic, 26:46091
uses of fiber-reinforced composites of, 26:46086
B₄C--graphite-base, radiation effects on boron-10-containing control materials of, fast neutron, 26:46077 (GULF-GA-A-12035)

GRAVITATION
black holes in Brans--Dicke theory of, gravitational mass loss in formation, equivalence principle violation, and absence of scalar gravitational radiation in collision of, 26:46192
constant of, effect of many-body forces and matter distribution on, 26:46765
energy emission by isolated material source, Einstein field equation solution for, (T), 26:46760
fermion system coupled by lines of, energy of basic state of many-body, (T), 26:46748 (JINR-P2-6270)
graviton exchange potentials, gauge transformation for, 26:46766
interactions in universal, repulsive, (T), 26:46749 (ORO-3992-94)
mass loss in black hole formation in Brans--Dicke theory, (T), 26:46192
screening from negative mass in cosmological model, (T), 26:46761

GRAVITATIONAL COLLAPSE
theory in magnetic field, exact solutions of Einstein equations for gravitational fields with sources of dust and magnetic fields in, (T), 26:46755

GRAVITATIONAL FIELDS
charged-particle motion in electromagnetic and, model for spinning, (T), 26:46767
Einstein equations for metrics with groups of motions on two-dimensional transitivity hypersurfaces in model with sources of dust and magnetic field, exact solutions of, (T), 26:46755
metrics with groups of motions on two-dimensional transitivity hypersurfaces in model with sources of dust and magnetic field, exact solutions of Einstein equations for, (T), 26:46755
neutrino interactions with spherically symmetric gravitational fields, Weyl equation for, 26:46772
particle motion in, canonical dynamics of spinning, 26:46753
relativistic Roche problem associated with black holes, (T), 26:46189
structure of spatial infinity, analysis of, 26:46752
theory for homogeneous rotating ring, null bundle transformations for Kerr solution for Einstein--Maxwell

equations in, (T), 26:46762

GRAVITATIONAL RADIATION
existence in isolated material source, Einstein field equation solution for, (T), 26:46760
spectra of high-frequency scalar, electromagnetic, and gravitational synchrotron, effective potential shape for differences among, (T), 26:46757
wave-induced elastic-body vibrations, equation of elastic waves for, (T), 26:46768

GROUND
motion from Cannikin Event, effects on Amchitka Island archaeological sites, (E), 26:45485 (SC-RR-720359)
motion from underground nuclear explosions, calculation of seismic-source energies from measurements of, 26:45490
motion of Hot Creek Valley, Nev., from underground nuclear explosions, methods for evaluating effects on soils and vegetation, 26:45478(R) (NVO-409-1)

GROUND WATERS
radioisotope transport by, from Rainier Event melt-glass, (E), 26:45482 (NVO-1229-177)

GROUP THEORY
conformal invariant bootstrap quantum field theory, matrix elements of internal symmetry currents, energy--momentum density tensor, and generalized unitarity in, 26:46697 (DESY-72/20)
Lie algebra in, conjugacy and stability theorems and classification for contraction of, 26:46666 (IC-72/49)
Lorentz transformations for electromagnetic field vectors for electric charge motion in constant and homogeneous electromagnetic fields, 26:46679
Poincare group for electric and magnetic form factors for NN ν vertex in, field theory on 8-dimensional homogeneous space of, 26:46514 (CTH-ITP-71-30)
Poincare group representations labelled by additional discrete quantum numbers, explicit, 26:46665 (CTH-ITP-71-14)
Poincare transformations for properties of one-particle wave functions of free arbitrary-spin particles in, 26:46667 (ITF-71-117-P)

GUANIDINE, 1-METHYL-3-NITRO-1-NITROSO-
effects on DNA in cultured human Xeroderma pigmentosum cells in comparison to effects of uv radiation, extent of repair synthesis following, 26:45742

GUINEA PIGS
immunization against equine encephalitis virus, effectiveness of gamma irradiated virus for, 26:45752

GUM NEBULA
origin and evolution of, (T), 26:46165

GYMNOSPERMIDAE
Cryptomeria japonica (Japanese cedar), radioinduced recessive mutations in, for albinism, gamma, 26:45771
Picea sp. (spruce), radiation effects on anatomical structure and chemical composition of wood in seedlings of, γ , 26:45774

H

H.B. ROBINSON POWER REACTOR, UNIT 2
boron injection tank for, incorrect dilution in, 26:47366 (DOCKET-50261-102)
steam line isolation valve for, malfunction of, 26:47365 (DOCKET-50261-101)

HI REGIONS
molecule abundances in, (E/T), 26:46244

HII REGIONS
distribution in spiral arm of Maffei 2, (E), 26:46237
fine structure in, (E/T), 26:46198
microwave emission from OH--H₂O sources associated with, (E/T), 26:46206
models for, (E/T), 26:46198

HADRONS
composite structure of, scaling law and size of urbaryons in, 26:46608
existence of isomeric states of, (T), 26:46588
interactions at high energies, dominance of one-photon-exchange mechanism in electromagnetic, (T), 26:46533
interactions at super-high energies, bounds on μ^+ -- μ^- pair and W^+ (W^-) production in, 26:46598
interactions of, consequences of vanishing of triple-pomeranchukon vertex for, 26:46597
interactions of, correlation functions and generalization in formulation of Feynman gas model from jet approach to deep inelastic, (T), 26:46586 (JINR-P2-6501)
interactions of, polarization in inclusive, (T), 26:46595
interactions of, space--time description of, 26:46603
interactions of, urbaryon rearrangement scheme for differential cross sections of inclusive, (T), 26:46606
interactions with deuterons, recoil effect in Glauber theory for quasi-elastic, 26:46653
interactions with hadrons at high energies and large angles, review of elastic, (E/T), 26:46591
interactions with hadrons at high energies, comparison of two-particle distribution functions in diffractive excitation and multiperipheral models for, (T), 26:46587 (RLO-2230-T-4-1)
interactions with hadrons at high energies, conference on, (E/T), 26:46590
interactions with hadrons at high energies, exchange degeneracy and dip mechanism in differential cross

- sections of, (E/T), 26:46592
interactions with hadrons in composite model, strong core in, 26:46607
interactions with hadrons, review of dual quark-parton model for, (T), 26:46515 (DAMTP-71/24)
interactions with leptons, relation between fixed poles in virtual Compton scattering and scale invariance in, 26:46559
interactions with photons, polarization in inclusive, (T), 26:46595
interactions with photons, scaling limit for virtual Compton, (T), 26:46530
interactions with protons at 10 and 14 GeV/c using polarized proton and deuteron targets, review of elastic, (E), 26:46436
interactions with virtual photons, relation between fixed poles and scale invariance in, 26:46559
production in $e + N \rightarrow e + \text{hadrons} + \text{anything}$ interactions via virtual photon $+ N \rightarrow \text{hadron} + \text{anything}$ interactions at high energies and momentum transfer, model-independent derivation of generalized scaling laws and multiplicity for, (T), 26:46521 (LBL-773)
production in semi-inclusive $e^+ + e^- \rightarrow \text{hadron} + \text{anything}$ interactions, review of dual quark-parton model for, (T), 26:46515 (DAMTP-71/24)
self-mass of, scaling behavior and finite electromagnetic, 26:46558
structure from neutrino interactions, 26:46584
structure model for, $SU(1,1)$ algebra classification for wave packet realization of lightlike particle states for formulation of parton, (T), 26:46589
structure model for, $SU(1,1)$ algebra representations for realization of lightlike particle states for formulation of parton wave packet, (T), 26:46681 (NYO-4076-19)
structure of, composite model and strong core for, 26:46607
structure of, quartet model in chiral $U(4) \times U(4)$, 26:46687
superfluid phase for system of, derivation of critical temperature for, (T), 26:46596
symmetry breaking of, mathematical characterization of internal, 26:46682
weak interactions of, electromagnetic divergences in, 26:46582
- HAFNIUM**
determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
effects on metallography of niobium-oxygen systems, electron microscopy studies on, 26:46039
neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)
production and uses of, review on, 26:45979
separation of protactinium from, in chloride and sulfate media by sorption on silica gel, 26:45390
- HAFNIUM ALLOYS AND SYSTEMS**
Hf-Nb-base, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)
Hf-Pt-base--Rh--Ti--W, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)
- HAFNIUM ISOTOPES**
x-ray spectra from, isotope shifts in, (E), 26:46973
- HAFNIUM ISOTOPES Hf-175**
energy-level scheme from tantalum-175 decay, (E), 26:46979
- HAFNIUM ISOTOPES Hf-176**
energy levels of, mixing of two-proton and two-neutron bands in, (T), 26:46992
- HAIR**
analysis for trace elements by neutron activation and γ spectroscopy, 26:45214
analysis for trace heavy metals by neutron and photon activation methods, 26:45198
- HAMSTERS**
radiation effects on ovarian function in pregnant, x, 26:45825
- HANFORD POWER REACTOR, UNIT 2**
design parameters of, requested additional data on, 26:47556 (DOCKET-50397-19)
operations aspects of, requested additional data on, 26:47556 (DOCKET-50397-19)
safety of, revised and additional data on, 26:47557 (DOCKET-50397-20)
safety of, revised and requested data on, 26:47558 (DOCKET-50397-21)
site for, revised and additional data on, 26:47556 (DOCKET-50397-19)
- HASTELLOY C**
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
properties for use as strontium fluoride waste capsules, 26:46099 (ARH-SA-117)
- HASTELLOY N**
radiation effects on high temperature properties of, neutron, 26:47302
- HASTELLOY X**
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
oxidation by air at 1000°C, 26:45701(R) (DP-1300)
- HAWAII**
radiation monitoring in marine environment of, during 1966 and 1968, 26:45534
- HAYNES 25 ALLOY**
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
- HAYNES 188 ALLOY**
corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
- HEART**
radiation effects on rate of, in mice, effects of dose, dose rate, and phenobarbital on gamma, 26:45846
radioinduced malformations in, of rat fetuses following sublethal whole-body exposure in utero, statistical analysis of relation between age, dose, and incidence of fast neutron, 26:45867
- HEART, MECHANICAL**
development of nuclear-fueled, 26:45700(R) (COO-2155-4)
efficiencies and losses in Kwan-Gett human, measurements of, 26:45700(R) (COO-2155-4)
heat sources for, French production of plutonium-238, 26:45706
power requirements for human, calculation of, 26:45700(R) (COO-2155-4)
testing of silastic, by implication in bovine calf, 26:45700(R) (COO-2155-4)
- HEAT EXCHANGERS**
design criteria for, for use with vapor generating equipment, 26:45436 (CEA-CONF-2010)
design for power plant using gas-cooled breeder reactor and gas turbine, parametric study of, 26:45439 (KFK-1566)
- HEAT PIPES**
design for controlling heat input to thermoelectric generator from radioisotope source, 26:45427(P)
design of, review on methods for, 26:45435 (CEA-CONF-2008)
fabrication of internally-finned, method for, 26:45449(P)
uses in industry, review on, 26:45435 (CEA-CONF-2008)
working fluids for cylindrical, calculation of vapor transport limits for, 26:45460
- HEAT-RESISTING ALLOYS AND METALS**
radioinduced density changes in, effects of stress on neutron, 26:46065 (CONF-710601-, pp 841-52)
- HEAT SOURCES**
encapsulation of radioisotope, method for, 26:45707(P)
production of radioisotope, for direct energy converters in ENEA countries, predictions to 1975 on, 26:45705
testing of radioisotope, for space uses, 26:45701(R) (DP-1300)
thermal energy transfer from radioisotope, to thermoelectric generator, design of heat pipe for controlling, 26:45427(P)
- HEAT TRANSFER**
analysis of multi-region, point-matching method for, 26:45465
- HEAT TRANSFER SYSTEMS**
design concepts for high-temperature liquid metal, review on, 26:45437 (CONF-720904-1)
- HEATED EFFLUENTS**
diffusion in streams, development of mathematical models for prediction of, 26:45575
discharge from power reactors, technical specifications for, 26:47437 (DOCKET-50271-92)
discharge into constant depth water channels, jet flow studies on, (E/T), 26:45582
discharge into large lakes, atmospheric effects of, 26:45584
dispersion of underwater-discharged, from power plants, method for acquiring and analyzing data on, 26:45579
droplet size in cooling tower and process stack, method for measuring, 26:45574 (CONF-720619-1)
effects of Candu reactor-discharged, on temperature structure of Lake Huron, 26:45581
effects of power plant-discharged, on temperature distribution in southwest Lake Michigan, 26:45580
effects of underwater-discharged nuclear power plant, on aquatic production in Lake Ontario, (EE), 26:45578
effects on aquatic environments, evaluation of, 26:45583
effects on environment of power reactor, 26:47397 (DOCKET-50213-152)
effects on environment of power reactor, 26:47424 (DOCKET-50255-96)
effects on environment of power reactor, 26:47404 (DOCKET-50237-249)
effects on environment of power reactor, 26:47446 (DOCKET-50275-22)
effects on environment of power reactor, 26:47401 (DOCKET-50220-98)
effects on environment of power reactor, 26:47473 (DOCKET-50309-61)
effects on environment of power reactor, 26:47532 (DOCKET-50368-31)
effects on environment of power reactor, 26:47435 (DOCKET-50271-88)
effects on environment of power reactor, 26:47471

(DOCKET-50305-45)
 effects on environment of power reactor, 26:47448
 (DOCKET-50277-52)
 effects on environment of power reactor, 26:47545
 (DOCKET-50387-19)
 effects on environment of power reactor, 26:47413
 (DOCKET-50250-84)
 effects on environment of power reactor, 26:47535
 (DOCKET-50373-24)
 effects on environment of power reactor, 26:47534
 (DOCKET-50369-28)
 use of US power plant, for heating small livestock shelters, proposal for, 26:45576
 uses of electric power plant, economic evaluation of, 26:45573 (AE-448)

HELIUM
 abundance variations in solar wind, dependence on velocity and proton flux, (E/T), 26:46328
 chemistry as coolant in HTGR, 26:45288 (TID-26106)
 diffusion in metals, activation energy and mechanisms of, 26:46050 (CONF-710601-, pp 230-47)
 effects of injected, on void formation in proton-irradiated Type 316 stainless steel, 26:46044 (AI-AEC-13033)
 electron density and radial distribution in, variational calculation of expectation values for, 26:46417
 electron reactions at 40 to 70 eV with, resonance in differential cross section for 2^3S excitation in, (E), 26:46405
 electron reactions at 500 eV and 1 keV with, angular and energy distributions of secondary electrons from direct ionization and autoionization in, (E), 26:46367
 electron reactions at zero angle with, second Born differential cross sections in high-energy limit for excitation of 2^3S and 2^3P states in, 26:46373
 electron reactions in eV range with, trapped-electron measurements of integral cross sections for production of secondary electrons of 0 to 1 eV by ionization in, 26:46390
 electron reactions with, structure in electron energy distribution, (E), 26:46407
 electron scattering by, calculation using method of effective potentials and variational formulations, 26:46400
 energy levels of diatomic, origin of energy barriers in ^1E excited states, (T), 26:46346
 energy levels of, coincidence measurement of lifetime of 4^1D level, 26:46345
 energy levels of, contributions from ^1P and ^3D intermediate states to second-order corrections to fine structure of 2^3P level, 26:46360
 energy levels of, relaxation processes in $n = 3$ states, (E), 26:46348
 energy levels of, Stark splitting in metastable 2^3S_1 state, (E), 26:46342
 energy levels of, variational calculation of upper and lower bounds to S and P excited states, 26:46413
 energy-level transitions in, electric-field effects on rate of, (E), 26:46342
 helium-atom elastic scattering at low energies by, retardation effects on differential and total cross sections for, (T), 26:46377
 hydrogen-ion (H_2^+ and H_3^+) reactions at 4 to 20 keV with, differential cross sections for formation of hydrogen $2s$ atoms by dissociation in, (E), 26:46388
 hydrogen-ion (H_2^+) reactions at 30 to 700 eV with, cross sections and polarization of optical emission arising from processes involving excitation, dissociation, and charge exchange in, (E), 26:46371
 interaction energy between two atoms of, at long range, calculation using Unsöld approximation, 26:46351
 interactions with helium and hydrogen atoms, retarded van der Waals potential for, as function of interatomic distance, 26:46357
 interatomic potentials for, calculation from total cross sections for elastic scattering, 26:46387
 lithium-ion reactions at 600 to 2800 eV with, oscillations of cross sections for charge exchange in, (E), 26:46368
 plasma of, electron energy distribution and striations in Langmuir paradox region of discharges, (E/T), 26:47133
 plasma of, far-infrared radiation from turbulent theta-pinch, (E), 26:47143
 plasma of, parallel magnetic multipole confinement of, (E), 26:47111
 positron scattering by, calculation using method of effective potentials and variational formulations, 26:46400
 proton reactions at 4 to 20 keV with, differential cross sections for formation of hydrogen $2s$ atoms by electron capture in, (E), 26:46388
 proton reactions at 4 to 20 keV with, differential cross sections for proton scattering and for formation of metastable and neutral hydrogen atoms by electron capture in, (E), 26:46397
 proton reactions at 75 to 350 keV with, absolute cross sections for formation of $3s$, $3p$, and $3d$ hydrogen atoms by charge exchange and excitation in, (E), 26:46378
 purity control in HTGR, 26:45288 (TID-26106)
 reactions between atoms of, in electric discharges, rates

for associative ionization, excitation transfer, and total neutral collisions in, (E), 26:46348
 reactions of 2^3S_1 atoms to produce ionization, structure in electron energy distribution, (E), 26:46407
 reactions with acetylene, argon, and nitrogen molecules at 50 to 600 eV of metastable atoms of, cross sections for de-excitation in, (E), 26:46394
 reactions with argon and nitrogen molecules at 150 to 1600 eV of metastable 2^1S and 2^3S atoms of, absolute cross sections for de-excitation in, (E), 26:46394
 reactions with argon at 10^{-3} to 10^4 eV of 2^3S and 2^1S atoms of, semi-empirical calculations of total cross sections for associative and Penning ionization in, 26:46395
 reactions with argon at thermal energies of metastable 2^1S and 2^3S atoms of, crossed-beam and gas-cell measurements of cross sections for ionization in, 26:46404
 scattering by deuterium molecules and hydrogen tritides, coupled-channel calculations of effects of molecular symmetry on cross sections for, 26:46379
 scattering by helium, total cross sections for elastic, (T), 26:46387
 spectral lines due to neutral, in planetary nebulae, (E), 26:46162
 stopping in matter at 0.1, 0.8, and 10 MeV, cross sections for electronic, as function of target atomic number, (T), 26:46365(R) (ORO-2911-14)
 x-ray scattering by ground-state, analytic calculation of form factor for coherent, 26:46359

HELIUM, SUPERFLUID
 current states in rotating, persistent, (T), 26:46801
 energy flux in between two baths of, separated by thin layer, (T), 26:46794
 energy transfer from pulsed thermal source to, below 0.3 K, 26:46797
 hydrodynamics of, consequences of microscopic symmetries and linear response theory for, 26:46796
 thermal capacity in, phonon dispersion and contribution to, 26:46795
 vortex ring dynamics and vortex core radius in, quantized, 26:46800

HELIUM IONS
 effects of 7-MeV/amu, on crystal structure of minerals, 26:46156
 energy levels of He^- , fine-structure intervals in metastable $1s2s2p^2P_{sub J}$ state, (E), 26:46343
 reactions with carbon foils at 40 keV, relative level populations excited by, as functions of quantum number and spin, (E), 26:46337
 reactions with cobalt, iron, tin, and zirconium at 5 MeV/amu, energy shift and relative intensity of K x rays from, (E), 26:46339
 reactions with hydrogen molecules at 30 to 700 eV, cross sections and polarization of optical emission arising from processes involving excitation, dissociation, and charge exchange in, (E), 26:46371
 reactions with hydrogen molecules below 700 eV, cross sections for emission and polarization of Lyman, Balmer, and He I lines in, (E), 26:46396
 spectral lines in O and Of stars due to, (E), 26:46180

HELIUM ISOTOPES He-3
 deuteron reactions (d, pn) at 27.5 MeV, analysis of $p-n$ final-state interaction in, (E), 26:46860 (LYCEN-7186)
 energy level at 14.5 MeV from deuterium (p, γ) reactions width of, (E), 26:46868
 impurities in liquid, microscopic theory of, 26:46803
 interactions $\gamma + ^3He + K^+ + A-^3He$, current algebra in soft kaon limit for differential cross sections and amplitudes of, (T), 26:46545
 meson reactions (π^-, π^+) ^3He with, analysis of nonresonant neutron attraction in, (T), 26:46862
 nuclear magnetic moment of, effects of two-pion contribution to proton electromagnetic structure on, (T), 26:46867
 reactions ($He-3, \alpha$) with carbon-13 at 29 to 42 MeV, excitation functions for, p. 31-2, (E), 26:46873(R) (CEA-N-1522)
 reactions ($He-3, \alpha$) with deuterium at 17.5 to 44.1 MeV, angular distributions for, (E), 26:46871
 reactions ($He-3, \alpha$) with tantalum-181, gold-197, and lead-208 up to 80 MeV, α spectra and cross sections for, (E), 26:47008
 reactions ($He-3, \alpha p$) with lithium-6 at 1.54 MeV, analysis of reaction mechanism in, (E), 26:46897
 reactions ($He-3, d$) with molybdenum-92 and -94 at 30.2 MeV, angular distribution and energy spectra from, p. 110-12, (E), 26:46873(R) (CEA-N-1522)
 reactions ($He-3, d$) with molybdenum-92 at 35 MeV, diffraction-model analysis of differential cross sections for, (T), 26:47075
 reactions ($He-3, d$) with neon-22 at 15 MeV, angular distributions for, (E), 26:46915
 reactions ($He-3, d$) with zinc-68 at 30.2 MeV, differential cross sections and spectrum from, p. 112-14, (E), 26:46873(R) (CEA-N-1522)
 reactions ($He-3, p$) with neon-21 at 18 MeV, angular distributions for, (E), 26:46915
 reactions ($He-3, t$) with argon-36 at 22.2 MeV, triton spectra and Q-value for, (E), 26:46904
 reactions ($He-3, t$) with titanium-46, chromium-50, iron-54, nickel-62, nickel-64, and zirconium-90 at 21.4 to 37.5 MeV with isobaric analog excitation, model

- analysis of, (E/T), 26:46930
 reactions with natural chromium, production of carrier-free manganese-54 from, 26:45394
 scattering by deuterium at 17.5 to 44.1 MeV, angular distributions for elastic, (E), 26:46871
 scattering by tantalum-181, gold-197, and lead-208 up to 80 MeV, energy spectra and cross sections for inelastic, (E), 26:47008
 spin-wave phenomena in liquid, 26:46802
 He-3--He-4, microscopic theory of impurities in liquid, 26:46803
 He-3--He-4, spin-wave phenomena in liquid systems, 26:46802
- HELIUM ISOTOPES He-4**
 deuteron reactions (d,pn) at 27.5 MeV, analysis of p-n final-state interaction in, (E), 26:46860 (LYCEN-7186)
 electrons on liquid, negative results for surface-bound, (E), 26:46798
 excitation spectrum of, absence of quadratic term in, (E/T), 26:46799
 interactions $\gamma + {}^4\text{He} + {}^4\text{He} + {}^4\text{He}$ at 2 to 5 GeV, proposed experiment for streamer chamber study of cross sections and π^0 lifetime in, (E), 26:46432(T) (SLAC-Trans-142)
 meson reactions ($\pi^+, 2p$) at 76 MeV, cross section for, 26:46870
 neutron inelastic scattering by solid, Fourier transform for, 26:46845 (BNL-16935)
 photon reactions (γ, n) and (γ, p) up to 34 MeV, analysis of charge-dependent effects using continuum shell model and coupled channel theory, (T), 26:46866
 photon reactions (γ, p^-) at 300 MeV, differential cross sections for, p. 182-3, (E), 26:46873(R) (CEA-N-1522)
 thermal capacity of, absorbed on bare copper, (E/T), 26:46804
 He-3--He-4, microscopic theory of impurities in liquid, 26:46803
 He-3--He-4, spin-wave phenomena in liquid systems, 26:46802
- HEMATOPOIESIS**
 radiation effects on, in newborn swine, effects of dose on gamma, 26:45852
- HEMATOPOIETIC SYSTEM**
 radiation effects on cells of, in bone marrow, lymph nodes, and spleen of newborn swine, effects of dose on gamma, 26:45853
- HIGH-ENERGY PHYSICS**
 lectures on, (T), 26:46509
- HIGH-TEMPERATURE SEPARATION PROCESSES**
 development of zone melting, in rotating container, 26:45276
- HIROSHIMA**
 muscular reflex time in residents of, age dependence of, 26:45795
- HISTAMINE**
 effects on respiratory exchange in rats, 26:45823
- HOLMIUM**
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 magnetic susceptibility of single crystals of, anisotropy in para-, (E), 26:47253
- HOLMIUM ALLOYS AND SYSTEMS**
 Fe--Ho, microstructure of intermetallic compounds of, 26:46016
- HOLMIUM ISOTOPES Ho-155**
 beta and electron-capture decay of, β , γ , and conversion electron spectra from, (E), 26:46986
- HOLMIUM ISOTOPES Ho-164**
 beta decay of, branching ratios, gamma and x-ray spectra, and half-life for, (E), 26:46998
 electron-capture decay of, branching ratios, gamma and x-ray spectra, and half-life for, (E), 26:46998
 energy-level transitions in, relative intensities of, (E), 26:46998
 isomeric transition in, half-life for, (E), 26:46998
- HOLMIUM ISOTOPES Ho-165**
 energy levels of, calculation using phenomenological model, (T), 26:47081
 neutron reactions (n, α) with, calculation of excitation functions for fast, 26:47066
 neutron strength functions at 10 to 1000 keV, S_0 and S_1 , p. 293-5, (T), 26:46873(R) (CEA-N-1522)
 nuclear hyperfine structure of, atomic-beam magnetic resonance measurements of, (E), 26:46989
 nuclear magnetic dipole moment and g factor for, atomic-beam magnetic resonance measurement of, (E), 26:46990
 photon reactions (γ, n) and ($\gamma, 3n$) at 90 MeV, relative cross sections for, (E), 26:46998
 photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
- HORMONES, ADRENAL**
 cortisone, effects on immune reactions to allogeneic tumor transplants in whole-body x-irradiated rats, 26:45820 (UR-3490-121)
 hydrocortisone, effects on incorporation of tritium-labeled-thymidine into rat liver DNA, 26:47646 (UR-3490-131)
 hydrocortisone, effects on plasma protein synthesis in isolated regenerating rat liver, 26:47646 (UR-3490-131)
- HORMONES, SALIVARY GLAND**
 parotin, effects on growth of bone tissues in irradiated rats, 26:45870
- HORMONES, SEX**
 androgen, radioinduced increase in production of, in fish, internal phosphorus-32 β , 26:45898
 dihydrotestosterone, radioimmunoassay in plasma of male and female subjects, 26:45928
 testosterone, radioimmunoassay in plasma of male and female subjects, 26:45928
- HOT CELLS**
 description of dual-mode operation, for alpha-active materials, 26:46107 (UCRL-73756)
 design and operation of, at Pakistan Institute of Nuclear Science and Technology, 26:45499
 design for transfer of alpha contaminated materials incorporating remote-controlled plastic bag sealer and cannister welder, 26:46108 (UCRL-73757)
 design modification of LASL plutonium, for nondestructive examination of irradiated fuels, 26:46104 (LA-DC-72-556)
- HUMBOLDT BAY POWER REACTOR**
 fuel assemblies for, design characteristics of Type 3 Batch 4, 26:47392 (DOCKET-50133-89)
 transmission line for, out of service condition for 115 kV, 26:47342 (DOCKET-50133-91)
- HUNGARY**
 well logging in, review on equipment and instruments for, 26:45560
- HUTCHINSON ISLAND POWER REACTOR, UNIT 1**
 circulating water system for, requested additional data on construction of intake and discharge pipelines for, 26:47506 (DOCKET-50335-32)
 environmental report for, cost-benefit data for, 26:47504 (DOCKET-50335-29)
 transmission line crossings of Indian River, alternatives to proposed, 26:47505 (DOCKET-50335-30)
- HYDRIODIC ACID**
 reactions of gaseous, with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
- HYDROBROMIC ACID**
 reactions of gaseous, with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
- HYDROCARBONS**
 radiolysis of liquid, theoretical study of electron scavenging process in pulse, 26:45333
- HYDROCHLORIC ACID**
 reactions of gaseous, with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
- HYDROFLUORIC ACID**
 positron lifetime in frozen aqueous solutions of, effects of temperature and positronium trapping on, (E), 26:46362 (RIS0-254)
 Hf--HNO₃, properties for etching germanium--silicon alloys, 26:45935 (SC-RR-710884)
- HYDROGEN**
 analysis of, allowance for background in mass spectrometric isotopic, 26:45241
 density and temperature of neutral, in interstellar dust clouds, (E), 26:46247
 deuteron elastic scattering at 27.5 MeV by polarized, spin correlation coefficients from vector-polarized, p. 142-5, (E), 26:46873(R) (CEA-N-1522)
 deuteron reactions (d,pn) at 27.5 MeV, analysis of p-n final-state interaction in, (E), 26:46860 (LYCEN-7186)
 electron scattering by molecular, close-coupling calculations of differential and total cross sections for, using square-well potentials, 26:46399
 energy-level transitions in presence of intense electromagnetic field, amplitudes of 1S--2P multiphoton, (T), 26:46356
 energy-level transitions in static and dynamic electric fields, resonance effects on shape of Lyman- α line from, 26:46361
 exchange with deuterium on aliphatic amines to produce heavy water, 26:45696(P)
 helium-ion reactions at 30 to 700 eV with molecular, cross sections and polarization of optical emission arising from processes involving excitation, dissociation, and charge exchange in, (E), 26:46371
 helium-ion reactions up to 700 eV with molecular, cross sections for emission and polarization of Lyman, Balmer, and He I lines in, (E), 26:46396
 hydrogen-ion (H₂⁺ and H₃⁺) reactions at 4 to 20 keV with molecular, differential cross sections for formation of hydrogen 2s atoms by dissociation in, (E), 26:46388
 hyperfine splitting of, structure functions from resonance model of inelastic e + p interactions for polarization contribution to ground-state, (T), 26:46523 (NYO-4076-17)
 hyperfine structure of ground-state, pressure shift in argon, krypton, and xenon, (E), 26:46347
 intensity ratio of H β /N₂⁺ 4709-A and vibrational enhancement of N₂⁺ first negative bands in proton-excited auroras, (E), 26:46276
 interaction energy between two atoms of, at long range, calculation using Unsöld approximation, 26:46351
 interactions between atoms of, calculation of higher multipole coefficients for, using Pade approximants, 26:46344

- interactions between atoms of, van der Waals constant for, (T), 26:46415
- interactions with helium and hydrogen atoms, retarded van der Waals potential for, as function of interatomic distance, 26:46357
- motion of, neutral, in spiral arms and local region, (E), 26:46246
- plasma of, electron density determination by laser Thomson scattering on discharge, (E), 26:47125
- plasma of, parallel magnetic multipole confinement of, (E), 26:47111
- plasma of, time-resolved diagnostic method for, 26:47127
- plasma of, validity of theory of dynamical linear Stark effect for interpretation of spectra of turbulent, 26:46361
- polarizability of atomic, variational bounds for, (T), 26:46415
- polarizability of molecular, molecular-beam magnetic-resonance measurement of anisotropy of static electric, 26:46358
- positron reactions with atomic, polarization effects in close-coupling calculations of elastic phase shifts and cross sections for inelastic scattering and positronium formation in, 26:46372
- positron scattering by, calculation using method of effective potentials and variational formulations, 26:46400
- production by water electrolysis for use in network power plants, evaluation of reactor systems for, 26:45429
- proton reactions at 75 to 350 keV with molecular, absolute cross sections for formation of 3s, 3p, and 3d hydrogen atoms by charge exchange and excitation in, (E), 26:46378
- proton scattering by atomic, coupled-state calculations of, in Reading representation, 26:46370
- reactions between atoms and molecules of, acoustic measurements of rotational energy transfer in, 26:46376
- reactions of atoms of, with aromatic carbonyl compounds, 26:45354
- reactions with alkali metal atoms, Born-approximation calculations of excitation to 2s, 2p, 3s, 3p, and 3d states in, 26:46403
- reactions with protactinium, x-ray powder diffraction studies on compounds produced by, 26:45992 (LBL-659)
- rotation--vibration properties of ground-state molecular, effects of polarization and screening on, (T), 26:46350
- separation from deuterium in formaldehyde by photopredissociation, 26:45307
- sorption by iron and steel during cathodic charging and chemical and electrochemical finishing, review on, 26:45986 (BARC-585)
- spectra of atomic, in rotating electric field, calculation of emission, 26:46352
- spectra of, review on Lamb shift, fine structure, and hyperfine structure in, 26:46710
- yields from γ radiolysis of ammonia--olefin mixtures, 26:45332
- yields from γ radiolysis of neutral and acidic ethanol, effects of pressure on scavengers effects on, 26:45351
- HYDROGEN IONS (H^+)**
- reactions with aluminum, Born-approximation calculations of energy losses and L-shell ionization in, 26:46393
- reactions with atomic and molecular gases at 75 to 350 keV, absolute cross sections for formation of 3s, 3p, and 3d hydrogen atoms by charge exchange and excitation in, (E), 26:46378
- reactions with atoms, classical calculations of cross sections for single-electron capture into ground and excited states in, 26:46398
- reactions with helium at 4 to 20 keV, differential cross sections for proton scattering and for formation of metastable and neutral hydrogen atoms by electron capture in, (E), 26:46397
- reactions with helium at 4 to 20 keV, differential cross sections for formation of hydrogen 2s atoms by electron capture in, (E), 26:46388
- reactions with RNase at 0.1 to 100 keV, cross sections for inactivation in, (T), 26:46365(R) (ORO-2911-14)
- scattering by argon and krypton, reduced cross sections for, (T), 26:46365(R) (ORO-2911-14)
- scattering by hydrogen atoms, coupled-state calculations of, in Reading representation, 26:46370
- HYDROGEN IONS (H_2^+)**
- diamagnetic susceptibility of, approximate variation--perturbation calculation of, 26:46353
- diamagnetic susceptibility of, effects of internuclear separation on, (T), 26:46354
- energy levels of, perturbation and variation--perturbation calculations of first-order wave functions and binding energies of, for internuclear separations of 0.2 to 10 a_0 , 26:46355
- nuclear magnetic shielding constant of, approximate variation--perturbation calculation of, 26:46353
- nuclear magnetic shielding constant of, effects of internuclear separation on, (T), 26:46354
- production for plasma injection experiments in cusped magnetic fields, 26:47153 (NP-19364)
- reactions with gases, formation of excited hydrogen atoms by dissociation in, (E), 26:46378
- reactions with helium and hydrogen molecules at 4 to 20 keV, differential cross sections for formation of hydrogen 2s atoms by dissociation in, (E), 26:46388
- reactions with helium at 30 to 700 eV, cross sections and polarization of optical emission arising from processes involving excitation, dissociation, and charge exchange in, (E), 26:46371
- HYDROGEN IONS (H_3^+)**
- production for plasma injection experiments in cusped magnetic fields, 26:47153 (NP-19364)
- reactions with gases, formation of excited hydrogen atoms by dissociation in, (E), 26:46378
- reactions with helium and hydrogen molecules at 4 to 20 keV, differential cross sections for formation of hydrogen 2s atoms by dissociation in, (E), 26:46388
- HYDROGEN ISOTOPES**
- determination of, allowance for background in mass spectrometric, 26:45241
- HYDROGEN PEROXIDES**
- photolysis of solid, ESR spectral studies of far-uv, 26:45323
- HYDROGEN SULFIDES**
- D_2S-H_2S , distillation of liquid, for production of deuterium, 26:45690(P)
- HYDROGEN SYSTEMS**
- H--V, structure of, 26:47263
- HYDROGEN TRITIDES**
- helium-atom scattering by, coupled-channel calculations of effects of molecular asymmetry on cross sections for, 26:46379
- HYDROLASES**
- radiation effects on activity of 5'nucleotidase in plasma membranes of rat liver, effects of glucagon and sodium fluoride on gamma, 26:45831
- radiation effects on deoxyribonuclease activity in rat liver, effects of cytostatic drugs on, 26:45732
- radiation effects on localization of acetylcholinesterase in neuromuscular junction and muscle of frogs, gamma, 26:45828
- radioinduced inhibition of glucose-6-phosphatase activity in erythrocytes of rabbits and rats following whole-body exposure, effects of cysteine and reduced glutathione on extent of recovery from x, 26:45848
- HYDROMAGNETIC WAVES**
- electron-plasma frequency fluctuations in oblique nonlinear, detection of solar-wind, 26:46329
- propagation in ionospheric F region, relation of spread F with Pc 1 micropulsations, Jan. 1, 1970 to June 30, 1971, (E/T), 26:46274
- HYDROXYL (HO)**
- emission in nightglow Nov. 1968--Oct. 1969, intensity ratio between (5,1) and (9,4) bands of, (E), 26:46286
- HYPERNUCLEI**
- decay of $A-^3H$, coupling constant of axial current for beta, (T), 26:46545
- production of $A-^3H$ in $\gamma + ^3He + K^+ + A-^3H$ interactions, current algebra in soft kaon limit for differential cross sections and amplitudes of, (T), 26:46545
- HYPERONS (Λ)**
- interactions $\Lambda + N$, two-pion exchange contribution to ΛN potential from amplitudes for, 26:46652
- production in $K + N \rightarrow \pi + \Lambda$ and $\pi + N \rightarrow K + \Lambda$ interactions at low energies, Veneziano duality for dips in cross sections for, (T), 26:46635
- HYPERONS ($\bar{\Lambda}$)**
- decay in bubble chamber, review of, (E), 26:46480
- production in $K^- + p$ interactions at 10 GeV/c in bubble chamber, review of, (E), 26:46480
- HYPOXIA**
- effects on metabolism of erythropoietin in whole-body irradiated rats, 26:45854
- ICE**
- photolysis of, ESR spectral studies of far-uv, 26:45323
- positron lifetime in, effects of temperature and positronium trapping on, (E), 26:46362 (RISO-254)
- radiolysis of doped single crystals of, role of fluoride ion in, 26:45345
- radiolysis of, optical absorption spectra of transient electrons from pulse, 26:45309(R) (AE-453)
- ICE-d₂**
- positron lifetime in, effects of temperature and positronium trapping on, (E), 26:46362 (RISO-254)
- radiolysis of, optical absorption spectra of transient electrons from pulse, 26:45309(R) (AE-453)
- ILLINOIS**
- cesium-137 content in human diets in Chicago, Jan. to June 1972, 26:45540 (HASL-257, pp III.45-52)
- IMAGE BRIGHTNESS INTENSIFIERS**
- development for neutron radiography, 26:45467 (BNWL-SA-4320)
- IMIDES**
- polymers, radioinduced photoconductivity in films of, x, (E/T), 26:45668 (UCRL-73974)
- IMMUNE REACTIONS**
- development in living cells against chemicals and radiation, 26:45845
- radioinduced suppression of, in rabbits to Salmonella typhimurium, protective effects of AET against γ ,

radioinduced suppression of, to allogeneic tumor transplants in rats, additive effects of cortisone on x, 26:45820 (UR-3490-121)

IN-PILE LOOPS

safety of CART, computer models for evaluation of, 26:47580 (CISE-R-259)

INCOLOY 800

corrosion cracking of, in oxygenated water at 288°C, stress, 26:45966 (BMI-1927)

INCONEL 600

compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)

corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)

corrosion cracking of, in oxygenated water at 288°C, stress, 26:45966 (BMI-1927)

INCONEL 625

corrosion cracking of, in oxygenated water at 288°C, stress, 26:45966 (BMI-1927)

INDEXES

KWIC index for numerical algebra, 26:47287 (ORNL-4778)

INDIA

nuclear energy in, developments in, 26:47300

INDIAN POINT POWER REACTOR, UNIT 1

control rod for, repair of stuck, 26:47320 (DOCKET-50003-118)

coolant for, excessive radioactivity in secondary, 26:47337 (DOCKET-50003-119)

emergency core cooling system for, performance evaluation of vertical induction pump motors for low pressure, 26:47388 (DOCKET-50003-120)

INDIAN POINT POWER REACTOR, UNIT 2

control rod drive line in, malfunction of, 26:47322 (DOCKET-50247-77)

valves for, installation and stress analysis for safety and pressure relief, 26:47412 (DOCKET-50247-76)

INDIAN POINT POWER REACTOR, UNIT 3

safety of, requested additional data on, 26:47456 (DOCKET-50286-47)

safety of, requested additional data on, 26:47458 (DOCKET-50286-49)

safety of, revised and requested additional data on, 26:47457 (DOCKET-50286-48)

safety of, revised and requested additional data on, 26:47455 (DOCKET-50286-46)

INDIUM

determination by fluorometry using Superchrome Garnet Y, 26:45236

determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218

indium-116 NMR in superconducting, (E), 26:46911

positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268

ultrasonic attenuation in superconducting, (E/T), 26:46816

INDIUM ALLOYS AND SYSTEMS

decomposition of supersaturated solid solutions of, effects of aging on mechanism of, 26:46032

Ag-base--In, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)

Bi--In, thermodynamic properties of intermetallic In_5Bi_3 , 26:46026

Cu-base--In, precipitation in, mechanisms of cellular, 26:46017

Hg--In--Sn, thermodynamic properties of amalgams in, at 150 to 340°C, 26:46004

In--La, nuclear magnetic resonance in LaIn_3 at 1.5 to 300°K, Knight shift and spin--lattice relaxation time, 26:47281

In--Sn-base, transitions of superconducting, resistive, (E/T), 26:46817

In-base--Sn, ultrasonic attenuation in superconducting, (E/T), 26:46816

INDIUM ISOTOPE IN-116

beta decay of polarized, asymmetry of, (E), 26:46911

nuclear magnetic resonance in indium antimonides and superconducting metal, (E), 26:46911

INDOLE, 3-(2-AMINOETHYL)-5-METHOXY-

radioprotective effects on mice, mixed γ and neutron, 26:45840

INDUSTRIAL WASTES

disposal of plastic-type, electron embrittlement--consolidation studies for, 26:46090

radiosterilization of liquid, design of cobalt-60 irradiation facility for, 26:45456(P)

radiosterilization of, description of spiral-flow gamma facility for, 26:45719(P)

INFECTIONS

effects of respiratory, on retention of chromium-51 in lungs of mice following intratracheal injection as oxide, 26:45910 (CONF-720505-2)

INFLAMMATIONS

dermatitis, radioinduced acute and chronic, in dogs following fractionated and single partial body exposure, effects of dose and fractionation interval on x, 26:45860

enteritis, radioinduced, in rats following abdominal exposure, effectiveness of cholestyramine resin for prevention and treatment of choleheic enteropathy resulting from x, 26:45821

glossitis, radioinduced, in rabbits, permeability of lingual mucosa during development and recovery from β , γ , and x, 26:45862

role of allergic, in pathogenesis of x radioinduced injuries in lungs of rabbits and rats, 26:45863

INFORMATION RETRIEVAL SYSTEMS

description of Canadian scientific and technical, 26:47652

INORGANIC COMPOUNDS

ionization energies of, photoelectron-spectroscopy study of chemical shifts in inner-shell, 26:46340 (ORNL-TM-3820)

preparation of high-purity, 26:45311(R) (BARC-599)

INSECTA

Acarus siro (grain mite), radiation effects on respiration rate in, effects of dose, life stage, sex, and temperature on γ , 26:45884

Dolichoderus quadripunctatus (ant), radiation effects on fecundity of female and worker, dose dependence of gamma, 26:45877

Drosophila melanogaster (fruit fly), radiation effects on frequency of chromosome loss in male germ cells of, enhancing effects of dose fractionation and lucanthone on x, 26:45887

Drosophila melanogaster (fruit fly), radioinduced chromosome translocations in male, effects on frequency of crossing over in chromosomes of F_1 females, x, 26:45876

Drosophila melanogaster (fruit fly), radioinduced mutations in, following ingestion of carbon-14, phosphorus-32, or phosphorus-33, beta, 26:45912(R) (ORD-3728-5)

Drosophila melanogaster (fruit fly), radioinduced non-disjunction in chromosomes of, method for quantitative study of dose effects on x, 26:45880

Drosophila sp. (fruit fly), radioinduced genetic recombination in, in relation to tergite development, x, 26:45881

Habrobracon sp. (wasp), radioinduced detrimental and lethal mutations in sperm of, effects on fecundity and life span of F_1 females, x, 26:45878

Heliothis zea (noctuid moth), radiation effects on visual sensitivity of, γ , 26:45873 (TID-26078)

Hemerocampa leucostigma (tussock moth), radiation effects on growth, development, and survival of larvae of, effects of dose on γ , 26:45885

Stomoxys calcitrans (stable fly), radiation effects on life span, mating competitiveness, and reproductive capacity of adult, following exposure of pupae, effects of dose and pupal age at exposure on γ , 26:45883

Tenebrio molitor (yellow mealworm), radioinduced injuries in midgut epithelium of adult and larval, dose and time dependence of γ , 26:45875

Tyroglyphus sp. (mite), radiation effects on survival of, infesting ham, electron, 26:45886

INTERNAL CONVERSION

spectra as function of atomic number from, Auger-electron and fluorescence-photon, (T), 26:47209

INTERSTELLAR MATERIAL

condensations of cooling, thermal, (T), 26:46245

light absorption by dust in, (E), 26:46171

molecule abundances in, (E/T), 26:46244

molecules in, complex organic, (T), 26:46240

INTESTINES

radioinduced rectovaginal fistula in human large, following combined external and intracavitary therapy of uterine carcinoma, surgical repair of γ , 26:45800

radioinduced stenosis of human small, following therapy of uterine carcinomas, symptoms and treatment of γ and x, 26:45802

INVARIANCE PRINCIPLES

charge conjugation invariance from asymmetry in $\eta \rightarrow \pi^+ \pi^-$ decay, (E), 26:46482

charge conjugation invariance from asymmetry in $\eta \rightarrow \pi^+ \pi^-$ decay, (E), 26:46483

conformal invariance of zero-mass Klein--Gordon-type equation, 26:46731

CP and T violation in weak interactions, 26:46584

CP violation test in charge asymmetry in $K_2^0 \rightarrow \pi^+ \pi^-$ decay in heavy-liquid bubble chamber, (E), 26:46494

CPT and Poincare invariance in determination of wave equations for arbitrary-spin particles, 26:46714

time reversal invariance in gamma transitions of oriented nuclear states, optimization of tests in, 26:46677

IODIDES

determination of nanogram quantities of, by isotope exchange, 26:45257

solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

IODINE

atom reactions with molecular, probabilities for vibrational transitions in, (T), 26:46375

chemistry in acidic and alkaline solutions, 26:45408(R) (ORNL-TM-2949)

chemistry in nitric acid solutions in fuel reprocessing, 26:45410(R) (ORNL-TM-3018)

chemistry in oxide fuel dissolution, 26:45407(R) (ORNL-TM-2918)

chemistry of uncommon volatile species of, 26:45415(R) (ORNL-TM-3217)

chemistry of uncommon volatile species of, 26:45409(R) (ORNL-TM-2996)

chemistry of uncommon volatile species of, 26:45414(R) (ORNL-TM-3180)

chemistry of uncommon volatile species of, 26:45416(R) (ORNL-TM-3250)

containment of fission product, in pyrochemical reprocessing of LMFBR fuels, 26:45421

distribution between aqueous and gaseous phases, 26:45413(R) (ORNL-TM-3127)

effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331

partition between aqueous and gaseous systems, 26:45415(R) (ORNL-TM-3217)

properties in nitric acid solutions, 26:45414(R) (ORNL-TM-3180)

reactions with liquid fins, 26:45421

release in fuel reprocessing cycle, 26:45412(R) (ORNL-TM-3095)

release in reprocessing of LMFBR fuels, 26:45406(R) (ORNL-TM-2871)

separation from air by cold trapping, 26:45416(R) (ORNL-TM-3250)

separation from aqueous solutions, 26:45417(R) (ORNL-TM-3281)

separation from aqueous solutions by sorption on silver iodide in presence of nonionogenic surface-active agents, 26:45279

separation from fuel reprocessing off-gas, 26:45411(R) (ORNL-TM-3072)

separation from fuel reprocessing off-gases, 26:45417(R) (ORNL-TM-3281)

separation from fuel reprocessing off-gases, 26:45409(R) (ORNL-TM-2996)

separation from fuel reprocessing off-gases by sorption on zeolites or nitrate mercury, 26:45407(R) (ORNL-TM-2918)

separation from gas streams by sorption on porous plastics or zeolites, 26:45412(R) (ORNL-TM-3095)

separation from gas streams in fuel reprocessing, 26:45413(R) (ORNL-TM-3127)

separation from gaseous streams by sorption and reaction on zeolites, 26:45387(P)

separation from gases, development of inorganic adsorbents for, 26:45405(R) (ICP-1006)

separation from off-gases in LMFBR fuel reprocessing, 26:45408(R) (ORNL-TM-2949)

separation of fission product, as methyl iodide from gas streams by sorption on styrene-divinylbenzene copolymers, 26:45388

sorption on porous plastic, 26:45414(R) (ORNL-TM-3180)

sorption on zeolites, 26:45410(R) (ORNL-TM-3018)

sorption on zeolites, 26:45409(R) (ORNL-TM-2996)

sorption on zeolites, 26:45411(R) (ORNL-TM-3072)

spectra in alkaline and nitrate solutions, 26:45417(R) (ORNL-TM-3281)

spectra in nitric acid solutions, 26:45413(R) (ORNL-TM-3127)

spectra in nitric acid solutions, 26:45415(R) (ORNL-TM-3217)

spectra in nitric acid solutions, 26:45416(R) (ORNL-TM-3250)

spectra of species of, in nitric acid solutions, 26:45412(R) (ORNL-TM-3095)

volatilization from nitric acid, 26:45415(R) (ORNL-TM-3217)

volatilization from nitric acid, 26:45411(R) (ORNL-TM-3072)

volatilization from nitric acid solutions, 26:45407(R) (ORNL-TM-2918)

volatilization from nitric acid solutions, 26:45417(R) (ORNL-TM-3281)

volatilization from nitric acid solutions, 26:45413(R) (ORNL-TM-3127)

volatilization from nitric acid solutions, 26:45416(R) (ORNL-TM-3250)

volatilization from spent oxide fuels in nitric acid dissolution, 26:45408(R) (ORNL-TM-2949)

IODINE ISOTOPES I-123

production of, evaluation of nuclear reactions for cyclotron, 26:45392 (N-72-15068)

properties as radiopharmaceutical, 26:45392 (N-72-15068)

IODINE ISOTOPES I-125

decay by electron capture and internal conversion, fluorescence-photon and Auger-electron spectra of ^{125}Te atoms from, (T), 26:47209

decay of, in relation to intracellular toxicity, 26:45893 (ORO-2690-12(Pt.B), Paper 13)

decay of, in relation to intracellular toxicity, 26:45892 (ORO-2690-12(Pt.B), Paper 4)

effectiveness for inducing breaks in DNA of eukaryote cells, 26:45894 (ORO-2690-12(Pt.B), Paper 14)

effects on cells following uptake, biophysical evaluation of toxic, 26:45893 (ORO-2690-12(Pt.B), Paper 13)

effects on cells of mice following injection as labeled leukemia cells, 26:45892 (ORO-2690-12(Pt.B), Paper 4)

purification of trace amounts of, by water vapor distillation in oxidizing atmosphere, 26:45309(R) (AE-453)

IODINE ISOTOPES I-128

chemical state of recoil, from neutron reactions (n, γ) with iodine-127 in gaseous iodine-butane mixture, 26:45361

chemical state of recoil, from neutron reactions (n, γ) with iodine-127 in mixture of iodine and butane in liquid phase, 26:45362

energy levels from neutron resonances in iodine-127, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

reactions of recoil, in neutron-irradiated iodopentacarbonylmanganese, 26:45357

IODINE ISOTOPES I-129

release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)

IODINE ISOTOPES I-131

content in milk from Nevada Test Site environs during July through Dec. 1971, 26:45515

effects on cells of mice following injection as labeled leukemia cells, 26:45892 (ORO-2690-12(Pt.B), Paper 4)

effects on growth of thyroid gland in goitrogen-stimulated adult and normal fetal mice, 26:45865

effects on mammals following uptake of, effects on pituitary and thyroid glands, review of carcinogenic, 26:45895 (UR-3490-102)

effects on thyroglobulin-induced stimulation of lymphocytes in man following uptake by thyroid gland, 26:45889

effects on thyroid gland of dogs, combined effects of x rays on neoplastic, 26:45819 (UR-3490-63)

release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)

removal from ventilation exhaust streams at Australian Atomic Energy Commission Research Establishment, performance of impregnated charcoal adsorption beds for, 26:45502

sorption by soils, effects of soil characteristics on, (E), 26:45528

uptake by human thyroid gland following administration, diagnostic value of determining rate of, 26:45906(T) (TT-54002/4, pp 462-70)

IODINE SYSTEMS

bromotrifluoromethane--I, radiolysis of, yields of products from, 26:45322

butane--I, neutron reactions (n, γ) with iodine-127 in gaseous mixtures of, chemical state of recoil iodine-128 from, 26:45361

butane--I, neutron reactions (n, γ) with liquid mixtures of, chemical state of recoil iodine-128 from, 26:45362

ION EXCHANGE

column packings for, gel, 26:45302(R) (COO-1222-55)

ION EXCHANGE MATERIALS

radiation effects on γ -type zeolites, proton ENDOR studies on γ , 26:45324

ION SOURCES

development for acceleration of multicharged ions of solid materials, 26:46133

IONOSPHERE

arc observations on August 8-9, 1970, satellite and ground-based SAR, (E), 26:46275

component distribution along slant column from observation point toward Sun, approximation to $\text{ch}(X/\text{sub } p, \text{chi}/\text{sub } p)$, (T), 26:46273

conjugate point phenomena in, review of, (E/T), 26:46323

coupling with protonosphere, relation with long-term variations of ionospheric electron content, (T), 26:46291

D region, correlation of hf absorption and 10 mb winds and temperature during IQSY, 26:46264 (COO-2195-5)

D region, electron photodetachment from major negative ions in, daytime, (E/T), 26:46299

D region, height-dependent electron density determination from one-hop ionospheric propagation of ordinary and quasi-transverse waves, (E), 26:46298

D region, photodetachment rates at sunrise, 26:46280

D region, time variations of negative ions and electron densities in, 26:46281

D region, time variations of ozone and atomic and molecular oxygen in, 26:46279

D region, vlf steep-incidence radiowave reflections from, rapid computation method of coefficients for, (E/T), 26:46263 (AD-734392)

E region, contribution of high-energy electron microbursts from daytime irregular geomagnetic pulsations to, 26:46317

E region, nonlinear mechanism for saturating Type-1 equatorial electrojet instability, (T), 26:46277

electron content at low and middle latitudes of, effects of solar and geomagnetic activity on 1965-1968, (E/T), 26:46287

electron temperature measurements in, by comparison method of differential currents to two identical probes, (E), 26:46295

electron-density irregularities during summer, post-midnight large-scale periodic, (E), 26:46290

F region, low-latitude OI 5577-A spectra of green-line tropical nightglow, (E), 26:46271

F region, neutral temperature at 200 to 320 km from 6300-A oxygen dayglow measurements, Sept. 1969 magnetic-storm-associated, (E), 26:46270

F region, small-scale fluctuations in critical frequency observed at Delhi, (E), 26:46289

F region, theoretical models of ionospheric storms,

(E/T), 26:46300
 F region, visible detection of rf heating of electrons from intensity of 6300-Å line of atomic oxygen nightglow, (E), 26:46292
 F2 layer, daily variation of equatorial anomaly in American and Asian sectors, (E/T), 26:46293
 F2 layer, effects of magnetic storm at Lima, Peru, March 8, 1970 on, 26:46282
 geomagnetic-field variations in polar cap due to critical geometric component of interplanetary magnetic field, (E), 26:46266
 ion density in, variations for large and small ions of time-dependent, (E/T), 26:46288
 oxygen O(¹D) quenching by molecular nitrogen using ionospheric modification experiment, 26:46278
 plasma experiments for, OGO-1 and OGO-3 satellite, 26:46265 (N-72-18715)
 solar-wind interactions with planetary, electrodynamic model for, (T), 26:46324 (N-72-18796)
 spread F, relation with Pc 1 micropulsations, Jan. 1, 1970 to June 30, 1971, (E/T), 26:46274
 wave losses in vlf duct associated with LHR frequency, collisional, (E/T), 26:46272
 whistler growth rate propagating obliquely in cold plasma with dilute energetic electron density, (E/T), 26:46314
 whistler side-band growth due to nonlinear wave--particle interaction, (E/T), 26:46315

IONS
 acceleration in solar atmosphere, effect of single- and multiple-ionization losses on, (T), 26:46230
 channeling in crystals, review of, 26:47228
 detection of heavy, in cosmic radiation, delta-ray theory for, 26:45612
 effects of accelerated heavy, on extracellular bacteriophages, lethal and mutagenic, 26:45746
 effects of energetic, on void formation in metals, models for, 26:46063 (CONF-710601-, pp 798-824)
 effects of energetic, on void nucleation in crystalline materials, 26:46064 (CONF-710601-, pp 825-40)
 effects on polyethylene sheets at 200 to 2000 eV, comparison with partial discharges, 26:46081
 effects on secondary emission and properties of solids, Russian book on, 26:47245
 electron pair correlation energies in, derivation from total correlation energies, (T), 26:46412
 energy levels of two-electron, variational calculation of upper and lower bounds to S and P excited states, 26:46413
 energy losses in matter, calculation of relative importance of electronic and nuclear processes in, (T), 26:46365(R) (ORO-2911-14)
 implantation in silicon, mass dependence of critical amorphizing dose in, (T), 26:47236
 photodetachment rates for negative, in ionospheric D region under sunrise conditions, 26:46280
 reactions with enzymes, binary encounter method for calculation of cross sections for inactivation in, 26:46365(R) (ORO-2911-14)
 reactions with nuclei, limits on angular momentum in fusion, (T), 26:47072
 scattering by germanium-74 and rhodium-103, time-dependent deorientation of recoils in vacuum from, (E), 26:46935
 stopping in matter, binary encounter method for calculation of cross sections for electronic, 26:46365(R) (ORO-2911-14)
 x-ray scattering by helium isoelectronic sequence of, analytic calculation of form factors for coherent, 26:46359

IRIDIUM
 determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
 determination of trace amounts of, in ores, concentrates, and sands by fire assay emission spectrography, 26:45229
 electric conductivity of pressed and fused, in range 100 to 1700°C, 26:47248
 ion implantation in, field-ion microscopic imaging of defects produced in, 26:47239
 ion implantation in, field-ion microscopic study of damage profiles produced in keV carbon, oxygen, iron, silver, and xenon, 26:47240
 microscopy of, imaging process in field ion, 26:46038
 properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)
 thermal conductivity of pressed and fused, in range 100 to 1700°C, 26:47248

IRIDIUM ALLOYS AND SYSTEMS
 Ir--Ni--Ta, phase studies on, 26:46010
 Ir-base--La, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248

IRIDIUM ISOTOPES Ir-179
 half-life of, (E), 26:46977

IRIDIUM ISOTOPES Ir-180
 half-life of, (E), 26:46977

IRIDIUM ISOTOPES Ir-181
 half-life of, (E), 26:46977

IRIDIUM ISOTOPES Ir-187
 electron-capture-decay of, branching ratios, gamma spectra, and log-ft values for, (E), 26:46984

IRIDIUM ISOTOPES Ir-192

availability and cost of radiation sources of, produced by Norway Institute for Atomic Energy, 26:45709 (INIS-mf-256)

IRON

abundance in primary cosmic radiation, (E), 26:46151
 abundance in solar-flare particles, (E), 26:46232
 analysis of high-purity, for manganese by neutron activation and preirradiation separation by ion exchange, 26:45216
 carbon-ion reactions at 10 MeV/amu with, energy shift and relative intensity of K x rays from, (E), 26:46339
 coatings for, structure of intermediate layers in diffusion, 26:46041
 determination in hair by neutron activation and γ spectroscopy, 26:45214
 determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
 effects on mechanical properties of molybdenum--uranium-base alloys at 20 to 640°C, 26:45995 (UJV-2802-M)
 effects on thermonuclear reactor breeding blanket performance, 26:47176
 ion reactions at 5 MeV/amu with, dependence of energy shift and relative intensity of K x rays from, on ion stopping power, (E), 26:46339
 muon (μ^+) range--momentum correlation in, 500- to 1500-MeV/c, (E), 26:46448 (LBL-791)
 muon (μ^-) range--momentum correlation in, 500- to 1500-MeV/c, (E), 26:46448 (LBL-791)
 neutron attenuation by, cross section for, (E), 26:46852
 neutron elastic scattering at 1.77 to 2.76 MeV, angular distributions for, (E), 26:46916 (AE-452)
 neutron total cross sections for, from 0.8 to 11 MeV, (E/T), 26:47220 (ORNL-TM-3867)
 neutron transmission measurements, (E), 26:46848
 radioinduced atomic displacement cascades in, statistical analysis of fast neutron, 26:46061 (CONF-710601-, pp 739-56)
 recrystallization of, effects of impurities on, 26:45982
 separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281
 separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
 sorptive properties for hydrogen during cathodic charging and chemical and electrochemical finishing, review on, 26:45986 (BARC-585)

IRON ALLOYS AND SYSTEMS
 Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009
 Al--Fe--Mo--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
 Al--Fe--Nb--Si, dissolution in, 26:46007
 Al--Fe--Si--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
 Al--Fe--U-base, mechanical properties of, at 20 to 640°C, effects of chromium and silicon additions on, 26:45995 (UJV-2802-M)
 Al--Fe--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
 Al--Fe--Y, phase studies on, 26:46005
 Al-base--Fe, superconductivity and localized spin fluctuations in, (E/T), 26:46838
 Co--Fe--W, crystal structure of, during aging, effects of molybdenum on, 26:46006
 Fe--Gd, magnetic compensation in films of, 26:47273
 Fe--Ho, microstructure of intermetallic compounds of, 26:46016
 Fe--Tb, magnetic compensation in films of, 26:47273

IRON CARBONYLS
 compaction of powdered, electrodynamic method for, 26:45951

IRON IONS
 implantation in iridium at keV energies, field-ion microscopic study of damage profiles produced in, 26:47240

IRON ISOTOPES Fe-54
 helium-3 reactions (He-3,t) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930
 neutron transmission measurements, (E), 26:46848
 oxygen-16 reactions (O-16, C-12) at 46 MeV, phase-shift analysis of differential cross sections for, p. 11-13, (E), 26:46873(R) (CEA-N-1522)
 oxygen-16 reactions (O-16, C-12) with, analysis using generator-coordinate method in DWBA framework, (T), 26:47069

IRON ISOTOPES Fe-55
 content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
 content in surface atmosphere at world sites, 1964 through Jan. 1970, 26:45512(R) (HASL-257(App.))

IRON ISOTOPES Fe-56
 alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932
 alpha reactions (α ,p) at 17 MeV, proton spectra from, (E), 26:46932
 energy-level density from alpha and proton reactions, (E), 26:46932
 neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909

proton inelastic scattering at 40 MeV, DWBA and coupled channel analysis of, (E), 26:46894

IRON ISOTOPES Fe-57
 Moessbauer effect in 3p-transition element matrix, background radiation in measurement of, 26:47282
 Moessbauer recoilless fraction and second-order Doppler shift of, pressure dependence of, (T), 26:47276
 Moessbauer spectra in β -rhombohedral boron, 26:47280

IRON ISOTOPES Fe-59
 distribution in tissues of newts, 26:45913(R) (RLO-2227-T-8-3)
 effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
 exchange between iron(II) and iron(III) in water--methyl sulfoxide systems, 26:45304

IRON NITRATES
 effects on corrosion of low-carbon 18/10 stainless steel by nitric acid, 26:45970

IRON OXIDES Fe₂O₃
 Al₂O₃--Fe₂O₃, phase studies on catalysts of, 26:45963

IRRADIATION FACILITIES
 atmosphere system for industrial, design of controlled, 26:45448(P)
 description of gamma, for industrial wastes and sewage, 26:45719(P)
 design and performance of, 26:45432
 design concepts for testing thermoelements at 2000°C, 26:45438 (KFK-1556)
 design for continuous irradiation of packaged materials, 26:45447(P)
 design for high initial intensity followed by low intensity irradiation, 26:45457(P)
 design of cobalt-60, for radiosterilization of liquid wastes, 26:45456(P)
 design of pool-type, with rotating radiation source, 26:45446(P)

ISOTOPE DILUTION ANALYSIS
 precision comparison for single and double, 26:45225

ISOTOPE SEPARATION METHODS
 centrifuge for, design of rotor for ultra-, 26:45697(P)
 development of equipment for, using supersonic gas jets, 26:45692(P)
 efficiency for uranium enrichment, report of Ad Hoc Committee on, 26:45689 (ORO-694)
 theory of Haeflner effect in electrotransport, 26:45694

ISOTOPES
 production using neutrons, design and performance of equipment for, 26:45432
 research progress in isotope chemistry, 26:47614(R) (COO-3498-2)

ITALY
 characteristics of Ispra site in, demographic, ecological, geologic, hydrologic, and meteorological, 26:45570 (EUR-4841)
 fallout deposition and content in atmosphere at Ispra, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
 fallout monitoring in atmosphere and milk at Ispra, 26:45511(R) (HASL-257)
 nuclear power plants for, site selection criteria for, 26:47574
 strontium-90 content in milk at Ispra, 1960 through 1971, 26:45519 (HASL-257, pp III.53-8)

J

JAMES A. FITZPATRICK POWER REACTOR
 environmental report for, requested additional data for, 26:47502 (DOCKET-50333-51)
 safety of, requested additional data on, 26:47501 (DOCKET-50333-50)

JAPAN
 nuclear power plants for, site selection criteria for, 26:47574

JAPANESE MATERIALS TESTING REACTOR
 analytical chemistry procedures at, 26:45182 (JAERI-M-4594)

JAWS
 radioinduced necrosis in human, following therapy of oral tumors, 26:45801

JETS
 use of supersonic gas, for separating heavy and light molecules from gaseous mixture, 26:45692(P)

JOSEPH M. FARLEY POWER REACTOR, UNIT 1
 construction of, deficiencies in inspection procedures for, 26:47519 (DOCKET-50348-42)

JOSEPH M. FARLEY POWER REACTOR, UNIT 2
 construction of, deficiencies in inspection procedures for, 26:47519 (DOCKET-50348-42)

K

KAOLINS
 kaolin--methyl methacrylate, radioinduced polymerization of composites of, γ , 26:45327

KEWAUNEE POWER REACTOR
 costs for, estimated operating and permanent shutdown, 26:47380 (DOCKET-50305-46)

effluents from, effects on environment of chemical, radioactive, and thermal, 26:47471 (DOCKET-50305-45)
 instrumentation for, installation of deficient controllers for Class 1, 26:47472 (DOCKET-50305-47)
 operation of, cost-benefit analysis for, 26:47471 (DOCKET-50305-45)
 safety of, revised and additional data on, 26:47470 (DOCKET-50305-44)
 site for, ecology, geology, hydrology, and meteorology of, 26:47471 (DOCKET-50305-45)

KIDNEYS
 radioinduced neoplasms in, following uptake of radioisotopes, review of pathology of, 26:45895 (UR-3490-102)

KRYPTON
 electron reactions at 1 keV with, angular and energy distributions of secondary electrons from direct ionization and autoionization in, (E), 26:46367
 proton scattering by, reduced cross sections for, (T), 26:46365(R) (ORO-2911-14)
 shipment of spent LMFBR, heat dissipation and shielding studies for, 26:45415(R) (ORNL-TM-3217)
 volatilization from oxide fuels, 26:45414(R) (ORNL-TM-3180)

KRYPTON ISOTOPES Kr-72
 nuclear properties of, compilation of, (E), 26:46939

KRYPTON ISOTOPES Kr-74
 energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo-SU₃ coupling schemes, 26:46931

KRYPTON ISOTOPES Kr-76
 energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo-SU₃ coupling schemes, 26:46931

KRYPTON ISOTOPES Kr-78
 energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo-SU₃ coupling schemes, 26:46931

KRYPTON ISOTOPES Kr-84
 reactions with thorium-232 at 450 to 494 Mev, excitation functions for production of thorium-225, thorium-226, actinium-224, and actinium-223 in, (E), 26:47035

KRYPTON ISOTOPES Kr-85
 release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)
 release from irradiated oxide fuels, 26:45414(R) (ORNL-TM-3180)
 release to atmosphere from nuclear facilities, need for controlling, 26:46102(T) (ORNL-tr-2369)

L

LA CROSSE BOILING-WATER REACTOR
 forced circulation pump for, fire in casing insulation of, 26:47382 (DOCKET-50409-6)
 fuel inventory for, 26:47570 (DOCKET-50409-5)
 operation of, flow anomaly in forced circulation system during, 26:47330 (DOCKET-50409-7)
 operations during June 1972, 26:47384 (DOCKET-50409-9)
 owners of, financial statements of, 26:47570 (DOCKET-50409-5)
 personnel for, revised organization of, 26:47569 (DOCKET-50409-4)
 radiation sources for, list of, 26:47570 (DOCKET-50409-5)
 shutdown condenser for, revised bolting requirements for partition for, 26:47382 (DOCKET-50409-6)
 steam condenser for, hydrogen detonation in off gas system of main, 26:47383 (DOCKET-50409-8)

LA SALLE COUNTY POWER REACTOR, UNIT 1
 effluents from, effects on environment of chemical, radioactive, and thermal, 26:47535 (DOCKET-50373-24)
 operation of, cost-benefit analysis for, 26:47535 (DOCKET-50373-24)
 safety of, revised and additional data on, 26:47536 (DOCKET-50373-25)
 site for, ecology, geology, hydrology, and meteorology of, 26:47535 (DOCKET-50373-24)

LA SALLE COUNTY POWER REACTOR, UNIT 2
 effluents from, effects on environment of chemical, radioactive, and thermal, 26:47535 (DOCKET-50373-24)
 operation of, cost-benefit analysis for, 26:47535 (DOCKET-50373-24)
 safety of, revised and additional data on, 26:47536 (DOCKET-50373-25)
 site for, ecology, geology, hydrology, and meteorology of, 26:47535 (DOCKET-50373-24)

LABORATORIES
 radiation accidents at University of Leeds, during 1958 to 1968, description of, 26:45791

LABORATORY EQUIPMENT
 design of denitrator, for conversion of uranyl nitrate to uranium trioxide, engineering drawings for, 26:45452

LAKE ERIE
 strontium content in waters of, concentration--time model for calculating, 26:45532

LAKE HURON
 conference on research on, Toronto, April 19 to 21, 1971, 26:45577
 strontium content in waters of, concentration--time model for calculating, 26:45532
 temperature structure of, effects of Candu reactor--

discharged thermal effluents on, 26:45581

LAKE MICHIGAN
 conference on research on, Toronto, April 19 to 21, 1971, 26:45577
 radioactivity monitoring in biota and surface waters of, near Big Rock Point power reactor during 1960 to 1969, gross beta, 26:45535
 radioisotope content of biota, sediments, and water in, near Big Rock Point power reactor during 1970, measurements of cesium-134, cesium-137, cobalt-60, and zinc-65, 26:45536
 strontium content in waters of, concentration--time model for calculating, 26:45532
 temperature distribution in waters of southwest, near power plant outfall, methods for measurement of, 26:45580

LAKE ONTARIO
 ecosystems in, effects of underwater-discharged nuclear power plant heated effluents on aquatic, (E), 26:45578
 strontium content in waters of, concentration--time model for calculating, 26:45532

LAKE SUPERIOR
 strontium content in waters of, concentration--time model for calculating, 26:45532

LAKES
 radioisotope content in Puget Sound basin, 26:45544 (RLD-2225-T-14-4)
 thermal effluent discharge into large, atmospheric effects of, 26:45584

LANTHANUM
 chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
 critical field behavior of superconducting, 26:46822
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 effects on grain boundary energy of nickel, 26:45999
 separation from other metals by combined ion exchange--solvent extraction chromatography, 26:45282
 separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
 solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

LANTHANUM ALLOYS AND SYSTEMS
 Al--Ca--La, critical field curves of Kondo superconducting, 26:46832
 Gd--La, critical-field and proximity-effect behavior of superconducting, many-body effects from, (E/T), 26:46822
 Ge--La, critical-field and proximity-effect behavior of superconducting, many-body effects from, (E/T), 26:46822
 In--La, nuclear magnetic resonance in LaIn_3 at 1.5 to 300°K, Knight shift and spin--lattice relaxation time, 26:47281
 Ir--base--La, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248
 La--Pb, nuclear magnetic resonance in LaPb_3 at 1.5 to 300°K, Knight shift and spin--lattice relaxation time, 26:47281
 La--Zn, phase studies and thermodynamic properties of, 26:46001

LANTHANUM ISOTOPES La-132
 beta and electron-capture decay of, γ and conversion electron spectra from, (E), 26:46948

LANTHANUM ISOTOPES La-133
 energy level at 535 keV, deformation parameter and quadrupole moment for, (E), 26:46960

LANTHANUM ISOTOPES La-138
 beta decay of, log-ft value for, (E), 26:46964
 electron-capture decay of, log-ft value for, (E), 26:46964
 energy levels from lanthanum-139 (d,t) reactions, spectroscopic factors and spin--parity assignments for, (E), 26:46964
 nuclear electric quadrupole moment of, from optical interference spectroscopy, (E), 26:46950
 nuclear spin and parity from lanthanum-139 (d,t) reactions, (E), 26:46964

LANTHANUM ISOTOPES La-139
 deuteron reactions (d,t) at 16 MeV, DWBA analysis of differential cross section and Q value for, (E), 26:46964
 energy-level transition from cerium-139 decay, internal conversion coefficient for 166-keV, (E), 26:46947
 nuclear magnetic resonance in LaIn_3 and LaPb_3 , 26:47281

LANTHANUM ISOTOPES La-140
 beta decay of, γ - γ angular correlations for, (E), 26:46958

LANTHANUM OXIDES
 CaO--La₂O₃, thermodynamics of solid electrolytes of, review on, 26:46013

LANTHANUM PEROXIDES
 preparation and properties of, 26:46085

LARVAE
 radiation effects on tussock moth, dose dependence of growth, development, and survival, γ , 26:45885

LARYNX
 radioinduced mucosal edema of human, following therapy of laryngeal carcinomas, incidence of x, 26:45781(T) (TT-54002/4, pp 437-41)

LASER RADIATION
 effects of beams of, on void concentration in nickel and

101 SUBJECTS

LEAD ISOTOPES Pb-208

vanadium foils, 26:46059 (CONF-710601-, pp 613-29)
 frequency of 633-nm line of, (E), 26:46733
 reactions with solid targets, ion energy spectra from, (E), 26:47154
 reactions with solid targets, momentum transfer in, 26:47155
 scattering on hydrogen discharges, electron density determination from Thomson, (E), 26:47125
 scattering on Tokamak plasma, magnetic field measurement from, 26:47120

LASERS
 design of compact solid-state, using decaying radioisotope as internal power supply, 26:45677(P)
 performance of nuclear-fueled, on earth satellites for energy transfer to lunar stations and space vehicles, 26:45428
 radiation effects on CO₂, alpha, 26:45669
 radiation effects on stimulated radiation properties of ruby, γ -, 26:47227
 use of pulsed gas, as high-intensity pulsed light sources for testing photomultiplier tubes, 26:45640

LEAD
 content in surface atmosphere at world sites, Oct. 1966 through Nov. 1971, 26:45512(R) (HASL-257(App.))
 determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
 determination in soils by photon activation methods, 26:45198
 determination of, chemical procedures for, 26:45181 (CISE-R-172)
 mass flow choking limit for heat pipe working fluid of, (E), 26:45460
 neutron elastic scattering at 8.05 MeV in natural and radiogenic, angular distributions for, (E), 26:46916 (AC-452)
 positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
 proton reactions at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

LEAD ALLOYS AND SYSTEMS
 La--Pb, nuclear magnetic resonance in LaPb_3 at 1.5 to 300°K, Knight shift and spin--lattice relaxation time, 26:47281
 Pb--Sn, photoelectric process in various shells in, effective atomic numbers in, (T), 26:47206

LEAD ISOTOPES
 photon reactions (γ , π^0) at 3.5 to 8.3 GeV, A dependence of, 26:46447
 photon reactions at 1.160 GeV at 150°, photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441
 x-ray spectra from, isotope shifts in, (E), 26:46973

LEAD ISOTOPES Pb-198
 energy levels from mercury (α ,xn) reactions, microscopic analysis of, (E/T), 26:47009

LEAD ISOTOPES Pb-200
 energy levels from mercury (α ,xn) reactions, microscopic analysis of, (E/T), 26:47009

LEAD ISOTOPES Pb-202
 energy levels from mercury (α ,xn) reactions, microscopic analysis of, (E/T), 26:47009

LEAD ISOTOPES Pb-204
 energy levels from mercury (α ,xn) reactions, microscopic analysis of, (E/T), 26:47009

LEAD ISOTOPES Pb-206
 energy level of, g-factor of 12⁺ isomeric, (E), 26:47002
 energy levels from mercury (α ,xn) reactions, microscopic analysis of, (E/T), 26:47009

LEAD ISOTOPES Pb-207
 energy level at 897 keV, Doppler-shift-attenuation measurement of lifetime of, p. 53-7, (E), 26:46873(R) (CEA-N-1522)
 nuclear magnetic resonance in LaPb_3 , 26:47281

LEAD ISOTOPES Pb-208
 alpha inelastic scattering up to 80 MeV, α spectra and cross section for, (E), 26:47008
 alpha reactions (α ,He-3) up to 80 MeV, energy spectra and cross section for, (E), 26:47008
 charge states of, formed by alpha decay of polonium-212, 26:45359
 electron elastic scattering at 400 and 600 MeV, phase-shift analysis of form factor for, p. 192-4, (E), 26:46873(R) (CEA-N-1522)
 energy levels of, configuration strengths of low-lying particle-hole, p. 38-41, (E), 26:46873(R) (CEA-N-1522)
 helium-3 inelastic scattering up to 80 MeV, energy spectra and cross section for, (E), 26:47008
 helium-3 reactions (He-3, α) up to 80 MeV, α spectra and cross section for, (E), 26:47008
 lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
 neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909
 photon reactions (γ ,n) and (γ ,p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
 photon reactions at 7.170 MeV, gamma spectrum from, p. 313-4, (E), 26:46873(R) (CEA-N-1522)
 proton reactions (p ,n) at 20 to 48 MeV with subsequent

proton decay, coupled channels calculations of energy dependence of, (T), 26:47073

LEAD ISOTOPES Pb-209
beta decay of, half-life, Kurie plot, log-ft value, and shape factor for, (E), 26:47007

LEAD ISOTOPES Pb-210
content in organisms, sediment, and water in freshwater lake and salt water sound in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
content in plants and soils in uraniferous area in New Zealand, 26:45529
retention in beagle dogs following intravenous injection, 26:45916 (UR-3490-159)

LEAD URANIUM SELENIDES
preparation and structure of PbU_2Se_8 , 26:45378

LEAD URANIUM SULFIDES
preparation and structure of PbU_2S_8 , 26:45378

LEAVES
analysis for cesium by neutron activation, 26:45192 (TID-25954, pp 373-5)

LEPTONS
conservation law for, (T), 26:46578
coupling of massless composite lepton-antilepton states to free, (T), 26:46570 (IC-72/54)
interactions anti $l + l$, consequences, spin, and coupling to free leptons for massless composite lepton-antilepton states from exchange of virtual neutrino pairs in, (T), 26:46570 (IC-72/54)
interactions $\gamma + l^-$ in Weinberg model, Drell-Hearn sum rule for Compton, (T), 26:46553
interactions $l + N$ using polarized proton and deuteron targets, review of, (E), 26:46436
interactions $l + N$, parton models with integral charge for deep-inelastic, 26:46574
interactions $\nu + l$, lepton charge theory for, (T), 26:46578
interactions with hadrons, relation between fixed poles in virtual Compton scattering and scale invariance in, 26:46559
magnetic moment of, second-order weak corrections to anomalous, (T), 26:46553
magnetic moments of charged, quantum electrodynamic contribution to anomalous, 26:46710
production of massless composite lepton-antilepton states from exchange of virtual neutrino pairs in anti $l + l$ interactions, (T), 26:46570 (IC-72/54)
spin of massless composite lepton-antilepton states in exchange of virtual neutrino pairs in anti $l + l$ interactions, (T), 26:46570 (IC-72/54)

LEUKOCYTES
radiation effects on phagocytic activity of macrophages co-cultured with tumor cells, x, 26:45738
radiation effects on response of mononuclear phagocytic, to intraperitoneal inflammation in mice, effects of time of exposure on x, 26:45833
radioinduced reduction of count of, in man following therapy of gynecologic neoplasms, protective effects of cepharanthine against, 26:45805

LEVEL INDICATORS
description of radiometric, for liquid metals heated by electron beam in crucible, 26:45655(P)

LICENSING AND REGULATION
procedures for U.S. nuclear power plants, 26:47571

LIGASES
role of DNA ligase in repair of uv-radioinduced damage in DNA in vitro, 26:45727

LIGHT
interactions with matter, review of, (T), 26:46508
velocity of, precision determination of, (E), 26:46733

LIGHT SOURCES
relativistic beaming from rotating, (T), 26:46201

LIMERICK POWER REACTOR, UNIT 1
environmental report on, requested additional data on, 26:47520 (DOCKET-50352-27)

LIMERICK POWER REACTOR, UNIT 2
environmental report on, requested additional data on, 26:47520 (DOCKET-50352-27)

LINEAR ACCELERATORS
beam extraction system for Kharkov 300-MeV electron, principles of operation of, 26:46131
beam handling system for CEA 600-MeV, description of, p. 178-81, 202-6, 26:46873(R) (CEA-N-1522)
beam stability in induction, solution of kinetic equation for longitudinal, 26:46138
description and operation of CEA 60-MeV, p. 257-61, 26:46873(R) (CEA-N-1522)
electron source for, secondary emission multipactoring 20-mA, 26:46134
experimental facilities at CEA 600-MeV, description of computer-based, p. 242-8, 26:46873(R) (CEA-N-1522)
experimental facilities at CEA 600-MeV, description of muon, p. 219-21, 26:46873(R) (CEA-N-1522)
experimental facilities at CEA 600-MeV, design of spectrometer, p. 206-10, 26:46873(R) (CEA-N-1522)
experimental facility at electron, description of bremsstrahlung, 26:46145
operation of CEA 60-MeV, description of proposed on-line, p. 263-4, 26:46873(R) (CEA-N-1522)
operation of CEA 600-MeV, Oct. 1, 1970-Oct. 15, 1971, p. 171-81, 26:46873(R) (CEA-N-1522)

performance of traveling-wave, Q-value effects of accelerating structure on, 26:46122
radiofrequency control system for, with automatic compensation for temperature changes in accelerating structures, 26:46127

LINEAR ENERGY TRANSFER
problems in use of, 26:45611

LIPIDS
radiation effects on peroxidation of, in tissues of mice following whole-body exposure, effects of dietary antioxidants on x, 26:45872

LIQUID JETS
flow characteristics of rectangular, horizontally entering constant depth basin, (E/T), 26:45582

LIQUID METAL FAST BREEDER REACTOR
cost of, effects of development delays on, 26:47316
fuel cycle development studies (April 1970) on, 26:45409(R) (ORNL-TM-2996)
fuel cycle development studies (August 1970) on, 26:45413(R) (ORNL-TM-3127)
fuel cycle development studies (Dec. 1970) on, 26:45417(R) (ORNL-TM-3281)
fuel cycle development studies (Feb. 1970) on, 26:45407(R) (ORNL-TM-2918)
fuel cycle development studies (July 1970) on, 26:45412(R) (ORNL-TM-3095)
fuel cycle development studies (June 1970) on, 26:45411(R) (ORNL-TM-3072)
fuel cycle development studies (March 1970) on, 26:45408(R) (ORNL-TM-2949)
fuel cycle development studies (May 1970) on, 26:45410(R) (ORNL-TM-3018)
fuel cycle development studies (Nov. 1970) on, 26:45416(R) (ORNL-TM-3250)
fuel cycle development studies (Oct. 1970) on, 26:45415(R) (ORNL-TM-3217)
fuel cycle development studies (Sept. 1970) on, 26:45414(R) (ORNL-TM-3180)
fuel cycle studies (Jan 1970) for, 26:45406(R) (ORNL-TM-2871)
fuel reprocessing for, containment of fission product iodine in pyrochemical, 26:45421
fuel reprocessing for, dissolution following shearing in, 26:45397 (CONF-720903-9)
fuels for, development of plutonium nitride-uranium nitride, 26:45960

LIQUIDS
interfaces of, design of gamma radiometric device for locating, 26:45659(P)
positron annihilation in crystals of, positronium-phase and temperature dependence of lifetimes in, (E), 26:46363

LITHIUM
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

LITHIUM CHLORIDES
effects on rats, relation to development of taste aversions, 26:45817 (ORO-2690-12(Pt.B), Paper 12)

LITHIUM DEUTERIDES
plasma of, ion energy spectra from laser-produced, (E), 26:47154

LITHIUM IONS
implantation in nickel at 1 to 10 keV, penetration and spatial distribution from, 26:47235
reactions with carbon foils at 40 keV, relative level populations excited by, as functions of quantum number and spin, (E), 26:46337
reactions with helium at 600 to 2800 eV, oscillations of cross sections for charge exchange in, (E), 26:46368

LITHIUM ISOTOPES Li-6
alpha elastic scattering at 36.6 MeV, angular distribution for, (E), 26:46883
energy level at 3.56 MeV from beryllium-9 (p, α) reactions, search for parity and isospin violating decay of, (E), 26:46900
helium-3 reactions ($He-3, \alpha p$) at 1.54 MeV, analysis of reaction mechanism in, (E), 26:46897
meson reactions ($w^+, 2p$) at 76 MeV, cross section for, 26:46870
meson reactions ($w^-, w^- p$) at 1 GeV/c, w -meson momenta from, (E), 26:46869
neutron cross sections for, review of ENDF/B III files for, 26:46876 (ORNL-TM-3859)
nuclear charge form factor and quadrupole moment of, cluster model calculation of, (T), 26:46878
proton reactions ($p, He-3$) up to 125 MeV (cms), calculation of differential cross sections using unified microscopic theory, (T), 26:47077
reactions ($Li-6, d$) with fluorine-19 at 16 MeV, angular distributions for, (E), 26:46915
scattering by $A = 12$ to 208 nuclei at 30 MeV, angular distributions for elastic, (E), 26:46883
scattering by boron-11 and carbon-13 at 28 MeV, optical-model analysis of angular distributions from elastic, p. 26-30, (E), 26:46873(R) (CEA-N-1522)

LITHIUM ISOTOPES Li-7
neutron cross sections for, review of ENDF/B III files for, 26:46876 (ORNL-TM-3859)
reactions ($Li-7, \alpha$) with neon-20 at 22 MeV, angular distributions for, (E), 26:46915
reactions with gold-197, mechanism for breakup, p. 24-6,

(E), 26:46873(R) (CEA-N-1522)

LITHIUM MOLYBDATE
 Li_2MoO_4 -- $\text{Pu}(\text{MoO}_4)_2$, phase diagram of, 26:45954
 Li_2MoO_4 -- $\text{Pu}(\text{MoO}_4)_2$, scheelite structure of phases in, 26:45954

LIVER
radiation effects on activity of glutathione peroxidase in, of swine, x, 26:45733
radiation effects on deoxyribonuclease activity in rat, effects of cytostatic drugs on, 26:45732
radioinduced injuries to human, following therapy of adjacent organs, dose and time dependence of γ , 26:45809
radioinduced neoplasms in, following uptake of radioisotopes, review of pathology of, 26:45895 (UR-3490-102)

LIVERMORE POOL-TYPE REACTOR
operation and repair of, Sept.--Dec. 1971, pp.14-16, 26:45673(R) (UCRL-50007-71-3)

LOW TEMPERATURE PHYSICS
book: Introduction to Low Temperature Physics (In Romanian), 26:46841

LUCANTHONE
radiosensitizing effects of, on male germ cells of *Drosophila*, x, 26:45887

LUNAR MATERIALS
fission track studies in, 26:46156
radiation effects on, search for, 26:46156

LUNGS
chromium-51 retention in, of mice following intratracheal injection as oxide, effects of respiratory infection on effective half-life for, 26:45910 (CONF-720505-2)
clearance of metastable silver-110 from, of dogs following inhalation, 26:45915 (UR-3490-115)
radioinduced injuries in, of rabbits and rats, role of allergic inflammations in pathogenesis of x, 26:45863

LUTETIUM
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination of, chemical procedures for, 26:45181 (CISE-R-172)

LUTETIUM BOROCARBIDES
preparation, properties, and structure of, 26:45964

LUTETIUM CARBIDES
vaporization of solid LuC_2 , thermodynamics of, 26:45961

LUTETIUM ISOTOPES Lu-171
energy levels from thulium-169 ($\alpha, 2n$) reactions, comparison with rotor model predictions, (E/T), 26:46995
energy levels from thulium-169 ($\alpha, 2n$) reactions, high-spin rotational, (E), 26:46980
energy levels from thulium-169 ($\alpha, 2n\gamma$) reactions, properties of rotational, (E), 26:46987
energy levels of, Coriolis coupling in rotational-band, (T), 26:46988

LUTETIUM ISOTOPES Lu-175
energy-level transitions in, nuclear part of parity nonconserving circular polarization of 343- and 396-keV γ -rays from, (T), 26:46982

LUTETIUM PEROXIDES
preparation and properties of, 26:46085

LYMPHOCYTES
radiation effects on potassium ion transport in rat thymic, following exposure in vitro, x, 26:45744
radiation effects on thyroglobulin-induced stimulation of, in man following uptake of iodine-131 by thyroid gland, β and γ , 26:45889

LYSOSOMES
radiation effects on function and structure of, in rat liver cells following lethal whole-body exposure, x, 26:45839

M

MC GUIRE POWER REACTOR, UNIT 1
construction of, cost data for, 26:47533 (DOCKET-50369-26)
effluents from, effects on environment of chemical, radioactive, and thermal, 26:47534 (DOCKET-50369-28)
operation of, cost-benefit analysis for, 26:47534 (DOCKET-50369-28)
site for, ecology, geology, hydrology, and meteorology of, 26:47534 (DOCKET-50369-28)

MC GUIRE POWER REACTOR, UNIT 2
construction of, cost data for, 26:47533 (DOCKET-50369-26)
effluents from, effects on environment of chemical, radioactive, and thermal, 26:47534 (DOCKET-50369-28)
operation of, cost-benefit analysis for, 26:47534 (DOCKET-50369-28)
site for, ecology, geology, hydrology, and meteorology of, 26:47534 (DOCKET-50369-28)

MACHINE TOOLS
wear of steel, post-cutting chip activation analysis method for measuring, 26:45713

MACHINING
method for deburring using thermal shock, 26:47609 (BDX-613-642(Rev.))

MAGNESIUM
determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
properties of extruded and recrystallized rods and tubes

of, for use as fuel element cladding, anisotropic, 26:46012
prospecting for deposits of, pulsed neutron well logging method for, 26:45559(P)
solution numbers from compressibilities and ionic vibration potentials measurements, 26:45369

MAGNESIUM ALLOYS AND SYSTEMS
Al--Mg, grain refining of, mechanism and method for, 26:45984
Cu--Mg, dislocation energy of impurity atoms in solid solutions of, internal friction method for studying, 26:46023
Mg--Zn, grain refining of, mechanism and method for, 26:45984
Mg--Zr, grain refining of, mechanism and method for, 26:45984

MAGNESIUM ISOTOPES Mg-24
energy levels from sodium-23 (p, γ) reactions, widths and spin-parity assignments for, (E), 26:46903(R) (COD-1120-149)
nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892
oxygen-16 reactions (O-16, C-12) at 42 MeV, differential cross sections and spectrum from, (E), 26:46913
photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
proton reactions (p, γ) at 200 to 1700 keV, gamma angular distributions and spectra and Q value for, (E), 26:46910
triton reactions (t, α) at 15 MeV, angular distributions for, (E), 26:46915

MAGNESIUM ISOTOPES Mg-25
deuteron reactions (d, α) at 11.57, 11.82, and 12.07 MeV, angular distributions for, (E), 26:46915
energy level at 975 keV, lifetime of, (E), 26:46908

MAGNESIUM ISOTOPES Mg-26
oxygen-16 reactions (O-16, α) at 62 MeV, spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
oxygen-16 reactions (O-16, C-12) at 42 MeV, peak cross sections for, (E), 26:46913
photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

MAGNESIUM OXIDES
properties for use in magnetohydrodynamic generators, 26:45426(P)
 Al_2O_3 -- CaO -- Cr_2O_3 -- F -- MgO -- MoO_3 -- TiO_2 -- ZrO_2 , use as welding flux, 26:45941(P)
 Al_2O_3 --MgO, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
CoO--MgO, corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
MgO-- ZrO_2 , phase studies on partially stabilized, 26:45945
MgO-- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

MAGNET COILS
charging or discharging superconducting, method and means of, 26:46842(P)

MAGNETIC FIELDS
charged-particle motion in two-dimensional neutral sheet with linear variation, Hamiltonian methods for oscillation frequency and analytic treatment of perturbed, (T), 26:46522 (N-72-17763)
design and production of antisymmetric 1/r, with periodic azimuthal symmetry, 26:45638
effects of critical geometric component of interplanetary, on large geomagnetic effects in solar cap, (E), 26:46266
effects of external wave, on electron polarization in magnetic field, (T), 26:46551
effects of transverse, on heat transfer from partially ionized argon, 26:45464
effects on synchronous rotation of multiple rotors, 26:45441 (SRARI-P-120)
electron at $n = 1$ level in, induced emission from, (T), 26:46551
electron polarization in, external magnetic field wave effects on, (T), 26:46551
electron ultra-luminous motion in medium in, spin effects on synchrotron radiation intensity and lifetime of ground and first excited states for, (T), 26:46548
gravitational collapse in, exact solutions of Einstein equations for gravitational fields with sources of dust and magnetic field for, (T), 26:46755
heat conduction in turbulent, effect of long wavelength, nonresonant magnetic-field fluctuations on, (T), 26:46326
neutron motion through medium in, coherent generation mechanism for induced electromagnetic waves from, (T), 26:46541
observations from HEOS-1 launched Dec. 5, 1968, (E), 26:46334

MAGNETIC MIRROR MACHINES
plasma confinement in parallel magnetic multipoles, (E), 26:47111
plasma confinement in, effect of nonadiabatic energy limit versus mirror ratio on, (E/T), 26:47106

MAGNETIC MONOPOLES

existence of, formulation of Einstein--Kaufman nonsymmetric unified field theory for, (T), 26:46694

MAGNETIC PROBES

optimization in transient plasmas, 26:47124

MAGNETIC RESONANCE

model for powder ENDOR, 26:47279

MAGNETIC STORMS

effect of Sept. 1969 on F-region neutral temperature at 200 to 320 km from 6300-A oxygen dayglow measurements, (E), 26:46270

electron pitch-angle distribution during substorms, relation of magnetotail structure with anisotropic, (E/T), 26:46306

growth phase of magnetospheric substorms, changes in magnetosphere configuration during, (E), 26:46305
observation at Lima, Peru, March 8, 1970, 26:46282
occurrence on Sept. 2, 1966, electron repopulation in inner radiation belts before and after, (E/T), 26:46310

radio-wave emission during post-breakup phase of polar substorms, fine structure of vlf, (E), 26:46285
relation of polar and magnetospheric substorms with Pld irregular pulsations in daytime auroral zone, (E), 26:46317

relation with solar-flare-associated interplanetary streams in solar wind, (E/T), 26:46332

MAGNETIC TESTING

method for welds in steel chemical munition containers, 26:45470

MAGNETOHYDRODYNAMIC GENERATORS

containment walls for, development of materials for, 26:45426(P)

economics of, for use with fusion reactors, 26:47170 (IPP-IV-32)
electric parameters of Hall-type, 26:45431

energy conversion in, analysis of optimum conditions for, 26:45430

MAGNETOHYDRODYNAMICS

equations for, two-dimensional nonsteady-state solution of, (T), 26:46425

MAGNETOSHEATH

plasma flow observations adjacent to bow shock at 20 to 60 R/sub E/, July to Oct. 1967 and Dec. 1967 to April 1968, (E), 26:46303

MAGNETOSPHERE

component distribution along slant column from observation point toward Sun, approximation to $\chi(X/\text{sub } p)$, $\chi(h/\text{sub } p)$, (T), 26:46273

conjugate point phenomena in, review of, (E/T), 26:46323
coupling between ionosphere and protonosphere, relation with long-term variations of ionospheric electron content, (T), 26:46291

electron density profile determination from plasmapause whistlers, ring-current effects on, (T), 26:46284

electron pitch-angle diffusion within plasmasphere, association with plasmaspheric whistler-mode waves, (E/T), 26:46311

electron spikes at high latitudes in, equal to or greater than 425 keV, (E), 26:46307

electrostatic electron cyclotron harmonic wave turbulence in, vlf nonlinear, (E), 26:46313

emission of intense bands of elf noise in association with low-energy auroral electron precipitation at low altitudes, (E), 26:46308

geomagnetic-field variations in polar cap due to critical geometric component of interplanetary magnetic field, (E), 26:46266

interaction with solar wind, magnetosphere compression variations and reflection and refraction of MHD waves at magnetosphere boundary during, (T), 26:46321

magnetic substorms in, relation with Pld irregular pulsations in daytime auroral zone, (E), 26:46317

magnetohydrodynamic effects from wave reflection and refraction at boundary of, (T), 26:46321

magnetotail configuration variations during growth phase of magnetospheric substorm, (E/T), 26:46305

observations from HEOS-1 launched Dec. 5, 1968, (E), 26:46334

plasmasphere formation and maintenance from light-ion trough characteristics in plasmapause, (E), 26:46301 (N-72-18362)

wave losses in vlf duct associated with LHR frequency, collisional, (E/T), 26:46272

wave observations near interplanetary discontinuities on Feb. 2, 1969, Pioneer-9 lf, (E), 26:46302

whistler growth rate propagating obliquely in cold plasma with dilute energetic electron density, (E/T), 26:46314

whistler side-band growth due to nonlinear wave--particle interaction, (E/T), 26:46315

whistler-mode wave emission between 3.9 and 10.4 kHz in high-altitude radiation-belt slot, vlf, (E), 26:46312

MAGNETOTAIL

proton entry mechanism at 100 to 300 MeV into distant, during Nov. 18, 1968 and Feb. 25, 1969 solar events, (E/T), 26:46304

structure in, relation to anisotropic electron pitch-angle distribution during substorms, (E/T), 26:46306

MAGNETS

design of superconducting, improvements in, 26:46843(P)
energy--force calculations for, FORGY computer program for, 26:47599 (ANL-TKK/RJL-2)

power supplies for superconducting, design of removable current leads for, 26:45679

MAINE YANKEE POWER REACTOR

coolant loss accident analysis for, revised, 26:47478 (DOCKET-50309-66)

effluents from, effects on environment of chemical, radioactive, and thermal, 26:47473 (DOCKET-50309-61)

electric cables for, qualification testing of, 26:47475 (DOCKET-50309-63)

electrical equipment for, seismic qualification of, 26:47477 (DOCKET-50309-65)

equipment for, earthquake stress analysis for, 26:47476 (DOCKET-50309-64)

instrumentation for, seismic qualification of, 26:47477 (DOCKET-50309-65)

operation of, cost-benefit analysis for, 26:47473 (DOCKET-50309-61)

safety of, revised and additional data on, 26:47474 (DOCKET-50309-62)

site for, ecology, geology, hydrology, and meteorology of, 26:47473 (DOCKET-50309-61)

structures for, earthquake stress analysis for, 26:47476 (DOCKET-50309-64)

MALEIC ACID

radiolysis of single crystals of deuterated, ESR spectral studies of trapped vinyl-type radical from γ , 26:45347

MALFORMATIONS

fistulas, radioinduced rectovaginal, in man following combined external and intracavitary therapy of uterine carcinoma, surgical repair of γ , 26:45800

radioinduced cardiovascular, in rat fetuses following sublethal whole-body exposure in utero, statistical analysis of relation between age, dose, and incidence of fast neutron, 26:45867

MALONIC ACID

radiolysis of single crystals of, deuterium isotope effects on yields of acyl radical from x, 26:45343

MAMMALIA

radiation effects on survival of, review of effects of dose and LET on, 26:45816 (FRNC-R-7)

radioinduced pathological changes in cells of, following internal irradiation, review of, 26:45890 (NYO-3970-29)

radioisotope uptake by small, at Nevada Test Site, fallout, 26:45914 (TID-25954, pp 58-64)

MAN

blood circulation in, calculation of power requirements for, 26:45700(R) (COO-2155-4)

book: Radiation Protection Measurement, 26:45808

cesium-137 content in diets of, in Chicago, Jan. to June 1972, 26:45540 (HASL-257, pp III.45-52)

iodine-131 uptake by thyroid gland in, following administration, diagnostic value of determining rate of, 26:45906(T) (TT-54002/4, pp 462-70)

muscular reflex time of, age dependence of, in residents of Hiroshima and Nagasaki, 26:45795

radiation dose distribution in pelvis of, from radium applicator implanted in uterus, method for calculating, 26:45784(T) (TT-54002/4, pp 456-61)

radiation dose distribution in, during therapy of Hodgkin's disease, role of clinical state of patient in planning x, 26:45786(T) (TT-54002/5, pp 590-3)

radiation dose rate to, during high-altitude supersonic flight, effects of solar flares on cosmic, 26:45806

radiation dose to, during supersonic flight, measurements on board the Concorde prototype, 26:45796

radiation dose to, evaluation of conservative and realistic quality factors used in expressing recommended limits for neutron, 26:45789

radiation dose to, following criticality accident, effects of age, diet, and sex on reliability of sulfur activation in hair and nails for determining fast neutron, 26:45788

radiation dose to, from diagnostic radiography, method and units for expressing cumulative x, 26:45799

radiation effects on count of mast cells in blood of, following mediastinal exposure, x, 26:45810

radiation effects on ESR spectra of blood plasma of, following therapy, 26:45723

radiation effects on incidence of leukemia in, following diagnostic radiography, effects of age, exposure site, number of exposures, and sex on x, 26:45798

radiation effects on survival of, following therapy of breast carcinomas, correlation between rate of growth and size of tumor and x, 26:45783(T) (TT-54002/4, pp 451-5)

radiation effects on survival of, following therapy of metastases to neck lymph nodes in tumors of larynx, nasopharynx, and tonsils, comparison of x, 26:45782(T) (TT-54002/4, pp 442-50)

radiation effects on thyroglobulin-induced stimulation of lymphocytes in, following iodine-131 uptake by thyroid gland, β and γ , 26:45889

radiation effects on, contribution of nuclear interactions with tissues to neutron, 26:45789

radiation effects on, following chronic low dose internal exposure, validity of animal data for evaluation of pathological, 26:45911 (COO-3013-25)

radiation hazards to health of, relation of operation of

nuclear power plants to, 26:45904 (C00-3013-1)
 radiation hazards to, following uptake of radioisotopes, relation of age and dose to neoplasia, 26:45888 (CONF-720520-1)
 radiation hazards to, in relation to operation of nuclear power plants, criteria for establishment of maximum permissible doses on basis of, 26:45778(T) (NP-tr-1941)
 radioimmunoassay of androgens in plasma of females and males, 26:45928
 radioinduced acneiform changes in skin of, following therapy of internal tumors, γ , 26:45793
 radioinduced carcinomas in, following exposure of pelvis, survey of x, 26:45807
 radioinduced chromosome aberrations in bone marrow cells of, following single local exposure, effects of dose and post-exposure time on frequency of γ , 26:45794
 radioinduced chromosome aberrations in cultured lymphocytes from Marshall Island, combined effects of age and sex on frequency of, 26:45792
 radioinduced edema of laryngeal mucosa in, following therapy of laryngeal carcinomas, incidence of x, 26:45781(T) (TT-54002/4, pp 437-41)
 radioinduced injuries in spinal cord of, following therapy of neoplasms, clinical course of latent recovery, x, 26:45797
 radioinduced injuries in, following exposure of pelvis, survey of x, 26:45807
 radioinduced injuries to liver of, following therapy of adjacent organs, dose and time dependence of γ , 26:45809
 radioinduced leukopenia in, following therapy of gynecologic neoplasms, protective effects of cepharanthine against, 26:45805
 radioinduced necrosis in mandible and soft oral tissues of, following therapy of oral tumors, 26:45801
 radioinduced neoplasms in, following uptake of plutonium radioisotopes, relevance of animal data for estimation of dose and time relationships for, 26:45888 (CONF-720520-1)
 radioinduced rectovaginal fistula in, following combined external and intracavitary therapy of uterine carcinoma, surgical repair of γ , 26:45800
 radioinduced stenosis of small intestine of, following therapy of uterine carcinomas, symptoms and treatment of γ and x, 26:45802
 radioisotope content in Argentina, 1960 through 1970, cesium-137 and strontium-90, 26:45520 (NP-19357)
 radioisotope distribution and retention in tissues of, following intake by various routes, compartmental model for determining relation between radiation dose and, 26:45905 (RD/B/N-2291)
 radium-226 excretion in feces of, following oral uptake as radium sulfate dust, effects of alginate ingestion on, 26:45907
 sodium-22 content of, residing in Sweden during 1961 to 1969, relation of consumption of reindeer meat to, 26:45909
 strontium-90 excretion in urine of children in Poland, relation between dietary calcium intake and, 26:45908
MANGANESE
 determination in high-purity iron by neutron activation and preirradiation separation by ion exchange, 26:45216
 determination in presence of cerium by solvent extraction using TTA and spectrophotometry, 26:45226
 determination in steels by neutron activation using plutonium-beryllium source, 26:45199
 effects on recrystallization of iron, 26:45982
MANGANESE, BROMOPENTACARBONYL-
 neutron reactions with, reactions of recoil bromine-82 and manganese-56 from, 26:45357
MANGANESE FLUORIDES
 point defects in, anelastic relaxation caused by, 26:47615 (CU-3194-22)
MANGANESE, IODOPENTACARBONYL-
 neutron reactions with, reactions of recoil iodine-128 and manganese-56 from, 26:45357
MANGANESE ISOTOPES Mn-54
 content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)
 content in surface atmosphere at world sites, 1963 through April 1966, 26:45512(R) (HASL-257(App.))
 distribution in tissues of newts, 26:45913(R) (RLO-2227-T-8-3)
 production of carrier-free, by alpha and helium-3 bombardment of natural chromium, 26:45394
MANGANESE ISOTOPES Mn-55
 neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909
 transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889
MANGANESE ISOTOPES Mn-56
 reactions of recoil, in neutron-irradiated bromopentacarbonylmanganese or iodopentacarbonylmanganese, 26:45357
MANY BODY PROBLEM
 algebraic method, 26:46671
 solutions for classical relativistic three-body, exact stationary, 26:46790
MASS SPECTROGRAPHY
 developments in, review on, 26:45248
 precision using isotope dilution, 26:45225

105 SUBJECTS

MESIC ATOMS

MASS SPECTROMETERS

mass peak in three-dimensional quadrupole, calculation of shape of, 26:45676
 sample admission system of MI-1309, reduction of isotopic fractionation of gaseous samples in, 26:45681
 secondary ion emission analysis using, review on, 26:45249

MATERIALS HANDLING

system for warehouse, computer simulation of automated, 26:47604 (BDX-613-456)

MATERIALS TESTING, NONDESTRUCTIVE

methods for radioactive waste capsules, remote, 26:46099 (ARH-SA-117)

MATHEMATICAL PHYSICS

functional-derivative equations in, method of characteristics for, 26:46728

MATHEMATICS

cobordism theory, introduction to, 26:47627 (SCL-RR-720270)

nonlinear analysis in, methods of, 26:47283 (CERN-72-8)
 numerical algebra, KWIC index for, 26:47287 (ORNL-4778)

MATTER

field theory for charged, formulation of Einstein-Kaufman nonsymmetric unified, (T), 26:46694
 superfluidity of hadronic, derivation of critical temperature for, (T), 26:46596

MEATS

ham, radiation processing of, for disinfection of mites, effectiveness of electron, 26:45886
 sausage, radiation processing of packaged, inhibitory effects in combination with heat treatment on sporulation of Clostridium, 26:45747

MECHANICAL ENGINEERING

radioisotope uses in, review on, 26:45710
 radioisotope uses in, review on, 26:45711

MEDICAL PERSONNEL

radiation dose to, during use of californium-252 for interstitial and intracavitary therapy, methods for reducing neutron, 26:45790
 radiation monitoring of diagnostic radiography, in Canada, 26:45811

MEDICAL SUPPLIES

radiosterilization of disposable, sensitivity of contaminating microorganisms to, 26:45748

MEDICINE

book: Masking and Demasking of Chemical Reactions. Theoretical Aspects and Practical Applications, 26:45188

radiation safety in Virginia dental offices, 26:45500
 translation of Polish Review of Radiology and Nuclear Medicine volume 35, number 4, 1971, 26:45779(T) (TT-54002/4)

translation of Polish Review of Radiology and Nuclear Medicine, volume 35; number 5, 1971, 26:45785(T) (TT-54002/5)

MEDITERRANEAN SEA

book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
 fallout content and distribution in silts in, 26:45531 (C00-3563-3)

MENDOCCINO POWER REACTOR, UNIT 1

owners of, Department of Justice opinion on monopolistic actions of, 26:47562 (DOCKET-50398-17)
 safety of, revised and additional data on, 26:47561 (DOCKET-50398-16)

safety of, revised and requested additional data on, 26:47559 (DOCKET-50398-11)

site for, requested data on geology and seismology of, 26:47560 (DOCKET-50398-15)

MENDOCCINO POWER REACTOR, UNIT 2

owners of, Department of Justice opinion on monopolistic actions of, 26:47562 (DOCKET-50398-17)
 safety of, revised and additional data on, 26:47561 (DOCKET-50398-16)

site for, requested data on geology and seismology of, 26:47560 (DOCKET-50398-15)

MERCAPTO GROUP

reactions with free radicals, deuterium and tritium isotope effects on hydrogen abstraction in, 26:45342

MERCURY

determination in hair by neutron activation and γ spectroscopy, 26:45214

determination of trace amounts of, in environmental samples by neutron activation, 26:45198

mass flow choking limit for heat pipe working fluid of, (E), 26:45460

positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268

MERCURY ALLOYS AND SYSTEMS

Hg--In--Sn, thermodynamic properties of amalgams in, at 150 to 340°C, 26:46004

MERCURY ISOTOPES

x-ray emission in muonic, test of quantum electrodynamic corrections from, 26:46708

MERCURY ISOTOPES Hg-197

production of carrier-free, in cyclotron by proton bombardment of gold target, 26:45395

MESIC ATOMS

gamma and x-ray spectra from thallium-203 and -205 μ^- ,

(E), 26:47031

MESONS

- interactions with baryons and mesons, duality properties of Regge cuts in, (T), 26:46600
- interactions with baryons, partial wave relations from fixed-u dispersion relations for, 26:46610
- interactions with baryons, test of equal-phase assumption for helicity amplitudes for, (T), 26:46621
- interactions with baryons, urbaryon rearrangement diagrams for differential cross sections and polarizations for, (T), 26:46647
- interactions with mesons, partial wave relations from fixed-u dispersion relations for, 26:46610
- interactions with mesons, space-time description of, 26:46603
- interactions with neutrons at high energies, electromagnetic effects in nucleon polarization in, (T), 26:46520 (JINR-P2-6438)
- interactions with spin-1/2 baryons using polarized targets, lecture on method for determination of spins of baryon resonances produced in quasi-two-body, (T), 26:46628
- mass formulas for, dynamic interpretation of, (T), 26:46594
- production in $e + e$, $e + p$, and $p + p$ interactions, invariant method for differential cross sections, angular distributions, and two-photon mechanism for, (T), 26:46518 (ITEF-848)
- reactions with nuclei at 5 GeV, change of mechanism for inelastic, (T), 26:47038(T) (BNL-tr-487)

MESONS (η)

- decay $\eta \rightarrow 2\gamma$, tests of Han-Nambu and Gell-Mann-Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)
- decay $\eta \rightarrow \pi^+ + \pi^- + \gamma$, spark chamber study of charge asymmetry in, (E), 26:46482
- decay $\eta \rightarrow \pi^+ + \pi^- + \pi^0$, spark chamber study of charge asymmetry in, (E), 26:46483
- production in $\gamma + p$ interactions, model predictions of backward differential cross section for, (T), 26:46529
- production in $\pi^- + p + \eta$ and n interactions, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- production of η in anti $p + p$ interactions at rest, (E), 26:46465

MESONS (K)

- decay $K \rightarrow \pi + l + \nu$, bounds for parameters of, 26:46575
- decay $K \rightarrow \pi + l + \nu$, scalar form factors and parameters for, 26:46583
- interactions anti $K + N \rightarrow \pi + A$ at low energies, Veneziano duality for dips in cross sections for, (T), 26:46635
- interactions anti $K + N \rightarrow \pi + Y$ above 2.5 GeV/c, review of backward and forward, (E), 26:46489
- interactions $K + N \rightarrow K + N$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- interactions $K + N$ at high energies, modifications in Regge pole model for phenomenology of total cross sections in forward, (T), 26:46633
- interactions $K + N$, low lying singularities for energy dependence of cross sections for angular momentum plane structure for amplitudes of line-reversed, (E/T), 26:46629
- interactions $K + N$, nature of absorption in ω -exchange amplitudes for, (T), 26:46623
- interactions $K + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462
- interactions $\pi + K \rightarrow \omega + K$, finite-energy sum rule for, (T), 26:46611
- interactions $\pi + K \rightarrow \phi + K$, finite-energy sum rule for, (T), 26:46611
- production in proton reactions with nuclei at 24 GeV/c, spectra of, (E), 26:46459

MESONS (K^-)

- capture by copper and nickel, gamma spectra from, (E), 26:46940
- capture by nuclei, $K^- + N$ scattering amplitudes for, (E/T), 26:46487
- interactions $K^- + n + A \rightarrow \pi^+ + 2\pi^- + \pi^0$ (MM) at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
- interactions $K^- + n + A \rightarrow \pi^- + K^+ (K_2^0) + K^- (K_1^0)$ at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
- interactions $K^- + n + A \rightarrow \pi^-$, differential cross section and polarization for backward, 26:46648
- interactions $K^- + n + A(\Sigma^0) \rightarrow \pi^+ + 2\pi^-$ at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
- interactions $K^- + N$ at intermediate energies, test of two-component theory of duality for, 26:46624
- interactions $K^- + N$, scattering amplitude for meson (K^-) capture by nuclei, (E/T), 26:46487
- interactions $K^- + p + \text{anti } K^0 + n$ with octet exchange in crossed channel, models for extrapolation of differential cross sections from experimental points for, (T), 26:46630
- interactions $K^- + p + \text{anti } K^0 + n$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642
- interactions $K^- + p + \text{anti } K^0 + n$, modified absorptive Regge cut model for, (T), 26:46622

interactions $K^- + p + \text{anti } K^0(K^-) + n(p)$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635

- interactions $K^- + p + K^- + p + \pi^+ + \pi^- + \gamma(\eta)$ at 3.9 and 4.6 GeV/c, production of $M(953)$ and $\phi(1019)$ in, (E), 26:46471 (BNL-16954)
- interactions $K^- + p + K^- + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492
- interactions $K^- + p + K^- + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + K^+ + K^-$ interactions and, (E), 26:46463
- interactions $K^- + p + A + \omega$ at 3.95 GeV/c, differential cross sections and ω density matrix elements for, (E), 26:46485
- interactions $K^- + p + A + \rho$ at 3.95 GeV/c, differential cross sections and ρ density matrix elements for, (E), 26:46485
- interactions $K^- + p + A + \rho$ at 3.95 GeV/c, differential cross sections and ρ density matrix elements for, (E), 26:46485
- interactions $K^- + p + \pi^- + \Sigma^+$ at 5 GeV/c, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut collision mechanism for cross sections for, (T), 26:46631
- interactions $K^- + p + \pi^- + \Sigma^+$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642
- interactions $K^- + p + \Sigma^0 + \phi$ at 3.95 GeV/c, differential cross sections and ϕ density matrix elements for, (E), 26:46485
- interactions $K^- + p + \Sigma^0 + \rho$ at 3.95 GeV/c, differential cross sections and ρ density matrix elements for, (E), 26:46485
- interactions $K^- + p$ at 10 GeV/c in bubble chamber, review of elastic and total cross sections, dynamics of two-body, quasi-two-body, and multibody reactions, and Ω^- and resonance production in, (E), 26:46480
- interactions $K^- + p$, Pomeranchuk theorem violation in total cross sections of nonforward, (T), 26:46634
- scattering by $A = 10$ to 35 nuclei, scattering lengths for, (E), 26:46912

MESONS (K_0)

- decay $K_1^0 \rightarrow 2\gamma/K_1^0 \rightarrow \text{all}$ in optical spark chamber, branching ratio for, (E), 26:46434
- decay $K_1^0 \rightarrow 2\mu$, limitations on width for, (T), 26:46572
- decay $K_2^0 \rightarrow 2\gamma/K_2^0 \rightarrow \text{all}$ in optical spark chamber, branching ratio for, (E), 26:46434
- decay $K_2^0 \rightarrow 3\pi^0$ in Wilson chamber, branching ratio for, (E), 26:46449(T) (SLAC-Trans-143)
- decay $K_2^0 \rightarrow \chi^{10} + \gamma$, existence of χ^{10} from, (T), 26:46577
- decay $K_2^0 \rightarrow \mu^+ + \mu^-$, CP-invariant contribution of χ^{10} intermediate state to, (T), 26:46577
- decay $K_2^0 \rightarrow \mu^+ + \mu^-$, CPT violation in, (E/T), 26:46680
- decay $K_2^0 \rightarrow \pi + e + \nu$ in Wilson chamber, branching ratio for, (E), 26:46449(T) (SLAC-Trans-143)
- decay $K_2^0 \rightarrow \pi + \mu + \nu$ in Wilson chamber, branching ratio for, (E), 26:46449(T) (SLAC-Trans-143)
- decay $K_2^0 \rightarrow \pi^+ + \pi^- + \pi^0$ in heavy-liquid bubble chamber, slope parameter and test of CP violation in charge asymmetry in, (E), 26:46494
- decay $K_2^0 \rightarrow \pi^+ + \pi^- + \pi^0$ in Wilson chamber, branching ratio for, (E), 26:46449(T) (SLAC-Trans-143)
- decay of combinations of states of, in time-dependent external field, frequency dependence of decay constants for parametric resonance in exponential, (T), 26:46571 (KFKI-72-15)
- decay of, mixing parameter in, (T), 26:46585
- lifetime of K_2^0 , Wilson chamber study of, (E), 26:46449(T) (SLAC-Trans-143)
- regeneration $K_2^0 \rightarrow K_1^0$ on neutrons, Pomeranchuk theorem violation in amplitudes of, (T), 26:46634
- regeneration $K_2^0 \rightarrow K_1^0$ on protons, modifications in Regge pole model for amplitude of, (T), 26:46633

MESONS (K^+)

- decay $K^+ \rightarrow e^+ + \pi^0 + \nu$, branching ratio for, (E), 26:46455
- decay $K^+ \rightarrow \mu^+ + \nu$, branching ratio for, (E), 26:46455
- decay $K^+ \rightarrow \mu^+ + \pi^0 + \nu$, branching ratio for, (E), 26:46455
- decay $K^+ \rightarrow \pi^0 + \pi^0 + e^+ + \nu$, branching ratio for, (E), 26:46454
- decay $K^+ \rightarrow \pi^+ + \gamma + \gamma$, upper limit on branching ratio for, (E), 26:46454
- decay $K^+ \rightarrow \pi^+ + \pi^0 + \gamma$, branching ratio for, (E), 26:46454
- decay $K^+ \rightarrow \pi^+ + \pi^0 + \pi^0$, branching ratio for, (E), 26:46455
- decay $K^+ \rightarrow \pi^+ + \pi^0$, branching ratio for, (E), 26:46455
- decay $K^+ \rightarrow \pi^+ + \pi^- + \pi^+$, branching ratio for, (E), 26:46455
- interactions $K^+ + n + K^0 + p$ with octet exchange in crossed channel, models for extrapolation of differential cross sections from experimental points for, (T), 26:46630
- interactions $K^+ + n + K^0 + p$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642
- interactions $K^+ + n + K^0 + p$, modified absorptive Regge cut model for, (T), 26:46622
- interactions $K^+ + n + K^0 + p$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- interactions $K^+ + N$ at intermediate energies, test of

- two-component theory of duality for, (T), 26:46624
 interactions $K^+ + p \rightarrow K^+ + p$ at 0.52 to 1.89 GeV/c, partial-wave amplitudes and exotic resonance in P/sub 3/2/ partial wave from phase shift analysis of, (T), 26:46632
 interactions $K^+ + p \rightarrow K^+ + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492
 interactions $K^+ + p \rightarrow K^+ + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p \rightarrow K^+ + K^-$ interactions and, (E), 26:46463
 interactions $K^+ + p \rightarrow p + K^+ + \omega$, $\omega \rightarrow \pi^- + \pi^+ + \pi^0$ at 12 GeV/c, cross sections, momentum-transfer spectra, and angular-decay spectra for, (E), 26:46484
 interactions $K^+ + p \rightarrow p + K^+ + \phi$, $\phi \rightarrow K^- + K^+$ at 12 GeV/c, cross sections, momentum-transfer spectra, and angular-decay spectra for, (E), 26:46484
 interactions $K^+ + p$ at 5 GeV/c in hydrogen bubble chamber, systematic methods for particle track analysis of, (E), 26:45651
 interactions $K^+ + p$, Pomeranchuk theorem violation in total cross sections of nonforward, (T), 26:46634
 production in $\gamma + N \rightarrow K^+ + N$ interactions at 5 to 15 GeV/c, Regge model for cross sections for, (T), 26:46534
 production in $\gamma + p \rightarrow K^+ + \Lambda$ interactions at 1 to 2 and 8 to 16 GeV, finite-energy sum rules for duality relations between, (T), 26:46535
- MESONS (ω)**
 decay $\omega \rightarrow 2\pi/\omega \rightarrow 3\pi$, lower limit on branching ratio for, (E), 26:46496
 decay $\omega \rightarrow e^+ + e^-$, width and branching ratio for, (E), 26:46427 (CONF-710816-13)
 decay $\omega \rightarrow \pi^0 + \gamma$, test of Han-Nambu and Gell-Mann-Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)
 decay $\omega \rightarrow \pi^+ + \pi^-$, branching ratio for, (E), 26:46428 (CONF-710816-14)
 decay $\omega^0 \rightarrow 2\pi^0 + \gamma$, branching ratio for, (E), 26:46472 (CONF-710816-12)
 decay $\omega^0 \rightarrow \eta + \gamma$, upper limit for branching ratio for, (E), 26:46472 (CONF-710816-12)
 decay $\omega^0 \rightarrow \pi^0 + \gamma$, branching ratio for, (E), 26:46472 (CONF-710816-12)
 decay $\omega^0 \rightarrow \pi^+ + \pi^- + \pi^0$, branching ratio for, (E), 26:46472 (CONF-710816-12)
 interference of $\rho^0 \rightarrow \omega$ in $\gamma + N \rightarrow \pi^+ + \pi^- + \Delta$ interactions, calculation of, (T), 26:46527
 interference of $\rho^0 \rightarrow \omega$ in $e^+ + e^- \rightarrow \pi^+ + \pi^-$ annihilation in colliding beams, amplitude and phase of, (E), 26:46428 (CONF-710816-14)
 interference of $\rho^0 \rightarrow \omega$ in $\pi^- + p \rightarrow \pi^+ + \pi^- + n$ interactions at 15 GeV/c, (E), 26:46496
 production in $\gamma + p$ interactions at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446
 production in $\gamma + p$ interactions, model predictions of backward differential cross section for, (T), 26:46529
 production in $K^- + n \rightarrow \Lambda + 2\pi^- + \pi^+ + \pi^0$ interactions at 4.5 GeV/c in deuterium bubble chamber, cross sections for, (E), 26:46474 (COO-1428-306)
 production in $K^- + p \rightarrow \Lambda + \omega$ interactions at 3.95 GeV/c, differential cross sections and density matrix elements for, (E), 26:46485
 production of ω^0 in anti $p + p$ interactions at rest, (E), 26:46465
 width in $e^+ + e^- \rightarrow \pi^+ + \pi^- + \pi^0$ interactions in storage ring, (E), 26:46427 (CONF-710816-13)
- MESONS (ϕ)**
 decay $\phi \rightarrow \eta + \gamma$ from production in $\pi^- + p \rightarrow \eta + \phi$ interactions at 1.8 GeV/c, branching ratio for, (E), 26:46435
 decay $\phi \rightarrow K_1^0(K^+) + K_2^0(K^-)$ in deuterium bubble chamber, branching ratio for, (E), 26:46474 (COO-1428-306)
 decay $\phi \rightarrow \eta + \gamma$, branching ratios for, (E), 26:46472 (CONF-710816-12)
 mass from $e^+ + e^- \rightarrow \phi + K_1^0 + K_2^0$, $K_1^0 \rightarrow \pi^+ + \pi^-$ interactions, (E), 26:46443
 production in $\gamma + p$ interactions, model predictions of backward differential cross section for, (T), 26:46529
 production in $K^- + n$ interactions at 4.5 GeV/c in deuterium bubble chamber, cross sections for, (E), 26:46474 (COO-1428-306)
 production in $K^- + p \rightarrow \Lambda(\Sigma^0) + \phi$ interactions at 3.95 GeV/c, differential cross sections and density matrix elements for, (E), 26:46485
- MESONS (π)**
 coupling constant for $A_1\rho\omega$ from analysis of $\pi + N \rightarrow A_1 + N$ interaction, relation of t-channel helicity conservation with, (T), 26:46638
 coupling constant for $\rho\omega\omega$, (E), 26:46428 (CONF-710816-14)
 coupling constants of $A_1\rho\omega$, use of collinear dispersion relations and equal-time commutators for, 26:46612
 coupling constants of $\rho\omega\omega$, use of collinear dispersion relations and equal-time commutators for, 26:46612
 exchange contributions to AN potential from $\Lambda + N$ interactions, 26:46652
 form factor determination from $e + N \rightarrow \pi + N$ interactions in first resonance region, model with axial vector form factor for, (T), 26:46513 (CONF-710816-17)
 form factor from $\pi + \pi$ interactions in Bethe-Salpeter model, electromagnetic, 26:46615
 form factor inside analytic region, upper bound for electromagnetic, 26:46528
 form factor of, contributions of heavy ρ' meson to, 26:46561
 interactions $\gamma + \pi \rightarrow \gamma + \pi$, pomeron coupling in forward Compton, (T), 26:46560
 interactions $\pi + K \rightarrow \omega + K$, finite-energy sum rule for, (T), 26:46611
 interactions $\pi + K \rightarrow \phi + K$, finite-energy sum rule for, (T), 26:46611
 interactions $\pi + N \rightarrow A_1 + N$, relation of $A_1\rho\omega$ coupling constant with t-channel helicity conservation in, (T), 26:46638
 interactions $\pi + N \rightarrow A_1 + N$, s-channel and t-channel helicity conservation in, (T), 26:46646
 interactions $\pi + N \rightarrow K + \Lambda$ at low energies, Veneziano duality for dips in cross sections for, (T), 26:46635
 interactions $\pi + N \rightarrow N + \text{virtual } \gamma$, quasi-threshold behavior of amplitudes and cross sections for, (T), 26:46620 (JINR-P4-6475)
 interactions $\pi + N \rightarrow \pi + N$, duality in parton model for, (T), 26:46599
 interactions $\pi + N \rightarrow \pi + N$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
 interactions $\pi + N$ at 7 to 20 GeV, amplitude analysis in Coulomb interference region for, (T), 26:46625
 interactions $\pi + N$ at high energies and small momentum transfer, review of near forward, (E/T), 26:46636
 interactions $\pi + N$ at high energies, modifications in Regge pole model for phenomenology of total cross sections in forward, (T), 26:46633
 interactions $\pi + N$, low-energy theorem from pion source dynamics for amplitude of, 26:46641
 interactions $\pi + N$, removal of ambiguities from phase-shifts for inelastic, 26:46640
 interactions $\pi + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462
 interactions $\pi + p$, low lying singularities for energy dependence of cross sections for angular momentum plane structure for amplitudes of line-reversed, (E/T), 26:46629
 interactions $\pi + \pi \rightarrow \pi + \pi$, explicit Poincare group representations labelled by additional discrete quantum numbers for, (T), 26:46665 (CTH-ITP-71-14)
 interactions $\pi + \pi$ in Bethe-Salpeter model, π electromagnetic form factor from, 26:46615
 mass of, amplitude dependence on photon mass in $\gamma + \gamma \rightarrow \pi^+ + \pi^-$ interactions for electromagnetic, (T), 26:46519 (JINR-P2-6333)
 production in anti $p + p$ interactions at 1.51 to 2.90 GeV/c in bubble chamber, multiplicity and cross sections for, (E), 26:46469
 production in $e + N \rightarrow \pi + N$ interactions near threshold, current algebra and PCAC hypothesis for amplitudes and cross section sensitivity to nucleon axial-vector form factor for, (T), 26:46549
 production in $\gamma + N \rightarrow \pi + N$ interactions, method for dispersion theory for, (T), 26:46511 (CONF-710816-9)
 production in $\mu + N \rightarrow \pi + \text{anything}$ interactions at 10.1 and 15.8 GeV, momentum distributions of, (E), 26:46452
 production in proton reactions with nuclei at 24 GeV/c, spectra of, (E), 26:46459
 production in $\pi^- + p$ interactions at 4 GeV, spectrum symmetry of longitudinal momenta in, (T), 26:46619 (JINR-P1-6461)
 reactions with neon, research at Duke Univ., July 1971 to June 1972, (KE), 26:46479(R) (ORO-3065-24)
 reactions with nuclei above 10 GeV, cascade-evaporation model for shower particle cascade multiplicity and many-particle interaction effects in inelastic, (T), 26:46649 (JINR-P2-6395)
 reactions with xenon at 2.34 to 9 GeV/c, ω^0 and resonance production in, (E), 26:46476 (JINR-P1-6408)
- MESONS (π^-)**
 interactions $\pi^- + d \rightarrow 2n + \gamma$ at 150 to 300 MeV, phenomenological calculation of soft-photon cross sections for, (T), 26:46531
 interactions $\pi^- + N \rightarrow \rho^0 + N$, interference of photon exchange with nucleon exchange in, (T), 26:46643
 interactions $\pi^- + p \rightarrow 2\pi^0 + n$ at 1.60 to 2.40 GeV/c, total cross sections, peripheral dipion production distribution and scattering, and $\Delta^0(1236)$ production in, (E), 26:46477 (LBL-378)
 interactions $\pi^- + p \rightarrow \eta + n$, duality-breaking effect in exchange-degenerate, (T), 26:46642
 interactions $\pi^- + p \rightarrow \eta + n$, modified absorptive Regge cut model for, (T), 26:46622
 interactions $\pi^- + p \rightarrow \eta + \phi$ at 1.8 GeV/c, branching ratio for $\phi \rightarrow \eta + \gamma$ decay from production in, (E), 26:46435
 interactions $\pi^- + p \rightarrow \eta(X) + n$ Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
 interactions $\pi^- + p \rightarrow \pi^0 + n$ at 1.6 and 2.4 GeV/c, scintillation counter study of resonance production in, (E), 26:46475 (IS-2886)
 interactions $\pi^- + p \rightarrow \pi + n$ at 2.5 GeV/c, enhancement at 968 MeV in, (E), 26:46430 (COO-1195-241)
 interactions $\pi^- + p \rightarrow \pi + \pi^0$ at 22.6 MeV, scintillation counter study of time-of-flight spectrum from, p. 221-3, (E), 26:46873(R) (CEA-N-1522)

interactions $\pi^- + p + p + \text{MM}^-$ at 25 and 40 GeV/c, cross section scaling in $s^{1/2}$ s and triple-Pomeranchuk coupling in, (T), 26:46616 (BNL-16928)

interactions $\pi^- + p + p + \pi^-$ at 6 GeV/c, polarization parameter in backward, (E), 26:46490

interactions $\pi^- + p + \pi^0 + n$ at 5.0 to 8.0 GeV/c, polarization parameter in, (E), 26:46491

interactions $\pi^- + p + \pi^0 + n$ at high energies, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut collision mechanism for cross sections and polarization for, (T), 26:46631

interactions $\pi^- + p + \pi^0 + n$ at high energies, total cross sections for, (T), 26:46618 (IFVE-STF-71-71)

interactions $\pi^- + p + \pi^0 + n$, 3-channel helicity amplitudes without pomeron for, (T), 26:46637

interactions $\pi^- + p + \pi^0 + n$, duality-breaking effect in exchange degenerate model for, (T), 26:46642

interactions $\pi^- + p + \pi^0 + n$, modified absorptive Regge cut model for, (T), 26:46622

interactions $\pi^- + p + \pi^0 + n$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645

interactions $\pi^- + p + \pi^0(\text{eta}) + n$ with octet exchange in crossed channel, models for extrapolation of differential cross sections from experimental points for, (T), 26:46630

interactions $\pi^- + p + \pi^0(\pi^-) + n(p)$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635

interactions $\pi^- + p + \pi^+ + 2\pi^- + p$ at 11 to 40 GeV in bubble and spark chambers, A_2 production in, (E), 26:46473 (COO-1195-239)

interactions $\pi^- + p + \pi^- + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions $\pi^- + p + \pi^- + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + \pi^+ + \pi^-$ interactions and, (E), 26:46463

interactions $\pi^- + p + \pi^+ + \pi^- + n$ at 15 GeV/c, vector dominance and ρ - ω interference in, (E), 26:46496

interactions $\pi^- + p + \pi^- + \pi^+(\pi^0) + n(p)$ at 4 GeV/c, effective mass and angular distributions and differential cross sections for $\Delta(1236)$ and $\Delta^+(1236)$ forward production in, (E), 26:46488

interactions $\pi^- + p + \rho^0 + n$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481

interactions $\pi^- + p + \rho^0 + p$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481

interactions $\pi^- + p$ at 4 GeV, spectrum symmetry of pion longitudinal momenta in, (T), 26:46619 (JINR-P1-6461)

interactions $\pi^- + p$ at 6 and 16 GeV, intermediate energy contribution to dispersion sum rules and rotation of polarization in Regge pole model for, (T), 26:46617 (IFVE-STF-71-15)

interactions $\pi^- + p$ at 60 GeV/c, two-particle correlations for inclusive, (E/T), 26:46460

interactions $\pi^- + p$ at high energies, interpolation curves for total cross sections for elastic, elastic with charge exchange, and inelastic, (E), 26:46478(T) (ORNL-tr-2544)

interactions $\pi^- + p$ in dual absorptive model, Pomeron and f exchange amplitudes in, (T), 26:46639

production in $e + p + e + p + \pi^+ + \pi^-$ interactions in rho resonance region, extension of diffraction--dissociation model for polarization, momentum transfer, and virtual photon and dipion mass dependence of cross sections for, (T), 26:46517 (DESY-72/23)

production in $e^+ + e^- + \pi^+ + \pi^-$ interactions in colliding beams, cross sections for, (E), 26:46428 (CONF-710816-14)

production in photon reactions with aluminum, carbon, lead, and tantalum at 1.160 GeV at 150° , photoemulsion study of kinetic energy and total yield for, (E), 26:46441

production of soft, in $e + N + \pi^- + \text{anything}$ interactions, PCAC hypothesis and soft pion theorems for cross sections for, (T), 26:46521 (LBL-773)

reactions $(\pi^+, \pi^-)_{3n}$ with helium-3, analysis of nonresonant neutron attraction in, (T), 26:46862

reactions (π^-, π^-n) at 140 to 240 MeV, cross sections for, (E), 26:46486

reactions (π^-, π^-n) with fluorine-19 at 140 to 240 MeV, cross sections for, (E), 26:46486

reactions (π^-, π^-p) with carbon-12 and lithium-6 at 1 GeV/c, π -meson momenta from, (E), 26:46869

MESONS (π^0)

decay of, one-fermion loop contributions to two-graviton, (T), 26:46501

decay $\pi^0 + 2\gamma$, tests of Han--Nambu and Gell-Mann--Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)

interactions $\pi^0 + \pi^0$ in $\pi^- + p + 2\pi^0 + n$ interactions at 1.60 to 2.40 GeV/c, (E), 26:46477 (LBL-378)

lifetime of, proposed experiment for streamer chamber study of cross sections for $\gamma + {}^4\text{He} + {}^4\text{He} + \pi^0$ interactions at 2 to 5 GeV for determination of, (E), 26:46432(T) (SLAC-Trans-142)

production in $e + N + \pi^0 + N$ interactions in first resonance region, multipole amplitude analysis of angular distributions for, (T), 26:46512 (CONF-710816-16)

production in $\gamma + {}^4\text{He} + {}^4\text{He} + \pi^0$ interactions at 2 to 5 GeV, proposed experiment for streamer chamber study of cross sections for, (E), 26:46432(T) (SLAC-Trans-142)

production in $\gamma + p + p + \pi^0$ interactions at 426 to 918 MeV, asymmetry measurements for, (E), 26:46442

production in photon reactions (γ, π^0) with complex nuclei at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447

production in π reactions with xenon at 2.34 to 9 GeV/c, (E), 26:46476 (JINR-P1-6408)

production of dipion in $\pi^- + p + 2\pi^0 + n$ interactions at 1.60 to 2.40 GeV/c, distribution for peripheral, (E), 26:46477 (LBL-378)

MESONS (π^+)

interactions $\pi^+ + d$ at high energies, research at Duke Univ., July 1971 to June 1972, (E), 26:46479(R) (ORO-3065-24)

interactions $\pi^+ + N + \rho + N$, interference of photon exchange with nucleon exchange in, (T), 26:46643

interactions $\pi^+ + p + K^+ + \pi^+$ at 5 GeV/c, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut mechanism for cross sections and polarization for, (T), 26:46631

interactions $\pi^+ + p + K^+ + \pi^+$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642

interactions $\pi^+ + p + p + \pi^+$ at 6 GeV/c, polarization parameter in backward, (E), 26:46490

interactions $\pi^+ + p + \pi^0 + \Delta^+(1236)$ at 1820 to 2090 MeV, partial-wave analysis of, (E), 26:46495

interactions $\pi^+ + p + \pi^+ + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions $\pi^+ + p + \pi^+ + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + \pi^+ + \pi^-$ interactions and, (E), 26:46463

interactions $\pi^+ + p + \pi^+ + p$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635

interactions $\pi^+ + p + \pi^+ + \pi^+ + \pi^- + \pi^0 + p$ at 5 GeV/c, evidence for double diffraction dissociation in, (E), 26:46493

interactions $\pi^+ + p + \pi^+ + \pi^+ + \pi^- + \pi^+ + n$ at 5 GeV/c, evidence for double diffraction dissociation in, (E), 26:46493

interactions $\pi^+ + p + \rho^0 + p$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481

interactions $\pi^+ + p$ at high energies, interpolation curves for total cross sections for elastic, elastic with charge exchange, and inelastic, (E), 26:46478(T) (ORNL-tr-2544)

interactions $\pi^+ + p$ in dual absorptive model, Pomeron and f exchange amplitudes in elastic, (T), 26:46639

production in $e + p + e + p + \pi^+ + \pi^-$ interactions in rho resonance region, extension of diffraction--dissociation model for polarization, momentum transfer, and virtual photon and dipion mass dependence of cross sections for, (T), 26:46517 (DESY-72/23)

production in $e^+ + e^- + \pi^+ + \pi^-$ interactions in colliding beams, cross sections for, (E), 26:46428 (CONF-710816-14)

production in $\gamma + p + n + \pi^+$ interactions at 390 to 909 MeV, asymmetry measurements for, (E), 26:46442

production in $\gamma + p + \pi^+ + n$ interactions at 300 to 900 MeV in polarized butanol target, asymmetry of cross sections for, (E), 26:46444

production in $\gamma + p + \pi^+ + n$ interactions at 5 and 16 GeV using polarized proton target, review of asymmetry in, (E), 26:46436

production in photon reactions with aluminum, carbon, lead, and tantalum at 1.160 GeV at 150° , photoemulsion study of kinetic energy and total yield for, (E), 26:46441

production of soft, in $e + N + \pi^+ + \text{anything}$ interactions, PCAC hypothesis and soft pion theorems for cross sections for, (T), 26:46521 (LBL-773)

reactions $(\pi^+, 2p)$ with $A = 2$ to 16 nuclei at 76 MeV, cross sections for, (E), 26:46870

reactions (π^+, π^0) with zirconium-91 at 30 MeV, cross section for, p. 224-5, (E), 26:46873(R) (CEA-N-1522)

MESONS (ρ)

coupling constant for $A_1\rho\omega$ from analysis of $\pi + N + A_1 + N$ interaction, relation of t -channel helicity conservation with, (T), 26:46638

coupling constant for γ - ρ in vector dominance model, 26:46445

coupling constant for $\rho\omega\omega$, (E), 26:46428 (CONF-710816-14)

coupling constants of $A_1\rho\omega$, use of collinear dispersion relations and equal-time commutators for, 26:46612

coupling constants of $\rho\omega\omega$, use of collinear dispersion relations and equal-time commutators for, 26:46612

decay $\rho \rightarrow e^+ + e^-$, branching ratio and width for, (E), 26:46428 (CONF-710816-14)

decay $\rho \rightarrow \pi^0 + \gamma$, test of Han--Nambu and Gell-Mann--Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)

interactions $\rho\omega^0 + N + \rho\omega^0 + N$, total and differential cross sections for, (E), 26:46445

interference of $\rho\omega^0$ - ω in $\gamma + N + \pi^+ + \pi^- + A$

- interactions, calculation of, (T), 26:46527
- interference of $\rho^0 \rightarrow \pi^+ + \pi^- \rightarrow \pi^+ + \pi^-$ annihilation in colliding beams, amplitude and phase of, (E), 26:46428 (CONF-710816-14)
- interference of $\rho^0 \rightarrow \pi^+ + \pi^- \rightarrow \pi^+ + \pi^-$ interactions at 15 GeV/c, (E), 26:46496
- mass and width in $\pi^+ + \pi^- \rightarrow \pi^+ + \pi^-$ annihilation in colliding beams, (E), 26:46428 (CONF-710816-14)
- production in $\pi^+ + \pi^- \rightarrow \pi^+ + \pi^-$ interactions in colliding beams, cross sections for, (E), 26:46428 (CONF-710816-14)
- production in $\gamma + d \rightarrow \rho^0 + d$ interactions at 6, 12, and 18 GeV, forward differential cross sections for, (E), 26:46445
- production in $\gamma + d \rightarrow \rho^0 + d$ interactions in Glauber theory, differential cross sections and density matrix elements for, (T), 26:46555
- production in $\gamma + N \rightarrow \rho^0 + A(1236)$ interactions, method for dispersion theory for, (T), 26:46511 (CONF-710816-9)
- production in $\gamma + p \rightarrow \rho^0 + p$ interactions in geometric diffraction picture, slope of cross sections for virtual, 26:46564
- production in $\gamma + p$ interactions at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446
- production in $\gamma + p$ interactions, model predictions of backward differential cross section for, (T), 26:46529
- production in $K^- + n$ interactions at 4.5 GeV/c in deuterium bubble chamber, cross sections for, (E), 26:46474 (COO-1428-306)
- production in $K^- + p \rightarrow \Sigma^0 + \rho^0$ interactions at 3.95 GeV/c, differential cross sections and density matrix elements for, (E), 26:46485
- production in $\pi^+ + p$ interactions at 6 GeV/c, meson (π) exchange in, (E), 26:46481
- production in $\pi^- + N \rightarrow \rho^0 + N$ and $\pi^+ + N \rightarrow \rho^0 + N$ interactions, interference of photon exchange with nucleon exchange in, (T), 26:46643
- production in $\pi^- + p \rightarrow \pi^0 + n$ interactions at 1.6 GeV/c, scintillation counter study of, (E), 26:46475 (IS-2886)
- production in $\pi^- + p \rightarrow \pi^+ + \pi^- + n$ at 15 GeV/c, differential cross sections and density matrix elements for, (E), 26:46496
- production of ρ^0 in anti $p + p$ interactions at rest, (E), 26:46465
- production of ρ^0 in $\gamma + p \rightarrow \rho^0 + \text{anything}$ interactions at 4 to 18 GeV, diffractive fragmentation in incident and target particles in, (E), 26:46437
- production of ρ^0 in $\gamma + p \rightarrow \rho^0 + p$ interactions at 2.8 and 4.7 GeV in hydrogen bubble chamber, t dependence of mass distributions, spin dependence, and cross sections for, (T), 26:46438
- Regge cuts in forward $p + p \rightarrow p + p$ interactions, inclusive finite mass sum rules for upper bound for, 26:46651 (RPP/T-18)
- MESONS, PSEUDOSCALAR
- interactions with baryons, duality constraints for baryon exchange degeneracy patterns and absence of parity doublets and backward peak behavior in octet pseudoscalar meson and decuplet baryon production in, (T), 26:46627
- MESONS, RESONANCE
- charge conjugation parity of N anti $N(1970)$, (E/T), 26:46458
- classification of spin--parity--charge conjugation parity = 3^- , in $SU(3)$ nonet, 26:46611
- decay angular distributions in photon interactions with spin- $1/2$ baryons, comparison of excited quark, interacting quarks, and additive quark models for constraints on, (T), 26:46526 (TPJU-14/71)
- decay $M(953) \rightarrow \pi^+ + \pi^- + \gamma(\eta)$, branching ratios for, (E), 26:46471 (BNL-16954)
- decay $M(953) \rightarrow \rho^0 + \gamma$, branching ratio for, (E), 26:46471 (BNL-16954)
- decay modes and masses of $g(1660)$, $\phi(1650)$, $\phi(1830)$, and $K(1760)$, (T), 26:46611
- decay of N anti $N(1970)$, branching ratio for, (E/T), 26:46458
- decay of, comparison of excited quark, interacting quarks, and additive quark models for constraints on angular distributions for, (T), 26:46569
- effect of $\rho^0(1700)$ on π form factor, 26:46561
- existence of 0^- χ , from quartet model in $U(4) \times U(4)$, 26:46687
- existence of exotic, in inclusive interactions in Regge pole model, 26:46605
- mass and width of $M^0(940)$ and $\delta(963)$ in $\pi^- + p \rightarrow \pi^0 + n$ interactions at 2.4 GeV/c, scintillation counter study of, (E), 26:46475 (IS-2886)
- mass of $M(953)$ in $K^- + p \rightarrow K^- + p + \pi^+ + \pi^- + \gamma(\eta)$ interactions at 3.9 and 4.6 GeV/c, (E), 26:46471 (BNL-16954)
- parity and spin of $M(953)$, (E), 26:46471 (BNL-16954)
- parity of N anti $N(1970)$, (E/T), 26:46458
- production in $\gamma + N$ and $\gamma + \Sigma$ interactions in broken nonlinear chiral $SU(3) \times SU(3)$ dynamics, differential cross sections for vector and axial-vector, (T), 26:46563
- production in photon interactions with baryons, comparison of excited quark, interacting quark, and additive quark models for constraints on angular decay distributions in, (T), 26:46569
- production in π reactions with xenon at 2.34 to 9 GeV/c, (E), 26:46476 (JINR-P1-6408)
- production of (968) enhancement in $\pi^- + p \rightarrow n + \pi^0$ interactions at 2.5 GeV/c, (E), 26:46430 (COO-1195-241)
- production of exotic, in P /sub 3/2/ partial wave in phase shift analysis of $K^+ + p \rightarrow K^+ + p$ interactions at 0.52 to 1.89 GeV/c, (T), 26:46632
- production of $M(953)$ in $K^- + p \rightarrow K^- + p + \pi^+ + \pi^- + \gamma(\eta)$ interactions at 3.9 and 4.6 GeV/c, (E), 26:46471 (BNL-16954)
- production of $M^0(940)$ and $\delta(963)$ in $\pi^- + p \rightarrow \pi^0 + n$ interactions at 2.4 GeV/c, scintillation counter study of, (E), 26:46475 (IS-2886)
- production of R in $K^- + n \rightarrow \Lambda + \pi^+ + 2\pi^- + \pi^0$ interactions at 4.5 GeV/c in deuterium bubble chamber, cross sections for, (E), 26:46474 (COO-1428-306)
- properties of exotic, phase shift analysis of $K^+ + p \rightarrow K^+ + p$ interactions at 0.52 to 1.89 GeV/c for, (T), 26:46632
- spin distribution in dual resonance model, statistical mechanics calculation of, (T), 26:46609 (ITF-72-13-P)
- spin of N anti $N(1970)$, (E/T), 26:46458
- MESONS, RESONANCE (η)
- coupling constant for $\epsilon\pi\pi\gamma$, canonical anomaly for, (T), 26:46556
- decay $\epsilon(750) \rightarrow 2\gamma$, tests of Han--Nambu and Gell-Mann--Zweig triplet quark models for, (E), 26:46516 (DESY-72/21)
- decay $\eta'(958) \rightarrow 2\gamma$, tests of Han--Nambu and Gell-Mann--Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)
- mass of $\eta'(958)$ in $\pi^- + p \rightarrow \pi^0 + n$ interactions at 2.4 GeV/c, scintillation counter study of, (E), 26:46475 (IS-2886)
- production of $\epsilon(750)$, $\epsilon(1200)$, and $f(1260)$ in anti $p + p$ interactions at rest, (E), 26:46465
- production of $\eta'(958)$ in $\pi^- + p \rightarrow \pi^0 + n$ interactions at 2.4 GeV/c, scintillation counter study of, (E), 26:46475 (IS-2886)
- production of $\eta(958)$ in $\pi^- + p \rightarrow n + X$ interactions, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- MESONS, RESONANCE (ϕ)
- production of $\phi(1019)$ in $K^- + p \rightarrow K^- + p + \pi^+ + \pi^- + \gamma(\eta)$ interactions at 3.9 and 4.6 GeV/c, (E), 26:46471 (BNL-16954)
- MESONS, RESONANCE (π)
- coupling constant for $A_1\rho\omega$ from analysis of $\pi + N \rightarrow A_1 + N$ interaction, relation of t -channel helicity conservation with, (T), 26:46638
- coupling constants of $A_1\rho\omega$, use of collinear dispersion relations and equal-time commutators for, 26:46612
- decay $A_2 \rightarrow \pi^+ + 2\pi^-$ in deuterium bubble chamber, (E), 26:46474 (COO-1428-306)
- decay $A_3 \rightarrow f^0 + \pi^-$, bubble and spark chamber study of branching ratio for, (E), 26:46473 (COO-1195-239)
- parity and spin of A_2^- in $K^- + n \rightarrow \Lambda(\Sigma^0) + \pi^+ + 2\pi^-$ interactions at 4.5 GeV/c in deuterium bubble chamber, (E), 26:46474 (COO-1428-306)
- production of A_2 in $K^- + n \rightarrow \Lambda(\Sigma^0) + \pi^+ + 2\pi^-$ interactions at 4.5 GeV/c in deuterium bubble chamber, cross sections for, (E), 26:46474 (COO-1428-306)
- production of $A_2(1300)$ in anti $p + p$ interactions at rest (E), 26:46465
- production of A_2^+ in $\gamma + p$ interactions at 4.3 to 7.5 GeV, (E), 26:46446
- production of A_3 in $\pi^- + p \rightarrow \pi^+ + 2\pi^- + p$ interactions at 11 to 40 GeV in bubble and spark chambers, (E), 26:46473 (COO-1195-239)
- properties of A_3 , bubble chamber and spark chamber study of, (E), 26:46473 (COO-1195-239)
- Regge cuts of A_2 in forward $p + p \rightarrow p + p$ interactions inclusive finite mass sum rules for upper bound for, 26:46651 (RPP/T-18)
- MESONS, RESONANCE (ρ)
- production of $B(1250)$ and $\rho^0(1250)$ in anti $p + p$ interactions at rest, (E), 26:46465
- production of ρ^0 in $\gamma + p$ interactions at 4.3 to 7.5 GeV, (E), 26:46446
- MESONS, TENSOR
- production in $\gamma + N \rightarrow \text{baryon} + \text{tensor meson}$ interactions, $SU(6)$ symmetry for, (T), 26:46684
- MESONS, VECTOR
- decay angular distributions from $\gamma + p$ inclusive interactions, quark model predictions for, (T), 26:46536
- decay of, restrictions on symmetry-breaking relations for octet of transition operators in algebraic hadron model for rates for e^+ and e^- emission in, (T), 26:46525 (ORO-3992-80)
- production in $\gamma + N$ interactions in broken nonlinear chiral $SU(3) \times SU(3)$ dynamics, differential cross sections for, (T), 26:46563
- METAL COATINGS
- structure on intermediate layers in diffusion, review on, 26:46041
- METALLURGICAL EQUIPMENT
- description of WAK radiochemical plant, 26:46106(T) (ORNL-tr-2372)

design for measuring zirconium evaporation from open surfaces at high temperatures, 26:46011

METALLURGY

book: Constitution of Binary Alloys, Second Supplement, 26:45997

METALS

annealing of dislocation loops and tetrahedra in, analysis of mechanisms of, 26:45989 (CONF-710601-, pp 647-62)

book: Modern Analytical Techniques for Metals and Alloys. Part 2, 26:45187

book: Physicochemical Measurements in Metals Research. Pt. 2, 26:46015

book: Physicochemical Measurements in Metals Research. Pt. 1, 26:46014

chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278

coating with radiopolymerizable resins, method for, 26:46089(P)

conference on radiation effects on, Albany, N.Y., June 9 to 11, 1971, 26:46045 (CONF-710601)

crystal-field splitting effects of rare earth ions in, (T), 26:47269

defects in, ultrasonic method using frequency analysis system for locating and sizing, 26:45469(P)

diffusion of helium in, ion implantation studies of, 26:46050 (CONF-710601-, pp 230-47)

grain boundary models for, relations with free volume, 26:46027

level of electron beam-heated liquid, radiometric gage for measuring, 26:45655(P)

microscopy of bcc, model for studying symmetry of self-interstitial atoms in field ion, 26:46029

microscopy of, imaging process in field ion, 26:46038

ordering of point defects in, relations of piezoresistance with stress-induced, 26:46066(R) (COO-3431-1)

preparation of high-purity, 26:45311(R) (BARC-599)

radiation effects on atomic displacement in, relations of recoil energy with neutron, 26:46051 (CONF-710601-, pp 397-429)

radiation effects on hardening of, neutron, 26:46071

radioinduced atomic displacement cascades in, computer simulation of neutron, 26:46061 (CONF-710601-, pp 739-56)

radioinduced crystal lattice defects in, review on fast electron, γ , and neutron, 26:46076

radioinduced void formation in, analysis of mechanisms of, 26:46058 (CONF-710601-, pp 563-604)

radioinduced void formation in, CEA research programs on neutron, 26:46046 (CONF-710601-, pp 31-83)

radioinduced void formation in, models for fast neutron and ion, 26:46063 (CONF-710601-, pp 798-824)

reactions of injected molten, with water, mechanisms of explosive, 26:45461

recrystallization of, effects of purity on, 26:45982

solute segregation and stresses around growing voids in, during cooling from high-temperature anneals, 26:45988 (CONF-710601-, pp 630-46)

thermodynamic properties of liquid, design concepts of high-temperature systems for studying, 26:45437 (CONF-720904-1)

void formation in quenched, analysis of mechanisms of, 26:46058 (CONF-710601-, pp 563-604)

METEORITES

analysis for calcium and titanium by proton activation, 26:45215

analysis for calcium by bremsstrahlung activation, 26:45215

argon isotope abundances in Lost City, (E), 26:46155

radiation effects on, search for, 26:46156

METEOROLOGICAL INSTRUMENTS

criteria for reactor accident evaluation, 26:47317 (BNWL-1635)

METHACRYLIC ACID

2-methyl-2-nitropropyl ester, radioinduced polymerization of, γ , 26:45325

hydroxyalkyl esters, radioinduced graft polymerization on polymers, hydrogels for biological uses from γ , 26:45312(R) (COO-2147-2)

METHACRYLIC ACID, METHYL ESTER

kaolin-methyl methacrylate, radioinduced polymerization of composites of, γ , 26:45327

methyl methacrylate-styrene-vinylphosphonic dichloride, radioinduced polymerization of liquid and solid, reaction mechanisms in, 26:45318

METHANE

neutron spectra in liquid and solid, cold, (E), 26:46858

proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s hydrogen atoms by charge exchange and excitation in, (E), 26:46378

METHANE, BROMOTRIFLUORO-

radiolysis of liquid, yield of products from, 26:45322

bromotrifluoromethane-I, radiolysis of, yields of products from, 26:45322

METHANE, CHLOROTRIFLUORO-

performance as fire extinguishing agent, evaluation by gamma attenuation, 26:45712

METHANE, DICHLORO-

radioinduced reaction with cyclohexanone, yields from γ ,

26:45335

METHANE, IODO-

reactions with nitric acid, 26:45417(R) (ORNL-TM-3281)

reactions with nitric acid, 26:45414(R) (ORNL-TM-3180)

reactions with nitric acid, 26:45415(R) (ORNL-TM-3217)

separation from air by cold trapping, 26:45416(R) (ORNL-TM-3250)

separation from fuel reprocessing off-gas, 26:45411(R) (ORNL-TM-3072)

separation from fuel reprocessing off-gases by sorption on zeolites or mercury nitrate, 26:45407(R) (ORNL-TM-2918)

separation from gaseous streams by sorption and reaction on zeolites, 26:45387(P)

separation from gases by sorption on styrene-divinylbenzene copolymers, 26:45388

separation from gases, development of inorganic adsorbents for, 26:45405(R) (ICP-1006)

separation from off-gases in LMFBR fuel reprocessing, 26:45408(R) (ORNL-TM-2949)

sorption from gas streams on solids, 26:45415(R) (ORNL-TM-3217)

sorption on porous plastic, 26:45414(R) (ORNL-TM-3180)

sorption on solids, research progress on, 26:45416(R) (ORNL-TM-3250)

sorption on zeolites, 26:45410(R) (ORNL-TM-3018)

sorption on zeolites, 26:45413(R) (ORNL-TM-3127)

sorption on zeolites, 26:45414(R) (ORNL-TM-3180)

sorption on zeolites, 26:45411(R) (ORNL-TM-3072)

METHANE, NITRO-

effects on quenching of radioinduced excited states in naphthalene and pyrene in cyclohexane, 26:45331

effects on yields of excited states from pulse radiolysis of deaerated naphthalene-cyclohexane solutions, 26:45330

effects on yields of singlet state from pulse radiolysis of naphthalene and pyrene in benzene and cyclohexane solutions, 26:45329

METHANOL

effects on sorption of protactinium on silica gel from aqueous solutions of hydrochloric and nitric acids, 26:45389

METHYLDIENE

abundance in circumstellar shells in brightest members of Pleiades, (E), 26:46190

METROLOGY

quantum proposition systems for ideal measurements of first kind in, covering law in, 26:47294

MICA

particle track counting in, using capacitive discharge device, 26:46083(P)

MICE

chromium-51 retention in lungs of, following intratracheal injection as oxide, effects of respiratory infection on effective half-life for, 26:45910 (CONF-720505-2)

immunization against equine encephalitis virus, effectiveness of gamma irradiated virus for, 26:45752

plutonium-239 distribution in tissues of, following intravenous injection of monomeric and polymeric forms as citrate and nitrate, comparison of, 26:45917

radiation dose to thyroid gland of adult and fetal, following injection of iodine-131, mathematical model for calculating β , 26:45902

radiation effects on colony formation by rat bone marrow cells in spleen of, x, 26:45834

radiation effects on DNA metabolism in intestine of, comparison of RBE for gamma and proton, 26:45842

radiation effects on enzyme activity and RNA synthesis in spinal neurons of, 26:45868

radiation effects on heart rate and survival time of, effects of dose, dose rate, and phenobarbital on gamma, 26:45846

radiation effects on hematopoietic stem cells of, effects of cell-free tissue extracts on recovery from x, 26:45847

radiation effects on host, containing diffusion chamber cultures of bone marrow and blood cells, effects of, on yield of granulocytes and macrophages, 26:45741

radiation effects on iron metabolism in, following whole-body exposure, protective effects of AET and mexamine against mixed γ and neutron, tracer study using iron-59, 26:45840

radiation effects on lipid peroxidation in heart, kidneys, liver, and spleen of, following whole-body exposure, effects of dietary antioxidants on x, 26:45872

radiation effects on litter size and reproduction span of, following chronic ingestion of cesium-139 and strontium-90 in drinking water, 26:45900

radiation effects on lymphomagenic properties of thymus gland of neonatal, subsequently transplanted into Gross murine leukemia virus-infected hosts, γ , 26:45818 (UCLA-12-867)

radiation effects on protein synthesis in spleen cell nuclei of, following whole-body exposure, comparison of effects of Friend virus infection with x, 26:45849

radiation effects on regeneration of bone marrow lymphoid cells in, effects of thymectomy on x, 26:45830

radiation effects on regeneration of lymphoid cells in bone marrow of germ free, x, 26:45843

radiation effects on response of phagocytes of, to intraperitoneal inflammation, effects of time of

- exposure on x, 26:45833
radiation effects on stem cell renewal in bone marrow and spleen of, following fractionated whole-body exposure, x, 26:45871
radiation effects on survival of hypoxic, following whole-body exposure, sensitizing effects of TAN on electron, 26:45869
radiation effects on survival of, following whole-body exposure, comparison of RBE for cobalt-60 γ and pulsed x, 26:45866
radiation effects on urinary content of 5-hydroxyindole acetic acid, volume of urine excreted, and water consumption of, following partial or whole-body exposure, x, 26:45826
radioinduced leukemia and solid tumors in, following perinatal exposure, effects of age and strain on incidence of, 26:45814(R) (COO-3017-43)
radioinduced leukemia in, in vitro culture of cells from x, 26:45855
radioisotope uptake and retention by, following gastric tube administration, effects of aluminum phosphate gel on calcium-47, radium-226, and strontium-85, 26:45918
radiosensitivity of thyroid gland in goitrogen-stimulated adult and normal fetal, comparison of iodine-131 β and γ and external x, 26:45865
radium-226 retention in, following injection, method for direct measurement of, 26:45920
radon-222 distribution in tissues of, following injection of radium-226, kinetics of, 26:45920
strontium-89 effects on bone marrow of, following intraperitoneal injection, relation to hematopoiesis and immune reactions, 26:45896
- MICROORGANISMS**
radioinduced mutations for radioresistance in spores of, following exposure during sterilization, probability of, 26:45750
radiosensitivity of, contaminating disposable medical supplies, method for determining, 26:45748
- MICROWAVES**
cosmic background, interaction with 10^{20} -eV cosmic particles, (T), 26:46257
cosmic background, isotropy at 6 cm, (E), 26:46255
effects of, thyroid--pituitary pathophysiologic, 26:47648 (UR-3490-137)
reflection through laboratory plasmas, boundary parameters from, (E), 26:47118
- MIDWEST FUEL RECOVERY PLANT**
licensing consideration and hearing notice, 26:45402 (DOCKET-50268-31)
radioactive waste processing at, proposed operation of evaporation system for low-level liquid, 26:45404 (DOCKET-50268-33)
uranium cold run testing at, request for use of service steam system in, 26:45403 (DOCKET-50268-32)
- MILK**
analysis of powdered, by neutron activation and radiochemical separation, 26:45196
fallout cesium-137 and strontium-90 content in, Mecklenburg, GR, 1969, 26:45546
iodine-131 content of, from off-site environs of Nevada Test Site during July through Dec. 1971, 26:45515
radioactivity monitoring of, in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
radioisotope content in Western Hemisphere, during Mar. 1971 to Mar. 1972, cesium-137 and strontium-90, 26:45552
strontium-90 content in, in New York City, June 1954 through March 1972, 26:45512(R) (HASL-257(App.))
strontium-90 content in, Ispra, Italy, 1960 through 1971, 26:45519 (HASL-257, pp IIL53-8)
- MILKY WAY**
clouds near center of, kinematics of molecular, (T), 26:46251
gamma background in, origin of, (E), 26:46154
hydrogen in dust clouds in, density and temperature of neutral, (E), 26:46247
hydrogen motion in spiral arms and local region, neutral, (E), 26:46246
interstellar cloud formation in, mechanism for, (T), 26:46245
positions of bright nebulae in, relation to high-velocity gas concentrations, (E), 26:46250
regions of faint H α emission, detection of two new large-diameter, (E), 26:46236
star birth and death processes in, survey of, (T), 26:46166
- MILLSTONE POINT POWER REACTOR, UNIT 1**
containment structure for, revised leak rate test requirements for primary, 26:47410 (DOCKET-50245-144)
control rod drives for, maintenance procedures for, 26:47411 (DOCKET-50245-145)
electric power source for, additional data on 4.16 kV auxiliary bus system connection to reserve off site, 26:47408 (DOCKET-50245-140)
instrumentation for, incorrect trip times for delay relays of average power range monitoring system, 26:47360 (DOCKET-50245-143)
operating procedures for, revised, 26:47409 (DOCKET-50245-141)
operations during January to June 1972, 26:47361 (DOCKET-50245-146)
personnel for, revised organization of, 26:47409
- (DOCKET-50245-141)
pressure vessel high pressure scram switches for, incorrect trip points for, 26:47407 (DOCKET-50245-139)
- MILLSTONE POINT POWER REACTOR, UNIT 2**
control systems for, final safety analysis report on, 26:47509 (DOCKET-50336-37)
cooling system for, final safety analysis on, 26:47508 (DOCKET-50336-36)
design criteria for, AEC, 26:47508 (DOCKET-50336-36)
electrical systems for, final safety analysis report on, 26:47509 (DOCKET-50336-37)
engineered safeguards for, final safety analysis report on, 26:47509 (DOCKET-50336-37)
instrumentation for, final safety analysis report on, 26:47509 (DOCKET-50336-37)
kinetics for, final safety analysis on, 26:47508 (DOCKET-50336-36)
owners of, financial data on, 26:47507 (DOCKET-50336-35)
pressure vessel internals for, stress analysis for, 26:47508 (DOCKET-50336-36)
quality assurance program for, 26:47508 (DOCKET-50336-36)
site for, final safety analysis on, 26:47508 (DOCKET-50336-36)
structures for, final safety analysis on, 26:47508 (DOCKET-50336-36)
- MINERALS**
analysis for potassium by neutron activation, 26:45197
radiation effects on crystal structure of, heavy-ion, 26:46156
- MITOCHONDRIA**
radiation effects on content of thiol groups in, of rat heart and liver, 26:45829
radiation effects on function and structure of rat liver, following lethal whole-body exposure, 26:45838
- MIXER-SETTLERS**
design for simultaneous mixing and separation of multi-phase fluids, 26:45451(P)
- MOESSBAUER EFFECT**
absorber thickness optimization for single-line, in presence of strong non-resonant absorption, 26:47274
background in transmission geometry using iron alloys with 3d-transition elements, 26:47282
pressure dependence of recoilless fraction and second-order Doppler shift of iron-57 14.4-keV gamma ray in host lattices, (T), 26:47276
spectral analysis of five-line hyperfine split absorbers by fitting sums of Lorentzians to numerical solutions of transmission spectra, 26:47275
- MOESSBAUER SPECTROMETERS**
description for diffractometry, 26:45683
- MOLECULAR PROPERTIES**
research programs at Florida State University, Tallahassee, during 1971 and 1972, 26:45891(R) (ORO-2690-12(Pt.B))
- MOLECULES**
abundances in interstellar clouds, (E/T), 26:46244
atom reactions with diatomic, comparison of S matrices calculated by various exponential approximations for rotationally inelastic, 26:46369
atom reactions with diatomic, optical-model calculations of probabilities for vibrational transitions in, 26:46375
atom reactions with, Monte Carlo trajectory calculations of three-body recombination and dissociation in, 26:46382
electron elastic scattering by, Glauber-theory calculations of amplitudes and differential cross sections for multiple, 26:46381
electron reactions with, third-derivative method for determination of ionization and appearance potentials for, (E/T), 26:46385
force constants for, utilization of very heavy atom isotopic frequency data for calculating, 26:45299
interactions between, inversion method for calculation of potentials for, from scattering cross sections, 26:46387
kinematics of clouds of, near galactic center, (T), 26:46251
properties of, infinite-order perturbation theory for calculation of, 26:46410
scattering of, bound states in quantum electrodynamics for, (T), 26:46721
separation of heavy and light, from gaseous mixtures, supersonic gas jet method for, 26:45692(P)
- MOLLUSCA**
Crassostrea gigas, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
fallout content in north Atlantic Ocean, ecological relations for cesium-137, plutonium-238, -239, and -240, and strontium-90, 26:45541 (NYO-2174-130)
Mya arenaria, radioisotope accumulation and loss by, model for food chains from kinetics of cesium-137 and cobalt-60, 26:45545 (UCRL-73744)
Mytilus californianus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
Mytilus californianus, zinc-65 content in Columbia River, following shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)
Mytilus edulis, lead-210 and polonium-210 content in, in

Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
 Mytilus edulis, zinc-65 content in Columbia River, following shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)

MOLTEN SALT BREEDER REACTOR
 development of, review of, 26:47302

MOLTEN SALT REACTOR EXPERIMENT
 operation of, review of, 26:47302

MOLYBDENUM
 argon-ion reactions at low energies with monocrystals of, mechanism of ion capture by surface in, (E/T), 26:46335 (FRNC-TH-198)
 brittle--ductile transition in sheets of, transition temperatures for, 26:45993(T) (ORNL-tr-2377)
 chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
 deformation of monocrystals of, during dynamic loading, effects of impurities and temperature on, 26:46043
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 determination by catalytic effect on Methylene Blue reaction with hydrazine sulfate, spectrophotometric, 26:46227
 determination by electron probe microanalysis methods, 26:45237
 determination by spectrography, spectrochemical buffers and volatilization mechanisms in, 26:45222 (JEN-229-DQ/I-79)
 determination of trace amounts of, in atmospheric particulates by atomic absorption and flame emission spectroscopy, 26:45244
 determination of, chemical procedures for, 26:45181 (CISE-R-172)
 determination using thiolactic acid, spectrophotometric, 26:45239
 dislocation energy of impurity atoms in solid solutions of, internal friction method for studying, 26:46023
 effects on microstructure and properties of nickel-base--niobium alloy, 26:46035
 effects on recrystallization of iron, 26:45982
 effects on structure of cobalt--iron--tungsten alloys during aging, 26:46006
 neon-ion reactions at low energies with monocrystals of, mechanism of ion capture by surface in, (E/T), 26:46335 (FRNC-TH-198)
 neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)
 radiation effects on length of, γ , 26:46075
 radiation effects on void lattice formation in, 2-MeV nitrogen ion, 26:46055 (CONF-710601-, pp 522-32)

MOLYBDENUM ALLOYS AND SYSTEMS
 decomposition of supersaturated solid solutions of, effects of aging on mechanism of, 26:46032
 Al_2O_3 --Mo, brittle--ductile transition in dispersions of, transition temperatures for, 26:45993(T) (ORNL-tr-2377)
 Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009
 Al--Fe--Mo--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
 Al--Mo--Si--U, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
 Mo--Nb, reactions with boron at 1100 to 1400°C, effects of composition on, 26:46002
 Mo--Nb-base--W, tensile properties at 900°K, athermal, 26:45991(R) (COO-916-25)
 Mo--Pt, superconducting phases in, identification of, (E), 26:46828
 Mo--ThO₂, porosity and tensile properties of test fuels of, 26:45703(R) (ORNL-4796)
 Mo--U-base, mechanical properties of, at 20 to 640°C, effects of aluminum, iron, and silicon additions on, 26:45995 (UJV-2802-M)
 Mo-base--Re, superconducting properties of, effect of dislocation tangles on, (T), 26:46836

MOLYBDENUM CARBONYLS
 compaction of powdered, electrodynamic method for, 26:45951

MOLYBDENUM ISOTOPES Mo-92
 electron inelastic scattering at 209 MeV, differential cross sections for, p. 189-91, (E), 26:46873(R) (CEA-N-1522)
 energy level at 2761 keV, g-factor for, (E), 26:46959
 energy levels of, unpaired-nucleon collective, (T), 26:46965
 helium-3 reactions (He-3,d) at 30.2 MeV, DWBA analysis of differential cross sections and energy spectrum from, p. 110-12, (E), 26:46873(R) (CEA-N-1522)
 helium-3 reactions (He-3,d) at 35 MeV, diffraction-model analysis of differential cross sections for, (T), 26:47075
 photon reactions (γ ,n) and (γ ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

MOLYBDENUM ISOTOPES Mo-94
 energy levels of, unpaired-nucleon collective, (T), 26:46965
 helium-3 reactions (He-3,d) at 30.2 MeV, energy spectrum from, p. 110-12, (E), 26:46873(R) (CEA-N-1522)

MOLYBDENUM ISOTOPES Mo-96

energy levels from neutron capture by molybdenum-95, (E), 26:46917(T) (BNL-tr-477)

MOLYBDENUM ISOTOPES Mo-97

neutron capture at thermal energies and 2 and 25 keV, gamma spectrum from, (E), 26:46917(T) (BNL-tr-477)

MOLYBDENUM ISOTOPES Mo-98

photon reactions (γ ,n) and (γ ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

MOLYBDENUM ISOTOPES Mo-99

range of fission fragments of, in aluminum--uranium alloys and uranium oxides, 26:46074

MOLYBDENUM OXIDES

Al_2O_3 --CaO--Cr₂O₃--F--MgO--MoO₃--TiO₂--ZrO₂, use as welding flux, 26:45941(P)
 Al_2O_3 --MoO₃, phase studies on, 26:45963

MONKEYS

radiation detection by eyes of anomic, γ , 26:45817 (ORO-2690-12(Pt.B), Paper 12)
 radiation injuries to central nervous system of, following local exposure, x, 26:45850
 radioinduced decrease in body weight of, effects of age, dose, energy, and sex on proton, 26:45841
 radioinduced injuries in visual cortex of, latent x, 26:45858

MONTICELLO POWER REACTOR, UNIT 1

core power range monitoring system for, calibration of average, 26:47323 (DOCKET-50263-141)
 emergency core cooling system for, failure of coolant injection turbine valve in, 26:47333 (DOCKET-50263-142)
 emergency core cooling system for, failure of high pressure coolant injection, 26:47368 (DOCKET-50263-139)
 emergency core cooling system for, fast start failure of high pressure coolant injection system in, 26:47334 (DOCKET-50263-143)
 residual heat removal system for, revised service water pump head requirements for, 26:47429 (DOCKET-50263-138)
 safety relief valve for, malfunction of steam line, 26:47369 (DOCKET-50263-140)
 safety relief valves for, failure of bellows leakage monitoring systems for, 26:47369 (DOCKET-50263-140)
 turbine for, unplanned trip at 100 percent power, 26:47367 (DOCKET-50263-137)

MOON

magnetic fields on, possible production mechanisms for, (E/T), 26:46262
 power stations for use on, description of laser-energized, 26:45428

MOSSES

radioisotope content in land, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)

MOUTH

radioinduced necrosis in soft tissues of, in man following therapy of oral tumors, 26:45801

MUCOSA

permeability of lingual, in rabbits during development and recovery from radioinduced glossitis, 26:45862
 radioinduced edema of human laryngeal, following therapy of laryngeal carcinomas, incidence of x, 26:45781(T) (TT-54002/4, pp 437-41)

MUONIC ATOMS

x-ray emission in mercury and thallium-203, test of quantum electrodynamic corrections from, 26:46708

MUONIUM

hyperfine splitting in, 26:46710

MUONS

decay of, radiative corrections to, 26:46576
 g-factor of, review and analysis of data on ratio to electron for, (E), 26:46439
 interactions in nuclear emulsions at 6 and 12 GeV, cross sections for inelastic, (E), 26:46451
 interactions $\mu + N \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV, momentum distributions of π in, (E), 26:46452
 magnetic moment of, quantum electrodynamic contribution to anomalous, 26:46710
 magnetic moment of, review and analysis of data on relation for, (E), 26:46439
 magnetic moment of, second-order weak corrections to anomalous, (T), 26:46553
 mass of, cosmological significance of difference of electron mass and, (T), 26:46259
 mass of, review and analysis of data on ratio to electron for, (E), 26:46439
 spectra of cosmic, bounds on production of μ^+ -- μ^- pairs in super-high energy hadron interactions from, (T), 26:46598
 stopping underground of atmospheric, pion-decay-produced, and neutrino-interaction-produced, (T), 26:46157

MUONS (μ^-)
 capture by oxygen-16, silicon-28, sulfur-32, and calcium-40, particle-hole calculation of capture-rate and neutron spectra for, (T), 26:46880
 production of heavy μ^+ - μ^- pairs in semi-inclusive $e^+ + p \rightarrow \mu^+ + \mu^- + \text{anything}$ interactions, review of dual quark-parton model for, (T), 26:46515 (DAMP-71/24)
 production of μ^+ -- μ^- pair in super-high energy hadron interactions, bounds on cross sections for, (T), 26:46598
 range--momentum correlation in graphite and iron for 500-

- to 1500-MeV/c, (E), 26:46448 (LBL-791)
 reactions (μ^- , e^+) with copper, upper limit on branching ratio relative to ordinary capture for, (E), 26:46941
- MUONS (μ^+)**
 production of heavy $\mu^+ - \mu^-$ pairs in semi-inclusive $e^+ + p + \mu^+ + \mu^-$ + anything interactions, review of dual quark-parton model for, (T), 26:46515 (DAMTP-71/24)
 production of $\mu^+ - \mu^-$ pair in super-high energy hadron interactions, bounds on cross sections for, (T), 26:46598
 range-momentum correlation in graphite and iron for 500- to 1500-MeV/c, (E), 26:46448 (LBL-791)
- MUSCLES**
 cesium-137 content in, of broilers, hens, and partridges, Warsaw, Poland, 1968 to 1969, 26:45548
 radiation effects on histochemistry of, of frogs, gamma, 26:45828
 radiation effects on regenerative capacity of rat, following traumatic injury, 26:45827
- MUTATIONS**
 radioinduced detrimental and lethal, in wasp sperm, effects on fecundity and life span of F_1 females, x, 26:45878
 radioinduced, for radioresistance in spores of microorganisms following exposure during sterilization, probability of, 26:45750
 radioinduced, in cultured human fibroblasts, effects of dose on frequency of x, 26:45740
 radioinduced, in *Drosophila* following ingestion of carbon-14, phosphorus-32, or phosphorus-33, beta, 26:45912(R) (ORO-3728-5)
 radioinduced, in extracellular bacteriophages, effects of LET on frequency of γ , heavy ion, and proton, 26:45746
 radioinduced, in Japanese cedars, for albinism, gamma, 26:45771
 radioinduced, in peas, review of, 26:45776
 radioinduced, in potato plants for protein and starch contents and yield, fast neutron and γ , 26:45770
- MYLAR**
 radioinduced electric conductivity of, effects of dose rate of pulsed γ - and x-, 26:47225 (AWRE-O-31/72)
- N**
- NAGASAKI**
 muscular reflex time in residents of, age dependence of, 26:45795
- NAPHTHALENE**
 radioinduced fluorescence in benzene and cyclohexane solutions, effects of concentration on, 26:45328
 radioinduced fluorescence in cyclohexane, effects of scavengers on electron pulse, 26:45331
 radiolysis in benzene and cyclohexane solutions, effects of intersystem crossing from excited singlet state on yield of triplet state from pulse, 26:45329
 radiolysis in benzene and cyclohexane solutions, effects of concentration on yield of excited triplet states from pulse, 26:45328
 radiolysis in benzene or cyclohexane solutions, yields of excited singlet state solute molecules from pulse, 26:45328
 radiolysis of deaerated cyclohexane solutions of, effects of scavengers on yield of excited states from pulse, 26:45330
- 2,7-NAPHTHALENEDISULFONIC ACID, 4,5-DIHYDROXY-3-[(4-PHENYL-2-THIAZOLYL)AZO]-**
 uses in photocolometric determination of thorium, 26:45240
- 2-NAPHTHALENESULFONIC ACID, 1,4-DIHYDRO-1,4-DIOXO-**
 reactions with hydrated electrons and hydrogen atoms from radiolysis of aqueous solutions, 26:45354
- 4-NAPHTHALENESULFONIC ACID, 1,2-DIHYDRO-1,2-DIOXO-**
 reactions with hydrated electrons from pulse radiolysis of aqueous solutions, 26:45354
- 1,4-NAPHTHOQUINONE**
 electron reactions with, cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- NATURAL BACKGROUND RADIATION**
 distribution in ground-level atmosphere, effects of meteorological conditions on, 26:45524
 levels in atmospheric aerosols in Poland, relations with aerosol dust content, (E), 26:45521
- NATURAL GAS**
 production stimulation by Project Wagon Wheel, prediction of explosion effects on surrounding sandstone from, 26:45494
 production stimulation using underground nuclear explosions, preliminary evaluation of Wagon Wheel site for testing, 26:45571 (NVO-55)
 storage in underground nuclear cavities, feasibility of, 26:45493
- NEBULAE**
 distance of eta Carinae, (T), 26:46160
 hydrogen in dust, density and temperature of, (E), 26:46247
 isophote chart of eta Carinae, (T), 26:46160
 positions of bright, relation to concentrations of high-velocity gas, (E), 26:46250
- NECROSIS**
 radioinduced, in human mandible and soft oral tissues following therapy of oral tumors, 26:45801

NEMATODA

radioisotope content in, in Puget Sound basin area, 26:45544 (RLO-2225-T-14-4)

NEODYMIUM

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 effects on grain boundary energy of nickel, 26:45999
 separation from europium, gadolinium, praseodymium, and samarium by anion exchange using acetic acid-methanol-nitric acid mixtures, 26:45280
 solubility in oxygen-containing chromium at 1000 to 1800°C, microhardness studies on, 26:46034
 solvent properties of oxygen-containing, for cerium, neodymium, and praseodymium at 1000 to 1800°C, microhardness studies on, 26:46034

NEODYMIUM ISOTOPES

x-ray spectra from, isotope shifts in, (E), 26:46973

NEODYMIUM ISOTOPES Nd-141

energy-level scheme from promethium-141 decay, (E), 26:46972

NEODYMIUM ISOTOPES Nd-144

energy levels from neutron resonances in neodymium-143, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

NEODYMIUM ISOTOPES Nd-145

proton inelastic scattering at 8 to 14 MeV, cross sections for, p. 34-5, (E), 26:46873(R) (CEA-N-1522)

NEODYMIUM ISOTOPES Nd-146

energy levels from neutron resonances in neodymium-145, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

NEON

abundance in solar-flare particles, (E), 26:46232
 electron reactions in eV range with, trapped-electron measurements of integral cross sections for production of secondary electrons of 0 to 1 eV by ionization in, 26:46390
 electron reactions with clusters of, multiple ionization in, (E), 26:46386
 electron scattering by, bound-state calculation of resonance due to Ne^- ion in, 26:46383
 interactions with argon and neon atoms, retarded van der Waals potential for, as function of interatomic distance, 26:46357
 plasma of, wave coupling in afterglows, (E), 26:47193
 reactions with argon at 50 to 600 eV of metastable atoms of, cross sections for de-excitation in, (E), 26:46394

NEON IONS

effects of 10.4-MeV/amu, on crystal structure of minerals, 26:46156
 reactions with cobalt, iron, tin, and zirconium at 5 MeV/amu, energy shift and relative intensity of K x rays from, (E), 26:46339
 reactions with molybdenum monocrystals at low energies, mechanism of ion capture by surface in, (E/T), 26:46335 (FRNC-TH-198)
 wave functions for $2p^3s^2 \ 2P$ state of Ne^- , 26:46383

NEON ISOTOPES

meson (π) reactions with, research at Duke Univ., July 1971 to June 1972, (E), 26:46479(R) (ORO-3065-24)

NEON ISOTOPES Ne-18

energy levels of, data compilation on, 26:46882

NEON ISOTOPES Ne-19

energy levels of, data compilation on, 26:46882

NEON ISOTOPES Ne-20

energy levels of, data compilation on, 26:46882
 lithium-7 reactions ($Li-7, \alpha$) at 22 MeV, angular distributions for, (E), 26:46915
 nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892
 nuclear quadrupole moment rearrangement due to proton removal, calculation of, (T), 26:46886
 proton separation and orbital rearrangement energies, calculations of, (T), 26:46886
 reactions with silver-107 at 90 to 210 MeV, calculation of fusion excitation function using equilibrium model with fission competition, (T), 26:47072

NEON ISOTOPES Ne-21

energy levels from oxygen-18 ($\alpha, n\gamma$) reactions, lifetimes and spin-parity assignments for, (E), 26:46907
 energy-level transitions from oxygen-18 ($\alpha, n\gamma$) reactions, multipole mixing ratios for, (E), 26:46907
 helium-3 reactions ($He-3, p$) at 18 MeV, angular distributions for, (E), 26:46915

NEON ISOTOPES Ne-22

helium-3 reactions ($He-3, d$) at 15 MeV, angular distributions for, (E), 26:46915

NEOPLASMS

carcinomas, radioinduced squamous cell, in eyelids of cattle, comparison with human xeroderma pigmentosum, uv, 26:45851
 carcinomas, radioinduced uterine, in parabiont and single rats following focal internal exposure, cobalt-60 γ , 26:45814(R) (COO-3017-43)
 carcinomas, radioinduced, in man following exposure of pelvis, survey of x, 26:45807
 carcinomas, radiotherapy of laryngeal, effects of clinical stage and histologic type of tumor on effectiveness of x, 26:45780(T) (TT-54002/4, pp 430-6)
 fibrosarcomas, radioinduced, in cats following chronic

- ingestion of strontium-89, dose dependence of β , 26:45901
- hemangiosarcomas, radioinduced, in cats following chronic ingestion of strontium-89, dose dependence of β , 26:45901
- leukemia, radioinduced, in man following exposure during diagnostic radiography, effects of age, exposure site, number of exposures, and sex on incidence of x , 26:45798
- leukemia, radioinduced, in mice following perinatal exposure, effects of age and strain on incidence of, 26:45814(R) (COO-3017-43)
- leukemia, radioinduced, in rats, lymphoid cell proliferation kinetics in thymus during development of x , 26:45822
- leukemia, transplantation of radioinduced canine myelomonocytic, into canine fetuses, hematologic changes in peripheral blood of fetuses following serial, 26:45897
- localization using cobalt-57 labeled bleomycin, 26:45929
- lymphomas, single cell transplants of Gross virus-induced mouse, 26:47629 (UCLA-12-866)
- osteosarcomas, radioinduced, in cats following chronic ingestion of strontium-89, dose dependence of β , 26:45901
- radioinduced, in animals following inhalation or injection of plutonium-239, dose and time dependence of, 26:45888 (CONF-720520-1)
- radioinduced, in mammals following radioisotope uptake, review of pathology of, 26:45895 (UR-3490-102)
- radioinduced, in man following uptake of plutonium radioisotopes, relevance of animal data for estimation of dose and time relationships for, 26:45888 (CONF-720520-1)
- radioinduced, in parabiont rats following acute supralethal whole-body exposure, high incidence of, 26:45814(R) (COO-3017-43)
- radioinduced, in thyroid gland of dogs, combined iodine-131 β and γ and external x , 26:45819 (UR-3490-63)
- sarcomas, radioinduced osteogenic, in parabiont and single rats following focal internal exposure, cobalt-60 γ , 26:45814(R) (COO-3017-43)
- tumors, radioinduced giant cell, in bones and soft tissues adjacent to bones in cats following chronic ingestion of strontium-89, dose dependence of β , 26:45901
- tumors, radioinduced solid, in mice following perinatal exposure, effects of age and strain on incidence of, 26:45814(R) (COO-3017-43)
- NEPTUNIUM CHLORIDES
- magnetic properties of, 26:47272
- magnetic properties of $NpCl_3$, 26:47272
- NEPTUNIUM ISOTOPES Np-231
- electron-capture decay of, gamma spectra from, (E), 26:47023 (KFK-1537)
- NEPTUNIUM ISOTOPES Np-232
- electron-capture decay of, gamma spectra from, (E), 26:47023 (KFK-1537)
- NEPTUNIUM ISOTOPES Np-233
- electron-capture decay of, gamma spectra from, (E), 26:47023 (KFK-1537)
- NERVES
- radiation effects on histochemistry of, of frogs, gamma, 26:45828
- NERVOUS SYSTEM
- radiation injuries to central, of monkeys following local exposure, x , 26:45850
- NEUTRINOS
- emission from supergiant stars, (T), 26:46177
- exchange of pairs of virtual, in anti $l + l$ interactions, consequences, spin, and coupling to free leptons for massless composite lepton-antilepton states from, (T), 26:46570 (IC-72/54)
- interactions at accelerator energies, review on, 26:46584
- interactions $\nu + d + \mu^- + p + p$ in impulse approximation, differential cross sections for, (T), 26:46581
- interactions $\nu + l$, lepton charge theory for, (T), 26:46578
- interactions $\nu + N$, $\Delta(1236)$ production in, 26:46584
- interactions $\nu + N$, parton models with integral charge for deep-inelastic, 26:46574
- interactions $\nu + N$, review of dual quark-parton model for structure functions and total cross sections for, (T), 26:46515 (DAMTP-71/24)
- interactions $\nu + p$ in propane bubble chamber, test of PCAC from differential cross sections for inelastic, (E/T), 26:46450
- interactions with spherically symmetric gravitational fields, Weyl equation for, 26:46772
- production of beams of, at proton accelerators, spectra and flux of, 26:46584
- reactions with carbon in propane bubble chamber, test of PCAC from differential cross sections for inelastic, (E/T), 26:46450
- NEUTRINOS (ANTI ν)
- interactions anti $\nu + N$, parton models with integral charge for deep-inelastic, 26:46574
- NEUTRINOS (ν /sub e/)
- interactions $\nu + \text{anti } \nu + e^- + e^+$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions for, (T), 26:46579
- interactions $\nu + e + \nu + e$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions for, (T), 26:46579
- interactions $\nu + \text{anti } \nu + \mu^- + e^+$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions for, (T), 26:46579
- interactions $\nu + e + \text{anti } \nu + e(\mu^-)$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions and energy dependence of ν energy loss and scattering angle in, (T), 26:46579
- NEUTRINOS (ν /sub μ /)
- interactions $\nu + e + \nu + \mu^-$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions and energy dependence of ν energy loss and scattering angle for, (T), 26:46579
- interactions $\nu + n + \nu + n + \nu^0$ in optical spark chamber, upper limit on neutral current in, (E), 26:46453
- interactions $\nu + n + p + \mu^-$, upper bound for tensor component of second-class current in, (T), 26:46573
- interactions $\nu + p + \nu + p + \nu^0$ in optical spark chamber, upper limit on neutral current in, (E), 26:46453
- interactions $\nu + p + \nu + p + \nu^0$ in Weinberg model, bound on cross sections for, 26:46614
- NEUTRINOS (anti ν /sub μ /)
- interactions $\nu + \text{anti } \nu + \mu^- + e^+$, relations among differential and total cross sections of nonzero-mass $\nu + l$ interactions for, (T), 26:46579
- NEUTRON CROSS SECTIONS
- determination of effective, least squares method for, 26:47595
- measurements of total, survey on, 26:46967
- measurements on capture and scattering, review on, 26:46968
- NEUTRON-DIFFRACTION ANALYSIS
- use to study structure of condensed matter, 26:47251
- NEUTRON SOURCES
- description of pad-mounted, for well logging, 26:45562(P)
- description using Van de Graaff accelerator and time-of-flight techniques, 26:46848
- fabrication of californium-252, description of method for, 26:45714 (CONF-720902-2)
- neutron flux amplification in subcritically-enhanced californium-252, 26:46855
- uses of subcritically-enhanced californium-252, 26:46855
- NEUTRON SPECTROMETERS
- control system for crystal, for inelastic scattering studies, 26:45636 (BARC-591)
- description of high-resolution time-of-flight, 26:46848
- description of ORELA modified time-of-flight, 26:46144 (CONF-720607-22)
- performance of pulsed accelerator time-of-flight, review of, 26:45641
- NEUTRON SPECTROSCOPY
- book: Experimental Neutron Resonance Spectroscopy, 26:46966
- NEUTRON STARS
- characteristics of, review of, 26:46466
- equation of state of, (T), 26:46170
- models for, (T), 26:46170
- NEUTRONS
- absorption resonance integral for americium-243 at 0.625 to 1000 eV, (E), 26:47010 (ANCR-1060)
- attenuation by iron and water, (E/T), 26:46852
- capture by nuclei, $N-Z$ dependence of cross sections for, (T), 26:47085
- capture by nuclei, review on γ spectra from resonance, 26:46969
- detection of, design and performance of NE 323 scintillation detector for, p. 267-8, 26:46873(R) (CEA-N-1522)
- diffusion in beryllium and graphite, temperature dependence of diffusion length for, (T), 26:46857
- distribution at large distances from source, generalized method for calculation, 26:47215
- dose to man, evaluation of conservative and realistic quality factors used in expressing recommended limits for, 26:45789
- dose to medical personnel during use of californium-252 for interstitial and intracavitary therapy, methods for reducing, 26:45790
- dosimetry of, calibration of Bonner sphere $^6LiI(Eu)$ scintillation detector for, 26:45599
- dosimetry of, characteristics of fission foil--mica track detectors for, 26:45617 (FEI-296)
- dosimetry of, design of tissue-equivalent liquid ionization chamber for, 26:45621
- dosimetry of, performance of Cs and dielectric track detectors for, 26:45630
- effects on atomic displacement cascades in metals, relations with recoil energy, 26:46051 (CONF-710601-, pp 397-429)
- effects on crystal lattice defect formation in alloys and metals, review on, 26:46076
- effects on density of reactor alloys and metals, relations with stress, 26:46065 (CONF-710601-, pp 841-

effects on hardening of structural metals, 26:46071
 effects on high temperature properties of graphite and Hastelloy N, 26:47302
 effects on internal friction of aluminum wires at room temperature, 26:46073
 effects on iron metabolism in mice following whole-body exposure, protective effects of AET and mexamine against pathological mixed γ and, tracer study using iron-59, 26:45840
 effects on man, contribution of nuclear interactions with tissues to, 26:45789
 effects on mechanical properties of Type 304 stainless steel, relations with titanium additions, 26:46068 (ORNL-TM-3733)
 effects on microwave bipolar transistors, (E/T), 26:45663 (SC-DC-721233)
 effects on oxide fuels, 26:45412(R) (ORNL-TM-3095)
 effects on oxide fuels, 26:45411(R) (ORNL-TM-3072)
 effects on oxide fuels, 26:45409(R) (ORNL-TM-2996)
 effects on p-n junctions, carrier trapping, (T), 26:45664 (SC-DC-721237)
 effects on plutonium dioxide-uranium dioxide fuels, 26:45408(R) (ORNL-TM-2949)
 effects on rabbits, effects on crystalline lens of, effects of dose on RBE for mixed γ and, in comparison with x radiation, 26:45864
 effects on seeds of tomatoes, effects of dose and moisture on rate of germination and plant growth, 26:45773
 effects on Type A302B steel, activation fluence method for analysis of, 26:46072
 effects on void formation in metals, CEA research program on, 26:46046 (CONF-710601-, pp 31-83)
 effects on void formation in nickel and steel, comparison of experimental and theoretical, 26:46062 (CONF-710601-, pp 769-97)
 emission from 24-kt nuclear explosions, calculation of time-dependent, 26:45497
 emission of bursts of, following solar flare, identification by measuring trapped-neutron-decay protons in radiation belts, 26:46320
 flux distribution in reactors, FLUB code for analysis of, 26:47594 (BMBW-FBK-72-6)
 interactions $K^+ + n \rightarrow K^0 + p$ with octet exchange in crossed channel, models for extrapolation of differential cross sections from experimental points for, (T), 26:46630
 interactions $K^+ + n \rightarrow K^0 + p$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642
 interactions $K^+ + n \rightarrow K^0 + p$, modified absorptive Regge cut model for, (T), 26:46622
 interactions $K^+ + n \rightarrow K^0 + p$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
 interactions $K^- + n \rightarrow \Lambda + \pi^+ + 2\pi^- + \pi^0$ (MM) at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
 interactions $K^- + n \rightarrow \Lambda + \pi^- + K^+ (K_2^0) + K^- (K_1^0)$ at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
 interactions $K^- + n \rightarrow \Lambda + \pi^-$, differential cross section and polarization for backward, 26:46648
 interactions $K^- + n \rightarrow \Lambda(\Sigma^0) + \pi^+ + 2\pi^-$ at 4.5 GeV/c in deuterium bubble chamber, cross sections and resonance production in, (E), 26:46474 (COO-1428-306)
 interactions $n + n$, calculation of scattering length from p-p data accounting for charge asymmetry, 26:47055
 interactions $n + n$, scattering length in final-state, (E), 26:46861
 interactions $n + p$ at 24 MeV, differential elastic, 26:46872
 interactions $\nu + n \rightarrow \nu + n + \pi^0$ in optical spark chamber, upper limit on neutral current in, (E), 26:46453
 interactions $\nu + n \rightarrow p + \mu^-$, upper bound for tensor component of second-class current in, (T), 26:46573
 interactions $p + n$ in final state of $d + p \rightarrow 2p + n$ interactions at 52.3 MeV, angular distributions and scattering lengths for, (E), 26:46457 (KFK-1509)
 interactions with mesons at high energies, electromagnetic effects in nucleon polarization in, (T), 26:46520 (JINR-P2-6438)
 meson (K_1^0) regeneration on, Pomeranchuk theorem violation in amplitudes of, 26:46634
 motion in central electric field, (T), 26:46547
 motion through medium in magnetic field, coherent generation mechanism for induced electromagnetic waves from, (T), 26:46541
 photoproduction of, conversion of bremsstrahlung photoneutron mean energies to monochromatic photoneutron mean energies for, 26:46953
 production of slow, at CEA EL3 reactor, p. 309-11, 26:46873(R) (CEA-N-1522)
 reactions (n,f) with $A = 235$ to 241 nuclei, resonance parameters for, p. 279-86, (E), 26:46873(R) (CEA-N-1522)
 reactions (n,f) with aligned uranium-233, angular distribution of fragments from, (E), 26:47026
 reactions (n,f) with californium-252 and uranium-235, gamma and neutron emission in, p. 314-8, (E), 26:46873(R) (CEA-N-1522)
 reactions (n,f) with plutonium-239 at 10 to 310 eV, multiplicity of prompt neutrons from, p. 287-8, (E),

26:46873(R) (CEA-N-1522)
 reactions (n, γ) with iodine-127 in gaseous mixtures of butane and iodine, 26:45361
 reactions (n, γ) with iodine-127 in mixtures of iodine and butane in liquid phase, chemical state of recoil iodine-128 from, 26:45362
 reactions with americium-243 at 0.5 to 1000 eV, resonance parameters, total cross sections, and transmission for, (E), 26:47010 (ANCR-1060)
 reactions with antimony at 0.05 to 150 eV, cross sections and resonance parameters for, (E), 26:46946
 reactions with bromopentacarbonylmanganese and iodopentacarbonylmanganese, reactions of recoil bromine-82, iodine-128, and manganese-56 from, 26:45357
 reactions with lithium-6 and -7, review of ENDF/B III cross sections for, 26:46876 (ORNL-TM-3859)
 reactions with potassium chromates, comparison of thermal annealing of chromium-51 dopant-atoms and chromium-51 recoil atoms from, 26:45364
 reactions with solid acetanilide, reactions of recoil carbon-14 from, 26:45358
 scattering by solid helium-4, Fourier transform for inelastic, 26:46845 (BNL-16935)
 scattering by vortex lattice in Type-II superconductors, London model for inelastic, (T), 26:46820
 scattering in studies of condensed matter, use of inelastic, 26:47251
 shielding against $^{239}\text{PuO}_2$, (E/T), 26:46852
 single-particle energies in momentum-dependent potentials, calculation of, (T), 26:47071
 spectra from plutonium dioxide, effects of spontaneous fission and (α ,n) reactions on, 26:45704
 spectrum from $^{239}\text{PuO}_2$, (E/T), 26:46852
 thermalization of, density of neutron lethargy probability distribution for, (T), 26:46856
 thermalization of, Green function in secondary model for, 26:46846 (JEN-221)
 total cross sections for iron, from 0.8 to 11 MeV, (E/T), 26:47220 (ORNL-TM-3867)
 total cross sections for nitrogen, from 0.8 to 10 MeV, (E/T), 26:47222 (ORNL-TM-3869)
 total cross sections for oxygen, from 1.9 to 8.6 MeV, (E/T), 26:47221 (ORNL-TM-3868)
 total cross sections for sodium, from 0.8 to 11 MeV, (E/T), 26:47223 (ORNL-TM-3870)
 total cross sections for sodium, from 1.2 to 11 MeV, (E/T), 26:47224 (ORNL-TM-3871)
 transport equation for moving medium, Green's function solution for, 26:46850
 transport equation for, finite-element solution of multidimensional, 26:46851
 transport equation for, parallel shooting method for solution of spherical harmonics approximation to, 26:46859
 transport in quarter space, treatment of two-dimensional, (T), 26:46854
 transport of, solution of time-dependent equation including delayed neutrons, (T), 26:46847
 use of pulsed, in well logging, description of control and transmission system for, 26:45563(P)
 NEUTRONS, COLD
 scattering by copper below 10^{-4} eV, scattering length for coherent, (E), 26:46853
 scattering by deuterium and oxygen at -100 to $+100^\circ\text{C}$, total cross sections for elastic, (E), 26:46887
 spectra in liquid and solid methane, (E), 26:46858
 transmission through carbon, copper, and gold films below 10^{-4} eV, interference effects in, (E), 26:46853
 NEUTRONS, EPITHERMAL
 reactions (n,f) at 6 to 34 eV, probability of long-range α emission in, p. 288-9, (E), 26:46873(R) (CEA-N-1522)
 NEUTRONS, FAST
 angular distribution of cosmic-ray, above 5×10^{11} eV, (E), 26:46153
 dose distributions and spectra in tissue-equivalent phantoms, 26:45626
 dose to man following criticality accident, effects of age, diet, and sex on reliability of sulfur activation in hair and nails for determining, 26:45788
 effects on atomic displacement cascades in metals, computer simulation of, 26:46061 (CONF-710601-, pp 739-56)
 effects on blood vessels and heart of fetal rats following sublethal whole-body exposure in utero, statistical analysis of relation between age, dose, and incidence of teratogenic, 26:45867
 effects on dimensions of boron-10-enriched boron carbide-graphite control materials, 26:46077 (GULF-GA-A-12035)
 effects on elasticity and swelling of aluminum oxide and yttrium oxide insulators, 26:45702(R) (LA-4978)
 effects on melting heat rating for plutonium dioxide-uranium dioxide-base fuels, effects of porosity on, 26:45930 (HEDL-TME-72-81)
 effects on potato plants, mutagenic, 26:45770
 effects on rabbits, effects on crystalline lens of, effects of dose on RBE for 14 MeV, in comparison with x radiation, 26:45864
 effects on size of defect aggregates in aluminum and

- copper, relations with temperature, 26:46060 (CONF-710601-, pp 672-83)
- effects on swelling and creep of stainless steels, review of, 26:47303
- effects on void formation in metals, models for, 26:46063 (CONF-710601-, pp 798-824)
- effects on void formation in nickel, Nimonic PE16, and Type 316 stainless steel, 26:46052 (CONF-710601-, pp 430-48)
- effects on void formation in niobium-stabilized Type 348 stainless steel, 26:46049 (CONF-710601-, pp 174-86)
- effects on void formation in Types 304, 316, and 347 stainless steel, relations with fluence and temperature, 26:46048 (CONF-710601-, pp 156-73)
- effects on void nucleation in crystalline materials, relations with interstitials and vacancy distribution, 26:46064 (CONF-710601-, pp 825-40)
- reactions (n,2n) with beryllium-9 at 2.37 to 3.24 MeV, absolute cross sections for, (E), 26:46874 (JUL-834-RG)
- reactions (n, α) and (n,p) with nuclei, systematics of excitation functions for, 26:47066
- reactions (n, α) with vanadium-51, arsenic-75, cesium-133, and holmium-165, calculation of excitation functions for, 26:47066
- reactions (n,f) with thorium-232, uranium-233, -235, and -238 at 14 MeV, independent and cumulative yields of rare-gas isotopes from, (E), 26:47030
- reactions (n,p) with deuterium at 50 MeV, proton spectra from, (E), 26:46861
- reactions with uranium-238 at 0.8 to 30 MeV, total cross section for, (E), 26:47013 (COO-3058-23)
- scattering by A = 27 to 209 nuclei at 0.6 to 3.4 MeV, γ spectra and cross sections for inelastic, (E), 26:46909
- scattering by chromium, iron, and nickel at 1.77 to 2.76 MeV, angular distributions for elastic, (E), 26:46916 (AE-452)
- scattering by deuterium below 3.34 MeV, effective-range analysis of elastic, (T), 26:46864
- scattering by nickel-58, nickel-60, zinc-64, zinc-66, and zinc-68 up to 1.7 MeV, γ spectra and excitation functions for, (E), 26:46936
- scattering by Z = 33 to 82 nuclei at 8.05 MeV, angular distributions for elastic, (E), 26:46916 (AE-452)
- spectra of cosmic-ray, above 5×10^{11} eV, (E), 26:46153
- NEUTRONS, FISSION**
- detection of spontaneous, design and performance of well coincidence counters for, 26:45587 (BNL-50332)
- reactions (n, α), (n,p), and (n,2n) in stainless steel, cross sections for, (E), 26:46918 (JEN-230-DME/I-25)
- spectra for, effects on fast reactor physics measurements of, 26:47585 (EURFNR-1008)
- NEUTRONS, INTERMEDIATE**
- capture by titanium-48 and molybdenum-97 at 2 and 25 keV, gamma spectra from, (E), 26:46917 (T) (BNL-tr-477)
- reactions (n,f) with uranium-235, time delayed fragment spectra from, (E), 26:47018 (INDC-7-U, pp 61-4)
- reactions with uranium-235 and plutonium-239 at 8 to 60 keV, capture-to-fission ratios for, (E), 26:47024 (KFK-1563)
- transmission by iron-54 and natural iron, measurements of, (E), 26:46848
- NEUTRONS, PILE**
- effects on void formation in Nimonic PE16 alloy, 26:46047 (CONF-710601-, pp 125-41)
- effects on void formation in Types M316, EN58B, and FV548 stainless steel, 26:46047 (CONF-710601-, pp 125-41)
- NEUTRONS, PROMPT**
- multiplicity from neutron fission of plutonium-239 at 10 to 310 eV, p. 287-8, (E), 26:46873 (R) (CEA-N-1522)
- yields in fission, review on, 26:47020 (INDC-7-U, pp 113-29)
- NEUTRONS, THERMAL**
- capture by zinc-68, germanium-70, and selenium-80, γ spectra from, (E), 26:46937
- capture in boron rods, cellulose track detector measurement of spatial distribution of, 26:45615
- effects on ruby and corundum crystals, 26:47244
- reactions (n,f) with curium-245, mass yields from, (E), 26:47027
- reactions (n,f) with uranium-235, calculation of prompt γ -ray yields using renormalized Fermi gas model, (T), 26:47025
- scattering by deuterium and oxygen at -100 to +100°C, total cross sections for elastic, (E), 26:46887
- theory for pulsed, 26:46849
- NEVADA**
- ecology of Hot Creek Valley, effects of underground nuclear explosions on soil and vegetation, 26:45478 (R) (NVO-409-1)
- NEVADA TEST SITE**
- cesium content of plants at, 26:45192 (TID-25954, pp 373-5)
- cesium cycling in ecosystem of, 26:45193 (TID-25954, pp 376-8)
- ecology of Artemisia species (sagebrush) at, relation of fallout deposition to, 26:45763 (TID-25954, pp 87-101)
- ecology of Ephedra (mormon tea) at, 26:45764 (TID-25954, pp 102-9)
- ecology of *Franseria dumosa* (burr sage or burro bush) at, 26:45768 (TID-25954, pp 153-61)
- ecology of *Larrea divaricata* (creosote bush) at, relation to radioreistance, 26:45765 (TID-25954, pp 115-32)
- ecology of *Lycium* species at, relation to radiosensitivity, 26:45766 (TID-25954, pp 133-46)
- ecology of *Salsola* (Russian thistle or tumbleweed) at, 26:45767 (TID-25954, pp 147-52)
- ecology of *Yucca* species at, relation to radiosensitivity, 26:45769 (TID-25954, pp 188-98)
- fallout effects on plants at, revegetation following severe injury, 26:45762 (TID-25954, pp 75-86)
- nuclear explosions at Pahute Mesa and Yucca Flat areas of, analysis and comparison of response spectra data from, 26:45480 (NVO-1163-TM-32)
- radiation effects on desert plant survival and reproduction at, fallout beta and γ , 26:45759 (TID-25954, pp 54-7)
- radiation effects on ecology of plants at, chronic γ , 26:45757 (TID-25954, pp 12-48)
- radiation effects on growth and reproduction of desert shrubs at, chronic γ , 26:45756 (TID-25954, pp 5-11)
- radiation effects on growth and reproduction of desert shrubs at, chronic γ , 26:45757 (TID-25954, pp 12-48)
- radiation monitoring at, during July through Dec. 1971, off-site, 26:45515
- radioecology and ecophysiology of desert plants at, conference on, 26:45755 (TID-25954)
- radioisotope transport through soil-plant-small mammal ecosystems at, fallout, 26:45914 (TID-25954, pp 58-64)
- NEW MEXICO**
- leasing of USAEC-controlled uranium-bearing lands in, environmental impact report on, 26:45572 (WASH-1523)
- NEW YORK**
- strontium-90 content in diets in New York City, Jan. to March 1972, 26:45539 (HASL-257, pp II.1-6)
- NEW ZEALAND**
- radioactivity of plants and soils in uraniferous area in, 26:45529
- NEUBOLD ISLAND POWER REACTOR, UNIT 1**
- pressure vessel for, inspection during fabrication of, 26:47521 (DOCKET-50354-25)
- NICKEL**
- determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)
- determination in nickel-63 solutions, spectrographic, 26:45223 (JEN-232-DQ/I-81)
- determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
- diffusion and outgassing of ion implanted-helium in, effects of temperature on, 26:46050 (CONF-710601-, pp 230-47)
- effects on recrystallization of iron, 26:45982
- grain boundary energy of, effects of cerium, lanthanum, neodymium, and praseodymium additives on, 26:45999
- ion implantation at 1 to 10 keV in, distribution of alkali-metal, 26:47235
- meson (K-) capture by, gamma spectra from, (E), 26:46940
- neutron elastic scattering at 1.77 to 2.76 MeV, angular distributions for, (E), 26:46916 (AE-452)
- radiation effects on length of, γ , 26:46075
- radiation effects on void formation in, ion simulation of fast neutron, 26:46052 (CONF-710601-, pp 430-48)
- radioinduced defect clusters in foils of, electron microscopy studies of high-temperature high-voltage electron, 26:46057 (CONF-710601-, pp 550-62)
- radioinduced void concentrations in foils of, laser-beam, 26:46059 (CONF-710601-, pp 613-29)
- radioinduced void formation and growth in, comparison of experimental and theoretical data on neutron, 26:46062 (CONF-710601-, pp 769-97)
- separation from other metals as β -diketonates by column liquid-liquid chromatography, 26:45281
- NICKEL 201**
- oxidation by air at 1000°C, 26:45701 (R) (DP-1300)
- NICKEL ALLOYS AND SYSTEMS**
- production of wrought superalloys of, vacuum induction melting method for, 26:45981
- radioinduced void growth in, use of high-voltage electron microscope in studies of electron, 26:46056 (CONF-710601-, pp 533-49)
- Al--Cr--Ni--ThO₂, properties of dispersion-strengthened, for use in space shuttle, 26:45934 (N-72-18585)
- Ir--Ni--Ta, phase studies on, 26:46010
- Nb--Ni-base, microstructure and properties of, effects of molybdenum and tungsten additions on, 26:46035
- Nb--Ni-base, phase studies on, 26:46035
- NICKEL ISOTOPES Ni-58**
- alpha inelastic scattering at 30 MeV, DWBA analysis of differential cross sections and spectrum from, (E), 26:46927
- alpha reactions (α ,p) at 19.3 MeV, zero-range DWBA analysis of angular distributions for, (E), 26:46934
- deuteron elastic scattering at 100 MeV, effect of diffuse-edge potential and Coulomb interaction on, (T), 26:46923
- energy levels from alpha inelastic scattering, deformations and spin-parity assignments for, (E), 26:46927
- lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
- neutron inelastic scattering up to 1.7 MeV, γ spectra and excitation function for, 26:46936

photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

proton inelastic scattering at 40 MeV, DWBA and coupled channel analysis of, (E), 26:46894

NICKEL ISOTOPES Ni-60

alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932

alpha reactions (α, p) at 17 MeV, proton spectra from, (E), 26:46932

alpha reactions (α, p) at 19.3 MeV, zero-range DWBA analysis of angular distributions for, (E), 26:46934

energy levels from proton inelastic scattering and cobalt-59 ($He-3, d, \gamma$) reactions, spins and parities of, p. 41-53, (E), 26:46873(R) (CEA-N-1522)

energy-level density from alpha and proton reactions, (E), 26:46932

neutron inelastic scattering up to 1.7 MeV, γ spectra and excitation function for, 26:46936

photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

NICKEL ISOTOPES Ni-62

alpha inelastic scattering at 17 MeV, α spectra from, (E), 26:46932

alpha reactions (α, p) at 17 MeV, proton spectra from, (E), 26:46932

alpha reactions (α, p) at 19.3 MeV, zero-range DWBA analysis of angular distributions for, (E), 26:46934

energy-level density from alpha and proton reactions, (E), 26:46932

helium-3 reactions ($He-3, t$) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930

proton inelastic scattering at 13 MeV, proton spectra from, (E), 26:46932

proton reactions (p, α) at 13 MeV, α spectra from, (E), 26:46932

NICKEL ISOTOPES Ni-63

analysis of solutions of, for calcium-45 and target element, spectrographic, 26:45223 (JEN-232-DQ/I-81)

NICKEL ISOTOPES Ni-64

alpha reactions (α, p) at 19.3 MeV, zero-range DWBA analysis of angular distributions for, (E), 26:46934

helium-3 reactions ($He-3, t$) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930

NIMONIC PE 16

radiation effects on void formation in, ion simulation of fast neutron, 26:46052 (CONF-710601, pp 430-48)

radioinduced void formation in, at 300 to 650°C, pile neutron, 26:46047 (CONF-710601, pp 125-41)

NINE MILE POINT POWER REACTOR, UNIT 1

containment atmosphere for, delay in delivery of nitrogen gas for inerting of primary, 26:47350 (DOCKET-50220-101)

core spray system for, malfunction of pumps of, 26:47349 (DOCKET-50220-100)

effluents from, effects on environment of chemical, radioactive, and thermal, 26:47401 (DOCKET-50220-98)

operation of, cost-benefit analysis for, 26:47401 (DOCKET-50220-98)

operation of, proposed changes in organization for, 26:47351 (DOCKET-50220-103)

site for, ecology, geology, hydrology, and meteorology of, 26:47401 (DOCKET-50220-98)

steam lines for, leak testing of isolation valves on main, 26:47575 (DOCKET-50220-102)

NIOBIUM

creep rate during twisting at 1200 to 1850°C, effects of oxygen on, 26:46000

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251

determination by electron probe microanalysis methods, 26:45237

determination in alloys, ores, and rocks by atomic absorption spectrophotometry, 26:45234

determination of, chemical procedures for, 26:45181 (CISE-R-172)

dissolution in iron--niobium alloys, 26:46007

electron interactions with crystals of, at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550

magnetic field distributions in ellipsoids of superconducting, (T), 26:46833

magnetic field distributions in superconducting, neutron diffraction at 4.2 K of local, 26:46826

mechanical properties and structure of, review on effects of dislocations, grain orientation, heat treatments, and plastic deformation on, 26:46019

positron interactions with crystals of, at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550

radioinduced desorption of carbon monoxide from (111) face of, temperature dependence of electron, 26:47242

separation from other metals by combined ion exchange--solvent extraction chromatography, 26:45282

separation of protactinium from, in chloride and sulfate media by sorption on silica gel, 26:45390

superconducting properties of, effects of surface conditions on, (E), 26:46805 (BNL-16913)

ultrasonic attenuation in pure superconducting, (E), 26:46818

NIOBIUM ALLOYS AND SYSTEMS

Al--Fe--Nb--Si, dissolution in, 26:46007

Al--Ge--Nb, critical temperature of superconducting intermetallic Nb₃Al/sub x/Ge/sub 1-x/, variations in, 26:46021

Al--Ge--Nb, phase diagram of niobium corner of, 26:46031

Al--Ge--Nb, superconducting properties of, 26:46031

Al--Nb, critical currents in superconducting Nb₃Al, (E), 26:46829

Al--Nb, critical temperature of superconducting intermetallic Nb₃Al, variations in, 26:46021

Al--Nb, phase transformations and decomposition in superconducting, effects of heat treatments on, 26:46020

Ga--Nb, phase diagram of, 26:46018

Hf--Nb-base, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)

Mo--Nb, reactions with boron at 1100 to 1400°C, effects of composition on, 26:46002

Mo--Nb-base--W, tensile properties at 900°K, athermal, 26:45991(R) (COO-916-25)

Nb--Ni-base, microstructure and properties of, effects of molybdenum and tungsten additions on, 26:46035

Nb--Ni-base, phase studies on, 26:46035

Nb--O, metallography of, electron microscopy studies on effects of hafnium and zirconium additions on, 26:46039

Nb--Ti, phase transformation and structure of solid solutions of, mechanism of $\beta \rightarrow \omega$, 26:46036

Nb--Ti-base, phase transformations in, relation of martensite crystallography to Burgers and Bogers--Burgers, 26:46037

Nb--Zr, phase transformation and structure of solid solutions of, mechanism of $\beta \rightarrow \omega$, 26:46036

Nb-base--Ta, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)

Nb-base--Ti, fabrication of aluminum-clad wires of, method for, 26:45972(R) (COO-3311-1)

Nb-base--W, hardening of high-purity studies of substitutional solid solution, 26:45991(R) (COO-916-25)

NIOBIUM CARBIDES

electric conductivity of, effects of temperature on, 26:47250

magnetic susceptibility of cubic, at low temperatures, 26:47266

properties of resistors of, 26:47250

NIOBIUM ISOTOPES Nb-91

energy levels from zirconium-90 (d,n) reactions, spectroscopic factors for, (E), 26:46956

NIOBIUM ISOTOPES Nb-93

neutron elastic scattering at 8.05 MeV, angular distributions for, (E), 26:46916 (AE-452)

neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra and cross section for, (E), 26:46909

photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

NIOBIUM ISOTOPES Nb-95

content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)

effects on gastrointestinal tract following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

NIOBIUM ISOTOPES Nb-97

energy levels from zirconium-96 (d,n) reactions, spectroscopic factors for, (E), 26:46956

NIOBIUM OXIDES

preparation of framed films of, for electrolytic membranes and radiation windows, 26:45958(P)

K₂O--Nb₂O₅--SrO, ferroelectric properties and structure of compounds in, 26:45962

NIOBIUM SELENIDES

magnetic flux flow in superconducting NbSe₂, discontinuous, 26:46831

NITRIC ACID

corrosive effects of iron nitrate-containing, on low-carbon 18/10 stainless steel, effects of heat treatments on, 26:45970

reactions with methyl iodide, 26:45414(R) (ORNL-TM-3180)

reactions with sodium, 26:45411(R) (ORNL-TM-3072)

Hf--HNO₃, properties for etching germanium--silicon alloys, 26:45935 (SC-RR-710884)

NITROGEN

analysis for carbonyl fluoride, chlorine fluoride, perchloryl fluoride, silicon fluoride, and uranium hexafluoride by automatic absorption spectrometry, 26:45243(P)

analysis of, allowance for background in mass spectrometric isotopic, 26:45241

effects on erosion resistance of thoriated tungsten cathode in electric current, 26:45968

electron elastic scattering by molecular, crossed-beam measurements of angular distribution and total elastic-scattering and momentum-transfer cross sections for, 26:46392

electron reactions at 0 to 2000 eV with molecular, Born-approximation calculations of cross sections for excitation in, effects of choice of wave functions on,

- 26:46391
helium-metastable-atom reactions at 150 to 1600 eV with molecular, absolute cross sections for de-excitation of 2^1S and 2^3S states in, (E), 26:46394
helium-metastable-atom reactions at 50 to 600 eV with molecular, cross sections for de-excitation in, (E), 26:46394
neutron total cross sections for, from 0.8 to 10 MeV, (E/T), 26:47222 (ORNL-TM-3869)
plasma of, sampling diagnostics using ion kinetic energy distributions, 26:47128
plasma of, spectral intensity measurements from high-pressure, (E), 26:47117
proton reactions at 75 to 350 keV with molecular, absolute cross sections for formation of $3s$, $3p$, and $3d$ hydrogen atoms by charge exchange and excitation in, (E), 26:46378
quenching of $O(1D)$ by molecular, from IME lifetime measurements of $O\ I\ 6300\text{-}\text{\AA}$ emission, (E/T), 26:46278
reactions with argon, Monte Carlo trajectory calculations of three-body recombination and dissociation in, 26:46382
reactions with carbon and uranium dioxide, 26:45417(R) (ORNL-TM-3281)
reactions with carbon and uranium dioxide, 26:45408(R) (ORNL-TM-2949)
reactions with protactinium, x-ray powder diffraction studies on copounds produced by, 26:45992 (LBL-659)
reactions with vanadium at 900 to 1400°C and 0.07 to 700 torr, kinetics of, 26:45969
reactions with zirconium at 1440°C, kinetics of, 26:46030
- NITROGEN IONS**
effects on void lattice formation in molybdenum at 2 MeV, 26:46055 (CONF-710601-, pp 522-32)
- NITROGEN IONS (N_2^+)**
intensity ratio of HP/N_2^+ 4709-A and vibrational enhancement of N_2^+ first negative bands in proton-excited auroras, (E), 26:46276
- NITROGEN ISOTOPES**
determination of, allowance for background in mass spectrometric, 26:45241
- NITROGEN ISOTOPES N-14**
meson reactions (π^+ , $2p$) at 76 MeV, cross section for, 26:46870
proton inelastic scattering at 3.1 to 3.5 MeV, excitation function for, (E), 26:46903(R) (COO-1120-149)
proton reactions (p , γ) at 3.1 to 3.5 MeV, γ spectra and excitation function for, (E), 26:46903(R) (COO-1120-149)
reactions (N-14, Li-6) with carbon-12 and oxygen-16 at 45 MeV, energy spectra from, p. 22-4, (E), 26:46873(R) (CEA-N-1522)
- NITROGEN ISOTOPES N-15**
proton inelastic scattering at 24.5 MeV, DWBA analysis of differential cross sections and asymmetry from polarized, p. 123-9, (E), 26:46873(R) (CEA-N-1522)
- NITROGEN ISOTOPES N-18**
energy levels of, data compilation on, 26:46882
- NITROGEN ISOTOPES N-19**
energy levels of, data compilation on, 26:46882
- NITROGEN ISOTOPES N-20**
energy levels of, data compilation on, 26:46882
- NITROGEN OXIDES**
electron photodetachment from NO_3^- in ionospheric D region, (E/T), 26:46299
- NITROGEN OXIDES NO**
proton reactions at 75 to 350 keV with, absolute cross sections for formation of $3s$ hydrogen atoms by charge exchange and excitation in, (E), 26:46378
- NITROGEN OXIDES NO_2 OR N_2O_4**
content variations of NO_2^- with time in ionospheric D region, 26:46281
electron photodetachment from negatively ionized, in lower ionospheric D region, (E/T), 26:46299
- NITROGEN OXIDES NO_3**
content variations of NO_3^- with time in ionospheric D region, 26:46281
- NITROGEN OXIDES N_2O**
effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331
effects on yields of excited states from pulse radiolysis of deaerated naphthalene-cyclohexane solutions, 26:45330
electron reactions with, cross sections for O^- production by dissociative electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- NORTH ANNA POWER REACTOR, UNIT 1**
cooling water reservoir for, environmental effects of, 26:47511 (DOCKET-50338-48)
environmental report for, cost-benefit analysis for alternate designs, 26:47510 (DOCKET-50338-47)
transmission line for, request to resume construction of 500 kV, 26:47512 (DOCKET-50338-49)
- NORTH ANNA POWER REACTOR, UNIT 2**
cooling water reservoir for, environmental effects of, 26:47511 (DOCKET-50338-48)
environmental report for, cost-benefit analysis for alternate designs, 26:47510 (DOCKET-50338-47)
- transmission line for, request to resume construction of 500 kV, 26:47512 (DOCKET-50338-49)
NORTH ANNA POWER REACTOR, UNIT 3
environmental report for, cost-benefit analysis for alternate designs, 26:47510 (DOCKET-50338-47)
licensing of, antitrust problems in, 26:47568 (DOCKET-50404-20)
safety of, requested additional data on, 26:47567 (DOCKET-50404-19)
safety of, revised and additional data on, 26:47566 (DOCKET-50404-18)
NORTH ANNA POWER REACTOR, UNIT 4
environmental report for, cost-benefit analysis for alternate designs, 26:47510 (DOCKET-50338-47)
licensing of, antitrust problems in, 26:47568 (DOCKET-50404-20)
safety of, requested additional data on, 26:47567 (DOCKET-50404-19)
safety of, revised and additional data on, 26:47566 (DOCKET-50404-18)
- NORWAY**
radioisotope production at Institute for Atomic Energy in, 26:45709 (INIS-mf-256)
- NOVO VORONEZH POWER REACTOR, UNIT 1**
control system for, analog simulation of dynamics of, 26:47324
- NOVO VORONEZH POWER REACTOR, UNIT 2**
control system for, analog simulation of dynamics of, 26:47324
- NOVO VORONEZH POWER REACTOR, UNIT 3**
control system for, analog simulation of dynamics of, 26:47324
- NOZZLES**
joints with cylinders, method for calculation of stress distribution in, 26:45445
- NUCLEAR AUXILIARY POWER SYSTEMS**
description of 160 kW(e) zirconium hydride reactor, for Brayton turboalternator-compressor, 26:45424 (N-72-15607)
- NUCLEAR BINDING ENERGY**
saturation of, self-consistent-field calculations of, (T), 26:47039(T) (BNL-tr-488)
- NUCLEAR ELECTRIC MOMENTS**
calculation of quadrupole, single-particle configurations for Hartree-Fock, (T), 26:47057
- NUCLEAR EMULSIONS**
meson (μ) interactions at 6 and 12 GeV, cross sections for inelastic, (E), 26:46451
- NUCLEAR EXPLOSIONS**
environmental contamination from, hazards to man from, 26:45555
gamma radiation and neutron escape rates from 24-kt, HENRE program for calculation of, 26:45497
radiation dose from, use of aluminum oxide bricks as thermoluminescent dosimeter for measuring, 26:45628
- NUCLEAR EXPLOSIONS, UNDERGROUND**
data from accelerometers used in, review of problems affecting validity of, 26:45674 (UCRL-73665)
discrimination between earthquakes and, effects of tensile faults on, (T), 26:45491
effects of Cannikin, on environment and on nuclear arms race, 26:45492
effects of Cannikin, on geology and hydrology of Amchitka Island area, (E), 26:45488 (USGS-474-148)
ground motion from Cannikin Event, effects on Amchitka Island archaeological sites, (E), 26:45485 (SC-RR-720359)
ground motion from, at Hot Creek Valley, Nev., methods for evaluating effects on soils and vegetation from, 26:45478(R) (NVO-409-1)
location from seismic data measured at single station, method for improving accuracy of, 26:45495
measurement errors from misinterpretation of accelerometer results in, 26:45674 (UCRL-73665)
radioactivity distribution in and near rubble chimney from Rainier, (E), 26:45483 (NVO-1229-180)
radioisotopes in chimney rubble from Rainier, ground water transport of, (E), 26:45482 (NVO-1229-177)
seismic effects of Cannikin and Milrow, on Amchitka Island during 1969 to 1972, seismograph monitoring of, 26:45479 (NVO-746-122)
seismic identification of, at East German Moxa Station, description of simple method for, 26:45496
seismic waves from, at Pahute Mesa and Yucca Flat, Nev., analysis and comparison of response spectra data on, 26:45480 (NVO-1163-TM-32)
seismic waves generated by Nevada Test Site, and recorded at Las Vegas, characteristics of, 26:45481 (NVO-1163-TM-33)
seismic-source energies of, calculations using ground motion measurements, 26:45490
use for construction of inland waterway between Orinoco and Rio Negro Rivers in Venezuela, feasibility study of, 26:45475 (NP-19361)
uses in natural gas and hydrocarbon production stimulation and storage, evaluation of, 26:45493
- NUCLEAR EXPLOSIONS, UNDERWATER**
water waves generated by cratering, method for estimating height of, 26:45486 (UCRL-51235)
- NUCLEAR FORCES**
charge asymmetry in, effect of mixing of isovector and isoscalar mesons on, 26:47055
nucleon-nucleon, parameters for effective, (T), 26:47094

saturation of, self-consistent-field calculations of, (T), 26:47039(T) (BNL-tr-488)
single-particle spin-orbit potential, analysis of, 26:47060

NUCLEAR MAGNETIC RESONANCE

pulse Fourier transform techniques, review of, 26:47277
review of developments in high-resolution, 26:47278

NUCLEAR MATERIALS MANAGEMENT

accountability of fuel reprocessing plant input for, using weight measurements, 26:46093 (ICP-1014)
book: Atomic Safeguards: A Study in International Verification, 26:46096
book: Selected Measurement Methods for Plutonium and Uranium in the Nuclear Fuel Cycle, 26:45183 (TID-7029(2nd Ed.))
fuel assembly identification for, standard for, 26:46097
inspection and sampling method for, based on hypergeometric distribution, 26:46094 (WASH-1210)
methods at ICPP, 26:45405(R) (ICP-1006)
methods for measurements on fuel cycle materials, 26:45184 (TID-7029(2nd Ed.)), pp 3-65
plutonium measurement methods for, 26:45186 (TID-7029(2nd Ed.)), pp 253-329
sampling method for, development of continuous, 26:46092 (BNL-16930)
uranium measurement methods for, 26:45185 (TID-7029(2nd Ed.)), pp 69-252)

NUCLEAR MATTER

review of neutron, in neutron stars, 26:46466
theory of, review on, 26:47096

NUCLEAR MODELS

optical, Hartree--Fock approximation with Skyrme interaction for, (T), 26:47064
optical, isospin dependence of potential for, (T), 26:47059
shell, determination of magic numbers using statistical method of Fermi, (T), 26:47040(T) (BNL-tr-489)
shell, neutron single-particle energies in momentum-dependent potential, (T), 26:47071

NUCLEAR PHYSICS

research at Technical University of Eindhoven, 26:46146

NUCLEAR POWER

uses in production of hydrogen for transmission to consumer outlets, evaluation of, 26:45429

NUCLEAR POWER PLANTS

cost of, trends in capital, 26:47385
development in Australia, radioactive waste management aspects of, 26:46098 (AAEC/TM-628)
economics of, U.S.S.R. and Western calculational methods for, 26:47387
effects on environment, methods for evaluating, 26:45510 (AAEC/TM-627)
environmental contamination from, hazards to man from, 26:45555
environmental effects of, benefit-cost analysis of, 26:47386
heat exchanger for, using gas-cooled breeder reactor and gas turbine, parametric design study of, 26:45439 (KFK-1566)
licensing of U.S., proposed procedures for, 26:47571
licensing of, procedures in, 26:47573
location of existing and proposed US, 26:45515
radiation hazards to man in relation to operation of, criteria for establishment of maximum permissible doses on basis of, 26:45778(T) (NP-tr-1941)
site selection for, safety criteria for, 26:47574
thermal effluents from, effects on Lake Ontario aquatic ecosystems of underwater-discharged, 26:45578
turbines for 300 to 1200 MW, development of valves for, 26:45455

NUCLEAR REACTION ANALYSIS

research at Technical University of Eindhoven, 26:46146

NUCLEAR REACTIONS

DWBA for deuteron stripping, formulation using Mitra three-body model, (T), 26:47054
energy dependence of (p,n), coupled-channels calculations of, (T), 26:47073
form factors for two-nucleon transfer, in deformed nuclei, (T), 26:47061
gamma ray emission in, role of exit states of compound nucleus on, (T), 26:47082
generator-coordinate formalism for transfer in heavy-ion, development using DWBA framework, 26:47069
geometrical effects in, using polarized beams and polarized targets, 26:47041 (CEA-N-1529)
giant resonances in, phenomenological theory for, 26:47098
mechanism of heavy-ion induced, analysis of intermediate, 26:47056
Monte Carlo intranuclear cascade calculations for, using diffuse nuclear boundary, 26:47047 (JINR-P2-6503)
resonances in, spectroscopy of isobaric analog, 26:47099
S-matrix theory of, structure of compound states in, (T), 26:47068
theory for direct, at high energies, 26:47101
theory of direct, unified microscopic, (T), 26:47077
theory of multi-nucleon transfer, quasi-classical, (T), 26:47076
theory of transfer, to unbound states, diffraction, (T), 26:47075

NUCLEAR STRUCTURE

calculations for collective single-particle states of deformed odd-A nuclei, (T), 26:47081

119 SUBJECTS

NUCLEI, EVEN-EVEN

calculations for low-lying collective states of even-even nuclei, 26:47092
calculations for 0^+ vibrations in deformed nuclei, using monopole plus quadrupole pairing, 26:47042 (COO-3001-12)
calculations for sd-shell nuclei using alpha cluster model, 26:47083
calculations using hyperspherical harmonics method, 26:47090
calculations using K-harmonics method, 26:47091
description using $U_{\text{sub}} 3(A-1)/$ scheme, (T), 26:47095
regularities in, calculations using self-consistent-field method, (T), 26:47039(T) (BNL-tr-488)
research on four and five nucleon systems, review of, 26:47089
rotational-band transition probabilities, $SU(3)$ calculation of, 26:47049 (JINR-P4-6458)
symmetry for, (T), 26:46508

NUCLEAR THEORY

Bethe--Goldstone equation in, solution in shell-model basis using N--N phase shift data, 26:47074
conference on, 26:47086
few nucleon systems in, bound states of, 26:47088
few nucleon systems in, review on integral equations for scattering states of, 26:47087
microscopic approach to, development of, 26:47093
quantum theory of, review of, 26:46508
three body problem in, solution using generalized spherical harmonics, 26:47103
three-body problem in, relaxation method for solution of Faddeev equations for, 26:47065
two-particle density matrix for p-shell nuclei in $SU(3)$ configurations, 26:47067
wave functions in, expansion coefficients for (44) and (60) + (06) interactions for, 26:47084

NUCLEAR WEAPONS

existence of, environmental threat from, 26:45492

NUCLEI

electron scattering with bremsstrahlung emission, effect of nuclear multipole moments on polarized, (T), 26:47052
energy levels of, design and performance of fast-timing system for measuring lifetimes of, p. 73-6, 26:46873(R) (CEA-N-1522)
heavy-ion fusion reactions with, limits on angular momentum in, (T), 26:47072
meson (K-) capture by, K- + N scattering amplitudes for, (E/T), 26:46487
meson (ω) inelastic reactions above 10 GeV, cascade--evaporation model for shower particle cascade multiplicity and many-particle interaction effects in, (T), 26:46649 (JINR-P2-6395)
meson (ω) inelastic reactions at 5 GeV, change of mechanism for, (T), 26:47038(T) (BNL-tr-487)
neutron capture by, N-Z dependence of cross sections for, (T), 26:47085
neutron radii of, calculation using Coulomb energy differences, (T), 26:47097
neutron reactions (n,α) and (n,p) with, systematics of excitation functions for fast, 26:47066
neutron single-particle energies in, calculations using momentum-dependent potential, (T), 26:47071
nucleon inelastic reactions above 10 GeV, cascade--evaporation model for shower particle cascade multiplicity and many-particle interaction effects in, (T), 26:46649 (JINR-P2-6395)
nucleon inelastic reactions at 5 GeV, change of mechanism for, (T), 26:47038(T) (BNL-tr-487)
nucleon reactions with, numerical aspects for solving partial wave equations in momentum space for strongly peaked potentials for, (T), 26:46787
particle scattering at high energies, formalism for, 26:47100
photon reactions with, conversion of bremsstrahlung photoneutron mean energies to monochromatic photoneutron mean energies for, 26:46953
photon reactions with, isotopic structure of giant resonance in, (T), 26:47053
proton elastic scattering at 1 to 3.3 MeV by $A = 20$ to 66, fine structure of isobaric analog resonances in, (E), 26:46914
proton reactions at high energy, survey on direct, 26:46891
proton spallation reactions at 660 MeV, electromagnetic mass separator for isolation of isobars from, 26:46985
reactions of iso-, contradictions between isonuclei and many-nucleon resonance theories for energy dependence of total and differential cross sections and number of reaction occurrences as function of invariant and deficient mass of system for, (T), 26:46658
triton reactions (t,p) with, repulsive core effects in, (T), 26:47078

NUCLEI, EVEN-EVEN

collective excitations in intermediate and heavy, analysis using phenomenological models, (T), 26:47102
energy level transitions in oriented, optimization of tests of validity of time reversal in, 26:46677
energy levels of deformed, calculation using rotational model, (T), 26:47080
energy levels of deformed, $SU(3)$ multiplets for, (T),

- 26:47048 (JINR-P4-6452)
energy levels of, calculations for low-lying collective,
26:47092
energy-level transitions in deformed, SU(3)
parametrization of E2, (T), 26:46975 (JINR-P4-6462)
- NUCLEI, HEAVY
alpha decay of A = 227 to 257, relative transition
probabilities in, (T), 26:47079
collective excitations in even-even, analysis using
phenomenological models, (T), 26:47102
nucleon reactions with cosmic complex, at high energies,
pion spectra and energy transfer to π^0 for study of
collective, (E), 26:46461
proton reactions (p,n) with subsequent proton decay,
coupled-channels calculations of energy dependence of,
(T), 26:47073
- NUCLEI, INTERMEDIATE
book: Isobaric Nuclei with Mass Number A = 72 (In
Russian), 26:46939
collective excitations in even-even, analysis using
phenomenological models, (T), 26:47102
deuteron reactions (d,p) with A = 80 to 100, analysis of
threshold anomalies in, (T), 26:47058
proton reactions (p,n) with subsequent proton decay,
coupled-channels calculations of energy dependence of,
(T), 26:47073
proton reactions (p,t) with K = 0 excitation, analysis
using pairing plus quadrupole model, (T), 26:47062
- NUCLEI, LIGHT
decay of, vertex parameters and coupling constants for
virtual, (T), 26:47063
energy levels of p-shell, two-particle density matrix in
SU(3) configuration for, 26:47067
energy levels of two-baryon Δ and three-baryon,
isotriplet and isoquintet, (T), 26:46644
orbital rearrangement due to proton removal, calculations
using Brink--Boeker force and Yale potential, (T),
26:46886
particle reactions with carbon-type, at 300 to above 1000
MeV, intranuclear cascade model for density effects on
cascade particle properties and density and cluster-
structure effects on isotope yield in, (T), 26:46664
theory for Δ , containing $\Delta(1236)$, (T), 26:46644
- NUCLEI, ODD-ODD
energy levels of, shell-model analysis of multiplet, (T),
26:47051
- NUCLEIC ACID
radiation chemistry of, 26:45315(R) (COO-3286-3)
radiosensitivity of, mechanisms of, 26:45726
synthesis of, mechanisms of, 26:47612(R) (COO-3078-6)
- NUCLEIC ACID, DEOXYRIBO-
degradation of Escherichia coli, bacteriophage T4
inhibition of colicin E2-induced, 26:47639 (UR-3055-2)
phosphorus-33 incorporation into, of Drosophila, tritium
tracer study of, 26:45912(R) (ORO-3728-5)
radiation effects on distribution of nucleotide blocks
in, of rat bone marrow in comparison to leukosis
effects, x, 26:45856
radiation effects on metabolism of, in intestine of mice
in vitro and in vivo, comparison of RBE for gamma and
proton, 26:45842
radiation effects on synthesis of, in recombination-
deficient strains of Escherichia coli, γ , 26:45753
radiation effects on unscheduled synthesis of, in HeLa
cells following microbeam exposure of half-nuclei,
effects of hydroxyurea on uv, 26:45737
radiation effects on, from Bacillus subtilis, role of DNA
ligase, DNA polymerase, and endonuclease in repair of
in vitro uv, 26:45727
radiation effects on, in cultured Chinese hamster cells,
dose dependence of repair processes following x,
26:45745
radiation effects on, in cultured human Xeroderma
pigmentosum cells in comparison to effects of chemical
carcinogens and mutagens, extent of repair synthesis
following uv, 26:45742
radiation effects on, in living cells, effects of
hydroxyl radicals on, 26:45845
radioinduced damage to, role of enzymes in repair of uv,
26:45722(R) (COO-3232-1)
radioinduced single-strand breaks in transforming, of
Bacillus subtilis, effectiveness of toluene treatment
for enhancing semi-in vitro repair of γ , 26:45731
radioinduced strand breaks in, comparison of
effectiveness of iodine-125 and tritium for, 26:45894
(ORO-2690-12(Pt.B), Paper 14)
radiolysis of iodine-125-labeled, 26:45893 (ORO-2690-
12(Pt.B), Paper 13)
- NUCLEIC ACID, RIBO-
radiation effects on in vitro coding activity of
bacteriophage R17, uv, 26:45728
radiation effects on synthesis of, in spinal neurons of
mice, 26:45868
- NUCLOGENESIS
oxygen burning, hydrostatic, (T), 26:46178
- NUCLEONS
charge spectrum of solar cosmic rays from nuclear
emulsions and Explorer-35 solid-state detector, (E),
26:46330
exchange in $\pi^- + N \rightarrow \rho^0 + N$ and $\pi^+ + N \rightarrow \rho^0 + N$
interactions, interference with photon exchange in,
(T), 26:46643
form factor of, t-dependence of magnetic and electric
isovector, 26:46557
form factors for NN γ vertex, field theory on 8-
dimensional homogeneous space of Poincare group for
electric and magnetic, (T), 26:46514 (CTH-ITP-71-30)
form factors in field theory on homogeneous space of
Poincare groups, 26:46554
interactions anti K + N $\rightarrow \pi + A$ at low energies,
Veneziano duality for dips in cross sections for,
26:46635
interactions anti K + N $\rightarrow \pi + Y$ above 2.5 GeV/c, review
of backward and forward, (E), 26:46489
interactions anti $\nu + N$, parton models with integral
charge for deep-inelastic, 26:46574
interactions $\Delta + N$, models for induced nonexchange and
exchange, (T), 26:46644
interactions $\Delta + N$, properties of $\pi + N + N \rightarrow \pi + N + N$
amplitude for amplitude of, (T), 26:46644
interactions $e + N \rightarrow e + \text{hadron} + \text{anything}$ via virtual
photon + N $\rightarrow \text{hadron} + \text{anything}$ interactions at high
energies and momentum transfer, model-independent
derivation of generalized scaling laws and produced-
particle multiplicity for, (T), 26:46521 (LBL-773)
interactions $e + N \rightarrow \pi + N$ in first resonance region,
model with axial vector form factor for pion form
factor from, (T), 26:46513 (CONF-710816-17)
interactions $e + N \rightarrow \pi + N$ near threshold, current
algebra and PCAC hypothesis for amplitudes and cross
section sensitivity to nucleon axial-vector form factor
for, (T), 26:46549
interactions $e + N \rightarrow \pi^0 + N$ in first resonance region,
multipole amplitude analysis of angular distributions
and transition form factor for $\Delta(1236)$ in, (T),
26:46512 (CONF-710816-16)
interactions $e + N \rightarrow \pi^+(\pi^-) + e + \text{anything}$, PCAC
hypothesis and soft pion theorems for cross sections
for, (T), 26:46521 (LBL-773)
interactions $e + N$ at 20 GeV, evidence for partons from
inelastic, (E), 26:46466
interactions $e + N$, formulation of field theory of quarks
for structure functions of nucleon in inelastic, (T),
26:46524 (NYO-4076-18)
interactions $e + N$, predictions of nonrelativistic
harmonic-oscillator quark model for cross sections for
resonance production in, 26:46544
interactions $\gamma + N \rightarrow \text{baryon} + \text{tensor meson}$, SU(6)
symmetry for tensor meson production in, (T), 26:46684
interactions $\gamma + N \rightarrow K^+ + N$ at 5 to 15 GeV/c, Regge model
for cross sections of, (T), 26:46534
interactions $\gamma + N \rightarrow N + \text{vector mesons}$, Pomeranchuk
theorem violation for quark model for, (T), 26:46645
interactions $\gamma + N \rightarrow \pi + N$, coupling of pion-exchange
pole to photons in, (T), 26:46560
interactions $\gamma + N \rightarrow \pi(\rho^0) + N(\Delta(1236))$, method for
dispersion theory for, (T), 26:46511 (CONF-710816-9)
interactions $\gamma + N \rightarrow \pi^+ + \pi^- + \Delta$, calculation of $\rho^0 \rightarrow \pi^0$
interference in, (T), 26:46527
interactions $\gamma + N$ in broken nonlinear chiral SU(3) x
SU(3) dynamics, differential cross sections of vector
and axial-vector meson production in, (T), 26:46563
interactions $\gamma + N$ in Van Hove-type model, scaling and
Regge behavior of amplitudes in, 26:46565
interactions K + N $\rightarrow K + N$, Pomeranchuk theorem violation
for differential cross sections for, (T), 26:46645
interactions K + N, low lying singularities for energy
dependence of cross sections for angular momentum plane
structure for amplitudes of line-reversed, (E/T),
26:46629
interactions K + N, nature of absorption in ω -exchange
amplitudes for, (T), 26:46623
interactions K $^+$ + N at intermediate energies, test of
two-component theory of duality for, (T), 26:46624
interactions K $^-$ + N at intermediate energies, test of
two-component theory of duality for, 26:46624
interactions K $^-$ + N, scattering amplitude for meson (K $^-$)
capture by nuclei, (E/T), 26:46487
interactions l + N using polarized proton and deuteron
targets, review of, (E), 26:46436
interactions l + N, parton models with integral charge
for deep-inelastic, 26:46574
interactions $\Lambda + N$, two-pion exchange contribution to AN
potential from amplitudes for, 26:46652
interactions $\mu + N \rightarrow \pi + \text{anything}$ at 10.1 and 15.8 GeV,
momentum distributions of π in, (E), 26:46452
interactions N + N, nature of absorption in ω -exchange
amplitudes for, (T), 26:46623
interactions N* + N, properties of $\pi + N + N \rightarrow \pi + N + N$
amplitude for amplitudes of, (T), 26:46644
interactions $\nu + N$, $\Delta(1236)$ production in, 26:46584
interactions $\nu + N$, parton models with integral charge
for deep-inelastic, 26:46574
interactions $\nu + N$, review of dual quark-parton model for
structure functions and total cross sections for, (T),
26:46515 (DAMTP-71/24)
interactions $\pi + N \rightarrow A_1 + N$, relation of A_1 ρ coupling
constant with t-channel helicity conservation in, (T),
26:46638
interactions $\pi + N \rightarrow A_1 + N$, s-channel and t-channel
helicity conservation in, (T), 26:46646
interactions $\pi + N \rightarrow K + A$ at low energies, Veneziano
duality for dips in cross sections for, (T), 26:46635

- interactions $\pi + N \rightarrow N + \text{virtual } \gamma$, quasi-threshold behavior of amplitudes and cross sections for, (T), 26:46620 (JINR-P4-6475)
- interactions $\pi + N \rightarrow \pi + N$, duality in parton model for, (T), 26:46599
- interactions $\pi + N \rightarrow \pi + N$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- interactions $\pi + N$ at 7 to 20 GeV, amplitude analysis in Coulomb interference region for, (T), 26:46625
- interactions $\pi + N$ at high energies and small momentum transfer, review of near forward, (E/T), 26:46636
- interactions $\pi + N$ at high energies, modifications in Regge pole model for phenomenology of total cross sections in forward, (T), 26:46633
- interactions $\pi + N$, low-energy theorem from pion source dynamics for amplitude of, 26:46641
- interactions $\pi + N$, removal of ambiguities from phase-shifts for inelastic, 26:46640
- interactions $\pi^+ + N \rightarrow \rho^0 + N$, interference of photon exchange with nucleon exchange in, (T), 26:46643
- interactions $\pi^- + N \rightarrow \rho^0 + N$, interference of photon exchange with nucleon exchange in, (T), 26:46643
- mass radius of, mean square gravitational, (T), 26:46626
- polarization in meson interactions with nucleons at high energies, electromagnetic effects in, (T), 26:46520 (JINR-P2-6438)
- reactions with cosmic complex heavy nuclei at high energies, pion spectra and energy transfer to π^0 for study of collective, (E), 26:46461
- reactions with nuclei above 10 GeV, cascade-evaporation model for shower particle cascade multiplicity and many-particle interaction effects in inelastic, (T), 26:46649 (JINR-P2-6395)
- reactions with nuclei at 5 GeV, change of mechanism for inelastic, (T), 26:47038(T) (BNL-tr-487)
- reactions with nuclei, numerical aspects for solving partial wave equations in momentum space for strongly peaked potentials for, (T), 26:46787
- structure functions in inelastic $e + N$ interactions, formulation of field theory of quarks for, (T), 26:46524 (NYO-4076-18)
- structure of, evidence for parton, (E), 26:46466
- NUCLEOTIDES**
- radiation effects on distribution of, in rat bone marrow DNA, x, 26:45856
- O**
- OAK RIDGE ELECTRON LINEAR ACCELERATOR**
- experimental facility for, modification of neutron time-of-flight, 26:46144 (CONF-720607-22)
- OCONEE POWER REACTOR, UNIT 1**
- construction of, extension of permit completion date for, 26:47431 (DOCKET-50269-56)
- management surveillance of, audit task force for, 26:47433 (DOCKET-50269-61)
- safety of, revised and additional data on, 26:47432 (DOCKET-50269-60)
- OCONEE POWER REACTOR, UNIT 2**
- management surveillance of, audit task force for, 26:47433 (DOCKET-50269-61)
- operation of, notice of license issuance consideration for, 26:47434 (DOCKET-50270-49)
- safety of, revised and additional data on, 26:47432 (DOCKET-50269-60)
- OCONEE POWER REACTOR, UNIT 3**
- management surveillance of, audit task force for, 26:47433 (DOCKET-50269-61)
- operation of, notice of license issuance consideration for, 26:47434 (DOCKET-50270-49)
- safety of, revised and additional data on, 26:47432 (DOCKET-50269-60)
- ORDNANCE MATERIEL**
- inspection of welds in steel chemical munition containers, magnetic method for, 26:45470
- ORES**
- analysis for gold or silver by crucible fusion and neutron activation, 26:45200
- analysis for niobium by atomic absorption spectrophotometry, 26:45234
- analysis for trace amounts of iridium, ruthenium, and osmium by fire assay emission spectrography, 26:45229
- analysis for zirconium using picramine-epsilon, photometric, 26:45233
- atomic number of, in presence of intermediate zone, γ - γ method for determination of effective, 26:45566
- density of, in presence of intermediate zone, γ - γ method for determination of, 26:45566
- uranium-235 content in, development of semiconductor probe for determining, 26:45658(P)
- ORGANIC ACIDS**
- sulfinyl-, radioprotective effects of, on *Escherichia coli* and mice, 26:45859
- sulfonyl-, radioprotective effects of, on *Escherichia coli* and mice, 26:45859
- thio-, radioprotective effects of, on *Escherichia coli* and mice, 26:45859
- ORGANIC COMPOUNDS**
- abundance of complex, in interstellar space, (T), 26:46240
- analysis of carbon-, hydrogen-, and oxygen-containing, for ultra-micro amounts of oxygen by pyrolysis and electrolytic hygrometry, 26:45221 (EUR-4822)
- biological properties of, with C-F bonds, 26:47637 (UR-49-1508)
- oxidation in fuel reprocessing cycle, 26:45410(R) (ORNL-TM-3018)
- oxidation in fuel reprocessing off-gas, 26:45409(R) (ORNL-TM-2996)
- oxidation in fuel reprocessing solutions, 26:45420(P)
- ORNL HIGH-FLUX ISOTOPE REACTOR**
- control cylinders for, radiation effects on absorber and structural materials in, (E), 26:47587
- OSCILLATORS**
- design and performance of high-peak-power high-efficiency microcircuit avalanche diode, 26:47624 (SC-CR-723142)
- OSMIUM**
- determination of trace amounts of, in ores, concentrates, and sands by fire assay emission spectrography, 26:45229
- electric conductivity of pressed and fused, in range 100 to 1700°C, 26:47248
- thermal conductivity of pressed and fused, in range 100 to 1700°C, 26:47248
- OSMIUM ALLOYS AND SYSTEMS**
- Os-base--Th, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248
- OSMIUM ISOTOPES Os-187**
- energy levels from iridium-187 electron-capture decay, Nilsson and spin--parity assignments for, (E), 26:46984
- energy-level transitions from iridium-187 electron-capture decay, internal conversion coefficients and multipolarities for, (E), 26:46984
- OSMIUM ISOTOPES Os-189**
- energy levels at 69, 95, and 219 keV, half-lives and g-factors for, (E), 26:46981
- OSMIUM ISOTOPES Os-191**
- isomeric transition in, K-conversion coefficient for, (E), 26:47005
- OSTEICHTHYES**
- anoplopora fimbria, lead-210 and polonium-210 content in fresh- and seawater, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Clupea pallasii, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- fallout content in north Atlantic Ocean, ecological relations for cesium-137, plutonium-238, -239, and -240, and strontium-90, 26:45541 (NYO-2174-130)
- Gasterosteus aculeatus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Heteropneustes fossilis, radiation effects on androgen production and weight of seminal vesicles of, internal phosphorus-32 β , 26:45898
- Hydrolagus collieri, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Merluccius productus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Mylocheilus caurinus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Oncorhynchus nerka, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Oncorhynchus tshawytscha, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Platycephalus stellatus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Porichthys notatus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- radioisotope content of Lake Michigan, near Big Rock Point power reactor during 1970, measurements of cesium-134, cesium-137, cobalt-60, and zinc-65, 26:45536
- Spirinchus thaleichthys, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Thaleichthys pacificus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Theragra chalcogrammus, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- Trichodon trichodon, lead-210 and polonium-210 content in, in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
- OVARIES**
- radiation effects on function of, in pregnant hamsters, x, 26:45825
- OXAZOLE, 2,5-DIPHENYL-**
- radioinduced fluorescence in benzene and cyclohexane solutions, effects of concentration on, 26:45328
- radiolysis in benzene and cyclohexane solutions, effects of concentration on yield of excited triplet states from pulse, 26:45328
- OXIDES**
- radiation effects on thin layers of, sputtering, energy loss, and luminescence from 5.6-keV noble-gas-ion, 26:47238
- OXIDOREDUCTASES**
- radiation effects on activity of glutathione peroxidase in blood and liver of swine, x, 26:45733
- OXYGEN**
- abundance in solar-flare particles, (E), 26:46232
- analysis of, allowance for background in mass

- spectrometric isotopic, 26:45241
- burning in stars, nucleosynthesis during, (T), 26:46178
- cesium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366
- collisional quenching of O(1D) by molecular nitrogen from lifetime measurements of O I 6300-A emission during ionospheric modification experiments, 26:46278
- content variations with time in ionospheric D region, atomic and molecular, 26:46279
- control of atmosphere of, in industrial irradiation facility, design of equipment for, 26:45448(P)
- density distribution at 95 to 120 km in thermosphere of molecular, from solar eu occultation measurements, (E), 26:46269
- determination in boreholes, neutron activation logging method for, 26:45567
- determination in irradiated fuels and cladding materials, development of remote apparatus for, 26:45224 (LA-DC-72-524)
- determination in stainless steel and Zircaloy by deuteron and proton activation, 26:45309(R) (AE-453)
- determination of trace amounts of, in organic coolants by pyrolysis and electrolytic hygrometry, 26:45221 (EUR-4822)
- determination of ultra-micro amounts of, in organic compounds containing carbon, hydrogen, and oxygen by pyrolysis and hygrometry, 26:45221 (EUR-4822)
- determination of, choice of activation methods for, 26:45212
- effects on creep rate of niobium during twisting at 1200 to 1850°C, 26:46000
- effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331
- effects on recrystallization of iron, 26:45982
- electron reactions with molecular, cross sections for electron attachment and dissociative electron attachment in, (E), 26:46364 (ORNL-TM-3839)
- intensity of 6300-A line of atomic, visible detection of rf heating of F-region electrons from nightglow, (E), 26:46292
- neutron elastic scattering at -100 to +100°C, total cross sections for cold and thermal, (E), 26:46887
- neutron total cross sections for, from 1.9 to 8.6 MeV, (E/T), 26:47221 (ORNL-TM-3868)
- potassium-atom reactions at 50 to 350 eV with molecular, energy losses and excitation functions for dissociation, excitation, and ionization in, (E), 26:46366
- reactions with protactinium, x-ray powder diffraction studies on copounds produced by, 26:45992 (LBL-659)
- spectra of 6300-A dayglow, determination of Sept. 1969 magnetic-storm effect on F-region neutral temperature from, (E), 26:46270
- spectra of tropical nightglow in ionospheric F region, low-latitude OI 5577-A, (E), 26:46271
- OXYGEN IONS**
- content variations of O₂⁻ with time in ionospheric D region, 26:46281
- implantation in iridium at keV energies, field-ion microscopic study of damage profiles produced in, 26:47240
- reactions with cobalt, iron, tin, and zirconium at 5 MeV/amu, energy shift and relative intensity of K x rays from, (E), 26:46339
- spectral lines of, under solar-flare conditions, time-dependent ionization equilibrium, (T), 26:46226
- OXYGEN IONS (O⁻)**
- content variations with time in ionospheric D region, 26:46281
- electron photodetachment from, in lower ionospheric D region, (E/T), 26:46299
- OXYGEN IONS (O₂⁻)**
- content variations with time in ionospheric D region, 26:46281
- electron photodetachment from, in lower ionospheric D region, (E/T), 26:46299
- OXYGEN IONS (O₃⁻)**
- content variations with time in ionospheric D region, 26:46281
- electron photodetachment from, in lower ionospheric D region, (E/T), 26:46299
- OXYGEN ISOTOPES**
- determination of, allowance for background in mass spectrometric, 26:45241
- OXYGEN ISOTOPES O-13**
- beta-delayed proton decay of, analysis for mirror symmetry in, (E), 26:46879
- OXYGEN ISOTOPES O-15**
- energy levels at 10.275, 10.459, and 10.466 MeV from nitrogen-14 (p,γ) reactions, spin-parity assignments and decay properties of, (E), 26:46903(R) (COO-1120-149)
- OXYGEN ISOTOPES O-16**
- deuteron reactions (d,n) at 3 to 4 MeV, DWBA analysis of angular distributions for, (E), 26:46899
- deuteron reactions (d,p) at 12 MeV, diffraction-model analysis of differential cross sections for, (T), 26:47075
- lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
- meson reactions (π⁺,2p) at 76 MeV, cross section for, 26:46870
- muon (μ⁻) capture by, particle-hole calculation of capture rate and neutron spectra for, (T), 26:46880
- nitrogen-14 reactions (N-14, Li-6) at 45 MeV, energy spectra from, p. 22-4, (E), 26:46873(R) (CEA-N-1522)
- nuclear quadrupole moment rearrangement due to proton removal, calculation of, (T), 26:46886
- oxygen-16 reactions (O-16,α) at 62 MeV, spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- oxygen-16 reactions (O-16,C-12) at 17.5 to 30 MeV, analysis of quasimolecular resonance structure in, (E), 26:46884
- photon reactions (γ,α) at 13.1 MeV, detailed balance test of time reversal invariance in, (E), 26:46881
- photon reactions (γ,n) and (γ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
- proton separation and orbital rearrangement energies, calculations of, (T), 26:46886
- reactions (O-16, C-12) with iron-54 at 46 MeV, phase-shift analysis of differential cross sections for, p. 11-13, (E), 26:46873(R) (CEA-N-1522)
- reactions (O-16,α) with carbon-14 at 35 MeV, cross sections and spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- reactions (O-16,α) with oxygen-16 and magnesium-26 at 62 MeV, spectra from, p. 13-8, (E), 26:46873(R) (CEA-N-1522)
- reactions (O-16,C-12) with iron-54, analysis using generator-coordinate method in DWBA framework, (T), 26:47069
- reactions (O-16,C-12) with magnesium-24 and -26 and silicon-28 at 42 MeV, differential cross sections for, (E), 26:46913
- reactions (O-16,C-12) with oxygen-16 at 17.5 to 30 MeV, analysis of quasimolecular resonance structure in, (E), 26:46884
- reactions with aluminum-27 at 50 to 170 MeV, calculation of fusion excitation function using equilibrium model with fission competition, (T), 26:47072
- reactions with carbon-12 in stars, rates for, (T), 26:46179
- reactions with oxygen-16 in stars, rates for, (T), 26:46179
- scattering by tin-117 and -119 at 45 MeV, γ-ray angular distributions and spectra for inelastic, (E), 26:46961
- OXYGEN ISOTOPES O-18**
- alpha reactions (α,γ) at 4.9 to 11.5 MeV, γ-ray angular distributions and γ-γ and n-γ angular correlations for, (E), 26:46907
- carbon-12 reactions (C-12, Be-10) at 46 MeV, energy spectra from, p. 20-1, (E), 26:46873(R) (CEA-N-1522)
- energy levels of, calculation using Bethe-Goldstone equation in shell-model basis, 26:47074
- energy levels of, data compilation on, 26:46882
- proton inelastic scattering at 24.4 MeV, DWBA analysis of differential cross section and asymmetry from polarized, p. 123-9, (E), 26:46873(R) (CEA-N-1522)
- proton reactions (p,d) at 24.5 MeV, DWBA analysis of differential cross sections and asymmetries from polarized, p. 116-9, (E), 26:46873(R) (CEA-N-1522)
- proton reactions (p,n) at 3.3 to 4.0 MeV, angular distributions from, (E), 26:46929
- OXYGEN ISOTOPES O-19**
- energy levels of, data compilation on, 26:46882
- OXYGEN ISOTOPES O-20**
- energy levels of, data compilation on, 26:46882
- OXYGEN SYSTEMS**
- Nb-O, metallography of, electron microscopy studies on effects of hafnium and zirconium additions on, 26:46039
- OYSTER CREEK POWER REACTOR, UNIT 1**
- containment spray system for, failure of pump for, 26:47348 (DOCKET-50219-204)
- control rod worth minimizer for, performance of, 26:47346 (DOCKET-50219-199)
- emergency service water pump for, failure of expansion joint for, 26:47345 (DOCKET-50219-197)
- isolation condenser for, failure of steam line high flow differential pressure switch for, 26:47347 (DOCKET-50219-203)
- protection system for, failure of main steam isolation valves and recirculation loop actuation relay of, 26:47344 (DOCKET-50219-196)
- radioactive waste storage tank for, excessive radioactivity in outside, 26:47399 (DOCKET-50219-201)
- site for, maximum hurricane flood analysis for, 26:47398 (DOCKET-50219-200)
- steam isolation valve for, maintenance for main, 26:47345 (DOCKET-50219-197)
- technical specifications for, revised, 26:47400 (DOCKET-50219-202)
- OZONE**
- content variations with time in ionospheric D region, 26:46279

P**PACIFIC OCEAN**

chemical explosion detection and location in northwest,
performance of hydrophone method for, 26:45489

PACKAGING

design for radioactive materials, 26:46115(P)

PAKISTAN

water requirements for, prospects of dual purpose plants
for, 26:47579

PALISADES POWER REACTOR

effluents from, effects on environment of radioactive and
thermal, 26:47424 (DOCKET-50255-96)
radioactive waste disposal system for, codes regulating
modifications to, 26:47422 (DOCKET-50255-94)
safety injection tanks for, revised operating limits for,
26:47423 (DOCKET-50255-95)
site for, ecology of, 26:47424 (DOCKET-50255-96)

PALLADIUM

diffusion and outgassing of ion implanted-helium in,
effects of temperature on, 26:46050 (CONF-710601-, pp
230-47)
electron-phonon mass enhancement factor from specific-
heat data, 26:47262
microscopy of, imaging process in field ion, 26:46038
separation from metals by ion exchange using ion
exchangers containing isonitrosoacetamide functional
groups, 26:45283
separation from spent fuels, 26:45423(P)

PARTICLE TRACKS

counting in sheet materials using capacitive discharge
devices, 26:46083(P)
studies of fission, in lunar samples, 26:46156

PARTICLES

motion of spinning, in gravitational and electromagnetic
fields, 26:46753
motion of test, in general relativity theory, covariant
canonical equations of, (T), 26:46770

PARTONS

charge distribution in protons for scaling in deep
inelastic e + p interactions, (T), 26:46566
evidence in electron inelastic scattering by protons at
20 GeV, (E), 26:46466

PEACH BOTTOM POWER REACTOR, UNIT 2

costs for, estimated operating and permanent shutdown,
26:47375 (DOCKET-50277-51)
effluents from, effects on environment of chemical and
thermal, 26:47448 (DOCKET-50277-52)
transmission lines for, requested additional data on,
26:47448 (DOCKET-50277-52)

PEACH BOTTOM POWER REACTOR, UNIT 3

costs for, estimated operating and permanent shutdown,
26:47375 (DOCKET-50277-51)
effluents from, effects on environment of chemical and
thermal, 26:47448 (DOCKET-50277-52)
transmission lines for, requested additional data on,
26:47448 (DOCKET-50277-52)

PEOPLE'S DEMOCRATIC REPUBLIC OF GERMANY

fallout cesium-137 and strontium-90 content in cereals,
coffee, fruits, milk, and tea in Mecklenburg, 1969,
26:45546

PERCHLORYL FLUORIDES

determination in nitrogen by automatic absorption
spectrometry, 26:45243(P)

PEROXIDES

radioinduced formation of, in blood plasma in vitro,
effects of water on x, 26:45725

PERSONNEL

postgraduate training of scientific, in European
Community, 26:47616 (EUR-4832)
radiation dose to, following uptake of plutonium-239 or
plutonium-241, methods for estimation of, 26:45888
(CONF-720520-1)
radiation dose to, in Federal Republic of Germany,
legislation on maximum permissible, 26:45803
radiation protection programs for nuclear, in Federal
Republic of Germany, medical aspects of, 26:45803
radiation protection programs for, at Marcoule nuclear
center, 26:45804

PERTURBATION THEORY

Lagrangian in higher orders of, finiteness of
nonpolynomial, 26:46738
S-matrix unitarity in third order of, 26:46738
scattering amplitudes in arbitrary order of, power of
increase of, 26:46738
series convergence for arbitrary values of coupling
constant, equations for S-matrix elements of quantum
electrodynamics for, 26:46739

PETROLEUM

prospecting for, pulsed neutron well logging method for,
26:45564
storage in underground nuclear cavities, feasibility of,
26:45493

PHAGOCYTOSIS

radiation effects on, by macrophages cultured with tumor
cells, x, 26:45738

PHANTOMS

radiation dose distribution in water, for 4.15- to 6.15-
GeV bremsstrahlung, 26:45625
radiation dose distributions in tissue-equivalent, for 3-
and 15-MeV neutrons, 26:45626

radiation dose measurements in, description of two-
channel scanner for, 26:45607

radiation dosimetry in water, method using
thermoluminescent dosimeters for x, 26:45622

PHARMACEUTICALS

availability and cost of radioactive, produced by Norway
Institute for Atomic Energy, 26:45709 (INIS-mf-256)

PHASE STUDIES

book: Physicochemical Measurements in Metals Research. Pt
1, 26:46014

PHOSPHATES

determination in nuclear materials by extraction of
molybdophosphoric acid in butyl acetate in presence of
EDTA and fluoride, spectrophotometric, 26:45219 (ARH-
SA-110)
determination of, chemical procedures for, 26:45181
(CISE-R-172)

PHOSPHITES

effects of organic, on radiation resistance of plastic
insulators, 26:45672(P)

PHOSPHONIC ACID, VINYL-

methyl methacrylate--styrene--vinylphosphonic dichloride,
radioinduced polymerization of liquid and solid,
reaction mechanisms in, 26:45318

PHOSPHORIC ACID, TRIBUTYL ESTER

solvent properties for plutonium, 26:45382 (ARH-SA-130)
solvent properties for rhodium from thiocyanate
solutions, use of neutron activation analysis for
studying, 26:45194
stability in Purex Process, 26:45407(R) (ORNL-TM-2918)

PHOSPHORS

efficiency of cathodoluminescent, absolute, (E),
26:45591

PHOSPHORUS

determination as impurity in vinyl chloride by neutron
activation, 26:45204
effects on void formation in quenched stainless steel,
26:45987 (CONF-710601-, pp 605-12)

PHOSPHORUS IONS

production and acceleration of multicharged, development
of ion source for, 26:46133

PHOSPHORUS ISOTOPES P-29

energy levels from proton resonances in silicon-28, spin-
parity assignments and widths for, (E), 26:46924

PHOSPHORUS ISOTOPES P-31

meson (K-) elastic scattering by, scattering lengths for,
(E), 26:46912
meson reactions (π^- , π^-n) at 140 to 240 MeV, cross
sections for, (E), 26:46486
photon reactions (γ,n) and (γ,p) with carbon-12 up to 35
MeV, dipole resonance structure in cross sections for,
(E), 26:46890

PHOSPHORUS ISOTOPES P-32

effects on *Drosophila* following ingestion, genetic, dose
dependence of β , 26:45912(R) (ORO-3728-5)
effects on fish following internal administration,
effects on androgen production and weight of seminal
vesicles, 26:45898
effects on lungs following uptake of, review of
carcinogenic, 26:45895 (UR-3490-102)

PHOSPHORUS ISOTOPES P-33

effects on *Drosophila* following ingestion, genetic, dose
dependence of β , 26:45912(R) (ORO-3728-5)

PHOTOCHEMISTRY

research progress on, India, 1970-71, 26:45311(R) (BARC-
599)
transient species in, absorption spectroscopy of,
26:45340

PHOTOELECTRIC CELLS

signal-to-noise characteristics compared to
photomultipliers, 26:45682

PHOTOGRAPHIC FILM

image quality of radiographic, sensitometric measurements
of, 26:45468

PHOTOMULTIPLIER TUBES

performance for scintillation detectors, use of pulsed
gas laser for testing, 26:45640
signal-to-noise characteristics compared to photodiodes,
26:45682

PHOTON BEAMS

production of continuous, using internal target in
electron storage ring, 26:46145
production of quasi-monochromatic, using annihilation in
flight of monochromatic positrons, p. 265-7,
26:46873(R) (CEA-N-1522)

PHOTONS

coupling constant for γ -rho in vector dominance model,
26:46445
dosimetry of, French national standards for, 26:47216
exchange in $\pi^- + N + \rho^0 + N$ and $\pi^+ + N + \rho^0 + N$
interactions, interference with nucleon exchange in,
(T), 26:46643
exchange mechanism in hadron- and photon-induced
electromagnetic interactions at high energies,
dominance of, (T), 26:46533
form factor for $\gamma p \Delta$ vertex determination from
longitudinal and transverse cross sections for
inelastic e + p interactions in $\Delta(1236)$ region at 20 to
810, magnetic dipole transition, (E), 26:46431 (NP-
19256)

- form factors for NN γ vertex, field theory on 8-dimensional homogeneous space of Poincare group for electric and magnetic, (T), 26:46514 (CTH-ITP-71-30)
- interactions at high energies, dominance of one-photon-exchange mechanism in electromagnetic, (T), 26:46533
- interactions $\gamma + {}^3\text{He} + \text{K}^+ + \Lambda\text{-}^3\text{H}$, current algebra in soft kaon limit for differential cross sections and amplitudes of, (T), 26:46545
- interactions $\gamma + {}^4\text{He} + {}^4\text{He} + \text{w}^0$ at 2 to 5 GeV, proposed experiment for streamer chamber study of cross sections and w^0 lifetime in, (E), 26:46432(T) (SLAC-Trans-142)
- interactions $\gamma + \text{d} + \text{rho}^0 + \text{d}$ at 6, 12, and 18 GeV, forward differential cross section for, (E), 26:46445
- interactions $\gamma + \text{d} + \text{rho}^0 + \text{d}$ in Glauber theory, differential cross sections and density matrix elements for, (T), 26:46555
- interactions $\gamma + \text{e}$, quantum-mechanical calculation of, (E), 26:46532
- interactions $\gamma + \gamma + \text{w}^+ + \text{w}^-$, dispersion relations for cross section dependence on phase shift and inelasticity and pion electromagnetic mass difference from amplitude dependence on photon mass, (T), 26:46519 (JINR-P2-6333)
- interactions $\gamma + \text{l}^-$ in Weinberg model, Drell--Hearn sum rule for Compton, (T), 26:46553
- interactions $\gamma + \text{N} + \text{baryon} + \text{tensor meson}$, SU(6) symmetry for tensor meson production in, (T), 26:46684
- interactions $\gamma + \text{N} + \text{K}^+ + \text{N}$ at 5 to 15 GeV/c, Regge model for cross sections of, (T), 26:46534
- interactions $\gamma + \text{N} + \text{N} + \text{vector mesons}$, Pomanchuk theorem violation for quark model for, (T), 26:46645
- interactions $\gamma + \text{N} + \text{w} + \text{N}$, coupling of pion-exchange pole to photons in, (T), 26:46560
- interactions $\gamma + \text{N} + \text{w}(\text{rho}) + \text{N}(\Delta(1236))$, method for dispersion theory for, (T), 26:46511 (CONF-710816-9)
- interactions $\gamma + \text{N} + \text{w}^+ + \text{w}^- + \Delta$, calculation of $\text{rho}^0 \rightarrow \text{w}$ interference in, (T), 26:46527
- interactions $\gamma + \text{N}$ in broken nonlinear chiral SU(3) x SU(3) dynamics, differential cross sections of vector and axial-vector meson production in, (T), 26:46563
- interactions $\gamma + \text{N}$ in Van Hove-type model, scaling and Regge behavior of amplitudes in, 26:46565
- interactions $\gamma + \text{p} + \text{A}_2^+ + \text{n}$ at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446
- interactions $\gamma + \text{p} + \text{K}^+ + \Lambda$ at 1 to 2 and 8 to 16 GeV, finite-energy sum rules for duality relations between, (T), 26:46535
- interactions $\gamma + \text{p} + \text{n} + \text{w}^+$ at 390 to 909 MeV, asymmetry measurements for w^+ from polarized, (E), 26:46442
- interactions $\gamma + \text{p} + \text{w} + \text{p}$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446
- interactions $\gamma + \text{p} + \text{p} + (\text{rho}^0 + \text{w})$, broken-SU(3) quark model and vector dominance predictions of backward differential cross sections for, (T), 26:46529
- interactions $\gamma + \text{p} + \text{p} + \text{eta}$, factorization of Regge residues for backward differential cross sections for, (T), 26:46529
- interactions $\gamma + \text{p} + \text{p} + \text{phi}$, broken-SU(3) quark model and vector dominance predictions of backward differential cross sections for, (T), 26:46529
- interactions $\gamma + \text{p} + \text{p} + \text{w}^0$ at 426 to 918 MeV, asymmetry measurements for w^0 from polarized, (E), 26:46442
- interactions $\gamma + \text{p} + \text{p} + \text{w}^+ + \text{w}^- + \text{w}^0$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production in, (E), 26:46446
- interactions $\gamma + \text{p} + \text{p} + \text{w}^+ + \text{w}^-$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production in, (E), 26:46446
- interactions $\gamma + \text{p} + \text{p} + \text{rho}^0$, factorization of Regge residues for backward differential cross sections for, (T), 26:46529
- interactions $\gamma + \text{p} + \text{w}^- + \Delta^+$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446
- interactions $\gamma + \text{p} + \text{w}^+ + \text{n}$ at 300 to 900 MeV in polarized butanol target, asymmetry of cross section for, (E), 26:46444
- interactions $\gamma + \text{p} + \text{w}^+ + \text{n}$ at 5 and 16 GeV using polarized proton target, review of asymmetry in, (E), 26:46436
- interactions $\gamma + \text{p} + \text{w}^+ + \text{w}^+ + \text{w}^- + \text{n}$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production, (E), 26:46446
- interactions $\gamma + \text{p} + \text{rho}^0 + \text{anything}$ at 4 to 18 GeV, diffractive fragmentation of incident and target particles in, (E), 26:46437
- interactions $\gamma + \text{p} + \text{rho}^0 + \text{p}$ at 2.8 and 4.7 GeV in hydrogen bubble chamber, t dependence of mass distributions, spin dependence, and cross sections for, (T), 26:46438
- interactions $\gamma + \text{p} + \text{rho}^0 + \text{p}$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446
- interactions $\gamma + \text{p} + \text{rho}^0 + \text{p}$ in geometric diffraction picture, slope of cross sections for virtual, 26:46564
- interactions $\gamma + \text{p} + \text{rho}^- + \Delta^+$ at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446
- interactions $\gamma + \text{p} + \text{vector mesons} + \text{anything}$, quark model predictions for decay angular distributions of vector mesons from, (T), 26:46536
- interactions $\gamma + \text{p}$ in $\text{e} + \text{p}$ interactions at 5 + 200 GeV in colliding beams in ISABELLE, feasibility of experiments on particle production in, (E), 26:46426 (BNL-16968)
- interactions $\gamma + \text{p}$, fixed pole in amplitude for virtual Compton, (T), 26:46537
- interactions $\gamma + \text{w} + \gamma + \text{w}$, pomeron coupling in forward Compton, (T), 26:46560
- interactions $\gamma + \Sigma$ in broken nonlinear chiral SU(3) x SU(3) dynamics, differential cross sections for axial-vector meson production in, (T), 26:46563
- interactions with baryons, comparison of excited quark, interacting quark, and additive quark models for constraints on angular decay distributions of resonances produced in, (T), 26:46569
- interactions with bosons, regularization method in S-matrix theory in p^* momentum representation for, (T), 26:46567
- interactions with hadrons, polarization in inclusive, (T), 26:46595
- interactions with hadrons, relation between fixed poles and scale invariance in virtual Compton, 26:46559
- interactions with hadrons, scaling limit for virtual Compton, (T), 26:46530
- interactions with photons, regularization method in S-matrix theory with p^* momentum representation for, (T), 26:46567
- interactions with spin-1/2 baryons, comparison of excited quark, interacting quarks, and additive quark models for constraints on angular decay distributions of resonances produced in, (T), 26:46526 (TPJU-14/71)
- production in $\text{e} + \text{e}$ and $\text{e}^+ + \text{e}^-$ interactions, tests of Han--Nambu and Gell-Mann--Zweig triplet quark models for, (T), 26:46516 (DESY-72/21)
- production in $\text{w} + \text{N} + \text{N} + \text{virtual } \gamma$ interactions, quasi-threshold behavior of amplitudes and cross sections for, (T), 26:46620 (JINR-P4-6475)
- production of retarding, in electron and positron interactions with niobium and silicon crystals at 1 GeV, width of minimum of total yield for, (E), 26:46550
- production of soft, in $\text{w}^- + \text{d} + 2\text{n} + \gamma$ at 150 to 300 MeV, phenomenological calculation of cross sections for, (T), 26:46531
- reactions (γ, n) and (γ, p) with helium-4 up to 34 MeV, analysis of charge-dependent effects using continuum shell model and coupled channel theory, (T), 26:46866
- reactions (γ, w^0) with aluminum at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^0) with beryllium at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^0) with carbon at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^0) with copper at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^0) with lead at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^0) with silver at 3.5 to 8.3 GeV, A dependence of, (E), 26:46447
- reactions (γ, w^+) with nuclei at intermediate energies, nuclear transparency to w^+ from, (T), 26:46889
- reactions (γ, w^-) with nuclei at intermediate energies, nuclear transparency to w^- from, (T), 26:46889
- reactions with aluminum, carbon, copper, lead, and tantalum at 1.160 GeV at 150 $^\circ$, photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441
- reactions with atoms, bound states in quantum electrodynamics for, (T), 26:46721
- reactions with nuclei, conversion of bremsstrahlung photoneutron mean energies to monochromatic photoneutron mean energies for, 26:46953
- reactions with nuclei, isotopic structure of giant dipole resonance in, (T), 26:47053
- theory for elementary length, Green function in nonlocal quantum electrodynamics in, (T), 26:46698 (ITF-71-128-P)
- theory for, corpuscular character of plane wave in, (T), 26:46693
- ### PHOTOSPHERE
- faculae in, effects on apparent solar oblateness, (E/T), 26:46224
- faculae in, effects on apparent solar oblateness, (E/T), 26:46225
- ### PHTHALOCYANINE
- complexes with copper, neutron reactions with, radiation and quenching effects on chemical behavior of recoil copper-64 from, 26:45356
- ### PHYSICS
- book: Physics in the Twentieth Century, 26:46508
- statistical mechanics, functional-derivative equations for correlation functions in, 26:46728
- ### PILGRIM POWER REACTOR
- coolant cleanup system for, inoperable high flow isolation differential pressure switches of, 26:47377 (DOCKET-50293-82)
- cooling system for, unplanned actuation of isolation valves for core isolation, 26:47376 (DOCKET-50293-80)
- recirculation pumps for, unplanned restart during loss of power startup test, 26:47378 (DOCKET-50293-83)
- ### PINCH DEVICES
- discharge tube design for zeta, metal, 26:47178(P)
- plasma equilibrium in Scyllac 5-meter theta-, (E), 26:47156 (LA-DC-72-706)
- plasma instabilities in shock-heated toroidal z-, (E),

- 26:47158 (LA-DC-72-726)
 plasma properties in Scyllac 5-meter linear theta-, (E),
 26:47141 (LA-DC-72-703)
 plasma stabilization in $l = 1$ theta-, feedback, 26:47157
 (LA-DC-72-707)
 properties of superposed fields theta-, (E), 26:47171
 radiations from turbulent helium in theta-, far-infrared,
 (E), 26:47143
- 4-PIPERIDONE, 2,2,6,6-TETRAMETHYL-, N-OXIDE
 radiosensitizing effects on hypoxic mice, electron,
 26:45869
- PIPES
 radioactivity of water, fabricated in blast furnace
 containing cobalt-60 lining-wear tracers, 26:45720
- PITUITARY GLAND
 radioinduced tumors of, in mammals following uptake of
 iodine-131, review, 26:45895 (UR-3490-102)
- PLANETARY NEBULAE
 distances of, (E), 26:46171
 formation of, mechanism for, (T), 26:46176
 infrared emission at 1.65 to 3.4 microns from, (E),
 26:46175
 spectra of, neutral helium triplet in, (E), 26:46162
- PLANKTON
 radioisotope content in fresh- and seawater zoo-, in
 Puget Sound basin, 26:45544 (RLD-2225-T-14-4)
- PLANTS
 analysis for cesium by neutron activation, 26:45193
 (TID-25954, pp 376-8)
 analysis for fluorine by neutron activation using
 plutonium-beryllium source, 26:45199
 analysis for potassium by potassium-40 γ spectroscopy,
 26:45193 (TID-25954, pp 376-8)
 analysis for sulfur-35 by oxygen flask combustion and
 liquid scintillation counting, 26:45266
 analysis for trace heavy metals by neutron and photon
 activation methods, 26:45198
 cesium content at Nevada Test Site, 26:45192 (TID-25954,
 pp 373-5)
 damage to Nevada Test Site, from close-in fallout from
 Schooner Event, (E), 26:45542 (PNE-529)
 fallout effects on ecology of, at Nevada Test Site,
 revegetation following severe injury, 26:45762 (TID-
 25954, pp 75-86)
 ground motion effects on Hot Creek Valley, Nev., from
 underground nuclear explosions, methods for evaluating,
 26:45478(R) (NVO-409-1)
 radiation effects on desert shrub, at Nevada Test Site,
 chronic γ , 26:45757 (TID-25954, pp 12-48)
 radiation effects on desert shrub, at Nevada Test Site,
 effects on growth and reproduction during chronic five-
 year exposure, γ , 26:45756 (TID-25954, pp 5-11)
 radiation effects on survival and reproduction of desert,
 at Nevada Test Site, fallout beta and γ , 26:45759
 (TID-25954, pp 54-7)
 radiation effects on survival of 30 species of desert,
 dose dependence of γ , 26:45760 (TID-25954, pp 65-8)
 radioactivity monitoring of, in vicinity of Lawrence
 Livermore Laboratory, Jan. through Dec. 1971, 26:45513
 (UCRL-51242)
 radioactivity of, in uraniferous area in New Zealand,
 26:45529
 radioecology and ecophysiology of desert, at Nevada Test
 Site, conference on, 26:45755 (TID-25954)
 radioisotope transport through, at Nevada Test Site,
 fallout, 26:45914 (TID-25954, pp 58-64)
 radioisotope uptake and retention by aquatic, relation to
 transport through ecosystem, 26:45926
- PLASMA
 acceleration processes from turbulence in, (T), 26:47134
 book: Plasma: The Fourth State of Matter, 26:47104
 boundary sheath around cold electrodes in, (T), 26:47130
 conductivity of partially ionized cesium, impedance from,
 (E/T), 26:47135
 confinement by external rf fields, (E), 26:47114
 confinement of magnetic-field-free, in parallel magnetic
 multipoles, (E), 26:47111
 confinement of toroidally resistive, viscosity and
 thermal conductivity effects on, (T), 26:47105
 confinement of, nonadiabatic energy limit versus mirror
 ratio for, (E/T), 26:47106
 electric discharges through magnetic neutral point, (E),
 26:47142
 electron beam interactions with, wave excitation from
 cold, (T), 26:47151
 electron energy distribution and striations in Langmuir
 paradox region of helium discharges, (E/T), 26:47133
 equilibrium in Scyllac 5-meter theta-pinch, (E),
 26:47156 (LA-DC-72-706)
 equilibrium in Tokamak and Harmonica 2, displacement in,
 (T), 26:47115
 heating by neutral beam injection into toroidal magnetic
 traps, 26:47108
 heating by transit-time magnetic pumping and ion
 cyclotron waves, collisionless, 26:47107
 heating near ion-cyclotron frequency and harmonics,
 (E/T), 26:47109
 heating of ions in toroidal discharge, turbulent, (E),
 26:47112
 injection and collision phenomena in octupole magnetic
 fields, (E), 26:47110
 kinetic theory of two-dimensional magnetized, 26:47146
 kinetic theory of, self-consistent pair correlations and
 dielectric response in, 26:47150
 properties in Scyllac 5-meter linear theta-pinch, (E),
 26:47141 (LA-DC-72-703)
 radiations from magnetically confined hot-electron,
 synchrotron, (E), 26:47144
 radiations from Maxwellian, bremsstrahlung, (T),
 26:47152
 radiations from turbulent helium theta-pinch, enhanced
 far-infrared, (E), 26:47143
 simulation of excited monochromatic waves in one-
 dimensional Maxwellian, (T), 26:47182
 simulation of, research on, 26:47145(R) (COO-3086-15)
 spectra of argon, near-ultraviolet and visible, 26:47132
 spectra of turbulent hydrogen, validity of theory of
 dynamical linear Stark effect for interpretation of,
 26:46361
 spectral line broadening in, effects of ion motion, (T),
 26:47148
 test-particle energy losses in collisional magnetized,
 charged-, (T), 26:47149
 thermodynamics of collisionless inhomogeneous,
 statistical, (T), 26:47163
 thermodynamics of collisionless inhomogeneous, stability
 theory from statistical, 26:47164
 thermonuclear combustion waves in bounded, 26:47172
 transport of charged particles in astrophysical, (T),
 26:46294
 wave coupling in cold, electromagnetic--space charge,
 (E/T), 26:46318
- PLASMA ACCELERATORS
 jet mass, energy, and momentum from MPD, 26:47137
 properties of dense plasma focus during second
 compression in final stage, 26:47139
- PLASMA DIAGNOSTICS
 ion energy distribution from microwave discharges for
 plasma sheath kinetics, 26:47128
 ionization rate in toroidal discharges, electron
 temperature from, (E), 26:47122
 laser beam scattering on Tokamak plasma, magnetic field
 measurement from, 26:47120
 laser Thomson scattering measurements of electron density
 in hydrogen discharges, (E), 26:47125
 optical measurements for laser-produced hydrogen plasma,
 time-resolved, 26:47127
 photography of plasma focus, high-speed, 26:47121
 probe characteristics in afterglow plasmas, technique for
 recording Langmuir, 26:47119
 probe measurements in transient plasmas, optimization of
 magnetic, 26:47124
 probe measurements of supersonic cesium in magnetic de
 Laval nozzle, (E/T), 26:47123
 probe signal interpretation from laser-produced plasmas,
 cylindrical Langmuir, 26:47126
 spectra measurements of electron density and temperature
 in argon--potassium plasma, 26:47116 (IPP-IV-12)
 spectral intensity measurements from high-pressure
 nitrogen, (E), 26:47117
 spectrographic measurement of temperature of plasma
 analyzed by statistical methods, (E), 26:47129
- PLASMA INSTABILITIES
 absolute and convective instabilities in infinite plasma,
 saddle-point method calculations for, (T), 26:47161
 convective and absolute instabilities in infinite plasma,
 saddle-point method calculations for, (T), 26:47161
 drift instabilities in Q devices, effects of parallel
 wavelength on collisional, (E), 26:47168
 electromagnetic instabilities in finite-pressure, with
 anisotropic electron velocity distribution, (T),
 26:47159(T) (UCRL-Trans-10630)
 electrostatic instabilities in Tokamak devices, (T),
 26:47166
 equilibrium and stability theory of magnetically confined
 nonneutral plasmas, (T), 26:47167
 flute modes in presence of nonuniform electric fields,
 (T), 26:47162
 interchange instabilities in gravitational fields,
 stabilization by combined resistivity and viscosity,
 (T), 26:47160
 ionization instabilities in unbounded low temperature
 plasma, threshold of, (T), 26:47169
 microinstabilities in high-pressure plasma, effects of
 ion viscosity on, (T), 26:47165
 properties in shock-heated toroidal z pinches, (E),
 26:47158 (LA-DC-72-726)
 stabilization of $l = 1$ theta-pinch, feedback, 26:47157
 (LA-DC-72-707)
 theory for collisionless inhomogeneous plasma,
 statistical thermodynamics, 26:47164
- PLASMA JETS
 flow parameters of cesium, in magnetic de Laval nozzle,
 (E/T), 26:47123
 photography of plasma focus, high-speed, 26:47121
- PLASMA PRODUCTION
 hydrogen-ion injection into cusped magnetic fields,
 density measurements of, 26:47153 (NP-19364)
 ionization in electromagnetic shock tubes, precursor,
 (E), 26:47131
 laser beam irradiation of solid target for, ion energy
 spectra from, (E), 26:47154
 laser beam irradiation of solid target, momentum transfer

in, 26:47155

PLASMA WAVE PHENOMENA

Alfven wave amplification in plasma, parametric, (E/T), 26:47187

cyclotron harmonic wave propagation in inhomogeneous plasma, (T), 26:47188

cyclotron harmonic wave propagation in inhomogeneous plasma, (E), 26:47189

cyclotron harmonic waves in reflex discharges, (E/T), 26:47184

electromagnetic wave propagation in dissipative plasma, collision effects on nonlinearities in, (T), 26:47183

electromagnetic wave self-focusing in isotropic plasma, nonlinear, (T), 26:47185

electron wave echoes in plasma, structure of, (E/T), 26:47195

electrostatic electron cyclotron harmonic wave turbulence in magnetosphere, vlf nonlinear, (E), 26:46313

ion and electron wave coupling in auto-resonant plasma, parametric, (E), 26:47190

microwave reflection through laboratory plasma, boundary parameters from, (E), 26:47118

shock wave production in exploding current sheet in plasma, (E), 26:47186

shock wave propagation in expanding discharge, low-density, (E/T), 26:47194

shock wave propagation in low-beta plasma, effects of reflected ions on, 26:47199

shock wave propagation in plasma, effect of thermal conductivity on, 26:47200

shock wave structure in collisionless plasma, large-amplitude stationary hydromagnetic, (E), 26:47196

shock waves structure propagating perpendicular to initial magnetic field, collisionless, (E/T), 26:47198

soliton propagation in nonisothermal collisionless plasma, (T), 26:47202

wave coupling processes in beam-plasma system, (E), 26:47193

wave emission from plasma with finite-amplitude waves, spontaneous, 26:47180

wave evolution in one-dimensional Maxwellian plasma, simulation of excited monochromatic, (T), 26:47182

wave excitation by modulated relativistic electron beams, (E), 26:47201

wave propagation and Landau damping mechanisms in plasma, longitudinal-, (T), 26:47191

wave propagation in cylindrical plasma, nonlocal drift, (T), 26:47181

wave propagation in presence of counterstreaming beams, modified ordinary, 26:47192

PLASMAPAUSE

electromagnetic-wave coupling to space-charge waves due to beaming electrons in ambient plasma outside, (E/T), 26:46318

ion trough in, characteristics from 1969-1970 Ogo-5 data on light-, (E), 26:46301 (N-72-18362)

SAR arc formation in, theory of wave-particle interactions as mechanism for supplying thermal energy to plasmasphere electrons, (E/T), 26:46275

PLASTICS

radioinduced polymerization in wood, materials and methods for, 26:45718

use as coating for uranium solvent extraction tanks, 26:46080

x-ray transmission through, low-energy, (E), 26:47205

PLATES

response to time-dependent boundary conditions, method for calculation of, 26:45443

PLATINUM

determination of trace amounts of, in glasses by neutron activation, 26:45217

electron-phonon mass enhancement factor from specific-heat data, 26:47262

PLATINUM ALLOYS AND SYSTEMS

Co--Pt, microscopy of, imaging process in field ion, 26:46038

Hf--Pt-base--Rh--Ti--W, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)

Mo--Pt, superconducting phases in, identification of, (E), 26:46828

Pt-base--Rh, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)

Pt-base--Rh--W, properties of sheets of, for fuel capsule fabrication, 26:45703(R) (ORNL-4796)

PLATINUM ISOTOPES Pt-194

energy levels from gold-194 decay, (E), 26:47003

PLATYHELMINTHES

Leucochloridiomorpha constantiae (trematode), radiation effects on infectivity and development of metacercariae of, in chick and chorioallantois, dose dependence of, 26:45879

PLOWSHARE PROJECT

bibliography of open-file reports on Gasbuggy, Rio Blanco, Rulison, and Wagon Wheel Projects in, 26:45477 (NVO-86(Rev.1))

effects of Project Wagon Wheel in, on surrounding sandstone, prediction of explosion, 26:45494

fallout from Schooner Event in, effects on vegetation of close-in, (E), 26:45542 (PNE-529)

reservoir performance of Rio Blanco Event in, computer

calculations for predicting, 26:45476 (NVO-38-33)

site for Wagon Wheel Event in, preliminary evaluation of, 26:45571 (NVO-55)

soil porosity at site of Rio Blanco Event in, well logging data on, 26:45476 (NVO-38-33)

PLUMES

diffusion of gaseous, about isolated structures of simple geometry, effects of building orientation and shape on, 26:45514

PLUTONIUM

analysis of, review on methods for isotopic, 26:45186 (TID-7029(2nd Ed.), pp 253-329)

determination in environmental materials by extraction and liquid scintillation counting, 26:45268

determination in fuel cycle nuclear materials, sampling, standards, and bulk measurements for, 26:45184 (TID-7029(2nd Ed.), pp 3-65)

determination in fuel cycle product material, review on methods for, 26:45186 (TID-7029(2nd Ed.), pp 253-329)

determination in fuel reprocessing and radioactive waste solutions, review on methods for, 26:45190 (TID-7029(2nd Ed.), pp 331-411)

determination in fuels, book on methods for, 26:45183 (TID-7029(2nd Ed.))

determination of, equipment and reagents for, 26:45191 (TID-7029(2nd Ed.), pp 413-33)

diffusion of cobalt in delta and epsilon phases in, 26:46024

effects on spectrophotometric determination of phosphate by extracting molybdophosphoric acid in butyl acetate in presence of EDTA and fluoride, 26:45219 (ARH-SA-110)

hazards to human populations from release to environment, 26:45888 (CONF-720520-1)

production of, development of plants for, 26:45974

purification in Purex Process, 26:45410(R) (ORNL-TM-3018)

purification in Purex Process, 26:45411(R) (ORNL-TM-3072)

purification in Purex Process, 26:45407(R) (ORNL-TM-2918)

purification in Purex Process, 26:45412(R) (ORNL-TM-3095)

purification in Purex Process, 26:45416(R) (ORNL-TM-3250)

reduction of hexavalent to tetravalent, using nitric oxide gas, 26:45406(R) (ORNL-TM-2871)

separation from plutonium-uranium scrap, solvent extraction flowsheet using TBP for, 26:45382 (ARH-SA-130)

separation from spent fuels, criticality in solvent extraction system at West Valley for, 26:45401 (DOCKET-50201-128)

separation from spent oxide fuels by solvent extraction using TBP, 26:45406(R) (ORNL-TM-2871)

shipping costs for, 26:45407(R) (ORNL-TM-2918)

utilization in high temperature gas cooled reactor fuels, physics aspects of, (T), 26:47313

PLUTONIUM ALLOYS AND SYSTEMS

Pu--U, separation of plutonium from scrap, solvent extraction flowsheet using TBP for, 26:45382 (ARH-SA-130)

PLUTONIUM CARBIDES

PuC--UC, dissolution of irradiated, in nitric acid, oxidation of organic compounds formed in, 26:45420(P)

PuC--UC, preparation by sol-gel process, 26:45406(R) (ORNL-TM-2871)

PLUTONIUM CARBONITRIDES

PuCN--UCN, preparation, 26:45415(R) (ORNL-TM-3217)

PuCN--UCN, preparation of, 26:45409(R) (ORNL-TM-2996)

PuCN--UCN, preparation of, 26:45411(R) (ORNL-TM-3072)

PuCN--UCN, preparation studies on, 26:45413(R) (ORNL-TM-3127)

PLUTONIUM CHLORIDES PuCl₃

magnetic properties of, 26:47272

PLUTONIUM FLUORIDES PuF₃

magnetic properties of, 26:47272

PLUTONIUM ISOTOPES

determination in fuel cycle product material, review on methods for, 26:45186 (TID-7029(2nd Ed.), pp 253-329)

determination in fuel reprocessing and radioactive waste solutions, review on methods for, 26:45190 (TID-7029(2nd Ed.), pp 331-411)

PLUTONIUM ISOTOPES Pu-236

fission of, angular distribution of fragments from spontaneous, (E), 26:47022 (JINR-P7-6466)

PLUTONIUM ISOTOPES Pu-237

fission of, angular distribution of fragments from spontaneous, (E), 26:47022 (JINR-P7-6466)

PLUTONIUM ISOTOPES Pu-238

alpha decay of, γ spectra from, (E), 26:47032

content and distribution in open ocean silts from Atlantic Ocean and Mediterranean Sea, particle-settling model for, 26:45531 (COO-3563-3)

content in atmosphere in Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)

content in marine organisms in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)

content in soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

content in surface atmosphere at world sites, 1965 through Nov. 1971, 26:45512(R) (HASL-257(App.))

deposition at Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)

effects on lungs following uptake, review of carcinogenic, 26:45895 (UR-3490-102)

fuel capsule for thermoelectric and thermionic generator heat sources of, design of, 26:45708(P)

production and uses in medicine in France, review of, 26:45706

PLUTONIUM ISOTOPES Pu-239

content and distribution in open ocean silts from Atlantic Ocean and Mediterranean Sea, particle-settling model for, 26:45531 (COO-3563-3)

content in atmosphere in Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)

content in marine organisms in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)

content in soils in Utah, 1971, effects of nuclear weapons testing on, 26:45526 (HASL-257, pp I.95-118)

content in soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

content in surface atmosphere at world sites, 1965 through Nov. 1971, 26:45512(R) (HASL-257(App.))

content in surface atmosphere in western US, 26:45517 (HASL-257, pp I.18-32)

deposition at Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)

determination in waste drums, development of NaI(Tl) scanning system for, pp.4-6, 26:45673(R) (UCRL-50007-71-3)

distribution in tissues of mice and rabbits following intravenous injection of monomeric and polymeric forms as citrate and nitrate, 26:45917

dose to personnel following uptake of, methods for estimation of, 26:45888 (CONF-720520-1)

effects on bones following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

effects on liver following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

neutron capture-to-fission ratio at 8 to 60 keV, (E), 26:47024 (KFK-1563)

neutron fission at 10 to 310 eV, multiplicity of prompt neutrons from, p. 287-8, (E), 26:46873(R) (CEA-N-1522)

neutron strength functions at 10 to 1000 keV, S_0 and S_1 , p. 293-5, (T), 26:46873(R) (CEA-N-1522)

shipping of, design of container for, 26:46116

uptake and retention by giant brown algae, correlation between content of seawater and, 26:45924

PLUTONIUM ISOTOPES Pu-240

content in marine organisms in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)

content in soils in Utah, 1971, effects of nuclear weapons testing on, 26:45526 (HASL-257, pp I.95-118)

determination of, design and performance of neutron well coincidence counters for, 26:45587 (BNL-50332)

PLUTONIUM ISOTOPES Pu-241

dose to personnel following uptake of, methods for estimation of, 26:45888 (CONF-720520-1)

PLUTONIUM MOLYBDATES

$\text{Li}_2\text{MoO}_4\text{--Pu}(\text{MoO}_4)_2$, phase diagram of, 26:45954

$\text{Li}_2\text{MoO}_4\text{--Pu}(\text{MoO}_4)_2$, scheelite structure of phases in, 26:45954

$\text{Na}_2\text{MoO}_4\text{--Pu}(\text{MoO}_4)_2$, scheelite structure of phases in, 26:45954

$\text{NaMoO}_4\text{--Pu}(\text{MoO}_4)_2$, phase diagram of, 26:45954

PLUTONIUM NITRATES

conversion to oxide in LMFBR fuel cycle, 26:45406(R) (ORNL-TM-2871)

PLUTONIUM NITRIDES

PuN--UN , preparation of reactor fuels of, review on, 26:45960

PuN--UN , properties for fast breeder reactor fuels, review on, 26:45960

PLUTONIUM OXIDES PuO_2

neutron spectra from, effects of spontaneous fission and (α, n) reactions on, 26:45704

neutron spectrum from $^{238}\text{PuO}_2$, (E/T), 26:46852

preparation from fuel reprocessing solutions, evaluation of processes for, 26:45408(R) (ORNL-TM-2949)

preparation from fuel reprocessing solutions, evaluation of methods for, 26:45409(R) (ORNL-TM-2996)

preparation from fuel reprocessing solutions, evaluation of methods for, 26:45411(R) (ORNL-TM-3072)

preparation from solutions from fuel reprocessing, 26:45417(R) (ORNL-TM-3281)

preparation from solutions from fuel reprocessing, 26:45413(R) (ORNL-TM-3127)

preparation from solutions from fuel reprocessing, 26:45415(R) (ORNL-TM-3217)

preparation from solutions from fuel reprocessing, 26:45414(R) (ORNL-TM-3180)

preparation from solutions from fuel reprocessing cycle, 26:45412(R) (ORNL-TM-3095)

preparation of sols of, by alcohol extraction, 26:45406(R) (ORNL-TM-2871)

production of sinterable powders of, from fuel reprocessing solutions, 26:45407(R) (ORNL-TM-2918)

reactions of sols of, with formic acid, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, analysis by γ spectroscopy, 26:45409(R) (ORNL-TM-2996)

$\text{PuO}_2\text{--UO}_2$, analysis of pellets of, for moisture, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, diffusion of fission products in, 26:45417(R) (ORNL-TM-3281)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45413(R) (ORNL-TM-3127)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45415(R) (ORNL-TM-3217)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45411(R) (ORNL-TM-3072)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45409(R) (ORNL-TM-2996)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45412(R) (ORNL-TM-3095)

$\text{PuO}_2\text{--UO}_2$, dissolution in nitric acid, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, dissolution of irradiated, in nitric acid, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, fabrication of fuel rods of, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, fabrication of pellets from sol-gel-derived, 26:45406(R) (ORNL-TM-2871)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45410(R) (ORNL-TM-3018)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45412(R) (ORNL-TM-3095)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45413(R) (ORNL-TM-3127)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45415(R) (ORNL-TM-3217)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45417(R) (ORNL-TM-3281)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45407(R) (ORNL-TM-2918)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45409(R) (ORNL-TM-2996)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45414(R) (ORNL-TM-3180)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, fabrication of sol-gel-derived, Sphere-Pac process for, 26:45411(R) (ORNL-TM-3072)

$\text{PuO}_2\text{--UO}_2$, oxidation of unirradiated pellets of, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, performance of fuels of, 26:45406(R) (ORNL-TM-2871)

$\text{PuO}_2\text{--UO}_2$, preparation and properties of microspheres of, 26:45411(R) (ORNL-TM-3072)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45410(R) (ORNL-TM-3018)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45413(R) (ORNL-TM-3127)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45417(R) (ORNL-TM-3281)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45407(R) (ORNL-TM-2918)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45414(R) (ORNL-TM-3180)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45411(R) (ORNL-TM-3072)

$\text{PuO}_2\text{--UO}_2$, preparation by sol-gel process, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres from sols of, 26:45409(R) (ORNL-TM-2996)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45410(R) (ORNL-TM-3018)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45412(R) (ORNL-TM-3095)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45415(R) (ORNL-TM-3217)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45417(R) (ORNL-TM-3281)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45414(R) (ORNL-TM-3180)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, 26:45408(R) (ORNL-TM-2949)

$\text{PuO}_2\text{--UO}_2$, preparation of microspheres of, by CUSP-APEX mixed sol process, 26:45406(R) (ORNL-TM-2871)

$\text{PuO}_2\text{--UO}_2$, preparation of sols of, 26:45407(R) (ORNL-TM-2918)

$\text{PuO}_2\text{--UO}_2$, preparation of sols of, 26:45409(R) (ORNL-TM-2996)

$\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, 26:45415(R) (ORNL-TM-3217)

$\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, 26:45417(R) (ORNL-TM-3281)

$\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, 26:45414(R) (ORNL-TM-3180)

$\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, 26:45416(R) (ORNL-TM-3250)

$\text{PuO}_2\text{--UO}_2$, radiation effects on sol-gel-derived, neutron, 26:45413(R) (ORNL-TM-3127)

- PuO₂--UO₂, radiation effects on, neutron, 26:45410(R)
(ORNL-TM-3018)
- PuO₂--UO₂, radiation effects on, neutron, 26:45412(R)
(ORNL-TM-3095)
- PuO₂--UO₂, radiation effects on, neutron, 26:45411(R)
(ORNL-TM-3072)
- PuO₂--UO₂, radiation effects on, neutron, 26:45409(R)
(ORNL-TM-2996)
- PuO₂--UO₂, radiation effects on, neutron, 26:45408(R)
(ORNL-TM-2949)
- PuO₂--UO₂, reprocessing of metallic-clad fuel elements
of, electrochemical method of cutting and leaching in,
26:45422(P)
- PuO₂--UO₂, solubility in nitric acid, 26:45417(R) (ORNL-
TM-3281)
- PuO₂--UO₂-base, radiation effects on melting heat rating
for, effects of porosity on fast neutron, 26:45930
(HEDL-TME-72-81)
- PLUTONIUM COMPOUNDS
- hydrolysis of, partition chromatography for studies on,
26:45366 (JINR-P6-6402)
- PLYWOOD
- x-ray transmission through, low-energy, (E), 26:47205
- PNEUMATIC SYSTEMS
- design for sealing flasks in hot cells, 26:46112(P)
- POINT BEACH POWER REACTOR, UNIT 1
- boric acid storage tank for, incorrect dilution in,
26:47372 (DOCKET-50266-73)
- operations of, additional data on Off-Site Review
Committee activities for, 26:47430 (DOCKET-50266-74)
- radiation monitoring requirements for, additional data
on, 26:47430 (DOCKET-50266-74)
- steam generator for, repair of tube sheet cladding for,
26:47373 (DOCKET-50266-75)
- POINT BEACH POWER REACTOR, UNIT 2
- boric acid storage tank for, incorrect dilution in,
26:47372 (DOCKET-50266-73)
- operation of, license for 300 MW(t) maximum, 26:47469
(DOCKET-50301-64)
- operations of, additional data on Off-Site Review
Committee activities for, 26:47430 (DOCKET-50266-74)
- radiation monitoring requirements for, additional data
on, 26:47430 (DOCKET-50266-74)
- POLAND
- cesium-137 content in muscles of broilers, hens, and
partridges in Warsaw area, 1968 to 1969, 26:45548
- radioactivity of atmospheric aerosols in, relation with
aerosol dust content, (E), 26:45521
- POLARIMETERS
- development of two-section Ge(Li) compton, for γ rays,
26:46893
- POLONIUM ISOTOPES Po-210
- content in organisms, sediment, and water in freshwater
lake and salt water sound in Puget Sound basin,
26:45544 (RLO-2225-T-14-4)
- effects on kidneys following uptake of, review of
carcinogenic, 26:45895 (UR-3490-102)
- effects on lungs following uptake of, review of
carcinogenic, 26:45895 (UR-3490-102)
- encapsulation of heat sources of, method for,
26:45707(P)
- energy levels from astatine-210 decay, lifetimes and
spin--parity assignments for, (E), 26:47006
- energy-level transitions from astatine-210 decay,
conversion coefficients and multipolarities for, (E),
26:47006
- uptake and retention by giant brown algae, correlation
between content of seawater and, 26:45924
- POLONIUM ISOTOPES Po-212
- alpha decay of, charge states of lead-208 from, 26:45359
- POLYMERS
- radiation effects on cords of Kapron, x-ray scattering
studies on electron or γ , 26:45338
- radiation resistance of insulators of, effects of organic
phosphite additives on, 26:45672(P)
- radioinduced graft polymerization of hydroxyalkyl
methacrylates on, hydrogels for biological uses from γ ,
26:45312(R) (COO-2147-2)
- concretes--polymers, properties of radiation-polymerized,
for use in low-porosity pipes, 26:45718
- POPULATIONS
- hazards to human, from release of plutonium to
environment, 26:45888 (CONF-720520-1)
- POROSIMETERS
- calibration of two-detector gamma--gamma logging,
26:45653(T) (CSIRO-Trans-10705)
- POSITRONIUM
- decay in time-dependent external field, perturbation
frequency dependence of decay constants for parametric
resonance in, (T), 26:46571 (KFKI-72-15)
- fine structure of, 26:46710
- POSITRONS
- annihilation in liquid crystals, positronium-phase and
temperature dependence of lifetimes in, (E), 26:46363
- effects on secondary emission and properties of solids,
Russian book on, 26:47245
- interactions $e^+ + e^- + 2e^+ + 2e^-$ at 990 MeV in colliding
beams, upper limit for cross sections for, (E),
26:46429 (CONF-710816-15)
- interactions $e^+ + e^- + e^+ + e^-$, acoplanarity
distributions and radiative corrections in Bhabha, (T),
26:46543
- interactions $e^+ + e^- + \gamma +$ hadrons, canonical anomaly for
high-energy behavior of, (T), 26:46556
- interactions $e^+ + e^- +$ hadron pairs at 1400 to 2400 MeV,
energy dependence of, (E/T), 26:46552
- interactions $e^+ + e^- +$ hadrons + anything, review of dual
quark-parton model for semi-inclusive, (T), 26:46515
(DAMTP-71/24)
- interactions $e^+ + e^- + \mu^+ + \mu^- +$ anything, review of dual
quark-parton model for semi-inclusive, (T), 26:46515
(DAMTP-71/24)
- interactions $e^+ + e^- + \phi + K_1^0 + K_2^0, K_1^0 + \pi^+ + \pi^-$,
mass of ϕ from, (E), 26:46443
- interactions $e^+ + e^- + \pi +$ hadrons, behavior of structure
functions and cross sections for, (T), 26:46540
- interactions $e^+ + e^- + \pi^+ + \pi^- + \pi^0$ ($2\pi^0$) at 990 MeV in
colliding beams, total cross sections for, (E),
26:46429 (CONF-710816-15)
- interactions $e^+ + e^- + \pi^+ + \pi^- + \pi^0$ in ω region in
storage ring, cross sections for, (E), 26:46427 (CONF-
710816-13)
- interactions $e^+ + e^- + \pi^+ + \pi^-$ in colliding beams, $\pi^+\pi^-$
excitation curve in rho region for amplitude and phase
of ω --rho interference and rho production cross
sections in, (E), 26:46428 (CONF-710816-14)
- interactions $e^+ + e^-$, tests of Han--Nambu and Gell-Mann--
Zweig triplet quark models for 2γ production in, (T),
26:46516 (DESY-72/21)
- interactions of, test of quantum electrodynamics from,
26:46710
- interactions with niobium and silicon crystals at 1 GeV,
width of minimum of total yield of retarding photons
from, (E), 26:46550
- lifetime in light and heavy ice and water and frozen
aqueous solutions, effects of temperature and
positronium trapping on, (E), 26:46362 (RISO-254)
- reactions with hydrogen atoms, polarization effects in
close-coupling calculations of elastic phase shifts and
cross sections for inelastic scattering and positronium
formation in, 26:46372
- scattering by helium and hydrogen, calculation using
method of effective potentials and variational
formulations, 26:46400
- spectra from electron reactions with tantalum at 12 to 30
MeV, effects of target thickness on, 26:46149
- POTASSIUM
- content in human diets in Chicago, Jan. to June 1972,
26:45540 (HASL-257, pp III.45-52)
- determination in animal, plant, and soil samples by
potassium-40 γ spectroscopy, 26:45193 (IID-25954, pp
376-8)
- determination in hair by neutron activation and γ
spectroscopy, 26:45214
- determination in minerals and rocks by neutron
activation, 26:45197
- determination in soils, 26:45263 (RT/PROT-72-11)
- electron--phonon unklapp scattering at low temperatures
in, (E), 26:47264
- mass flow choking limit for heat pipe working fluid of,
(E), 26:45460
- reactions with bromine, chlorine, and oxygen molecules at
50 to 350 eV, energy losses and excitation functions
for dissociation, excitation, and ionization in, (E),
26:46366
- solvation numbers from compressibilities and ionic
vibration potentials measurements, 26:45369
- POTASSIUM ALLOYS AND SYSTEMS
- Ar--K, spectra measurements of electron density and
temperature in plasma of, 26:47116 (IPP-IV-12)
- POTASSIUM CHLOROPALLADATES
- radiolysis of single crystals of, ESR studies on products
from electron or γ , 26:45341
- POTASSIUM CHLOROPLATINATES
- radiolysis of single crystals of, ESR studies on products
from electron or γ , 26:45341
- POTASSIUM CHROMATES
- neutron reactions with, comparison of thermal annealing
of chromium-51 dopant-atoms and chromium-51 recoil
atoms from, 26:45364
- POTASSIUM CYANOIRIDATES
- radiation effects on, in potassium chloride lattice, ESR
studies on, 26:45317
- POTASSIUM FLUORIDES
- KF--UF₆, spectra of complex compounds in, ir, 26:45230
- POTASSIUM IONS
- implantation in nickel at 1 to 10 keV, penetration and
spatial distribution from, 26:47235
- POTASSIUM ISOTOPES K-36
- energy levels from argon-36 (He-3,t) reactions, (E),
26:46904
- POTASSIUM ISOTOPES K-39
- energy levels from proton inelastic scattering, low-lying
doublet, (E), 26:46928
- POTASSIUM ISOTOPES K-40
- content in fallout deposited on Dallas, Texas soil during
Jan. to July 1964, (E), 26:45527 (IID-26103)
- content in milk, soils, and vegetation in vicinity of
Lawrence Livermore Laboratory, Jan. through Dec. 1971,
26:45513 (UCRL-51242)
- POTASSIUM ISOTOPES K-41
- proton reactions (p,n) at 3.7 and 3.9 MeV, angular
distributions from, (E), 26:46929

POTASSIUM ISOTOPES K-42
energy levels of, spin determinations from γ - γ angular correlations, 26:46901

POTASSIUM NITRATES
decomposition in fluidized-bed processing of radioactive wastes, 26:45383 (ICP-1008)

POTASSIUM OXIDES
 K_2O - Nb_2O_5 - SrO , ferroelectric properties and structure of compounds in, 26:45962
 K_2O - SiO_2 - SrO , diffusion coefficients for, at 630 to 810°C, measurement of inter-, 26:45946

POTASSIUM URANIUM CHLORIDES
 K_2UCl_6 - Rb_2UCl_6 , spectra of, effects of chemical state of uranium on, 26:45377

POWDERS
compaction of, electrodynamic method for, 26:45951
electron nuclear double resonance spectra of, 26:47279

POWER CYCLES
performance of nuclear Brayton, for compressor and turboalternator using helium-xenon working fluid, 26:45424 (N-72-15607)

POWER PLANTS
effects on environment of fossil-fueled, methods for evaluating, 26:45510 (AAEC/TM-627)
thermal effluent discharge from, into large lakes, effects on atmosphere, 26:45584
thermal effluents from, economic evaluation of, 26:45573 (AE-448)
thermal effluents from, effects on aquatic environment, 26:45583
thermal effluents from, effects on temperature distribution in southwest Lake Michigan waters from underwater-discharged, (E), 26:45580
thermal effluents from, method for acquiring and analyzing data on dispersion of underwater-discharged, 26:45579
turbines for 1000 MW steam, development of 50-Hz 44-in. blades for, 26:45454
waste heat from US, use for heating small livestock shelters, 26:45576

POWER SUPPLIES
current leads for superconducting magnets, design of removable, 26:45679

PRAIRIE ISLAND POWER REACTOR, UNIT 1
construction of, application for extension of completion date for, 26:47450 (DOCKET-50282-59)
refueling water storage tanks for, corrosion of stainless steel, 26:47452 (DOCKET-50282-68)
safety of, revised and additional data on, 26:47451 (DOCKET-50282-67)
safety of, revised and requested data on, 26:47453 (DOCKET-50282-69)

PRAIRIE ISLAND POWER REACTOR, UNIT 2
refueling water storage tanks for, corrosion of stainless steel, 26:47452 (DOCKET-50282-68)
safety of, revised and additional data on, 26:47451 (DOCKET-50282-67)
safety of, revised and requested data on, 26:47453 (DOCKET-50282-69)

PRASEODYMIUM
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
effects on grain boundary energy of nickel, 26:45999
separation from europium, gadolinium, neodymium, and samarium by anion exchange using acetic acid-methanol-nitric acid mixtures, 26:45280
solubility in oxygen-containing chromium at 1000 to 1800°C, microhardness studies on, 26:46034
solvent properties of oxygen-containing, for cerium, neodymium, and praseodymium at 1000 to 1800°C, microhardness studies on, 26:46034
uptake of radioactive, by larval and hatchling amphibians and reptiles, 26:45815(R) (COO-3486-1)

PRASEODYMIUM ISOTOPES Pr-143
beta decay of, γ spectra from hindered, (E), 26:46955

PRASEODYMIUM ISOTOPES Pr-144
accumulation by organisms and plants in littoral freshwater microecosystem, 26:45553

PREAMPLIFIERS
circuits for charge, p. 77-8, 26:46873(R) (CEA-N-1522)
circuits for Cherenkov spectrometers, 26:45608

PRESSURE VESSELS
development of reactor prestressed concrete, review of, 26:47311
fracture of heavy-section steel, mechanics of, 26:45996 (WCAP-7841)
fracture toughness of steel, use of J integral for analysis of, 26:45990 (CONF-720806-2)

PRINCETON SYNCHROTRON
operation for accelerating argon ions to 11.7 GeV, 26:46124

PROMETHIUM ISOTOPES Pm-141
beta and electron-capture decay of, γ singles and coincidence spectra from, (E), 26:46972
energy-level scheme from samarium-141 decay, (E), 26:46972

PROMETHIUM ISOTOPES Pm-144
content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (IID-26103)

PROMETHIUM ISOTOPES Pm-147
beta decay of, energy-dependent K-shell internal ionization probability in, (T), 26:46962
distribution and retention in tissues of rats following

intravenous injection as chloride solution, effects of DTPA treatment on removal, 26:45922

PROMETHIUM ISOTOPES Pm-148
distribution and retention in tissues of rats following intravenous injection as chloride solution, effects of DTPA treatment on removal, 26:45922

PROPANE
proton reactions at 75 to 350 keV with, absolute cross sections for formation of 3s hydrogen atoms by charge exchange and excitation in, (E), 26:46378

1-PROPANOL
effects on sorption of protactinium on silica gel from aqueous solutions of hydrochloric and nitric acids, 26:45389

2-PROPANOL
effects on sorption of protactinium on silica gel from aqueous solutions of hydrochloric and nitric acids, 26:45389

2-PROPANONE, 1,3-DIHYDROXY-
reactions with hydrated electrons from pulse radiolysis of aqueous solutions, 26:45354

2-PROPANONE, PHENYL-
reactions with hydrated electrons from pulse radiolysis of aqueous solutions, 26:45354

PROPENE
oxidation to acrolein, structure and activity relation of antimony uranium oxide catalyst used in, 26:45374
reactions with deuterium over zinc oxide, mechanisms of, 26:45303
 NH_2 -propene, radiolysis of mixtures of, yields of hydrogen from γ , 26:45332

PROPENE POLYMERS
radioinduced graft polymerization of unsaturated monomer containing hydrophilic groups in, for inducing dyeability, γ , 26:45336(P)

PROTACTINIUM
crystal structure of, 26:45992 (LBL-659)
properties of, magnetic and thermal, 26:45992 (LBL-659)
reactions with gases, crystal structure of compounds produced by, 26:45992 (LBL-659)
separation from hafnium, niobium, and zirconium by sorption on silica gel from chloride and sulfate media, 26:45390
separation from thorium salts and uranium ore by sorption on silica gel, 26:45390
separation of tetravalent and pentavalent, in sulfuric acid and hydrochloric acid by sorption on silica gel and elution with chromium(II), 26:45391
sorption from aqueous hydrochloric and nitric acid solutions, effects of alcohols, acetone, and dioxane on, 26:45389

PROTACTINIUM CHLORIDES
magnetic properties of $PaCl_3$, 26:47272

PROTACTINIUM FLUORIDES
magnetic susceptibility of, 26:45992 (LBL-659)

PROTEINS
radiation effects on synthesis of, in mouse spleen cell nuclei following whole-body exposure, comparison of effects of Friend virus infection with x, 26:45849
synthesis during cycle of L5178Y cells, 26:47640 (UR-3490-18)
synthesis during cycle of L5178Y cells, 26:47651 (UR-3490-154)
synthesis of G_1 , in L5178Y-cell, effects of dactinomycin on, 26:47650 (UR-3490-153)
synthesis of, mechanisms of, 26:47612(R) (COO-3078-6)
synthesis of, regulation of early steps in, 26:47611(R) (COO-3078-5)

PROTON SOURCES
design and performance of, 26:45432

PROTON SPECTROMETERS
description of electrostatic, with channel electron multipliers for protons at 0.03 to 10 keV, 26:45642

PROTONS
channeling in paraelectric and ferroelectric barium titanate crystals at 3.8 MeV, 26:47234
diffusion in troposphere and stratosphere, relation with atmospheric circulation, (E/T), 26:46297
dosimetry of, design of tissue-equivalent liquid ionization chamber for, 26:45621
effects of 22-MeV, on silicon single-junction and silicon-controlled rectifiers, 26:45661 (N-72-17158)
effects on auroral excitation, HP/N_2^+ 4709-A intensity ratio and vibrational enhancement of N_2^+ first negative bands in, (E), 26:46276
effects on body weight of monkeys, effects of age, dose, energy, and sex on, 26:45841
effects on extracellular bacteriophages, lethal and mutagenic, 26:45746
effects on mechanical properties of copper and tungsten at 10 MeV and low temperatures, 26:46066(R) (COO-3431-1)
effects on mice, effects on DNA metabolism in intestine of, in vitro and in vivo, RBE in comparison to gamma, 26:45842
effects on n-type silicon at keV energies, electrical properties of shallow donors produced in, 26:47237
effects on sodium chloride crystals, intensity dependence of F-center accumulation in, 26:47246
effects on swelling of Type 316 stainless steel at 0.75

to 1.0 MeV and 500 to 600°C, 26:46054 (CONF-710601-, pp 511-21)

effects on void formation in Type 316 stainless steel at 400 to 600°C and 1 MeV, 26:46044 (AI-AEC-13033)

flux in solar wind, dependence of helium abundance on, (E/T), 26:46328

fluxes in nighttime aurorae, Fermi acceleration mechanism for keV, (E/T), 26:46267

form factor for $\gamma p \Delta$ vertex determination from longitudinal and transverse cross sections for inelastic $e + p$ interactions in $\Delta(1236)$ region at 20 to 810, magnetic dipole transition, (E), 26:46431 (NP-19256)

form factor from elastic $p + p$ interactions at high energies, strong-interaction, 26:46654

form factors from $e + p$ interactions, validity of scale relation for electromagnetic, (E), 26:46433

implantation in gallium arsenide at 300 keV, for optical waveguide production, 26:47232

interactions anti $p + p + \text{anti } n + n$ at 1.00 to 3.17 GeV/c, forward differential cross section in, (E), 26:46470

interactions anti $p + p + \text{anti } p + p$ at 1.6 and 2.5 GeV/c, asymmetries in polarized, (E), 26:46468

interactions anti $p + p + \text{anti } p + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions anti $p + p + K^+ + K^-$ at 5 GeV/c, crossing relations from differential cross sections for backward $K^+ + p + K^+ + p$ and $K^- + p + K^- + p$ interactions and, (E), 26:46463

interactions anti $p + p + \pi^+ + \pi^-$ pions, limiting behavior of partially inclusive, (T), 26:46613

interactions anti $p + p + \pi^+ + \pi^-$ at 1.6 GeV/c, asymmetries in polarized, (E), 26:46468

interactions anti $p + p + \pi^+ + \pi^-$ at 5 GeV/c, crossing relations from differential cross sections for backward $\pi^+ + p + \pi^+ + p$ and $\pi^- + p + \pi^- + p$ interactions and, (E), 26:46463

interactions anti $p + p$ at 1.51 to 2.90 GeV/c in bubble chamber, cross sections for, (E), 26:46469

interactions anti $p + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462

interactions anti $p + p$ at rest, resonance production in, (E), 26:46465

interactions anti $p + p$, low lying singularities for energy dependence of cross sections for angular momentum plane structure for amplitudes of line-reversed, (E/T), 26:46629

interactions at super-high energies, bounds on $\mu^+ - \mu^-$ pair and $W^+(W^-)$ production in, 26:46598

interactions $d + p + 2p + n$ at 52.3 MeV, angular distributions and scattering lengths for $p + p$ and $p + n$ final-state interactions and model for $p + p$ final-state interactions in, (E/T), 26:46457 (KFK-1509)

interactions $d + p + 2p + n$ at 52.3 MeV, differential cross sections and scattering length for $p + p$ final-state interactions for amplitudes of, (T), 26:46650 (KFK-1547)

interactions $e + p + e + p + \pi^+ + \pi^-$ in rho resonance region, extension of diffraction--dissociation model for polarization, momentum transfer, and virtual photon and dipion mass dependence of cross sections for, (T), 26:46517 (DESY-72/23)

interactions $e + p$ at 5 + 200 GeV in colliding beams in ISABELLE, feasibility of photoproduction experiments from total cross sections for, (E), 26:46426 (BNL-16968)

interactions $e + p$ at large angles, parity violation in elastic, (E/T), 26:46562

interactions $e + p$ in $\Delta(1236)$ region at 20 to 810, magnetic dipole transition form factor of $\gamma p \Delta$ vertex determination from longitudinal and transverse cross sections for inelastic, (E), 26:46431 (NP-19256)

interactions $e + p$ in parton model, scaling behavior of structure functions for deep inelastic, (T), 26:46566

interactions $e + p$, invariant method for differential cross sections, angular distributions, and two-photon mechanism for meson production in, (T), 26:46518 (ITEF-848)

interactions $e + p$, polarization contribution to ground-state hyperfine splitting of hydrogen from structure functions from resonance model of inelastic, (T), 26:46523 (NYO-4076-17)

interactions $e^- + p + e^- + \mu^+ + \mu^- + p$, scaling limit for virtual Compton scattering off forward direction from, (T), 26:46530

interactions $\gamma + p + A_2^+ + n$ at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446

interactions $\gamma + p + K^+ + A$ at 1 to 2 and 8 to 16 GeV, finite-energy sum rules for duality relations between, (T), 26:46535

interactions $\gamma + p + n + \pi^+$ at 390 to 909 MeV, asymmetry measurements for π^+ from polarized, (E), 26:46442

interactions $\gamma + p + \omega + p$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446

interactions $\gamma + p + p + (\rho^0 + \omega)$, broken-SU(3) quark model and vector dominance predictions of backward

differential cross sections for, (T), 26:46529

interactions $\gamma + p + p + \text{eta}$, factorization of Regge residues for backward differential cross sections for, (T), 26:46529

interactions $\gamma + p + p + \phi$, broken-SU(3) quark model and vector dominance predictions of backward differential cross sections for, (T), 26:46529

interactions $\gamma + p + p + \pi^0$ at 426 to 918 MeV, asymmetry measurements for π^0 from polarized, (E), 26:46442

interactions $\gamma + p + p + \pi^+ + \pi^- + \pi^0$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production in, (E), 26:46446

interactions $\gamma + p + p + \pi^+ + \pi^-$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production in, (E), 26:46446

interactions $\gamma + p + p + \rho^0$, factorization of Regge residues for backward differential cross sections for, (T), 26:46529

interactions $\gamma + p + \pi^- + \Delta^{++}$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446

interactions $\gamma + p + \pi^+ + n$ at 300 to 900 MeV in polarized butanol target, asymmetry of cross section for, (E), 26:46444

interactions $\gamma + p + \pi^+ + n$ at 5 and 16 GeV using polarized proton target, review of asymmetry in, (E), 26:46436

interactions $\gamma + p + \pi^+ + \pi^+ + \pi^- + n$ at 4.3 to 7.5 GeV, cross sections for nonstrange boson resonance production, (E), 26:46446

interactions $\gamma + p + \rho^0 + \text{anything}$ at 4 to 18 GeV, diffractive fragmentation of incident and target particles in, (E), 26:46437

interactions $\gamma + p + \rho^0 + p$ at 2.8 and 4.7 GeV in hydrogen bubble chamber, t dependence of mass distributions, spin dependence, and cross sections for, (T), 26:46438

interactions $\gamma + p + \rho^0 + p$ at 4.3 to 7.5 GeV, cross section for, (E), 26:46446

interactions $\gamma + p + \rho^0 + p$ in geometric diffraction picture, slope of cross sections for virtual, 26:46564

interactions $\gamma + p + \rho^- + \Delta^{++}$ at 4.3 to 7.5 GeV, cross sections for, (E), 26:46446

interactions $\gamma + p + \text{vector mesons} + \text{anything}$, quark model predictions for decay angular distributions of vector mesons from, (T), 26:46536

interactions $\gamma + p$ in $e + p$ interactions at 5 + 200 GeV in colliding beams in ISABELLE, feasibility of experiments on particle production in, (E), 26:46426 (BNL-16968)

interactions $\gamma + p$, fixed pole in amplitude for virtual Compton, (T), 26:46537

interactions $K + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462

interactions $K^- + p + \text{anti } K^0 + n$ with octet exchange in crossed channel, models, for extrapolation of differential cross sections from experimental points for, (T), 26:46630

interactions $K^- + p + \text{anti } K^0 + n$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642

interactions $K^- + p + \text{anti } K^0 + n$, modified absorptive Regge cut model for, (T), 26:46622

interactions $K^- + p + \text{anti } K^0(K^-) + n(p)$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635

interactions $K^- + p + K^- + p + \pi^+ + \pi^- + \gamma(\text{eta})$ at 3.9 and 4.6 GeV/c, production of $M(953)$ and $\phi(1019)$ in, (E), 26:46471 (BNL-16954)

interactions $K^+ + p + K^+ + p$ at 0.52 to 1.89 GeV/c, partial-wave amplitudes and exotic resonance in $P_{\text{sub}} 3/2$ partial wave from phase shift analysis of, (T), 26:46632

interactions $K^+ + p + K^+ + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions $K^- + p + K^- + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions $K^+ + p + K^+ + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + K^+ + K^-$ interactions and, (E), 26:46463

interactions $K^- + p + K^- + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + K^+ + K^-$ interactions and, (E), 26:46463

interactions $K^- + p + K^- + \omega$ at 3.95 GeV/c, differential cross sections and ω density matrix elements for, (E), 26:46485

interactions $K^- + p + \Lambda + \rho^0$ at 3.95 GeV/c, differential cross sections and ρ^0 density matrix elements for, (E), 26:46485

interactions $K^- + p + \Lambda + \rho^0$ at 3.95 GeV/c, differential cross sections and ρ^0 density matrix elements for, (E), 26:46485

interactions $K^+ + p + p + K^+ + \omega$, $\omega + \pi^- + \pi^+ + \pi^0$ at 12 GeV/c, cross sections, momentum-transfer spectra, and angular-decay spectra for, (E), 26:46484

interactions $K^+ + p + p + K^+ + \phi$, $\phi + K^- + K^+$ at 12 GeV/c, cross sections, momentum-transfer spectra, and angular-decay spectra for, (E), 26:46484

interactions $K^- + p + \pi^- + \Sigma^+$ at 5 GeV/c, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut collision mechanism for cross sections for, (T), 26:46631

interactions $K^- + p + \pi^- + \Sigma^+$, duality-breaking effect in

- exchange-degenerate model for, (T), 26:46642
- interactions $K^- + p + E^0 + \phi$ at 3.95 GeV/c, differential cross sections and ϕ density matrix elements for, (E), 26:46485
- interactions $K^- + p + E^0 + \rho$ at 3.95 GeV/c, differential cross sections and ρ density matrix elements for, (E), 26:46485
- interactions $K^- + p$ at 10 GeV/c in bubble chamber, review of elastic and total cross sections, dynamics of two-body, quasi-two-body, and multibody reactions, and Ω^- and resonance production in, (E), 26:46480
- interactions $K^+ + p$ at 5 GeV/c in hydrogen bubble chamber, systematic methods for particle track analysis of, (E), 26:45651
- interactions $K^- + p$, Pomeranchuk theorem violation in total cross sections of nonforward, (T), 26:46634
- interactions $K^+ + p$, Pomeranchuk theorem violation in total cross sections of nonforward, (T), 26:46634
- interactions $n + p$ at 24 MeV, differential elastic, 26:46872
- interactions $\nu + p + \nu + p + \nu^0$ in optical spark chamber, upper limit on neutral current in, (E), 26:46453
- interactions $\nu + p + \nu + p + \nu^0$ in Weinberg model, bound on cross sections for, 26:46614
- interactions $\nu + p$ in propane bubble chamber, test of PCAC from differential cross sections for inelastic, (E/T), 26:46450
- interactions $p + d$ in Glauber theory, differential cross sections, polarization, and asymmetry for, (T), 26:46655
- interactions $p + n$ in final state of $d + p + 2p + n$ interactions at 52.3 MeV, angular distributions and scattering lengths for, (E), 26:46457 (KFK-1509)
- interactions $p + p + \text{anti } p + \text{anything}$ at 24 and 1500 GeV/c, triple-Regge couplings for inclusive distributions in central region for, (T), 26:46661
- interactions $p + p + \text{anti } p + \text{anything}$, Regge analysis of inclusive spectra in, (T), 26:46659
- interactions $p + p + \gamma + \text{anything}$ at high energies, average ν multiplicities from, (T), 26:46662
- interactions $p + p + K^- + \text{anything}$ at 24 GeV/c, triple-Regge couplings for inclusive distributions in central region for, (T), 26:46661
- interactions $p + p + p + \text{anything}$, Regge analysis of inclusive spectra in, (T), 26:46659
- interactions $p + p + p + p + \pi^+ + \pi^- + X^0$ at 19 GeV/c, production of $p\pi^-X^0(2700)$ enhancement in, (E), 26:46467
- interactions $p + p + p + p + \pi^+ + \pi^-$ at 13.1 GeV/c, multi-Regge models for, (T), 26:46656
- interactions $p + p + p + p$ at 200 GeV at high transverse momentum, proposed experiment for spectrometer study of upper limit for cross section for determination of regions of, (E), 26:46456 (COO-1112-75)
- interactions $p + p + p + p$ at high energies, Pomeron amplitude for, (T), 26:46660
- interactions $p + p + p + p$, inclusive finite mass sum rules for upper bound for magnitude of ρ and A_2 Regge cuts in forward, (T), 26:46651 (RPP/T-18)
- interactions $p + p + \pi^0 + \text{anything}$ from $p + p + \gamma + \text{anything}$ interactions, invariant cross sections for, (T), 26:46662
- interactions $p + p + \pi^- + \text{anything}$ at 24 to 1500 GeV/c, triple-Regge couplings for inclusive distributions in central region for, (T), 26:46661
- interactions $p + p$ at 67 GeV/c, two-particle correlations for inclusive, (E/T), 26:46460
- interactions $p + p$ at asymptotic energies, comparison of two-particle distribution functions in diffractive-excitation and multiperipheral models for pion-pion correlation in, (T), 26:46587 (RLD-2230-T-4-1)
- interactions $p + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462
- interactions $p + p$ at high energies, strong-interaction proton form factor from elastic, 26:46654
- interactions $p + p$ in final state of $d + p + 2p + n$ interactions at 52.3 MeV, scattering length for, (T), 26:46650 (KFK-1547)
- interactions $p + p$ in final state of $d + p + 2p + n$ interactions at 52.3 MeV, angular distributions, scattering lengths, and model for, (E/T), 26:46457 (KFK-1509)
- interactions $p + p$ in GeV range, review of differential cross sections and angular distributions and cross sections for resonance production in elastic and inelastic, (E), 26:46464
- interactions $p + p$ in two-fireball model, Reggeon exchange contributions to, 26:46663
- interactions $p + p$, asymptotic behavior of cross section for inclusive, (T), 26:46657
- interactions $p + p$, invariant method for differential cross sections, angular distributions, and two-photon mechanism for meson production in, (T), 26:46518 (ITEF-848)
- interactions $\pi + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462
- interactions $\pi + p$, low lying singularities for energy dependence of cross sections for angular momentum plane structure for amplitudes of line-reversed, (E/T), 26:46629
- interactions $\pi^- + p + 2\pi^0 + n$ at 1.60 to 2.40 GeV/c, total cross sections, peripheral dipion production distribution and scattering, and $A^0(1236)$ production in, (E), 26:46477 (LBL-378)
- interactions $\pi^- + p + \pi + n$, duality-breaking effect in exchange-degenerate, (T), 26:46642
- interactions $\pi^- + p + \pi + n$, modified absorptive Regge cut model for, (T), 26:46622
- interactions $\pi^- + p + \pi + \phi$ at 1.8 GeV/c, branching ratio for $\phi \rightarrow \pi + \gamma$ decay from production in, (E), 26:46435
- interactions $\pi^- + p + \pi(X) + n$ Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- interactions $\pi^+ + p + K^+ + \pi^+$ at 5 GeV/c, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut mechanism for cross sections and polarization for, (T), 26:46631
- interactions $\pi^+ + p + K^+ + \pi^+$, duality-breaking effect in exchange-degenerate model for, (T), 26:46642
- interactions $\pi^- + p + \pi^0 + n$ at 1.6 and 2.4 GeV/c, scintillation counter study of resonance production in, (E), 26:46475 (IS-2886)
- interactions $\pi^- + p + n + \pi^0$ at 2.5 GeV/c, enhancement at 968 MeV in, (E), 26:46430 (COO-1195-241)
- interactions $\pi^- + p + n + \pi^0$ at 22.6 MeV, scintillation counter study of time-of-flight spectrum from, p. 221-3, (E), 26:46873(R) (CEA-N-1522)
- interactions $\pi^- + p + p + \pi^0$ at 25 and 40 GeV/c, cross section scaling in $s^{1/2}$ and triple-Pomeranchuk coupling in, (T), 26:46616 (BNL-16928)
- interactions $\pi^+ + p + p + \pi^+$ at 6 GeV/c, polarization parameter in backward, (E), 26:46490
- interactions $\pi^- + p + p + \pi^-$ at 6 GeV/c, polarization parameter in backward, (E), 26:46490
- interactions $\pi^+ + p + \pi^0 + \Delta^{++}(1236)$ at 1820 to 2090 MeV, partial-wave analysis of, (E), 26:46495
- interactions $\pi^- + p + \pi^0 + n$ at 5.0 to 8.0 GeV/c, polarization parameter in, (E), 26:46491
- interactions $\pi^- + p + \pi^0 + n$ at high energies, properties of complex-conjugate Regge poles from multiperipheral unitarity for pole-cut collision mechanism for cross sections and polarization for, (T), 26:46631
- interactions $\pi^- + p + \pi^0 + n$ at high energies, total cross sections for, (T), 26:46618 (IFVE-STF-71-71)
- interactions $\pi^- + p + \pi^0 + n$, 3-channel helicity amplitudes without pomeron for, (T), 26:46637
- interactions $\pi^- + p + \pi^0 + n$, duality-breaking effect in exchange degenerate model for, (T), 26:46642
- interactions $\pi^- + p + \pi^0 + n$, modified absorptive Regge cut model for, (T), 26:46622
- interactions $\pi^- + p + \pi^0 + n$, Pomeranchuk theorem violation for differential cross sections for, (T), 26:46645
- interactions $\pi^- + p + \pi^0(\pi) + n$ with octet exchange in crossed channel, models for extrapolation of differential cross sections from experimental points for, (T), 26:46630
- interactions $\pi^- + p + \pi^0(\pi^-) + n(p)$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635
- interactions $\pi^- + p + \pi^+ + 2\pi^- + p$ at 11 to 40 GeV in bubble and spark chambers, A_3 production in, (E), 26:46473 (COO-1195-239)
- interactions $\pi^+ + p + \pi^+ + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492
- interactions $\pi^- + p + \pi^- + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492
- interactions $\pi^+ + p + \pi^+ + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + \pi^+ + \pi^-$ interactions and, (E), 26:46463
- interactions $\pi^- + p + \pi^- + p$ at 5 GeV/c, crossing relations from differential cross sections for backward anti $p + p + \pi^+ + \pi^-$ interactions and, (E), 26:46463
- interactions $\pi^+ + p + \pi^+ + p$ at low energies, Veneziano formula for dips in cross sections for, (T), 26:46635
- interactions $\pi^- + p + \pi^+ + \pi^- + n$ at 15 GeV/c, vector dominance and ρ - ω interference in, (E), 26:46496
- interactions $\pi^+ + p + \pi^+ + \pi^+ + \pi^- + \pi^0 + p$ at 5 GeV/c, evidence for double diffraction dissociation in, (E), 26:46493
- interactions $\pi^+ + p + \pi^+ + \pi^+ + \pi^- + \pi^+ + n$ at 5 GeV/c, evidence for double diffraction dissociation in, (E), 26:46493
- interactions $\pi^- + p + \pi^- + \pi^0 + n(p)$ at 4 GeV/c, effective mass and angular distributions and differential cross sections for $\Delta^-(1236)$ and $\Delta^+(1236)$ forward production in, (E), 26:46488
- interactions $\pi^- + p + \rho^0 + n$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481
- interactions $\pi^- + p + \rho^- + p$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481
- interactions $\pi^+ + p + \rho^+ + p$ at 6 GeV/c in bubble chamber, effects of meson (π) exchange on differential cross sections and density-matrix elements from, (E), 26:46481
- interactions $\pi^- + p$ at 4 GeV, spectrum symmetry of pion

longitudinal moments in, (T), 26:46619 (JINR-P1-6461)

interactions $\pi^- + p$ at 6 and 16 GeV, intermediate energy contribution to dispersion sum rules and rotation of polarization in Regge pole model for, (T), 26:46617 (IFVE-STF-71-15)

interactions $\pi^- + p$ at 60 GeV/c, two-particle correlations for inclusive, (E/T), 26:46460

interactions $\pi^- + p$ at high energies, interpolation curves for total cross sections for elastic, elastic with charge exchange, and inelastic, (E), 26:46478(T) (ORNL-tr-2544)

interactions $\pi^+ + p$ at high energies, interpolation curves for total cross sections for elastic, elastic with charge exchange, and inelastic, (E), 26:46478(T) (ORNL-tr-2544)

interactions $\pi^+ + p$ in dual absorptive model, Pomeron and f exchange amplitudes in elastic, (T), 26:46639

interactions $\pi^- + p$ in dual absorptive model, Pomeron and f exchange amplitudes in, (T), 26:46639

interactions with hadrons at 10 and 14 GeV/c using polarized proton and deuteron targets, review of elastic, (E), 26:46436

magnetic moment of, review and analysis of data on relation for, (E), 26:46439

mass of, review and analysis of data on ratio to electron for, (E), 26:46439

meson (K_1^0) regeneration on, modifications in Regge pole model for amplitude of, (T), 26:46633

observations of solar, from HEOS-1 launched Dec. 5, 1968, (E), 26:46334

production in proton reactions with nuclei at 24 GeV/c, spectra of, (E), 26:46459

reactions (p, α) with cobalt-59, nickel-62, copper-63, and copper-65 at 13 MeV, α spectra from, (E), 26:46932

reactions (p, α) with sodium-23 at 1.4 to 3 MeV, resonance structure in excitation functions for, (E), 26:46903(R) (COO-1120-149)

reactions (p, d) with chlorine-35 at 40 MeV, DWBA analysis of angular distributions for, (E), 26:46906

reactions (p, d) with oxygen-18 at 24.5 MeV, DWBA analysis of differential cross sections and asymmetries from polarized, p. 116-9, (E), 26:46873(R) (CEA-N-1522)

reactions (p, d) with scandium-45 at 24.5 MeV, nonlocal finite-range DWBA analysis of differential cross sections and spectrum from, p. 107-9, (E), 26:46873(R) (CEA-N-1522)

reactions (p, f) with uranium-238 at 16 and 42 MeV, angular distribution of long range alpha particles from, (E), 26:47037

reactions (p, γ) with calcium-44 at 1658 keV, gamma spectra from, (E), 26:46920

reactions (p, γ) with carbon-12 in steel, use in measurement of carbon distribution, 26:45466 (AE-451)

reactions (p, γ) with deuterium at 10 to 17.5 MeV, excitation function for, (E), 26:46868

reactions (p, γ) with magnesium-24 at 200 to 1700 keV, gamma angular distributions and spectra and Q value for, (E), 26:46910

reactions (p, γ) with nitrogen-14 at 3.1 to 3.5 MeV, γ spectra and excitation function for, (E), 26:46903(R) (COO-1120-149)

reactions (p, γ) with sodium-23 at 1.8 to 2.8 MeV, γ yields for, (E), 26:46903(R) (COO-1120-149)

reactions (p, γ) with sulfur-32 at 3.36 to 5.41 MeV, excitation functions and gamma spectra from, (E), 26:46902 (BARC-598)

reactions ($p, He-3$) and (p, t) with fluorine-19 at 45 MeV, cross sections and angular distributions for, (E), 26:46895

reactions ($p, He-3$) with lithium-6 up to 125 MeV (cms), calculation of differential cross sections using unified microscopic theory, (T), 26:47077

reactions (p, n) with aluminum-27 at 20.3 MeV, DWBA analysis of differential asymmetry from polarized, p. 120-3, (E), 26:46873(R) (CEA-N-1522)

reactions (p, n) with gold-197, production of carrier-free mercury-197 by, 26:45395

reactions (p, n) with intermediate and heavy nuclei with subsequent proton decay, coupled-channels calculations of energy dependence of, (T), 26:47073

reactions (p, n) with oxygen-18, chlorine-37, potassium-41, chromium-53, and cobalt-59 at 2.4 to 4.0 MeV, angular distributions from, (E), 26:46929

reactions (p, n) with tin isotopes at subthreshold energies, shell-model analysis of, (T), 26:46952

reactions (p, t) with intermediate nuclei with $K = 0$ excitation, analysis using pairing plus quadrupole model, (T), 26:47062

reactions with aluminum at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

reactions with beryllium at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

reactions with boron carbide at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

reactions with copper at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

reactions with lead at 24 GeV/c, π , K, p, and anti p spectra from, (E), 26:46459

reactions with nuclei at 660 MeV, electromagnetic mass separator for isolation of isobars from spallation,

26:46985

reactions with nuclei at high energy, survey on direct, 26:46891

scattering by $A = 20$ to 66 nuclei at 1 to 3.3 MeV, fine structure of isobaric analog resonances in elastic, (E), 26:46914

scattering by beryllium-9 at 6.36 and 6.48 MeV, angular distributions for elastic and inelastic, (E), 26:46885

scattering by calcium-40 at 2.36 to 2.78 MeV, polarization from polarized elastic, (E), 26:46924

scattering by carbon-12 and -13 at 24.5 MeV, optical-model analysis of angular distributions and polarizations from polarized elastic, p. 134-7, (E), 26:46873(R) (CEA-N-1522)

scattering by carbon-12 at 24.5 MeV, coupled-channels and DWBA analyses of differential cross sections and asymmetries from polarized inelastic, p. 138-42, (E), 26:46873(R) (CEA-N-1522)

scattering by carbon-12, calcium-40, iron-56, and nickel-58 at 40 MeV, DWBA and coupled channel analysis of inelastic, (E), 26:46894

scattering by chromium-50, -52, and -54 at 6 MeV, proton spectra and angular distributions for elastic and inelastic, (E), 26:46922

scattering by cobalt-59, nickel-62, copper-63, and copper-65 at 13 MeV, proton spectra from inelastic, (E), 26:46932

scattering by deuterium at 14 to 17.6 MeV, differential cross sections for elastic, (E), 26:46863

scattering by neodymium-145 at 8 to 14 MeV, cross sections for inelastic, p. 34-5, (E), 26:46873(R) (CEA-N-1522)

scattering by nitrogen-14 at 3.1 to 3.5 MeV, excitation function for inelastic, (E), 26:46903(R) (COO-1120-149)

scattering by nitrogen-15 and oxygen-18 at 24 MeV, DWBA analysis of differential cross sections and asymmetries from polarized inelastic, p. 123-9, (E), 26:46873(R) (CEA-N-1522)

scattering by silicon-28 at 1.5 to 3.0 MeV, polarization from polarized elastic, (E), 26:46924

scattering by sodium-23 at 1.4 to 3 MeV, resonance structure in excitation functions for elastic and inelastic, (E), 26:46903(R) (COO-1120-149)

scattering by sulfur-32 and -34 and samarium-148 and -152 at 24.5 MeV, coupled-channels analysis of differential cross sections and asymmetry from polarized inelastic, p. 129-34, (E), 26:46873(R) (CEA-N-1522)

structure of bound, two-pion contribution to electromagnetic, (T), 26:46867

PROTONS (anti p)

interactions anti $p + p + \text{anti } n + n$ at 1.00 to 3.17 GeV/c, forward differential cross section in, (E), 26:46470

interactions anti $p + p + \text{anti } p + p$ at 1.6 and 2.5 GeV/c, asymmetries in polarized, (E), 26:46468

interactions anti $p + p + \text{anti } p + p$ at 5 GeV at large angles, wire spark chamber telescope study of complete angular distributions and cross sections for, (E), 26:46492

interactions anti $p + p + K^+ + K^-$ at 5 GeV/c, crossing relations from differential cross sections for backward $K^+ + p + K^+ + p$ and $K^- + p + K^- + p$ interactions and, (E), 26:46463

interactions anti $p + p + \pi^+ + \pi^-$ pions, limiting behavior of partially inclusive, (T), 26:46613

interactions anti $p + p + \pi^+ + \pi^-$ at 1.6 GeV/c, asymmetries in polarized, (E), 26:46468

interactions anti $p + p + \pi^+ + \pi^-$ at 5 GeV/c, crossing relations from differential cross sections for backward $\pi^+ + p + \pi^+ + p$ and $\pi^- + p + \pi^- + p$ interactions and, (E), 26:46463

interactions anti $p + p$ at 1.51 to 2.90 GeV/c in bubble chamber, cross sections for, (E), 26:46469

interactions anti $p + p$ at high energies and large angles, polarization parameters for backward, (E/T), 26:46462

interactions anti $p + p$ at rest, resonance production in, (E), 26:46465

interactions anti $p + p$, low lying singularities for energy dependence of cross sections for angular momentum plane structure for amplitudes of line-reversed, (E/T), 26:46629

magnetic moment of, determination from fine-structure splittings in atoms formed by antiprotons with lead and uranium, (E), 26:46440

production in proton reactions with nuclei at 24 GeV/c, spectra of, (E), 26:46459

PUBLIC HEALTH

radiation hazards to human, from environmental radioactivity, 26:45911 (COO-3013-25)

radiation hazards to, from operation of nuclear power reactors, assessment of, 26:45904 (COO-3013-1)

radiation hazards to, from radioisotopes in human environment, 26:45890 (NYO-3970-29)

PULSARS

gamma-ray pulses from NP 0532, (E), 26:46199

glitches in NP 0532, gamma radiation following, (E), 26:46214

mass of Crab Nebula and Vela, repulsive interaction in universal gravitation for, (T), 26:46749 (ORO-3992-94)

mass of, repulsive interaction in universal gravitation for, (T), 26:46749 (ORO-3992-94)

circuits for twin time-to-amplitude, for Ge(Li)
detectors, p. 70-3, 26:46873(R) (CEA-N-1522)
dead time of amplitude, calculation of, 26:45684
design and performance of twin time-to-amplitude, for
Ge(Li) detectors, p. 70-3, 26:46873(R) (CEA-N-1522)

PULSE DISCRIMINATORS

design and performance of, using MECL III integrated
circuits, p. 233-4, 26:46873(R) (CEA-N-1522)

PULSE GENERATORS (ELECTRONICS)

circuits for, p. 156, 26:46873(R) (CEA-N-1522)
design for calibration of charge integrators, p. 234-6,
26:46873(R) (CEA-N-1522)

PUMPS

flow through high-head, radioisotope tracer measurements
of, 26:45721

PUREX PROCESS

flowsheet development for LMFBR fuels, 26:45406(R)
(ORNL-TM-2871)

flowsheet development for LMFBR fuels, 26:45408(R)
(ORNL-TM-2949)

flowsheet development for LMFBR fuels, 26:45409(R)
(ORNL-TM-2996)

solvent stability in, 26:45407(R) (ORNL-TM-2918)

PYRENE

radioinduced fluorescence in benzene and cyclohexane
solutions, effects of concentration on, 26:45328
radioinduced fluorescence in cyclohexane, effects of
scavengers on electron pulse, 26:45331
radiolysis in benzene and cyclohexane solutions, effects
of intersystem crossing from excited singlet state on
yield of triplet state from pulse, 26:45329
radiolysis in benzene and cyclohexane solutions, effects
of concentration on yield of excited triplet states
from pulse, 26:45328

PYRIMIDINES

spectra of derivatives of, absorption and fluorescence,
26:45606



Q DEVICES

drift instabilities in, effects of parallel wavelength on
collisional, (E), 26:47168

QUAD CITIES POWER REACTOR, UNIT 1

circulating water system for, operation of discharge
diffuser pipes of, 26:47421 (DOCKET-50254-112)

condenser pump room for, modifications to, 26:47420
(DOCKET-50254-110)

containment structure drywell for, inspection of torus
section headers of, 26:47402 (DOCKET-50237-244)

control rod worth minimizer for, performance of,
26:47358 (DOCKET-50237-254)

core isolation cooling system for, failure of motor
operated valve for, 26:47363 (DOCKET-50254-105)

safety of, requested additional data on, 26:47418
(DOCKET-50254-107)

steam condenser pump room for, modifications for,
26:47364 (DOCKET-50254-108)

technical specifications for, revised and additional data
on, 26:47419 (DOCKET-50254-109)

QUAD CITIES POWER REACTOR, UNIT 2

circulating water system for, operation of discharge
diffuser pipes of, 26:47421 (DOCKET-50254-112)

containment structure drywell for, inspection of torus
section headers of, 26:47402 (DOCKET-50237-244)

control rod worth minimizer for, performance of,
26:47358 (DOCKET-50237-254)

core isolation cooling system for, failure of containment
isolation valve in, 26:47335 (DOCKET-50265-100)

core isolation cooling system for, failure of inboard
steam supply valve for, 26:47371 (DOCKET-50265-96)

electrical cable trays for, fire in, 26:47370 (DOCKET-
50265-95)

safety of, requested additional data on, 26:47418
(DOCKET-50254-107)

steam condenser pump room for, modifications for,
26:47364 (DOCKET-50254-108)

technical specifications for, revised and additional data
on, 26:47419 (DOCKET-50254-109)

waste processing system for, failure of timer in off gas,
26:47325 (DOCKET-50265-101)

QUALITY CONTROL

method for fuel cladding, 26:45472(P)
method for procurement of equipment for WAK radiochemical
plant, 26:46105(T) (ORNL-tr-2371)

QUANTUM ELECTRODYNAMICS

bound states in, for scattering of electrons, atoms, and
molecules, 26:46721

elementary length from photon Green function in nonlocal,
26:46698 (ITF-71-128-P)

elementary length value, 26:46711

gauge invariant regularization of, method of, 26:46719

Green function in nonlocal, form factors and photon,
26:46711

Hamiltonian in scalar, lower unboundedness of, 26:46702
(JINR-P2-6474)

photon Green function in nonlocal, for elementary length,
26:46698 (ITF-71-128-P)

regularization of, Fermi weak-coupling constant for,

26:46722

review on recent developments in, 26:46710

S-matrix elements of Euclidean, perturbation series
convergence for arbitrary values of coupling constant
from equations for, 26:46739

test from x-ray emission in muonic mercury and thallium-
203, 26:46708

transformation properties of operators in, 26:46745

vector potentials in, measurement of, 26:46542

QUANTUM FIELD THEORY

Bethe-Salpeter equation for T matrix, crossing
symmetric, 26:46727

commutation relations in $(\phi^4)_3$ model of, independence
of space cutoff for algebras for equivalence of non-
Fock representations of canonical, 26:46715

commutator between dilatation charge and field in
lambdaphi⁴, equal-time and equal light-cone, 26:46743

coupling constant in, $(\phi^4/\sup 2n)_2$ field Hamiltonian
for complex, 26:46718

dilatations in Minkowski space in, unitary
representations on null plane for unitary
representations of, 26:46716

dual amplitudes in, 26:46726

eikonal representation of amplitudes in scalar,
corrections to, 26:46699 (JINR-P2-6394)

energy-momentum density tensor in conformal invariant
bootstrap, 26:46697 (DESY-72/20)

equation curl $W = 0$ for fields with vanishing curl,
necessary and sufficient conditions for existence of
local operator solution of, 26:46717

field operators on null plane in relativistic, existence
of, 26:46716

fields on 8-dimensional homogeneous space of Poincare
group for electric and magnetic form factors of NN γ
vertex in, 26:46514 (CTH-ITP-71-30)

formulation for elementary particles with nonzero mass,
covariant, 26:46723

formulation of lightlike quark, current densities in,
26:46524 (NYO-4076-18)

formulation of translation-invariant ϕ^4 , in three
dimensions, independence of space cutoff for algebras
for, 26:46715

gauge problem in, vanishing of electromagnetic field
tensor in, 26:46720

Haag theorem in null plane formulation of relativistic,
26:46716

Hilbert spaces in, charge operators for rigged, 26:46737

Klein-Gordon type equations in, conformal invariance of
zero-mass, 26:46731

Lagrangians for free massless fields in, third-order
perturbation theory for exponential, 26:46706

Lagrangians in, factor-ordering problems in nonlinear,
26:46744

Lorentz invariance of Wightman functions in, relation
between spectrum conditions and, 26:46707

particle-field interaction in, Bogoliubov transformation
in strong coupling theory for, 26:46691

potential equations in, conception of straight-line paths
in high energy limits for operator method of solving
quasi-, 26:46700 (JINR-P2-6437)

potential parameters in local, asymptotic estimates of
quasi-, 26:46703 (JINR-P2-6482)

Reggeon propagator, vertex function, form factor, and
renormalization constants in Zachariasen model

formulation, perturbation theory for intermediate,
26:46747

renormalization by continuous space dimension, analytic,
26:46729

renormalization of exponential interaction Lagrangians
in, analytic, 26:46712

renormalized ϕ^4 interactions in three-dimensional
space-time in, independence of space cutoff in
formulation of algebra of observables for, 26:46715

representations of Weyl algebra for model, equivalence of
dressing transformations for, 26:46713

scalar fields in relativistic, dressed operators for
charged and neutral ϕ , 26:46701 (JINR-P2-6454)

scattering amplitudes in scalar, corrections to eikonal
representation of, 26:46699 (JINR-P2-6394)

scattering operator in $(\phi^4)_2$, unitarity, causality,
and generalization to test function space for
Bogoliubov, 26:46740

Schwinger equation in, method of characteristics for
functional-derivative equation for, 26:46728

spinor fields in, functional definition and calculation
of global observables of nonlinear, 26:46746

state vectors in VTheta sector, congruence transformation
for algebraic solution for Kaelen-Pauli, 26:46705

symmetry breakdown in interaction picture for preferred
fields, 26:46676

symmetry currents in conformal invariant bootstrap,
matrix elements of internal, 26:46697 (DESY-72/20)

symmetry problems in, 26:46674

Thirring model in, exact expansions of operator products
in terms of C-number functions singular on light cone
and regular operators for calculation of Wightman
functions in terms of current matrix elements for
complete solution to, 26:46696 (CONF-710816-19)

unitarity in conformal invariant bootstrap, generalized,
26:46697 (DESY-72/20)

- unitary equivalence of different fields in null plane formulation of relativistic, 26:46716
vacuum polarization calculations in regularization in, 26:46742
vertex functions from local commutativity in scalar, one-dimensional dispersion relations for, 26:46736
vertex functions from local commutativity in, two-dimensional dispersion relations for, 26:46735
Wick--Cutkosky model in, Regge poles for large coupling constants in, 26:46725
Zachariasen model formulation for perturbation theory for intermediate Reggeon propagator, vertex function, form factor, and renormalization constants in, 26:46747

QUANTUM MECHANICS

- algebraic method for solvable problems in, 26:46671
axioms of, lattice postulate in, 26:47295
book: Lectures on Quantum Statistics. Volume 2. Quasi-Averages, 26:47299
expectation values from hypercharacteristic function in, 26:47289
formulation of nonrelativistic, incompatibility of gauge invariance and locality in path integral, 26:47292
formulation of nonrelativistic, with deterministic quantum processes and usual statistical properties, review of generalization in, 26:47297
Hartree--Fock equations in, quasi-classical approximation for solution of system of integro-differential, (T), 26:47288
Lagrangian in, variational principle for nonlinear, 26:47298
many-body systems in, unitary operator form of general Bogolubov--Tyablikov transformation for, 26:47291
measurements of first kind in, covering law in quantal proposition systems for ideal, 26:47294
parameters in, theorem on impossibility of introducing hidden, 26:47285 (ITF-71-101-P)
Schrödinger equation in, accuracy and efficiency improvements for method for numerical solution for one-dimensional potential in bound-state, 26:47290
Schrödinger equation in, differential equations as limiting cases and correspondence principle for finite-difference, 26:47293
Schrödinger equation of three-body problem in, method avoiding group theory for solutions in form of power series expansions for, 26:47296
Schrödinger group algebra for many-body system of interacting particles, realizations of, (T), 26:46673
three-body problem in, method avoiding group theory for solutions in form of power series expansions for Schrödinger equation for, 26:47296
wave functions in, formulation of algebra with power asymptotic number elements for operations with nonnormalizable, 26:47284 (IC-72/45)

QUARKS

- existence of, implications of quark light-cone current algebra for, 26:46604
existence of, SU(3) properties for, 26:46683
search for, pseudo cloud chamber tracks for, 26:45647 (UCRL-51214)

QUARTZ

- radiation effects on oscillation frequency of AT-cut, cobalt-60 gamma, 26:45667(T) (SC-T-722519)
radioinduced frequency change in 125 MHz AT-cut oscillators of, rapid annealing of, 26:45665 (SC-DC-721240)

QUASARS

- associations between galaxies and, physical, (E/T), 26:46258
evolution of, (E/T), 26:46208
luminosity--spectral index relation for, (E/T), 26:46248
luminosity-volume relation for, (E/T), 26:46208
radio polarization at 49 and 73 cm, (E), 26:46204
relation of Ton 155 and 156, (E), 26:46210
relation to nearby galaxies, (T), 26:46234
spectra of, in test of time variation of fine-structure constant, (T), 26:46254 (COO-1112-51)
spectral redshifts of Ton 155 and 156, (E), 26:46210
spectral redshifts of, (E/T), 26:46208

QUINOLINE

- 8-mercapto-, radioprotective effects of, on mice, combined effects of AET on, 26:45859
QUINOLINE, 4-NITRO-, 1-OXIDE
effects on DNA in cultured human Xeroderma pigmentosum cells in comparison to effects of uv radiation, extent of repair synthesis following, 26:45742

R

RABBITS

- intravenous catheterization in, procedures for chronic, 26:47641 (UR-3490-21)
plutonium-239 distribution in tissues of, following intravenous injection of monomeric and polymeric forms as citrate and nitrate, comparison of, 26:45917
radiation effects on bones of, pathological changes in blood vessels and bone marrow following exposure of mandible, effects of dose on gamma, 26:45835
radiation effects on enzyme activity in neuroglia of brain of developing, x, 26:45836

- radiation effects on lens cell proliferation and regeneration in eyes of, during injury response and wound healing, x, 26:45813(R) (COO-1427-41)
radioinduced glossitis in, permeability of lingual mucosa during development and recovery from β , γ , and x, 26:45862
radioinduced inhibition of glucose-6-phosphatase activity in erythrocytes of, following whole-body exposure, effects of cysteine and reduced glutathione on extent of recovery from, 26:45848
radioinduced injuries in lungs of, role of allergic inflammations in pathogenesis of, 26:45863
radioinduced injuries to crystalline lens of, effects of dose on RBE for fast neutron, γ , mixed γ -neutron, and x, 26:45864
radioinduced suppression of immune reactions in, to Salmonella typhimurium, protective effects of AET against γ , 26:45824

RADIATION

- book: Radiation Physics of Crystals and p-n Junctions (In Russian), 26:47214
book: Radiation Technology (In Russian), 26:45432
dose distribution in human pelvis from radium applicator implanted in uterus, methods for calculating, 26:45784(T) (TT-54002/4, pp 456-61)
dose to man from radioisotopes in body, compartmental model for determining relation between tissue distribution and retention and, 26:45905 (RD/B/N-2291)
dose to nuclear personnel, correlation between blood picture and absorbed, 26:45787(T) (TT-54002/5, pp 602-4)
dose to passengers and personnel of supersonic aircraft, measurements on board Concorde prototype, 26:45796
dose to personnel in Federal Republic of Germany, legislation on maximum permissible, 26:45803
effects on activity and regulatory functions of enzymes, 26:45726
effects on albumins, mechanisms of sensitivity, 26:45726
effects on animal feeds, pathological effects on rats following ingestion, 26:45899
effects on animals following chronic low dose internal exposure, repair mechanisms at cellular and molecular level for pathological, 26:45911 (COO-3013-25)
effects on bromotrifluoromethane and iodine--bromotrifluoromethane mixtures, yield of products from, 26:45322
effects on butyrophene in benzene solutions, yields from, 26:45326
effects on chromosomes in lymphocytes of Marshall Island human population, combined effects of age and sex on frequency of aberrant, 26:45792
effects on cultured cells, environmental and intrinsic factors affecting sensitivity to lethal, 26:45739
effects on deformation of fuel elements, relations of stress and temperature with in-pile, 26:46070 (UJV-2798-M)
effects on DNA in cells of higher animals and plants in relation to organizational properties of genome, 26:45894 (ORD-2690-12(Pt.B), Paper 14)
effects on doped ice, effects of fluoride ion on yields of trapped electrons from, 26:45345
effects on enzyme activity and RNA synthesis in spinal neurons of mice, 26:45868
effects on ESR spectra of human blood plasma following therapy, 26:45723
effects on grain growth in uranium, relations with statistical fluctuations in texture, 26:46069 (UJV-2795-M)
effects on growth of bone marrow and blood cells in diffusion chamber cultures in host mice, effects on yield of granulocytes and macrophages, 26:45741
effects on growth of bone tissues of rats, effects of dose and parotin on, 26:45870
effects on human marrow cells, protective effects of taurine against, use of chromosome fragmentation counts for determining, 26:45734
effects on leukocyte count in man following therapy of gynecologic neoplasms, protective effects of cepharanthine against, 26:45805
effects on living cells, review of sensitization of, 26:45845
effects on man following therapy of oral tumors, relation to necrosis of mandible and soft oral tissues in, 26:45801
effects on materials, method of correlating data on damaging, 26:46072
effects on mechanical properties of rubber for use in vacuum, 26:46087
effects on metabolism of erythropoietin in rats following whole-body exposure, additive effects of hypoxia on, 26:45854
effects on mice following perinatal exposure, effects of age and strain on incidence of neoplastic, 26:45814(R) (COO-3017-43)
effects on microorganisms contaminating disposable medical supplies, determination of sensitivity to sterilizing dose, 26:45748
effects on mitochondria in rat heart and liver, effects on content of thiol groups in, 26:45829
effects on mitochondria in rat liver, effects on function and structure following lethal whole-body exposure, 26:45838
effects on muscles of rats, effects on capacity to

- regenerate following traumatic injury, 26:45827
effects on nucleic acids, mechanisms of sensitivity, 26:45726
effects on parabiont rats following acute supralethal whole-body exposure, high incidence of neoplastic, 26:45814(R) (COO-3017-43)
effects on peas, review of genetic and toxic, 26:45776
effects on poly(vinyl chloride) and poly(vinylidene fluoride) coatings, 26:45352(P)
effects on rats, effects on deoxyribonuclease activity in liver, effects of cytostatic drugs on, 26:45732
effects on regeneration of blood vessels in muscle transplants in rats, 26:45857
effects on reproduction of Clostridium in packaged sausage following processing, combined effects of heat treatment on inhibitory, 26:45747
effects on spores of microorganisms during use for sterilization, probability of mutagenic, 26:45750
effects on spores, biochemical mechanisms of resistance to, 26:45751
effects on sprout inhibition, storage, and wholesomeness of potatoes, comparison with chemical processing, 26:45716 (BFL-2/1972)
effects on survival of cultured cells, predictions of delta-ray theory of track effects for, 26:45611
effects on survival of cultured cells, predictions of delta-ray theory of track structure for, 26:45612
effects on survival of mammals, review of effects of dose and LET on, 26:45816 (FRNC-R-7)
effects on yeasts, factors affecting sensitivity to lethal, 26:45749
emission from electromagnetic interactions at high energies in condensed media, book on, (T), 26:46538
emission from systems of N two-level atoms, Green-function calculation of cooperative effects in spontaneous, 26:46418
emission from two-level atoms confined within sphere or right circular cylinder, far-field patterns for, (T), 26:46411
hazards to man in relation to operation of nuclear power plants, criteria for establishment of maximum permissible doses on basis of, 26:45778(T) (NP-tr-1941)
metrology of ionizing radiations, 26:47219
theory for total pure, special class of solutions of Lichnerowicz equations and Maxwell equations for singular source-free electromagnetic field for nonspecial recurrent space time for general relativistic, (T), 26:46771
- RADIATION CHEMISTRY**
developments in, 26:45313(R) (COO-3029-1)
research progress on, India, 1970-71, 26:45311(R) (BARC-599)
solvated electron dielectric models in, 26:45344
transient species in, absorption spectroscopy of, 26:45340
- RADIATION DETECTORS**
description and use for measurement of low-level radioactivity, 26:45261 (ICRU-22)
description of resonance, for detection of 3-GeV electrons, 26:45648
response of, delta-ray theory for, 26:45612
response of, delta-ray theory for, 26:45611
- RADIATION DETECTORS, BUBBLE CHAMBER**
bubble density in, determination of, 26:45649
particle tracks in hydrogen, systematic methods for analysis of $K^+ + p$ interaction, (E), 26:45651
- RADIATION DETECTORS, CHERENKOV**
performance testing of threshold plastic, with wavelength shifter and quencher, 26:45646 (RPP/H-89)
preamplifiers for, circuits for, 26:45608
- RADIATION DETECTORS, CLOUD CHAMBER**
particle track analysis in search for quarks, method for pseudo, 26:45647 (UCRL-51214)
- RADIATION DETECTORS, CRYSTAL**
description of diamond, 26:45597(P)
- RADIATION DETECTORS, DIELECTRIC TRACK**
characteristics of fission foil--mica track detectors for neutron dosimetry, 26:45617 (FEI-296)
particle track analysis in, prolonged etching correction factor for, 26:45604
particle track etching conditions for mica and nitrocellulose, for detecting alpha particles and fission fragments, 26:45610
particle tracks in, description of optical system for measuring density of, 26:45598
performance for neutron dosimetry, 26:45630
performance of cellulose-based, for alpha detection, 26:45615
- RADIATION DETECTORS, ELECTRON MULTIPLIER**
description with electrostatic analyzer for electron and proton spectroscopy at 0.03 to 10 keV, 26:45642
operation and performance of, used with electrostatic analyzers for charged particle spectroscopy, 26:45645
- RADIATION DETECTORS, FISSION FOIL**
characteristics of fission foil--mica track detectors for neutron dosimetry, 26:45617 (FEI-296)
- RADIATION DETECTORS, GEIGER-MUELLER**
description of miniature low-background, for measuring argon-37 and argon-39 activities, 26:45614
- RADIATION DETECTORS, IONIZATION CHAMBER**
calibration and operation of flowthrough, for monitoring tritium in urine based on acetylene carrier from calcium carbide reactions, 26:45585 (AERE-HP/M-74)
calibration and performance of thimble, for dosimetry of electron beams at 8 to 30 MeV, 26:45623
description for dosimetry, 26:45635(P)
description for monitoring and controlling radiation exposure from dental x-ray units, 26:45603
design of tissue-equivalent liquid, for dosimetry, 26:45621
drift in dc, device for controlling slow, 26:45609(P)
insulators for pocket, radiation resistant plastic, 26:45672(P)
operation and performance of tissue-equivalent liquid, for dosimetry, 26:45634
performance for dosimetry of pulsed electrons at 10 to 30 MeV, 26:45627
performance for dosimetry of pulsed x rays at 10 to 35 MV, 26:45627
radiation and temperature effects on Radocon, for gamma dose rate measurements, 26:45620
reliability and sensitivity of, for personnel dosimetry, 26:45624
temperature effects on ionization current from, 26:45594
- RADIATION DETECTORS, PROPORTIONAL**
circuits for multi-wire, logic, p. 198-200, 26:46873(R) (CEA-N-1522)
data acquisition system for, circuits for, p. 236-8, 26:46873(R) (CEA-N-1522)
description of miniature low-background, for measuring argon-37 and argon-39 activities, 26:45614
description of spherical tissue-equivalent, for determining depth--dose distributions, 26:45626
design and testing of multiwire, p. 210-7, 26:46873(R) (CEA-N-1522)
design, operation, and performance of moderated array of coincidence BF₃ or ³He, for plutonium-240 determination, 26:45587 (BNL-50332)
development for 1-GeV particles, p. 160-3, 26:46873(R) (CEA-N-1522)
development of multiwire, p. 227-30, 26:46873(R) (CEA-N-1522)
- RADIATION DETECTORS, SCINTILLATION (LIQUID)**
design and performance of NE 323, for photoneutron detection, p. 267-8, 26:46873(R) (CEA-N-1522)
photomultipliers for, use of pulsed gas laser for determining properties of, 26:45640
scintillators for, efficiency of pyrimidine-derivative, 26:45606
- RADIATION DETECTORS, SCINTILLATION (SOLID)**
calibration of Bonner sphere ⁶LiI(Eu), for neutron dosimetry, effects of scattering on, 26:45599
circuits for, logic, p. 197-8, 26:46873(R) (CEA-N-1522)
design for continuous measurement of active uptake of radioactive substances by cells, 26:45592
development of NaI(Tl), for determining plutonium-239 in waste drums, pp.4-6, 26:45673(R) (UCRL-50007-71-3)
directional properties of cylindrical, with sleeve-type lead screen for gamma detection, 26:45605
efficiency of NE110, Pilot F, and Pilot Y, 26:45588 (COO-2114-2)
photomultipliers for, use of pulsed gas laser for determining properties of, 26:45640
scintillator composition for tissue-equivalent plastic, for x rays, 26:45619
- RADIATION DETECTORS, SEMICONDUCTOR**
book: Radiation Protection Measurement, 26:45808
charge collection in, average pulse-amplitude meter for measuring, 26:45685
development of CdTe, GaAs, and HgI₂, 26:45600
- RADIATION DETECTORS, SEMICONDUCTOR (GERMANIUM)**
development of Ge(Li) probe for gamma well logging, 26:45658(P)
performance and recovery of Ge(Li), 26:45586 (ARH-2432)
time measurements with, preamplifier and pulse shaper circuit for fast, 26:45616
- RADIATION DETECTORS, SEMICONDUCTOR (SILICON)**
development of Si(Li) probe for gamma well logging, 26:45658(P)
resolution of Si(Li), for x-ray spectroscopy, 26:45644
- RADIATION DETECTORS, SPARK CHAMBER**
particle track drift compensation using high-voltage radiofrequency pulses, 26:45652
readout system for, on-line performance logic for computer-based, 26:45650
- RADIATION DOSE**
calculations outside shielding at high-energy proton accelerators, Moyer method for, 26:46121
estimation of contribution to, of fluorescence-photon and Auger-electron spectra from electron-capture- and internal conversion decaying nuclei, 26:47209
maximum permissible for human populations resulting from operation of nuclear power plants, criteria for, 26:45778(T) (NP-tr-1941)
recommendations for maximum permissible, to man from environmental contamination, 26:45554
recommendations for maximum permissible, to man from environmental contamination, 26:45555
- RADIATION DOSIMETERS**
description and uses for electron dosimetry at 1 to 50 MeV, 26:45618 (ICRU-21)
for Gemini spacecraft, 26:47203 (N-72-18797)

- response of, delta-ray theory for, 26:45611
response of, delta-ray theory for, 26:45612
- RADIATION DOSIMETERS, BIOLOGICAL**
effectiveness of sulfur activation in human hair and nails for fast neutron, following criticality accident, effects of age, diet, and sex on, 26:45788
- RADIATION DOSIMETERS, CHEMICAL**
container materials for methyl orange, 26:45633
description of FeSO_4 , for dosimetry of bremsstrahlung in GeV range, 26:45625
dose--flux relation for FeSO_4 , for 6.2-GeV bremsstrahlung, 26:45625
readout of alcoholic chlorobenzene, radiofrequency method for, 26:45632
- RADIATION DOSIMETERS, PHOTOGRAPHIC FILM**
calibration and use for radiotherapy treatment verification, 26:45631
characteristics of personal, for x radiation at 10 to 500 keV, 26:45629
performance for gamma dosimetry, 26:45630
reliability and sensitivity of, for personnel dosimetry, 26:45624
- RADIATION DOSIMETERS, PHOTOLUMINESCENT**
performance for gamma dosimetry, 26:45630
reliability and sensitivity of, for personnel dosimetry, 26:45624
- RADIATION DOSIMETERS, THERMOLUMINESCENT**
development of Al_2O_3 , for criticality accidents and nuclear explosions, 26:45628
performance of CaS and LiF , for gamma and neutron dosimetry, 26:45630
performance of LiF and CaF_2 , for x-ray dosimetry in water phantoms, 26:45622
reliability and sensitivity of, for personnel dosimetry, 26:45624
sensitivity to fluorescent and ultraviolet light, pp.13,14, 26:45673(R) (UCRL 50007-71-3)
- RADIATION DOSIMETRY**
book: Radiation Protection Measurement, 26:45808
for Gemini program, 26:47203 (N-72-18797)
methods in radiodiagnosis and radiation therapy rooms, 26:45629
standards for photon, French national, 26:47216
- RADIATION MONITORING**
book: Radiation Protection Measurement, 26:45808
instrumentation criteria for, following reactor accidents, 26:47317 (BNWL-1635)
programs for environmental, near nuclear facilities, development of, 26:45556
- RADIATION PROTECTION**
book: Radiation Protection Measurement, 26:45808
methods for personnel at hot facilities of Pakistan Institute of Nuclear Science and Technology, 26:45499
methods for reducing neutron dose to medical personnel during use of californium-252 for interstitial and inter-cavitary therapy, 26:45790
methods for Virginia dental offices, 26:45500
programs for diagnostic radiography facilities in Canada, 26:45811
programs for environment and personnel at Marcoule nuclear center, 26:45804
programs for nuclear personnel in Federal Republic of Germany, medical aspects of, 26:45803
- RADIATION-SCATTERING ANALYSIS**
developments in elastic, review on, 26:45271
- RADIATION SOURCES**
availability and cost of, produced by Norway Institute for Atomic Energy, 26:45709 (INIS-mf-256)
design of rotating, for pool-type irradiator, 26:45446(P)
uses in industry, medicine, and science, radiation hazards to man from, 26:45555
- RADIATION STANDARDS AND UNITS**
quality factor, evaluation of conservative and realistic quantities for use in expressing recommended limits for neutron exposures of man, 26:45789
use of kilogram-rads as units for expressing cumulative x-ray dose to man from diagnostic radiography, 26:45799
- RADIATION TARGET CANS**
design for in-pile use at 2000°C, 26:45438 (KFK-1556)
testing under EBR-II conditions, design of facility for, 26:45702(R) (LA-4978)
- RADIATION TARGET CHAMBERS**
design for recoil-distance measurements of nuclear lifetimes, p. 66-8, 26:46873(R) (CEA-N-1522)
- RADIATION TARGETS**
design of aluminum-backed and self-supporting carbon-14, p. 93-4, 26:46873(R) (CEA-N-1522)
development of differentially pumped gaseous, for heavy-ion reactions, p. 59-64, 26:46873(R) (CEA-N-1522)
development of polarized proton and deuteron, 26:46436
polarization of protons in polarized, corrections for nonlinearity of dc Q-meter for measuring, 26:45680
thickness optimization of tantalum, for positron production by electron reactions, 26:46149
use of polarized proton and deuteron, in high-energy physics experiments, 26:46436
- RADIO GALAXIES**
luminosity--spectral index relation for, (E/T), 26:46248
- polarization at 49 and 73 cm, (E), 26:46204
- RADIO WAVES**
emission at high and low latitudes, association of micropulsation with quasi-periodic vlf, (E), 26:46319
emission during post-breakup phase of polar substorms, fine structure of vlf, (E), 26:46285
reflection from ionospheric D region, rapid computation method of coefficients for vlf steep-incidence, (E/T), 26:46263 (AD-734392)
- RADIOACTIVE CONTAMINATION**
book: Radiation Protection Measurement, 26:45808
hazards to man from environmental, control and evaluation of, 26:45554
hazards to man from environmental, from peaceful uses of nuclear energy, evaluation of, 26:45555
- RADIOACTIVE MATERIALS**
concentration in solutions, design of equipment for, 26:46111(P)
flow of solutions containing, design of electromagnetic valve for control of, 26:46109(P)
handling of liquid, design of shielded container for visible syringe for, 26:46113(P)
packaging for neutron-emitting, design of, 26:46115(P)
shipping of, German regulations for rail and road, 26:46117
transfer of vials containing, from loading platform to light-tight counting chamber, description of equipment for, 26:46110(P)
- RADIOACTIVE WASTE**
activity of aqueous, in environment and surface waters at Nuclear Research Institute, Rez, Czechoslovakia, 26:45565
analysis for phosphates by extraction of molybdophosphoric acid in butyl acetate in presence of EDTA and fluoride, spectrophotometric, 26:45219 (ARH-SA-110)
analysis for plutonium and uranium, review on methods for, 26:45190 (TID-7029(2nd Ed.), pp 331-411)
analysis for plutonium-239, development of NaI(Tl) scanning system for, pp.4-6, 26:45673(R) (UCRL-50007-71-3)
calcination using in-bed combustion of fuel oil, 26:45405(R) (ICP-1006)
disposal in salt deposits, methods and siting parameters for, 26:46100 (CONF-720903-4)
disposal in sea, International Laboratory of Marine Radioactivity research on contamination from, 26:46101 (IAEA-136, pp 10-17)
encapsulation of high-level, design of remote handling equipment for, 26:46099 (ARH-SA-117)
liquidation at Nuclear Research Institute, Rez, Czechoslovakia, 26:45565
management at ICPP, 26:45405(R) (ICP-1006)
management at Pakistan Institute of Nuclear Science and Technology, 26:45499
management of fuel-reprocessing, in Australia, plans for, 26:46098 (AAEC/TM-628)
processing by fluidized-bed calcination, 26:45405(R) (ICP-1006)
processing of low-level aqueous, at Midwest Fuel Recovery Plant, proposed operation of evaporation system for, 26:45404 (DOCKET-50268-33)
processing of low-level liquid, at West Valley Processing Plant, 26:45400 (DOCKET-50201-127)
processing of organic solutions of, oxidation process for, 26:45386(P)
processing of, decomposition of potassium and sodium nitrates in fluidized-bed, 26:45383 (ICP-1008)
properties of solid calcined, 26:45405(R) (ICP-1006)
release of gaseous, to atmosphere, problems associated with, 26:46102(T) (ORNL-tr-2369)
separation of cesium from liquid, using colloidal suspension, 26:45384(P)
shipping of, design and testing of drum-type shielded container for, 26:46114 (MLM-1911)
solidification in phosphate glass, Ester program research progress on, 26:45385
solidification of concentrated high-salt solution of, using phosphoric acid, 26:45381 (ARH-SA-120)
storage of, description of equipment and methods for, 26:46103(P)
- RADIOACTIVITY**
control and effects of artificial, on environment, 26:45554
distribution in and near Rainier rubble chimney, (E), 26:45483 (NVO-1229-180)
monitoring in Pearl Harbor marine environment during 1966 and 1968, 26:45534
monitoring in US surface waters during Oct. 1971, 26:45537
- RADIOBIOLOGY**
book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
research programs at Florida State University, Tallahassee, during 1971 and 1972, 26:45891(R) (ORO-2690-12(Pt.B))
- RADIOCHEMICAL ANALYSIS**
book: Radiation Protection Measurement, 26:45808
- RADIOCHEMICAL PROCESSING PLANTS**
environmental protection studies at ICPP, 26:45405(R) (ICP-1006)
equipment for German WAK, description of, 26:46106(T) (ORNL-tr-2372)

equipment for German WAK, quality control and specifications for, 26:46105(T) (ORNL-tr-2371)
evaluation for LMFBR fuel reprocessing, 26:45411(R) (ORNL-TM-3072)

evaluation for LMFBR spent fuel reprocessing, 26:45410(R) (ORNL-TM-3018)
evaluation for LMFBR spent fuel reprocessing, 26:45413(R) (ORNL-TM-3127)
evaluation for reprocessing LMFBR spent fuels, 26:45408(R) (ORNL-TM-2949)
fuel storage at ICPP, 26:45405(R) (ICP-1006)
instrumentation for input weight measurements at ICPP, 26:46093 (ICP-1014)
nuclear materials safeguards program at ICPP, 26:45405(R) (ICP-1006)
radioactive gas releases from, problems associated with atmospheric contamination from, 26:46102(T) (ORNL-tr-2369)
radioactive waste management at ICPP, 26:45405(R) (ICP-1006)
research programs and operating experience at ICPP, 26:45405(R) (ICP-1006)

RADIOCHEMISTRY

research progress on, India, 1970-71, 26:45311(R) (BARC-599)

RADIOFREQUENCY OSCILLATORS

radiation effects on AT-cut quartz, cobalt-60 gamma, 26:45667(T) (SC-T-722519)
radioinduced frequency change in 125 MHz AT-cut quartz, rapid annealing of x, 26:45665 (SC-DC-721240)

RADIOGRAPHY, BIOMEDICAL

radiation dose to man from, method and units for expressing cumulative x, 26:45799
radiation effects on incidence of leukemia in man following diagnostic, effects of age, exposure site, number of exposures, and sex on x, 26:45798
radiation monitoring of facilities and personnel for diagnostic, in Canada, 26:45811
safety factors for, in Virginia dental offices, 26:45500

RADIOGRAPHY, INDUSTRIAL

development of neutron, using californium-252 sources, 26:45467 (BNWL-SA-4320)
equipment for, design and performance of, 26:45432
films for gamma and x-ray, sensitometric measurements of image quality of, 26:45468
methods for steel weld crack detection, factors affecting gamma and x-ray, 26:45471

RADIOISOTOPE SCANNERS

description of isosensitive, with multichannel analyzer for quantitative evaluation of radioisotope distribution, 26:45601
description of two-channel Scinticart Numeric T, 26:45607
operation and performance of dual-headed whole-body, for subtraction and positron scanning, 26:45593

RADIOISOTOPES

book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
book: Radiation Protection Measurement, 26:45808
content in Amchitka Island surface waters, effects of Cannikin Event on, (E), 26:45488 (USGS-474-148)
content in nuclear geothermal steam, effects on design and operation of steam-turbine power plants, 26:45487 (UCRL-73939)
determination in soil surface-deposited fallout, performance of continuously operating gamma spectrometer for, 26:45527 (TID-26103)
determination of low-activity, recommendations for, 26:45261 (ICRU-22)
distribution and retention in human tissues following intake by various routes, compartmental model for determining relation between radiation dose and, 26:45905 (RD/B/N-2291)
distribution and retention of aerosol incorporated, in respiratory tract, use of biological model and radioautography for measurements of kinetics of, 26:45921
effects on mammals following uptake, review of carcinogenic, 26:45895 (UR-3490-102)
excitation energies of daughters of beta-decay, tables for mean, 26:46875 (KFK-EXT-10/72-1)
production at Pakistan Institute of Nuclear Science and Technology using swimming-pool reactor, program for, 26:45396
transport by ground water of, contained in Rainier Event melt-glass, measurement of long-term, 26:45482 (NVO-1229-177)
transport in aquatic food chains, mathematical model for calculation of, 26:45547
transport through aquatic ecosystems, relation of retention in plants to, 26:45926
uptake by mammals, pathological changes in cells induced by, review of, 26:45890 (NYO-3970-29)
use as heat sources in direct energy converters in ENEA countries, predictions to 1975 on demand for, 26:45705
use as internal power supply for compact solid-state laser, 26:45677(P)
uses in mechanical engineering, review on, 26:45710
uses in mechanical engineering, review on, 26:45711

RADIOLOGY

translation of Polish Review of Radiology and Nuclear Medicine, volume 35; number 5, 1971, 26:45785(T) (TT-54002/5)

137 SUBJECTS

RARE EARTHS

translation of Polish Review of Radiology and Nuclear Medicine volume 35, number 4, 1971, 26:45779(T) (TT-54002/4)

RADIOMETRIC TESTING

equipment for measuring wear of materials and machine parts, description of two-channel automatic, 26:45657
method for inspection of welded joints in large-capacity boilers, 26:45473

RADIOPROTECTIVE COMPOUNDS

effects of DEAE Sephadex on survival of x-irradiated rats, 26:45844
properties of organic disulfide, 26:45290

RADIOTHERAPY

dosimetry for treatment verification using photographic film, method for, 26:45631
equipment for, design and performance of, 26:45432

RADIUM

separation from thorium, method for, 26:47014 (FRNC-TH-184)

RADIUM ISOTOPES Ra-226

content in plants and soils in uraniferous area in New Zealand, 26:45529
effects on bones following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
energy-level transitions from thorium-230 decay, (E), 26:47028
excretion in human feces following oral uptake as radium sulfate dust, effects of alginate ingestion on, 26:45907
retention in mice following injection, method for direct measurement of, 26:45920
uptake and retention in mice following gastric tube administration, effects of aluminum phosphate gel on, 26:45918

RADIUM ISOTOPES Ra-228

beta decay of, γ and conversion electron spectra from, (E), 26:47014 (FRNC-TH-184)
effects on bones following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
preparation of, by chemical separation from thorium-232, 26:47014 (FRNC-TH-184)

RADON

distribution in atmosphere, use for calculating turbulent diffusion coefficient and vertical component of wind velocity, 26:45522
polarizability of ground-state, electric dipole, (T), 26:46349
release from soils to atmosphere, (E/T), 26:45524

RADON ISOTOPES Rn-206

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-207

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-208

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-209

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-210

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-211

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-212

alpha spectra from, (E), 26:47004

RADON ISOTOPES Rn-222

distribution in tissues in mice following injection of radium-226, kinetics of, 26:45920
dose to respiratory tract epithelial cells following inhalation as aerosol, relation of content of daughter products in air to α , 26:45861

RANCHO SECO POWER REACTOR

environmental report for, revised and additional data for, 26:47479 (DOCKET-50312-41)
pipes for, imperfections in welds for, 26:47481 (DOCKET-50312-43)
piping for, defects in welds for, 26:47484 (DOCKET-50312-47)
safety of, requested additional data on, 26:47482 (DOCKET-50312-45)
safety of, revised and requested additional data on, 26:47480 (DOCKET-50312-42)
steam generator for, weld defects in lateral support for, 26:47485 (DOCKET-50312-48)
steam generator for, weld defects in lateral support for, 26:47483 (DOCKET-50312-46)

RARE EARTH IONS

crystal field splitting effects in metals, (T), 26:47269
polarizabilities of ground-state, electric dipole, (T), 26:46349

RARE EARTH ISOTOPES

neutron strength functions for, J dependence of S_0 , p. 298-9, (T), 26:46873(R) (CEA-N-1522)

RARE EARTH OXIDES

decomposition by mass spectroscopy, 26:45259
rare earth oxides-- UO_2 , crystal structure of solid solutions in, 26:45414(R) (ORNL-TM-3180)

RARE EARTHS

analysis for terbium by luminescent method using pyrogallolsulfonic acid, 26:45231
determination in artificial fluorites, use of paper chromatography in, 26:45232
determination of, evaluation of mass spectroscopy for,

26:45259
electron band structure of, 4f levels in, (T), 26:47260
electronic properties, effect of 4f levels on, 26:47260
polarizabilities of ground-state, electric dipole, (T),
26:46349
separation by anion exchange using acetic acid--methanol--
nitric acid mixture, 26:45280
separation by paper chromatography, evaluation of
developers for, 26:45275

RARE GAS IONS
effects on glasses and thin oxide layers at 5.6 keV,
sputtering, energy loss, and luminescence from,
26:47238

RARE GASES
electron scattering by, scattering lengths for, (T),
26:46401

RATS
effects of irradiated animal feeds on, following
ingestion, 26:45899
iodine-125-labeled transplantation antibody distribution
in syngeneic and allogeneic, 26:47649 (UR-3490-147)
promethium-147 distribution and retention in tissues of,
following intravenous injection as chloride solution,
effects of DTPA treatment on removal, 26:45922
promethium-148 distribution and retention in tissues of,
following intravenous injection as chloride solution,
effects of DTPA on, 26:45922
radiation detection by eyes of anomic, γ , 26:45817
(ORO-2690-12(Pt.B), Paper 12)
radiation effects on activity of adenylylase and
5'nucleotidase of plasma membranes of liver of, effects
of glucagon and sodium fluoride on gamma, 26:45831
radiation effects on activity of deoxyribonuclease in
liver of, effects of cytostatic drugs on, 26:45732
radiation effects on blood vessel regeneration in muscle
transplants in, 26:45857
radiation effects on bone marrow of, following lethal
whole-body exposure, role of transplanted slowly
proliferating bone marrow cells in regeneration of x,
26:45837
radiation effects on capacity of injured muscle of, to
regenerate following traumatic injury, 26:45827
radiation effects on content of thiol groups in heart and
liver mitochondria of, 26:45829
radiation effects on function and structure of liver
mitochondria in, following lethal whole-body exposure,
26:45838
radiation effects on growth of bone tissues of, effects
of dose and parotin on, 26:45870
radiation effects on lysosomes in liver cells of,
following lethal whole-body exposure, x, 26:45839
radiation effects on metabolism of erythropoietin in,
following whole-body exposure, additive effects of
hypoxia on, 26:45854
radiation effects on nucleotide distribution in DNA of
bone marrow of, x, 26:45856
radiation effects on survival of, protective effects of
DEAE Sephadex against x, 26:45844
radioinduced cardiovascular malformations in fetuses of,
following sublethal whole-body exposure in utero,
statistical analysis of relation between age, dose, and
incidence of fast neutron, 26:45867
radioinduced enteritis in, following abdominal exposure,
effectiveness of cholestyramine resin for prevention
and treatment of cholelithic enteropathy resulting from
x, 26:45821
radioinduced inhibition of glucose-6-phosphatase activity
in erythrocytes of, following whole-body exposure,
effects of cysteine and reduced glutathione on extent
of recovery from x, 26:45848
radioinduced injuries in lungs of, role of allergic
inflammations in pathogenesis of x, 26:45863
radioinduced leukemia in, lymphoid cell proliferation
kinetics in thymus during development of x, 26:45822
radioinduced neoplasms in parabiont, following acute
supralethal whole-body exposure, high incidence of,
26:45814(R) (COO-3017-43)
radioinduced osteogenic sarcomas and uterine carcinomas
in parabiont and single, following focal internal
exposure, cobalt-60 γ , 26:45814(R) (COO-3017-43)
radioinduced suppression and recovery of colony-forming
units of bone marrow cells of, low dose x, 26:45834
radioinduced suppression of immune reactions in, to
allogeneic tumor transplants, additive effects of
cortisone on x, 26:45820 (UR-3490-121)
radioinduced taste aversions in, comparison of lithium-
chloride-induced taste aversions with gamma, 26:45817
(ORO-2690-12(Pt.B), Paper 12)
respiratory exchange in, effects of radioprotective
compounds on, 26:45823

REACTIVITY
feedback analysis by rod drop in fast reactors, digital
reduction procedure for on-line, 26:47307

REACTOR CONTAINMENT STRUCTURES
airlocks for, standards for gaskets for, 26:47593 (RDT-
E-14-6-(I-6-72))

REACTOR CONTAINMENT SYSTEMS
transient pressure and temperature in pressure
suppression type, DECAT code for long term, 26:47318
(RT/PROT-72-4)

REACTOR CONTROL ELEMENTS

reactivity worth of high temperature gas cooled reactor,
computer simulation of pulsed neutron experiments for,
26:47319 (GULF-GA-8-10959)

REACTOR CONTROL SYSTEMS
design for cooling systems with once-through steam
generators, 26:47308

REACTOR COOLANTS
analysis of organic, for trace amounts of oxygen by
pyrolysis and electrolytic hygrometry, 26:45221 (EUR-
4822)

REACTOR EXPERIMENTAL FACILITIES
description of CEA EL3, for fission measurements, p. 311-
3, 26:46873(R) (CEA-N-1522)
description of CEA EL3, for production of slow neutrons
and monochromatic gamma rays, p. 309-11, 26:46873(R)
(CEA-N-1522)

REACTOR FUEL ELEMENTS
analysis for uranium-235, delayed neutron interrogation
method for, 26:45269
centering parts for gas cooled, effects of flow
instabilities on vibration of, 26:47314
cladding for, ultrasonic quality control method for,
26:45472(P)
cladding mechanical interaction with fuel in,
26:45416(R) (ORNL-TM-3250)
cladding mechanical interactions with fuel in,
26:45412(R) (ORNL-TM-3095)
cladding mechanical interactions with oxide fuel in,
FMODEL code prediction of, 26:45415(R) (ORNL-TM-3217)
cladding of oxide, economic and performance evaluation of
materials for, 26:45410(R) (ORNL-TM-3018)
cladding reaction with oxide fuels in, 26:45409(R)
(ORNL-TM-2996)
cladding reactions with oxide fuel in, 26:45410(R)
(ORNL-TM-3018)
coolant flow in bundles of rod, method for calculating
cross-flow component of, 26:47596
criticality studies on spent LMFBR, 26:45416(R) (ORNL-
TM-3250)
decladding of stainless steel-clad oxide, containment of
fission product iodine by pyrochemical, 26:45421
dissolution of stainless steel-clad, research progress on
electrolytic, 26:45405(R) (ICP-1006)
fission product release in oxide, 26:45412(R) (ORNL-TM-
3095)
fuel pellet densification in, 26:45411(R) (ORNL-TM-3072)
heat transfer characteristics of gas cooled tubes,
effects of twisted tapes on, (E), 26:47312 (CONF-
720802-1)
heat transfer to fluid flowing parallel to square arrays
of, point-matching method for analysis of, 26:45465
identification of, marking standards for, 26:46097
performance evaluation of clad oxide, 26:45416(R) (ORNL-
TM-3250)
performance modeling of clad oxide, 26:45414(R) (ORNL-
TM-3180)
performance of clad oxide, 26:45417(R) (ORNL-TM-3281)
performance studies on stainless steel-clad oxide,
26:45411(R) (ORNL-TM-3072)
physics aspects of plutonium containing HTGR-type,
effects of plutonium loading on space-energy dependent,
(T), 26:47313
radiation effects on deformation of uranium, relations of
stress and temperature with in-pile, 26:46070 (UJV-
2798-M)
reprocessing of clad oxide, head-end steps in,
26:45416(R) (ORNL-TM-3250)
reprocessing of metallic-clad oxide, electrochemical
method of cutting and leaching in, 26:45422(P)
reprocessing of oxide, voloxidation fission product
removal in head-end, 26:45413(R) (ORNL-TM-3127)
reprocessing of sheared LMFBR, continuous or
semicontinuous leacher for dissolution in, 26:45397
(CONF-720903-9)
reprocessing of stainless steel-clad oxide, shearing
studies on, 26:45407(R) (ORNL-TM-2918)
reprocessing of stainless steel-clad oxide, shearing
studies for, 26:45408(R) (ORNL-TM-2949)
reprocessing of, fission product removal by voloxidation
in head-end, 26:45414(R) (ORNL-TM-3180)
reprocessing of, volatile fission product removal in
head-end, 26:45410(R) (ORNL-TM-3018)
reprocessing of, volatilization of fission products in
head-end, 26:45412(R) (ORNL-TM-3095)
shipment of spent LMFBR, cask design studies for,
26:45411(R) (ORNL-TM-3072)
shipment of spent LMFBR, cask evaluation for,
26:45417(R) (ORNL-TM-3281)
shipment of spent LMFBR, cask evaluation studies for,
26:45412(R) (ORNL-TM-3095)
shipment of spent LMFBR, cask evaluation studies for,
26:45410(R) (ORNL-TM-3018)
shipment of spent LMFBR, cask evaluation studies on,
26:45413(R) (ORNL-TM-3127)
shipment of spent LMFBR, cask evaluations for,
26:45414(R) (ORNL-TM-3180)
shipment of spent LMFBR, criticality and cask evaluation
for, 26:45406(R) (ORNL-TM-2871)
shipment of spent LMFBR, criticality and heat dissipation
studies on, 26:45407(R) (ORNL-TM-2918)
shipment of spent LMFBR, criticality and heat dissipation
studies for, 26:45408(R) (ORNL-TM-2949)

shipment of spent LMFBR, criticality and heat dissipation studies on, 26:45409(R) (ORNL-TM-2996)
 shipment of spent LMFBR, heat dissipation and shielding studies for, 26:45415(R) (ORNL-TM-3217)
 shipment of spent LMFBR, heat dissipation studies for, 26:45416(R) (ORNL-TM-3250)
 swelling and gas release in stainless steel-clad oxide, 26:45408(R) (ORNL-TM-2949)
 thermal conductivity of sheared stainless steel-clad oxide, 26:45412(R) (ORNL-TM-3095)

REACTOR FUELS

analysis of irradiated, for oxygen, development of remote apparatus for, 26:45224 (LA-DC-72-524)
 analysis of reprocessing solutions of, for uranium and plutonium, review on methods for, 26:45190 (TID-7029(2nd Ed.), pp 331-411)
 book: Selected Measurement Methods for Plutonium and Uranium in the Nuclear Fuel Cycle, 26:45183 (TID-7029(2nd Ed.))
 burnup analysis of fast breeder, 26:45405(R) (ICP-1006)
 chemistry of graphite, in HTGR, 26:45288 (TID-26106)
 criticality studies on LMFBR, 26:45411(R) (ORNL-TM-3072)
 cycle costs for power, effects of batch and continuous refueling on, (T), 26:47315 (CISE-R-178)
 dissolution of spent oxide, in nitric acid, 26:45409(R) (ORNL-TM-2996)
 fission product release from oxide, 26:45411(R) (ORNL-TM-3072)
 inspection of irradiated, modification of LASL plutonium cells for nondestructive, 26:46104 (LA-DC-72-556)
 neutron cross section ratios for uranium-235 enriched, determination of effective, (E/T), 26:47595
 neutron source strength of spent LMFBR, 26:45408(R) (ORNL-TM-2949)
 performance evaluation of oxide, 26:45415(R) (ORNL-TM-3217)
 performance evaluation of oxide, 26:45409(R) (ORNL-TM-2996)
 performance of oxide, 26:45410(R) (ORNL-TM-3018)
 performance of oxide, 26:45412(R) (ORNL-TM-3095)
 performance of oxide, 26:45413(R) (ORNL-TM-3127)
 performance studies on oxide, 26:45407(R) (ORNL-TM-2918)
 performance studies on oxide, 26:45408(R) (ORNL-TM-2949)
 photon spectra of spent LMFBR, 26:45408(R) (ORNL-TM-2949)
 preparation of coated-particle, incorporating fertile materials, 26:45959(P)
 processing of molten salt, development of procedures for, 26:47302
 radiation effects on restructuring of ceramic, model for, 26:45417(R) (ORNL-TM-3281)
 radiation properties and shielding of LMFBR, 26:45410(R) (ORNL-TM-3018)
 reprocessing of carbide, oxidation of organic compounds following nitric acid dissolution step in, 26:45420(P)
 reprocessing of graphite, combustion-dissolution studies for head-end, 26:45405(R) (ICP-1006)
 reprocessing of LMFBR, progress report (Jan 1970) on, 26:45406(R) (ORNL-TM-2871)
 reprocessing of LMFBR, research progress (April 1970) on aqueous, 26:45409(R) (ORNL-TM-2996)
 reprocessing of LMFBR, research progress (August 1970) on aqueous, 26:45413(R) (ORNL-TM-3127)
 reprocessing of LMFBR, research progress (Dec 1970) on, 26:45417(R) (ORNL-TM-3281)
 reprocessing of LMFBR, research progress (Feb. 1970) on aqueous, 26:45407(R) (ORNL-TM-2918)
 reprocessing of LMFBR, research progress (July 1970) on, 26:45412(R) (ORNL-TM-3095)
 reprocessing of LMFBR, research progress (June 1970) on, 26:45411(R) (ORNL-TM-3072)
 reprocessing of LMFBR, research progress (March 1970) on aqueous, 26:45408(R) (ORNL-TM-2949)
 reprocessing of LMFBR, research progress (May 1970) on, 26:45410(R) (ORNL-TM-3018)
 reprocessing of LMFBR, research progress (Nov. 1970) on aqueous, 26:45416(R) (ORNL-TM-3250)
 reprocessing of LMFBR, research progress (Oct 1970) on aqueous, 26:45415(R) (ORNL-TM-3217)
 reprocessing of LMFBR, research progress (Sept 1970) on aqueous, 26:45414(R) (ORNL-TM-3180)
 reprocessing of oxide, continuous dissolution in, 26:45406(R) (ORNL-TM-2871)
 reprocessing of oxide, decay heat dissipation in head-end, 26:45406(R) (ORNL-TM-2871)
 reprocessing of spent, separation of platinum metals in, 26:45423(P)
 reprocessing of zirconium, by solvent extraction at ICPP, 26:45405(R) (ICP-1006)
 reprocessing of, Australian plans for management of radioactive wastes from, 26:46098 (AAEC/TM-628)
 reprocessing of, evaluation of fluoride volatility method for, 26:45419
 reprocessing of, research progress at ICPP on, 26:45405(R) (ICP-1006)
 reprocessing of, uses of fluidized beds in, 26:45418

REACTOR MODERATORS
 materials for, properties of heavy water, 26:45698

REACTOR SAFETY
 instrumentation criteria for reactor accident evaluation, 26:47317 (BNWL-1635)

REACTOR SIMULATORS
 development for operator training, summary of, 26:47572

139 SUBJECTS

REACTORS, RESEARCH

use of computer, for pulsed neutron experiments in high temperature gas cooled reactors, 26:47319 (GULF-GA-B-10959)

REACTORS
 development of, progress review on, 26:47301
 fission products in effluent from, design of remote-controlled sequential sampler for sampling, 26:45613(P)
 neutron flux distribution in, FLUB code for analysis of, 26:47594 (BMBW-FBK-72-6)

REACTORS, AIRCRAFT
 coolant blowers for, effects of power take-off design on performance of helium, 26:45425 (N-72-16724)

REACTORS, BOILING
 coolant loss accidents in, DECAT code for long term pressure and temperature transients in containment following, 26:47318 (RT/PROT-72-4)

REACTORS, BREEDER
 power shift during long-term operation of, effects on breeding ratio, core power density, and reactivity coefficients from, (T), 26:47306

REACTORS, FAST
 design of, review of stainless steel swelling and creep effects on, 26:47303
 performance of, review of stainless steel swelling and creep effects on, 26:47303
 physics measurements for plutonium-graphite and uranium-graphite mixtures in, (E), 26:47581 (CEA-N-1483)
 physics measurements for power, effects of fission neutron spectra on, 26:47585 (EURFNR-1008)
 power shift during long-term operation of, effects on breeding ratio, core power density, and reactivity coefficients from, (T), 26:47306

REACTORS, GAS-COOLED
 chemistry problems in high-temperature, 26:45288 (TID-26106)
 control materials for, effects of fast neutrons on dimensional stability of, 26:46077 (GULF-GA-A-12035)
 cores for high temperature, mechanical and thermal behavior of, 26:47304
 fuel element centering parts for, effects of flow instabilities on vibration of, 26:47314
 fuel elements for high temperature, effects of plutonium loading on space-energy dependent physics of, (T), 26:47313
 fuel tubes for, effects of twisted tapes on heat transfer characteristics of, (E), 26:47312 (CONF-720802-1)
 steam generators for high temperature, control system for once-through, 26:47308

REACTORS, MARITIME
 book: Analog Models of Marine Nuclear Steam Generating Plants, in Russian, 26:47578
 operation of, analog models for simulation of, 26:47578

REACTORS, MOLTEN-SALT
 coolant salts for, development of, 26:47302
 design parameters for 350 MW(e) demonstration, 26:47305 (ORNL-TM-3832)
 fuel salt for, development of, 26:47302
 materials development for, review of, 26:47302
 tritium control in, development of procedures for, 26:47302

REACTORS, POWER
 accidents in, criteria for instrumentation for evaluation of, 26:47317 (BNWL-1635)
 containment vessel airlocks for, standards for gaskets for, 26:47593 (RDT-E-14-6-T(6-72))
 cooling system for CIRENE, PECVAR and CRIP codes for thermohydraulic analysis of, 26:47309 (CISE-R-257)
 cooling systems for, effects of pressure fluctuations on vibrations in, 26:47310
 design parameters for 350 MW(e) demonstration molten salt, 26:47305 (ORNL-TM-3832)
 development of dual purpose, desalination and, 26:47579
 effects on public health of operation of, assessment of, 26:45904 (COO-3013-1)
 fluid flow in parallel channels of, studies on stability of, (E), 26:45459 (CISE-R-275)
 fuel cycle costs for CIRENE, effects of batch and continuous refueling on, (T), 26:47315 (CISE-R-178)
 materials analysis for CIRENE program, 26:45181 (CISE-R-172)
 operator training for, development of simulators for, 26:47572
 physics measurements for fast, effects of fission neutron spectra on, 26:47585 (EURFNR-1008)
 pressure vessels for, development of prestressed concrete, 26:47311
 radioactive gas releases from, problems associated with atmospheric contamination from, 26:46102(T) (ORNL-tr-2369)

REACTORS, PROCESS HEAT
 development of dual purpose, for desalination and power, 26:47579

REACTORS, RESEARCH
 development of Apsara, history of, 26:47588
 experimental channels for, performance of expansion bellows for, 26:47591
 in-pile loop in ESSOR, computer models for evaluation of safety of CART fuel test facility in, 26:47580 (CISE-R-259)

- operation of Minerve, history of, 26:47589
- REACTORS, ROCKET**
- fuels for, development of nitride, 26:45960
- fuels for, shock and vibration testing of uranium nitride, 26:45933 (N-72-15609)
- REACTORS, SWIMMING-POOL**
- design and operation of, at Pakistan Institute of Nuclear Science and Technology, 26:45499
- radioisotope production in, at Pakistan Institute of Nuclear Science and Technology, program for, 26:45396
- REACTORS, WATER-COOLED**
- coolant chemistry in, review on, 26:45310 (BARC-595)
- pressure vessels for, development of prestressed concrete, 26:47311
- REACTORS, WATER-MODERATED**
- coolant chemistry in, review on, 26:45310 (BARC-595)
- RECOILS**
- reactions of, variation of energy loss parameter in Estrup--Wolfgang theory with energy in, 26:45363
- RECTIFIERS**
- radiation effects on silicon single-junction and silicon-controlled, 22-MeV proton, 26:45661 (N-72-17158)
- RED SEA**
- book: Problems of Production, Sanitation, and Technical Hydrobiology of the Southern Seas, 26:45919
- REGGE POLE THEORY**
- absorptive cuts for meson--baryon interactions, modified model of, 26:46622
- cuts in forward $p + p + p + p$ interactions, inclusive finite mass sum rules for upper bound for magnitude of ρ and A_2 , 26:46651 (RPP/T-18)
- cuts in meson--meson and meson--baryon interactions, duality properties of, 26:46600
- dual amplitude with Mandelstam analyticity in, threshold behavior of trajectory and, 26:46774 (ITF-72-44-E)
- exchange degeneracy in strong interactions at high energies in, 26:46592
- interactions in, duality and exotic resonances in inclusive, 26:46605
- interactions $p + p + p + p + \pi^+ + \pi^-$ at 13.0 GeV/c in multi-Regge models, 26:46656
- model for cross sections of $\gamma + N \rightarrow K^+ + N$ interactions at 5 to 15 GeV/c, 26:46534
- model for rotation of polarization in $\pi^- + p$ interactions at 6 and 16 GeV, 26:46617 (IFVE-STF-71-15)
- phenomenology of total cross sections in forward $\pi + N$ and $K + N$ interactions at high energies and K_1^0 regeneration amplitude in, modifications in, 26:46633
- poles from multiperipheral unitarity for pole--cut collision mechanism in, properties of complex-conjugate, 26:46631
- Pomeranchuk theorem violation for differential cross sections for $K + N$ and $\pi + N$ interactions and quark model for $\gamma + N \rightarrow N +$ vector mesons interactions, 26:46645
- Pomeranchuk theorem violation in total cross sections of nonforward $K^- + p$ and $K^+ + p$ interactions and amplitudes of $K_2^0 + K_1^0$ regeneration on neutrons, 26:46634
- pomeranchukon vertex for hadron interactions, consequences of vanishing of, 26:46597
- Reggeon propagator, vertex function, form factor, and renormalization constants in Zachariassen model formulation, perturbation theory for intermediate, 26:46747
- trajectories for large coupling constants in Wick--Cutkosky, 26:46725
- trajectory threshold behavior for dual amplitude with Mandelstam analyticity in, 26:46774 (ITF-72-44-E)
- triple-Regge model with Pomeron for inclusive distributions in central region, 26:46661
- RELATIVITY THEORY, GENERAL**
- black holes in, Kerr solution and surface area for upper limits on amounts of energy extractable from, 26:46193
- black holes in, upper limits on amounts of extractable energy for spin-dependent force between two, 26:46193
- Einstein equation solutions with cosmological term for homogeneous anisotropic model of universe in, 26:46759
- Einstein equations for gravitational fields with groups of motions on two-dimensional transitivity hypersurfaces in model with sources of dust and magnetic field, exact solutions of, 26:46755
- Einstein equations in vacuum in, null bundle transformations for solutions to, 26:46763
- Einstein equations in, validity of spatially homogeneous universes with rotation in variations in action principle for, 26:46758
- Einstein field equations for formulation of homogeneous anisotropic model for rotating universe with flat space, 26:46261
- Einstein--Maxwell equations in, null bundle transformations for Kerr solution for gravitational field of homogeneous rotating ring for, 26:46762
- Einstein--Maxwell equations in, null bundle transformations for solutions to, 26:46762
- Einstein--Maxwell equations in, null bundle transformations for solutions to, 26:46763
- Einstein--scalar space-times, characteristic development of trapped surfaces in, 26:46764
- energy--momentum determination in, 26:46756
- equations in Minkowski space in, integration of Newman--Penrose, 26:46769
- field equations for static, spherically symmetric metric for charged point mass in Brans--Dicke Theory, 26:46750
- field equations in formulation of Einstein--Kaufman nonsymmetric unified field theory in, static, spherically symmetric solutions of, 26:46694
- field equations in, cylindrical wave solutions of Bonner and Schroedinger unified, 26:46695
- field theory in, formulation of Einstein--Kaufman nonsymmetric unified, 26:46694
- geometrodynamics in, bubble-time canonical formalism for, 26:46751
- gravitational fields with groups of motions on two-dimensional transitivity hypersurfaces in model with sources of dust and magnetic field, exact solutions of Einstein equations for, 26:46755
- Minkowski space in, integration of Newman--Penrose equations in, 26:46769
- motion of test particle in, covariant canonical equations of, 26:46770
- radiation in, special class of solutions of Lichnerowicz equations and Maxwell equations for singular source-free electromagnetic field for nonspecial recurrent space--time for total pure, 26:46771
- tachyon dust metric in, field equations for, 26:46754
- RELATIVITY THEORY, SPECIAL**
- causality group in finite four-dimensional space--time, 26:46724
- group structure for, 26:46669
- Lorentz transformations in, derivation of linearity and scale factor for, 26:46732
- Lorentz-invariant formulation of Cherenkov radiation emission by tachyons in, 26:46704 (N-72-19766)
- REMOTE-CONTROL EQUIPMENT**
- description for plastic bag sealing to control alpha contamination in hot cells, 26:46108 (UCRL-73757)
- REMOTE-HANDLING EQUIPMENT**
- design for sealing flasks in hot cells, 26:46112(P)
- design for transfer of radioactive sample vials from loading platform to light-tight counting chamber, 26:46110(P)
- design for welding and testing of radioactive waste capsules, 26:46099 (ARH-SA-117)
- RESINS**
- effectiveness of cholestyramine, for prevention and treatment of cholestatic enteropathy in x-irradiated rats, 26:45821
- effects of synthetic polythiol, on absorption and excretion of methylmercury by animals, 26:47643 (UR-3490-80)
- radioinduced deposition--polymerization on metal substrates, method for, 26:46089(P)
- RESISTORS**
- properties of niobium-carbide, 26:47250
- RESPIRATION**
- simulation of human, calibration and performance of mechanical apparatus for, pp.7-11, 26:45673(R) (UCRL-50007-71-3)
- RESPIRATORS**
- efficiency of cartridges for, effects of steady state and pulsating flow on, pp.7, 26:45673(R) (UCRL-50007-71-3)
- RESPIRATORY TRACT**
- radiation dose to epithelial cells of, in animals following inhalation of radon-222 aerosols, relation of content of daughter products in air to α , 26:45861
- radioisotope uptake and retention in, following inhalation as aerosols, use of biological model and autoradiography for measurements of kinetics of, 26:45921
- RHENIUM**
- electric conductivity of pressed and fused, in range 100 to 1700°C, 26:47248
- microscopy of, imaging process in field ion, 26:46038
- thermal conductivity of pressed and fused, in range 100 to 1700°C, 26:47248
- RHENIUM ALLOYS AND SYSTEMS**
- Mo-base--Re, superconducting properties of, effect of dislocation tangles on, (T), 26:46836
- Re--W-base, compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
- Re--W-base, corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
- Re-base--Th, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248
- RHENIUM COUPLES**
- Re/W, radiation effects on properties of, at 2000°C, in-pile, 26:45438 (KFK-1556)
- RHENIUM FLUORIDES**
- electron elastic scattering by, Glauber-theory calculations of amplitudes and differential cross sections for multiple, 26:46381
- RHENIUM ISOTOPES Re-179**
- electron-capture decay scheme, 26:46976 (JINR-P6-6492)
- RHENIUM ISOTOPES Re-182**
- electron-capture decay of ground-state and isomeric, conversion electron spectra from, (E), 26:46978
- energy levels of, Nilsson-model interpretation of ground-state and isomeric, (T), 26:46983
- RHODIUM**
- determination by neutron activation to determine

efficiency of TBP extraction from thiocyanate, 26:45194

separation from spent fuels, 26:45423(P)
RHODIUM ALLOYS AND SYSTEMS

Hf--Pt-base--Rh--Ti--W, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)
Pt-base--Rh, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)
Pt-base--Rh--W, properties of sheets of, for fuel capsule fabrication, 26:45703(R) (ORNL-4796)

RHODIUM ISOTOPES Rh-103

heavy ion scattering by, time-dependent deorientation of recoils in vacuum from, (E), 26:46935
neutron reactions (n,p) with, calculation of excitation functions for fast, 26:47066
ROBERT E. GINNA POWER REACTOR, UNIT 1
control rod logic relay system for, loss of power supply for, 26:47321 (DOCKET-50244-109)
personnel for, excessive radiation exposure of, 26:47359 (DOCKET-50244-106)
pressurizer safety valve discharge piping for, hydraulic and stress analyses of, 26:47576 (DOCKET-50244-108)
radioactive wastes for, measurements for June 23 to July 25, 1972, 26:47406 (DOCKET-50244-107)
steam safety valve supports for, modification to, 26:47576 (DOCKET-50244-108)

ROCKET REACTORS (NERVA)

control system for, design parameters and performance of, (T), 26:47577

ROCKS

analysis for niobium by atomic absorption spectrophotometry, 26:45234
analysis for potassium by neutron activation, 26:45197
atomic number of, in presence of intermediate zone, γ - γ method for determination of effective, 26:45566
density of, in presence of intermediate zone, γ - γ method for determination of, 26:45566
physical properties of, surrounding underground nuclear explosions, effects on seismic wave propagation, (E/T), 26:45490
radioisotope content of, in Rainier rubble chimney, measurement of antimony-125, cerium-144, cobalt-60, cesium-134, cesium-137, europium-152 and -154, iron-55, ruthenium-106, and strontium-90, 26:45483 (NVO-1229-180)
radioisotopes in melt-glass, surrounding Rainier Event cavity, ground water transport of, (E), 26:45482 (NVO-1229-177)
uranium formation in, under oxidizing conditions, geochemistry of, 26:45504

RODS

heat transfer to laminar flow through bundles of, effects of rod displacement on, 26:45462
impact attenuation equation for, longitudinal, 26:47626(T) (SC-T-722520)

ROTORS

rotation of multiple, suspended in magnetic field, method for synchronous, 26:45441 (SRARI-P-120)

RUBBER

radiation effects on mechanical properties of, for use in vacuum, 26:46087

RUBIDIUM

determination in hair by neutron activation and γ spectroscopy, 26:45214
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

RUBIDIUM IONS

implantation in nickel at 1 to 10 keV, penetration and spatial distribution from, 26:47235

RUBIDIUM ISOTOPES Rb-77

half-life of, (E), 26:46985

RUBIDIUM ISOTOPES Rb-85

effects on electromigration of rubidium ions in molten rubidium nitrate, effects of temperature on isotopic, 26:45294

RUBIDIUM ISOTOPES Rb-86

distribution in tissues of maize following uptake from soil, effects of calcium oxide and potassium content of soil on, 26:45925

RUBIDIUM ISOTOPES Rb-87

effects on electromigration of rubidium ions in molten rubidium nitrate, effects of temperature on isotopic, 26:45294

RUBIDIUM URANIUM CHLORIDES

K_2UCl_6 -- Rb_2UCl_6 , spectra of, effects of chemical state of uranium on, 26:45377

RUBIES

radiation effects on crystals of, thermal-neutron, γ , and x, 26:47244

RUPTURE

analysis of inelastic, using J integral, 26:45990 (CONF-720806-2)

RUTHENIUM

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination of trace amounts of, in ores, concentrates, and sands by fire assay emission spectrography, 26:45229
separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281
separation from spent fuels, 26:45423(P)

RUTHENIUM ISOTOPES Ru-94

energy levels of, unpaired-nucleon collective, (T), 26:46965

RUTHENIUM ISOTOPES Ru-96

energy levels of, unpaired-nucleon collective, (T), 26:46965

RUTHENIUM ISOTOPES Ru-100

transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889

RUTHENIUM ISOTOPES Ru-102

energy levels of, unpaired-nucleon collective, (T), 26:46965

RUTHENIUM ISOTOPES Ru-103

content in atmosphere and vegetation in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

RUTHENIUM ISOTOPES Ru-106

content in atmosphere in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)

effects on gastrointestinal tract following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

S

S-MATRIX

cross section relations with, formulation of algebra with power asymptotic number elements for operations with nonnormalizable wave functions in quantum mechanics for, 26:47284 (IC-72/45)
crossing symmetry and unitarity in Bethe--Salpeter equation for T matrix, 26:46727
elements of Euclidean quantum electrodynamics, equations for, 26:46739
formulation for spin-0 bosons and spin-1/2 fermions in external electric fields, (T), 26:46741
pole trajectories in N/D method for potential scattering, 26:46789
unitarity in third order of perturbation theory, 26:46738
unitarity in third order of perturbation theory in G in theories with nonpolynomial Lagrangians, role of anomalous singularities and scattering amplitude in fourth order in G in, 26:46775 (JINR-P2-6393)

SAHARA DESERT

tritium content in aquifer in central and northwest Algeria area of, 26:45558

SALEM POWER REACTOR, UNIT 1

environmental report on, additional data for, 26:47445 (DOCKET-50272-33)

safety of, requested additional data on, 26:47443 (DOCKET-50272-31)

safety of, requested additional data on, 26:47444 (DOCKET-50272-32)

safety of, requested additional data on, 26:47442 (DOCKET-50272-30)

safety of, revised and additional data on, 26:47441 (DOCKET-50272-29)

SALEM POWER REACTOR, UNIT 2

environmental report on, additional data for, 26:47445 (DOCKET-50272-33)

safety of, requested additional data on, 26:47442 (DOCKET-50272-30)

safety of, requested additional data on, 26:47444 (DOCKET-50272-32)

safety of, requested additional data on, 26:47443 (DOCKET-50272-31)

safety of, revised and additional data on, 26:47441 (DOCKET-50272-29)

SALT DEPOSITS

radioactive waste disposal in, methods and siting parameters for, 26:46100 (CONF-720903-4)

SAMARIUM

compounds, paramagnetic susceptibility in metallic, effects of conduction-electron polarization on, (T), 26:47259

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
separation from europium, gadolinium, neodymium, and praseodymium by anion exchange using acetic acid--methanol--nitric acid mixtures, 26:45280

SAMARIUM ALLOYS AND SYSTEMS

Co-base--Sm, sintering of Co₂Sm permanent magnet alloys, mechanism of, 26:45976

SAMARIUM ISOTOPES

x-ray spectra from, isotope shifts in, (E), 26:46973

SAMARIUM ISOTOPES Sm-141

beta and electron-capture decay of ground-state and isomeric, γ singles and coincidence spectra from, (E), 26:46972

SAMARIUM ISOTOPES Sm-148

energy levels from neutron resonances in samarium-147, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)

proton inelastic scattering at 24.5 MeV, coupled-channels analysis of differential cross sections and asymmetry from polarized, p. 129-34, (E), 26:46873(R) (CEA-N-

- 1522)
SAMARIUM ISOTOPES Sm-150
 energy levels from neutron resonances in samarium-149, spin assignments for, p. 295-8, (E), 26:46873(R) (CEA-N-1522)
SAMARIUM ISOTOPES Sm-151
 beta decay of, energy-dependent K-shell internal ionization probability in, (T), 26:46962
SAMARIUM ISOTOPES Sm-152
 electron inelastic scattering at 76 MeV, electron spectra from, (E), 26:46993
 energy-level transitions in deformed, SU(3) parametrization of E2, (T), 26:46975 (JINR-P4-6462)
 nuclear deformation parameters from electron scattering, (E), 26:46993
 proton inelastic scattering at 24.5 MeV, coupled-channels analysis of differential cross sections and asymmetry from polarized, p. 129-34, (E), 26:46873(R) (CEA-N-1522)
SAMARIUM ISOTOPES Sm-153
 energy-level transitions from neutron capture by samarium-152, internal conversion coefficient and mixing ratio for, (E), 26:46994
 nuclear spin gyromagnetic ratio for, (E), 26:46994
SAMARIUM OXIDES
 density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
 Al₂O₃-Sm₂O₃, density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
SAMARIUM PEROXIDES
 preparation and properties of, 26:46085
SAMPLING
 method for attribute, based on hypergeometric distribution, 26:46094 (WASH-1210)
 method for continuous, to obtain physical inventories of special nuclear materials, 26:46092 (BNL-16930)
SAN ONOFRE POWER REACTOR, UNIT 2
 owners of, financial statements of, 26:47522 (DOCKET-50361-25)
 safety of, revised and requested additional data on, 26:47523 (DOCKET-50361-26)
SAN ONOFRE POWER REACTOR, UNIT 3
 owners of, financial statements of, 26:47522 (DOCKET-50361-25)
 safety of, revised and requested additional data on, 26:47523 (DOCKET-50361-26)
SANDS
 analysis for trace amounts of iridium, ruthenium, and osmium by fire assay emission spectrography, 26:45229
SANDSTONE
 age of cuprous Udokan series (Olekmo-Vitim Mountains), uranium-lead method for estimation of, 26:45569
 nuclear explosion effects on, surrounding Project Wagon Wheel, prediction of underground, 26:45494
SATELLITES
 data from ultraviolet sky-scan telescope on TD1, 26:46186
 description of nuclear-fueled laser, for energy transfer to lunar bases and other space vehicles, 26:45428
SCANDIUM
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 determination by fluorometry using Superchrome Garnet Y, 26:45236
 determination in hair by neutron activation and γ spectroscopy, 26:45214
 determination in scandium-46 solutions, spectrographic, 26:45223 (JEN-232-DQ/I-81)
 determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
 electron-phonon mass enhancement factor from specific-heat data, 26:47262
 separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285
SCANDIUM HYDRIDES
 elastic properties of ScH_{1.99}, from acoustic velocity measurements, 26:45950
SCANDIUM ISOTOPES Sc-41
 energy levels from proton resonances in calcium-40, spin-parity assignments and widths for, (E), 26:46924
SCANDIUM ISOTOPES Sc-44
 energy levels from scandium-45 (p,γ) reactions, spectroscopic factors for, p. 107-9, (E), 26:46873(R) (CEA-N-1522)
SCANDIUM ISOTOPES Sc-45
 energy levels from calcium-44 (p,γ) reactions, spin-parity assignments for, (E), 26:46920
 energy-level transitions from calcium-44 (p,γ) reactions, branching ratios for, (E), 26:46920
 proton reactions (p,γ) at 24.5 MeV, nonlocal finite-range DWBA analysis of differential cross sections and spectrum from, p. 107-9, (E), 26:46873(R) (CEA-N-1522)
SCANDIUM ISOTOPES Sc-46
 analysis of solutions of, for calcium-45 and target element, spectrographic, 26:45223 (JEN-232-DQ/I-81)
SCATTERING
 amplitudes for identical mass, solution of Faddeev equations for one-dimensional system for, 26:46782
 amplitudes in arbitrary order of perturbation theory, power of increase of, 26:46738
 amplitudes in fourth order in G in S-matrix unitarity in third order of perturbation theory in G in theories with nonpolynomial Lagrangians, 26:46775 (JINR-P2-6393)
 amplitudes of large-angle high-energy, solution of quasipotential equation in quasipotential approximation for, 26:46776 (JINR-P2-6442)
 amplitudes of spinning-particle, relativistic generalization of Faxen-Holtzmark method for, 26:46793
 amplitudes of, relativistic eikonal approximation for inelastic effects in, 26:46791
 amplitudes with Mandelstam analyticity for, threshold behavior of trajectory and dual, 26:46774 (ITF-72-44-E)
 cross sections for, variational methods for construction of Jost functions for interpolation of, 26:46402
 eikonal formalism for production processes in, 26:46778
 formalism in Fock space, 26:46783
 inverse problem of, free Riemann-Green function for, 26:46785
 operator in (φ₁)₂ theory, unitarity, causality, and generalization to test function space for Bogoliubov, 26:46740
 phase shifts for r/sup -2n/ potential, Born-approximation calculation of, and application to dipole-quadrupole r² potential, 26:46380
 potential equation in quasipotential approximation for amplitudes of large-angle high-energy, solution of quasi-, 26:46776 (JINR-P2-6442)
 potential, Glauber approximation from multiple-scattering expansion in, 26:46792
 potential, numerical aspects for solving partial wave equations in momentum space for strongly peaked, 26:46787
 potential, probabilistic interpretation of cross sections for quantum, 26:46777
 potential, relativistic three-particle equations for arbitrary, (T), 26:46779
 potential, resonance calculations in, using Jost function theory, 26:46400
 potential, Riemann method for inverse problem of, 26:46788
 potential, S-matrix pole trajectories in N/D method for, 26:46789
 potential, solution of inverse, 26:46784
 potential, solution of standing-wave integral equation for, 26:46786
 potential, time-dependent and time-independent theory for asymptotic Coulomb, 26:46780
 potential, von Neumann algebras for nonrelativistic time-dependent, 26:46781
 Reggeon amplitude in, formulation of Zachariasen model for intermediate, 26:46747
 S-matrix theory of, structure of compound states in, (T), 26:47068
 theory for three-body, exact stationary solutions for classical relativistic, 26:46790
 theory for three-body, in eikonal approximation, breakdown of simple cancellation in operator multiple scattering series in, 26:46773 (CTH-ITP-71-17)
SCINTILLATION CAMERAS
 collimator for improved peripheral response, 26:45596(P)
 collimator for, convergent multihole focusing, 26:45602
SEALS AND GLANDS
 standards for reactor containment vessel airlock gaskets, 26:47593 (RDT-E-14-6-T(6-72))
SEAS
 beryllium-7 deposition and distribution in, relations of fission products to, 26:45530 (BNWL-SA-4205(Rev.))
 fission product deposition in, relations with atmospheric beryllium-7 content, 26:45530 (BNWL-SA-4205(Rev.))
SEAWATER
 cobalt-60 content in samples of Pearl Harbor, during 1966 and 1968, 26:45534
 contamination by radioactive wastes, International Laboratory of Marine Radioactivity research on, 26:46101 (IAEA-136, pp 10-17)
 corrosive effects on refractory metal fuel capsules, 26:45701(R) (DP-1300)
 desalting of, methods for large-scale, 26:47579
 radioisotope content in Puget Sound basin, 26:45544 (RLO-2225-T-14-4)
 radioisotope content of, correlation between content of giant brown algae and polonium-210 and plutonium-239, 26:45924
SEED
 mutations induced in barley, by ethylenimine, protective effects of cysteine against, 26:45772
 radiation effects on germination and growth of desert plant, dose and dose rate dependence of γ, 26:45758 (TID-25954, pp 49-53)
SEEDLINGS
 radiation effects on survival of desert plant, following seed exposure, dose dependence of γ, 26:45758 (TID-25954, pp 49-53)
SEISMIC WAVES
 characteristics of surface, generated by Nevada Test Site underground nuclear explosions and recorded at Las Vegas, 26:45481 (NVO-1163-TM-33)
 discrimination between earthquake- and underground nuclear explosion-generated, at East German Moxa Station, description of simple method for, 26:45496

- effects of Cannikin and Milrow Events-generated, on seismicity of Amchitka Island area during 1969 to 1972, (E), 26:45479 (NVO-746-122)
- measurements using continental and large aperture arrays, 26:45498
- propagation from low-magnitude tensile fault in noise field, (T), 26:45491
- propagation from underground nuclear explosions, effects of surrounding media on, (E/T), 26:45490
- spectra of underground nuclear explosion-generated, at Pahute Mesa and Yucca Flat, Nev., analysis and comparison of response, 26:45480 (NVO-1163-TM-32)
- travel-times of, method for correcting errors in measuring single-station, 26:45495
- SEISMOLOGY**
- seismic arrays for research in, performance of continental and large aperture, 26:45498
- SELENIUM**
- determination in hair by neutron activation and γ spectroscopy, 26:45214
- determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
- SELENIUM ISOTOPES Se-72**
- energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo+SU₂ coupling schemes, 26:46931
- nuclear properties of, compilation of, (E), 26:46939
- SELENIUM ISOTOPES Se-74**
- energy levels from nickel (HI,x) reactions, comparison with pseudo-LS and pseudo+SU₂ coupling schemes, 26:46931
- SELENIUM ISOTOPES Se-76**
- energy levels from decay of bromine-76 decay, (E), 26:46938
- SELENIUM ISOTOPES Se-80**
- neutron capture by, γ spectra from thermal, (E), 26:46937
- SELENIUM ISOTOPES Se-81**
- energy levels from selenium-80 neutron capture, (E), 26:46937
- SEMICONDUCTOR DEVICES**
- injection characteristics of symmetrical ion-implanted PIN diodes, double, 26:45678
- radiation effects on passivation, 26:45662 (SC-DC-721232)
- radioinduced carrier trapping effects in p-n, neutron, (T), 26:45664 (SC-DC-721237)
- radioinduced current pulses in circuits of, degradation by gamma, 26:45666 (SC-RR-720515)
- SEMICONDUCTOR JUNCTIONS**
- book: Radiation Physics of Crystals and p-n Junctions (in Russian), 26:47214
- defect centers in p-n, calculation of spatial distribution of deep, (E/T), 26:47243
- profile determination by wafer angle lapping, 26:47608 (BDX-613-604(Rev.))
- radiation effects on carrier lifetime and isochronal annealing of silicon p-n, γ , 26:45671
- radiation effects on MAS, x, 26:45670
- radioinduced carrier trapping effects in p-n, neutron, (T), 26:45664 (SC-DC-721237)
- SEMICONDUCTORS**
- book: Radiation Physics of Crystals and p-n Junctions (in Russian), 26:47214
- SEMINAL VESICLES**
- radioinduced hypertrophy of, in fish, internal phosphorus-32 β , 26:45898
- SEPARATION PROCESSES**
- efficiency of combined ion exchange and solvent extraction, on paper chromatographs, 26:45282
- SERPUKHOV SYNCHROTRON**
- shielding for, intranuclear cascades in concrete, (E/T), 26:46126(T) (ORNL-tr-2617)
- SEWAGE**
- radioactivity monitoring of California, during 1970, gross beta, 26:45537
- radioactivity monitoring of, at Brookhaven National Laboratory during Jan. to July 1968, 26:45515
- radiosterilization of, description of spiral-flow gamma facility for, 26:45719(P)
- radiosterilization of, design of cobalt-60 irradiation facility for, 26:45456(P)
- SEX**
- effects on frequency of chromosome aberrations in lymphocytes of irradiated Marshall Island population, 26:45792
- SHEARON HARRIS POWER REACTOR, UNIT 1**
- design parameters of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- environmental report for, requested additional data for, 26:47564 (DOCKET-50400-20)
- operations aspects of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- safety of, requested additional data on, 26:47565 (DOCKET-50400-21)
- SHEARON HARRIS POWER REACTOR, UNIT 2**
- design parameters of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- environmental report for, requested additional data for, 26:47564 (DOCKET-50400-20)
- operations aspects of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- safety of, requested additional data on, 26:47565 (DOCKET-50400-21)
- SHEARON HARRIS POWER REACTOR, UNIT 3**
- design parameters of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- environmental report for, requested additional data for, 26:47564 (DOCKET-50400-20)
- operations aspects of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- safety of, requested additional data on, 26:47565 (DOCKET-50400-21)
- SHEARON HARRIS POWER REACTOR, UNIT 4**
- design parameters of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- environmental report for, requested additional data for, 26:47564 (DOCKET-50400-20)
- operations aspects of, revised and additional data on, 26:47563 (DOCKET-50400-19)
- safety of, requested additional data on, 26:47565 (DOCKET-50400-21)
- SHEETS**
- particle track counting in, using capacitive discharge device, 26:46083(P)
- SHELLS (GEOMETRY)**
- consistent theory for thin spherical, in equilibrium, 26:45453
- SHELTERS**
- heating of small livestock, using power plant waste heat, 26:45576
- SHIELDED CONTAINERS**
- design and testing of drum-type, for shipping fissionable materials and radioactive wastes, 26:46114 (MLM-1911)
- design for enclosing visible syringe-contained radioactive liquid, 26:46113(P)
- design for shipping fissile materials, engineering drawing for, 26:46116
- SHIELDING**
- data test for, iron neutron total cross sections at 0.8 to 11 MeV, (E/T), 26:47220 (ORNL-TM-3867)
- data test for, nitrogen neutron total cross sections at 0.8 to 10 MeV, (E/T), 26:47222 (ORNL-TM-3869)
- data test for, oxygen neutron total cross sections at 1.9 to 8.6 MeV, (E/T), 26:47221 (ORNL-TM-3868)
- data test for, sodium neutron total cross sections at 0.8 to 11 MeV, (E/T), 26:47223 (ORNL-TM-3870)
- data test for, stainless-steel neutron total cross sections at 1.2 to 11 MeV, (E/T), 26:47224 (ORNL-TM-3871)
- design and performance for reducing background radiation in whole-body counters, 26:45590(T) (ORNL-tr-2532)
- radiation dose rates outside, Moyer method for estimating, 26:46121
- SHIPPING**
- container for fissile material, design of, 26:46116
- regulations for radioactive materials, description of German, 26:46117
- SHOCK TUBES**
- plasma ionization in electromagnetic, precursor, (E), 26:47131
- precursor ionization in electrically driven, (E), 26:47197
- simulation of, research on, 26:47145(R) (COO-3086-15)
- SHOCK WAVES**
- fine structure of interplanetary, HEOS-1 data of March 8, 1970, (E), 26:46334
- production by exploding current sheet in plasma, (E), 26:47186
- propagation in low-beta plasma, effects of reflected ions on, 26:47199
- propagation in plasma, effect of thermal conductivity on structure and critical parameters of, 26:47200
- structure of large-amplitude stationary hydromagnetic, in collisionless plasma, (E), 26:47196
- structure propagating perpendicular to magnetic fields, collisionless, (E/T), 26:47198
- SILICATES**
- abundance in circumstellar shells, (E), 26:46174
- accretion of grains in early solar system, (E), 26:46229
- analysis of, containing phosphorus, titanium, and zirconium, for silica by precipitation of oxine 12-molybdisilicate, 26:45255
- SILICIDES**
- structure of binary, data compilation on, 26:45965
- SILICON**
- abundance in solar-flare particles, (E), 26:46232
- determination in boreholes, neutron activation logging method for, 26:45567
- determination in coal and coal ash by neutron activation, 26:45195
- dissolution in iron, 26:46007
- effects on mechanical properties of aluminum-iron-uranium-base and molybdenum-uranium-base alloys at 20 to 640°C, 26:45995 (UJV-2802-M)
- electron interactions with crystals of, at 1 GeV, width of minimum of total yield of retarding photons from, (E), 26:46550
- impurity effects on gamma-irradiated crystals of, 26:45671
- ion implantation in, mass dependence of critical amorphizing dose in, (T), 26:47236
- positron interactions with crystals of, at 1 GeV, width

- of minimum of total yield of retarding photons from, (E), 26:46550
radioinduced shallow donors in n-type, electrical properties of proton-, 26:47237
- SILICON FLUORIDES**
determination in nitrogen by automatic absorption spectrometry, 26:45243(P)
- SILICON IONS**
spectral lines of, under solar-flare conditions, time-dependent ionization equilibrium, (T), 26:46226
- SILICON ISOTOPES Si-28**
lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883
muon (μ^-) capture by, particle-hole calculation of capture rate and neutron spectra for, (T), 26:46880
nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892
oxygen-16 reactions (O-16,C-12) at 42 MeV, differential cross sections for, (E), 26:46913
photon reactions (γ,n) and (γ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
proton elastic scattering at 1.5 to 3.0 MeV, polarization from polarized, (E), 26:46924
- SILICON JUNCTIONS**
radiation effects on, 22-MeV proton, 26:45661 (N-72-17158)
radioinduced deep defect centers in p-n, determination of 300-keV proton, (E/T), 26:47243
- SILICON OXIDES**
crystallization from lithium molybdate--molybdenum oxide flux, effects of solute composition on, 26:45943
determination in silicates containing phosphorus, titanium, and zirconium by precipitation of oxine 12-molybdosilicate, 26:45255
BaO--SiO₂, radioinduced optical absorption and paramagnetic centers in glassy, spectral studies on γ , 26:47226
K₂O--SiO₂--SrO, diffusion coefficients for, at 630 to 810°C, measurement of inter-, 26:45946
SiO₂--SrO, radioinduced optical absorption and paramagnetic centers in glassy, spectral studies on γ , 26:47226
- SILICON SYSTEMS**
Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009
Al--Fe--Nb--Si, dissolution in, 26:46007
Al--Fe--Si--U-base, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Al--Mo--Si--U, mechanical properties of, at 20 to 640°C, 26:45995 (UJV-2802-M)
Ge--Si-base, etching by hydrofluoric acid--nitric acid solutions, 26:45935 (SC-RR-710884)
Ge--Si-base, oxidation by air and water, effects of temperature on, 26:45935 (SC-RR-710884)
- SILICONE RUBBERS**
fabrication of thin slabs of Y-3333, 26:47603(R) (BDX-613-440(Rev.))
- SILOXANES**
radioinduced deposition--polymerization on metal substrates, method for, 26:46089(P)
- SILTS**
cobalt-60 content in samples of Pearl Harbor, during 1966 and 1968, 26:45534
fallout content and distribution in Atlantic Ocean and Mediterranean Sea, 26:45531 (COO-3563-3)
plutonium isotope content and distribution in Atlantic Ocean and Mediterranean Sea, particle-settling model for, 26:45531 (COO-3563-3)
plutonium isotope content in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)
radioisotope content of Lake Michigan, near Big Rock Point power reactor during 1970, measurements of cesium-134, cesium-137, cobalt-60, and zinc-65, 26:45536
trace metal content and distribution in Arctic Ocean, during June 1971 to June 1972, 26:47623(R) (RLO-2229-T-1-18)
- SILVER**
determination in hair by neutron activation and γ spectroscopy, 26:45214
determination in ores by lead crucible fusion and neutron activation, 26:45200
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
- SILVER ALLOYS AND SYSTEMS**
Ag-base--In, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)
- SILVER IODIDES**
sorpitive properties for iodine in aqueous solutions, 26:45279
- SILVER IONS**
implantation in iridium at keV energies, field-ion microscopic study of damage profiles produced in, 26:47240
- SILVER ISOTOPES**
photon reactions (γ,ν^0) at 3.5 to 8.3 GeV, A dependence of, 26:46447
- SILVER ISOTOPES Ag-107**
neon-20 reactions at 90 to 210 MeV, calculation of fusion excitation function using equilibrium model with fission competition, (T), 26:47072
- SILVER ISOTOPES Ag-110**
distribution of metastable, in tissues of dogs following inhalation, 26:45915 (UR-3490-115)
uptake and retention of metastable, in lungs of dogs following inhalation, 26:45915 (UR-3490-115)
- SKIN**
radioinduced acneform changes in human, following therapy of internal tumors, γ , 26:45793
radioinduced injuries to, of dogs following fractionated or single partial body exposure, effects of dose and fractionation interval on x, 26:45860
- SODIUM**
determination in hair by neutron activation and γ spectroscopy, 26:45214
effects on recrystallization of iron, 26:45982
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
neutron total cross sections for, from 0.8 to 11 MeV, (E/T), 26:47223 (ORNL-TM-3870)
reactions of, review on, 26:45417(R) (ORNL-TM-3281)
reactions of, survey on, 26:45416(R) (ORNL-TM-3250)
reactions with nitric acid, 26:45411(R) (ORNL-TM-3072)
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
uptake of radioactive, by larval and hatchling amphibians and reptiles, 26:45815(R) (COO-3486-1)
- SODIUM CHLORIDES**
chlorine-38 NMR in, (E), 26:46911
radioinduced F centers in crystals of, intensity dependence of proton-, 26:47246
- SODIUM HYDROXIDES**
effects of etch solutions of, on thickness of aluminum hydroxides, 26:47605 (BDX-613-515(Rev.))
- SODIUM IONS**
implantation in nickel at 1 to 10 keV, penetration and spatial distribution from, 26:47235
- SODIUM ISOTOPES Na-19**
energy levels of, data compilation on, 26:46882
- SODIUM ISOTOPES Na-20**
energy levels of, data compilation on, 26:46882
- SODIUM ISOTOPES Na-22**
content of lichen-reindeer-man food chain in Sweden during 1961 to 1969, fallout, 26:45909
content of man residing in Sweden during 1961 to 1969, whole-body counting of fallout, 26:45909
- SODIUM ISOTOPES Na-23**
energy levels from charged particle reactions, spin-parity assignment and transition strengths for, (E), 26:46915
energy levels of, spin determinations from γ -- γ angular correlations, 26:46901
proton elastic and inelastic scattering at 1.4 to 3 MeV, resonance structure in excitation functions for, (E), 26:46903(R) (COO-1120-149)
proton reactions (p,α) at 1.4 to 3 MeV, resonance structure in excitation functions for, (E), 26:46903(R) (COO-1120-149)
proton reactions (p,γ) at 1.8 to 2.8 MeV, γ yields for, (E), 26:46903(R) (COO-1120-149)
- SODIUM MOLYBDATES**
Na₂MoO₄--Pu(MoO₄)₂, scheelite structure of phases in, 26:45954
NaMoO₄--Pu(MoO₄)₂, phase diagram of, 26:45954
- SODIUM NITRATES**
decomposition in fluidized-bed processing of radioactive wastes, 26:45383 (ICP-1008)
decomposition of, 26:45405(R) (ICP-1006)
- SOILS**
analysis for calcium, cesium-137, potassium, and strontium-90, 26:45263 (RT/PROT-72-11)
analysis for cesium by neutron activation, 26:45193 (TID-25954, pp 376-8)
analysis for potassium by potassium-40 γ spectroscopy, 26:45193 (TID-25954, pp 376-8)
analysis for trace heavy metals by neutron and photon activation methods, 26:45198
characteristics of Hot Creek Valley, Nev., methods for evaluating effects of underground nuclear explosion-generated ground motion on, 26:45478(R) (NVO-409-1)
plutonium isotope content in Utah, 1971, effects of nuclear weapons testing on, 26:45526 (HASL-257, pp 1.95-118)
porosity of Rio Blanco Site, well logging data on, 26:45476 (NVO-38-33)
radioactivity in, coring sampler for depth profile measurements of, pp.18-19, 26:45673(R) (UCRL-50007-71-3)
radioactivity monitoring of, in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
radioactivity of, in uraniferous area in New Zealand, 26:45529
radioisotope transport through, at Nevada Test Site, fallout, 26:45914 (TID-25954, pp 58-64)
radon release from, to atmosphere, (E/T), 26:45524
sorpitive properties for iodine-131, effects of soil

- characteristics on, (E), 26:45528
strontium-90 deposition and content in Salt Lake City, Utah, 1957 to 1971, effects of stratospheric fallout on, 26:45525 (HASL-257, pp I.71-94)
strontium-90 uptake by wheat in podzolic, effects of calcium on, 26:45550
- SOL-GEL PROCESS**
development for LMFBR fuel recycle, 26:45406(R) (ORNL-TM-2871)
- SOLAR CORONA**
dynamics of, (E/T), 26:46233
electron densities in, from occultations of NP 0532, (E), 26:46227
heating of transition region with chromosphere, (E/T), 26:46228
properties of, (E/T), 26:46233
structure of transition region with chromosphere, (E/T), 26:46228
temperature at various heights, (E), 26:46221
- SOLAR FLARES**
effects on cosmic radiation dose rate to man during high-altitude supersonic flight, 26:45806
effects on generation of interplanetary streams in solar wind, relation with geomagnetic storms and Forbush decreases, (E/T), 26:46332
electron densities and temperatures during, (E/T), 26:46226
emission of various areas and intensities, relation with common-type 4995- and 8800-MHz solar radio bursts, Nov. 1966 to Jan. 1970, (E), 26:46333
gamma energy spectra and flux from, review of, 26:46231
ionization equilibrium and line radiation in, time-dependent, (E/T), 26:46226
longitudes during 20th cycle of, active, (E), 26:46222
nuclei in, composition and energy spectra of heavy, (E), 26:46232
proton events of Nov. 18, 1968 and Feb. 25, 1969, entry mechanism at 100 to 300 MeV into distant magnetotail during, (E/T), 26:46304
radio emission associated with, of Nov. 15 and 16, 1970, (E), 26:46223
x-ray emission associated with, of Nov. 15 and 16, 1970, (E), 26:46223
- SOLAR PARTICLES**
acceleration in solar atmosphere, effect of single- and multiple-ionization losses on, (T), 26:46230
emission of bursts of neutrons, detection by measuring trapped-neutron-decay protons in radiation belts, 26:46320
- SOLAR RADIATION**
charge spectrum of cosmic rays from nuclear emulsions and Explorer-35 solid-state detector, (E), 26:46330
detection above 75 GeV, (E), 26:46219
- SOLAR RADIO BURSTS**
emission during chromospheric flares of Nov. 15 and 16, 1970, (E), 26:46223
exciter of Type III, characteristics of, (E), 26:46221
infrared radiation from, far-, (E), 26:47143
occurrences of 4995- and 8800-MHz common types from Nov. 1966 to Jan. 1970, relation to flares of various areas and intensities, (E), 26:46333
spectral observations of Type-III and spike, (E), 26:46217
time profiles of Type III, (E), 26:46221
- SOLAR SYSTEM**
accretionary processes in early, (E), 26:46229
- SOLAR WIND**
charged-particle transport in turbulent plasma of, (T), 26:46294
cosmic-particle modulation by, integration of steady Fokker-Planck equation for, (T), 26:46152
electron heat conduction in, inhibition mechanisms of, (T), 26:46326
electron-plasma frequency fluctuations in oblique nonlinear MHD wave, detection of, (E), 26:46329
helium abundance variations in, dependence on velocity and proton flux, (E/T), 26:46328
interaction with magnetosphere, magnetosphere compression variations and reflection and refraction of MHD waves at magnetosphere boundary during, (T), 26:46321
interactions with planetary ionosphere, electrodynamic model for, (T), 26:46324 (N-72-18796)
model for, two-fluid, 26:46331
relation of solar-flare-associated interplanetary streams in, with geomagnetic storms and Forbush decreases, (E/T), 26:46332
- SOLAR X-RAY BURSTS**
emission during chromospheric flares of Nov. 15 and 16, 1970, (E), 26:46223
- SOLID STATE PHYSICS**
research progress on, India, 1970-71, 26:45311(R) (BARC-599)
- SOLIDS**
book: Secondary Emission and Structural Properties of Solid Bodies (in Russian), 26:47245
ion mass spectrometric studies of, 26:47265
radiation effects on secondary emission and properties of, Russian book on, 26:47245
radioinduced nucleation of voids in crystalline, mechanisms of high-energy ion or neutron, 26:46064 (CONF-710601-, pp 825-40)
radioinduced void formation and growth in, theory of, 26:46062 (CONF-710601-, pp 769-97)
- structure of, use of neutron scattering in studies of, 26:47251
thermal properties of, conference on, 26:45463
- SOLUTIONS**
analysis of, emission spectrographic methods for, 26:45181 (CISE-R-172)
- SOLVENT EXTRACTION PROCESSES**
evaluation for LMFBR fuel reprocessing, 26:45414(R) (ORNL-TM-3180)
evaluation for reprocessing LMFBR fuels, 26:45415(R) (ORNL-TM-3217)
tanks for uranium, plastic coating for corrosion protection of, 26:46080
- SOUTH AMERICA**
radioactivity monitoring of atmospheric precipitation and ground-level atmosphere in, during Feb. 1972, gross beta, 26:45523
radioisotope content of milk in, during Mar. 1971 to Mar. 1972, cesium-137 and strontium-90, 26:45552
- SOUTHWEST EXPERIMENTAL FAST OXIDE REACTOR**
decommissioning plan for, modifications for, 26:47582 (DOCKET-50231-120)
dismantling of, request for order for, 26:47583 (DOCKET-50231-121)
personnel for, site security, 26:47584 (DOCKET-50231-122)
- SOVIET UNION**
age of cuprous Udokan series (Olekmo-Vitim Mountains) in, uranium-lead method for estimation of, 26:45569
radioactivity in coal deposits in Pechora River basin in, distribution and genesis of, 26:45505
- SPACE**
conference on thermophysical aspects of, Washington, D. C., Dec. 1, 1971, 26:45463
discontinuities in interplanetary, Pioneer-9 lf-wave observations on Feb. 2, 1969 near, (E), 26:46302
- SPACE FLIGHT**
book: Radiation Protection Measurement, 26:45808
radiation hazards to astronauts during, cosmic, 26:45812
- SPACE VEHICLES**
power conversion systems for, description of laser-energized, 26:45428
- SPECTROMETERS**
design of large-angle, for high-energy particles, p. 165-6, 26:46873(R) (CEA-N-1522)
filters for amplitude, optimization of electronic, 26:45637 (JINR-P13-6428)
operation and performance of electrostatic charged-particle, using electron multiplier detectors, 26:45645
pole pieces for magnetic, design and fabrication of, with antisymmetric 1/r fields with periodic azimuthal symmetry, 26:45638
- SPECTROSCOPY**
book: Analytical Emission Spectroscopy, Part I, 26:45252
book: Analytical Emission Spectroscopy, Part II, 26:45253
book: Physical Methods of Chemistry. Vol. I, Part IIB, 26:45250
chemical shifts in photoelectron, relation to inner-shell ionization energies of inorganic compounds, 26:46340 (ORNL-TM-3820)
development of flame emission, for trace amounts of elements in atmospheric particulates, 26:45244
developments in emission, review on, 26:45247
methods for semiquantitative, element volatilization mechanism in, 26:45222 (JEN-229-DQ/1-79)
- SPERM**
radioinduced detrimental and lethal mutations in wasp, effects on fecundity and life span of F_1 females, x, 26:45878
- SPHERES**
motion in electrically conducting rotating fluid, combined effect of magnetic field and rotation on, (T), 26:46424
- SPINAL CORD**
radiation effects on enzyme activity and RNA synthesis in neurons of, in mice, 26:45868
radioinduced injuries in human, following therapy of neoplasms, clinical course of latent recovery, x, 26:45797
- SPLEEN**
radiation effects on stem cell renewal in mouse, following fractionated whole-body exposure, x, 26:45871
- SPORES**
radioinduced mutations for radioresistance in microorganism, following sterilization, probability of, 26:45750
radiosensitivity of, biochemical mechanisms of, 26:45751
- STACK DISPOSAL**
monitoring of effluents from reactor, design of remote-controlled sequential sampler for, 26:45613(P)
- STAINLESS STEEL**
analysis for carbon and oxygen by deuteron and proton activation, 26:45309(R) (AE-453)
analysis of irradiated, for oxygen, development of remote apparatus for, 26:45224 (LA-DC-72-524)
analysis of, emission spectrographic methods for, 26:45181 (CISE-R-172)

- carbide precipitation reactions in austenitic, 26:45994(R) (UCLA-34-P-177-10)
- corrosion cracking of Types 304, 304L, 309, 316, and 347, in oxygenated water at 288°C, stress, 26:45966 (BMI-1927)
- corrosion of low-carbon 18/10, by nitric acid in presence of iron, effects of heat treatments on, 26:45970
- creep of tubing of, 26:45417(R) (ORNL-TM-3281)
- diffusion and outgassing of ion implanted-helium in Types 304L, 309, and 316, effects of temperature on, 26:46050 (CONF-710601-, pp 230-47)
- fracture modes of Types 304 and 316, relations with microstructure, 26:45994(R) (UCLA-34-P-177-10)
- grain boundary ledge in Type 304 austenitic, contrast effects at, 26:47630 (UCLA-34-P-177-11)
- neutron reactions (n, α), (n, p), and ($n, 2n$) in fission spectrum, cross sections for, (E), 26:46918 (JEN-230-DME/I-25)
- neutron total cross sections for type-310, from 1.2 to 11 MeV, (E/T), 26:47224 (ORNL-TM-3871)
- passivity of Type 304, effects of plastic deformation on, 26:45967
- production of wrought, vacuum induction melting method for, 26:45981
- properties of Type 316, for use as cesium chloride waste capsules, 26:46099 (ARH-SA-117)
- radiation effects on mechanical properties of Type 304, effects of titanium additions on neutron, 26:46068 (ORNL-TM-3733)
- radiation effects on swelling and creep of, review of fast neutron, 26:47303
- radiation effects on void formation in annealed titanium-modified Type 316, at 400 to 600°C, 1-MeV proton, 26:46044 (AI-AEC-13033)
- radiation effects on void formation in Type 316, at 500°C, relations of helium content with 1-MeV proton, 26:46044 (AI-AEC-13033)
- radiation effects on void formation in Type 316, ion stimulation of fast neutron, 26:46052 (CONF-710601-, pp 430-48)
- radiation effects on void formation in Types 304 and 321, deuteron, 26:46053 (CONF-710601-, pp 479-98)
- radiation effects on void formation in Types 304, 316, and 347, at 525 to 550°C, fast neutron, 26:46048 (CONF-710601-, pp 156-73)
- radioinduced displacement and damage in Type 316, at 500 to 600°C, 0.75 to 1.0 MeV proton, 26:46054 (CONF-710601-, pp 511-21)
- radioinduced void formation in Type 348, effects of niobium content on fast neutron, 26:46049 (CONF-710601-, pp 174-86)
- radioinduced void formation in Types M316, EN58B, and FV548, pile neutron, 26:46047 (CONF-710601-, pp 125-41)
- radioinduced void growth in, use of high-voltage electron microscope in studies of electron, 26:46056 (CONF-710601-, pp 533-49)
- tensile properties of cold worked Type 316, effects of stress and temperature on, 26:46067 (HEDL-TME-71-126)
- void formation in quenched, effects of phosphorus content on, 26:45987 (CONF-710601-, pp 605-12)
- STANDARDS**
- fuel assembly identification, ANSI N18.3-1972, 26:46097
- gaskets for reactor containment vessel airlocks, 26:47593 (RDT-E-14-6-T(6-72))
- photon dosimetry, French national, 26:47216
- specifications for WAK radiochemical plant equipment, 26:46105(T) (ORNL-tr-2371)
- STANNANE, TRIETHYLENETHACRYLOXY-**
- radioinduced polymerization in wood, γ , 26:45355(P)
- STAR CLUSTERS**
- mass of NGC 6656 and NGC 7078, repulsive interaction in universal gravitation for, (T), 26:46749 (ORO-3992-94)
- stars in Pleiades, CH⁺ in circumstellar shells of, (E), 26:46190
- stars in, variation with mass of red and blue supergiant, 26:46177
- STARS**
- angular momentum loss from contact binary, (T), 26:46167
- atmospheres of, convection in, (T), 26:46172
- carbon burning in supergiant, effects of neutrino emission on, (T), 26:46177
- color excess-distance relations for early-type, near planetary nebulae, (), 26:46171
- density of white-dwarf, central, (E/T), 26:46163
- detection of metallic-line, in Orion Ic association, (E), 26:46181
- envelopes of red-giant, dynamic instabilities in, (T), 26:46176
- equation of state for, (T), 26:46170
- evolution of, survey of birth and death processes in, (T), 26:46166
- formation of, influence of local conditions in interstellar medium on, (T), 26:46169
- infrared radiation from circumstellar shells of cool, polarization of 10-micron, (E), 26:46185
- infrared spectroscopy of circumstellar shells, 10-micron, (E), 26:46174
- isophote chart of eta Carinae, (T), 26:46160
- light curves of 32 Cygni, blue and ultraviolet, (E/T), 26:46184
- luminosities of F- and early G-type metal-deficient, (E), 26:46183
- luminosity of red-giant, (T), 26:46176
- masses of circumstellar shells around infrared, (T), 26:46169
- metallicism in border regions of Am domain, (E), 26:46181
- microwave emission from water vapor associated with late M-type, (E/T), 26:46206
- models for differentially rotating, general relativistic analysis of linear pulsations and stability of, 26:46196
- models for differentially rotating, Newtonian analysis of linear pulsations and stability of, 26:46195
- models for uniformly rotating synchronous close binary, (T), 26:46173
- models for white-dwarf, (T), 26:46170
- models for white-dwarf, viscous effects in rapidly rotating, (E/T), 26:46163
- motions of Type A, in Ursa Major stream, proper, (E), 26:46164
- neutrino emission from supergiant, (T), 26:46177
- occurrence of red and blue supergiant, in young open clusters and subgroups of associations, 26:46177
- optical counterpart for Cen X-3, search for, (E), 26:46213
- oxygen burning at balanced power in, (T), 26:46178
- photometry of M- and S-Type Mira variable, near-infrared, (E), 26:46194
- photometry of, in LMC, (E), 26:46238
- proper motion of BD + 4° 3561, periodic perturbations of, 26:46197
- pulsations of long-period variable, mechanism for, (T), 26:46176
- relation of bright, to cosmic x-ray sources, (E), 26:46212
- rotational velocities of Ap, (E), 26:46182
- space motions of F- and early G-Type metal deficient, (E), 26:46183
- spectra of eclipsing binary LR Cen, primary, (E), 26:46191
- spectra of O and Of, He II λ 10124 line in, (E), 26:46180
- spectra of, 6284-A diffuse interstellar line in, (E), 26:46168
- stability of, relativistic velocity-potential hydrodynamics in, (T), 26:46161
- structure of close binary, equilibrium, (T), 26:46173
- thermonuclear reaction rates for heavy ions in, (T), 26:46179
- turbulence in atmospheres of, hydrodynamic character of micro-, (E/T), 26:46188
- ultraviolet spectra of, TD1 data on, 26:46186
- velocities of Type A, in Ursa Major stream, radial, (E), 26:46164
- STEAM**
- corrosive effects of nuclear geothermal, on components in steam-turbine power plants, 26:45487 (UCRL-73939)
- production for Bruce Heavy Water Plant, description of auxiliary generator for, 26:45695
- quality of nuclear geothermal, effects on design and operation of steam-turbine power plants, 26:45487 (UCRL-73939)
- radioisotope content of nuclear geothermal, effects on design and operation of steam-turbine power plants, 26:45487 (UCRL-73939)
- separation from water, performance of steam separators for, 26:45442
- H₂O--steam, density, energy, and pressure in boiler circuits, computer program for analysis of, 26:45458 (CISE-R-245)
- STEAM POWER PLANTS**
- corrosion and scaling in nuclear geothermal, 26:45487 (UCRL-73939)
- cost of, trends in capital, 26:47385
- design and operation of nuclear geothermal, effects of steam quality on, 26:45487 (UCRL-73939)
- design and operation of nuclear geothermal, effects of radioisotope content of steam on, 26:45487 (UCRL-73939)
- STEEL**
- analysis for manganese by neutron activation using plutonium--beryllium source, 26:45199
- carbon distribution in surfaces of, measurement by charged particle and proton activation, 26:45466 (AE-451)
- coatings for, structure of intermediate layers in diffusion, 26:46041
- fracture of heavy-section pressure vessel, mechanics of, 26:45996 (WCAP-7841)
- fracture toughness of pressure vessels of A533B, use of J integral for analysis of, 26:45990 (CONF-720806-2)
- oxide inclusions in, radiometric and activation methods for detection of, 26:46008
- radiation effects on ductility transition temperature of Type A302B, activation fluence method for analysis of neutron, 26:46072
- radioinduced void formation and growth in, comparison of experimental and theoretical data on neutron, 26:46062 (CONF-710601-, pp 769-97)
- sorptive properties for hydrogen during cathodic charging and chemical and electrochemical finishing, review on,

- 26:45986 (BARC-585)
weld crack detection in, factors affecting gamma and x-ray radiographic methods for, 26:45471
weld inspection in chemical munitions containers of, magnetic method for, 26:45470
- STELLAR FLARES**
detection of hard x-ray, from Cyg X-1, (E), 26:46211
- STORAGE RINGS**
description of CERN ISR, 26:46143
electron bundle longitudinal dimensions and phase motion in, particle collective interaction effects on relativistic, (T), 26:46546
experimental facility at electron, description of bremsstrahlung, 26:46145
experimental program at ISABELLE, photoproduction, 26:46426 (BNL-16968)
radiation dose rates outside shielding at CERN ISR, estimates using Moyer method for, 26:46121
- STRATOSPHERE**
circulation in troughs and crests in, relation with high-energy proton behavior, (E/T), 26:46297
fallout content in, to July 1971, effects of nuclear weapons testing on, 26:45518 (HASL-257, pp I.33-50)
planetary transport processes in, research programs on, 26:45503(R) (COO-2195-6)
- STREAMS**
diffusion of heated effluents in, development of mathematical models for prediction of, 26:45575
- STRESS**
distribution in nozzle-to-cylinder connections, finite element method for calculation of, 26:45445
- STRONTIUM**
chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination in rat plasma by neutron activation and radiochemical separation, 26:45202
mass flow choking limit for heat pipe working fluid of, (E), 26:45460
solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369
sorption on barium sulfate--calcium sulfate system, kinetics of, 26:45273 (UJV-2820-Ch)
- STRONTIUM ISOTOPES Sr-85**
uptake and retention in mice following gastric tube administration, effects of aluminum phosphate gel on, 26:45918
- STRONTIUM ISOTOPES Sr-89**
content in atmosphere in Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
content in surface atmosphere at world sites, 1963 through Nov. 1971, 26:45512(R) (HASL-257(App.))
deposition at Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
deposition at weather stations in Atlantic Ocean, June 1965 through 1971, 26:45512(R) (HASL-257(App.))
deposition at world land sites, rate of monthly, 26:45512(R) (HASL-257(App.))
effects on bone marrow of mice following intraperitoneal injection, effects on hematopoiesis and immune reactions, 26:45896
effects on bones and adjacent soft tissues in cats following chronic ingestion, dose dependence of neoplastic, 26:45901
- STRONTIUM ISOTOPES Sr-90**
accumulation and distribution in natural plant associations of alpine, forest, and subalpine belts in Carpathians, 26:45551
accumulation in wheat in podzolic soils, effects of calcium on, 26:45550
content and distribution in open ocean silts from Atlantic Ocean and Mediterranean Sea, 26:45531 (COO-3563-3)
content and distribution in the Caribbean Antilles Arc during 1970, vertical profiles of, 26:45533
content in atmosphere and man in Argentina, 1960 through 1970, 26:45520 (NP-19357)
content in atmosphere in Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
content in cereals, coffee, fruits, milk, and tea in Mecklenburg, GDR, 1969, 26:45546
content in diets in New York City and San Francisco, Jan. to March 1972, 26:45539 (HASL-257, pp II.1-6)
content in drinking water in New York City, Aug. 1954 through Mar. 1972, 26:45512(R) (HASL-257(App.))
content in food products processed from contaminated bovine bones, 26:45549
content in Great Lakes waters, concentration--time model for calculating, 26:45532
content in human bones in New York City and San Francisco, 1971, effects of diet on, 26:45538 (HASL-257, pp I.51-70)
content in marine organisms in north Atlantic Ocean, ecological relations in, 26:45541 (NYO-2174-130)
content in milk at Ispra, Italy, 1960 through 1972, 26:45519 (HASL-257, pp III.53-8)
content in milk in New York City, June 1954 through March 1972, 26:45512(R) (HASL-257(App.))
content in precipitation and surface atmosphere in western US, 26:45517 (HASL-257, pp I.18-32)
content in Rainier rubble chimney rocks, (E), 26:45483 (NVO-1229-180)
- content in sewage and soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
content in stratosphere to July 1971, effects of nuclear weapons testing on, 26:45518 (HASL-257, pp I.33-50)
content in surface atmosphere at world sites, 1963 through Nov. 1971, 26:45512(R) (HASL-257(App.))
content in US surface waters during Jan. to Apr. 1971, 26:45537
content in Western Hemisphere milk samples, during Mar. 1971 to Mar. 1972, 26:45552
content of flour and wheat in France during 1960 to 1969, 26:45903
deposition and content in soils in Salt Lake City, Utah, 1957 to 1971, effects of stratospheric fallout on, 26:45525 (HASL-257, pp I.71-94)
deposition at Ispra, Italy, Sept. 1971 through March 1972, 26:45519 (HASL-257, pp III.53-8)
deposition at weather stations in Atlantic Ocean, June 1965 through 1971, 26:45512(R) (HASL-257(App.))
deposition at world land sites, rate of monthly, 26:45512(R) (HASL-257(App.))
determination in soils, 26:45263 (RT/PROT-72-11)
effects on bones following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
effects on mice following chronic ingestion, effects on litter size and reproduction span of, effects of concentration on, 26:45900
encapsulation of waste, remote welding methods for, 26:46099 (ARH-SA-117)
excretion in urine of children in Poland, relation between dietary intake and, 26:45908
release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)
solubility of Rainier Event rubble-contained, measurement of long-term, 26:45482 (NVO-1229-177)
transport in troposphere, meridional, 26:45516 (HASL-257, pp I.2-17)
transport through soil-plant-small mammal ecosystems at Nevada Test Site, 26:45914 (TID-25954, pp 58-64)
uptake and distribution in tissues of desert plants, relation of content of strontium and other cations in soil to, 26:45923 (TID-25954, pp 69-72)
- STRONTIUM OXIDES**
electric, thermionic, and thermoelectric properties of crystals of, 26:47249
 $K_2O-Nb_2O_5-SrO$, ferroelectric properties and structure of compounds in, 26:45962
 K_2O-SiO_2-SrO , diffusion coefficients for, at 630 to 810°C, measurement of inter-, 26:45946
 SiO_2-SrO , radioinduced optical absorption and paramagnetic centers in glassy, spectral studies on, 26:47226
- STRONTIUM PHOSPHATES**
cathodoluminescence efficiency of doped, absolute, (E), 26:45591
- STRONTIUM URANIUM SELENIDES**
preparation and structure of SrU_2Se_8 , 26:45378
- STRONTIUM ZIRCONATES**
properties for use in magnetohydrodynamic generators, 26:45426(P)
- STRUCTURAL ENGINEERING**
plate response to time-dependent boundary conditions, analysis of, 26:45443
- STRUCTURES**
damage to residential chimney, in Glendale, Calif. from ground motion generated by Feb. 9, 1971 San Fernando earthquake, 26:45474 (JAB-99-88)
gaseous plume diffusion about isolated, of simple geometry, effects of structure orientation and shape on, 26:45514
- STYRENE**
radioinduced polymerization of, effects of peroxide sensitizers on dose rate in γ , 26:45325
butyl acrylate--styrene, radioinduced graft polymerization on cellulose and cellulose triacetate, γ , 26:45320
methyl methacrylate--styrene--vinylphosphonic dichloride, radioinduced polymerization of liquid and solid, reaction mechanisms in, 26:45318
- STYRENE POLYMERS**
radiation resistance of insulators of, effects of organic phosphite additives on, 26:45672(P)
x-ray transmission through, low-energy, (E), 26:47205
polydivinylbenzene--polystyrene, sorptive properties for iodomethane, 26:45388
- SULFATES**
determination of, chemical procedures for, 26:45181 (CISE-R-172)
- SULFUR**
abundance in solar-flare particles, (E), 26:46232
content in atmosphere, isotope ratio tracer method for determining source of, 26:45568
- SULFUR FLUORIDES**
effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331
effects on yields of excited states from pulse radiolysis of deaerated naphthalene--cyclohexane solutions, 26:45330

electron reactions with SF_6 , electron cross sections for electron attachment in, (E), 26:46364 (ORNL-TM-3839)

SULFUR ISOTOPES S-32
meson (K^-) elastic scattering by, scattering lengths for, (E), 26:46912
muon (μ^-) capture by, particle-hole calculation of capture rate and neutron spectra for, (T), 26:46880
nuclear quadrupole moment of, calculation using effective N-N interactions, 26:46892
photon reactions (γ, n) and (γ, p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
proton inelastic scattering at 24.5 MeV, coupled-channels analysis of differential cross sections and asymmetry from polarized, p. 129-34, (E), 26:46873(R) (CEA-N-1522)
proton reactions (p, γ) at 3.36 to 5.41 MeV, excitation functions and gamma spectra from, (E), 26:46902 (BARC-598)
separation from sulfur-33 and -36 by chemical exchange, coefficients for, 26:45688(T) (MLM-1936(TR))

SULFUR ISOTOPES S-33
separation from sulfur-32 by chemical exchange, coefficients for, 26:45688(T) (MLM-1936(TR))

SULFUR ISOTOPES S-34
effects on microwave rotational spectra of ethylene sulfone, 26:45291
proton inelastic scattering at 24.5 MeV, coupled-channels analysis of differential cross sections and asymmetry from polarized, p. 129-34, (E), 26:46873(R) (CEA-N-1522)

SULFUR ISOTOPES S-35
determination in plants by oxygen flask combustion and liquid scintillation counting, 26:45266
effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

SULFUR ISOTOPES S-36
separation from sulfur-32 by chemical exchange, coefficients for, 26:45688(T) (MLM-1936(TR))

SUN
activity of, correlation of 12-month wave in geomagnetic activity with, 26:46316
activity of, effect on 1965-1968 electron content in low and middle latitudes of ionosphere, (E/T), 26:46287
atmosphere of, effect of single- and multiple-ionization losses on charged-particle acceleration in, (T), 26:46230
cycles of 11 years from 1884 to 1967, correlation of geomagnetic-field variations with, 26:46322
eclipse of, thermospheric molecular oxygen density distribution at 95 to 120 km from solar eu occultation measurements, (E), 26:46269
gamma-radiation spectrum from, review of solar-flare-associated production mechanisms of flux and energy, 26:46231
oblateness of, effect of photospheric faculae on, (E/T), 26:46224
oblateness of, effects of faculae on, (E/T), 26:46225
spectra of, continuum near H and K lines of Ca⁺ in, (E/T), 26:46218

SUNSPOTS
cycles of, modulation effect on 20-MeV to 15-GeV cosmic-ray spectra near maximum, (E), 26:46325
longitudes during 20th cycle of, active, (E), 26:46222

SUPERCONDUCTING JUNCTIONS
current-phase relation for thin-film microbridges, one-dimensional Ginzburg-Landau theory treatment of, 26:46806
electron tunneling in N/I/S systems with paramagnetic impurities, (T), 26:46837
thermodynamic properties of Josephson, with normal-metal barrier, (T), 26:46808
tunneling into proximity-effect sandwiches with magnetic impurities on normal side, 26:46822

SUPERCONDUCTIVITY
book: Lectures on Quantum Statistics. Volume 2. Quasi-Averages, 26:47299
theory of time-dependent, and quantum dissipation, 26:46821

SUPERCONDUCTORS
cryogenic problems of forced-cooled systems of, 26:46839
crystal-field splitting effects of rare earth ions in, (T), 26:47269
electric potentials induced in, in absence of transport currents, (T), 26:46835
electron-hole imbalance in, tunneling generation, relaxation, and tunneling detection of, (T), 26:46811
intermediate state of Type-I, resistivity of, (E/T), 26:46823
Josephson effect in, model of, (T), 26:46834
kinetic equation for alloys in weak static fields, (T), 26:46825
magnetic field distributions in Type-II, (T), 26:46833
magnetic flux flow in nonideal Type-II, (T), 26:46819
magnetic states in, localized, (T), 26:46809
magnetic-flux-flow noise in Type-II, (E/T), 26:46815
neutron scattering by vortex lattice in Type-II, London model for inelastic, (T), 26:46820
phenomena in weakly coupled, time-dependent, (T), 26:46827

surface excitations of Type-I, in static magnetic field, (T), 26:46830
thermodynamic properties of small particles of, (T), 26:46813
transitions in microwave region, phase-sensitive method for detecting superconducting, (E), 26:46840
transitions of, resistive, (E/T), 26:46817
vortex lattice of Type-II, calculation of Debye-Waller factor of, 26:46820
vortices in, graphs for B approximately $H/\text{sub } c2/$ in dynamic structure of, (T), 26:46814
vortices in, numerical constants for isolated, (T), 26:46812

SUPERFLUIDITY
book: Lectures on Quantum Statistics. Volume 2. Quasi-Averages, 26:47299

SUPERNOVA REMNANTS
radio emission from five southern, polarization and total power at 2.7 GHz of, (E), 26:46216
radio map of IC 443 at 1.42 GHz, (E), 26:46203
radio polarization aperture synthesis of five possible, (E), 26:46207

SUPERNOVAE
model for Type I, (T), 26:46163
observations of Type I, in NGC 5253, optical, (E), 26:46187

SURFACE-ACTIVE AGENTS
determination of anionic, in water using iron-59, radiometric, 26:45272

SURFACE PROPERTIES
measurement of roundness, method for separating spindle and test ball error in, 26:47635 (UCRL-73805)

SURFACE WATERS
contamination of Amchitka Island, effects of Cannikin Event on, (E), 26:45488 (USGS-474-148)
radioactivity from waste discharge at Nuclear Research Institute, Rez, Czechoslovakia, 26:45565
radioactivity monitoring in US, Oct. 1971, 26:45537
radioactivity monitoring of, in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
strontium-90 content in US, during Jan. to Apr. 1971, 26:45537

SURFACES
bibliography on surface and vacuum physics, 26:46336
properties of, method for determination of, 26:47636 (UCRL-73925)

SURGERY
effectiveness for treatment of radioinduced stenosis of human small intestine, 26:45802
technique for repair of radioinduced rectovaginal fistula in man, 26:45800

SURRY POWER REACTOR, UNIT 1
containment recirculation spray system for, performance evaluation of vertical induction pump motor for, 26:47449 (DOCKET-50280-77)
cooling system for, authorization request to drain, 26:47328 (DOCKET-50280-78)

SURRY POWER REACTOR, UNIT 2
containment recirculation spray system for, performance evaluation of vertical induction pump motor for, 26:47449 (DOCKET-50280-77)

SUSQUEHANNA POWER REACTOR, UNIT 1
effluents from, effects on environment of chemical, radioactive, and thermal, 26:47545 (DOCKET-50387-19)
operation of, cost-benefit analysis for, 26:47545 (DOCKET-50387-19)
safety of, requested additional data on, 26:47548 (DOCKET-50387-22)
safety of, revised data on, 26:47547 (DOCKET-50387-21)
safety of, revised data on, 26:47549 (DOCKET-50387-23)
site for, ecology, geology, hydrology, and meteorology of, 26:47545 (DOCKET-50387-19)
site for, ecology, geology, hydrology, and meteorology of, 26:47546 (DOCKET-50387-20)
site for, ecology, geology, hydrology, and meteorology of, 26:47545 (DOCKET-50387-19)

SUSQUEHANNA POWER REACTOR, UNIT 2
effluents from, effects on environment of chemical, radioactive, and thermal, 26:47545 (DOCKET-50387-19)
operation of, cost-benefit analysis for, 26:47545 (DOCKET-50387-19)
safety of, requested additional data on, 26:47548 (DOCKET-50387-22)
safety of, revised data on, 26:47547 (DOCKET-50387-21)
safety of, revised data on, 26:47549 (DOCKET-50387-23)
site for, ecology, geology, hydrology, and meteorology of, 26:47546 (DOCKET-50387-20)
site for, ecology, geology, hydrology, and meteorology of, 26:47545 (DOCKET-50387-19)

SWEDEN
sodium-22 content of lichen-reindeer-man food chain in, during 1961 to 1969, fallout, 26:45909

SWINE
radiation effects on activity of glutathione peroxidase in blood and liver of, x, 26:45733
radioinduced anemia and leukopenia in newborn, effects of dose on gamma, 26:45852
radioinduced hemorrhage and pneumonitis in newborn, effects of dose and whole-body exposure on gamma, 26:45853

SYMMETRY GROUPS
breaking in quantum theory of preferred fields, 26:46676
breaking relations for octet of transition operators in algebraic hadron model for rates for e^+ and e^- emission

in vector meson decay, restrictions on, 26:46525 (ORO-3992-80)
conformal, for zero-mass component in evolution of universe, 26:46674
Lie algebra for solvable quantum problem, 26:46671
Lie algebra of Virasoro group in dual resonance models, 26:46686
Lorentz, for commutation relations for quantized fields and phase space, 26:46674
Lorentz, for superluminal velocities, four-dimensional, 26:46734
O(3,2) de Sitter, for harmonic oscillator, nonrelativistic contraction of, 26:46670
Poincare group algebra for charged-particle interactions with electromagnetic fields, 26:46678
Poincare, nucleon form factors in field theory on homogeneous space of, 26:46554
relativity groups, structure of, 26:46669
Schroedinger group algebra for many-body system of interacting particles, realizations of, (T), 26:46673
SL(2,c), amplitudes and propagators for hadrons in representations of, 26:46603
SL(2,R), for generalized dual resonance model, 26:46593
Sp(2n,R) algebra, irreducible representations for para-Bose, 26:46507
SU(2,2), irreducible representations of nondegenerate complementary series of, 26:46672
SU(3) internal symmetry breaking by hadrons, mathematical characterization of, 26:46682
SU(3), algebraic method with Bose operators for unitary representations of, 26:46675
SU(3), nonet structure of spin--parity--charge conjugation parity = 3-- mesons in, 26:46611
SU(3), properties for existence of quarks, 26:46683
SU(6), exchange degeneracy for baryon mixing in 70 supermultiplet of, 26:46685
SU(6), for tensor meson production in $\gamma + N \rightarrow$ baryon + tensor meson interactions, 26:46684
SU(n), eigenvalues of invariant operators of unitary unimodular, 26:46668
theory for nuclear, atomic, and complex structures, review of, (T), 26:46508
U(4) x U(4), quartet scheme of hadrons in chiral, 26:46687

SYNCHROCYCLOTRONS
space charge for different beam shapes in central region of, calculation of, 26:46140

SYNCHROTRON RADIATION
emission from magnetically confined hot-electron plasma, (E), 26:47144
emission in electron ultra-luminous motion in medium in magnetic field, spin effects on intensity of, (T), 26:46548

SYNCHROTRONS
description of 2-MeV electron undulator, at Tomsk Polytechnic Inst, 26:46120
radiation dose rates outside shielding at high-energy proton, Moyer method for estimating, 26:46121

T

T-111 (TANTALUM ALLOY)
mechanical properties of, at high temperatures, effects of gaseous impurities on, 26:45703(R) (ORNL-4796)
properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)

TACHYONS
annihilation by antitachyons, Lorentz-invariant formulation of Cherenkov radiation emission for, (T), 26:46704 (N-72-19766)
Cherenkov radiation emission by, Lorentz-invariant formulation for, (T), 26:46704 (N-72-19766)
field equations for dust metric of, in general relativity theory, 26:46754
Lorentz group transformations for, 26:46734
relativity group for, representations of, 26:46724
theory for, Lorentz-invariant formulation of Cherenkov radiation emission for Cauchy data in, (T), 26:46704 (N-72-19766)
theory of, one-dimensional causal, 26:46499

TANKS
coating with plastic for corrosion protection in solvent extraction processes, 26:46080

TANTALUM
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination by electron probe microanalysis methods, 26:45237
determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
electron reactions at 12 to 30 MeV, target thickness effects on positron spectra from, (E), 26:46149
photon reactions at 1.160 GeV at 1500, photoemulsion study of kinetic energy and total yield of charged pions from, (E), 26:46441
separation from other metals by combined ion exchange--solvent extraction chromatography, 26:45282

TANTALUM ALLOYS AND SYSTEMS
Ir--Ni--Ta, phase studies on, 26:46010
Nb-base--Ta, hardening of high-purity, studies of substitutional solid solution, 26:45991(R) (COO-916-25)

149 SUBJECTS

TERBIUM ISOTOPES Tb-150

Ta--Ti-base, phase transformations in, relation of Martensite crystallography to Burgers and Bogers--Burgers, 26:46037

TANTALUM CARBIDES
magnetic susceptibility of cubic, at low temperatures, 26:47266

TANTALUM ISOTOPES Ta-175
electron-capture decay of, γ and conversion electron spectra from, (E), 26:46979

TANTALUM ISOTOPES Ta-181
alpha inelastic scattering up to 80 MeV, α spectra and cross section for, (E), 26:47008
alpha reactions (α ,He-3) up to 80 MeV, energy spectra and cross section for, (E), 26:47008
boron-10 reactions at 50 to 100 MeV, analysis of compound nucleus deexcitation using statistical evaporation model, (E/T), 26:46974 (FRNC-TH-207)
boron-11 reactions at 50 to 100 MeV, analysis of compound nucleus deexcitation using statistical evaporation model, (E/T), 26:46974 (FRNC-TH-207)
energy-level transition in, nuclear part of parity nonconserving circular polarization of 482 keV γ -ray from, (T), 26:46982
helium-3 inelastic scattering up to 80 MeV, energy spectra and cross section for, (E), 26:47008
helium-3 reactions (He-3, α) up to 80 MeV, α spectra and cross section for, (E), 26:47008
photon reactions (γ ,n) and (γ ,p) up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890
use as label for determining migration of salamanders, 26:45815(R) (COO-3486-1)

TANTALUM ISOTOPES Ta-182
energy level of, Nilsson-model interpretation of ground-state, (T), 26:46983
use as label for determining migration of salamanders, 26:45815(R) (COO-3486-1)

TANTALUM OXIDES
preparation of framed films of, for electrolytic membranes and radiation windows, 26:45958(P)

TAURINE
radioprotective effects of, on human marrow cells, use of chromosome fragmentation counts for determining, 26:45734

TD-NICKEL
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)

TD-NICKEL CHROMIUM
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
properties for use in space shuttle, 26:45934 (N-72-18585)

TECHNETIUM
magnetic properties of superconducting, with hexagonal symmetry, (T), 26:46807
separation from spent fuels, 26:45423(P)

TECHNETIUM ISOTOPES Tc-99
energy levels from molybdenum-99 decay, lifetimes of, (E), 26:46951
preparation of human albumin-incorporated metastable, method for, 26:45927(T) (TT-54002/4, pp 471-4)

TELLURIUM
chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
purification by zone melting in rotating container, 26:45276

TELLURIUM ISOTOPES Te-125
energy level at 462 keV, g-factor for, (E), 26:46957

TELLURIUM ISOTOPES Te-126
energy-level scheme from antimony-126 decay, (E), 26:46944

TELLURIUM ISOTOPES Te-128
energy-level scheme from antimony-128 decay, (E), 26:46944

TELLURIUM ISOTOPES Te-130
energy-level scheme from antimony-130 decay, (E), 26:46944

TELLURIUM ISOTOPES Te-132
range of fission fragments of, in aluminum--uranium alloys and uranium oxides, 26:46074

TERBIUM
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination by luminescent method using pyrogallolsulfonic acid, 26:45231
magnetic susceptibility of single crystals of, anisotropy in para-, (E), 26:47253

TERBIUM ALLOYS AND SYSTEMS
Cr--Tb, hardness and metallography of, 26:46033
Cr--Tb, phase diagram at 600 to 1900°C, 26:46033
Fe--Tb, magnetic compensation in films of, 26:47273

TERBIUM BOROCARBIDES
preparation, properties, and structure of, 26:45964

TERBIUM ISOTOPES Tb-148
beta and electron-capture decay of, γ and conversion electron spectra from, (E), 26:46942 (JINR-P6-6512)

TERBIUM ISOTOPES Tb-150
beta and electron-capture decay of, γ and conversion

electron spectra and half-life for, (E), 26:46942 (JINR-P6-6512)

TERBIUM ISOTOPES Tb-160
 beta decay of, $e - \gamma$ angular correlations in, (E), 26:46991
 energy level of, Nilsson-model interpretation of ground-state, (T), 26:46983

TEREPHTHALIC ACID, POLYETHYLENE ESTERS
 radioinduced photoconductivity in films of, x, (E/T), 26:45668 (UCRL-73974)

THALLIUM
 mass flow choking limit for heat pipe working fluid of, (E), 26:45460
 positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268

THALLIUM ISOTOPES Tl-189
 decay of, half-life and gamma spectrum from, (E), 26:46999

THALLIUM ISOTOPES Tl-203
 gamma and x-ray spectra from μ -mesic atoms of, (E), 26:47031
 x-ray emission in muonic, test of quantum electrodynamic corrections from, 26:46708

THALLIUM ISOTOPES Tl-205
 gamma and x-ray spectra from μ -mesic atoms of, (E), 26:47031

THALLIUM ISOTOPES Tl-208
 charge state of, formed by alpha decay of bismuth-212, 26:45359
 content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)

THERMAL CONDUCTIVITY
 conference on, Washington, D. C., Dec. 1, 1971, 26:45463

THERMAL EXPANSION
 measurement of, conference on, 26:45463

THERMAL INSULATION
 fabrication of low-density, using Fiberfrax paper and rigidizers, 26:47601(R) (BDX-613-288(Rev.))

THERMAL PROPERTIES
 measurement of, conference on, 26:45463

THERMIONIC CONVERTERS
 heat sources for plutonium-238-fueled, design of fuel capsules for, 26:45708(P)

THERMOELECTRIC GENERATORS
 design and performance of, 26:45432
 fuel systems for radioisotope-fueled, development of, 26:45701(R) (DP-1300)
 heat source for, neutron spectra characteristics of plutonium-238 dioxide, 26:45704
 heat sources for plutonium-238-fueled, design of fuel capsules for, 26:45708(P)
 thermal input to, from radioisotope heat source, design of heat pipe for controlling, 26:45427(P)

THERMONUCLEAR REACTIONS
 combustion waves in bounded plasma, 26:47172

THERMONUCLEAR REACTORS
 blanket calculations for, iron effects on neutron, 26:47176
 breeding ratios and inventory calculations for deuterium-tritium fueled, 26:47173
 design of toroidal, with electron sheath, 26:47174(P)
 design using electrostatic confinement, 26:47113(P)
 development of, review of, 26:47179
 energy conversion scheme for losses from, 26:45433(P)
 magnetohydrodynamic generator use with, economics of, 26:47170 (IPP-IV-32)
 plasma displacement calculations for Harmonica 2, nonlinear, (T), 26:47115

THERMOSPHERE
 oxygen density distribution at 95 to 120 km in, from solar oc occultation measurements, molecular, (E), 26:46269

THICKNESS GAGES
 description of beta backscattering, for thin films, 26:45656(P)
 description of radiometric, for wire thickness measurements, 26:45717
 description of radiometric, with automatic control, 26:45654(P)

THIOLS
 effects on survival of x-irradiated sea urchin eggs, 26:45882
 radiation effects on content of, in mitochondria in rat heart and liver, 26:45829
 reactions with free radicals, deuterium and tritium isotope effects on hydrogen abstraction in, 26:45342

THORIUM
 chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
 complexes, stability constants for acetate, 26:45373
 complexes, stability constants for propionate, 26:45373
 compounds, ionization in mass spectrometry, 26:45258
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 determination of, chemical procedures for, 26:45181 (CISE-R-172)
 determination using phenylthiazolylazochromotropic acid, photocolometric, 26:45240
 effects on spectrophotometric determination of phosphate

by extracting molybdophosphoric acid in butyl acetate in presence of EDTA and fluoride, 26:45219 (ARR-SA-110)
 separation from other metals by combined ion exchange-solvent extraction chromatography, 26:45282
 separation of protactinium from, by sorption on silica gel, 26:45390
 separation of radium from, method for, 26:47014 (FRNC-TH-184)
 solvation numbers from compressibilities and ionic vibration potentials measurements, 26:45369

THORIUM ALLOYS AND SYSTEMS
 Os-base--Th, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248
 Re-base--Th, electric and thermal conductivities of, effects of temperature in ranges 100 to 1700°C and 100 to 1200°C, respectively, 26:47248
 Th--U--Zr, phase studies on liquidus and solidus surfaces in, 26:46003

THORIUM ISOTOPES
 determination of ratios of, by mass spectroscopy, 26:45258

THORIUM ISOTOPES Th-228
 content in plants and soils in uraniferous area in New Zealand, 26:45529
 effects on bones following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
 energy-level scheme from actinium-228 decay, (E), 26:47014 (FRNC-TH-184)

THORIUM ISOTOPES Th-230
 alpha decay of, conversion electron spectra from, (E), 26:47028
 determination relative to thorium-232 in thorium nitrate by mass spectroscopy, 26:45258
 deuteron reactions (d,pf) at 15 MeV, comparison of subbarrier resonances with predictions of Strutinsky theory, (E/T), 26:47033
 triton reactions (t,pf) at 15 MeV, comparison of subbarrier resonances with predictions of Strutinsky theory, (E/T), 26:47033

THORIUM ISOTOPES Th-231
 energy levels from thorium-232 (d,t) reactions, spin-parity assignments for, (E), 26:47029

THORIUM ISOTOPES Th-232
 determination relative to thorium-230 in thorium nitrate by mass spectroscopy, 26:45258
 deuteron reactions (d,p) and (d,t) at 13.1 MeV, Q-value and angular distributions for, (E), 26:47029
 effects on lungs following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)
 krypton-84 reactions at 450 to 494 MeV, excitation functions for production of thorium-225 thorium-226, actinium-223, and actinium-224 in, (E), 26:47035
 neutron fission at 14 MeV, independent and cumulative yields of rare-gas isotopes from, (E), 26:47030
 photofission of cross section for, (E), 26:46890
 triton reactions (t,pf) at 15 MeV, comparison of subbarrier resonances with predictions of Strutinsky theory, (E/T), 26:47033

THORIUM ISOTOPES Th-233
 energy levels from thorium-232 (d,p) reactions, spin-parity assignments for, (E), 26:47029

THORIUM NIOBIUM TUNGSTEN BRONZE
 crystal structure of high-temperature phases in, 26:45953

THORIUM NITRATES
 analysis for thorium isotope ratios by mass spectroscopy, 26:45258

THORIUM OXALATES
 precipitation of, optimization studies on, 26:45365 (IEA-183)

THORIUM OXIDES ThO₂
 microstructure of transparent yttrium oxide-doped, 26:45948
 preparation of transparent yttrium oxide-doped, method for, 26:45948
 transport properties of, at 500 to 1600°C, effects of oxygen pressure on, 26:45938
 Al--Cr--Ni--ThO₂, properties of dispersion-strengthened, for use in space shuttle, 26:45934 (N-72-18585)
 Mo--ThO₂, porosity and tensile properties of test fuels of, 26:45703(R) (ORNL-4796)
 ThO₂--UO₂, radiation effects on crystal lattice of fuels of, fission fragment, 26:45939
 ThO₂--Y₂O₃, transport properties of, at 500 to 1600°C, effects of oxygen pressure on, 26:45938

THORIUM TANTALUM TUNGSTEN BRONZE
 crystal structure of high-temperature phases in, 26:45953

THREE MILE ISLAND POWER REACTOR, UNIT 1
 containment structure ring girder for, additional data on construction of, 26:47459 (DOCKET-50289-61)
 safety of, revised and additional data on, 26:47461 (DOCKET-50289-63)
 valves for, deficient wall thickness for cast stainless steel, 26:47460 (DOCKET-50289-62)

THULIUM
 determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
 separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285

THULIUM ALLOYS AND SYSTEMS

Sb--Tm, magnetization of TmSb at 1.5 and 20.4°K, anisotropic dc, 26:47258

THULIUM BOROCARBIDES

preparation, properties, and structure of, 26:45964

THULIUM ISOTOPES Tm-169

alpha reactions ($\alpha,2n\gamma$) at 28 MeV, γ -ray angular distributions and excitation functions for, (E), 26:46995

alpha reactions ($\alpha,2n\gamma$) with, γ spectra from, (E), 26:46980

alpha reactions ($\alpha,2n\gamma$) with, γ - γ coincidence spectra from, (E), 26:46987

use as internal power supply for compact solid-state laser, 26:45677(P)

THULIUM ISOTOPES Tm-173

energy levels from erbium-173 beta decay, Nilsson and spin--parity assignments for, (E), 26:46997

energy-level transitions from erbium-173 beta decay, branching ratios, K-conversion coefficients, and multipolarities for, (E), 26:46997

isomeric transition from erbium-173 beta decay half-life for, (E), 26:46997

THYMINE

hydroperoxides, effects of radiolytic, on compounds, 26:45315(R) (COO-3286-3)

radiation chemistry of, 26:45315(R) (COO-3286-3)

radiation chemistry of aqueous solutions of, thymine glycols in, 26:45314 (COO-3286-1)

THYMUS GLAND

effects of surgical removal of, on regeneration of bone marrow lymphoid cells in x-irradiated mice, 26:45830

radiation effects on lymphomagenic properties of neonatal mouse, subsequently transplanted into Gross murine leukemia virus-infected hosts, γ , 26:45818 (UCLA-12-867)

radiosensitivity of human, relation to count of mast cells in blood following exposure of mediastinum, x, 26:45810

THYROID GLAND

iodine-131 uptake by human, following administration, diagnostic value of determining rate of, 26:45906(T) (TT-54002/4, pp 462-70)

radiation dose to, of adult and fetal mice following injection of iodine-131, mathematical model for calculating β , 26:45902

radiation effects on growth of, in goitrogen-stimulated adult and normal fetal mice, comparison of iodine-131 β and γ and external x, 26:45865

radiation effects on human, following iodine-131 uptake, effects on thyroglobulin-induced stimulation of lymphocytes, β and γ , 26:45889

radioinduced neoplasms in, of dogs, combined iodine-131 β and γ and external x, 26:45819 (UR-3490-63)

radioinduced tumors of, in mammals following uptake of iodine-131, review, 26:45895 (UR-3490-102)

TIMING CIRCUITS

design for measuring energy-level lifetimes, p. 73-6, 26:46873(R) (CEA-N-1522)

TIN

carbon-ion reactions at 10 MeV/amu with, energy shift and relative intensity of K x rays from, (E), 26:46339

determination in aluminum-, copper-, and zirconium-base alloys by alternating current polarographic method, 26:45256

determination in high-purity cadmium oxide, spectroscopic, 26:45220 (BARC-593)

determination of, chemical procedures for, 26:45181 (CISE-R-172)

ion reactions at 5 MeV/amu with, dependence of energy shift and relative intensity of K x rays from, on ion stopping power, (E), 26:46339

positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268

ultrasonic attenuation in superconducting, effect of strain on, 26:46824

TIN ALLOYS AND SYSTEMS

Hg--In--Sn, thermodynamic properties of amalgams in, at 150 to 340°C, 26:46004

In--Sn-base, transitions of superconducting, resistive, (E/T), 26:46817

In-base--Sn, ultrasonic attenuation in superconducting, (E/T), 26:46816

Pb--Sn, photoelectric process in various shells in, effective atomic numbers in, (T), 26:47206

TIN ISOTOPES

proton reactions (p,n) at subthreshold energies, shell-model analysis of, (T), 26:46952

TIN ISOTOPES Sn-116

electron inelastic scattering at 209 MeV, differential cross sections for, p. 189-91, (E), 26:46873(R) (CEA-N-1522)

TIN ISOTOPES Sn-117

alpha inelastic scattering at 10, 10.5, and 11 MeV, γ -ray angular distributions and spectra for, (E), 26:46961

energy levels from alpha and oxygen-16 Coulomb excitation, lifetimes, spin--parity assignments, and transition probabilities for, (E), 26:46961

oxygen-16 inelastic scattering at 45 MeV, γ -ray angular distributions and spectra for, (E), 26:46961

TIN ISOTOPES Sn-119

alpha inelastic scattering at 10, 10.5, and 11 MeV, γ -ray

151 SUBJECTS

TRACE ELEMENTS

angular distributions and spectra for, (E), 26:46961

energy levels from alpha and oxygen-16 Coulomb excitation, lifetimes, spin--parity assignments, and transition probabilities for, (E), 26:46961

oxygen-16 inelastic scattering at 45 MeV, γ -ray angular distributions and spectra for, (E), 26:46961

proton reactions (p,n) at 14 to 44 MeV with subsequent proton decay, coupled-channels calculations of energy dependence of, (T), 26:47073

TIN ISOTOPES Sn-120

electron inelastic scattering at 209 MeV, differential cross sections for, p. 189-91, (E), 26:46873(R) (CEA-N-1522)

energy levels from deuteron inelastic scattering, particle-hole, (E), 26:46943 (N-72-18705)

lithium-6 elastic scattering at 30 MeV, angular distribution for, (E), 26:46883

TIN ISOTOPES Sn-124

electron inelastic scattering at 209 MeV, differential cross sections for, p. 189-91, (E), 26:46873(R) (CEA-N-1522)

TISSUES

radiation effects on mammalian, model for recovery from, 26:45816 (FRNC-R-7)

TITANIUM

determination by spectrography, spectrochemical buffers and volatilization mechanisms in, 26:45222 (JEN-229-DQ/I-79)

determination in meteorites by proton activation, 26:45215

effects on mechanical properties of neutron-irradiated Type 304 stainless steel, 26:46068 (ORNL-TM-3733)

TITANIUM ALLOYS AND SYSTEMS

decomposition of supersaturated solid solutions of, effects of aging on mechanism of, 26:46032

Al--Cr--Fe--Mo--Si--Ti-base, fatigue strength of, effects of mechanical surface hardening on low-cycle, 26:46009

Al-base--Ti, superconductivity and localized spin fluctuations in, (E/T), 26:46838

Hf--Pt-base--Rh--Ti--W, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)

Nb--Ti, phase transformation and structure of solid solutions of, mechanism of $\beta + \alpha$, 26:46036

Nb--Ti-base, phase transformations in, relation of martensite crystallography to Burgers and Bogers--Burgers, 26:46037

Nb-base--Ti, fabrication of aluminum-clad wires of, method for, 26:45972(R) (COO-3311-1)

Ta--Ti-base, phase transformations in, relation of martensite crystallography to Burgers and Bogers--Burgers, 26:46037

Ti--Zr, diffusion in β -phase of, mechanism of, 26:45985 (BARC-577)

TITANIUM CARBIDES

Co--TiC--WC, metallography of, methods for, 26:45955

TiC--WC, metallography of, methods for, 26:45955

TITANIUM HYDRIDES

phase studies of, 26:45942

use as hydrogen source in neutron columns, 26:45942

TITANIUM ISOTOPES Ti-46

helium-3 reactions (He-3,t) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930

TITANIUM ISOTOPES Ti-48

neutron capture at 25 keV, gamma spectrum from, (E), 26:46917(T) (BNL-tr-477)

TITANIUM ISOTOPES Ti-49

energy levels from neutron capture by titanium-48, (E), 26:46917(T) (BNL-tr-477)

TITANIUM OXIDES

effects on thermal expansion of powdered uranium dioxide lattice during sintering, 26:45956

energy level diagrams for, spectroscopic studies on high-temperature, 26:45932 (LBL-450)

thermodynamic properties of, at high temperatures, 26:45932 (LBL-450)

Al₂O₃--CaO--Cr₂O₃--F--MgO--MoO₃--TiO₂--ZrO₂, use as welding flux, 26:45941(P)

TOKAMAK DEVICES

blanket calculations for, iron effects on neutron, 26:47176

design of air-core, 26:47177(P)

drift-kinetic equation and electrostatic instabilities in, (T), 26:47166

magnetic field measurement using scattered laser beams, 26:47120

plasma displacement calculations for, nonlinear, (T), 26:47115

TOLUENE

effectiveness for enhancing semi-in vitro repair of γ -radioinduced single-strand breaks in transforming DNA of Bacillus subtilis, 26:45731

radiolysis of aqueous acid solutions of, spectra of benzyl radicals from pulse, 26:45309(R) (AE-453)

TOXINS

biological properties of organic, with C--F bonds, 26:47637 (UR-49-1508)

TRACE ELEMENTS

content and distribution of metallic, in arctic and subarctic marine environment during June 1971 to June

1972, 26:47623(R) (RLO-2229-T-1-18)
determination in food by neutron activation and radiochemical separation, 26:45189(R) (BFL-1/1972)
determination in hair by neutron activation and γ spectroscopy, 26:45214
localization in biological specimens, development of scanning electron microscope for, 26:45893 (ORO-2690-12(Pt.B), Paper 13)

TRACER TECHNIQUES
methods for measuring flow through high-head pumps and turbines using radioisotopes, 26:45721

TRACHEA
mucociliary clearance mechanism in rat, effects of chronic low-level nitrogen dioxide on, 26:47644 (UR-3490-85)

TRANSACTINIDE ELEMENTS
existence of superheavy, in solar system, 26:47016 (INDC-7-U, pp 23-5)

TRANSUDERS
development of pressure, for measurement of fuel reprocessing plant input weight, 26:46093 (ICP-1014)

TRANSFERASES
proton reactions with RNase, cross sections for inactivation in, (T), 26:46365(R) (ORO-2911-14)
role of DNA polymerase and endonuclease in repair of uv-radioinduced damage in DNA in vitro, 26:45727
role of DNA polymerase in repair of uv-radioinduced damage to DNA, 26:45722(R) (COO-3232-1)
role of endonuclease in repair of uv-radioinduced damage to DNA of *Micrococcus luteus*, 26:45722(R) (COO-3232-1)

TRANSISTORS
radiation effects on microwave bipolar, neutron and gamma, (E/T), 26:45663 (SC-DC-721233)
radiation effects on passivation silicon bipolar, 26:45662 (SC-DC-721232)

TRANSPLANTS
radiation effects on regeneration of blood vessels in muscle, in rats, 26:45857

TRANSPORT THEORY
Boltzmann equation in, random method for, 26:47208
calculation for neutral particle, using double Fourier transformation and Wiener--Hopf technique, 26:46854
discrete-ordinates methods for numerical solution of, review of, 26:47211

TRANSURANIUM ELEMENTS
decay of, ratios of x-ray to alpha intensities in, effects of isotopic composition on, (E), 26:47011 (BNWL-1648)
determination of small amounts of, by x-ray fluorescence analysis using α -x coincidence technique, 26:45270
hazards of, effects of changing isotopic composition on biological, (E), 26:47011 (BNWL-1648)
hydrolysis of, partition chromatography for studies on, 26:45366 (JINR-P6-6402)

TRANSURANIUM PROCESSING PLANT
iodine evolution and trapping studies at, 26:45409(R) (ORNL-TM-2996)
iodine evolution from dissolver solutions in, 26:45407(R) (ORNL-TM-2918)

TRIETHYLAMINE
deuterium concentration by, for production of heavy water, 26:45696(P)

TRIMETHYLAMINE
deuterium concentration by, for production of heavy water, 26:45696(P)

TRITIUM
content in central and northwest Algerian aquifers, 26:45558
content in milk, sewage, vegetation, and water in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
distribution in fuel reprocessing cycle at ICPP, 26:45405(R) (ICP-1006)
distribution in Zircaloy-clad uranium dioxide fuel rods, 26:45414(R) (ORNL-TM-3180)
effectiveness for inducing breaks in DNA of eukaryote cells, 26:45894 (ORO-2690-12(Pt.B), Paper 14)
effects on abstraction of hydrogen from thiols by free radicals, isotope, 26:45342
effects on cells of mice following injection as labeled leukemia cells, 26:45892 (ORO-2690-12(Pt.B), Paper 4)
hyperfine structure of ground-state, pressure shift in argon, (E), 26:46347
migration in environment, radiation burden from, 26:45509 (KFK-1560)
monitoring in urine, using acetylene from calcium carbide reactions, flowthrough ionization chamber for, 26:45585 (AERE-HP/M-74)
production in molten salt reactors, control of, 26:47302
reactions of recoil, variation of energy loss parameter in Estrup--Wolfgang theory with energy in, 26:45363
release at West Valley Processing Plant, April-June, 1972, 26:45398 (DOCKET-50201-125)
release in reprocessing of oxide fuels, 26:45416(R) (ORNL-TM-3250)
separation from air by cold trapping, 26:45416(R) (ORNL-TM-3250)
volatilization from oxide fuels, 26:45409(R) (ORNL-TM-2996)

TRITIUM SYSTEMS

D--T, breeding ratios and inventory calculations for use in thermonuclear reactors, 26:47173

TRITONS

nuclear magnetic moment of, effects of two-pion contribution to proton electromagnetic structure on, (T), 26:46867
reactions (t, α) with magnesium-24 at 15 MeV, angular distributions for, (E), 26:46915
reactions (t,p) with nuclei, repulsive core effects in, (T), 26:47078
reactions (t,pf) with thorium-230 and -232 at 15 MeV, comparison of subbarrier resonances with predictions of Strutinsky theory, (E/T), 26:47033

TROJAN POWER REACTOR, UNIT 1

construction of, licensing board decision on, 26:47515 (DOCKET-50344-29)
environmental report for, revised and additional data for, 26:47516 (DOCKET-50344-30)

TROPOSPHERE

circulation in troughs and crests in, relation with high-energy proton behavior, (E/T), 26:46297
circulation in, nature of cosmic-related semi-secular cycle in, (E/T), 26:46296
planetary transport processes in, research programs on, 26:45503(R) (COO-2195-6)
strontium-90 transport in, meridional, 26:45516 (HASL-257, pp 1.2-17)

TRYPTOPHAN

content of Ehrlich ascites cells in relation to x-radioinduced increase in uv fluorescence, 26:45736

TUNGSTEN

brittle--ductile transition in sheets of, transition temperatures for, 26:45993(T) (ORNL-tr-2377)
compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)
damping of elastic vibrations in monocrystals of, at 100 to 700°C, effects of orientation on, 26:46022
determination by spectrography, spectrochemical buffers and volatilization mechanisms in, 26:45222 (JEN-229-DQ/I-79)
dislocation energy of impurity atoms in solid solutions of, internal friction method for studying, 26:46023
effects on microstructure and properties of nickel-base--niobium alloy, 26:46035
effects on recrystallization of iron, 26:45982
electron density distribution on (001) plane of, from neutron-diffraction and x-ray-scattering measurements, 26:47254
erosion of thoriated cathode of, by electric current, protective effects of nitrogen against, 26:45968
microscopy of bcc, model for studying symmetry of self-interstitial atoms in field ion, 26:46029
microscopy of, imaging process in field ion, 26:46038
radiation effects on mechanical properties of, at low temperatures, 10-MeV proton, 26:46066(R) (COO-3431-1)
radiation effects on thermionic fuel cladding of, fission fragment, 26:45702(R) (LA-4978)
texture of annealed and rolled foils of, effects of impurities on, 26:46042
work function of barium sorbed on (110) single crystals of, temperature dependence of, (T), 26:47257

TUNGSTEN ALLOYS AND SYSTEMS

decomposition of supersaturated solid solutions of, effects of aging on mechanism of, 26:46032
Co--Fe--W, crystal structure of, during aging, effects of molybdenum on, 26:46006
Co--W, deposition of, effects of current density and temperature on electrolytic, 26:45977
Co--W, electrodeposition of, effects of temperature on, 26:45983
Hf--Pt-base--Rh--Ti--W, properties for use in fuel capsule fabrication, 26:45703(R) (ORNL-4796)
Mo--Nb-base--W, tensile properties at 900°K, athermal, 26:45991(R) (COO-916-25)
Nb-base--W, hardening of high-purity studies of substitutional solid solution, 26:45991(R) (COO-916-25)
Pt-base--Rh--W, properties of sheets of, for fuel capsule fabrication, 26:45703(R) (ORNL-4796)
Re--W-base, compatibility of fuel capsules of, with cobalt at 850 to 1000°C, 26:45701(R) (DP-1300)
Re--W-base, corrosion by marine environment, weight change data on, 26:45701(R) (DP-1300)

TUNGSTEN CARBIDES

Co--TiC--WC, metallography of, methods for, 26:45955
Co--WC, metallography of, methods for, 26:45955
TiC--WC, metallography of, methods for, 26:45955

TUNGSTEN COUPLES

Re/W, radiation effects on properties of, at 2000°C, in pile, 26:45438 (KFK-1556)

TUNGSTEN ISOTOPES W-170

half-life of, (E), 26:47001

TUNGSTEN ISOTOPES W-171

half-life of, (E), 26:47001

TUNGSTEN ISOTOPES W-179

energy levels from rhenium-179 decay, 26:46976 (JINR-P6-6492)

TUNGSTEN ISOTOPES W-181

distribution in tissues of newts, 26:45913(R) (RLO-2227-T-8-3)

energy-level scheme from rhenium-182 decay, (E),
26:46978

TUNGSTEN OXIDES

preparation of framed films of, for electrolytic
membranes and radiation windows, 26:45958(P)

TURBINE BLADES

development of 50-Hz 44-in., for 1000-MW class steam
turbines, 26:45454

TURBINES

flow through high-head, radiosotope tracer measurements
of, 26:45721

valves for high-rated conventional and nuclear steam,
development of, 26:45455

TURKEY POINT POWER REACTOR, UNIT 3

effluents from, effects on environment of chemical,
radioactive, and thermal, 26:47413 (DOCKET-50250-84)

emergency power system for, malfunction of busses in,
26:47415 (DOCKET-50250-86)

operation of, cost-benefit analysis for, 26:47413
(DOCKET-50250-84)

operation of, license for, 26:47414 (DOCKET-50250-85)

power tilt limits for, evaluation of, 26:47416 (DOCKET-
50250-88)

site for, ecology, geology, hydrology, and meteorology
of, 26:47413 (DOCKET-50250-84)

technical specifications for, 26:47414 (DOCKET-50250-85)

TURKEY POINT POWER REACTOR, UNIT 4

containment structure for, structural integrity test for,
26:47417 (DOCKET-50251-72)

effluents from, effects on environment of chemical,
radioactive, and thermal, 26:47413 (DOCKET-50250-84)

operation of, cost-benefit analysis for, 26:47413
(DOCKET-50250-84)

operation of, license for, 26:47414 (DOCKET-50250-85)

power tilt limits for, evaluation of, 26:47416 (DOCKET-
50250-88)

site for, ecology, geology, hydrology, and meteorology
of, 26:47413 (DOCKET-50250-84)

technical specifications for, 26:47414 (DOCKET-50250-85)

TZM (MOLYBDENUM ALLOY)

mechanical properties of, at high temperatures, effects
of gaseous impurities on, 26:45703(R) (ORNL-4796)

properties for use in fuel capsule fabrication,
26:45703(R) (ORNL-4796)

U

ULTRASONIC TESTING

method for determining location and size of defects in
metals using frequency analysis system, 26:45469(P)

method for fuel cladding, 26:45472(P)

ULTRAVIOLET RADIATION

effects on alkylation of caffeine by amines, 26:45326

effects on cerium oxide crystals, ESR spectra of hole
centers produced in, 26:47233

effects on coding activity of RNA of bacteriophage R17 in
vitro, 26:45728

effects on cultured squamous cells of eyelids of cattle,
26:45851

effects on DNA in cultured human Xeroderma pigmentosum
cells in comparison to effects of chemical carcinogens
and mutagens, extent of repair synthesis following,
26:45742

effects on DNA of *Micrococcus luteus*, role of enzymes in
repair of, 26:45722(R) (COO-3232-1)

effects on DNA synthesis in HeLa cells following
microbeam exposure of half-nuclei, effects of
hydroxyurea on, 26:45737

effects on DNA, role of DNA ligase, DNA polymerase, and
endonuclease in repair of in vitro, 26:45727

UNITED KINGDOM

nuclear power plants for, site selection criteria for,
26:47574

UNITED STATES

fallout content in precipitation and surface atmosphere
in western, 26:45517 (HASL-257, pp I.18-32)

fallout monitoring in atmosphere, bones, diets, and soils
in, 26:45511(R) (HASL-257)

nuclear power plants for, site selection criteria for,
26:47574

radioactivity monitoring of atmospheric precipitation and
ground-level atmosphere in, during Feb. 1972, gross
beta, 26:45523

radioactivity monitoring of surface waters in, during
1971, 26:45537

radioisotope content of milk in, during Mar. 1971 to Mar.
1972, cesium-137 and strontium-90, 26:45552

strontium-90 content in human bones in New York City and
San Francisco, 1971, effects of diet on, 26:45538
(HASL-257, pp I.51-70)

UNIVERSE

Einstein equations for spatially homogeneous, with
rotation, variations in action principle for, (T),
26:46758

mass of, repulsive interaction in universal gravitation
for, (T), 26:46749 (ORO-3992-94)

model formulation for rotating, with flat space, Einstein
field equations for homogeneous anisotropic, (T),
26:46261

model of, Einstein equation solutions with cosmological
term for homogeneous anisotropic, (T), 26:46759

URANIUM

analysis of, review on methods for isotopic, 26:45185
(TID-7029(2nd Ed.), pp 69-252)

chromatographic behavior on microcrystalline cellulose
impregnated with potassium chloride, 26:45278

complexes, electronic structure of insular uranyl,
26:45380

complexes, precipitation of peroxo, by using
hexamminecobalt(III) ion, 26:45370

compounds, ionization in mass spectrometer, 26:45258

content in Pechora River basin coal deposits,
distribution and genesis of, 26:45505

determination as uranyl in mine waters by extraction
photometry, 26:45242

determination by atomic absorption spectrometry,
sensitivities and detection limits for, 26:45251

determination in fuel cycle nuclear materials, sampling,
standards, and bulk measurements for, 26:45184 (TID-
7029(2nd Ed.), pp 3-65)

determination in fuel cycle product materials, review on
methods for, 26:45185 (TID-7029(2nd Ed.), pp 69-252)

determination in fuel reprocessing and radioactive waste
solutions, review on methods for, 26:45190 (TID-
7029(2nd Ed.), pp 331-411)

determination in fuels, book on methods for, 26:45183
(TID-7029(2nd Ed.))

determination in samples from water--uranium
compatibility test loop, radiochemical, 26:45181
(CISE-R-172)

determination in uranium ores by x-ray fluorescence,
preparation of pressed fused disc for, 26:45262 (NIM-
1427)

determination of trace amounts of, in germanium by
instrumental neutron activation, 26:45218

determination of, chemical procedures for, 26:45181
(CISE-R-172)

determination of, equipment and reagents for, 26:45191
(TID-7029(2nd Ed.), pp 413-33)

effects on spectrophotometric determination of phosphate
by extracting molybdophosphoric acid in butyl acetate
in presence of EDTA and fluoride, 26:45219 (ARH-SA-
110)

electron binding energy of, in metals and oxides,
photoelectron spectroscopy study of inner-shell,
26:46340 (ORNL-TM-3820)

energy from, relations with world balance of power,
26:46095

enrichment in Europe, plans for developing international
civilian gaseous diffusion plant for, 26:45693

enrichment in France, economic studies of gaseous
diffusion process for, 26:45691

enrichment of, Ad Hoc Committee report on efficiency of
processes for, 26:45689 (ORO-694)

ion exchange properties of uranyl, on ion exchangers
containing isonitrosoacetamide functional groups,
26:45283

radiation effects on deformation of fuel elements of,
relations of stress and temperature with in-pile,
26:46070 (UJV-2798-M)

radiation effects on grain growth in, model of effects
relations with statistical fluctuations in texture,
26:46069 (UJV-2795-M)

reactions between atoms of, potentials for calculation of
average energy going to electronic processes in,
26:46365(R) (ORO-2911-14)

recovery as by-product from uranium dioxide production,
method for, 26:45371(P)

reduction of hexavalent to tetravalent, in sulfuric acid
solutions, electrochemical, 26:45978

separation from spent oxide fuels by solvent extraction
using TBP, 26:45406(R) (ORNL-TM-2871)

spectra in chlorine- and oxygen-containing compounds,
chemical shifts of, 26:45377

spectra of, ascertaining valence states from x-ray
photoelectron, 26:45367

surface preparation prior to electroplating, method for,
26:45980(P)

URANIUM ALLOYS AND SYSTEMS

surface preparation prior to electroplating, method for,
26:45980(P)

Al--Fe--Mo--U-base, mechanical properties of, at 20 to
640°C, 26:45995 (UJV-2802-M)

Al--Fe--Si--U-base, mechanical properties of, at 20 to
640°C, 26:45995 (UJV-2802-M)

Al--Fe--U-base, mechanical properties of, at 20 to 640°C,
26:45995 (UJV-2802-M)

Al--Fe--U-base, mechanical properties of, at 20 to 640°C,
effects of chromium and silicon additions on, 26:45995
(UJV-2802-M)

Al--Mo--Si--U, mechanical properties of, at 20 to 640°C,
26:45995 (UJV-2802-M)

Al-base--U, fission fragment ranges in, 26:46074

Mo--U-base, mechanical properties of, at 20 to 640°C,
effects of aluminum, iron, and silicon additions on,
26:45995 (UJV-2802-M)

Pu--U, separation of plutonium from scrap, solvent
extraction flowsheet using TBP for, 26:45382 (ARH-SA-
130)

Th--U--Zr, phase studies on liquidus and solidus surfaces
in, 26:46003

U--UO₂, spectra of, effects of chemical state of uranium on, 26:45377

URANIUM ARSENIDES

UAs--US, magnetic structure of solid solutions, neutron-diffraction study of, 26:47261

URANIUM BROMIDES UBr_x

magnetic properties of, 26:47272

URANIUM CARBIDES

PuC--UC, dissolution of irradiated, in nitric acid, oxidation of organic compounds formed in, 26:45420(P)

PuC--UC, preparation by sol-gel process, 26:45406(R) (ORNL-TM-2871)

URANIUM CARBONITRIDES

preparation of, 26:45416(R) (ORNL-TM-3250)

PuCN--UCN, preparation, 26:45415(R) (ORNL-TM-3217)

PuCN--UCN, preparation of, 26:45409(R) (ORNL-TM-2996)

PuCN--UCN, preparation of, 26:45411(R) (ORNL-TM-3072)

PuCN--UCN, preparation studies on, 26:45413(R) (ORNL-TM-3127)

URANIUM CHLORIDES UCl₃

magnetic properties of, 26:47272

UCl₃--UO₂, spectra of, effects of chemical state of uranium on, 26:45377

URANIUM CHLORIDES UCl₄

magnetic properties of, 26:47272

Ca₂UCl₆--UCl₄, spectra of, effects of chemical state of uranium on, 26:45377

CaUCl₅--UCl₄, spectra of, effects of chemical state of uranium on, 26:45377

URANIUM DEPOSITS

exploration for Yugoslavian, status of, 26:45508

formation in rocks under oxidizing conditions, geochemistry of, 26:45504

mineralization in carbonate rocks, epigenetic zonality in, 26:45506

prospecting for Australian, aerial gamma spectrometric method for, 26:45507

URANIUM FLUORIDES UF₆

KF--UF₆, spectra of complex compounds in, ir, 26:45230

URANIUM FLUORIDES UF₄

analysis for volatile fluorides by ir spectrophotometry, 26:45238

determination in nitrogen by automatic absorption spectrometry, 26:45243(P)

production of, environmental studies on facility at Barnwell Nuclear Fuel Plant for, 26:45308 (DOCKET-701327-1)

URANIUM ISOTOPIES

determination in fuel cycle product materials, review on methods for, 26:45185 (TID-7029(2nd Ed.), pp 69-252)

determination in fuel reprocessing and radioactive waste solutions, review on methods for, 26:45190 (TID-7029(2nd Ed.), pp 331-411)

determination of ratios of, by mass spectroscopy, 26:45258

URANIUM ISOTOPIES U-231

energy levels from neptunium-231 electron-capture decay, (E), 26:47023 (KFK-1537)

energy-level transitions from neptunium-231 electron-capture decay, branching ratios for, (E), 26:47023 (KFK-1537)

URANIUM ISOTOPIES U-232

energy levels from neptunium-232 electron-capture decay, spins and parities of, (E), 26:47023 (KFK-1537)

energy-level transitions from neptunium-232 electron-capture decay, branching ratios for, (E), 26:47023 (KFK-1537)

URANIUM ISOTOPIES U-233

energy levels from neptunium-233 electron-capture decay, spins and parities of, (E), 26:47023 (KFK-1537)

energy-level transitions from neptunium-233 electron-capture decay, branching ratios for, (E), 26:47023 (KFK-1537)

neutron fission at 0.4 to 2000 eV using aligned target, angular distribution of fragments from, (E), 26:47026

neutron fission at 14 MeV, independent and cumulative yields of rare-gas isotopes from, (E), 26:47030

production of, spectrophotometric determination of phosphate in process solutions from, 26:45219 (ARH-SA-110)

shipping of, design of container for, 26:46116

URANIUM ISOTOPIES U-234

content in soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

determination relative to uranium-235 in uranyl nitrate by mass spectroscopy, 26:45258

energy levels from (p,t) and (t,p) reactions, calculation using monopole plus quadrupole pairing, 26:47042 (COO-3001-12)

energy-level scheme from plutonium-234 decay, (E), 26:47032

URANIUM ISOTOPIES U-235

content in ores, development of semiconductor probe for determining, 26:45658(P)

content in plants and soils in uraniferous area in New Zealand, 26:45529

content in soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

determination in power reactor fuels, delayed neutron interrogation method for, 26:45269

determination relative to uranium-234 in uranyl nitrate by mass spectroscopy, 26:45258

neutron capture-to-fission ratio at 8 to 60 keV, (E), 26:47024 (KFK-1563)

neutron fission at 14 MeV, independent and cumulative yields of rare-gas isotopes from, (E), 26:47030

neutron fission at 6 to 34 eV, probability of long-range α emission in, p. 288-9, (E), 26:46873(R) (CEA-N-1522)

neutron fission of, calculation of prompt gamma yields using improved renormalized Fermi gas model for thermal, (T), 26:47025

neutron fission of, gamma and neutron emission in, p. 314-8, (E), 26:46873(R) (CEA-N-1522)

neutron fission up to 3 keV, time delayed fragment spectra from, (E), 26:47018 (INDC-7-U, pp 61-4)

photo-fission of cross section for, (E), 26:46890

shipping of, design of container for, 26:46116

URANIUM ISOTOPIES U-236

fission of isomeric, half-life for, (E), 26:47018 (INDC-7-U, pp 61-4)

URANIUM ISOTOPIES U-238

content in plants and soils in uraniferous area in New Zealand, 26:45529

content in soils in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)

neutron fission at 14 MeV, independent and cumulative yields of rare-gas isotopes from, (E), 26:47030

neutron total cross section at 0.8 to 30 MeV, (E), 26:47013 (COO-3058-23)

photo-fission of cross section for, (E), 26:46890

proton fission at 16 and 42 MeV, angular distribution of long range alpha particles from, (E), 26:47037

transparency to photoproduced meson (π^+) and meson (π^-) at intermediate energies, (T), 26:46889

URANIUM NITRIDES

preparation of reactor fuels of, review on, 26:45960

properties for space reactor fuels, review on, 26:45960

testing of fuel pellets of, shock and vibration, 26:45933 (N-72-15609)

PuN--UN, preparation of reactor fuels of, review on, 26:45960

PuN--UN, properties for fast breeder reactor fuels, review on, 26:45960

URANIUM ORES

analysis for uranium by x-ray fluorescence, preparation of pressed fused disc for, 26:45262 (NIM-1427)

mining and milling of, on AEC-controlled lands in Colorado, New Mexico, and Utah, environmental impact report on, 26:45572 (WASH-1523)

processing of, use of plastic-coated tanks in solvent extraction method for, 26:46080

separation of protactinium from, by sorption on silica gel, 26:45390

URANIUM OXIDES

electron binding energies of uranium in, photoelectron-spectroscopy study of inner-shell, 26:46340 (ORNL-TM-3820)

preparation of sinterable, from uranyl nitrate solutions, equipment and method for, 26:45940(P)

URANIUM OXIDES UO₂

compaction of powdered, electrodynamic method for, 26:45951

crystal lattice expansion constant for powdered, during sintering, effects of titanium dioxide additions on, 26:45956

fission fragment ranges in, 26:46074

preparation by sol-gel process, 26:45410(R) (ORNL-TM-3018)

preparation by sol-gel process, 26:45408(R) (ORNL-TM-2949)

preparation from fuel reprocessing solutions, evaluation of methods for, 26:45409(R) (ORNL-TM-2996)

preparation from fuel reprocessing solutions, evaluation of processes for, 26:45408(R) (ORNL-TM-2949)

preparation from fuel reprocessing solutions, evaluation of methods for, 26:45411(R) (ORNL-TM-3072)

preparation from solutions from fuel reprocessing, 26:45417(R) (ORNL-TM-3281)

preparation from solutions from fuel reprocessing, 26:45413(R) (ORNL-TM-3127)

preparation from solutions from fuel reprocessing, 26:45415(R) (ORNL-TM-3217)

preparation from solutions from fuel reprocessing, 26:45414(R) (ORNL-TM-3180)

preparation from solutions from fuel reprocessing, 26:45416(R) (ORNL-TM-3250)

preparation from solutions from fuel reprocessing cycle, 26:45412(R) (ORNL-TM-3095)

preparation of microspheres of, 26:45412(R) (ORNL-TM-3095)

preparation of particulate fuels of, embedded in zirconium oxide matrix, 26:45957(P)

preparation of sols of, 26:45406(R) (ORNL-TM-2871)

preparation of sols of, CUSP process for, 26:45412(R) (ORNL-TM-3095)

preparation of sols of, CUSP process for, 26:45413(R) (ORNL-TM-3127)

preparation of sols of, CUSP process for, 26:45417(R) (ORNL-TM-3281)

preparation of sols of, CUSP process for, 26:45407(R)

(ORNL-TM-2918)
 preparation of sols of, CUSP process for, 26:45415(R)
 (ORNL-TM-3217)
 preparation of sols of, CUSP process for, 26:45409(R)
 (ORNL-TM-2996)
 preparation of sols of, CUSP process for, 26:45414(R)
 (ORNL-TM-3180)
 preparation of sols of, CUSP process for, 26:45411(R)
 (ORNL-TM-3072)
 preparation of sols of, CUSP process for, 26:45416(R)
 (ORNL-TM-3250)
 production of sinterable powders of, from fuel
 reprocessing solutions, 26:45407(R) (ORNL-TM-2918)
 production of, method for recovering uranium and other
 by-products from, 26:45371(P)
 reactions with carbon and nitrogen, 26:45417(R) (ORNL-
 TM-3281)
 reactions with carbon and nitrogen, 26:45408(R) (ORNL-
 TM-2949)
 reactions with gaseous bromine trifluoride, gravimetric
 study of, 26:45379
 spectra of, ascertaining valence states from x-ray
 photoelectron, 26:45367
 tritium distribution in Zircaloy-clad fuel rods of,
 26:45414(R) (ORNL-TM-3180)
 PuO₂--UO₂, analysis by γ spectroscopy, 26:45409(R)
 (ORNL-TM-2996)
 PuO₂--UO₂, analysis of pellets of, for moisture,
 26:45416(R) (ORNL-TM-3250)
 PuO₂--UO₂, diffusion of fission products in, 26:45417(R)
 (ORNL-TM-3281)
 PuO₂--UO₂, dissolution in nitric acid, 26:45415(R)
 (ORNL-TM-3217)
 PuO₂--UO₂, dissolution in nitric acid, 26:45413(R)
 (ORNL-TM-3127)
 PuO₂--UO₂, dissolution in nitric acid, 26:45411(R)
 (ORNL-TM-3072)
 PuO₂--UO₂, dissolution in nitric acid, 26:45409(R)
 (ORNL-TM-2996)
 PuO₂--UO₂, dissolution in nitric acid, 26:45412(R)
 (ORNL-TM-3095)
 PuO₂--UO₂, dissolution in nitric acid, 26:45416(R)
 (ORNL-TM-3250)
 PuO₂--UO₂, dissolution of irradiated, in nitric acid,
 26:45408(R) (ORNL-TM-2949)
 PuO₂--UO₂, fabrication of fuel rods of, 26:45408(R)
 (ORNL-TM-2949)
 PuO₂--UO₂, fabrication of pellets from sol-gel-derived,
 26:45406(R) (ORNL-TM-2871)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45410(R)
 (ORNL-TM-3018)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45412(R)
 (ORNL-TM-3095)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45413(R)
 (ORNL-TM-3127)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45415(R)
 (ORNL-TM-3217)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45417(R)
 (ORNL-TM-3281)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45407(R)
 (ORNL-TM-2918)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45409(R)
 (ORNL-TM-2996)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45414(R)
 (ORNL-TM-3180)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45408(R)
 (ORNL-TM-2949)
 PuO₂--UO₂, fabrication of sol-gel-derived, 26:45416(R)
 (ORNL-TM-3250)
 PuO₂--UO₂, fabrication of sol-gel-derived, Sphere-Pac
 process for, 26:45411(R) (ORNL-TM-3072)
 PuO₂--UO₂, oxidation of unirradiated pellets of,
 26:45416(R) (ORNL-TM-3250)
 PuO₂--UO₂, performance of fuels of, 26:45406(R) (ORNL-
 TM-2871)
 PuO₂--UO₂, preparation and properties of microspheres of,
 26:45411(R) (ORNL-TM-3072)
 PuO₂--UO₂, preparation by sol-gel process, 26:45410(R)
 (ORNL-TM-3018)
 PUO₂--UO₂, preparation by sol-gel process, 26:45417(R)
 (ORNL-TM-3281)
 PuO₂--UO₂, preparation by sol-gel process, 26:45413(R)
 (ORNL-TM-3127)
 PuO₂--UO₂, preparation by sol-gel process, 26:45414(R)
 (ORNL-TM-3180)
 PuO₂--UO₂, preparation by sol-gel process, 26:45411(R)
 (ORNL-TM-3072)
 PuO₂--UO₂, preparation by sol-gel process, 26:45407(R)
 (ORNL-TM-2918)
 PuO₂--UO₂, preparation by sol-gel process, 26:45408(R)
 (ORNL-TM-2949)
 PuO₂--UO₂, preparation by sol-gel process, 26:45416(R)
 (ORNL-TM-3250)
 PuO₂--UO₂, preparation of microspheres, 26:45416(R)
 (ORNL-TM-3250)
 PuO₂--UO₂, preparation of microspheres from sols of,
 26:45409(R) (ORNL-TM-2996)
 PuO₂--UO₂, preparation of microspheres of, 26:45410(R)
 (ORNL-TM-3018)
 PuO₂--UO₂, preparation of microspheres of, 26:45412(R)
 (ORNL-TM-3095)
 PuO₂--UO₂, preparation of microspheres of, 26:45415(R)
 (ORNL-TM-3217)

155 SUBJECTS

URINE

PuO₂--UO₂, preparation of microspheres of, 26:45417(R)
 (ORNL-TM-3281)
 PuO₂--UO₂, preparation of microspheres of, 26:45414(R)
 (ORNL-TM-3180)
 PuO₂--UO₂, preparation of microspheres of, 26:45408(R)
 (ORNL-TM-2949)
 PuO₂--UO₂, preparation of microspheres of, by CUSP-APEX
 mixed sol process, 26:45406(R) (ORNL-TM-2871)
 PuO₂--UO₂, preparation of sols of, 26:45407(R) (ORNL-TM-
 2918)
 PuO₂--UO₂, preparation of sols of, 26:45409(R) (ORNL-TM-
 2996)
 PuO₂--UO₂, radiation effects on sol-gel-derived,
 26:45415(R) (ORNL-TM-3217)
 PuO₂--UO₂, radiation effects on sol-gel-derived,
 26:45417(R) (ORNL-TM-3281)
 PuO₂--UO₂, radiation effects on sol-gel-derived,
 26:45414(R) (ORNL-TM-3180)
 PuO₂--UO₂, radiation effects on sol-gel-derived,
 26:45416(R) (ORNL-TM-3250)
 PuO₂--UO₂, radiation effects on sol-gel-derived, neutron,
 26:45413(R) (ORNL-TM-3127)
 PuO₂--UO₂, radiation effects on, neutron, 26:45412(R)
 (ORNL-TM-3095)
 PuO₂--UO₂, radiation effects on, neutron, 26:45410(R)
 (ORNL-TM-3018)
 PuO₂--UO₂, radiation effects on, neutron, 26:45411(R)
 (ORNL-TM-3072)
 PuO₂--UO₂, radiation effects on, neutron, 26:45409(R)
 (ORNL-TM-2996)
 PuO₂--UO₂, radiation effects on, neutron, 26:45408(R)
 (ORNL-TM-2949)
 PuO₂--UO₂, reprocessing of metallic-clad fuel elements
 of, electrochemical method of cutting and leaching in,
 26:45422(P)
 PuO₂--UO₂, solubility in nitric acid, 26:45417(R) (ORNL-
 TM-3281)
 PuO₂--UO₂-base, radiation effects on melting heat rating
 for, effects of porosity on fast neutron, 26:45930
 (HEDL-TME-72-81)
 rare earth oxides--UO₂, crystal structure of solid
 solutions in, 26:45414(R) (ORNL-TM-3180)
 ThO₂--UO₂, radiation effects on crystal lattice of fuels
 of, fission fragment, 26:45939
 U--UO₂, spectra of, effects of chemical state of uranium
 on, 26:45377
 URANIUM OXIDES UO₂
 production by fluidized-bed denitration of uranyl
 nitrate, 26:45405(R) (ICP-1006)
 production from uranyl nitrates, design of denitrator
 for, 26:45452
 spectra of, ascertaining valence states from x-ray
 photoelectron, 26:45367
 UCl₃--UO₃, spectra of, effects of chemical state of
 uranium on, 26:45377
 URANIUM OXIDES U₃O₈
 analysis of irradiated, for oxygen, development of remote
 apparatus for, 26:45224 (LA-DC-72-524)
 fission fragment ranges in, 26:46074
 reactions with gaseous bromine trifluoride, gravimetric
 study of, 26:45379
 spectra of, ascertaining valence states from x-ray
 photoelectron, 26:45367
 URANIUM SULFIDES
 UAs--US, magnetic structure of solid solutions, neutron-
 diffraction study of, 26:47261
 URANYL COMPOUNDS
 hydrolysis of, partition chromatography for studies on,
 26:45366 (JINR-P6-6402)
 URANYL FORMATES
 crystal structure of monohydrate of, x-ray diffraction
 studies on, 26:45368
 URANYL NITRATES
 analysis for uranium isotope ratios by mass spectroscopy,
 26:45258
 conversion to uranium trioxide, design of denitrator for,
 26:45452
 denitration of, fluidized-bed process for, 26:45405(R)
 (ICP-1006)
 vanadyl ion electron paramagnetic resonance in doped
 hexahydrate single crystal of, at room and liquid
 nitrogen temperatures, 26:45372
 UREA
 separation from cellular Y-3333 silicone rubber by
 aqueous extraction, 26:47603(R) (BDX-613-440(Rev.))
 UREA, 1,3-DIMETHYL-2-THIO-
 radiolysis of single crystals of, at 4.2°K, ESR spectral
 study of radical ion saturation in x, 26:45348
 UREA, HYDROXY-
 effects on DNA synthesis in uv-irradiated HeLa cells,
 26:45737
 UREA, THIO-
 effects on respiratory exchange in rats, 26:45823
 URIDINE, 2'-DEOXY-5-iodo-
 radiation effects on incorporation of, into DNA of mouse
 intestine, iodine-125 tracer study of gamma and proton,
 26:45842
 URINE
 monitoring for tritium, using acetylene from calcium
 carbide reactions, flowthrough ionization chamber for,

26:45585 (AERE-HP/M-74)
strontium-90 excretion in, of children in Poland,
relation between dietary intake of calcium and,
26:45908

UTAH

leasing of USAEC-controlled uranium-bearing lands in,
environmental impact report on, 26:45572 (WASH-1523)
plutonium isotope content in soils in, 1971, effects of
nuclear weapons testing on, 26:45526 (HASL-257, pp
I.95-118)
strontium-90 deposition and content in soils in Salt Lake
City, 1957 to 1971, effects of stratospheric fallout
on, 26:45525 (HASL-257, pp I.71-94)

V

VACUUM

bibliography on surface and vacuum physics, 26:46336

VAGINA

radioinduced rectovaginal fistula in human, following
combined external and intracavitary therapy of uterine
carcinoma, surgical repair of, 26:45800

VALVES

design of electromagnetic, for corrosive or radioactive
fluids, 26:46109(P)
development for high-rated conventional and nuclear steam
turbines, review on, 26:45455

VAN ALLEN RADIATION BELTS

alpha-particle distribution spectra in outer, June--July
1969, (E), 26:46309
electron pitch-angle diffusion within plasmasphere,
association with plasmaspheric whistler-mode waves,
(E/T), 26:46311
electron repopulation in inner, by radial diffusion
during and after Sept. 2, 1966 geomagnetic storm,
(E/T), 26:46310
proton flux in, identification of solar-neutron events
from increase above 20 MeV of, (E/T), 26:46320
whistler-mode wave emission between 3.9 and 10.4 kHz in
high-altitude radiation-belt slot, vlf, (E), 26:46312

VAN DE GRAAFF ACCELERATORS

control system for Cadarache, description of on-line
computer, 26:46123
design, operation, and performance of Hungarian 5-MeV,
26:46118 (FKFI-71-47)
operation of CEA tandem, Oct. 1, 1970-Oct. 15, 1971, p.
3-7, 26:46873(R) (CEA-N-1522)

VANADIUM

determination by spectrography, spectrochemical buffers
and volatilization mechanisms in, 26:45222 (JEN-229-
DQ/I-79)
determination of, chemical procedures for, 26:45181
(CISE-R-172)
determination of, in biological materials, comparison of
neutron activation and spectrophotometric methods for,
26:45203
electron-phonon mass enhancement factor from specific-
heat data, 26:47262
magnetic-flux-flow noise in superconducting, (E/T),
26:46815
oxidation by water vapor in air, 26:45971 (HMI-B-121, pp
VIII.1-37)
radioinduced void concentrations in foils of, laser-beam,
26:46059 (CONF-710601-, pp 613-29)
reactions with nitrogen at 900 to 1400°C and 0.07 to 700
torr, kinetics of, 26:45969
separation from other metals by combined ion exchange--
solvent extraction chromatography, 26:45282
vacancy defect production in quenched, mechanism of,
26:46025

VANADIUM ALLOYS AND SYSTEMS

Al-base--V, superconductivity and localized spin
fluctuations in, (E/T), 26:46838
H--V, structure of, 26:47263

VANADIUM CARBIDES

order--disorder transitions in, at 1152 to 1250°C,
thermodynamic studies of, 26:45937

VANADIUM HYDRIDES

ordering in, 26:47263

VANADIUM ISOTOPES V-51

energy-level transitions in, internal conversion
coefficients for, (T), 26:46921
neutron inelastic scattering at 0.6 to 3.4 MeV, γ spectra
and cross section for, (E), 26:46909
neutron reactions (n,α) with, calculation of excitation
functions for fast, 26:47066
neutron reactions (n,p) with, calculation of excitation
functions for fast, 26:47066
photon reactions (γ,n) and (γ,p) with carbon-12 up to 35
MeV, dipole resonance structure in cross sections for,
(E), 26:46890

VANADIUM OXIDES

energy level diagrams for, spectroscopic studies on high-
temperature, 26:45932 (LBL-450)
thermodynamic properties of, at high temperatures,
26:45932 (LBL-450)

VANADYL IONS

electron paramagnetic resonance in doped uranyl nitrate
hexahydrate single crystals at room and liquid nitrogen

temperatures, 26:45372

VEGETABLES

potatoes, radiation processing effects on sprout
inhibition, storage, and wholesomeness of, comparison
with chemical methods, 26:45716 (BFL-2/1972)

VENEZUELA

canal construction in, feasibility study of nuclear
excavation method for, 26:45475 (NP-19361)

VENTILATION

system at LLL, fire protection for, pp.16,17,
26:45673(R) (UCRL-50007-71-3)

VERMONT YANKEE POWER REACTOR

effluents from, effects on environment of chemical,
radioactive, and thermal, 26:47435 (DOCKET-50271-88)
effluents from, technical specifications for chemical and
thermal, 26:47437 (DOCKET-50271-92)
operation of, cost-benefit analysis for, 26:47435
(DOCKET-50271-88)
owners of, financial statements of, 26:47374 (DOCKET-
50271-90)
protection system for, revised scram bypass conditions
for, 26:47439 (DOCKET-50271-94)
radioactive waste processing system for, revised
technical specifications for off gas, 26:47440
(DOCKET-50271-95)
site for, ecology, geology, hydrology, and meteorology
of, 26:47435 (DOCKET-50271-88)
steam tunnel for, modifications to ventilation system of,
26:47438 (DOCKET-50271-93)
tests for, proposed 1 percent low power license,
26:47436 (DOCKET-50271-91)

VESSELS

sealing in hot cells, design of remote controlled
equipment for, 26:46112(P)

VIRGIL C. SUMMER POWER REACTOR, UNIT 1

safety of, additional data on, 26:47554 (DOCKET-50395-
24)
safety of, revised and additional data on, 26:47555
(DOCKET-50395-25)
safety of, revised and requested additional data on,
26:47553 (DOCKET-50395-22)

VIRUSES

Friend, effects on protein synthesis in mouse spleen cell
nuclei in comparison to effects of whole-body x-
irradiation, 26:45849
Gross murine leukemia, effects on lymphomagenic
properties of irradiated neonatal mouse thymus gland
transplanted into infected hosts, 26:45818 (UCLA-12-
867)

Venezuelan equine encephalitis, radioinduced inactivation
of purified, immunogenicity following γ , 26:45752

VITAMIN E GROUP

effects of dietary, on rate of lipid peroxidation in
tissues of whole-body x-irradiated mice, 26:45872

VOLTMETERS

description for measuring average pulse amplitude for
measuring charge collection efficiency in semiconductor
detectors, 26:45685

W

WATER

analysis for anionic surfactants using iron-59,
radiometric, 26:45272
analysis for uranium as uranyl by extraction photometry,
26:45242
chemistry in reactors, review on, 26:45310 (BARC-595)
corrosive effects of oxygenated, on Incoloy 800, Inconel
600 and 625, and stainless steel Types 304, 304L, 309,
316, and 347 at 288°C, 26:45966 (BMI-1927)
corrosive effects on germanium silicides, effects of
temperature on, 26:45935 (SC-RR-710884)
mass flow choking limit for heat pipe working fluid of,
(E), 26:45460
measurement of droplets of, in effluents from cooling
towers and process stacks, description of method for,
26:45574 (CONF-720619-1)
neutron attenuation by, (E/T), 26:46852
positron lifetime in, effects of temperature and
positronium trapping on, (E), 26:46362 (RISO-254)
reactions with molten metals, mechanisms of explosive,
26:45461
separation from steam, performance of steam separators
for, 26:45442
separation of entrained particles of, from air, survey
and testing of equipment for, 26:45440 (MSAR-71-45)
H₂O--steam, density, energy, and pressure in boiler
circuits, computer program for analysis of, 26:45458
(CISE-R-245)

WATER-d₂

determination in flowing water by measurement of
photoneutrons from high-energy photon bombardment,
26:45267(P)
positron lifetime in, effects of temperature and
positronium trapping on, (E), 26:46362 (RISO-254)
production at Bruce Heavy Water Plant, description of
auxiliary steam generator for, 26:45695
production by hydrogen isotopic exchange, 26:45696(P)
production of, methods for, 26:45698
properties and uses of, review on, 26:45698

WATER SUPPLIES

cesium-137 content in New York City, 1965 through 1971,

26:45512(R) (HASL-257(App.))
radioactivity monitoring of California, during 1970,
gross beta, 26:45537
strontium-90 content in New York City, Aug. 1954 through
March 1972, 26:45512(R) (HASL-257(App.))

WATER VAPOR
microwave emission from celestial sources of, (E/T),
26:46206
reactions with protactinium, x-ray powder diffraction
studies on compounds produced by, 26:45992 (LBL-659)
reactions with vanadium in air, oxidation, 26:45971
(HMI-B-121, pp VIII.1-37)

WATER WAVES
height of underwater cratering explosion-generated,
method for estimating, 26:45486 (UCRL-51235)

WATERFORD POWER REACTOR, UNIT 3
construction of, request for exemptions prior to permit
for, 26:47541 (DOCKET-50382-29)
environmental report for, requested additional data for,
26:47539 (DOCKET-50382-27)
environmental report for, revised and requested data for,
26:47543 (DOCKET-50382-31)
radioactive effluents from, data for source term
calculations for, 26:47542 (DOCKET-50382-30)
safety of, requested additional data on, 26:47538
(DOCKET-50382-25)
safety of, revised and additional data on, 26:47540
(DOCKET-50382-28)
safety of, revised and requested data on, 26:47544
(DOCKET-50382-32)

WATTS BAR POWER REACTOR, UNIT 1
construction permit for, additional data for application
for, 26:47552 (DOCKET-50390-19)
safety of, revised and additional data on, 26:47551
(DOCKET-50390-18)
site for, maximum possible flood conditions for,
26:47550 (DOCKET-50390-17)

WATTS BAR POWER REACTOR, UNIT 2
construction permit for, additional data for application
for, 26:47552 (DOCKET-50390-19)
safety of, revised and additional data on, 26:47551
(DOCKET-50390-18)
site for, maximum possible flood conditions for,
26:47550 (DOCKET-50390-17)

WEAR
measurement of machine parts, by activation using
accelerated charged particles or neutrons, 26:45711
measurements in machine parts, description of two-channel
automatic radiometric unit for, 26:45657

WEIGHT
measurement of fuel reprocessing plant input, development
of instruments for, 26:46093 (ICP-1014)

WEISWEILER POWER REACTOR
fuels for, development of production plant for, 26:45974

WELDED JOINTS
inspection in large-capacity boilers, radiometric methods
for, 26:45473

WELDING
equipment and methods for remote encapsulation of
radioactive waste, 26:46099 (ARH-SA-117)
equipment for radioactive waste cannister, description of
remote controlled, 26:46108 (UCRL-73757)
flux for submerged arc, composition of, 26:45941(P)

WELL LOGGING
equipment and instruments for, review of Hungarian,
26:45560
equipment for gamma, development of high-resolution
semiconductor, 26:45658(P)
method for elemental composition of formations by neutron
activation, 26:45561(P)
method for hydrocarbon prospecting, pulsed neutron,
26:45564
method for neutron, incorporating compensation for
borehole diameter and fluid density, 26:45562(P)
method for pulsed neutron, control and transmission
system for, 26:45563(P)
method for pulsed neutron, for aluminum and magnesium
prospecting, 26:45559(P)

WEST VALLEY PROCESSING PLANT
criticality in intercycle evaporators and plutonium
extraction system in, 26:45401 (DOCKET-50201-128)
equipment and materials technical specifications
revisions for, 26:45399 (DOCKET-50201-126)
evaluation for handling LMFBR fuels, 26:45407(R) (ORNL-
TM-2918)
radiation protection at, revisions of technical
specifications for, 26:45399 (DOCKET-50201-126)
radioactive waste management at, revisions of technical
specifications for, 26:45399 (DOCKET-50201-126)
radioactive waste processing at, low-level, 26:45400
(DOCKET-50201-127)
radioactivity releases at, April-June, 1972, 26:45398
(DOCKET-50201-125)
surveillance tests, April-June, 1972, 26:45398 (DOCKET-
50201-125)

WHISTLERS
growth of side-band, due to nonlinear wave--particle
interaction, (E/T), 26:46315
propagation in cold plasma with dilute energetic electron
density, stability of oblique, (E/T), 26:46314
wave emissions between 3.9 and 10.4 kHz in high-altitude
radiation-belt slot, vlf, (E), 26:46312

WHOLE-BODY COUNTERS

background spectra in, measured, 26:45590(T) (ORNL-tr-
2532)
shielding performance for reduction of background
radiation in, 26:45590(T) (ORNL-tr-2532)

WIND

velocity of, use of radon distribution for calculating
turbulent diffusion coefficient and vertical component
of, 26:45522

WIRES

losses in assembly of superconducting, transposed in
conducting matrix and subjected to pulsed magnetic
fields, 26:46810
thickness of, radiometric gage for measuring, 26:45717

WOOD

radiation effects on anatomical structure and chemical
composition of, in spruce seedlings, γ , 26:45774
radioinduced polymerization of plastic-impregnated,
materials and methods for, 26:45718
radioinduced polymerization of trialkylacryloxystannanes
in, γ , 26:45355(P)
x-ray transmission through lawanit and pine, low-energy,
(E), 26:47205

WYOMING

site selection for Wagon Wheel Event in, preliminary
evaluation of, 26:45571 (NVO-55)

X**X RADIATION**

attenuation in matter, method for determining mean energy
of secondary electrons from, 26:47213
cosmic background, surface brightness caused by
clustering or superclustering of sources, (E/T),
26:46260
dose distribution in man during therapy of Hodgkin's
disease, role of clinical state of patient in planning,
26:45786(T) (TT-54002/5, pp 590-3)
dose to man from diagnostic radiography, method and units
for expressing cumulative, 26:45799
dosimetry at 10 to 500 keV, characteristics of personal
photodosimeters for, 26:45629
dosimetry in water phantoms using thermoluminescent
dosimeters, 26:45622
dosimetry of pulsed, at 10 to 35 MV, performance of
ionization chambers for, 26:45627
dosimetry of, design of tissue-equivalent liquid
ionization chamber for, 26:45621
dosimetry of, plastic tissue-equivalent scintillation
detectors for, 26:45619
effectiveness for therapy of laryngeal carcinomas,
effects of clinical stage and histologic type of tumor
on, 26:45780(T) (TT-54002/4, pp 430-6)
effects of pulsed, on electric conductivity of polymers,
26:47225 (AWRE-G-31/72)
effects on activity of glucose-6-phosphatase in
erythrocytes of rabbits and rats following whole-body
exposure, effects of cysteine and reduced glutathione
on extent of recovery from inhibitory, 26:45848
effects on activity of glutathione peroxidase in blood
and liver of swine, 26:45733
effects on aqueous solutions of nickel
ethylenedinitrilotetraacetate, 26:45350
effects on AT-cut quartz radiofrequency oscillators,
rapid annealing of, 26:45665 (SC-DC-721240)
effects on blood plasma in vitro, effects on formation of
lipid peroxides, effects of water on, 26:45725
effects on bone marrow cells of rats, effects on colony-
forming ability following transplantation into mice,
low dose, 26:45834
effects on brain of rabbits, effects on enzyme activity
in neuroglia of developing, 26:45836
effects on chick mesonephros in vitro, effects on phenol
red uptake by, autoradiography study of, 26:45832
effects on chromosomes in cultured Chinese hamster ovary
cells, effects of post-exposure temperature on repair
of aberrant, 26:45743
effects on chromosomes in male *Drosophila* in relation to
suppression of crossing over in chromosomes of F_1
females, aberrant, 26:45876
effects on chromosomes of *Drosophila*, method for
quantitative study of autosomal non-disjunction,
26:45880
effects on content of 5-hydroxyindole acetic acid, volume
of urine excreted, and water consumption of mice
following partial or whole-body exposure, 26:45826
effects on count of mast cells in blood of man following
mediastinal exposure, 26:45810
effects on cultured human fibroblasts, effects of dose on
frequency of mutagenic, 26:45740
effects on DNA complex in cultured Chinese hamster cells,
dose dependence of repair processes following,
26:45745
effects on dogs, effects on skin following fractionated
or single partial body exposure, effects of dose and
fractionation interval on acute and chronic
pathological, 26:45860
effects on *Drosophila*, effects on genetic recombination
in relation to tergite development, 26:45881

effects on *Escherichia coli*, protective effects of chelating compounds against, 26:45859

effects on eyes of rabbits, effects on cellular proliferation and regeneration during injury response and wound healing of lens, 26:45813(R) (COO-1427-41)

effects on fatty acids of blood plasma in vitro, effects of water on, 26:45725

effects on films of dielectrics, induced photoconductivity, (E/T), 26:45668 (UCRL-73974)

effects on hamsters, effects on ovarian function in pregnant, 26:45825

effects on hematopoietic stem cells in mice, effects of tissue extracts on recovery of, 26:45847

effects on human laryngeal mucosa following therapy of laryngeal carcinomas, incidence of pathological, 26:45781(T) (TT-54002/4, pp 437-41)

effects on human spinal cord following therapy of neoplasms, clinical course of late pathological, 26:45797

effects on immune reactions of rats to allogeneic tumor transplants following whole-body exposure, additive effects of cortisone on suppressive, 26:45820 (UR-3490-121)

effects on incidence of leukemia in man following diagnostic radiography, effects of age, exposure site, number of exposures, and sex on, 26:45798

effects on lungs of rabbits and rats, role of allergic inflammations in pathological, 26:45863

effects on lysosomes in rat liver cells following lethal whole-body exposure, 26:45839

effects on male germ cells of *Drosophila*, relation to frequency of chromosome loss, enhancing effects of dose fractionation and lucanthone on, 26:45887

effects on malonic acid single crystals, deuterium isotope effects on yields of acyl radical from, 26:45343

effects on man following exposure of pelvis, survey of neoplastic and pathological, 26:45807

effects on man, effects on small intestine following therapy of uterine carcinomas, symptoms and treatment of pathological, 26:45802

effects on MAS diodes, 26:45670

effects on mice following acute pulsed exposure, comparison of RBE with cobalt-60 γ rays for lethal, 26:45866

effects on mice, effects on growth of thyroid gland in goitrogen-stimulated adult and normal fetal, 26:45865

effects on mice, effects on lipid peroxidation in tissues following whole-body exposure, effects of dietary antioxidants on dose and time dependent, 26:45872

effects on mice, effects on regeneration of bone marrow lymphoid cells in, effects of thymectomy on, 26:45830

effects on mice, effects on regeneration of lymphoid cells in bone marrow of germ-free, 26:45843

effects on mice, effects on stem cell renewal in bone marrow and spleen following fractionated whole-body exposure, 26:45871

effects on mice, in vitro culture of leukemic cells induced by, 26:45855

effects on mice, protective effects of chelating compounds against, 26:45859

effects on monkeys, effects on central nervous system following local exposure, pathological, 26:45850

effects on monkeys, effects on visual cortex in brain of, latent pathological, 26:45858

effects on nucleotide distribution in DNA of rat bone marrow in comparison to leukosis effects, 26:45856

effects on phagocytic activity of macrophages co-cultured with tumor cells, 26:45738

effects on potassium ion transport in rat thymocytes following exposure in vitro, inhibitory, 26:45744

effects on protein synthesis in mouse spleen cell nuclei following whole-body exposure, comparison of effects of Friend virus infection with, 26:45849

effects on rabbits, effects on crystalline lens in comparison to effects of fast neutron, γ , and mixed γ and neutron radiations, 26:45864

effects on rat bone marrow following lethal whole-body exposure, role of transplanted slowly proliferating bone marrow cells in regeneration, 26:45837

effects on rat intestine following abdominal exposure, effectiveness of cholestyramine resin for prevention and treatment of cholestatic enteropathy resulting from pathological, 26:45821

effects on rats, lymphoid cell proliferation kinetics in thymus during development of leukemogenic, 26:45822

effects on response of mononuclear phagocytes to thioglycollate-induced intraperitoneal inflammation, effects of time of exposure on, 26:45833

effects on ruby and corundum crystals, 26:47244

effects on single crystals of benzoyl disulfide and 1,3-dimethyl-2-thiourea, ESR spectral study of radical ion saturation in, 26:45348

effects on sperm of wasps in relation to fecundity and life span of F_1 females, mutagenic, 26:45878

effects on survival of cultured mammalian cells, relation of mathematical analysis of, to radiotherapy, 26:45735

effects on survival of man following therapy of metastases to neck lymph nodes from tumors of larynx, nasopharynx, and tonsils, comparison of, 26:45782(T)

(TT-54002/4, pp 442-50)

effects on survival of rats, protective effects of DEAE Sephadex against, 26:45844

effects on survival of sea urchin eggs, effects of cell phase and thiol content on, 26:45882

effects on thyroid gland of dogs, combined effects of internal iodine-131 on neoplastic, 26:45819 (UR-3490-63)

effects on tongue of rabbits, permeability of lingual mucosa during development and recovery from pathological, 26:45862

effects on trematode metacercariae, effects on infectivity and development in chick and chorioallantois, dose dependence of, 26:45879

effects on uv fluorescence of Ehrlich ascites cells, 26:45736

luminosity of clusters of galaxies, (T), 26:46249

scattering by helium isoelectronic sequence, analytic calculation of form factors for coherent, 26:46359

spectrometry using free-air ionization chamber, 26:47218

transmission through materials locally available in Philippines, (E), 26:47205

use for therapy of breast carcinomas in man, correlation between rate of growth and size of tumor and survival following, 26:45783(T) (TT-54002/4, pp 451-5)

use in production of organic aerosols, 26:46082

X-RAY EQUIPMENT

radiation exposure control system for dental, ionization chamber, 26:45603

X-RAY FLUORESCENCE ANALYSIS

description for determination of small amounts of transuranium element using α - x coincidence technique, 26:45270

radioisotope sources in nondispersive, for forensic investigation, 26:45213

X-RAY SOURCES

development of tube, for fluorescence analysis of trace elements, pp-2,3, 26:45673(R) (UCRL-50007-71-3)

X-RAY SPECTROMETERS

resolution of cooled Si(Li), 26:45644

X-RAY SPECTROSCOPY

use in determination of physical-chemical properties of matter, expanded, 26:47618 (JINR-P14-6407)

XENON

effects on quenching of excited states from pulse radiolysis of naphthalene and pyrene in cyclohexane, 26:45331

effects on yields of singlet state from pulse radiolysis of naphthalene and pyrene in benzene and cyclohexane solutions, 26:45329

electron reactions with clusters of, multiple ionization in, (E), 26:46386

XENON IONS

implantation in iridium at keV energies, field-ion microscopic study of damage profiles produced in, 26:47240

XENON ISOTOPES

meson (π) reactions at 2.34 to 9 GeV/c, π^0 and resonance production in, (E), 26:46476 (JINR-P1-6408)

XENON ISOTOPES Xe-133

diffusion of fission-product, from graphite to 1400°C, 26:46078 (NYO-1710-106)

Y

YANKEE POWER REACTOR

fuels for, effective cross section ratios for, (E/T), 26:47595

operations during June 1972, 26:47341 (DOCKET-50029-189)

technical specifications for, withdrawal of proposed change for, 26:47391 (DOCKET-50029-190)

YEASTS

effects of extracts of, on recovery of γ -irradiated carnation cuttings, 26:45777

radiosensitivity of, factors affecting, 26:45749

YTTERBIUM

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251

YTTERBIUM BOROCARBIDES

preparation, properties, and structure of, 26:45964

YTTRIUM

determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251

electron-phonon mass enhancement factor from specific-heat data, 26:47262

preparation of high purity monocrystals of, by zone melting, electron transfer during, 26:45975

separation of gallium from, by solvent extraction using methyl isobutyl ketone, 26:45285

YTTRIUM ALLOYS AND SYSTEMS

Al-Fe-Y, phase studies on, 26:46005

YTTRIUM FLUORIDES

CaF₂-YF₃, thermodynamics of solid electrolytes of, review on, 26:46013

YTTRIUM HYDRIDES

elastic properties of YH_{1.93}, from acoustic velocity measurements, 26:45950

YTTRIUM ISOTOPES Y-89

photon reactions (γ,n) and (γ,p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

beta decay of, energy-dependent K-shell internal ionization probability in, (T), 26:46962
effects on gastrointestinal tract following uptake of, review of carcinogenic, 26:45895 (UR-3490-102)

YTTRIUM ISOTOPES Y-91

range of fission fragments of, in aluminum--uranium alloys and uranium oxides, 26:46074

YTTRIUM OXIDES

analysis for terbium by luminescent method using pyrogallolsulfonic acid, 26:45231
cathodoluminescence efficiency of doped, absolute, (E), 26:45591
properties of plasma-sprayed, on multilayer niobium--zirconium substrate insulators, 26:45702(R) (LA-4978)
radiation effects on elasticity and swelling of, fast neutron, 26:45702(R) (LA-4978)
 Al_2O_3 -- Y_2O_3 , density, surface tension, and volume change on melting of, x-radiographic studies on, 26:45949
 ThO_2 -- Y_2O_3 , transport properties of, at 500 to 1600°C, effects of oxygen pressure on, 26:45938
 Y_2O_3 -- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

YTTRIUM OXSULFIDES

cathodoluminescence efficiency of doped, absolute, (E), 26:45591

YTTRIUM PHOSPHATES

cathodoluminescence efficiency of doped, absolute, (E), 26:45591

YTTRIUM VANADATES

cathodoluminescence efficiency of doped, absolute, (E), 26:45591

YUGOSLAVIA

uranium deposits in, exploration for, 26:45508

Z

ZINC

chemical behavior in Chesapeake Bay, 26:47613 (COO-3292-1)
determination in hair by neutron activation and γ spectroscopy, 26:45214
determination in high-purity cadmium oxide, spectrographic, 26:45220 (BARC-593)
determination of trace amounts of, in environmental samples by neutron activation, 26:45198
determination of trace amounts of, in germanium by instrumental neutron activation, 26:45218
effects on recrystallization of iron, 26:45982
positron annihilation radiation in, correlation of Fermi energy and effective electron mass with Doppler-broadened, 26:47268
reactions of liquid, with iodine, 26:45421
separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281
surface electron band structure of single crystals of, magnetically induced, 26:47256

ZINC ALLOYS AND SYSTEMS

La--Zn, phase studies and thermodynamic properties of, 26:46001
Mg--Zn, grain refining of, mechanism and method for, 26:45984

ZINC IODIDES

reactions with molten chlorides, 26:45421

ZINC IONS

production and acceleration of multicharged, development of ion source for, 26:46133

ZINC ISOTOPES Zn-64

neutron inelastic scattering up to 1.7 MeV, γ spectra and excitation function for, 26:46936

ZINC ISOTOPES Zn-65

content in biota, sediments, and water in Lake Michigan near Big Rock Point power reactor during 1970, 26:45536
content in marine mussels in Columbia River following shutdown of Hanford reactors, 26:45543 (RLO-2225-T-1-9)

ZINC ISOTOPES Zn-66

neutron inelastic scattering up to 1.7 MeV, γ spectra and excitation function for, 26:46936

ZINC ISOTOPES Zn-68

helium-3 reactions (He-3,d) at 30.2 MeV, differential cross sections and spectrum from, p. 112-4, (E), 26:46873(R) (CEA-N-1522)
neutron capture by, γ spectra from thermal, (E), 26:46937

neutron inelastic scattering up to 1.7 MeV, γ spectra and excitation function for, 26:46936

ZINC ISOTOPES Zn-69

energy levels from zinc-68 neutron capture, (E), 26:46937

ZINC ISOTOPES Zn-72

energy levels and properties of, compilation of, (E), 26:46939

ZINC SILICATES

cathodoluminescence efficiency of doped, absolute, (E), 26:45591

ZINC SULFIDES

cathodoluminescence efficiency of doped, absolute, (E), 26:45591

ZION POWER REACTOR, UNIT 1

components for, effects of warehouse fire on

acceptability of, 26:47463 (DOCKET-50295-58)
construction of, additional data on welding procedures for, 26:47462 (DOCKET-50295-57)
electric cables for, additional data on installation and inspection of, 26:47464 (DOCKET-50295-59)
instrumentation for, additional data on installation and inspection of, 26:47464 (DOCKET-50295-59)
personnel for, qualifications for nondestructive testing, 26:47462 (DOCKET-50295-57)
steam line isolation valve for, quality assurance requirements for main, 26:47462 (DOCKET-50295-57)
ZION POWER REACTOR, UNIT 2
components for, effects of warehouse fire on acceptability of, 26:47463 (DOCKET-50295-58)
construction of, additional data on welding procedures for, 26:47462 (DOCKET-50295-57)
control rod drives for, leaks in canopies of, 26:47379 (DOCKET-50304-59)
electric cables for, additional data on installation and inspection of, 26:47464 (DOCKET-50295-59)
instrumentation for, additional data on installation and inspection of, 26:47464 (DOCKET-50295-59)
personnel for, qualifications for nondestructive testing, 26:47462 (DOCKET-50295-57)
steam line isolation valve for, quality assurance requirements for main, 26:47462 (DOCKET-50295-57)

ZIRCALOY

analysis for carbon and oxygen by deuteron and proton activation, 26:45309(R) (AE-453)
analysis of, emission spectrographic methods for, 26:45181 (CISE-R-172)

ZIRCONATES

production and uses of, review on, 26:45979

ZIRCONIUM

carbon-ion reactions at 10 MeV/amu with, energy shift and relative intensity of K x rays from, (E), 26:46339
chromatographic behavior on microcrystalline cellulose impregnated with potassium chloride, 26:45278
determination by atomic absorption spectrometry, sensitivities and detection limits for, 26:45251
determination by electron probe microanalysis methods, 26:45237
determination in geological materials using picramine-epsilon, photometric, 26:45233
effects on metallography of niobium--oxygen systems, electron microscopy studies on, 26:46039
effects on recrystallization of iron, 26:45982
evaporation from open surfaces at high temperatures, design of equipment for measuring, 26:46011
ion reactions at 5 MeV/amu with, dependence of energy shift and relative intensity of K x rays from, on ion stopping power, (E), 26:46339
mechanical properties of α -, effects of strain rate and temperature on, 26:46040
nitridation at 1440°C, kinetics of, 26:46030
production and uses of, review on, 26:45979
separation from other metals as β -diketonates by column liquid--liquid chromatography, 26:45281
separation from other metals by combined ion exchange--solvent extraction chromatography, 26:45282
separation from solution by precipitation as tetrafluoride, 26:45274(P)
separation of protactinium from, in chloride and sulfate media by sorption on silica gel, 26:45390

ZIRCONIUM ALLOYS AND SYSTEMS

analysis for tin by alternating current polarographic method, 26:45256
production and uses of, review on, 26:45979
Mg--Zr, grain refining of, mechanism and method for, 26:45984
Nb--Zr, phase transformation and structure of solid solutions of, mechanism of β + ω , 26:46036
Th--U--Zr, phase studies on liquidus and solidus surfaces in, 26:46003
Ti--Zr, diffusion in β -phase of, mechanism of, 26:45985 (BARC-577)

ZIRCONIUM BORIDES

production and uses of, review on, 26:45979

ZIRCONIUM CARBIDES

production and uses of, review on, 26:45979

ZIRCONIUM FLUORIDES

preparation of anhydrous, 26:45274(P)

ZIRCONIUM HYDRIDES

phase studies of, 26:45942
plasma of, ion energy spectra from laser-produced, (E), 26:47154

production and uses of, review on, 26:45979

use as hydrogen source in neutron columns, 26:45942

ZIRCONIUM ISOTOPES Zr-90

deuteron reactions (d,n) at 12 MeV, DWBA analysis of angular distributions for, (E), 26:46956
electron elastic scattering at 302 MeV, differential cross sections for, p. 186-8, (E), 26:46873(R) (CEA-N-1522)
electron inelastic scattering at 209 MeV, differential cross sections for, p. 189-91, (E), 26:46873(R) (CEA-N-1522)
helium-3 reactions (He-3,t) at 21.4 to 37.5 MeV with isobaric analog excitation, model analysis of, (E/T), 26:46930

nuclear charge distribution from electron elastic scattering, p. 186-8, (E), 26:46873(R) (CEA-N-1522)
photon reactions (γ, n) and (γ, p) with carbon-12 up to 35 MeV, dipole resonance structure in cross sections for, (E), 26:46890

ZIRCONIUM ISOTOPES Zr-91

meson reactions (π^+, π^0) at 30 MeV, cross section for, p. 224-5, (E), 26:46873(R) (CEA-N-1522)
proton reactions (p, n) at 13 to 40 MeV with subsequent proton decay, coupled-channels calculations of energy dependence of, (T), 26:47073

ZIRCONIUM ISOTOPES Zr-95

content in atmosphere, soils, and vegetation in vicinity of Lawrence Livermore Laboratory, Jan. through Dec. 1971, 26:45513 (UCRL-51242)
content in fallout deposited on Dallas, Texas soil during Jan. to July 1964, (E), 26:45527 (TID-26103)
content in surface atmosphere at world sites, 1963 through Feb. 1972, 26:45512(R) (HASL-257(App.))
content stratosphere to July 1971, effects of nuclear weapons testing on, 26:45518 (HASL-257, pp I.33-50)
range of fission fragments of, in aluminum--uranium alloys and uranium oxides, 26:46074

ZIRCONIUM ISOTOPES Zr-96

deuteron reactions (d, n) at 12 MeV, DWBA analysis of angular distributions for, (E), 26:46956

ZIRCONIUM NITRIDES

production and uses of, review on, 26:45979

ZIRCONIUM OXIDES

electron spin resonance at 35 GHz of single crystals of yttria-stabilized, effects of blackening by electric current on, measurements at 293 to 77°K on, 26:47252
phase transformation in, differential thermal analysis studies on burst phenomenon and kinetics of, 26:45947
preparation of framed films of, for electrolytic membranes and radiation windows, 26:45958(P)
production and uses of, review on, 26:45979
properties for use as matrix for particulate uranium dioxide fuels, 26:45957(P)
 Al_2O_3 -- CaO -- Cr_2O_3 -- F -- MgO -- MoO_3 -- TiO_2 -- ZrO_2 , use as welding flux, 26:45941(P)

CaO -- ZrO_2 , phase studies on partially stabilized, 26:45945

CaO -- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

MgO -- ZrO_2 , phase studies on partially stabilized, 26:45945

MgO -- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

Y_2O_3 -- ZrO_2 , thermodynamics of solid electrolytes of, review on, 26:46013

ZIRCONIUM SELENIDES

production and uses of, review on, 26:45979

ZIRCONIUM SILICATES

production and uses of, review on, 26:45979

ZIRCONIUM SULFIDES

production and uses of, review on, 26:45979

ZIRCONIUM TELLURIDES

production and uses of, review on, 26:45979

REPORT NUMBER INDEX

Reports, conference papers and proceedings, numbered translations, and patents are included in this index. Each item is listed by report number; this is followed by the abstract number and a brief notation regarding public availability. The index entries show which reports are on file in the institutions listed on the back of the front cover (Dep.), which reports have been published in open-literature journals (journal citation), and which reports can be purchased (source and price). Whenever a report is identified by more than one number, the availability is given under the primary number, and a cross reference is given under the secondary report number.

The report number codes apply to USAEC reports unless designated otherwise. Non-USAEC reports are identified by parenthetical notes after the report codes; e.g., (non-AEC), (British), (Canadian), (French). The note (Misc.) identifies the codes used for both USAEC and non-USAEC reports. Patents are listed under the heading "PATENTS"; the country of origin is included in parentheses.

A comprehensive list of both USAEC and non-USAEC report codes is contained in the latest revision of TID-85, *Report Number Series Used by the USAEC Technical Information Center in Cataloging Reports*. This information is also included in the Cumulative Report Number Index (issued annually).

Availability of USAEC Reports

All USAEC Reports abstracted in *Nuclear Science Abstracts* are available through one or more of the following sources:

(1) University and public libraries in the United States. A list of these organizations appears on the inside front cover of each issue of *NSA*.

(2) NCR Microcard Editions in Microcard and micronegative form. Orders and inquiries should be sent to: Order Processing Department, 365 South Oak Street, West Salem, Wisconsin 54669.

(3) National Technical Information Service, U. S. Department of Commerce, Springfield, Virginia 22151.

(4) Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402.

(5) Microsurance, Inc., P. O. Box 3522, Oak Ridge, Tenn. 37830, in micronegative form. Purchase details available on request.

Availability of Non-USAEC Reports

U. S. Reports. Reports of U. S. Government agencies are available to the public from the National Technical Information Service (NTIS). If specific availability is not shown, inquiries may be directed to NTIS.

Non-U. S. Reports. Reports of atomic energy organizations of other countries are available from NTIS for U. S. sale only. Specific availability in the country of origin is shown, when known, at the report number entry. Otherwise, inquiries should be made to the organizations listed below:

AUSTRALIA

Australian Atomic Energy Commission Research Establishment. Available upon request from the Library, Australian Atomic Research Establishment, Private Mail Bag, Sutherland, New South Wales, Australia.

BELGIUM

Centre d'Étude de l'Energie Nucléaire. Available upon request to individuals and organizations from the Centre d'Étude de l'Energie Nucléaire (C.E.N.), Studiecentrum voor Kernenergie (S.C.K.), Technical Information Dept., MOL-DONK (Belgium).

CERN

CERN European Organization for Nuclear Research. Available without charge from CERN European Organization for Nuclear Research, Geneva 23, Switzerland. Copies are deposited in libraries throughout the world.

FRANCE

Commissariat à l'Energie Atomique. Inquiries for reports numbered 1-2199 should be addressed to Service Central de Documentation du C.E.A., Centre d'Etudes Nucléaires de Saclay, Boîte postale n° 2, 91. GIF-sur-YVETTE, FRANCE. Reports numbered 2200 and above are available from La Documentation Française, 31 quai Voltaire, 75, PARIS (7e), FRANCE.

Service Central de Protection contre les Rayonnements Ionisants. Copies are available from Ministère de la Santé Publique, Service Central de Protection contre les Rayonnements Ionisants, B.P. n° 78 Le Vesinet (France) Telex 25667.

GERMANY

Deutsches Elektronen-Synchrotron DESY. Copies are available from Deutsches Elektronen-Synchrotron DESY, Dokumentation, 2000 Hamburg 52, Notkestieg 1, Germany.

Gesellschaft für Kernforschung mbH (KFK). Copies are available from Gesellschaft für Kernforschung mbH, 75 Karlsruhe, Postfach 947, Germany. Copies are also deposited in many national nuclear centers in the world.

Kernforschungsanlage. Copies may be ordered from Zentralbibliothek der Kernforschungsanlage Jülich, Jülich, Bundesrepublik Deutschland.

Hahn-Meitner-Institute für Kernforschung. Available without charge from Hahn-Meitner-Institute für Kernforschung Ber-

lin, Bibliothek, Glienicke Strasse 100, 1 Berlin 39, Germany. Copies are deposited in many national nuclear centers in the world.

Johann-Wolfgang-Goethe Universität, Frankfurt Am Main, Institut für Kernphysik, (IKF). Copies are available from Gmelin Institut, 40-42 Varrentrappstrasse, Frankfurt (Main), West Germany.

ISRAEL

Israel Atomic Energy Commission. Available without charge from Israel Atomic Energy Commission, Soreq Nuclear Research Centre, Yavne, Israel.

ITALY

Centro Informazioni Studi Esperienze (CISE). For sale at CISE Press Office, Casella Postale 3986, Milano, Italy. Copies are deposited in the libraries of various nuclear centers in Italy and abroad.

Comitato Nazionale per l'Energia Nucleare. Copies are available from Comitato Nazionale per l'Energia Nucleare, Divisione Affari Internazionali e Studi Economici, Ufficio Pubblicazioni, Via Belisario, 15, Roma, Italy.

JAPAN

Japan Atomic Energy Research Institute (JAERI). Available from the Staff Supply Cooperative of Japan Atomic Energy Research Institute (JAERI), Tokai-mura, Naka-gun, Ibaraki-ken, Japan. Copies are deposited in depositories in Tokyo, Nagoya, and Osaka.

Nagoya University. Copies are available from Research Information Center, Institute of Plasma Physics, Nagoya University, Nagoya, Japan.

University of Tokyo. Available free of charge from Institute for Nuclear Study, University of Tokyo, Tanashi-Machi, Kitatama-gun, Tokyo, Japan.

NETHERLANDS

Reactor Centrum Nederland. For sale by Reactor Centrum Nederland, 112 Scheveningseweg, 's-Gravenhage, The Netherlands. Copies are deposited in university libraries in Holland and in many national nuclear centers in the world.

NEW ZEALAND

Institute of Nuclear Sciences. Copies are available from Librarian, N. Z. Institute of Nuclear Sciences, D.S.I.R., Private Bag, Lower Hutt, New Zealand. Copies are deposited in other countries.

PAKISTAN

Atomic Energy Center. Copies are available from Atomic Energy Center, P. O. Box 658, Lahore, (West Pakistan).

POLAND

Reports of the Polish nuclear institutes are published by the Nuclear Energy Information Center of the Government Commissioner for Use of Nuclear Energy. Copies are available free of charge from Nuclear Energy Information Center, Palace of Culture and Science, Warsaw, Poland.

SPAIN

Junta de Energia Nuclear. "JEN" reports are available without charge from Servicio de Documentación, Junta de Energia Nuclear, Ciudad Universitaria, Madrid (3), España.

SWITZERLAND

Eidgenössisches Institut für Reaktorforschung. "EIR" reports are available from Eidgenössisches Institut für Reaktorforschung, c/o Bibliothek, 5303 Würenlingen/Switzerland.

TURKEY

Turkish Atomic Energy Commission. Copies are available free of charge from T.A.E.C., Çekmece Nuclear Research Center, P. O. Box 1, Airport, Istanbul, Turkey.

Conferences

Published conference proceedings as well as individual unpublished papers from conferences are included in this index. A CONF number is assigned to the conference if published proceedings are received and also if only a single paper is received. The base number used is year-month-sequence. Thus, the number CONF-650101 indicates that the conference was held in 1965, in January, and that this was the first January 1965 conference so received. This number would be assigned to complete published proceedings. If individual papers are received, the base number is extended to include a paper number, e.g., CONF-650101-1. As additional papers are received, they will be assigned consecutive numbers in the series, CONF-650101-1, CONF-650101-2, etc. This numbering system ensures that all papers received from a conference will be associated with that conference in this index.

If published proceedings are received after the receipt of individual papers, the complete proceedings will be abstracted and the abstract number will appear in this index with the base CONF number.

Abbreviations Used in Availability Column

AECL	Reports of the Atomic Energy of Canada Limited are for sale by the Scientific Document Distribution Office, Atomic Energy of Canada Limited, Chalk River, Ontario, Canada.
CFSTI	See NTIS
DENM	Reports of the Atomic Energy Commission Research Establishment, Risö, are available on exchange from the Library, Danish Atomic Energy Commission, Risö, Roskilde, Denmark. Also for sale at the price shown from Jul. Gjellerup, Booksellers, 87 Sölvgade, København K., Denmark.
	Reports of Danatom are available on loan from Research Establishment, Risö, Roskilde, Denmark. Copies are available from Danatom, Aurehøjvej 2, Hellerup, Denmark.
Dep.	AEC reports so designated are maintained in the organizations listed on the inside front cover of NSA. Prior to July 1, 1965, reports were deposited either in printed copy form or in microform. After that time, all reports have been in the form of micronegatives.
	Reports from other countries so designated are available in the United States at the organizations listed on the inside front cover of NSA.

DTIE	Available from the U. S. Atomic Energy Commission, USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830. (Conference papers are available on loan only.)	Documents over 900 pages have special prices which may be obtained from NTIS. Microfiche copy of each separately bound document can be purchased for \$0.95.
EM	Blueline prints of AEC engineering drawings are available from NTIS, U. S. Department of Commerce, Springfield, Virginia 22151. Current prices of the CAPE-packages may be obtained from this address. TID-4100, available from the USAEC Technical Information Center, P. O. Box 62, Oak Ridge, Tennessee 37830, contains information on all the CAPE-packages for sale.	Reports issued by organizations outside the United States will be sold by NTIS <i>only to purchasers within the United States</i> . Purchasers outside the United States may acquire such reports from the sources described elsewhere in these columns.
EUR	For sale by Centrale de vente des publications des Communautés européennes 37, rue Glesener-Luxembourg.	OECD
GPO	For sale by the Superintendent of Documents, Government Printing Office, Washington, D. C. 20402.	Available from O. E. C. D., Publications 2, rue, André-Pascal, 75 Paris 16 ^e , France, and from its sales agencies in various countries throughout the world. A list of these agencies is found in each OECD publication.
IAEA	Available from the International Atomic Energy Agency, Vienna, Austria, and from its sales agencies in various countries throughout the world. A list of these agencies is found in each STI/DOC and STI/PUB publication.	RSO
INIS	Available on microfiche at \$0.65 per document from INIS Section, International Atomic Energy Agency, P. O. Box 590 A-1011, Vienna, Austria	Available <i>only to requesters within the United States</i> from the Reactor Standards Office, Oak Ridge National Laboratory, P. O. Box X, Oak Ridge, TN 37830
LMEC	Available <i>only to requesters within the United States</i> from the Liquid Metal Engineering Center, P. O. Box 1449, Canoga Park, CA 91304	SWED
NORW	Reports from Forsvarets Forskningsinstitut are available from Norwegian Defense Research Establishment, Library, P. O. Box 25, Kjeller, Norway. "KR" reports are available on an exchange basis from the Library, Institutt for Atomenergi, Box 40, Kjeller, Norway. Reports of the Norwegian Reactor School are available from the Library, Institutt for Atomenergi, P. O. Box 40, Kjeller, Norway.	"AE" reports (full size or microcopies) are for sale from Aktiebolaget Atomenergi, Studsvik, Library, Fack, S-611 01 NYKÖPING 1, Sweden. Other series are not generally available to the public. Organizations in the Atomic Energy field may obtain copies from Aktiebolaget Atomenergi.
NRC	For sale by the National Research Council of Canada, Ottawa, Ontario, Canada.	UK
NTIS	For sale by the National Technical Information Service (NTIS), U. S. Department of Commerce, Springfield, Virginia 22151 at the following prices for eye legible copy: 1-300 pages \$3.00 301-600 pages \$6.00 601-900 pages \$9.00	United Kingdom Atomic Energy Authority reports generally may be purchased from Her Majesty's Stationery Office, P. O. Box 569, London S.E.1. Copies of many UKAEA reports (particularly earlier ones) may be purchased in microcopy form from Micro Methods Ltd., East Ardsley, Wakefield, Yorkshire, England. Copies of UKAEA reports are deposited with the National Lending Library for Science and Technology and 10 public libraries in England; with the USAEC and (through USAEC) with NTIS; and in most national nuclear centers throughout the world. In the USA, UKAEA reports are available on a subscription basis, by subject category, from UNIPUB, Inc., P. O. Box 433, New York, NY 10016, USA; and from Pendragon House, Inc., 1093 Charter Avenue, Redwood City, CA 94063, USA.
		UN
		Mail orders should be directed to the Sales Section, Room 1059, United Nations Headquarters, New York, New York 10017.
		ZAED
		Available from ZAED, Kernforschungszentrum, 7501 Leopoldshafen, West Germany

Report No. Abstract No. Availability

AAEC/TM- (Australian)
627 26-45510 Dep. NTIS (U. S. Sales Only), 20p.
628 26-46098 Dep. NTIS (U. S. Sales Only), 19p.

AD- (Misc.)
734392 26-46263 NTIS, 25p.

AE- (Non-AEC)
448 26-45573 Dep. NTIS (U. S. Sales Only), 101p.
451 26-45466 Dep. NTIS (U. S. Sales Only), 47p.
452 26-46916 Dep. NTIS (U. S. Sales Only), 98p.
453 26-45309 Dep. NTIS (U. S. Sales Only), 21p.

AEEP-
3930-109- 26-46365 See ORO-2911-14
72-U

AERE-HP/M- (British)
74 26-45585 Dep. NTIS (U. S. Sales Only), 9p.

AI-AEC-
13033 26-46044 Dep. NTIS, 31p.

ANCR-
1060 26-47010 Dep. NTIS, 95p.

ANL-TKK/RJL-
2 26-47599 Dep. NTIS, 26p.

ARH-
2432 26-45586 Dep. NTIS, 35p.

ARH-SA-
110 26-45219 Dep. NTIS, 18p.
117 26-46099 Dep. NTIS, 22p.
120 26-45381 Dep. NTIS, 14p.
130 26-45382 Dep. NTIS, 22p.

AWRE-O- (British)
31/72 26-47225 Dep. NTIS (U. S. Sales Only). UK 50
pence, 22p.

BARC- (Non-AEC)
577 26-45985 Dep. NTIS (U. S. Sales Only), 25p.
585 26-45986 Dep. NTIS (U. S. Sales Only), 40p.
591 26-45636 Dep. NTIS (U. S. Sales Only), 15p.
593 26-45220 Dep. NTIS (U. S. Sales Only), 14p.
595 26-45310 Dep. NTIS (U. S. Sales Only), 36p.
598 26-46902 Dep. NTIS (U. S. Sales Only), 26p.
599 26-45311 Dep. NTIS (U. S. Sales Only), 163p.

BDX-
613-253 26-47600 Dep. NTIS, 54p.
(Rev.)
613-288 26-47601 Dep. NTIS, 32p.
(Rev.)
613-392 26-47602 Dep. NTIS, 34p.
(Rev.)
613-440 26-47603 Dep. NTIS, 32p.
(Rev.)
613-456 26-47604 Dep. NTIS, 79p.
613-515 26-47605 Dep. NTIS, 20p.
(Rev.)
613-543 26-47606 Dep. NTIS, 11p.
(Rev.)
613-579 26-47607 Dep. NTIS, 24p.
(Rev.)
613-604 26-47608 Dep. NTIS, 17p.
(Rev.)
613-642 26-47609 Dep. NTIS, 45p.
(Rev.)

BFL- (West German)
1/1972 26-45189 Dep. NTIS (U. S. Sales Only), 23p.
2/1972 26-45716 Dep. NTIS (U. S. Sales Only), 51p.

BMW-FBK- (West German)
72-6 26-47594 Dep. NTIS (U. S. Sales Only), 81p.

BMI- (Misc.)
1927 26-45966 Dep. NTIS, 66p.

BNL-
16913 26-46805 Dep. NTIS, 6p.
16928 26-46616 Dep. NTIS, 6p.
16930 26-46092 Dep. NTIS, 13p.
16935 26-46945 Dep. NTIS, 47p.
16954 26-46471 Dep. NTIS, 17p.
16968 26-46426 Dep. NTIS, 7p.
50332 26-45587 Dep. NTIS, 47p.

BNL-tr-
477 26-46917 Dep. NTIS
487 26-47038 Dep. NTIS
488 26-47039 Dep. NTIS
489 26-47040 Dep. NTIS

BNWL-
1511 26-47247 Dep. NTIS, 21p.

Report No. Abstract No. Availability
1635 26-47317 Dep. NTIS, 125p.
1648 26-47011 Dep. NTIS, 28p.

BNWL-SA-
4205(Rev.) 26-45530 Dep. NTIS, 18p.
4320 26-45467 Dep. NTIS, 4p.

BRH/DMRE- (Non-AEC)
70-4 26-45392 See N-72-15068

CALT-
767-P-3-22 26-46510 Dep. NTIS, 31p.

CAPE-
724 26-45452 EM
1657 26-46116 EM

CEA-CONF- (French)
1998 26-45699 Dep. NTIS (U. S. Sales Only), 30p.
2008 26-45435 Dep. NTIS (U. S. Sales Only), 16p.
2010 26-45436 Dep. NTIS (U. S. Sales Only), 13p.

CEA-N- (French)
1483 26-47581 Dep. NTIS (U. S. Sales Only), 132p.
1522 26-46873 Dep. NTIS (U. S. Sales Only), 358p.
1526 26-47012 Dep. NTIS (U. S. Sales Only), 272p.
1529 26-47041 Dep. NTIS (U. S. Sales Only), 118p.

CERN- (European Organization for Nuclear Research)
72-8 26-47283 Dep. NTIS (U. S. Sales Only), 47p.

CISE-R- (Italian)
172 26-45181 Dep. NTIS (U. S. Sales Only), 116p.
178 26-47315 Dep. NTIS (U. S. Sales Only), 59p.
245 26-45458 Dep. NTIS (U. S. Sales Only), 175p.
257 26-47309 Dep. NTIS (U. S. Sales Only), 221p.
259 26-47580 Dep. NTIS (U. S. Sales Only), 29p.
275 26-45459 Dep. NTIS (U. S. Sales Only), 57p.

CONF-610904- (Conference on Elementary Particles, Aix-en-
1 Provence, France, Sept. 14-20, 1961)
26-46449 See SLAC-Trans-143

CONF-700635- (2nd Symposium on the Problems of Nuclear
Physics, Novosibirsk, U.S.S.R., June 12-
19, 1970)
26-47086 Moscow; Izdatel'stvo Nauka (1971).
544p

CONF-701151- (Meeting on Metrology, Paris, France, Nov.
17, 1970)
1 26-47217 pp 65-75 of Proceedings of the
Meeting on Metrology, Paris;
Bureau National de la Metrologie
(1971)
2 26-47218 pp 181-8 of Proceedings of the
Meeting on Metrology, Paris;
Bureau National de la Metrologie
(1971)
3 26-47219 pp 175-80 of Proceedings of the
Meeting on Metrology, Paris;
Bureau National de la Metrologie
(1971)

CONF-701229- (2nd International Clean Air Congress
Conference, Washington, D.C. Dec. 6-11,
1970)
26-45205 New York; Academic Press, Inc.
(1971). 1378p. \$85.00

CONF-710345- (The Moriond Meeting on the Phenomenology of
Weak and Electromagnetic Interactions,
Meribel-les-Allues, France, Mar. 7, 1971)
(Vol.2) 26-46590 Paris; Laboratoire de Physique
Theorique et Hautes Energies
(1971). 559p

CONF-710431- (14th Conference on Great Lakes Research,
Toronto, Ontario, Canada, Apr. 19-21,
1971)
26-45577 Ann Arbor, Mich.; International
Association for Great Lakes
Research (1971). 844p

CONF-710501- (3rd National Symposium on Radioecology, Oak
Ridge, Tenn., May 10-12, 1971)
55 26-45541 See NYO-2174-130

CONF-710601- (International Conference on Radiation-
Induced Vacancies in Metals, Albany, N.Y.,
June 9-11, 1971)
26-46045 Dep. NTIS, 884p.

CONF-710651- (International Conference on Heat Exchangers,
Paris, France, June 16-18, 1971)
2 26-45436 See CEA-CONF-2010
3 26-45435 See CEA-CONF-2008

CONF-710710- (16th Annual Health Physics Society Meeting,
New York, N.Y., July 12-16, 1971)
12 26-45790 Health Phys.; 23: No. 2, 187-92(Aug
1972)

CONF-710816-	(International Symposium on Electron and Photon Interactions at High Energies, Ithaca, N.Y., Aug. 23-27, 1971)
9	26-46511 Dep. NTIS (U. S. Sales Only), 21p.
11	26-46526 See TPJU-14/71
12	26-46472 Dep. NTIS (U. S. Sales Only), 18p.
13	26-46427 Dep. NTIS (U. S. Sales Only), 12p.
14	26-46428 Dep. NTIS (U. S. Sales Only), 16p.
15	26-46429 Dep. NTIS (U. S. Sales Only), 8p.
16	26-46512 Dep. NTIS (U. S. Sales Only), 29p.
17	26-46513 Dep. NTIS (U. S. Sales Only), 8p.
18	26-46515 See DAMTP-71/24
19	26-46696
CONF-710913-	(162nd National Meeting of the American Chemical Society, Washington, D.C., Sept. 12-17, 1971)
3	26-47637 See UR-49-1508
CONF-710953-	(European Conference of Geochronology, Brussels, Belgium, Sept. 6, 1971)
2	26-45225 Ann. Soc. Geol. Belg.; 94: No. 2, 105-6(1971)
4	26-45197 Ann. Soc. Geol. Belg.; 94: No. 2, 132(1971)
CONF-710958-	(International Conference on the Noise in Active Semiconductor Devices, Toulouse, France, Sept. 21, 1971)
I	26-45660 See FRNC-CONF-29
CONF-711008-	(19th Conference on Remote Systems Technology, Miami Beach, Fla., Oct. 17-21, 1971)
5	26-45699 See CEA-CONF-1998
CONF-711218-	(Recent Developments in Thermophysical Properties Research Meeting, Washington, D.C., Dec. 1, 1971)
	26-45463 New York; American Society of Mechanical Engineers (1971). 18p. \$3.50
CONF-720119-	(American Physical Society Meeting, San Francisco, CA., Jan. 31-Feb. 3, 1972)
1	26-46943 See N-72-18705
CONF-720301-	(Spring Conference of the American Society for Nondestructive Testing, Los Angeles, Calif., Mar. 13-17, 1972)
5	26-45934 See N-72-18585
CONF-720312-	(Underground Nuclear Test Measurement Symposium II, Menlo Park, Ca., Mar. 21-22, 1972)
11	26-45674 See UCRL-73665
CONF-720409-	(3rd International Conference on Experimental Meson Spectroscopy, Philadelphia, PA., Apr. 28-29, 1972)
3	26-46616 See BNL-16928
4	26-46473 See COO-1195-239
5	26-46471 See BNL-16954
6	26-46430 See COO-1195-241
7	26-46475 See IS-2886
CONF-720420-	(Conference on Reactor Materials Performance, Richland, WA., Apr. 24-26, 1972)
8	26-46077 See GULF-GA-A-12035
CONF-720505-	(12th Hanford Biology Symposium on Radionuclide Carcinogenesis, Richland, WA., May 11-13, 1972)
2	26-45910 Dep. NTIS, 22p.
3	26-45819 See UR-3490-63
4	26-45895 See UR-3490-102
CONF-720513-	(Applied Superconductivity Meeting, Annapolis, MD., May 1-3, 1972)
8	26-46805 See BNL-16913
CONF-720520-	(Medical Planning and Care in Radiation Accidents Seminar, Oak Ridge, TN., May 21-24, 1972)
1	26-45888 Dep. NTIS, 14p.
CONF-720529-	(13th Annual Meeting of the Institute of Nuclear Materials Management, Boston, MA., May 31--June 2, 1972)
3	26-46092 See BNL-16930
CONF-720541-	(International Clean Air Conference, Melbourne, Australia, May 15-18, 1972)
	26-45501 Parkville, Australia; Clean Air Society of Australia and New Zealand (1972). 308p
CONF-720607-	(18th Annual American Nuclear Society

Report No.	Abstract No.	Availability
		Conference, Las Vegas, NV., June 18-22, 1972)
19	26-45382	See ARH-SA-130
22	26-46144	Dep. NTIS, 4p.
24	26-45487	See UCRL-73939
CONF-720619-	(65th Meeting of the Air-Pollution Control Association, Miami Beach, FL., June 18-22, 1972)	
1	26-45574	Dep. NTIS, 21p.
CONF-720625-	(4th Symposium on Damage in Laser Materials, Boulder, CO., June 14-15, 1972)	
1	26-47636	See UCRL-73925
CONF-720627-	(Symposium on the Role of Nuclear Energy in Australia's Development, Canberra, Australia, June 1-2, 1972)	
1	26-46098	See AAEC/TM-628
2	26-45510	See AAEC/TM-627
CONF-720628-	(27th ACS Northwest Regional Meeting, Corvallis, OR., June 15-16, 1972)	
2	26-45381	See ARH-SA-120
3	26-45219	See ARH-SA-110
CONF-720706-	(Power Engineering Society Summer Meeting, San Francisco, CA., July 9-14, 1972)	
1	26-47634	See UCRL-73587
CONF-720707-	(Annual Conference on Nuclear and Space Radiation Effects, Seattle, WA., July 24-27, 1972)	
2	26-45662	See SC-DC-721232
3	26-45663	See SC-DC-721233
4	26-45664	See SC-DC-721237
5	26-45665	See SC-DC-721240
10	26-45668	See UCRL-73974
CONF-720708-	(Symposium on the Interaction of Radioactive Contaminants with The Constituents of the Marine Environment, Seattle, WA., July 10-14, 1972)	
2	26-45544	See RLD-2225-T-14-4
3	26-45543	See RLD-2225-T-1-9
4	26-45531	See COO-3563-3
6	26-45530	See BNWL-SA-4205(Rev.)
7	26-45545	See UCRL-73744
CONF-720710-	(2nd Topical Conference on Pulsed High-Beta Plasmas, Garching, Germany, July 3-6, 1972)	
1	26-47158	See LA-DC-72-726
3	26-47156	See LA-DC-72-706
4	26-47157	See LA-DC-72-707
5	26-47141	See LA-DC-72-703
CONF-720802-	(13th National Heat Transfer Conference, Denver, CO., Aug. 6-9, 1972)	
1	26-47312	Dep. NTIS, 23p.
CONF-720806-	(6th National Symposium on Fracture Mechanics, Philadelphia, PA., Aug. 28-30, 1972)	
2	26-45990	Dep. NTIS, 29p.
CONF-720807-	(5th International Symposium on Discharges and Electrical Insulation in Vacuum, Poznan, Poland, Aug. 31-Sept. 1, 1972)	
1	26-47625	See SC-DC-722267
CONF-720809-	(23rd Meeting of the International Society of Electrochemistry, Stockholm, Sweden, Aug. 27-Sept. 2, 1972)	
1	26-47635	See UCRL-73805
CONF-720902-	(National Topical Meeting on the Applications of Californium-252, Austin, TX., Sept. 11-13, 1972)	
1	26-45467	See BNWL-SA-4320
2	26-45714	Dep. NTIS, 5p.
CONF-720903-	(20th Remote Systems Technology Conference, Idaho Falls, ID., Sept. 17-21, 1972)	
2	26-46108	See UCRL-73757
3	26-46104	See LA-DC-72-556
4	26-46100	Dep. NTIS, 15p.
5	26-46107	See UCRL-73756
8	26-45224	See LA-DC-72-524
9	26-45397	Dep. NTIS, 32p.
11	26-46099	See ARH-SA-117
CONF-720904-	(Conference on The Impact of Aerospace Technology on Society in The 70's, Los Angeles, CA., Sept. 11-13, 1972)	
1	26-45437	Dep. NTIS, 27p.
COO-916-25	26-45991	Dep. NTIS, 27p.

Report No.	Abstract No.	Availability
1112-51	26-46254	Dep. NTIS, 7p.
1112-75	26-46456	Dep. NTIS, 22p.
1120-149	26-46903	Dep. NTIS, 91p.
1195-239	26-46473	Dep. NTIS, 9p.
1195-241	26-46430	Dep. NTIS, 7p.
1222-55	26-45302	Dep. NTIS, 19p.
1427-41	26-45813	Dep. NTIS, 31p.
1428-306	26-46474	Dep. NTIS, 59p.
2114-2	26-45588	Dep. NTIS, 6p.
2118-34	26-47610	Dep. NTIS, 94p.
2147-2	26-45312	Dep. NTIS, 25p.
2155-4	26-45700	Dep. NTIS, 42p.
2195-5	26-46264	Dep. NTIS, 18p.
2195-6	26-45503	Dep. NTIS, 16p.
3001-12	26-47042	Dep. NTIS, 16p.
3013-1	26-45904	Dep. NTIS, 12p.
3013-25	26-45911	Dep. NTIS, 48p.
3017-43	26-45814	Dep. NTIS, 29p.
3029-1	26-45313	Dep. NTIS, 6p.
3058-23	26-47013	Dep. NTIS, 16p.
3078-5	26-47611	Dep. NTIS, 26p.
3078-6	26-47612	Dep. NTIS, 11p.
3086-15	26-47145	Dep. NTIS, 10p.
3232-1	26-45722	Dep. NTIS, 6p.
3286-1	26-45314	Dep. NTIS, 7p.
3286-3	26-45315	Dep. NTIS, 10p.
3292-1	26-47613	Dep. NTIS, 113p.
3311-1	26-45972	Dep. NTIS, 7p.
3431-1	26-46066	Dep. NTIS, 6p.
3486-1	26-45815	Dep. NTIS, 19p.
3498-2	26-47614	Dep. NTIS, 7p.
3563-3	26-45531	Dep. NTIS, 22p.
CPT-	(Non-AEC)	
134	26-46525	See ORO-3992-80
CSIRO-Trans-	(Australian)	
10705	26-45653	Dep. NTIS (U. S. Sales Only)
CTH-ITP-	(Swedish)	
71-14	26-46665	Dep. NTIS (U. S. Sales Only), 39p.
71-17	26-46773	Dep. NTIS (U. S. Sales Only), 6p.
71-30	26-46514	Dep. NTIS (U. S. Sales Only), 41p.
CU-	(Misc.)	
3194-22	26-47615	Dep. NTIS, 92p.
DAMTP-	(British)	
71/24	26-46515	Dep. NTIS (U. S. Sales Only), 24p.
DESY-	(West German)	
72/20	26-46697	Dep. NTIS (U. S. Sales Only), 53p.
72/21	26-46516	Dep. NTIS, 11p.
72/23	26-46517	Dep. NTIS (U. S. Sales Only), 54p.
DOCKET-1151-	(Elk River Reactor) (AEC)	
57	26-47336	Dep. NTIS, 32p.
DOCKET-50003-	(Indian Point Station, Unit 1; See also DOCKET-50247 and DOCKET-50286) (Non-AEC)	
118	26-47320	Dep. NTIS, 9p.
119	26-47337	Dep. NTIS, 3p.
120	26-47388	Dep. NTIS, 177p.
DOCKET-50010-	(Dresden Nuclear Power Station, Unit 1; See also DOCKET-50237 and DOCKET-50249) (Non-AEC)	
113	26-47389	Dep. NTIS, 8p.
114	26-47338	Dep. NTIS, 8p.
DOCKET-50016-	(Enrico Fermi Atomic Power Plant) (Non-AEC)	
160	26-47327	Dep. NTIS, 1p.
163	26-47390	Dep. NTIS, 43p.
164	26-47339	Dep. NTIS, 12p.
165	26-47340	Dep. NTIS, 1p.
DOCKET-50029-	(Yankee Atomic Electric Co.) (Non-AEC)	
189	26-47341	Dep. NTIS, 9p.
190	26-47391	Dep. NTIS, 1p.
DOCKET-50133-	(Humboldt Bay Power Plant) (Non-AEC)	
89	26-47392	Dep. NTIS, 13p.
91	26-47342	Dep. NTIS, 2p.
DOCKET-50155-	(Big Rock Point Reactor) (Misc.)	
117	26-47393	Dep. NTIS, 2p.
119	26-47394	Dep. NTIS, 6p.
120	26-47395	Dep. NTIS, 22p.
DOCKET-50201-	(West Valley Reprocessing Plant--Formerly Spent Fuel Processing Plant) (Misc.)	
125	26-45398	Dep. NTIS, 8p.
126	26-45399	Dep. NTIS, 48p.
127	26-45400	Dep. NTIS, 2p.
128	26-45401	Dep. NTIS, 3p.
DOCKET-50213-	(Haddam Neck Plant, Unit 1) (Misc.)	
150	26-47396	Dep. NTIS, 32p.
152	26-47397	Dep. NTIS, 77p.

DOCKET-50219-	(Oyster Creek Nuclear Power Plant) (Non-AEC)	
196	26-47344	Dep. NTIS, 1p.
197	26-47345	Dep. NTIS, 2p.
199	26-47346	Dep. NTIS, 3p.
200	26-47398	Dep. NTIS, 31p.
201	26-47399	Dep. NTIS, 2p.
202	26-47400	Dep. NTIS, 38p.
203	26-47347	Dep. NTIS, 2p.
204	26-47348	Dep. NTIS, 2p.
DOCKET-50220-	(Nine Mile Point Nuclear Station) (Non-AEC)	
98	26-47401	Dep. NTIS, 440p.
100	26-47349	Dep. NTIS, 4p.
101	26-47350	Dep. NTIS, 2p.
102	26-47575	Dep. NTIS, 2p.
103	26-47351	Dep. NTIS, 18p.
DOCKET-50231-	(Southwest Experimental Fast Oxide Reactor) (Non-AEC)	
120	26-47582	Dep. NTIS, 5p.
121	26-47583	Dep. NTIS, 2p.
122	26-47584	Dep. NTIS, 1p.
DOCKET-50237-	(Dresden Nuclear Power Station, Unit 2; See also DOCKET-50010 and DOCKET-50249)	
244	26-47402	Dep. NTIS, 8p.
245	26-47352	Dep. NTIS, 2p.
246	26-47403	Dep. NTIS, 6p.
247	26-47353	Dep. NTIS, 1p.
248	26-47354	Dep. NTIS, 2p.
249	26-47404	Dep. NTIS, 45p.
250	26-47331	Dep. NTIS, 2p.
251	26-47355	Dep. NTIS, 6p.
252	26-47356	Dep. NTIS, 2p.
253	26-47357	Dep. NTIS, 4p.
254	26-47358	Dep. NTIS, 5p.
255	26-47405	Dep. NTIS, 5p.
DOCKET-50244-	(Robert Emmett Ginna Nuclear Power Plant, Formerly Brookwood Nuclear Station) (Non-AEC)	
106	26-47359	Dep. NTIS, 2p.
107	26-47406	Dep. NTIS, 2p.
108	26-47576	Dep. NTIS, 207p.
109	26-47321	Dep. NTIS, 2p.
DOCKET-50245-	(Millstone Point Nuclear Power Station) (Non-AEC)	
139	26-47407	Dep. NTIS, 1p.
140	26-47408	Dep. NTIS, 9p.
141	26-47409	Dep. NTIS, 24p.
143	26-47360	Dep. NTIS, 1p.
144	26-47410	Dep. NTIS, 34p.
145	26-47411	Dep. NTIS, 3p.
146	26-47361	Dep. NTIS, 63p.
DOCKET-50247-	(Indian Point Nuclear Generating, Unit 2; See also DOCKET-50286 and DOCKET-50003) (Non-AEC)	
76	26-47412	Dep. NTIS, 22p.
77	26-47322	Dep. NTIS, 9p.
DOCKET-50249-	(Dresden Nuclear Power Station, Unit 3; See also DOCKET-50010 and DOCKET-50237) (Non-AEC)	
141	26-47402	See DOCKET-50237-244
143	26-47352	See DOCKET-50237-245
144	26-47403	See DOCKET-50237-246
145	26-47353	See DOCKET-50237-247
146	26-47354	See DOCKET-50237-248
147	26-47404	See DOCKET-50237-249
148	26-47332	Dep. NTIS, 2p.
149	26-47358	See DOCKET-50237-254
150	26-47405	See DOCKET-50237-255
151	26-47362	Dep. NTIS, 2p.
DOCKET-50250-	(Turkey Point, See also DOCKET-50251) (Non-AEC)	
84	26-47413	Dep. NTIS, 368p.
85	26-47414	Dep. NTIS, 214p.
86	26-47415	Dep. NTIS, 2p.
88	26-47416	Dep. NTIS, 4p.
DOCKET-50251-	(Turkey Point, See also DOCKET-50250) (Non-AEC)	
70	26-47413	See DOCKET-50250-84
71	26-47414	See DOCKET-50250-85
72	26-47417	Dep. NTIS, 142p.
73	26-47415	See DOCKET-50250-86
75	26-47416	See DOCKET-50250-88
DOCKET-50254-	(Quad-Cities Station; See also DOCKET-50265) (Non-AEC)	
104	26-47402	See DOCKET-50237-244
105	26-47363	Dep. NTIS, 2p.
107	26-47418	Dep. NTIS, 12p.
108	26-47364	Dep. NTIS, 2p.
109	26-47419	Dep. NTIS, 110p.
110	26-47420	Dep. NTIS, 8p.
111	26-47358	See DOCKET-50237-254

Report No. Abstract No. Availability
112 26-47421 Dep. NTIS, 1p.

DOCKET-50255- (Palisades Plant) (Non-AEC)
94 26-47422 Dep. NTIS, 2p.
95 26-47423 Dep. NTIS, 13p.
96 26-47424 Dep. NTIS, 360p.

DOCKET-50259- (Browns Ferry Nuclear Power Station. See also
DOCKET-50260 and DOCKET-50296) (Non-AEC)
57 26-47425 Dep. NTIS, 22p.
58 26-47426 Dep. NTIS, 25p.
59 26-47427 Dep. NTIS, 108p.
60 26-47428 Dep. NTIS, 15p.

DOCKET-50260- (Browns Ferry Nuclear Power Station. See also
DOCKET-50259 and DOCKET-50296) (Non-AEC)
53 26-47425 See DOCKET-50259-57
54 26-47426 See DOCKET-50259-58
55 26-47427 See DOCKET-50259-59
56 26-47428 See DOCKET-50259-60

DOCKET-50261- (H. B. Robinson) (Non-AEC)
101 26-47365 Dep. NTIS, 1p.
102 26-47366 Dep. NTIS, 1p.

DOCKET-50263- (Monticello Nuclear Generating Plant) (Non-
AEC)
137 26-47367 Dep. NTIS, 7p.
138 26-47429 Dep. NTIS, 5p.
139 26-47368 Dep. NTIS, 2p.
140 26-47369 Dep. NTIS, 4p.
141 26-47323 Dep. NTIS, 2p.
142 26-47333 Dep. NTIS, 1p.
143 26-47334 Dep. NTIS, 1p.

DOCKET-50265- (Quad-Cities Station; See also DOCKET-50254)
(Non-AEC)
91 26-47402 See DOCKET-50237-244
92 26-47418 See DOCKET-50254-107
93 26-47364 See DOCKET-50254-108
94 26-47419 See DOCKET-50254-109
95 26-47370 Dep. NTIS, 6p.
96 26-47371 Dep. NTIS, 1p.
97 26-47420 See DOCKET-50254-110
98 26-47358 See DOCKET-50237-254
99 26-47421 See DOCKET-50254-112
100 26-47335 Dep. NTIS, 2p.
101 26-47325 Dep. NTIS, 1p.

DOCKET-50266- (Point Beach Nuclear Plant) (Non-AEC)
73 26-47372 Dep. NTIS, 3p.
74 26-47430 Dep. NTIS, 10p.
75 26-47373 Dep. NTIS, 58p.

DOCKET-50268- (Midwest Fuel Recovery Plant) (Non-AEC)
31 26-45402 Dep. NTIS, 5p.
32 26-45403 Dep. NTIS, 5p.
33 26-45404 Dep. NTIS, 6p.

DOCKET-50269- (Oconee Nuclear Station, See also DOCKET-
50270, and DOCKET-50287) (Non-AEC)
56 26-47431 Dep. NTIS, 2p.
60 26-47432 Dep. NTIS, 224p.
61 26-47433 Dep. NTIS, 3p.

DOCKET-50270- (Oconee Nuclear Station, See also DOCKET-
50269, and DOCKET-50287) (Non-AEC)
48 26-47432 See DOCKET-50269-60
49 26-47434 Dep. NTIS, 5p.
50 26-47433 See DOCKET-50269-61

DOCKET-50271- (Vermont Yankee Nuclear Power Station) (Non-
AEC)
88 26-47435 Dep. NTIS, 357p.
90 26-47374 Dep. NTIS, 195p.
91 26-47436 Dep. NTIS, 7p.
92 26-47437 Dep. NTIS, 26p.
93 26-47438 Dep. NTIS, 4p.
94 26-47439 Dep. NTIS, 3p.
95 26-47440 Dep. NTIS, 18p.

DOCKET-50272- (Salem Nuclear Generating Station--Formerly
Burlington Nuclear Generating Station, See
also Docket-50311) (Non-AEC)
29 26-47441 Dep. NTIS, 173p.
30 26-47442 Dep. NTIS, 154p.
31 26-47443 Dep. NTIS, 109p.
32 26-47444 Dep. NTIS, 213p.
33 26-47445 Dep. NTIS, 206p.

DOCKET-50275- (Diablo Canyon Nuclear Power Plant, See also
DOCKET-50323) (Non-AEC)
22 26-47446 Dep. NTIS, 484p.
23 26-47447 Dep. NTIS, 716p.

DOCKET-50277- (Peach Bottom Atomic Power Station; See also
DOCKET-50171 and DOCKET-50278) (Non-AEC)
51 26-47375 Dep. NTIS, 4p.
52 26-47448 Dep. NTIS, 11p.

DOCKET-50278- (Peach Bottom Atomic Power Station; See also
DOCKET-50171 and DOCKET-50277) (Non-AEC)

167 REPORTS

DOCKET-50311

Report No. Abstract No. Availability
51 26-47375 See DOCKET-50277-51
52 26-47448 See DOCKET-50277-52

DOCKET-50280- (Surry Power Station, See also DOCKET-50281)
(Non-AEC)
77 26-47449 Dep. NTIS, 48p.
78 26-47328 Dep. NTIS, 2p.

DOCKET-50281- (Surry Power Station, See also DOCKET-50280)
(Non-AEC)
75 26-47449 See DOCKET-50280-77

DOCKET-50282- (Prairie Island Nuclear Generating Plant, See
also DOCKET-50306) (Non-AEC)
59 26-47450 Dep. NTIS, 3p.
67 26-47451 Dep. NTIS, 713p.
68 26-47452 Dep. NTIS, 4p.
69 26-47453 Dep. NTIS, 252p.

DOCKET-50285- (Fort Calhoun Station) (Non-AEC)
68 26-47454 Dep. NTIS, 99p.

DOCKET-50286- (Indian Point Nuclear Plant, Unit 3; See also
DOCKET-50003 and DOCKET-50247)
46 26-47455 Dep. NTIS, 36p.
47 26-47456 Dep. NTIS, 51p.
48 26-47457 Dep. NTIS, 104p.
49 26-47458 Dep. NTIS, 400p.

DOCKET-50287- (Oconee Nuclear Station; See also DOCKET-
50269 and DOCKET-50270) (Non-AEC)
43 26-47432 See DOCKET-50269-60
44 26-47434 See DOCKET-50270-49
45 26-47433 See DOCKET-50269-61

DOCKET-50289- (Three Mile Island Nuclear Station) (Non-AEC)
61 26-47459 Dep. NTIS, 4p.
62 26-47460 Dep. NTIS, 4p.
63 26-47461 Dep. NTIS, 136p.

DOCKET-50293- (Pilgrim Nuclear Power Station) (Non-AEC)
80 26-47376 Dep. NTIS, 3p.
82 26-47377 Dep. NTIS, 3p.
83 26-47378 Dep. NTIS, 3p.

DOCKET-50295- (Zion Station Nuclear Power Plant. See also
DOCKET-50304) (Non-AEC)
57 26-47462 Dep. NTIS, 10p.
58 26-47463 Dep. NTIS, 8p.
59 26-47464 Dep. NTIS, 7p.

DOCKET-50296- (Browns Ferry Nuclear Power Station. See
also DOCKET-50259 and DOCKET-50260) (Non-
AEC)
50 26-47425 See DOCKET-50259-57
51 26-47426 See DOCKET-50259-58
52 26-47427 See DOCKET-50259-59
53 26-47428 See DOCKET-50259-60

DOCKET-50298- (Cooper Nuclear Station) (Non-AEC)
43 26-47465 Dep. NTIS, 46p.
46 26-47466 Dep. NTIS, 256p.
47 26-47467 Dep. NTIS, 8p.
48 26-47468 Dep. NTIS, 41p.

DOCKET-50301- (Point Beach Nuclear Plant) (Non-AEC)
64 26-47469 Dep. NTIS, 11p.
65 26-47372 See DOCKET-50266-73
66 26-47430 See DOCKET-50266-74

DOCKET-50304- (Zion Station Nuclear Power Plant. See also
DOCKET-50295) (Non-AEC)
56 26-47462 See DOCKET-50295-57
57 26-47463 See DOCKET-50295-58
58 26-47464 See DOCKET-50295-59
59 26-47379 Dep. NTIS, 2p.

DOCKET-50305- (Kewaunee Nuclear Power Plant) (Non-AEC)
44 26-47470 Dep. NTIS, 197p.
45 26-47471 Dep. NTIS, 249p.
46 26-47380 Dep. NTIS, 5p.
47 26-47472 Dep. NTIS, 6p.

DOCKET-50306- (Prairie Island Nuclear Generating Plant, See
also DOCKET-50282) (Non-AEC)
64 26-47451 See DOCKET-50282-67
65 26-47452 See DOCKET-50282-68
66 26-47453 See DOCKET-50282-69

DOCKET-50309- (Maine Yankee Atomic Power Company) (Non-AEC)
61 26-47473 Dep. NTIS, 231p.
62 26-47474 Dep. NTIS, 96p.
63 26-47475 Dep. NTIS, 66p.
64 26-47476 Dep. NTIS, 113p.
65 26-47477 Dep. NTIS, 391p.
66 26-47478 Dep. NTIS, 33p.

DOCKET-50311- (Salem Nuclear Generating Station--Formerly
Burlington Nuclear Generating Station, See

Report No.	Abstract No.	Availability
also DOCKET-50272) (Non-AEC)		
29	26-47441	See DOCKET-50272-29
30	26-47442	See DOCKET-50272-30
31	26-47443	See DOCKET-50272-31
32	26-47444	See DOCKET-50272-32
33	26-47445	See DOCKET-50272-33

DOCKET-50312- (Rancho Seco Nuclear Generating Station) (Non-AEC)

41	26-47479	Dep. NTIS, 196p.
42	26-47480	Dep. NTIS, 497p.
43	26-47481	Dep. NTIS, 39p.
45	26-47482	Dep. NTIS, 43p.
46	26-47483	Dep. NTIS, 17p.
47	26-47484	Dep. NTIS, 39p.
48	26-47485	Dep. NTIS, 19p.

DOCKET-50313- (Arkansas Nuclear One, Unit 1) (Non-AEC)

44	26-47486	Dep. NTIS, 44p.
45	26-47487	Dep. NTIS, 91p.
46	26-47488	Dep. NTIS, 167p.

DOCKET-50315- (Donald C. Cook Nuclear Power Plant, See also DOCKET-50316) (Non-AEC)

65	26-47489	Dep. NTIS, 30p.
68	26-47490	Dep. NTIS, 176p.
69	26-47491	Dep. NTIS, 5p.
70	26-47492	Dep. NTIS, 7p.
71	26-47493	Dep. NTIS, 13p.

DOCKET-50316- (Donald C. Cook Nuclear Power Plant, See also DOCKET-50315) (Non-AEC)

65	26-47489	See DOCKET-50315-65
68	26-47490	See DOCKET-50315-68
69	26-47491	See DOCKET-50315-69
70	26-47492	See DOCKET-50315-70
71	26-47493	See DOCKET-50315-71

DOCKET-50317- (Calvert Cliffs Nuclear Power Plant, See also DOCKET-50318) (Non-AEC)

53	26-47381	Dep. NTIS, 6p.
----	----------	----------------

DOCKET-50318- (Calvert Cliffs Nuclear Power Plant, See also DOCKET-50317) (Non-AEC)

51	26-47381	See DOCKET-50317-53
----	----------	---------------------

DOCKET-50321- (Edwin I. Hatch Nuclear Plant) (Non-AEC)

46	26-47494	Dep. NTIS, 119p.
48	26-47495	Dep. NTIS, 8p.

DOCKET-50323- (Diablo Canyon Nuclear Power Plant, See also DOCKET-50275) (Non-AEC)

20	26-47446	See DOCKET-50275-22
21	26-47447	See DOCKET-50275-23

DOCKET-50324- (Brunswick Steam Electric Plant, Units 1 and 2, See also DOCKET-50325) (Non-AEC)

39	26-47496	Dep. NTIS, 3p.
40	26-47497	Dep. NTIS, 111p.

DOCKET-50325- (Brunswick Steam Electric Plant, Units 1 and 2, See also DOCKET-50324) (Non-AEC)

40	26-47496	See DOCKET-50324-39
41	26-47497	See DOCKET-50324-40

DOCKET-50331- (Daune Arnold Energy Center) (Non-AEC)

48	26-47498	Dep. NTIS, 93p.
49	26-47499	Dep. NTIS, 518p.
51	26-47500	Dep. NTIS, 172p.

DOCKET-50333- (James A. Fitzpatrick Nuclear Power Plant) (Non-AEC)

50	26-47501	Dep. NTIS, 172p.
51	26-47502	Dep. NTIS, 22p.

DOCKET-50334- (Beaver Valley Power Station) (Non-AEC)

31	26-47503	Dep. NTIS, 8p.
----	----------	----------------

DOCKET-50335- (Hutchinson Island Plant) (Non-AEC)

29	26-47504	Dep. NTIS, 34p.
30	26-47505	Dep. NTIS, 17p.
32	26-47506	Dep. NTIS, 11p.

DOCKET-50336- (Millstone Nuclear Power Station) (Non-AEC)

35	26-47507	Dep. NTIS, 55p.
36	26-47508	Dep. NTIS, 802p.
37	26-47509	Dep. NTIS, 720p.

DOCKET-50338- (North Anna Power Station; See also DOCKET-50339) (Non-AEC)

47	26-47510	Dep. NTIS, 42p.
48	26-47511	Dep. NTIS, 187p.
49	26-47512	Dep. NTIS, 3p.

DOCKET-50339- (North Anna Power Station; See also DOCKET-50338) (Non-AEC)

44	26-47510	See DOCKET-50338-47
45	26-47511	See DOCKET-50338-48
46	26-47512	See DOCKET-50338-49

Report No.	Abstract No.	Availability
DOCKET-50341- (Enrico Fermi Atomic Power Plant) (Non-AEC)		
42	26-47513	Dep. NTIS, 3p.
43	26-47514	Dep. NTIS, 12p.
44	26-47329	Dep. NTIS, 3p.

DOCKET-50344- (Trojan Nuclear Plant) (Non-AEC)

29	26-47515	Dep. NTIS, 27p.
30	26-47516	Dep. NTIS, 50p.

DOCKET-50346- (Davis-Besse Nuclear Power Station) (Non-AEC)

27	26-47517	Dep. NTIS, 5p.
29	26-47518	Dep. NTIS, 74p.
30	26-47326	Dep. NTIS, 4p.

DOCKET-50348- (Joseph M. Farley Nuclear Plant, Units 1 and 2, Formerly Seala Nuclear Plant, See also DOCKET-50364) (Non-AEC)

42	26-47519	Dep. NTIS, 13p.
----	----------	-----------------

DOCKET-50352- (Limerick Generating Station) (See also DOCKET-50353) (Non-AEC)

27	26-47520	Dep. NTIS, 56p.
----	----------	-----------------

DOCKET-50353- (Limerick Generating Station; See also DOCKET-50352) (Non-AEC)

27	26-47520	See DOCKET-50352-27
----	----------	---------------------

DOCKET-50354- (Newbold Island Nuclear Generating Station; See also DOCKET-50355) (Non-AEC)

25	26-47521	Dep. NTIS, 11p.
----	----------	-----------------

DOCKET-50361- (San Onofre Nuclear Generating Station, Units 2 and 3; See also DOCKET-50362 and DOCKET-50206, Unit 1) (Non-AEC)

25	26-47522	Dep. NTIS, 179p.
26	26-47523	Dep. NTIS, 285p.

DOCKET-50362- (San Onofre Nuclear Generating Station, Units 2 and 3; See also DOCKET-50361 and DOCKET-50206, Unit 1) (Non-AEC)

24	26-47522	See DOCKET-50361-25
25	26-47523	See DOCKET-50361-26

DOCKET-50363- (Forked River Nuclear Station, Unit 1) (Non-AEC)

30	26-47524	Dep. NTIS, 74p.
31	26-47525	Dep. NTIS, 50p.
32	26-47526	Dep. NTIS, 1p.
34	26-47527	Dep. NTIS, 3p.

DOCKET-50364- (Joseph M. Farley Nuclear Plant, Units 1 and 2, See also DOCKET-50348) (Non-AEC)

36	26-47519	See DOCKET-50348-42
----	----------	---------------------

DOCKET-50366- (Edwin I. Hatch Nuclear Plant, Unit 2) (Non-AEC)

29	26-47528	Dep. NTIS, 12p.
31	26-47529	Dep. NTIS, 182p.
32	26-47530	Dep. NTIS, 12p.
33	26-47531	Dep. NTIS, 6p.

DOCKET-50368- (Arkansas Nuclear One, Unit 2) (Non-AEC)

31	26-47532	Dep. NTIS, 210p.
----	----------	------------------

DOCKET-50369- (McGuire Nuclear Station, Units 1 and 2; See also DOCKET-50370) (Non-AEC)

26	26-47533	Dep. NTIS, 7p.
28	26-47534	Dep. NTIS, 302p.

DOCKET-50370- (McGuire Nuclear Station, Units 1 and 2; See also DOCKET-50369) (Non-AEC)

26	26-47533	See DOCKET-50369-26
28	26-47534	See DOCKET-50369-28

DOCKET-50373- (LaSalle County Station, Units 1 and 2; See also DOCKET-50374) (Non-AEC)

24	26-47535	Dep. NTIS, 133p.
25	26-47536	Dep. NTIS, 87p.

DOCKET-50374- (LaSalle County Station, Units 1 and 2; See also DOCKET-50373) (Non-AEC)

23	26-47535	See DOCKET-50373-24
24	26-47536	See DOCKET-50373-25

DOCKET-50376- (Aguirre Nuclear Plant) (Non-AEC)

34	26-47537	Dep. NTIS, 125p.
----	----------	------------------

DOCKET-50382- (Waterford Steam Electric Station; See also DOCKET-50383) (Non-AEC)

25	26-47538	Dep. NTIS, 173p.
27	26-47539	Dep. NTIS, 87p.
28	26-47540	Dep. NTIS, 61p.
29	26-47541	Dep. NTIS, 27p.
30	26-47542	Dep. NTIS, 21p.
31	26-47543	Dep. NTIS, 75p.
32	26-47544	Dep. NTIS, 85p.

DOCKET-50387- (Susquehanna Steam Electric Station, See also DOCKET-50388) (Non-AEC)

19	26-47545	Dep. NTIS, 339p.
20	26-47546	Dep. NTIS, 365p.
21	26-47547	Dep. NTIS, 2p.

Report No. Abstract No. Availability
 22 26-47548 Dep. NTIS, 29p.
 23 26-47549 Dep. NTIS, 28p.

DOCKET-50388- (Susquehanna Steam Electric Station, See also
 DOCKET-50387) (Non-AEC)
 19 26-47545 See DOCKET-50387-19
 20 26-47546 See DOCKET-50387-20
 21 26-47547 See DOCKET-50387-21
 22 26-47548 See DOCKET-50387-22
 23 26-47549 See DOCKET-50387-23

DOCKET-50390- (Watts Bar Nuclear Plant; See also DOCKET-
 50391) (Non-AEC)
 17 26-47550 Dep. NTIS, 82p.
 18 26-47551 Dep. NTIS, 319p.
 19 26-47552 Dep. NTIS, 51p.

DOCKET-50391- (Watts Bar Nuclear Plant; See also DOCKET-
 50390) (Non-AEC)
 17 26-47550 See DOCKET-50390-17
 18 26-47551 See DOCKET-50390-18
 19 26-47552 See DOCKET-50390-19

DOCKET-50395- (Virgil C. Summer Nuclear Station) (Non-AEC)
 22 26-47553 Dep. NTIS, 494p.
 24 26-47554 Dep. NTIS, 97p.
 25 26-47555 Dep. NTIS, 45p.

DOCKET-50397- (Hanford No. 2 Nuclear Power Plant--NON-AEC)
 19 26-47556 Dep. NTIS, 465p.
 20 26-47557 Dep. NTIS, 46p.
 21 26-47558 Dep. NTIS, 138p.

DOCKET-50398- (Mendocino Power Plant, Units 1 and 2; See
 also DOCKET-50399) (Non-AEC)
 11 26-47559 Dep. NTIS, 257p.
 15 26-47560 Dep. NTIS, 22p.
 16 26-47561 Dep. NTIS, 313p.
 17 26-47562 Dep. NTIS, 16p.

DOCKET-50399- (Mendocino Power Plant, Units 1 and 2; See
 also DOCKET-50398) (Non-AEC)
 11 26-47559 See DOCKET-50398-11
 15 26-47560 See DOCKET-50398-15
 16 26-47561 See DOCKET-50398-16
 17 26-47562 See DOCKET-50398-17

DOCKET-50400- (Shearon Harris Nuclear Power Plant; See also
 DOCKET-50401, 50402, 50403) (Non-AEC)
 19 26-47563 Dep. NTIS, 640p.
 20 26-47564 Dep. NTIS, 105p.
 21 26-47565 Dep. NTIS, 350p.

DOCKET-50401- (Shearon Harris Nuclear Power Plant, Units 1,
 2, 3 and 4; See also DOCKET-50400, 50402,
 50403) (Non-AEC)
 19 26-47563 See DOCKET-50400-19
 20 26-47564 See DOCKET-50400-20
 21 26-47565 See DOCKET-50400-21

DOCKET-50402- (Shearon Harris Nuclear Power Plant, Units 1,
 2, 3, and 4; See also DOCKET-50400, 50401,
 50403) (Non-AEC)
 19 26-47563 See DOCKET-50400-19
 20 26-47564 See DOCKET-50400-20
 21 26-47565 See DOCKET-50400-21

DOCKET-50403- (Shearon Harris Nuclear Power Plant, Units 1,
 2, 3, and 4; See also DOCKET-50400, 50401,
 50402) (Non-AEC)
 19 26-47563 See DOCKET-50400-19
 20 26-47564 See DOCKET-50400-20
 21 26-47565 See DOCKET-50400-21

DOCKET-50404- (North Anna Power Station, Units 3 and 4; See
 also DOCKET-50405) (Non-AEC)
 17 26-47510 See DOCKET-50338-47
 18 26-47566 Dep. NTIS, 125p.
 19 26-47567 Dep. NTIS, 19p.
 20 26-47568 Dep. NTIS, 9p.

DOCKET-50405- (North Anna Power Station, Units 3 and 4; See
 also DOCKET-50404) (Non-AEC)
 17 26-47510 See DOCKET-50338-47
 18 26-47566 See DOCKET-50404-18
 19 26-47567 See DOCKET-50404-19
 20 26-47568 See DOCKET-50404-20

DOCKET-50409- (LaCrosse Boiling Water Reactor--Formerly
 DOCKET-1155) (AEC)
 4 26-47569 Dep. NTIS, 3p.
 5 26-47570 Dep. NTIS, 3p.
 6 26-47382 Dep. NTIS, 9p.
 7 26-47330 Dep. NTIS, 2p.
 8 26-47383 Dep. NTIS, 5p.
 9 26-47384 Dep. NTIS, 31p.

DP-
 1300 26-45701 Dep. NTIS, 19p.

DPC-
 72-5 26-47382 See DOCKET-50409-6

169 REPORTS

IS

Report No. Abstract No. Availability
 72-6 26-47382 See DOCKET-50409-6
 72-7 26-47383 See DOCKET-50409-8
 851-44 26-47384 See DOCKET-50409-9

E- (Misc.)
 5811 26-45425 See N-72-16724
 6584 26-45933 See N-72-15609
 6772 26-46943 See N-72-18705
 6830 26-45934 See N-72-18585

ENDF-
 166 26-47220 See ORNL-TM-3867
 167 26-47221 See ORNL-TM-3868
 168 26-47222 See ORNL-TM-3869
 169 26-47223 See ORNL-TM-3870
 170 26-47224 See ORNL-TM-3871

EUR- (European Atomic Energy Community)
 4725 26-45439 See KFK-1566
 4806 26-47597 Dep. NTIS (U. S. Sales Only). EUR
 B. Fr. 85.-, 62p.
 4822 26-45221 Dep. NTIS (U. S. Sales Only). EUR
 B. Fr. 40.-, 20p.
 4832 26-47616 Dep. NTIS (U. S. Sales Only). EUR
 FB 225.-, 172p.
 4841 26-45570 Dep. NTIS (U. S. Sales Only). EUR
 FB 125.-, 98p.

EURFNR- (European Atomic Energy Community)
 1008 26-47585 Dep. NTIS (U. S. Sales Only), 34p.

FEI- (Russian)
 296 26-45617 Dep. NTIS (U. S. Sales Only), 30p.

FRNC-CONF- (French)
 29 26-45660 Dep. NTIS (U. S. Sales Only), 15p.

FRNC-R- (French)
 7 26-45616 Dep. NTIS (U. S. Sales Only), 58p.

FRNC-TH- (French)
 184 26-47014 Dep. NTIS (U. S. Sales Only), 70p.
 198 26-46335 Dep. NTIS (U. S. Sales Only), 100p.
 207 26-46974 Dep. NTIS (U. S. Sales Only), 87p.

GESP- (Non-AEC)
 493 26-45424 See N-72-15607

GULF-GA-A-
 12035 26-46077 Dep. NTIS, 27p.

GULF-GA-B-
 10959 26-47319 Dep. NTIS, 40p.

HASL-
 257 26-45511 Dep. NTIS, 201p.
 257(App.) 26-45512 Dep. NTIS, 437p.

HEDL-TME-
 71-126 26-46067 Dep. NTIS, 25p.
 72-42 26-47586 Dep. NTIS, 68p.
 72-81 26-45930 Dep. NTIS, 41p.

IC- (Italian)
 72/45 26-47284 Dep. NTIS (U. S. Sales Only), 14p.
 72/49 26-46666 Dep. NTIS (U. S. Sales Only), 20p.
 72/54 26-46570 Dep. NTIS (U. S. Sales Only), 14p.

ICP-
 1006 26-45405 Dep. NTIS, 82p.
 1008 26-45383 Dep. NTIS, 32p.
 1014 26-46093 Dep. NTIS, 33p.

ICRU- (Non-AEC)
 21 26-45618 ICRU Publications, Washington, D.
 C. \$4.00
 22 26-45261 ICRU Publications, Washington, D.
 C. \$3.50

IEA- (Brazilian)
 19 26-47617 Dep. NTIS (U. S. Sales Only), 50p.
 183 26-45365 Dep. NTIS (U. S. Sales Only), 28p.

IFVE-STF- (Russian)
 71-15 26-46617 Dep. NTIS (U. S. Sales Only), 21p.
 71-36 26-46497 Dep. NTIS (U. S. Sales Only), 11p.
 71-71 26-46618 Dep. NTIS (U. S. Sales Only), 8p.

INDC-
 7-U 26-47015

INIS-mf- (International Atomic Energy Agency)
 256 26-45709 INIS

IPP-IV- (German)
 12 26-47116 Dep. NTIS (U. S. Sales Only), 128p.
 32 26-47170 Dep. NTIS (U. S. Sales Only), 58p.

IS-

IS			REPORTS 170	Report No.	Abstract No.	Availability
Report No.	Abstract No.	Availability		450	26-45932	Dep. NTIS, 143p.
2886	26-46475	Dep. NTIS, 9p.		659	26-45992	Dep. NTIS, 98p.
				773	26-46521	Dep. NTIS, 93p.
				791	26-46448	Dep. NTIS, 25p.
ITEF-	(Russian)			LYCEN-	(French)	
848	26-46518	Dep. NTIS (U. S. Sales Only), 23p.		7186	26-46860	Dep. NTIS (U. S. Sales Only), 55p.
				7202	26-47286	Dep. NTIS (U. S. Sales Only), 27p.
ITF-	(Russian)			MIT-	(Misc.)	
71-101-P	26-47285	Dep. NTIS (U. S. Sales Only), 44p.		144	26-47620	
71-117-P	26-46667	Dep. NTIS (U. S. Sales Only), 14p.				
71-128-P	26-46698	Dep. NTIS (U. S. Sales Only), 28p.		MLM-		
72-13-P	26-46609	Dep. NTIS (U. S. Sales Only), 7p.		1911	26-46114	Dep. NTIS, 45p.
72-44-E	26-46774	Dep. NTIS (U. S. Sales Only), 11p.		1913	26-45286	Dep. NTIS, 205p.
				1936(TR)	26-45688	Dep. NTIS
JAB-	(Non-AEC)			MSAR-	(Misc.)	
99-88	26-45474	Dep. NTIS, 36p.		71-45	26-45440	Dep. NTIS, 163p.
JAERI-M-				N-	(Misc.)	
4594	26-45182	NTIS (U. S. Sales Only), 211p.		72-15068	26-45392	NTIS, 45p.
JEN-	(Spanish)			72-15607	26-45424	NTIS, 526p.
221	26-46846	Dep. NTIS (U. S. Sales Only), 170p.		72-15609	26-45933	NTIS, 31p.
229-DQ/I-79	26-45222	Dep. NTIS (U. S. Sales Only), 18p.		72-16724	26-45425	NTIS, 29p.
230-DME/I-	26-46918	Dep. NTIS (U. S. Sales Only), 35p.		72-17158	26-45661	NTIS, 31p.
25				72-17763	26-46522	NTIS, 30p.
232-DQ/I-81	26-45223	Dep. NTIS (U. S. Sales Only), 19p.		72-18362	26-46301	NTIS, 35p.
247	26-45589	Dep. NTIS (U. S. Sales Only), 47p.		72-18585	26-45934	NTIS, 33p.
JINR-	(Russian)			72-18705	26-46943	NTIS, 9p.
P1-6408	26-46476	Dep. NTIS (U. S. Sales Only), 13p.		72-18715	26-46265	NTIS, 208p.
P1-6461	26-46619	Dep. NTIS (U. S. Sales Only), 7p.		72-18796	26-46324	NTIS, 37p.
P2-4302	26-47038	For translation see BNL-tr-487		72-18797	26-47203	NTIS, 16p.
P2-5924	26-46478	For translation see ORNL-tr-2544		72-19766	26-46704	NTIS, 34p.
P2-6270	26-46748	Dep. NTIS (U. S. Sales Only), 13p.		NASA-CR-	(Misc.)	
P2-6333	26-46519	Dep. NTIS (U. S. Sales Only), 26p.		113925	26-45424	See N-72-15607
P2-6393	26-46775	Dep. NTIS (U. S. Sales Only), 18p.		122351	26-46265	See N-72-18715
P2-6394	26-46699	Dep. NTIS (U. S. Sales Only), 16p.		125673	26-46324	See N-72-18796
P2-6395	26-46649	Dep. NTIS (U. S. Sales Only), 15p.		NASA-TM-X-	(Misc.)	
P2-6437	26-46700	Dep. NTIS (U. S. Sales Only), 13p.		2237	26-45425	See N-72-16724
P2-6438	26-46520	Dep. NTIS (U. S. Sales Only), 24p.		2493	26-45933	See N-72-15609
P2-6442	26-46776	Dep. NTIS (U. S. Sales Only), 15p.		65821	26-46522	See N-72-17763
P2-6454	26-46701	Dep. NTIS (U. S. Sales Only), 47p.		65832	26-46301	See N-72-18362
P2-6474	26-46702	Dep. NTIS (U. S. Sales Only), 11p.		65851	26-46704	See N-72-19766
P2-6482	26-46703	Dep. NTIS (U. S. Sales Only), 11p.		67594	26-45392	See N-72-15068
P2-6501	26-46586	Dep. NTIS (U. S. Sales Only), 14p.		68004	26-46943	See N-72-18705
P2-6503	26-47047	Dep. NTIS (U. S. Sales Only), 9p.		68024	26-45934	See N-72-18585
P4-6452	26-47048	Dep. NTIS (U. S. Sales Only), 15p.		NASA-TN-D-	(Misc.)	
P4-6458	26-47049	Dep. NTIS (U. S. Sales Only), 16p.		6631	26-45661	See N-72-17158
P4-6462	26-46975	Dep. NTIS (U. S. Sales Only), 18p.		6695	26-47203	See N-72-18797
P4-6475	26-46620	Dep. NTIS (U. S. Sales Only), 8p.		NIM-	(Non-AEC)	
P6-6402	26-45366	Dep. NTIS (U. S. Sales Only), 15p.		1427	26-45262	Dep. NTIS (U. S. Sales Only), 8p.
P6-6492	26-46976	Dep. NTIS (U. S. Sales Only), 20p.		NP-	(AEC File No. for Non-AEC Reports)	
P6-6512	26-46942	Dep. NTIS (U. S. Sales Only), 28p.		11592 (Vol. 1)	26-46449	For translation see SLAC-Trans-143
P7-6466	26-47022	Dep. NTIS (U. S. Sales Only), 15p.		19256	26-46431	Dep. NTIS (U. S. Sales Only), 13p.
P9-6421	26-46136	Dep. NTIS (U. S. Sales Only), 13p.		19357	26-45520	Dep. NTIS (U. S. Sales Only), 11p.
P9-6514	26-46137	Dep. NTIS (U. S. Sales Only), 18p.		19360	26-47652	Philadelphia; National Federation of Science Abstracting and Indexing Services (1972). 28p
P13-6428	26-45637	Dep. NTIS (U. S. Sales Only), 23p.		19361	26-45475	
P14-6407	26-47618	Dep. NTIS (U. S. Sales Only), 5p.		19364	26-47153	Dep. NTIS (U. S. Sales Only), 86p.
JUL-	(German)			NP-tr-	(AEC File No. for Non-AEC Translations)	
834-RG	26-46874	Dep. NTIS (U. S. Sales Only), 101p.		1941	26-45778	Dep. NTIS (U. S. Sales Only)
KAPL-				NVO-		
3164	26-45931	Dep. NTIS, 8p.		38-33	26-45476	Dep. NTIS, 88p.
KFK-	(German)			55	26-45571	Dep. NTIS, 31p.
1492	26-45951	Elektrotech. Z., A; 92: No. 8, 486-8(Aug 1971)		86(Rev.1)	26-45477	Dep. NTIS, 33p.
1509	26-46457	Dep. NTIS (U. S. Sales Only), 77p.		409-1	26-45478	Dep. NTIS, 135p.
1537	26-47023	Dep. NTIS (U. S. Sales Only), 54p.		746-122	26-45479	Dep. NTIS, 36p.
1547	26-46650	Dep. NTIS (U. S. Sales Only), 20p.		1163-TM-32	26-45480	Dep. NTIS, 78p.
1556	26-45438	Dep. NTIS (U. S. Sales Only), 29p.		1163-TM-33	26-45481	Dep. NTIS, 72p.
1560	26-45509	Dep. NTIS (U. S. Sales Only), 26p.		1229-177	26-45482	Dep. NTIS, 21p.
1561	26-47585	See EURFNR-1008		1229-180	26-45483	Dep. NTIS, 91p.
1563	26-47024	Dep. NTIS (U. S. Sales Only), 71p.		NYO-		
1566	26-45439	Dep. NTIS (U. S. Sales Only), 32p.		1710-106	26-46078	Dep. NTIS, 28p.
KFK-EXT-	(West German)			2174-124	26-45533	J. Mar. Res.; 30: No. 1, 112-20(5 Jan 1972)
10/72-1	26-46875	Dep. NTIS (U. S. Sales Only), 51p.		2174-130	26-45541	Dep. NTIS, 19p.
KFKI-	(Hungarian)			3038-13	26-46150	Dep. NTIS, 25p.
71-47	26-46118	Dep. NTIS (U. S. Sales Only), 122p.		3144-72	26-47343	See DOCKET-50213-154
72-15	26-46571	Dep. NTIS (U. S. Sales Only), 11p.		3970-29	26-45890	Dep. NTIS, 71p.
KY-				4076-17	26-46523	Dep. NTIS, 13p.
642	26-47619	Dep. NTIS, 10p.		4076-18	26-46524	Dep. NTIS, 23p.
LA-				4076-19	26-46681	Dep. NTIS, 14p.
4978	26-45702	Dep. NTIS, 15p.		ORNL-		
5000	26-46119	Dep. NTIS, 68p.		4778	26-47287	Dep. NTIS, 538p.
LA-DC-				4796	26-45703	Dep. NTIS, 38p.
72-524	26-45224	Dep. NTIS, 18p.		ORNL-RMIC-		
72-556	26-46104	Dep. NTIS, 7p.		12	26-45973	Dep. NTIS, 142p.
72-703	26-47141	Dep. NTIS, 4p.		ORNL-TM-		
72-706	26-47156	Dep. NTIS, 4p.				
72-707	26-47157	Dep. NTIS, 4p.				
72-726	26-47158	Dep. NTIS, 4p.				
LBL-						
378	26-46477	Dep. NTIS, 172p.				

Report No.	Abstract No.	Availability
2871	26-45406	Dep. NTIS, 102p.
2918	26-45407	Dep. NTIS, 89p.
2949	26-45408	Dep. NTIS, 97p.
2996	26-45409	Dep. NTIS, 82p.
3018	26-45410	Dep. NTIS, 94p.
3072	26-45411	Dep. NTIS, 94p.
3095	26-45412	Dep. NTIS, 76p.
3127	26-45413	Dep. NTIS, 78p.
3180	26-45414	Dep. NTIS, 92p.
3217	26-45415	Dep. NTIS, 88p.
3250	26-45416	Dep. NTIS, 93p.
3281	26-45417	Dep. NTIS, 85p.
3733	26-46068	Dep. NTIS, 30p.
3810	26-47621	Dep. NTIS, 38p.
3820	26-46340	Dep. NTIS, 103p.
3832	26-47305	Dep. NTIS, 69p.
3839	26-46364	Dep. NTIS, 87p.
3859	26-46876	Dep. NTIS, 60p.
3867	26-47220	Dep. NTIS, 20p.
3868	26-47221	Dep. NTIS, 18p.
3869	26-47222	Dep. NTIS, 18p.
3870	26-47223	Dep. NTIS, 18p.
3871	26-47224	Dep. NTIS, 26p.

ORNL-tr-

2369	26-46102	Dep. NTIS
2371	26-46105	Dep. NTIS
2372	26-46106	Dep. NTIS
2377	26-45993	Dep. NTIS
2532	26-45590	Dep. NTIS
2544	26-46478	Dep. NTIS
2617	26-46126	Dep. NTIS

ORO-

694	26-45689	Dep. NTIS, 68p.
2690-12(Pt. B)	26-45891	Dep. NTIS, 75p.
2911-14	26-46365	Dep. NTIS, 27p.
3065-24	26-46479	Dep. NTIS, 6p.
3728-5	26-45912	Dep. NTIS, 8p.
3909-3	26-45316	Dep. NTIS, 84p.
3992-80	26-46525	Dep. NTIS, 25p.
3992-94	26-46749	Dep. NTIS, 7p.
4245-4	26-47622	Dep. NTIS, 8p.

PATENTS

(British)

1,269,297 26-45355

PATENTS

(Canadian)

891,958	26-45371
892,730	26-45336
893,684	26-45690
893,931	26-45267
895,492	26-45692
897,930	26-45719
899,029	26-45274

PATENTS

(French)

2,067,743	26-45384
2,070,479	26-45658
2,070,481	26-45243
2,070,925	26-45656
2,076,863	26-45339
2,079,708	26-45696
2,081,129	26-46111
2,081,176	26-45422
2,082,274	26-45697
2,082,735	26-45426
2,083,752	26-45609
2,088,028	26-46115
2,088,094	26-46109
2,088,170	26-45940
2,088,610	26-45289
2,088,806	26-46112
2,088,972	26-45472

PATENTS

(Indian)

114,500 26-45941

PATENTS

(U. S.)

3,657,137	26-45957
3,657,920	26-45613
3,658,467	26-45387
3,659,107	26-45708
3,662,172	26-45562
3,662,173	26-45559
3,662,179	26-45563
3,662,589	26-45469
3,662,718	26-45450
3,663,178	26-45451
3,663,360	26-45433
3,663,361	26-47177
3,663,362	26-47174
3,663,504	26-45672
3,663,815	26-46083
3,663,816	26-46110
3,663,817	26-46103
3,664,859	26-46088
3,664,920	26-47113
3,664,921	26-47175
3,665,183	26-45446
3,665,195	26-45561

171 REPORTS

SCL-RR

Report No.	Abstract No.	Availability
------------	--------------	--------------

3,665,351	26-46843	
3,665,573	26-45449	
3,666,560	26-45434	
3,666,673	26-45386	
3,666,846	26-45707	
3,667,029	26-46842	
3,667,059	26-47178	
3,667,068	26-45677	
3,668,386	26-45655	
3,668,392	26-45659	
3,668,395	26-45596	
3,668,396	26-45595	
3,668,398	26-45654	
3,668,400	26-45597	
3,669,796	26-46089	
3,669,832	26-45959	
3,671,741	26-45456	
3,672,443	26-45427	
3,672,848	26-45420	
3,672,875	26-45423	
3,673,409	26-45457	
3,673,411	26-46113	
3,674,655	26-45980	
3,675,021	26-45958	
3,676,192	26-45352	
3,676,673	26-45448	
3,676,675	26-45447	
3,676,682	26-45635	

PIB-

(West Germany)

1-157 26-46431 See NP-19256

PNE-

529 26-45542 Dep. NTIS, 67p.

PNE-RB-

26 26-45476 See NVO-38-33

PNE-WW-

2 26-45571 See NVO-55

PRDC-EF-

104 26-47339 See DOCKET-50016-164

R-

(Misc.)
72-8 26-47623 See RLO-2229-T-1-18
72-10 26-47620 See MIT-144

RD/B/N-

(British)
2291 26-45905 Dep. NTIS (U. S. Sales Only), 38p.

RDT-E-

14-6-T(6-72) 26-47593 RSO

RISO-

(Danish)
254 26-46362 Dep. NTIS (U. S. Sales Only), 115p.

RLO-

2225-T-1-9	26-45543	Dep. NTIS, 15p.
2225-T-14-4	26-45544	Dep. NTIS, 29p.
2227-T-8-3	26-45913	Dep. NTIS, 17p.
2229-T-1-18	26-47623	Dep. NTIS, 136p.
2230-T-4-1	26-46587	Dep. NTIS, 12p.

RPP/H-

(British)
89 26-45646 Dep. NTIS (U. S. Sales Only), 27p.

RS-

(Misc.)
34/66 26-45520 See NP-19357

RT/PROT-

(Italian)
72-4 26-47318 Dep. NTIS (U. S. Sales Only), 33p.
72-11 26-45263 Dep. NTIS (U. S. Sales Only), 27p.

SC-CR-

723142 26-47624 Dep. NTIS, 84p.

SC-DC-

721232	26-45662	Dep. NTIS, 19p.
721233	26-45663	Dep. NTIS, 11p.
721237	26-45664	Dep. NTIS, 25p.
721240	26-45665	Dep. NTIS, 34p.
722267	26-47625	Dep. NTIS, 6p.
722296	26-45484	Dep. NTIS, 28p.

SC-R-

722769 26-45287 Dep. NTIS \$12.00, 1149p.

SC-RR-

710884	26-45935	Dep. NTIS, 27p.
720359	26-45485	Dep. NTIS, 19p.
720515	26-45666	Dep. NTIS, 34p.

SC-T-

722519	26-45667	Dep. NTIS
722520	26-47626	Dep. NTIS

SCL-RR-

Report No. Abstract No. Availability
720270 26-47627 Dep. NTIS, 23p.

SLAC-Trans-

142 26-46432 Dep. NTIS
143 26-46449 Dep. NTIS

SM-

(Misc.)
158/16 26-45543 See RLO-2225-T-1-9
158/47 26-45544 See RLO-2225-T-14-4

SRARI-P-

(Russian)
120 26-45441 Dep. NTIS (U. S. Sales Only), 24p.
123 26-45264 Dep. NTIS (U. S. Sales Only), 28p.

SZS-

(German)
17/69 26-46102 For translation see ORNL-tr-2369

TID-

3767 26-47628 TIC Free
7029 (2nd Ed.) 26-45183 Dep. NTIS, 440p.
25954 26-45755 Dep. NTIS, 446p.
26078 26-45873 Dep. NTIS, 14p.
26103 26-45527 Dep. NTIS, 75p.
26106 26-45288 Dep. NTIS, 44p.

TM-

(Misc.)
72-2015-1 26-46324 See N-72-18796

TPJU-

(Polish)
14/71 26-46526 Dep. NTIS (U. S. Sales Only), 11p.

TT-

(Misc.)
54002/4 26-45779 NTIS
54002/5 26-45785 NTIS

UCID-

30047 26-47598 Dep. NTIS, 301p.

UCLA-

12-866 26-47629 Dep. NTIS, 15p.
12-867 26-45818 Dep. NTIS, 19p.
34-P-177-10 26-45994 Dep. NTIS, 13p.
34-P-177-11 26-47630 Dep. NTIS, 4p.

UCRL-

50007-71-3 26-45673 Dep. NTIS, 27p.
51061 (Vol. 1) 26-45265 Dep. NTIS, 78p.
51210 26-47631 Dep. NTIS, 262p.
51214 26-45647 Dep. NTIS, 27p.
51227 26-47632 Dep. NTIS, 40p.
51231 26-46341 Dep. NTIS, 62p.
51235 26-45486 Dep. NTIS, 10p.
51239 26-47633 Dep. NTIS, 9p.
51242 26-45513 Dep. NTIS, 47p.

Report No. Abstract No. Availability

73587 26-47634 Dep. NTIS, 19p.
73665 26-45674 Dep. NTIS, 12p.
73744 26-45545 Dep. NTIS, 35p.
73756 26-46107 Dep. NTIS, 10p.
73757 26-46108 Dep. NTIS, 11p.
73805 26-47635 Dep. NTIS, 11p.
73925 26-47636 Dep. NTIS, 19p.
73939 26-45487 Dep. NTIS, 36p.
73974 26-45668 Dep. NTIS, 55p.

UCRL-Trans-

10622 26-46079 Dep. NTIS
10630 26-47159 Dep. NTIS
10632 26-45936 Dep. NTIS

UJV-

(Czechoslovakian)
2795-M 26-46069 Dep. NTIS (U. S. Sales Only), 11p.
2798-M 26-46070 Dep. NTIS (U. S. Sales Only), 25p.
2802-M 26-45995 Dep. NTIS (U. S. Sales Only), 46p.
2820-Ch 26-45273 Dep. NTIS (U. S. Sales Only), 29p.

UR-

49-1508 26-47637 Dep. NTIS, 36p.
3055-1 26-47638 Dep. NTIS, 157p.
3055-2 26-47639 Dep. NTIS, 98p.
3490-18 26-47640 Dep. NTIS, 105p.
3490-21 26-47641 Dep. NTIS, 13p.
3490-47 26-47642 Dep. NTIS, 144p.
3490-63 26-45819 Dep. NTIS, 16p.
3490-80 26-47643 Dep. NTIS, 13p.
3490-85 26-47644 Dep. NTIS, 38p.
3490-102 26-45895 Dep. NTIS, 25p.
3490-107 26-47645 Dep. NTIS, 27p.
3490-115 26-45915 Dep. NTIS, 33p.
3490-121 26-45820 Dep. NTIS, 110p.
3490-131 26-47646 Dep. NTIS, 59p.
3490-132 26-47647 Dep. NTIS, 88p.
3490-137 26-47648 Dep. NTIS, 114p.
3490-147 26-47649 Dep. NTIS, 14p.
3490-153 26-47650 Dep. NTIS, 12p.
3490-154 26-47651 Dep. NTIS, 26p.
3490-159 26-45916 Dep. NTIS, 20p.

USGS-

474-148 26-45488 Dep. NTIS, 87p.

WASH-

1210 26-46094 Dep. GPO \$1.75
1523 26-45572 Dep. NTIS, 130p.

WCAP-

(Misc.)
7841 26-45996 Dep. NTIS, 24p.

X-

(Misc.)
621-72-44 26-46301 See N-72-18362
641-72-22 26-46522 See N-72-17763
641-72-73 26-46704 See N-72-19766

Appendix I

JOURNAL LIST

This list of journals supplements the listing which appeared as Appendix II in Volume 26, Number 1. The journals are arranged alphabetically by abbreviation.

The title abbreviations are based on the National Clearinghouse for Periodical Title Word Abbreviations, *Word Abbreviation List* published by the Standards Committee Z-39 of the American National Standards Institute.

TID-4579 (6th Ed.) *Serial Titles Cited in Nuclear Science Abstracts* lists the titles and abbreviations of the journals regularly scanned for NSA as well as those from which an occasional item is taken. This publication is available from the National Technical Information Service (NTIS), U. S. Department of Commerce, Springfield, Virginia, 22151, for \$6.00 for full size or reproduction copy and \$0.95 for microfiche copy.

A

Accident Anal. Prev. (Elmsford, N.Y.)
Accident Analysis and Prevention (Elmsford, New York) (U.S.)

Acta Radiobot. Genet.
Acta Radiobotanica et Genetica (Japan)

Acta Univ. Agr. (Brno), Ser. C
Acta Univ. Agr. (Brno), Series C (Facultas Silviculturae)
(Czechoslovakia)

Acta Univ. Upsal., Abstr. Uppsala Diss. Fac. Sci.
Acta Universitatis Upsaliensis, Abstracts of Uppsala
Dissertations from the Faculty of Science (Sweden)
Supersedes Acta Univ. Upsal., Abstr. Uppsala Diss. Sci.

Advan. Exp. Med. Biol.
Advances in Experimental Medicine and Biology (U.S.)

Advan. Instrum.
Advances in Instrumentation (U.S.)

Aerotec.-Missili Spazio
Aerotecnica-Missili e Spazio (Italy)
Formed by merger of Aerotecnica and Missili e Spazio

Ag Chem Commer. Fert.
Ag Chem and Commercial Fertilizer (U.S.)
Supersedes Agr. Chem.

Agra Univ. J. Res.
Agra University Journal of Research (India)

AIP (Amer. Inst. Phys.) Conf. Proc.
AIP (American Institute of Physics) Conference Proceedings
(U.S.)

Alimenta (Zurich)
Alimenta (Zurich) (Switzerland)

Allianz Ber. Betriebstech. Schadenverhuetzung
Allianz Berichte fuer Betriebstechnik und
Schadenverhuetzung (West Germany)

Amer. Chem. Soc., Div. Water, Air Waste Chem., Gen. Pap.
American Chemical Society, Division of Water, Air and Waste
Chemistry, General Papers (U.S.)

Amer. Fam. Physician
American Family Physician (U.S.)
Supersedes American Family Physician - GP

An. Asoc. Quim. Argent.
Anales de la Asociacion Quimica Argentine (Argentina)

Ann. Acad. Sci. Fenn., Ser. A III
Annales Academiae Scientiarum Fennicae, Series A III.
Geologica-Geographica (Finland)

Ann. Acad. Sci. Fenn., Ser. A V
Annales Academiae Scientiarum Fennicae, Series A V.
Medical (Finland)

Ann. Agron.
Annales Agronomiques (France)

Ann. Sci. Forest.
Annales des Sciences Forestieres (France)

Ann. Univ. Sarav., Math.-Naturwiss. Fak.
Annales Universitatis Saravienis, (Reihe), Mathematisch-
Naturwissenschaftliche Fakultät (West Germany)

Annu. Rep. Progr. Chem., Sect. A
Annual Reports on the Progress of Chemistry, Section A.
General, Physical and Inorganic Chemistry (England)

Annu. Rev. Mater. Sci.
Annual Review of Materials Science (U.S.)

An. Univ. Bucuresti, Chim.
Analele Universitatii Bucuresti, Chimie (Romania)

Appl. Elec. Phenomena (Engl. Transl.)
Applied Electrical Phenomena (English Translation of
Elektronnaya Obrabotka Materialov) (U.S.)

Appl. Solar Energy (USSR) (Engl. Transl.)
Applied Solar Energy (USSR) (English Translation of
Geliotekhnika) (U.S.)

App. Metody Rentgenovskogo Anal.
Apparatura i Metody Rentgenovskogo Analiza (U.S.S.R.)

Araneta J. Agr.
Araneta Journal of Agriculture (Philippines)

Arch. Dermatol. Forsch.
Archiv fuer Dermatologische Forschung (West Germany)
Supersedes Arch. Klin. Exp. Dermatol.

Arch. Fund. Roux-Ocefa
Archivos de la Fundacion Roux-Ocefa (Argentina)

Arch. Julius Klaus-Stift. Vererbungsforsch., Sozialanthropol.
Rassenhyg.
Archiv der Julius Klaus-Stiftung fuer Vererbungsforschung,
Sozialanthropologie und Rassenhygiene (Switzerland)

Arch. Sci. Biol. (Bologna)
Archivio di Scienze Biologiche (Bologna) (Italy)

Arctic
Arctic (U.S. - Canada)

Ark. Mat., Astron. Fys.
Arkiv for Matematik, Astronomi och Fysik (Sweden)

Astronautyka
Astronautyka (Poland)

At. Jap.
Atoms in Japan (Japan)

Aufbereit.-Tech.
Aufbereitungs-Technik (West Germany)

Aufschluss
Aufschluss (West Germany)

Aust. Elec. World
Australian Electrical World, The (Australia)

Automatica
Automatica (England - U.S.)

Awamia
Awamia, Al (Morocco)

B

Ber. Geol. Ges. Deut. Demokrat. Repub. Gesamtgeb. Geol. Wiss.
Berichte der Geologischen Gesellschaft in der Deutschen
Demokratischen Republik fuer das Gesamtgebiet der
Geologischen Wissenschaft (East Germany)
Divided into Ber. Deut. Ges. Geol. Wiss., Reihe A and
Ber. Deut. Ges. Geol. Wiss., Reihe B

Ber. Getreidechem.-Tag., Detmold
Bericht ueber die Getreidechemiker-Tagung, Detmold
(West Germany)

Beton, Herstellung, Verwend.
Beton, Herstellung, Verwendung (West Germany)

Biol. Conserv.
Biological Conservation (England)

Biol. Neonate
Biology of the Neonate (Switzerland)
Supersedes Biol. Neonatorum

Bitki Koruma Bul.
Bitki Koruma Bulteni (Plant Protection Bulletin) (Turkey)

Bitumen, Teere, Asphalte, Pech
Bitumen, Teere, Asphalte, Pech und Verwandte Stoffe
(West Germany)

Boll. Mal. Orecchio, Gola Naso
Bollettino delle Malattie dell'Orecchio, della
Gola e del Naso (Italy)

Boll. Soc. Geol. Ital.
Bollettino della Societa Geologica Italiana (Italy)

Boll. Unione Mat. Ital.
Bollettino dell'Unione Matematica Italiana (Italy)

Boll. Unione Mat. Ital., Ser. IV
Bollettino delle Unione Matematica Italiana, Series IV
(Italy)

Bot. Rev.
Botanical Review, The (U.S.)

Brit. Chem. Eng. Process Technol.
British Chemical Engineering and Process Technology
(England)
Supersedes British Chemical Engineering

Brit. Engine Tech. Rep.
British Engine Technical Reports (England)

Bull. Ass. Anciens Eleves Inst. Ind. Ferment. Bruxelles
Bulletin de l'Association des Anciens Eleves de
l'Institut des Industries de Fermentation de Burxelles
(Belgium)

Bull. Bot. Soc. Bengal
Bulletin of the Botanical Society of Bengal (India)

Bull. Doc., Cent. Inform. Chrome Dur
Bulletin de Documentation, Centre d'Information du
Chrome Dur (France)

Bull. Grain Technol.
Bulletin of Grain Technology (India)

Bull. Inform. Generateurs Isotop.
Bulletin d'Information sur les Generateurs Isotopiques
(France)
Issued in English version as Information Bulletin
on Isotopic Generators

Bull. Sci., Cons. Acad. Sci. Arts RSF Yougoslavie, Sect. A
Bulletin Scientifique, Conseil des Academies des Sciences
et des Arts de la RSF de Yougoslavie, Section A. Sciences
Naturelles, Techniques et Medicales (Yugoslavia)

Byull. Inst. Astrofiz., Akad. Nauk Tadzh. SSR
Byulleten Instituta Astrofiziki, Akademiya Nauk
Tadzhikskoi SSR (Tadzhik S.S.R.)

Byull. Nauch.-Tekh. Inform. Agron. Fiz.
Byulleten Nauchno-Tekhnicheskoi Informatsii po
Agronomicheskoi Fizike (U.S.S.R.)

Byull. Tsentra Yadern. Dannym
Byulleten Tsentra po Yadernym Dannym (U.S.S.R.)

C

Can. Geogr.
Canadian Geographer (Canada)

Can. J. Pharm. Sci.
Canadian Journal of Pharmaceutical Sciences (Canada)

Can. J. Spectrosc.
Canadian Journal of Spectroscopy (Canada)
Supersedes Can. Spectrosc.

Cardiology
Cardiology (Switzerland)
Supersedes Cardiologia

Cell. Immunol.
Cellular Immunology (U.S.)

Cem. Concr. (Delhi)
Cement and Concrete (Delhi)

Chagyo Gijutsu Kenkyu
Chagyo Gijutsu Kenkyu (Study of Tea) (Japan)

Chem. Chron., Epistem. Ekdosis
Chemika Chronika, Epistemonike Ekdosis (Greece)

Chem. Stosow., Ser. B
Chemia Stosowana, Seria B. Kwartalnik Poswiecony
Zagadnieniom Inzynierii i Aparatury Chemicznej
(Poland)

Chem.-Ztg., Chem. App.
Chemiker-Zeitung, Chemische Apparatur (West Germany)
Supersedes Chemiker-Zeitung

Chislennyye Metody Mekh. Sploshnoi Sredy
Chislennyye Metody Mekhaniki Sploshnoi Sredy (U.S.S.R.)

Clin. Otorinolaringoiat.
Clinica Otorinolaringoiatrica (Italy)

Collect. Treatises Fac. Hum., Univ. Fukuoka
Collection of Treatises Published by Faculty of Humanity,
University of Fukuoka (Japan)

Commentat. Biol.
Commentationes Biologicae, Societas Scientiarum Fennica
(Finland)

Comments Plasma Phys. Controlled Fusion
Comments on Plasma Physics and Controlled Fusion (England)

Computing
Computing (Austria)

Confructa
Confructa (West Germany)

Contam. Contr.
Contamination Control (U.S.)

Cryog. Eng. (Tokyo)
Cryogenic Engineering (Tokyo) (Japan)

Curr. Top. Radiat. Res. Quart.
Current Topics in Radiation Research Quarterly (Netherlands)
Supersedes Curr. Top. Radiat. Res.

Cutting Tool Eng.
Cutting Tool Engineering (U.S.)

D

Dainichi Nippon Densen Jiho
Dainichi Nippon Densen Jiho (Dainichi-Nippon Cables
Review) (Japan)

Delt. Hellen. Mikrobiol. Hyg. Hetair.
Deltion Hellenikes Mikrobiologikes kai Hygieinologikes
Hetaireias (Greece)

Demag Nachr.
Demag Nachrichten (West Germany)

Denki Keisan
Denki Keisan (Electrical Calculaton) (Japan)

Denshi Gijutsu Sogo Kenkyusho Chosa Hokoku
Denshi Gijutsu Sogo Kenkyusho Chosa Hokoku (Japan)

Dermatol. Monatsschr.
Dermatologische Monatsschrift (East Germany)

Doboku Gijutsu Shiryo
Doboku Gijutsu Shiryo (Public Works Review) (Japan)

Draht
Draht (West Germany)

Draht-Welt
Draht-Welt (West Germany)

Drug Ther. Bull.
Drug and Therapeutics Bulletin (England)

E

Earth Sci. Bull.
Earth Science Bulletin (U.S.)

Ecology (USSR) (Engl. Transl.)
Ecology (USSR) (English Translation of Ekologiya) (U.S.)

Econ. Int. Fonti Energ.
Economia Internazionale delle Fonti di Energia (Italy)

Econ. Trentina
Economia Trentina (Italy)

Eisenbahn-Ing.
Eisenbahn-Ingenieur (West Germany)

Ekologiya
Ekologiya (U.S.S.R.)

Elec. News Eng.
Electrical News and Engineering (Canada)

Elektro-Anz.
Elektro-Anzeiger (West Germany)

Elektronik (Munich)
Elektronik (Munich) (West Germany)

Elektron.-Ztg.
Elektronik-Zeitung (West Germany)

Elelmiszervizsgalati Kozlem.
Elelmiszervizsgalati Kozlemenyek (Hungary)

Energ. Elek.
Energetika i Elektrifikatsiya (Ukrainian S.S.R.)

Energiewirt. Tagesfragen
Energiewirtschaftliche Tagesfragen (West Germany)

Energ. Stroit.
Energeticheskoe Stroitel'stvo (U.S.S.R.)

Eng. Cybern. (USSR) (Engl. Transl.)
Engineering Cybernetics (USSR) (English Translation of
Izvestiya Akademii Nauk SSSR, Tekhnicheskaya Kibernetika)
(U.S.)

Environ. Aff.
Environmental Affairs (U.S.)

Environ. Contr. Saf. Manage.
Environmental Control and Safety Management (U.S.)

Environ. Physiol.
Environmental Physiology (Denmark)

Enzyme
Enzyme (Switzerland)
Supersedes Enzymol. Biol. Clin.

EOS, Trans., Amer. Geophys. Union
EOS, Transactions, American Geophysical Union (U.S.)
Supersedes Trans. Amer. Geophys. Union

Eur. J. Clin. Pharm.
European Journal of Clinical Pharmacology Continues
Pharmacol. Clin.

Europplast. Mon.
Europlastics Monthly (England)
Supersedes British Plastics

Eur. S. Observ., Bull.
European Southern Observatory, Bulletin (West Germany)

F

Farmacia (Bucharest)
Farmacia (Bucharest) (Romania)

Farm J. (Calcutta)
Farm Journal (Calcutta) (India)

Fertigung
Fertigung (Switzerland)

Festkoerperprobleme
Festkoerperprobleme (West Germany)

Fiz. Elem. Chastits At. Yadra
Fizika Elementarnykh Chastits i Atomnogo Yadra (U.S.S.R.)

Fiz. Szk.
Fizyka w Szkole (Poland)

Folia Forest. (Helsinki)
Folia Forestalia (Helsinki) (Finland)

G

Galvano
Galvano (France)

Gartenbauwissenschaft
Gartenbauwissenschaft (West Germany)

Geliotekhnika
Geliotekhnika (Uzbek S.S.R.)
Issued in English translation as Applied Solar Energy (USSR)

Geokhim. Sb.
Geokhimicheskii Sbornik (U.S.S.R.)

Geol. Tec.
Geologia Tecnica (Italy)

Gesuido Kyokai-Shi
Gesuido Kyokai-Shi (Journal of Japan Sewage Works Association) (Japan)

Gidromekhanika
Gidromekhanika (Ukrainian S.S.R.)

Glastech. Ber.
Glastechnische Berichte (West Germany)

God. Sofii. Univ., Biol. Fak.
Godishnik na Sofiiskiya Universitet, Biologicheski Fakultet (Bulgaria)

God. Sofii. Univ., Fiz. Fak.
Godishnik na Sofiiskiya Universitet, Fizicheski Fakultet (Bulgaria)

God. Zb. Med. Fak. Skopje
Godisen Zbornik na Medicinskiot Fakultet vo Skopje (Yugoslavia)

Gorskostop. Nauka
Gorskestopanska Nauka (Yugoslavia)

Gravitatsiya Teor. Otnositel'nosti
Gravitatsiya i Teoriya Otnositel'nosti (U.S.S.R.)

H

Haematol. Bluttransfus.
Haematologie und Bluttransfusion (Suppl. to Blut) (West Germany)

Han'guk Sikp'um Kwahakhoe Chi
Han'guk Sikp'um Kwahakhoe Chi (Journal of the Korean Society of Food Science and Technology) (South Korea)

Health Serv. Rep.
Health Services Reports (U.S.)
Supersedes HSMHA Health Rep.

Heat Transfer - Jap. Res.
Heat Transfer - Japanese Research (U.S.)

Heft Unfallheilk.
Heft zur Unfallheilkunde; Beiheft zur Monatsschrift fuer Unfallheilkunde, Versicherungs-, Versorgungs- und Verkehrsmedizin (West Germany)

Hiroshima Daigaku Genbaku Hoshano Igaku Kenkyusho Nenpo
Hiroshima Daigaku Genbaku Hoshano Igaku Kenkyusho Nenpo (Proceedings of the Research Institute for Nuclear Medicine and Biology, Hiroshima University) (Japan)

Hiroshima Nogyo Tanki Daigaku Kenkyu Hokoku
Hiroshima Nogyo Tanki Daigaku Kenkyu Hokoku (Bulletin of the Hiroshima Agricultural College) (Japan)

Hokkaido Daigaku Kogakubu Kenkyu Hokoku
Hokkaido Daigaku Kogakubu Kenkyu Hokoku (Bulletin of the Faculty of Engineering, Hokkaido University) (Japan)

HortScience
HortScience (U.S.)

Hoshasen Butsuri Kenkyu
Hoshasen Butsuri Kenkyu (Japan)

Hum. Pathol.
Human Pathology (U.S.)

I

Ind. At. Spatiales
Industries Atomiques et Spatiales (France)
Supersedes Ind. At. and Sci. Ind. Spatiales, beginning Jan. 1972

Ind.-Elek. Elektron.
Industrie-Elektrik und Elektronik (West Germany)

Indiana Univ. Math. J.
Indiana University Mathematics Journal (U.S.)
Supersedes J. Math. Mech.

Indian Eng.
Indian Engineer (India)

Indian Geotech. J.
Indian Geotechnical Journal (India)

Indian J. Anim. Sci.
Indian Journal of Animal Sciences (India)

Indian J. Biochem. Biophys.
Indian Journal of Biochemistry and Biophysics (India)
Continues Indian J. Biochem.

Indian J. Vet. Sci. Anim. Husb.
Indian Journal of Veterinary Science and Animal Husbandry (India)
Continued as Indian J. Anim. Sci.

Indian Mining Eng. J.
Indian Mining and Engineering Journal (India)

Ingegneria
Ingegneria (Italy)

Instrum. Power Ind.
Instrumentation in the Power Industry (U.S.)

- Int. Comp. Law Quart.
International and Comparative Law Quarterly, The
(England)
- Int. J. Environ. Anal. Chem.
International Journal of Environmental Analytical Chemistry
(England)
- Int. J. Mech. Sci.
International Journal of Mechanical Sciences (England)
- Int. Ver. Theor. Angew. Limnol., Mitt.
International Vereinigung fuer Theoretische und Angewandte
Limnologie, Mitteilungen (West Germany)
- Iwate Igaku Zasshi
Iwate Igaku Zasshi (Journal of the Iwate Medical Association)
(Japan)
- Izotoptekhnika
Izotoptekhnika (Hungary)
Supersedes Atomtech. Tajek.
- Izv. Akad. Nauk Tadzh. SSR, Otd. Fiz.-Mat. Geol.-Khim. Nauk
Izvestiya Akademii Nauk Tadzhikskoi SSR, Otdelenie
Fiziko-Matematicheskikh i Geologo-Khimicheskikh Nauk
(Tadzhik S.S.R.)
- Izv. Estestvennonauch. Inst. Perm. Gos. Univ.
Izvestiya Estestvennonauchnogo Instituta pri Permskom
Gosudarstvennom Universitete (U.S.S.R.)
- Izv. Leningrad. Elektrotekh. Inst.
Izvestiya Leningradskogo Elektrotekhnicheskogo Instituta
(U.S.S.R.)
- Izv. Vyssh. Ucheb. Zaved., Neft Gaz
Izvestiya Vysshikh Uchebnykh Zavedenii, Neft i Gaz
(Azerbaijani S.S.R.)
- J. Allergy Clin. Immunol.
Journal of Allergy and Clinical Immunology (U.S.)
Supersedes J. Allergy
- J. Amer. Concr. Inst.
Journal of the American Concrete Institute (U.S.)
- J. Chem. Soc., Chem. Commun.
Journal of the Chemical Society (London),
Chemical Communications (England)
Supersedes J. Chem. Soc., D
- J. Chem. Soc., Dalton Trans.
Journal of the Chemical Society (London), Dalton
Transactions (England)
Supersedes in part J. Chem. Soc., A
- J. Chem. Soc., Faraday Trans., I
Journal of the Chemical Society (London), Faraday
Transactions, I (England)
Supersedes in part J. Chem. Soc., A and Trans. Faraday Soc.
- J. Chem. Soc., Faraday Trans., II
Journal of the Chemical Society (London), Faraday
Transactions, II (England)
Supersedes in part J. Chem. Soc., A and Trans. Faraday Soc.
- J. Chem. Soc., Perkin Trans., II
Journal of the Chemical Society (London), Perkin
Transactions, II (England)
Supersedes J. Chem. Soc., B
- J. Chromatogr. Sci.
Journal of Chromatographic Science (U.S.)
Supersedes Journal of Gas Chromatography
- J. Conn. State Dent. Ass.
Journal of the Connecticut State Dental Association (U.S.)
- J. Coord. Chem.
Journal of Coordination Chemistry (England)
- J. Eur. Toxicol.
Journal Européen de Toxicologie (France)
- J. Fla. Med. Ass.
Journal of the Florida Medical Association (U.S.)
- J. Ga. Entomol. Soc.
Journal of the Georgia Entomological Society (U.S.)
- J. Geol. Soc. India
Journal of the Geological Society of India (India)
- Jinruigaku Zasshi
Jinruigaku Zasshi (The Journal of the Anthropological
Society of Japan) (Japan)
- J. Jap. Soc. Appl. Phys., Suppl.
Journal of the Japan Society of Applied Physics, Supplement
(Japan)
- J. Korean Inst. Electron. Eng.
Journal of the Korean Institute of Electronics
Engineers, The (Jeonja-gonghakhoe-ji) (South Korea)
- J. Microscopy
Journal of Microscopy (England)
Supersedes J. Roy. Microscop. Soc.
- J. Non-Cryst. Solids
Journal of Non-Crystalline Solids (Netherlands)
- Joekull (Reykjavik)
Joekull (Reykjavik) (Iceland)
- J. Ont. Dent. Ass.
Journal of the Ontario Dental Association, The (Canada)
- J. Paint Technol.
Journal of Paint Technology (U.S.)
- J. Periodontal Res.
Journal of Periodontal Research (Denmark)
- J. Phys. Chem. Ref. Data
Journal of Physical and Chemical Reference Data (U.S.)
- J. Soc. Dyers Colour.
Journal of the Society of Dyers and Colourists (England)
- J. Zool.
Journal of Zoology (England)
- Kagaku Keizai
Kagaku Keizai (Chemical Industrial Economy) (Japan)
- KFK Nachr.
KFK (Kernforschungszentrum Karlsruhe) Nachrichten
(West Germany)
- Kibern. Vychisl. Tekh.
Kibernetika i Vychislitel'naya Tekhnika (Ukrainian S.S.R.)
- Kim. Sanayi
Kimya ve Sanayi (Turkey)

Koku Igaku Jikkentai Hokoku
Koku Igaku Jikkentai Hokoku (The Reports of the Aero
Medical Lab.) (Japan)

Komety Meteory
Komety i Meteory (Tadzhik S.S.R.)

Konstr. Ing. Ber.
Konstruktiver Ingenieurbau Berichte (West Germany)

Konstr. Mater. Osn. Grafita
Konstruktsionnye Materialy na Osnove Grafita (U.S.S.R.)

Korean Biochem. J.
Korean Biochemical Journal, The (Hankuk Saenghwahakhoe-ji)
(South Korea)

Korean J. Hort. Sci.
Korean Journal of Horticultural Science (Hankuk
Wonyehakhoe-ji) (South Korea)

Koryo
Koryo (Aromatics) (Japan)

Kotsu Igaku
Kotsu Igaku (The Journal of Transportation Medicine) (Japan)

KSB Tech. Ber.
KSB (Klein, Schanzlin, Becker) Technische Berichte
(West Germany)

Kuki Chowa To Reito
Kuki Chowa To Reito (Air Conditioning and Refrigeration)
(Japan)

Kvantovaya Elektron.
Kvantovaya Elektronika (Ukrainian S.S.R.)

L

Lesovedenie
Lesovedenie (U.S.S.R.)

Life Sci. Space Res.
Life Sciences and Space Research (Netherlands)

Log Anal.
Log Analyst (U.S.)

M

Magy. Allami Eotvos Lorand Geofiz. Int. Evi Jelentese
Magyar Allami Eotvos Lorand Geofizikai Intezet Evi
Jelentese (Hungary)

Magy. Asvanyolaj Foldgaz Kiserl. Intez. Kiadv.
Magyar Asvanyolaj es Foldgaz Kiserleti Intezet Kiadvanyai
(Hungary)

Magy. Textiltech.
Magyar Textiltechnika (Hungary)

Magy. Tud. Akad. Biol. Tud. Oszt. Kozlem.
Magyar Tudomanyos Akademia Biologiai Tudomanyok
Osztalyanak Kozlemenyei (Hungary)

Maintenance
Maintenance (France)

Mannesmann Forschungsber.
Mannesmann Forschungsberichte (West Germany)

Mashinostroene
Mashinostroene (Bulgaria)

Mater. Tech.
Materiaux et Techniques (France)

Medd. Nor. Skogforsoeksv.
Meddelelser fra det Norske Skogforsoeksvesen (Norway)

Med. Lab. Technol.
Medical Laboratory Technology (England)
Supersedes J. Med. Lab. Technol.

Med. Microbiol. Immunol.
Medical Microbiology and Immunology (West Germany)
Supersedes Z. Med. Mikrobiol. Immunol.

Mem. Accad. Sci. Torino, Cl. Sci. Fis., Mat. Natur.
Memoire della Accademia delle Scienze di Torino, Classe
di Scienze Fisiche, Matematiche e Naturali (Italy)

Mem. Real Acad. Cienc. Artes Barcelona
Memorias de la Real Academia de Ciencias y Artes de
Barcelona (Spain)

Metallges., Rev. Aktiv.
Metallgesellschaft, Review of the Activities (West German)
German Edition issued as Metallges., Mitt. Arbeitsbereich

Metallofizika
Metallofizika (Ukrainian S.S.R.)

Metall-Reinig. Vorbehandl.
Metall-Reinigung und Vorbehandlung (West Germany)

Metallurgie (Mons, Belg.)
Metallurgie (Mons, Belgium) (Belgium)

Metalwork. Econ.
Metalworking Economics (U.S.)

Methods Biochem. Anal.
Methods of Biochemical Analysis (U.S.)

Mich. Med.
Michigan Medicine (U.S.)

Miner. Sci. Eng.
Minerals Science and Engineering (South Africa)

Minneapolis Dist. Dent. J.
Minneapolis District Dental Journal, The (U.S.)

Mitsubishi Seiko Giho
Mitsubishi Seiko Giho (Japan)

Mitt. Astron. Ges.
Mitteilungen der Astronomischen Gesellschaft (West German)

Modif. Polim. Mater.
Modifikatsiya Polimernykh Materialov (Latvian S.S.R.)

Moessbauer Eff. Methodol., Proc. Symp.
Moessbauer Effect Methodology, Proceedings of a
Symposium (U.S.)

Morsk. Hidrofiz. Issled.
Morskie Hidrofizicheskie Issledovaniya (U.S.S.R.)

Munca Sanit.
Munca Sanitara (Romania)

N

Nagoya Daigaku Kankyo Igaku Kenkyusho Nempo
Nagoya Daigaku Kankyo Igaku Kenkyusho Nempo (Japan)
English Edition issued as Annu. Rep. Res. Inst. Environ.
Med., Nagoya Univ.

Nagoya-Shi Kogyo Kenkyusho Kenkyu Hokoku
Nagoya-Shi Kogyo Kenkyusho Kenkyu Hokoku (Research Reports
for the Nagoya Municipal Industrial Research Institute)
(Japan)

Nat. Bur. Stand. (U.S.), Spec. Publ.
National Bureau of Standards (U.S.), Special
Publication (U.S.)

Natura (Plovdiv)
Natura (Plovdiv) (Bulgaria)

Nauch. Tr. Mosk. Inzh.-Ekon. Inst.
Nauchnye Trudy Moskovskogo Inzhenerno-Ekonomicheskogo
Instituta (U.S.S.R.)

Nippon Hifuka Gakkai Zasshi
Nippon Hifuka Gakkai Zasshi (Journal of the Japanese
Dermatological Association) (Japan)

Nogyo Gijutsu Kenkyusho Hokoku, B
Nogyo Gijutsu Kenkyusho Hokoku, B. Dojo Hiryo
(Bulletin of the National Institute of Agricultural Science,
Series B, Soils and Fertilizers) (Japan)

Nord. Hydrol.
Nordic Hydrology (Denmark)

Norsk Vet.-Tidsskr.
Norsk Veterinaer-Tidsskrift (Norway)

Nouv. Presse Med.
Nouvelle Presse Medicale, La (France)
Supersedes Presse Medicale, La

Novosti Med. Priborostr.
Novosti Meditsinskogo Priborostroeniya (U.S.S.R.)

Nuisances Environ.
Nuisances et Environment (France)

N. Z. J. Agr. Res.
New Zealand Journal of Agricultural Research (New Zealand)

O

Obz. Mat. Fiz.
Obzornik za Matematiko in Fiziko (Yugoslavia)

Occup. Health Bull.
Occupational Health Bulletin (Canada)

Occup. Health Rev.
Occupational Health Review (Canada)

P

Paraplegia
Paraplegia (Scotland)

Particles Nucl.
Particles and Nuclei (U.S.)

Perspect. Powder Met.
Perspectives in Powder Metallurgy (U.S. - England)

Pharm. Ztg.
Pharmazeutische Zeitung (West Germany)

Planta Med.
Planta Medica (West Germany)

Polimery
Polimery (Poland)

Poluprov. Tekh. Mikroelektron.
Poluprovodnikovaya Tekhnika i Mikroelektronika
(Ukrainian S.S.R.)

Polyar. Siyaniya
Polyarnye Siyaniya (U.S.S.R.)

Polym. J.
Polymer Journal (Japan)

Polytech. Tijdschr., Elektrotech./Electron.
Polytechnisch Tijdschrift, Elektrotechniek/Elektronica
(Netherlands)

Polytech. Tijdschr., Procestech.
Polytechnisch Tijdschrift, Procestechiek (Netherlands)

Praeventivmedizin
Praeventivmedizin (Switzerland)

Prax. Penumol.
Praxis der Penumologie (West Germany)

Probl. Fiz. Elem. Chastits At. Yadra
Problemy Fisiki Elementarnykh Chastits i Atomonogo
Yadra (U.S.S.R.)

Progr. Fish-Cult.
Progressive Fish-Culturist, The (U.S.)

Pr. Stud. Zakl. Badan Nauk. Gorn. Okregu Przem. Pol. Akad. Nauk
Prace i Studia Zakladu Badan Naukowych Gornoslaskiego Okregu
Przemyslowego Polskiej Akademii Nauk (Poland)

Publ. Tartu. Astrofiz. Observ.
Publikatsii Tartuskoj Astrofizicheskoi Observatorii
(Estonian S.S.R.)

Pulp Pap. Mag. Can.
Pulp and Paper Magazine of Canada (Canada)

Purasuchikkusu
Purasuchikkusu (Japan Plastics) (Japan)

R

Radiat. Data Rep.
Radiation Data and Reports (U.S.)
Supersedes Radiological Health Data and Reports

Radiats. Tekh.
Radiatsionnaya Tekhnika (U.S.S.R.)

Rayonnements Ionis. Tech. Mes. Prot.
Rayonnements Ionisants Techniques de Mesures et de
Protection (France)

Rend. Semin. Fac. Sci. Univ. Cagliari
Rendiconti del Seminario della Facolta di Scienze della
Universita di Cagliari (Italy)

Rend. Soc. Ital. Mineral. Petrologia
Rendiconti della Societa Italiana de Mineralogia e
Petrologia (Italy)

Res. Commun. Chem. Pathol. Pharmacol.
Research Communications in Chemical Pathology and
Pharmacology (U.S.)

Res. Rep. Fish. Comm. Oreg.
Research Reports of the Fish Commission of Oregon (U.S.)

Rev. A
Revue A (Institut Belge de Regulation et d'Automatisme,
Brussels) (Belgium)

Rev. Anal. Chem.
Reviews in Analytical Chemistry. International
Quarterly Scientific Reviews Journal (Israel)

Rev. Coatings Corros.
Reviews on Coatings and Corrosion (Israel)

Rev. Coroz.
Revista de Corozilune (Romania)

Rev. Epidemiol. Med. Soc. Sante Publique
Revue d'Epidemiologie Medecine Sociale et Sante Publique
(France)
Continues Rev. Hyg. Med. Soc.

Rev. Gaucha Odontol.
Revista Gaucha de Odontologia (Brazil)

Rev. High-Temp. Mater.
Reviews on High-Temperature Materials (Israel)

Rev. Soc. Roy. Belge Ing. Ind.
Revue de la Societe Royale Belge des Ingenieurs et des
Industriels (Belgium)

Ric. Mat.
Ricerche di Matematica (Italy)

Rinsho Hinyokika
Rinsho Hinyokika (Japan)

Rinsho Kagaku
Rinsho Kagaku (Japan)

Rocz. Inst. Przem. Miecz.
Roczniki Instytutu Przemyslu Mleczarskiego (Poland)

Russ. Eng. J. (Engl. Transl.)
Russian Engineering Journal (English Translation of
Vestnik Mashinostroeniya) (England)

S

Sanit. Heizungstech.
Sanitaer - Heizungstechnik (West Germany)

Sanka To Fujinka
Sanka To Fujinka (Obstetrics and Gynecology, The) (Japan)

Sb. Nauch. Tr., Fiz.-Tekh. Inst. Nizk. Temp., Akad. Nauk
Ukr. SSR
Sbornik Nauchnykh Trudov, Fiziko-Tekhnicheskii Institut
Nizhikh Temperatur, Akademiya Nauk Ukrainskoi SSR
(Ukrainian S.S.R.)

Sb. Tr. Klyuchevsk. Zavoda Ferrosplavov
Sbornik Trudov Klyuchevskogo Zavoda Ferrosplavov (U.S.S.R.)

Scand. J. Dent. Res.
Scandinavian Journal of Dental Research (Denmark)

Schiff Hafen
Schiff und Hafen (West Germany)

Schweisstechnik (Berlin)
Schweisstechnik (Berlin) (East Germany)

Schweiz. Z. Hydrol.
Schweizerische Zeitschrift fuer Hydrologie (Switzerland)

Sci. Publ., Res. Inst. Radiol. Radiat. Hyg.
Scientific Publications, Research Institute of Radiology
and Radiation Hygiene (Bulgaria)

Sediment. Geol.
Sedimentary Geology (Netherlands)

Seifen-Dele-Fette-Wachse
Seifen-Dele-Fette-Wachse (West Germany)

Selec. Annu. Rev. Anal. Sci.
Selected Annual Reviews of the Analytical Sciences
(England)

Shigen
Shigen (Resources) (Japan)

Shinryo To Shinyaku
Shinryo To Shinyaku (Medical Consultation and New
Remedies) (Japan)

Shizuoka-Ken Kogyo Shikenjo Hokoku
Shizuoka-Ken Kogyo Shikenjo Hokoku (Japan)

Shokuhin Kogyo
Shokuhin Kogyo (Food Industry) (Japan)

Shonika Rinsho
Shonika Rinsho (Japanese Journal of Pediatrics) (Japan)

Silk Rayon Ind. India
Silk and Rayon Industries of India (India)

Silva Fenn.
Silva Fennica (Finland)

Slevarenstvi
Slevarenstvi (Czechoslovakia)

Solvent Extr. Rev.
Solvent Extraction Reviews (U.S.)

Sov. Rubber Technol. (Engl. Transl.)
Soviet Rubber Technoigy (English Translation of
Kauchuk i Rezina) (England)

Statens Offentliga Utredn.
Statens Offentliga Utredningar (Sweden)

Stimulation Newslett.
Stimulation Newsletter (West Germany)

Stoma
Stoma (West Germany)

Stoma (Thessaloniki)
Stoma (Thessaloniki) (Greece)

Stomatol. Glas. Srb.
Stomatoloski Glasnik Srbije (Yugoslavia)

Struct. Eng.
Structural Engineer, The (England)

Stud. Univ. Babes-Bolyai, Ser. Geol.-Mineral.
Studia Universitatis Babes-Bolyai, Series Geologia-
Mineralogia (Romania)

Sumitomo Jukikai Giho
Sumitomo Jukikai Giho (Japan)

Sv. Kraftverksfoereningens Publ.
Svenska Kraftverksfoereningens Publikationer (Sweden)

Swed. J. Agr. Res.
Swedish Journal of Agricultural Research (Sweden)
Supersedes Lantbrukshogs. Ann.

Syoni Shikagaku Zasshi
Syoni Shikagaku Zasshi (Japan)

T

Taehan Saengri Hakhoe Chi
TTaehan Saengri Hakhoe Chi (Korean Journal of Physiology)
(South Korea)

Taiki Osen Kenkyu
Taiki Osen Kenkyu (Journal of the Japan Society of
Air Pollution) (Japan)

Takeda Kenkyusho-Ho
Takeda Kenkyusho-Ho (Journal of the Takeda Research
Laboratories) (Japan)

Tenki (Meteorol. Soc. Jap.)
Tenki (Weather (Meteorological Society of Japan))
(Japan)

Tenmon Geppo
Tenmon Geppo (The Astronomical Herald) (Japan)

Tenn. Law Rev.
Tennessee Law Review (U.S.)

Tensai Gijutsu Renraku Kenkyukai Happyo Ronbunshu
Tensai Gijutsu Renraku Kenkyukai Happyo
Ronbunshu (Japan)

Tenside
Tenside (West Germany)

TESLA Electron.
TESLA Electronics (Czechoslovakia)

Ti Ch'iu Wu Li Hsueh Pao
Ti Ch'iu Wu Li Hsueh Pao (Journal of Geophysics) (China)
Discontinued

Tokushuko
Tokushuko (Special Steel) (Japan)

Toyoda Kenkyu Hokoku
Toyoda Kenkyu Hokoku (Japan)

Toyo Shokuhin Kogyo Tanki Daigaku, Toyo Shokuhin
Kenkyusho Kenkyu Hokokusho
Toyo Shokuhin Kogyo Tanki Daigaku, Toyo Shokuhin Kenkyusho
Kenkyu Hokokusho (Toyo Junior College of Food Technology
and Toyo Institute of Food Technology, Research Report)
(Japan)

Trans. Amer. Microsc. Soc.
Transactions of the American Microscopical Society (U.S.)

Trans., Gulf Coast Ass. Geol. Soc.
Transactions, Gulf Coast Association of Geologic Societies
(U.S.)

Transp.-Med. Vesti
Transportno-Meditsinski Vesti (Bulgaria)

Transp. Theory Statist. Phys.
Transport Theory and Statistical Physics (U.S.)

Tr. Gor'k. Politekh. Inst.
Trudy Gor'kovskogo Politekhicheskogo Instituta (U.S.S.R.)

Tr. Inst. Mikrobiol. Virusol., Akad. Nauk Kaz. SSR
Trudy Instituta Mikrobiologii i Virusologii Akademiya
Nauk Kazakhskoi SSR (Kazakh S.S.R.)

Tr. Inst. Org. Katal. Elektrokhim., Akad. Nauk Kaz. SSR
Trudy Instituta Organicheskogo Kataliza i Elektrokhimii,
Akademiya Nauk Kazakhskoi SSR (Kazakh S.S.R.)

Tr. Khim. Khim. Tekhnol.
Trudy po Khimii i Khimicheskoi Tekhnologii (U.S.S.R.)

Tr. Leningrad. Khim.-Farm. Inst.
Trudy Leningradskogo Khimicheskogo-Farmatsevticheskogo
Instituta (U.S.S.R.)

Tr. Nauch.-Issled. Gidrometeorol. Inst. (Alma-Ata)
Trudy Nauchno-Issledovatel'skogo Gidrometeorologicheskogo
Instituta (Alma-Ata) (Kazakh S.S.R.)

Tr. Novocherkassk. Politekh. Inst.
Trudy Novocherkasskogo Politekhicheskogo Instituta
(U.S.S.R.)

Tr. Radiotekh. Inst.
Trudy Radiotekhnicheskogo Instituta (U.S.S.R.)

Tr. Severokavkaz. Gornomet. Inst.
Trudy Severokavkazskogo Gornometallurgicheskogo Instituta
(U.S.S.R.)

Tr. Tambov. Inst. Khim. Mashinostr.
Trudy Tambovskogo Instituta Khimicheskogo
Mashinostroeniya (U.S.S.R.)

Tr. Ufim. Nauch.-Issled. Inst. Gig. Profzabol.
Trudy Ufimskogo Nauchno-Issledovatel'skogo Instituta
Gigieny i Profzabolevani (U.S.S.R.)

Tr. Voronezh. Gos. Univ.
Trudy Voronezhskogo Gosudarstvennogo Universiteta (U.S.S.R.)

Tr., Vses. Nauch.-Issled. Inst. Prir. Gazov
Trudy, Vsesoyuznyi Nauchno-Issledovatel'skii Institut
Prirodnykh Gazov (U.S.S.R.)

Tsuchi To Kiso
Tsuchi To Kiso (Soil Mechanics and Foundation Engineering)
(Japan)

U

Uch. Zap., Mosk. Gos. Pedagog. Inst.
Uchenye Zapiski, Moskovskii Gosudarstvennyi Pedagogicheskii
Institut (U.S.S.R.)

Umwelt
Umwelt (West Germany)

Union Agr.
Union Agriculture (France)

Utsunomiya Daigaku Kyoikugakubu Kiyo
Utsunomiya Daigaku Kyoikugakubu Kiyo (Japan)

Uzb. Geol. Zh.
Uzbekskii Geologicheskii Zhurnal (Uzbek S.S.R.)

V

Vaigyanik
Vaigyanik (India)

Va. Dent. J.
Virginia Dental Journal (U.S.)

VDI (Ver. Deut. Ing.) Ber.
VDI (Verein Deutscher Ingenieure) Berichte (West Germany)

VDI (Ver. Deut. Ing.)-Forsch.
VDI (Verein Deutscher Ingenieure)-Forschungsheft (West
Germany)

Versicherungswirtschaft
Versicherungswirtschaft (West Germany)

Vestn. Khar'kov. Univ., Astron.
Vestnik Khar'kovskogo Universiteta, Astronomiya
(Ukrainian S.S.R.)

Vestn. Khar'kov. Univ., Ser. Biol.
Vestnik Khar'kovskogo Universiteta, Seriya
Biologicheskaya (Ukrainian S.S.R.)

Vestn. Khar'kov. Univ., Ser. Geol.
Vestnik Khar'kovskogo Politehnicheskogo Instituta, Seriya
Geologicheskaya (Ukrainian S.S.R.)

Vestn. Khar'kov. Univ., Ser. Khim.
Vestnik Khar'kovskogo Politehnicheskogo Instituta, Seriya Khimicheskaya
(Ukrainian S.S.R.)

Vestn. Kiev. Politekh. Inst., Ser. Mashinostr.
Vestnik Kievskogo Politehnicheskogo Instituta,
Seriya Mashinostroeniya (Ukrainian S.S.R.)

Vestn. Mashinostr.
Vestnik Mashinostroeniya (U.S.S.R.)
Issued in English translation as Russ. Eng. J.

Vestn. Zool.
Vestnik Zoologii (Ukrainian S.S.R.)

Vestsi Akad. Navuk Belarus. SSR, Ser. Fiz.-Energ. Navuk
Vestsi Akademii Navuk Belaruskai SSR, Seriya Fizika-
Energetichnykh Navuk (Belorussian S.S.R.)

Vet. Pathol.
Veterinary Pathology (Switzerland)
Continues Pathologia Veterinaria

Vie Milieu
Vie et Milieu (France)

Vop. Tekh. Teplofiz.
Voprosy Tekhnicheskoi Teplofiziki (Ukrainian S.S.R.)

W

Wasser Boden
Wasser und Boden (West Germany)

Weltraumfahrt Raketentech.
Weltraumfahrt und Raketentechnik (West Germany)

World Miner. Metals
World Minerals and Metals (England)

Y

Yale Sci. Mag.
Yale Scientific Magazine (Title varies; may be Yale
Scientific or Yale Scientific Magazine) (U.S.)

Yamaguchi Igaku
Yamaguchi Igaku (Yamaguchi Medicine) (Japan)

Z

Zavarivac
Zavarivac (Yugoslavia)

Zb. Meteorol. Hidrol. Rad.
Zbornik Meteoroloskih i Hidroloskih Radova (Yugoslavia)

Z. Erkr. Atmungsorgane Folia Bronchol.
Zeitschrift fuer Erkrankungen der Atmungsorgane mit
Folia Bronchologica (East Germany)

Zesz. Nauk. Akad. Gorn.-Hutn., Cracow, Mat., Fiz., Chem.
Zeszyty Naukowe Akademii Gorniczo-Hutniczej, Cracow,
Matematyka, Fizyka, Chemia (Poland)

Zhivotnovud. Nauki
Zhivottnovudni Nauki (Bulgaria)

Z. Immunitaetsforsch., Exp. Klin. Immunol.
Zeitschrift fuer Immunitaetsforschung, Experimentelle und
Klinische Immunologie (West Germany)
Supersedes Z. Immunitaetsforsch., Allergie Klin. Immunol.

Z. Neurol.
Zeitschrift fuer Neurologie (West Germany)
Supersedes Deut. Z. Nervenheilk.

Zool. Pol.
Zoologica Poloniae (Poland)

Z. Psychosom. Med.
Zeitschrift fuer Psychosomatische Medizin (West Germany)

Z. Werkstofftech.
Zeitschrift fuer Werkstofftechnik (West Germany)

Z. Wiss. Biol., Abt. A
Zeitschrift fuer Wissenschaftliche Biologie, Abteilung A.
Zeitschrift fuer Morphologie und Oekologie der Tiere
(West Germany)



AEC SYMPOSIUM SERIES

Available for \$6.00, except as noted, from the National Technical Information Service, U. S. Department of Commerce, Springfield, Virginia 2215

- 1 Progress in Medical Radioisotope Scanning (TID-7673), 1963
- 2 Reactor Kinetics and Control (TID-7662), 1964
- 3 Dynamic Clinical Studies with Radioisotopes (TID-7678), 1964
- 4 Noise Analysis in Nuclear Systems (TID-7679), 1964
- 5 Radioactive Fallout from Nuclear Weapons Tests (CONF-765), 1965
- 6 Radioactive Pharmaceuticals (CONF-651111), 1966
- 7 Neutron Dynamics and Control (CONF-650413), 1966
- 8 Luminescence Dosimetry (CONF-650637), 1967
- 9 Neutron Noise, Waves, and Pulse Propagation (CONF-660206), 1967
- 10 Use of Computers in Analysis of Experimental Data and the Control of Nuclear Facilities (CONF-660527), 1967
- 11 Compartments, Pools, and Spaces in Medical Physiology (CONF-661010), 1967
- 12 Thorium Fuel Cycle (CONF-660524), 1968
- 13 Radioisotopes in Medicine: In Vitro Studies (CONF-671111), 1968
- 14 Abundant Nuclear Energy (CONF-680810), 1969
- 15 Fast Burst Reactors (CONF-690102), 1969
- 16 Biological Implications of the Nuclear Age (CONF-690303), 1969
- 17 Radiation Biology of the Fetal and Juvenile Mammal (CONF-690501), 1969
- 18 Inhalation Carcinogenesis (CONF-691001), 1970
- 19 Myeloproliferative Disorders of Animals and Man (CONF-680529), 1970, \$9.00
- 20 Medical Radionuclides: Radiation Dose and Effects (CONF-691212), 1970
- 21 Morphology of Experimental Respiratory Carcinogenesis (CONF-700501), 1970
- 22 Precipitation Scavenging (1970) (CONF-700601), 1970
- 23 Neutron Standards and Flux Normalization (CONF-701002), 1971
- 24 Survival of Food Crops and Livestock in the Event of Nuclear War (CONF-700909), 1971, \$9.00
- 25 Biomedical Implications of Radiostrontium Exposure (CONF-710201), 1972
- 26 Radiation-Induced Voids in Metals (CONF-710601), 1972, \$9.00